THE COVARIANT LATTICE CONSTRUCTION OF FOUR-DIMENSIONAL STRINGS^{*}

W. Lerche[†]

and

A. N. Schellekens

CERN, 1211 Geneva 23, Switzerland

ABSTRACT

An elementary introduction is given to the covariant lattice construction of lower dimensional heterotic strings, and to the concepts on which it is based. This includes a review of properties of lattices, torus compactification and conformal field theory. Some more advanced topics, such as the bosonization of the superghost system, are also reviewed.

^{*} Based on lectures by the second author at the Summer School on Strings and Superstrings, Poiana Braşov, Romania, September 1987.

[†] Supported by the Max-Planck-Gesellschaft, Munich, FRG.

Address after 1st September 1988: California Institute of Technology, Pasadena, CA 91125, USA.

THE COVARIANT LATTICE CONSTRUCTION OF FOUR-DIMENSIONAL STRINGS[★]

W. Lerche[†]

and

A. N. Schellekens

CERN, 1211 Geneva 23, Switzerland

ABSTRACT

An elementary introduction is given to the covariant lattice construction of lower dimensional heterotic strings, and to the concepts on which it is based. This includes a review of properties of lattices, torus compactification and conformal field theory. Some more advanced topics, such as the bosonization of the superghost system, are also reviewed.

★ Based on lectures by the second author at the Summer School on Strings and Superstrings, Poiana Braşov, Romania, September 1987.

† Supported by the Max-Planck-Gesellschaft, Munich, FRG.

Address after 1st September 1988: California Institute of Technology, Pasadena, CA 91125, USA.

THE COVARIANT LATTICE CONSTRUCTION OF FOUR-DIMENSIONAL STRINGS

W. LERCHE AND A.N. SCHELLEKENS

CERN, 1211 Geneva 23, Switzerland

1. Introduction

The main goal of string theory is an ambitious one: to obtain a consistent theory of all interactions, including gravity. Just a few years ago it seemed that gravity was all it could describe, but that has changed drastically with the discovery of the heterotic string [1]. Although at first it seemed that the theory would fail its first confrontation with experiment by predicting the wrong space-time dimension, that too changed very quickly after its birth. Indeed, what once may have seemed an insurmountable obstacle is now a gaping hole: not only can we construct string theories in four dimensions, but it is also so easy that we can make them in embarrassing abundance. Meanwhile the miracles that started the excitement have survived, although we are now so used to them that we tend to forget them: chiral string theories in four dimensions exist and their spectra have the same general features (gauge groups, representations and family replication) as that of the standard model. If all this is realized in a class of theories that holds the promise of offering a finite and consistent theory of quantum gravity, then there is certainly reason for excitement. The dream of having a *unique* theory of everything must be shelved for the moment, but fortunately uniqueness is the aspect that we can most easily live without. On the other hand, giving up consistency would be fatal.

The floodgates of four-dimensional string construction were first opened in [2]. The initial trickle of Calabi-Yau compactifications quickly grew into a raging torrent, which broadened into orbifold compactifications [3]. The theories that are now usually called four-dimensional strings are mathematically not very different from these early ideas, and it doesn't make much sense to try to make a distinction. If there is a difference, then it is one of philosophy rather than physics. Whereas initially one tried to compactify ten-dimensional strings, nowadays the emphasis is more on constructing string theories directly in four dimensions. We believe that the idea of compactification is too limited, and too much linked to notions of space-time that may be unsuitable for string theory, to be of lasting value. But all this is just a matter of interpretation, and does not influence the facts, such as the spectrum and the S-matrix.

The change of philosophy from compactification to four-dimensional strings started with the seminal paper of Narain [4], and gained momentum due to the work of Kawai et. al. Soon thereafter many ways of constructing heterotic strings in four dimensions were discovered [5-8], the oldest being fermionic constructions, the covariant lattice construction, and asymmetric orbifolds, which are now understood to be close relatives.

Several other ideas have surfaced more recently [9-12]. For recent reviews of aspects of string vacua see e.g. [13], [14].

These lectures are devoted to a detailed description of the covariant lattice construction and its underlying principles. The construction is based on the bosonic formulation of the fermionic string and its ghost system [15]. The possibility of using this formalism for the construction of new modular invariant string theories was first discussed in [16], and it was used for four-dimensional strings in [6]. A related idea is the “charge lattice” of [5], which these authors obtained by bosonizing their fermionic construction. The result is a light-cone version of covariant lattices.

Although we tried to keep these notes as elementary as possible, some basic knowledge of string theory is assumed, and may be gotten from one of the many reviews of the subject, some of which are listed under [17]. In the next chapter we begin with an introduction to the principles of string construction. This section serves also a guide to the rest of the paper.

2. Basic principles of string construction

The most fundamental quantity in any string theory is a function $X^\mu(\sigma^0, \sigma^1)$ that specifies the space-time position of every point of the string (denoted by σ^1) for any value of the proper-time, σ^0 . Here μ labels the space-time coordinates. The space-time dimension is d , and our interest will ultimately be in the value $d = 4$. Throughout this paper we will only consider *closed* strings, which means that $X^\mu(\sigma^0, 0) = X^\mu(\sigma^0, \pi)$ (the choice of the σ_1 -interval is a matter of convention). When it propagates through space-time the string sweeps out a two-dimensional surface parametrized by σ^0 and σ^1 . This surface is called the *world-sheet*, in analogy to world-lines for point particles. For non-interacting closed strings it has the shape of a tube. If strings interact they do so by splitting and joining, and as a result they sweep out world-sheets that look like complicated pieces of plumbing.

The modern way of describing strings is to regard them as two-dimensional field theories. One regards $X^\mu(\sigma_0, \sigma_1)$ as a set of d scalar fields, which are functions of a two-dimensional coordinate $\sigma^\alpha, \alpha = 0, 1$. Once one realizes that one is dealing with two-dimensional field theories, one is more or less naturally led to the questions: Which conditions should be imposed on such a field theory for it to describe a consistent string theory? And which two-dimensional field theories satisfy these conditions?

One has to be careful in formulating these conditions, because many claims about uniqueness of string theory were based on implicitly or explicitly imposed conditions which were unnecessarily strong. So we should begin by carefully listing and examining the conditions. Our list of what we believe to be necessary conditions for d -dimensional strings is:

1. The presence of d unconstrained scalar fields X^μ
2. Reparametrization invariance
3. Conformal invariance
4. Modular invariance

5. World sheet supersymmetry and superconformal invariance (for fermionic strings).

These are all conditions imposed on the world-sheet field theory. Remarkably, if one satisfies them one is guaranteed not to be confronted with most of the disasters that can occur in field theory in d -dimensional space-time. Reparametrization invariance means that the description of string propagation does not depend on how one chooses the coordinates σ_0 and σ_1 on the world-sheet. This has a simple interpretation in two-dimensional field theory language: it's just two-dimensional general coordinate invariance. The first two points are so basic to string theory that one would not even want to consider what would go wrong if they were not satisfied. For the last three conditions (which we will discuss in more detail in a moment) one can identify some desirable space-time properties that follow from them.

Conformal invariance guarantees Lorentz-invariance and absence of ghosts, general coordinate invariance in space-time and gauge invariance (if the theory has massless vector bosons, as is usually the case). Although it has been the subject of debate whether one can make sense of string theory without it, it seems clear that one gets into serious technical complications if one abandons it. The attempts to abandon it were motivated by the fact that conformal invariance restricts, naively at least, the space-time dimension to 26 for bosonic strings, and to 10 for fermionic ones. Since there are now other ways to get to lower dimensions, it seems highly preferable to keep it.

One of the nice consequences of modular invariance is that it ensures that chiral fermion representations coming from string theory are anomaly free. This is highly non-trivial, especially in higher dimensions, and the discovery of a rather spectacular cancellation in 10 dimensions [18] was the starting point of the current wave of interest in the subject. The precise relation with modular invariance was only understood much later [19]. Anomalies are certainly not the only problems one can expect in the absence of modular invariance, but it is the easiest one to identify.

World-sheet supersymmetry is essential for fermionic strings, and one of the problems one has in its absence is a violation of Lorentz invariance. We will discuss this in chapter 7. Here we will continue with a more detailed look at conformal and modular invariance.

2.1 CONFORMAL INVARIANCE

A conformal transformation is a coordinate dependent rescaling of the metric on the world-sheet. It is this invariance which makes string theory so simple in comparison with, for example, membranes. The standard two-dimensional action for free bosons,

$$S = \int d^2\sigma \sqrt{-g} g^{\alpha\beta} \partial_\alpha X^\mu \partial_\beta X_\mu,$$

where $g_{\alpha\beta}$ is the world-sheet metric and g its determinant, is not only general coordinate invariant, but has an additional invariance under $g_{\alpha\beta} \rightarrow \Lambda(\sigma^\alpha) g_{\alpha\beta}$. This invariance makes it possible to choose a gauge (light-cone gauge) in which one gets an action which depends only on the $d - 2$ “transverse” (i.e. transforming in the vector representation of the little group $SO(d - 2)$ of a massless particle) degrees of freedom X^i , and which

is non-interacting and manifestly ghost-free. In the absence of conformal invariance there is an extra mode of the metric, the Liouville mode, which enters into the action, and which makes life considerably more complicated. Thus we would like to maintain conformal invariance even if other terms are added to the action.

Although conformal invariance is a symmetry of the classical action, it is afflicted by an anomaly in the quantized theory. There are several contributions to this anomaly. In units in which a scalar contributes 1, the relevant ones are $\frac{1}{2}$ for a Majorana-Weyl fermion, -26 for the ghosts of general coordinate invariance, and +11 for the ghosts of local supersymmetry. Notice that though the graviton has no physical components in two dimensions, it still contributes via its ghosts. There are two obvious ways of cancelling these anomalies :

$N = 0$: 26 scalars + gravity ghosts

$N = 1$: 10 scalars and Majorana Weyl fermions + gravity ghosts + superghosts

Since there is always general coordinate invariance on the world sheet, one always has the corresponding ghost. In these solutions one can replace a scalar by two fermions or vice versa, but that amounts just to two-dimensional bose-fermi equivalence and does not really bring anything new. We have made our choice in such a way that supersymmetry is manifest, but that is not necessary and we will in fact choose a purely bosonic realization in our construction. One can also try to have more supersymmetries, but the resulting theories turn out to be relevant only in two space-time dimensions or less.

At this point we should make a trivial but crucial remark. The traditional interpretation of these solutions is to associate all the scalars with space-time coordinates. In that case the $N = 0$ system lives in 26 dimensions, and the $N = 1$ system in $d = 10$. This is not necessary however. One can equally well associate just a subset of the scalars with space-time coordinates, and regard the others as some sort of “internal” degrees of freedom. Then the numbers 26 and 10 are “maximal” rather than “critical” dimensions. In some cases, and in particular for bosonic strings, this means nothing more than that some dimensions are “compactified”, but for fermionic strings there are more interesting possibilities, as we will see.

A special feature of conformally invariant closed strings is that the modes of vibration of the string can be unambiguously separated into *left-moving* ones and *right-moving* ones. Since the world-sheet theory is non-interacting, and because there are no endpoints of the string that modes can bounce back from, a left-moving mode will move to the left forever. This left/right-mover separation goes so far that one can actually choose *different* conformal systems for them. This leads in an obvious way to three kinds of closed string theory:

Bosonic :	$N = 0$ left-movers;	$N = 0$ right-movers
Heterotic :	$N = 0$ left-movers;	$N = 1$ right-movers
Type II :	$N = 1$ left-movers;	$N = 1$ right-movers

This also specifies our convention for left- and right-movers in the heterotic string.

We will mainly consider heterotic strings in this paper, although everything we say is valid for type II strings as well. The latter seem however phenomenologically less promising at the moment [20], [21]. The heterotic string forces us to make a previous condition more precise. Since it has 26 bosons on the left and only 10 on the right, it is not *a priori* obvious in which dimensions it can exist. The precise requirement on the presence of a set of scalars X^μ is however that both its left and right-moving components are present in the theory. One cannot associate an unpaired left-moving mode with a flat space-time dimension (this would be in conflict with modular invariance). Thus the maximal dimension for the heterotic string is 10. Psychologically it was perhaps this heterotic string asymmetry that lead to a re-evaluation of the role of world-sheet scalars, making four-dimensional strings possible.

2.2 MODULAR INVARIANCE

So far we have imposed conditions to ensure that string theory is well-defined locally on the world-sheet. In the perturbation expansion of string theory one does however encounter world-sheets with non-trivial topology, where the *global* properties of the theory become important. The simplest of these diagrams is the torus, roughly corresponding to a one-loop diagram in field theory. Topologically the description of the multiloop diagrams in the expansion of closed (oriented) strings is very simple: they are all equivalent to multiple tori, and there is just one topology for each order in the expansion. These loops may have external lines (tubes) attached to them. Such surfaces are known as *Riemann surfaces*.

By "modular invariance" we mean absence of global anomalies on any of these surfaces. We also mean *factorization*, i.e. the amplitude for a multiloop diagram should equal the product of its subdiagrams in a limit where it is pulled apart into these subdiagrams. Although one should impose modular invariance as a condition on diagrams with and without external lines, it is sufficient to examine the ones without external lines, because all other diagrams can always be obtained from them as a subdiagram of some higher loop surface.

The exact conditions for modular invariance beyond one loop are not known for fermionic strings, since one does not know enough about the dependence on the "supermoduli", but reasonable *ansätze* exist, and it is under these assumptions that one usually checks modular invariance. For pedagogical reasons we will not say much about higher loop modular invariance except that it is a property of the class of theories we will consider (under the aforementioned assumptions, which have to be made for any fermionic string theory).

One-loop modular invariance is however rather easy to understand. A one-loop diagram is a torus, and it can be represented by points in the complex plane modulo a two-dimensional lattice. This corresponds to a choice of coordinates and a flat, constant metric on the torus. To regard the complex plane modulo a lattice simply means that points that differ by a lattice vector are identified with each other. A two dimensional lattice has two basis vectors, chosen so that all other vectors are integral linear combinations of them. By rotations and scale transformations one can always align one of the basis vectors with the real axis, and scale it to have length 1. The second one can

always be put into the positive upper half plane, where it can be specified by a complex number τ , with $Im(\tau) > 0$. We have now used all the transformations at our disposal to bring the metric in the simplest possible form. The remaining free parameter of the torus, τ , parametrizes metrics that cannot be mapped into each other by general coordinate transformations or Weyl transformations.

Not all these metrics are inequivalent however. Although they cannot be related by *local* transformations, some can be related by *global* ones. For example, if one defines a torus by $\tau' = \tau + 1$ one makes the same identification in the complex plane, but the metric would look rather different, i.e. have a different but equivalent parametrization. On a real-life, donut-shaped torus this transformation corresponds to cutting the torus along a cycle which one cannot be continuously deformed to a point, rotating one edge by 2π , and glueing the edges back together. A torus has two independent cycles of this type, and hence there is a second transformation of τ which leads to equivalent metrics, namely $\tau \rightarrow -\frac{1}{\tau}$. Together these transformations generate the group of *modular transformations*. Although one can write down an explicit expression for all its elements, one needs in practice usually only its two generators,

$$\begin{aligned}\tau &\rightarrow \tau + 1 \\ \tau &\rightarrow -\frac{1}{\tau}\end{aligned}\tag{2.1}$$

To calculate loop-diagrams à la Polyakov [22] one performs a path-integral over all fields in the two-dimensional action (for a given metric on the world-sheet), and furthermore one integrates over all inequivalent metrics. The expression for a one loop diagram is (after some gauge fixing) of the form

$$\int \frac{d^2\tau}{(Im\tau)^2} \mathcal{F}(\tau),$$

where $\mathcal{F}$ is some function which depends on the theory, and the other factor is the invariant integration volume. The function $\mathcal{F}$ (or rather a slightly modified function, defined more precisely in section (4.3)) is called the *partition function*. The issue of modular invariance centers on the boundaries of the τ -integral. If one were to integrate over the entire positive half plane, one would encounter the same metric an infinite number of times. One would like to restrict the integration to a *fundamental domain*, which is defined as a region in the upper half plane which contains every metric just once (the canonical choice is $-1 \leq Re\tau < 1, |\tau| \geq 1$), but this can only be done in a meaningful way if $\mathcal{F}(\tau)$ does not depend on the region one chooses, since otherwise the result would be ambiguous. Thus $\mathcal{F}$ should be invariant under (2.1). This is what is meant by modular invariance.

For an n -loop surface the problem is very similar. Instead of a single complex parameter τ one gets now a complex $n \times n$ matrix Ω , the *period matrix*. For discussions of modular transformations for higher genus Riemann surfaces we refer to [23].

2.3 FOUR DIMENSIONAL STRINGS: THE BASIC IDEA

There is a fairly logical path towards four-dimensional string constructions of the type we are going to consider, and it might be useful to have some idea about this before starting with the more technical aspects. Terms not explained below will be discussed in detail in the remaining chapters.

Consider first the ten-dimensional heterotic string. Its construction has been explained in much detail and with great clarity in [1], which is obligatory literature for anyone interested in string constructions. Here we concentrate on just one aspect, namely the role of the 16 left-moving bosons that are not paired with right-movers into space-time coordinates. Instead one regards them as compactified space-time coordinates (at least in the mathematical sense of the word, because as a physical concept this is somewhat inadequate (and unnecessary)). The center-of-mass coordinate of the string in these 16 dimensions is put in a periodic space (a torus) with periodicities specified by a 16 dimensional lattice. Modular invariance restricts the choice of this lattice (and in particular does not allow an infinite radius torus, which would correspond to flat space). One finds that it must be even and *self-dual*, and such 16-dimensional lattices have been classified, the only possibilities being $E_8 \times E_8$ or a certain subset of the $SO(32)$ weight lattice.

Narain took this construction one logical step further. Instead of insisting on 10-dimensional Lorentz-invariance he required only Lorentz-invariance in $d < 10$ dimensions (we will choose $d = 4$ here, for definiteness and other reasons), and consequently paired only 4 left- and right-moving bosons into space-time coordinates. The remaining ones, 22 left-movers and 6 right-movers are “compactified” (the physical meaning of this terminology has become less and less clear throughout the historical development, but the mathematical description is unambiguous) on a torus of total dimension $22+6$. The condition for modular invariance is that the lattice defining this torus must be even and self-dual with respect to a *lorentzian metric*. There is a continuous infinity of such lattices!

A problem with this approach is that it does not lead to chiral theories. This is related to the fact that the 10 right-moving Majorana-Weyl fermions of the heterotic string have remained untouched. Since they are responsible for the existence of space-time fermions, what happens is that all ten-dimensional space-time fermions which are chiral are trivially compactified on a torus, and it is a well-known fact that this leads to non-chiral theories in lower dimensions.

The next logical step in the development avoids this problem by making the Majorana-Weyl fermions participate in the process. To do so, they have to be on equal footing with the other fields, which can be achieved by bosonizing them. Actually we are not going to bosonize anything, but we will just start with 15 bosons (rather than with 10 bosons and 10 fermions) to cancel the conformal anomaly. The five bosons replacing the 10 fermions are included in the same torus, which now has dimension $22+11$. There is a small technical complication however: to get a lattice description of the theory one has to bosonize also the ghosts of local supersymmetry, and enlarge the lattice to include the charge of the resulting boson.

Now one can check the conditions for modular invariance, and the result is again that the torus has to correspond to a self-dual lattice, only this time it is not even but odd. This is all that needs to be said about modular invariance in these theories. It can be shown that (within the usual assumptions) this self-duality guarantees not just one-loop, but also higher-loop modular invariance.

Before we can make all this more precise we have to explain the corner stones of constructions of this type, and we will do so in the next three chapters on lattices, torus compactification and conformal field theory. These chapters are more or less independent, and readers sufficiently familiar with any of these subjects may want to skip some of them. Chapter 6 discusses the Neveu-Schwarz-Ramond model and its bosonic form, which is the heart of our construction. In particular we have to explain in some detail the properties of its bosonized ghost system. It is unfortunately not possible to explain this without the sophisticated techniques of conformal field theory, and although we have tried to supply the necessary information in chapter 5, we are aware of the fact that it may not be adequate to readers unfamiliar with it. Fortunately one finally arrives at a formulation of the theory where most of this can be forgotten again, at least as long as one is only interested in model building and spectrum calculations. For this reason we have formulated chapters 8 and 9 in such a way that they are self-contained, so that even readers who didn't fully grasp the ghost system should be able to apply the rules of model building. Meanwhile chapter 7 contains the discussion of world-sheet supersymmetry promised above.

3. Lattices

In this section we collect most of the known facts about self-dual lattices. In the mathematics literature, most work on this subject appears in journals on Number Theory, which are unfortunately not always easily available in physics libraries. Some of the relevant papers are listed under Ref. [24-26], and readers interested in proofs not provided below should consult these papers. We also recommend the reviews by Goddard and Olive [27].

3.1 DEFINITIONS

A lattice Λ is defined as a set of points in a vector space V of dimension N , of the form

$$\Lambda = \left\{ \sum_{i=1}^N n_i \vec{e}_i \mid n_i \in \mathbb{Z} \right\}, \quad (3.1)$$

where $\vec{e}_i$ ($i = 1, \dots, N$) is a set of basis vectors of V . The vector spaces we will consider are R^N , with a euclidean inner product, and $R^{p,q}$, ($p + q = N$), with a “lorentzian” inner product. In terms of the components v^a of vectors $\vec{v} \in V$ we define these inner products as $\vec{v} \cdot \vec{w} = v^a w^b d_{ab}$, with $d_{ab} = \delta_{ab}$ for R^N , and $d_{ab} = \epsilon(a) \delta_{ab}$, $\epsilon(a) = 1$ for $1 \leq a \leq p$, $\epsilon(a) = -1$ otherwise, for $R^{p,q}$. In the following we will denote these inner products as $(+)^N$ and $(+)^p(-)^q$ respectively, and we will refer to d_{ab} as the *signature*.

The *unit cell* of a lattice is the set of points $\{\vec{x} = \sum_i x_i e_i | 0 \leq x_i < 1\}$. It contains precisely one lattice point (i.e $\vec{x} = 0$), and every vector of V has precisely one “image” in the unit cell, obtained by adding a lattice vector to it. The *volume* $vol(\Lambda)$ of the unit cell of Λ is obtained by defining a matrix

$$g_{ij} = \vec{e}_i \cdot \vec{e}_j, \quad (3.2)$$

in terms of which

$$vol(\Lambda) = \sqrt{|\det g|}. \quad (3.3)$$

The matrix g_{ij} is called the *metric* of the lattice. It contains all information regarding the angles of the basis vectors and their lengths. Notice that if we change the signature from lorentzian to euclidean (by changing $(+)^p(-)^q$ to $(+)^{p+q}$) then $\det g$ changes at most by a sign, so that the volume of the unit cell remains the same.

The *dual* Λ^* of Λ is defined as

$$\Lambda^* = \{\vec{w} \in V | \vec{w} \cdot \vec{v} \in Z, \forall \vec{v} \in \Lambda\}. \quad (3.4)$$

A basis for Λ^* is the set of vectors $\vec{e}_i^*$, with $\vec{e}_i^* \cdot \vec{e}_j = \delta_{ij}$. It is easy to see that this is indeed a basis for Λ^* , i.e.

$$\Lambda^* = \left\{ \sum_{i=1}^N m_i \vec{e}_i^* | m_i \in Z \right\}. \quad (3.5)$$

To calculate the volume of Λ^* we define

$$g_{ij}^* = \vec{e}_i^* \cdot \vec{e}_j^*, \quad (3.6)$$

which is the metric of the dual lattice. Using the definition of the basis vectors one can easily show that $\sum_i e_i^{*a} e_i^b = d^{ab}$, and that therefore g^* is the inverse of g . Thus we have

$$vol(\Lambda) = (vol(\Lambda^*))^{-1}. \quad (3.7)$$

A lattice Λ is called

- *unimodular* if $vol(\Lambda) = 1$
- *integral* if $\vec{v} \cdot \vec{w} \in Z, \forall \vec{v}, \vec{w} \in \Lambda$
- *even* if Λ is integral, and $\vec{v}^2 = 0 \bmod 2, \forall \vec{v} \in \Lambda$
- *odd* if Λ is integral, but not even
- *self-dual* if $\Lambda = \Lambda^*$

Obviously Λ is integral if and only if $\Lambda \subset \Lambda^*$.

3.2 COSETS

If Λ_s is a sublattice of Λ , then we can decompose Λ in cosets with respect to Λ_s . To do so, choose a set of coset representatives $\vec{\beta}_\alpha$, so that

$$\begin{aligned}\vec{\beta}_\alpha &\in \Lambda, \vec{\beta}_\alpha \notin \Lambda_s \\ \vec{\beta}_\alpha - \vec{\beta}_{\alpha'} &\notin \Lambda_s \text{ if } \alpha \neq \alpha'.\end{aligned}\tag{3.8}$$

If we extend this set with the vector $\vec{\beta} = 0$ to represent Λ_s itself, we get

$$\Lambda = \sum_{\alpha} (\vec{\beta}_\alpha + \Lambda_s),\tag{3.9}$$

i.e. every vector in Λ can be written as $\vec{\beta}_\alpha + \vec{v}_s, \vec{v}_s \in \Lambda_s$. Every coset $\vec{\beta}_\alpha + \Lambda_s$ has precisely one point in the unit cell of Λ_s . Therefore, if there are N_s cosets (including Λ_s), we find

$$vol(\Lambda) = \frac{1}{N_s} vol(\Lambda_s).\tag{3.10}$$

This is a fairly trivial, but very useful observation. A direct consequence is that a lattice is self-dual if and only if it is both integral and unimodular.

3.3 LIE ALGEBRA LATTICES

It is often convenient to express a self-dual lattice in terms of Lie algebra lattices. In the following when we speak of a Lie algebra lattice we mean any sublattice of the weight lattice of a semi-simple Lie algebra (for a discussion of Lie algebras, roots and weights we refer to [28]). In particular we need the *simply laced* lattices, belonging to semi-simple Lie algebras which are products of D_n, A_n and E_6, E_7, E_8 . They have the property that all their roots have the same length. If one normalizes them so that they have all norm 2, the weight lattice Λ_W is the dual of the root lattice Λ_R , which is itself a sublattice of the weight lattice (the norm of a vector is its squared length)

$$\Lambda_W = \Lambda_R^* ; \Lambda_W \supset \Lambda_R\tag{3.11}$$

Therefore we can decompose Λ_W into cosets with respect to Λ_R . In the case of Lie algebra lattices these cosets are called *conjugacy classes*. In general cosets form a group under addition, with the sublattice Λ_s as the unit element. In this case this group is called the *center* of the Lie algebra.

We will make extensive use of D_n lattices, for which we denote the conjugacy classes as follows

$$\begin{aligned}(0) &= \Lambda_R = \{(k_1, \dots, k_n), k_i \in \mathbb{Z}, \sum_{i=1}^n k_i = 0 \bmod 2\} \\ (v) &= \Lambda_R + (1, 0, 0, \dots, 0) \\ (s) &= \Lambda_R + (\frac{1}{2}, \frac{1}{2}, \dots, \frac{1}{2}) \\ (c) &= \Lambda_R + (-\frac{1}{2}, \frac{1}{2}, \dots, \frac{1}{2})\end{aligned}\tag{3.12}$$

One may check that the center is isomorphic to Z_4 if n is odd, and to $Z_2 \times Z_2$ if n is even.

The irreducible representations of the semi-simple Lie algebras are characterized by their weight vectors, which for a given representation all belong to the same conjugacy class of the lattice. Hence we can specify conjugacy classes also by giving representations that belong to them. In the case of D_n , (0) contains all even rank tensor products of the vector representation (which includes the singlet and the adjoint), (v) contains all odd-rank tensors, and (s) and (c) contain spinors of opposite chirality.

A useful property of conjugacy classes is that their mutual inner products modulo integers depend only on the class, and not on the individual vectors. If $\vec{\beta}_1, \vec{\beta}_2 \in \Lambda_W$, and $\vec{\lambda}_1, \vec{\lambda}_2 \in \Lambda_R$, we have

$$(\vec{\beta}_1 + \vec{\lambda}_1) \cdot (\vec{\beta}_2 + \vec{\lambda}_2) = \vec{\beta}_1 \cdot \vec{\beta}_2 \bmod 1,$$

because $\vec{\beta}_i \cdot \vec{\lambda}_j \in Z$, and because the root lattice of simply laced algebras is integral (this follows from the fact that it is contained in its own dual, see (3.11)). In fact Λ_R is an even lattice, because its basis vectors are roots, which have norm 2. Using this property one can show by a similar argument that the norms of two vectors in the same conjugacy class are always equal modulo 2. These properties are clearly of great help in constructing even integral lattices, because one only needs to check the lengths and inner products of conjugacy class representatives. For this reason we have collected some information concerning the conjugacy classes of the (simple) simply laced algebras in the following table.

Lie algebra	conjugacy class	smallest representation	weight norm (mod 2)
D_n	(0)	singlet	0
	(v)	vector	1
	(s)	spinor	$\frac{n}{4}$
	(c)	conjugate spinor	$\frac{n}{4}$
A_n	(p)	rank p tensor ($0 \leq p < n$)	$\frac{p(n+1-p)}{n+1}$
E_6	(0)	singlet	0
	(1)	(27)	$\frac{4}{3}$
	(2)	($\overline{27}$)	$\frac{4}{3}$
E_7	(0)	singlet	0
	(1)	(56)	$\frac{3}{2}$
E_8	(0)	singlet	0

The volume of the unit cells of Λ_R and Λ_W are easy to obtain. Since they are each

other's dual

$$\text{vol}(\Lambda_R) = \text{vol}(\Lambda_W)^{-1}$$

If we denote the order of the center (i.e. the number of conjugacy classes) by N_c , we learn from (3.10) that

$$\text{vol}(\Lambda_W) = \frac{1}{N_c} \text{vol}(\Lambda_R),$$

so that

$$\begin{aligned} \text{vol}(\Lambda_R) &= \sqrt{N_c} \\ \text{vol}(\Lambda_W) &= \frac{1}{\sqrt{N_c}} \end{aligned}$$

All these results are valid for simple as well as semi-simple Lie algebras. The order of the center of a semi-simple Lie algebra is simply the product of the order of the centers of its simple factors. The same is true for the volume of the unit cell.

To specify a Lie algebra lattice completely, we have to supply two pieces of information: the semi-simple Lie algebra, and a list of the conjugacy classes that appear. For a given semi-simple Lie algebra there are in general many possible choices of conjugacy classes. This choice is restricted only by the requirement that the list must close under addition, which in particular means that the root lattice must always be present. The possible sets of conjugacy classes are thus in one-to-one correspondence with the subgroups of the center. Instead of giving a full list of conjugacy classes, one can give a shorter list of generators, which is a subset of conjugacy classes from which all others can be obtained by addition.

Since the volume of the unit cell of the root lattice is $\sqrt{N_c}$, it follows immediately that a Lie algebra lattice is unimodular if and only if it contains precisely $\sqrt{N_c}$ conjugacy classes. This is only possible if we choose a Lie algebra for which N_c is a perfect square. The lattice will be integral if and only if all conjugacy class generators have integral norm and integral mutual inner product. It will be even if and only if they have even norm and integral inner products.

Using the table one can now easily construct many examples of euclidean self-dual lattices. The reader is invited to check that

- D_n with conjugacy classes (0) and (v) is odd and self-dual (this lattice is also known as Z^n).
- D_n with (0) and (s) is odd self-dual for $n = 4 \bmod 8$. and even self-dual for $n = 0 \bmod 8$
- A_{n^2-1} with (0) and (n) is odd self-dual for n even and even self-dual for n odd.

In general we call vectors of norm 2 on euclidean lattices *roots*. The roots include in any case those of the Lie algebras explicitly used in the construction of the lattice (i.e. those in the conjugacy class (0)), but in some cases there may be additional norm 2 vectors in the other conjugacy classes. In that case we can always write the lattice in a different way, using larger Lie algebras and fewer conjugacy classes, although this is not always convenient. For example, the lattices D_8 with (0) + (s) and A_8 with (0) + (3)

defined above are both equal (up to rotations) to E_8 , which has only one conjugacy class, namely (0) (and which is therefore automatically self-dual!).

A large class of even lorentzian self-dual lattices in $R^{p,p}$ can be constructed as follows. Take a semi-simple Lie algebra of rank p . Consider the set of all vectors in $R^{p,p}$ of the form $(\vec{v}_1, \vec{v}_2)$, so that $\vec{v}_1$ and $\vec{v}_2$ belong to the same conjugacy class of the Lie algebra. With respect to the lorentzian metric any such vector has integral inner product with any other such vector, any furthermore they all have even length. It is also easy to see that the resulting lorentzian lattice is unimodular, so that it is a self-dual lattice. Lattices of this type were first described by Englert and Neveu [29].

To conclude this discussion of Lie algebra lattices, we want to introduce a minor, but useful extension of this set, namely the lattice D_1 . Although it is not part of Cartan's classification of simple Lie algebra's, it is easy enough to define by simply extending the definition of D_n to $n = 1$. The resulting one-dimensional root lattice consists simply of the even integers. The shortest non-trivial vector has norm four, so there are no roots on this root lattice. The vector conjugacy class consists of the odd integers, and the two spinor conjugacy classes are obtained by shifting the root lattice by $+\frac{1}{2}$ and $-\frac{1}{2}$.

3.4 LATTICE ENGINEERING

There are several tricks for making new self-dual lattices out of already known ones. These tricks are all variations on the same basic theme: one adds new conjugacy classes and removes old ones, in such a way that their total number stays the same (i.e. the lattice remains unimodular), and that all inner products remain integral.

A very general method is the following. Let Λ be a self-dual lattice. Choose a vector $\vec{s}$ so that $k\vec{s} \notin \Lambda, 1 \leq k \leq p-1$, but $p\vec{s} \in \Lambda$. For convenience we will assume that p is prime for the moment. Now we can define a lattice Λ_0 as the sublattice of Λ which has integer inner product with $\vec{s}$. Since $\vec{s} \notin \Lambda$ there must be a vector $\vec{t} \in \Lambda$ such that $\vec{s} \cdot \vec{t}$ is not integer. But then $p\vec{t}$ belongs to Λ_0 , and is the smallest multiple of $\vec{t}$ in Λ_0 . Hence we can decompose Λ in cosets

$$\Lambda = \sum_{k=0}^{p-1} (k\vec{t} + \Lambda_0)$$

Consider now the new lattice

$$\Lambda' = \sum_{k=0}^{p-1} (k\vec{s} + \Lambda_0)$$

clearly Λ' is unimodular. If furthermore $\vec{s}$ has integral norm, it is self-dual. If Λ is even and $\vec{s}$ has even length, Λ' is even self-dual.

Another useful trick is the following. Consider a Lie algebra lattice containing, among others, a D_n factor. Now replace this D_n factor by D_m , where $m = n + 8l, l \in \mathbb{Z}$. Since the unit cells of D_n and D_m have the same volume for any n and m , the new lattice is unimodular. Furthermore one can easily check that if $n = m \bmod 8$ all inner products and norms of the four conjugacy classes remain unchanged modulo two. Hence the new lattice will be (even) self-dual if the original lattice was (even) self-dual.

This transformation is of course meaningless if it results in $m = 0$. But one can give a meaning to it if m becomes negative. Consider for example a euclidean even self-dual lattice $\Lambda_p \times D_n$ with inner product with signature $(+)^{p+n}$. For $m < 0$ (with m defined as above) we can consider the lorentzian lattice $\Lambda_p \times D_m$ with signature $(+)^p(-)^{-m}$. This lattice is unimodular for the reasons explained above. Furthermore the lorentzian inner products of the new lattice are equal to the euclidean ones of the original one modulo two. Thus we see that shifting n by multiples of eight to negative values has the effect of changing a euclidean self-dual lattice to a lorentzian one.

3.5 LATTICE CLASSIFICATION THEOREMS

A lot is known about the possibilities for constructing self-dual lattices. One should keep in mind here that euclidean lattices can only be classified up to rotations, because any rotation respects the self-duality conditions. Likewise, lorentzian lattices can only be classified modulo Lorentz transformations. The known results are as follows:

1. *Lorentzian, odd.* All such lattices are Lorentz transformations of the lattice $Z^{p,q}$, which is the set of points in $R^{p,q}$ defined as $\{(n_1, \dots, n_p, m_1, \dots, m_q), n_i, m_j \in \mathbb{Z}\}$.

2. *Lorentzian, even.* Lattices of this type exist only for $|p - q| = 0 \bmod 8$. For $p = q + 8n$ they are Lorentz transformations of the lattice $nE_8 \times qP^{1,1}$. The E_8 lattice is most easily described as D_8 with conjugacy classes (0) and (s) , while $P^{1,1}$ is a lorentzian lattice of signature $(+)(-)$, which may be described as $D_1 \times D_1$ with conjugacy classes $(0, 0) + (v, v) + (s, s) + (c, c)$ (this is a lattice of the Englert-Neveu type, discussed above).

3. *Euclidean, odd.* A class of such lattices which exists in any dimension is $\mathbb{Z}^n$. In general, if any odd self-dual lattice has vectors of length one, they can only have mutual inner products $\pm 1, 0$. Thus they form an orthonormal basis for a subspace of R^n , which must be part of the basis of the lattice. Any other basis vector of the lattice must be orthogonal to this subspace. Thus if there are $2p$ length one vectors, the lattice is of the form $\mathbb{Z}^p \otimes \Lambda^{n-p}$, where the second factor does not contain any vectors of length one.

An odd euclidean self-dual lattice Λ can be related to an even self-dual one of higher dimension in the following way [24]. Consider the sublattice Δ_0 of even norm vectors. Obviously Δ_0 has only one non-trivial coset in Λ , because the sum of two odd norm vectors is even. Thus

$$\Lambda = \Delta_0 + \Delta_v,$$

where Δ_v denotes the odd-norm vectors. Since $\text{vol}(\Delta_0)/\text{vol}(\Delta_0^*) = 4$, the dual of Δ_0 can be decomposed into four cosets,

$$\Delta_0^* = \Delta_0 + \Delta_v + \Delta_s + \Delta_c$$

These four cosets generate an abelian group of order four, which can only be $\mathbb{Z}_4$ or $\mathbb{Z}_2 \times \mathbb{Z}_2$. This allows us to determine the mutual inner products of the four cosets and their norms modulo 2. If the group is $\mathbb{Z}_4$, one finds $\Delta_c = -\Delta_s$ and $2\Delta_s \subset \Delta_v$, so that the norm of all vectors in Δ_s and Δ_c is equal to $\frac{k}{4} \bmod 2, k$ odd. In the latter case $2\Delta_s = 2\Delta_c \subset \Delta_0$, so that the norm of all vectors in Δ_s and Δ_c is half-integer. One can then use $\Delta_s + \Delta_v = \Delta_c$, plus the fact that vectors in Δ_s cannot have integer inner

products with those in Δ_v , to show that the norm of vectors in Δ_s and Δ_c must be equal modulo 2. Thus for both possible abelian groups the vectors in Δ_s and Δ_c have, modulo two, norms equal to each other and to $\frac{k}{4}$, $0 \leq k < 7$. Consider then the lattice obtained by adding a D_{8-k} factor, so that the entire lattice consists of the vectors

$$(\Delta_0, 0) + (\Delta_v, v) + (\Delta_s, s) + (\Delta_c, c)$$

This lattice is unimodular, integral and even, so that it is even self-dual. So all odd self-dual lattice can be obtained from the even self-dual ones by inverting this procedure to remove D_n factors. This leaves us with the classification problem of

4. *Euclidean, even lattices.* These exist only in dimensions which are a multiple of 8, and are known explicitly for 8, 16 and 24 dimensions. In 8 dimensions, the only possibility is E_8 ; in 16 dimensions one can have $E_8 \otimes E_8$ and D_{16} with conjugacy classes (0) and (s). These two lattices cannot be rotated into each other. In 24 dimensions there are 24 even self-dual lattices, which are usually called the *Niemeyer lattices* [25], after the German mathematician who first classified them. A list of their conjugacy classes can be found in the first paper of Ref. [24].

One of the 24 lattices is the Leech-lattice [30], which differs from the other 23 because it contains no vectors of norm 2. Nevertheless, we can in a certain sense regard it as a Lie algebra lattice, which is a product of 24 D_1 factors, on which one has 2^{24} conjugacy classes. A list of these classes (or rather 12 classes that generate them) will appear in [31].

There is a rather amusing way of estimating the number of lattices in $8n$ dimensions. Every lattice has a discrete automorphism group. This is a discrete subgroup of $O(8n)$ which takes the lattice into itself. It consists of at least two elements, the identity and the reflection through the origin. Denote the total number of elements of this group for a lattice Λ as $g(\Lambda)$. Then the following identity can be proved [26], [32].

$$\sum \frac{1}{g(\Lambda)} = \frac{B_{4n}}{8n} \prod_{j=1}^{4n-1} \frac{B_{2j}}{4j},$$

where B_{2j} are the Bernoulli numbers. This formula is known as the Minkowski-Siegel mass formula. The sum is over all lattices in $8n$ dimensions, and adds up to a number that is less than half the total number of lattices in this dimension. Thus the right hand side, multiplied by 2, is a lower limit of the number of lattices. Calculating it for the lowest dimensions, one finds

dimension	lower limit	actual number
8	2.8×10^{-9}	1
16	4.9×10^{-18}	2
24	15.8×10^{-15}	24
32	8.0×10^7	?

This explains why the lattices of dimension 32 have not yet been classified. The explosive growth of the number of lattices that one observes here is related to the reason why the number of chiral four dimensional string theories is so much larger than the number of ten-dimensional ones, as we will see.

4. Torus Compactification of the Bosonic String

Armed with all relevant facts about lattices, we can now attack the problem of the bosonic string, propagating through a 26 dimensional space-time of which some dimensions have been "compactified". Of course our ultimate interest is in fermionic strings, but since we will use a bosonic approach to fermionic strings, a thorough understanding of the compactified bosonic string is essential. The presentation given here is essentially the one of [4], [33], although these authors compactified the heterotic string rather than the bosonic string. To warm up, and to introduce some notations, we will begin with a very brief reminder of the basics, the quantization of the uncompactified string.

4.1 THE UNCOMPACTIFIED CLOSED BOSONIC STRING

The propagation of a string through d dimensional space-time is governed by the action

$$S = -\frac{1}{4\pi\alpha'} \int d\sigma_0 \int_0^\pi d\sigma_1 \partial_\alpha X^\mu \partial^\alpha X^\nu \eta_{\mu\nu}, \quad (4.1)$$

where σ_0 and σ_1 are the world-sheet coordinates, indicating respectively the time direction and the direction along the string (one often denotes these variables as τ and σ , but we won't do this to prevent confusion with the usual notation for the one-loop modular parameter). In (4.1) $\eta_{\mu\nu}$ is a flat metric and $X^\mu(\sigma_0, \sigma_1)$ is a map from the world-sheet to space-time, which specifies the position of every point of the string at any time. The parameter α' (the "Regge slope parameter") is usually set to the value $\frac{1}{2}$, but we will refrain from that for a while. We shall only consider closed strings, for which one has the boundary condition

$$X^\mu(\sigma_0, 0) = X^\mu(\sigma_0, \pi). \quad (4.2)$$

From the action one can in the usual way derive the operators H and P which generate translations in the σ_0 and σ_1 directions on the world-sheet:

$$H = \frac{1}{4\pi\alpha'} \int_0^\pi d\sigma_1 [(\partial_0 X^\mu)(\partial_0 X_\mu) + (\partial_1 X^\mu)(\partial_1 X_\mu)] \quad (4.3)$$

$$P = \frac{1}{2\pi\alpha'} \int_0^\pi d\sigma_1 (\partial_0 X^\mu \partial_1 X_\mu) \quad (4.4)$$

The quantization of this system is simplest in light-cone gauge, in which the dynamical variables are $X^i, i = 1, \dots, d-2$. Their canonical momenta Π_i are easily obtained from the action

$$\Pi_i = \frac{1}{2\pi\alpha'} \partial_0 X_i.$$

The solutions of the equations of motion have the form

$$X^i(\sigma_0, \sigma_1) = q^i + 2\alpha' p^i \sigma_0 + \frac{i}{2} \sqrt{2\alpha'} \sum_{n \neq 0} \frac{1}{n} (\alpha_n^i e^{-2in(\sigma_0 + \sigma_1)} + \tilde{\alpha}_n^i e^{-2in(\sigma_0 - \sigma_1)}). \quad (4.5)$$

Using the canonical commutator

$$[X^i(\sigma_0, \sigma_1), \Pi^j(\sigma_0, \sigma'_1)] = i\delta^{ij}\delta(\sigma_1 - \sigma'_1) \quad (4.6)$$

one finds the commutation relations for the quantized modes

$$\begin{aligned} [q^i, p^j] &= i\delta^{ij} \\ [\alpha_m^i, \alpha_n^i] &= m\delta^{ij}\delta_{n+m,0} \\ [\tilde{\alpha}_m^i, \tilde{\alpha}_n^i] &= m\delta^{ij}\delta_{n+m,0} \\ [\alpha_m^i, \tilde{\alpha}_n^i] &= 0 \end{aligned} \quad (4.7)$$

The spectrum is obtained by requiring that both H and P vanish. This leads to the conditions

$$2p_+p_- - p_i^2 \quad (= p^2 = m^2) = \frac{2}{\alpha'}(\mathcal{N} + \tilde{\mathcal{N}} - 2), \quad (4.8)$$

where the -2 is due to the zero mode fluctuations of the oscillators, and we have defined the number operators

$$\mathcal{N} = \sum_{n=1}^{\infty} \alpha_{-n}^i \alpha_n^i$$

and similarly for $\tilde{\mathcal{N}}$. From the vanishing of P we get

$$\mathcal{N} = \tilde{\mathcal{N}}.$$

In the quantized theory this is imposed as a condition on the physical spectrum.

It is convenient to define an operator H' , which is equal to H , but without the zero-mode terms (i.e. the operators p^i and q^i). It is also useful to define the linear combinations

$$\begin{aligned} H_L &= \frac{1}{2}(H' - P) = 2(\tilde{\mathcal{N}} - 1) \\ H_R &= \frac{1}{2}(H' + P) = 2(\mathcal{N} - 1) \end{aligned}$$

which are the generators of world-sheet translations in the $\sigma_0 - \sigma_1$ and $\sigma_0 + \sigma_1$ directions respectively. Their eigenstates are called left- and right-moving modes respectively. Denoting the eigenvalues of H'_L and H'_R on a state $|\psi\rangle$ as $\frac{1}{2}\alpha'm_L^2$ and $\frac{1}{2}\alpha'm_R^2$ we can write the result obtained above as^{*}

$$\begin{aligned} H_L |\psi\rangle &= \frac{1}{2}\alpha'm_L^2 |\psi\rangle \\ H_R |\psi\rangle &= \frac{1}{2}\alpha'm_R^2 |\psi\rangle \\ m^2 &= \frac{1}{2}(m_L^2 + m_R^2) \end{aligned}$$

with the condition

$$m_L^2 = m_R^2.$$

^{*} For comparison with [6] note that we have reversed the assignment of $\tilde{\alpha}$ and $\tilde{\tau}$ here from right to left-movers here. This is more convenient in heterotic strings, where we will choose the NSR system to be right-moving, just as in [6].

4.2 THE COMPACTIFIED CLOSED BOSONIC STRING

Now we turn to the case of a torus compactified space-time. A torus in N dimensions is simply R^N modulo a lattice Λ . This means that points in R^N which differ by a vector on Λ are identified. If a string propagates in a space-time with this topology it is not necessary anymore that it closes into the same space-time point, as in (4.2). We should allow for the possibility that the points $\sigma_1 = 0$ and $\sigma_1 = \pi$ of the string are identified points in R^N :

$$X^i(\pi) = X^i(0) + 2\pi L^i,$$

where $L^i \in \Lambda$ and the factor 2π is put in for convenience. Physically, this corresponds to a string winding around the torus a certain number of times. To accommodate this possibility one modifies the string field X^i in the following way

$$X^i(\sigma_0, \sigma_1) = q^i + 2\alpha' p^i \sigma_0 + 2L^i \sigma_1 + \frac{i}{2} \sqrt{2\alpha'} \sum_{n \neq 0} \frac{1}{n} (\alpha_n^i e^{-2in(\sigma_0 + \sigma_1)} + \tilde{\alpha}_n^i e^{-2in(\sigma_0 - \sigma_1)}), \quad (4.9)$$

where L^i is an operator whose eigenvalues are the lattice vectors.

Having done this, we go now through the same exercise as for the uncompactified string. It is easy to verify that nothing changes in the canonical commutators (4.7) except for the trivial addition that L^i should commute with all other operators. The mass formula replacing (4.8) acquires on the right hand side an extra term $\frac{1}{\alpha'^2} L^2$. Furthermore, to interpret the left hand side as a mass in $d - N$ dimensions we have to move the internal components of p^i to the right, to get

$$p^2 = m^2 = \sum_{I=1}^N p_I^2 + \frac{1}{\alpha'^2} L^2 + \frac{2}{\alpha'} (\mathcal{N} + \tilde{\mathcal{N}} - 2).$$

Here we have chosen the first N components as the compactified ones. The momenta $p_I, I = 1, \dots, N$ are canonical momenta of a periodic coordinate, and therefore (for reasons very familiar from solid state physics) they are quantized. Indeed, if we write their eigenfunctions of momentum k_I as $e^{ik_I x_I}$, then we see that these functions are periodic on the boundaries of the unit cell of Λ if and only if $\vec{k} \cdot \vec{l} \in \mathbb{Z}$ for every $\vec{l} \in \Lambda$. This means that the eigenvalues of k_I must lie on the dual lattice Λ^* of Λ .

The condition obtained from the vanishing of P reads now

$$\mathcal{N} - \tilde{\mathcal{N}} + p_I L_I = 0$$

It is again useful to define left and right hamiltonians H' exactly as before, but with the understanding that only the space-time momenta are removed from H to get H' , not the internal ones. Then we get

$$\begin{aligned} \frac{1}{4} \alpha' m_R^2 &= \frac{1}{2} p_R^2 + \mathcal{N} - 1 \\ \frac{1}{4} \alpha' m_L^2 &= \frac{1}{2} p_L^2 + \tilde{\mathcal{N}} - 1, \end{aligned} \quad (4.10)$$

where as before left and right masses must be equal to each other and to m^2 . The left

and right momenta are defined as

$$\begin{aligned} p_R^I &= \frac{1}{\sqrt{2}}(\sqrt{\alpha'} p^I + \frac{1}{\sqrt{\alpha'}} L^I) \\ p_L^I &= \frac{1}{\sqrt{2}}(\sqrt{\alpha'} p^I - \frac{1}{\sqrt{\alpha'}} L^I) \end{aligned}$$

Before discussing the meaning of these left and right momenta, we want to mention a simple generalization, obtained by adding a background antisymmetric tensor field. This background field is coupled to X^μ via an additional term in the action

$$S_B = \frac{1}{2\pi\alpha'} \int d\sigma_0 \int_0^\pi d\sigma_1 \epsilon^{\alpha\beta} B_{\mu\nu} \partial_\alpha X^\mu \partial_\beta X^\nu$$

We take $B_{\mu\nu}$ to be a constant background field, which has non-vanishing components only in the compactified dimensions. The extra term is then a total derivative, which does not affect H or P . So how can it make any difference? It can, because there is a change in the canonical momentum operator, which now becomes

$$\begin{aligned} \Pi_\mu &= \frac{1}{2\pi\alpha'} (\partial_0 + B_{\mu\nu} \partial_1) X^\nu \\ &= \frac{1}{\pi} (p_\mu + \frac{1}{\alpha'} B_{\mu\nu} L^\nu + \text{oscillators}) \end{aligned}$$

For the oscillator contributions, everything remains the same. Although $\partial_1 X^\nu$ contains oscillator terms, there is a cancellation between left- and right-mover contributions to $[\Pi, X]$, so that the canonical commutators of the oscillators remain unchanged. The difference occurs for the zero modes, where p_μ is replaced as the generator of translations by $\tilde{p}_\mu = p_\mu + \frac{1}{\alpha'} B_{\mu\nu} L^\nu$. So it is $\tilde{p}$ rather than p which has its eigenvalues quantized on the dual of Λ . Although the mass formula remains the same as a function of p , it makes now more sense to express it in terms of $\tilde{p}$. The result is the same as (4.10) above, but with

$$\begin{aligned} p_R^I &= \frac{1}{\sqrt{2}}(\sqrt{\alpha'} \tilde{p}^I + \frac{1}{\sqrt{\alpha'}} (L^I - B^{IJ} L_J)) \\ p_L^I &= \frac{1}{\sqrt{2}}(\sqrt{\alpha'} \tilde{p}^I - \frac{1}{\sqrt{\alpha'}} (L^I + B^{IJ} L_J)) \end{aligned} \tag{4.11}$$

where $\tilde{p} \in \Lambda^*$ and $L \in \Lambda$. In the following we will omit the tilde on p . It is clear that the dependence on α' is not very significant. We can absorb it by changing the length of all vectors on Λ by $\sqrt{\alpha'}$ (and hence those on Λ^* by $\frac{1}{\sqrt{\alpha'}}$). This amounts to a rescaling of the size of the torus. From now on we will use this freedom to set $\alpha' = \frac{1}{2}$.

To interpret the result it is convenient to express the left and right momenta in terms of the basis vectors e_a^I of the lattice:

$$\begin{aligned} L^I &= e_a^I n_a \\ p^I &= e_a^I m_a = e_a^I g_{ab}^* m_b \end{aligned}$$

Then we get

$$\begin{aligned} p_R^I &= e_a^I \left(\frac{1}{2} g_{ab}^* m_b - g_{ab}^* B_{bc} n_c + n_a \right) \\ p_L^I &= e_a^I \left(\frac{1}{2} g_{ab}^* m_b + g_{ab}^* B_{bc} n_c - n_a \right) \end{aligned}$$

(upper and lower indices have no special significance here.) One can regard g (or its inverse g^*) as a constant background field, just as B . This becomes evident if one performs this change of basis directly in the action (4.1) : the effect is to change the flat metric δ_{IJ} (in the compactified dimensions) to g_{IJ} .

It is now instructive to calculate the inner product of the $2N$ dimensional vectors^{*} $P = (p_L; p_R)$ and $P' = (p'_L; p'_R)$ (the latter is expressed in terms of n' and m') with respect to the *lorentzian* signature $[(+)^N; (-)^N]$. The result is

$$\vec{P} \cdot \vec{P}' = nm' + mn'$$

From this expression we can learn many things. First of all, the lattice $\Gamma_{N;N}$ consisting of the vectors P defined above is an integral and even *lorentzian* lattice. Secondly, the dot-products do not depend on the background fields g and B . Therefore they must be parameters of a Lorentz transformation. Finally, using this background independence, we can make the choice $e_a^I = \delta_a^I$, $g_{IJ} = \delta_{IJ}$, $B_{IJ} = 0$. In this special case the lattice is $(D_1 \times D_1)^N$ with conjugacy classes $(0, 0) + (v, v) + (s, c) + (c, s)$ in each factor, which is self-dual. Because Lorentz transformations preserve self-duality, we conclude that $\Gamma_{N;N}$ is self-dual for any value of the background fields.

Thus compactification of the bosonic string on a torus R^N/Λ , with an additional background field B_{IJ} , can be described in terms of a new lattice $\Gamma_{N;N}$ which for any Λ and B is *lorentzian* self-dual. It should be clear that the *lorentzian* signature is a mathematical device for characterizing the lattice; it has nothing to do with the metric of space-time and does not lead to ghosts.

The number of parameters of g and B is $\frac{1}{2}N(N+1) + \frac{1}{2}N(N-1) = N^2$. One can verify that they are precisely the N^2 parameters of the coset $\frac{SO(N,N)}{SO(N) \times SO(N)}$, which is the set of Lorentz transformations of the lattice modulo the rotations of the left and right sector. These rotations are a symmetry of the theory, because all masses depend on p_L and p_R only via their norms. The genuine *lorentz* transformations mixing p_L and p_R on the other hand are not symmetries. Although the self-duality condition is invariant under such Lorentz transformations, the spectrum is not. So we have an N^2 parameter family of different string theories.

As seems usual in string theory, these results were originally discovered in the opposite order: First Narain [4] constructed the *lorentzian* lattices, and afterwards Narain, Sarmadi and Witten [33] gave the background field interpretation from which we started. We still have not discussed the relevance of the self-dual lattice which emerges from this construction. To do so, we have to consider the partition function of the compactified theory.

* We separate left- and right-moving sectors by a semicolon in the following.

4.3 PARTITION FUNCTIONS

Consider first the uncompactified bosonic string at one loop. The amplitude is an expression of the form

$$A_{1-loop} = \int \frac{\mathcal{D}X \mathcal{D}g}{V_{Diff} V_C} \exp(-S_E(g, X)), \quad (4.12)$$

where V_{Diff} and V_C denote the volume of the set of local diffeomorphism and of the conformal transformations respectively, and S_E is the bosonic string action (4.1), continued to euclidean space. What is meant by this expression is that one should integrate over all values of X and the world-sheet metric on the torus that are not equivalent by reparametrizations and conformal transformations. This is the analog of the process of gauge-fixing in gauge theories, and this problem can be tackled with very similar techniques. We will not (need to) explain this procedure here, but simply state the result (the reader interested in the details is referred to [34]):

$$A_{1-loop} = V \int \frac{d\tau}{(Im\tau)^2} (Im\tau)^{\frac{2-d}{2}} \int \mathcal{D}'X \exp[-S_E(g(\tau), X)] \quad (4.13)$$

The factor V denotes the volume of space-time, which of course is infinite. This infinity is easy to understand: since the action vanishes for constant X , one of the infinitely many integrals in $\mathcal{D}X$ is simply an integral of a constant over all of space-time. This infinite factor is nothing to worry about, and is simply to be regarded as a momentum conservation delta function with argument zero, which appears because we are considering a diagram without external lines. However, we have to make sure that we factor it out correctly, which can be achieved by putting the system in a box in order to regulate the infinity. After taking out the zero mode integral, we denote the remaining integral by $\mathcal{D}'X$. After gauge fixing, the components of the world-sheet metric are constants (i.e. they are independent of the world-sheet coordinates), which depend on the modular parameter τ . The remaining X integral is therefore a simple gaussian, and the result may be written as

$$\int \mathcal{D}'X e^{-S_E} = \frac{1}{\det'(-\Delta)}, \quad (4.14)$$

where Δ is the laplacian on the torus, $\Delta = g_{\alpha\beta} \partial_\alpha \partial_\beta$, and the prime on the determinant indicates that the zero-mode should be omitted. For every component of X^μ that is factored out one is left with a factor $\frac{1}{\sqrt{Im\tau}}$, which is nothing but the normalization of the zero-mode of the laplacian on the skewed torus (which has volume $Im\tau$). This does not quite account for all of the factors of $Im\tau$ in (4.13), but the extra one is a remnant of the gauge fixing procedure. The thing to remember is that every uncompactified X^i contributes a factor $(Im\tau)^{-\frac{1}{2}}$ to the integrand.

The factor (4.14) of (4.13) is called the (one-loop) *partition function*, and will be denoted as $\mathcal{F}(\tau)$. It is possible to evaluate it directly by regularization of the infinite product of eigenvalues of Δ (see the appendix of [34]), but at one loop there is a more intuitive way. Let us first recall a result for a one-dimensional bosonic system, specified

by a coordinate $X(t)$ and an action $S(X(t))$. One can prove (if the action satisfies certain conditions) the following relation between the path integral for a euclidean action S_E and the trace of the exponential of the hamiltonian H of the system

$$\int_{PBC} \mathcal{D}X e^{-S_E} = \text{Tr} e^{-\beta H}.$$

Here “PBC” means that $X(t)$ must satisfy periodic boundary conditions on the interval $[0, \beta]$. For a derivation of this relation see e.g. [35]. The result is easily generalized to X ’s having more components, or even an infinite number of components. The latter case is relevant for us, because the scalar field $X(\sigma_0, \sigma_1)$ appearing in string theory can be regarded as having an infinite number of components labeled by σ_1 (times a finite number of components because of the space-time index, which we omitted here). To use this relation, one should think of $Im\tau$ as β . There is a small complication if $Re\tau \neq 0$, because then the periodicity of X in $Im\tau$ depends on $Re\tau$. This can be taken into account by including a σ_1 shift operator in the trace, so that states at $\sigma_0 = 0$ are not identified with the same states at $\sigma_0 = Im\tau$, but with states of a string which is rotated in the σ_1 direction by an amount $Re\tau$. The generator of infinitesimal σ_1 translations is of course P , so that one gets

$$\begin{aligned} \mathcal{F}(\tau) &= \int_{T(\tau)} \mathcal{D}'X e^{-S_E} = \text{Tr} e^{\pi Im\tau H'} e^{i\pi Re\tau P} \\ &= \text{Tr} e^{i\pi\tau H_R} e^{-i\pi\bar{\tau} H_L} \end{aligned} \tag{4.15}$$

Here we have rescaled the variables σ_0 and σ_1 by factors of π . The integral is over all two-dimensional fields X that are periodic on a torus T specified by a modular parameter τ .

This way of writing the partition function admits a simple interpretation. Expanding it in terms of $q = e^{i\pi\tau}$ one gets a power series of the form $\mathcal{F}(\tau) = \sum d_{nm} q^n \bar{q}^m$. Obviously, d_{nm} is simply the number of states of left mass m and right mass n . (At this point we should add a word of warning. The states appearing in the spectrum are only those for which $n = m$. Notice however that these are *not* the only states contributing to the one-loop path integral. The reader who thinks that states with $n \neq m$ are projected out by the $Re\tau$ integral should realize that the lower boundary of the standard modular domain is not a straight line.)

It is usually a simple combinatorial exercise to read off the partition function from the expressions for H_L and H_R . Every term in the hamiltonian yields a factor in the partition function, so that one can derive the partition function term-by-term. Take for example a single bosonic oscillator α_n . It contributes to the right Hamiltonian via a term $2\alpha_{-n}\alpha_n$, and when it acts it increases the mass of a state by $2n$. Thus it contributes to the partition function a factor

$$(1 + q^{2n} + q^{4n} + q^{6n} + \dots) = \frac{1}{(1 - q^{2n})},$$

where on the right the term q^{2kn} corresponds to the contribution of $(\alpha_{-n})^k$. The partition function of the bosonic string contains such a factor for every oscillator.

For the uncompactified string we leave out the zero modes in H_L and H_R as discussed before. The remaining trace, i.e. the contribution from the bosonic oscillators to the partition function, is equal to

$$\mathcal{F}_B(\tau) = \text{Tr} q^{H'_R} \bar{q}^{H'_L} = |\eta(\tau)|^{-48} \quad (4.16)$$

where η is the famous Dedekind η -function

$$\eta(\tau) = q^{\frac{1}{12}} \prod_{n=1}^{\infty} (1 - q^{2n})$$

(as demonstrated in these equations, we will freely mix τ 's and q 's in the following.)

Since the oscillator contribution to the mass formula is the same for compactified and uncompactified coordinates, the partition function for the compactified string contains the same factor. The only difference is that the zero modes of the compactified coordinates are now included in the determinant as well as the hamiltonian. So first of all we have to multiply the uncompactified string result by $\sqrt{Im\tau}$ for every compactified coordinate to remove the zero-mode normalizations in (4.12). Secondly, we have to take into account the contribution to the trace due to the extra momentum terms in the mass formula (4.10). Since this is simply an extra term in the mass formulas, it produces an extra factor in the partition function, which together with the $\sqrt{Im\tau}$ factors is called the *correction factor*, because it gives the correction in the bosonic partition function for the fact that part of the zero-modes are compactified. From the mass formula (4.10) we get for the for a lattice $\Gamma_{N;N}$

$$C(\Gamma_{N;N}) = \mathcal{L}_{N;N}(\tau) (Im\tau)^{N/2}, \quad (4.17)$$

where $\mathcal{L}_{N;N}$ is the *lattice partition function*:

$$\mathcal{L}_{N;N} = \sum_{(p_L; p_R)} e^{i\pi\tau p_R^2} e^{-i\pi\bar{\tau} p_L^2}. \quad (4.18)$$

Here the sum is over all the vectors on the lattice.

4.4 MODULAR INVARIANCE

We can now discuss the modular transformations of these functions. As explained in chapter 2, we have two transformations to check, namely $\tau \rightarrow \tau + 1$ and $\tau \rightarrow -\frac{1}{\tau}$. The first one leaves the partition function invariant if and only if it is an even function of q . This is clearly the case for both compactified and uncompactified strings, as a consequence of the fact that there are 24 transverse dimensions, and because $\Gamma_{N;N}$ is even. The second modular transformation is a bit harder to check, and requires a trick, which goes by the name of “Poisson resummation formula”.

Define instead of (4.18) the function

$$P(\tau, \vec{z}) = \sum_{(p_L; p_R)} e^{i\pi\tau(\vec{p}_R + \vec{z}_R)^2} e^{-i\pi\bar{\tau}(\vec{p}_L + \vec{z}_L)^2}, \quad (4.19)$$

where $\vec{z} = (\vec{z}_L, \vec{z}_R)$ is a $2N$ vector, which is not restricted to lie on the lattice. Since we sum over all vectors on the lattice, the function P is obviously invariant if we shift $\vec{z}$ by a lattice vector. In other words, it is periodic with the periodicity of the lattice, and hence we can Fourier transform it. Normally the Fourier analysis of lattice functions would involve the dual lattice, but because our lattice is self-dual, we don't have to bother with that. The expression for P in terms of its Fourier transform $\tilde{P}$ is

$$P(\tau, \vec{z}) = \sum_{\vec{q}} e^{-2\pi i \vec{z} \cdot \vec{q}} \tilde{P}(\tau, \vec{q}) \quad (4.20)$$

The inverse transformation is of course

$$\tilde{P}(\tau, \vec{q}) = \int_{\text{unit cell}} d^{2N}z e^{-2\pi i \vec{z} \cdot \vec{q}} P(\tau, \vec{z}),$$

i.e. we have a multiple integral over all components of $\vec{z}$ with boundaries given by the unit cell of the lattice. The function P contains a sum over all lattice vectors. We can take the exponential multiplying it inside the sum, and furthermore we can shift the variables $\vec{z}$ in this exponential factor by $\vec{p}$. This shift introduces only a factor $\exp(2\pi i \vec{q} \cdot \vec{p})$, which is equal to one because both $\vec{q}$ and $\vec{p}$ are lattice vectors. The result is thus

$$\tilde{P}(\tau, \vec{q}) = \sum_{\vec{p}} \int_{\text{unit cell}} d^{2N}z e^{-2\pi i \vec{q} \cdot (\vec{z} + \vec{p})} e^{i\pi\tau(\vec{p}_R + \vec{z}_R)^2} e^{-i\pi\bar{\tau}(\vec{p}_L + \vec{z}_L)^2}$$

The sum combined with the integral gives as an integral over *all* unit cells of the lattice, i.e. over the entire space. Thus we get

$$\tilde{P}(\tau, \vec{q}) = \int_{R^{N;N}} d^{2N}x e^{-2\pi i \vec{q} \cdot \vec{x}} e^{i\pi\tau(x_R)^2} e^{-i\pi\bar{\tau}(x_L)^2},$$

where we have introduced a new integration variable $\vec{x}$ representing, in each cell, $\vec{p} + \vec{z}$. This integral is a simple gaussian, which yields

$$\tilde{P}(\tau, \vec{q}) = |\tau|^{-N} e^{-\frac{i\pi}{\tau}(q_R)^2} e^{\frac{i\pi}{\bar{\tau}}(q_L)^2}$$

Notice that we have made use here of the fact that the lattice is unimodular, so that the usual determinant factors that one expects from gaussian integrals are equal to one. Substituting this result into (4.20) and putting $\vec{z}$ equal to zero again we get the identity

$$P(-\frac{1}{\tau}) = |\tau|^N P(\tau) \quad (4.21)$$

One can assemble all these ingredients into the one-loop partition function of the compactified bosonic string. To check the modular invariance of the complete answer one

has to verify that the factors $|\tau|$ (which are called *modular weight factors*) cancel. They do, because there are also contribution from the transformation of $Im\tau$ and $\eta(\tau)$. The transformation of $Im\tau$ is straightforward: $Im(-\frac{1}{\tau}) = |\tau|^{-2}Im\tau$, which can be used to show that the correction factor (4.17) is invariant. To check the invariance of the uncompactified bosonic string, one needs furthermore

$$\eta(-\frac{1}{\tau}) = \sqrt{-i\tau} \eta(\tau).$$

The reader who checks this will notice that again $d = 26$ in (4.13) is essential.

It is useful to know that all of this extends in several ways. First of all there is an extension to lattices with different left and right dimensions, which appear in compactifications of heterotic strings, and also in the covariant lattice construction to be discussed in the following. The derivations are completely analogous, and the only difference is that in (4.21) one gets a modular weight factor which has different powers of τ and $\bar{\tau}$. Since these lattice functions appear in the partition functions with different powers of $\eta(\tau)$ and $\eta(\bar{\tau})$, that is no problem.

The extra argument $\vec{z}$ which we added to P in the derivation given above is of interest in its own right. It defines a more general *character valued* partition function, which has been used to prove anomaly cancellation. We refer to [19] for an extensive discussion of these issues.

Finally we want to point out that there is a straightforward extension of the correction factor to higher loops. The η - functions are replaced by the determinant of the laplacian on higher genus Riemann surfaces. The modular properties of this object are such that the uncompactified bosonic string is modular invariant for any genus. For the correction factor we can write down an explicit expression (see e.g. [36])

$$C_g(\Gamma) = \sum_{p_i \in \Gamma} \exp(i\pi p_R^i \Omega_{ij} p_R^j - i\pi p_L^i \bar{\Omega}_{ij} p_L^j) \det(Im\Omega)^{\frac{N}{2}}.$$

Although we will not do very much with this object in these lectures, it is good to know that it can be shown, by very similar techniques, to be modular invariant [37]. Since the construction of fermionic strings which follows is heavily based on self-dual lattices, it is very easy to prove that they are also multi-loop modular invariant.

5. Conformal Field Theory and Bosonization

5.1 GENERALITIES

In this section we recall elementary concepts of conformal field theory [38] [39] which will be relevant in our context. The list of more detailed reviews contains [40–45]. We emphasize that conformal field theory is *a priori* a more general notion than string theory. Rather, strings moving in a specific d -dimensional space-time represent particular examples of conformal field theories. We will of course restrict the discussion to that types of (free) conformal field theories we need for our particular construction of heterotic strings.

The basic objects are *conformal* or *primary fields* $\phi(z, \bar{z})$, which are defined to transform as follows under coordinate reparametrizations,

$$\phi(z, \bar{z}) = \left(\frac{\partial w}{\partial z} \right)^h \left(\frac{\partial \bar{w}}{\partial \bar{z}} \right)^{\bar{h}} \phi(w, \bar{w}), \quad (5.1)$$

such that $\phi(z, \bar{z})(dz)^h(d\bar{z})^{\bar{h}}$ is conformally invariant. The complex variable parametrizes either the world-sheet cylinder continued to euclidean space (in which case we use instead the variable $w = 2\sigma_0 - 2i\sigma_1$) or the conformally equivalent punctured complex plane (with $z = e^w$). Fields on the cylinder and on the plane are related by

$$\phi_{cyl}(z, \bar{z}) = z^h \bar{z}^{\bar{h}} \phi_{plane}(w, \bar{w}). \quad (5.2)$$

Any conformal field is thus characterized by a *conformal weight* $(h, \bar{h})$; the scaling weight is then given by $h + \bar{h}$ and the two-dimensional spin by $h - \bar{h}$. We will call fields with half-integer spin two-dimensional fermions, those with integer spin bosons (there exist however also objects which are neither bosons nor fermions). If $\bar{h} = 0$ ($h = 0$), a field is called *chiral* or *right-(left)-moving*; it depends then only (anti-)analytically on z . In the following, we will mainly be concerned with chiral fields. All relevant (local) properties of a given conformal field are characterized by operator product expansions with arbitrary other fields:

$$\phi_i(z) \cdot \phi_j(w) = \sum_k (z - w)^{-h_i - h_j + h_k} \alpha_{ijk} \phi_k(w). \quad (5.3)$$

The non-zero coefficients α_{ijk} define the *fusion rules*. Unless all powers in $(z - w)$ in all possible operator products are integral ($h_k - h_j - h_i \in \mathbf{Z}$), the theory is ill-defined (*non-local*), that is, the correlation functions cannot be unambiguously integrated because of branch cuts. As we will see, imposing locality is a quite strong constraint.

The conformal weight h of any field ϕ_h is determined by the operator product with the stress-energy tensor $T = T_{zz}$, as it generates (5.1):

$$T(z) \cdot \phi_h(w) = h \frac{\phi_h(w)}{(z - w)^2} + \frac{\partial_w \phi_h(w)}{(z - w)} + \text{regular terms} . \quad (5.4)$$

Conformal fields have the special property that they obey (5.4) without creating higher poles in $(z - w)$. For example, the fields (basically derivatives of ϕ_h) appearing in the

non-leading and regular terms in (5.4) are not conformal fields. Quite generally, taking the operator product (5.3) of any two conformal fields creates a series of other operators, of which however only a part are conformal fields.

$T(z)$ itself has weight $h = 2$,

$$T(z) \cdot T(w) = \frac{\frac{1}{2}c}{(z-w)^3} + 2 \frac{T(w)}{(z-w)^2} + \frac{\partial_w T(w)}{(z-w)} + \text{regular terms} , \quad (5.5)$$

but for $c \neq 0$ it is not a conformal field due to the 'anomalous' higher pole. To understand the meaning of the parameter c , let us expand $T(z)$ in modes

$$T(z) = \sum_n z^{-n-2} L_n \quad (5.6)$$

according to the general rule

$$\phi_h(z) = \sum_n z^{-n-h} \phi_{h,n} , \quad \phi_{h,n} = \frac{1}{2\pi i} \oint dz z^{n+h-1} \phi(z) \quad (5.7)$$

for expansions on the plane. The values of n must be such that the integral is well-defined. For example if $h \in \mathbb{Z}$ and $\phi(z)$ is single valued around the origin, then n takes all integral values. Contour integrating (5.5) leads now to the following equivalent expression:

$$[L_n, L_m] = \frac{c}{12} n(n^2 - 1) \delta_{n,-m} + (n - m) L_{n+m}. \quad (5.8)$$

Thus, c represents a central charge (anomaly) in the Virasoro algebra, and the zero mode L_0 is precisely one-half the Hamiltonian of the right-moving (analytic) sector of the theory (of course, the full theory contains in addition an isomorphic Virasoro algebra in the left-moving sector).

Quite generally, as above, the information encoded in operator product expansions is equivalent to the collection of all (anti-)commutation relations among the modes; operator products are just an economic way for writing these.

One of the most important fields is the derivative of a free boson, ∂X , which has $h = 1$. An example of X is the string functional described in sect. 2.1, mapped to the complex plane (called the *Fubini-Veneziano field*). X is itself not a conformal field, a fact which is related to its highly non-local operator product with itself (basically the two-dimensional propagator on the complex plane):

$$X(z) \cdot X(w) = -\log(z - w). \quad (5.9)$$

In contrast, ∂X is a local field:

$$\partial X(z) \cdot \partial X(w) = -\frac{1}{(z-w)^2} + \text{regular terms} . \quad (5.10)$$

For free bosons, the corresponding stress energy tensor is

$$T(z) = -\frac{1}{2} : \partial X \partial X : = -\frac{1}{2} \lim_{w \rightarrow z} \left(\partial X(z) \cdot \partial X(w) + \frac{1}{(z-w)^2} \right), \quad (5.11)$$

where the right hand side indicates what the normal ordering dots mean (similarly,

in the following, products of operators taken at the same point are always implicitly understood to be normal ordered, ie., all singularities are subtracted). Inserting (5.11) into (5.5) one learns that in this case $c = 1$.

Another important type of conformal field is given by $\phi_\lambda = :e^{i\lambda X}:$. From $T(z)\phi_\lambda(w) = \frac{1}{2}\lambda^2 \frac{1}{(z-w)^2}\phi_\lambda(w) + \dots$ follows $h = \frac{1}{2}\lambda^2$. Furthermore, we have from (5.9)

$$\begin{aligned} :e^{i\lambda \cdot X}:(z) \cdot :e^{i\lambda' \cdot X}:(w) &= (z-w)^{\lambda \cdot \lambda'} :e^{i\lambda \cdot X(z) + i\lambda' \cdot X(w)}: \\ &= (z-w)^{\lambda \cdot \lambda'} :e^{i(\lambda + \lambda') \cdot X(w)}(1 + i(z-w)\lambda \cdot \partial X(w) + \dots): \end{aligned} \quad (5.12)$$

Nothing in the above formally changes if we allow $\lambda \rightarrow \lambda^\mu$ and $X \rightarrow X^\mu$ to become vectors (for instance, the space-time coordinates of a string). For $2n$ bosons, $\mu = 1, \dots, 2n$, $T(z) = -\frac{1}{2}\partial X^\mu \partial X_\mu$ and accordingly $c = 2n$.

In *fermionic* string theories, there is a supersymmetry between $2n$ bosons X^μ and $2n$ fermions ψ^μ . The latter obey

$$\psi^\mu(z) \cdot \psi^\nu(w) = \frac{\delta^{\mu\nu}}{(z-w)} + \text{regular terms} , \quad (5.13)$$

which, defining^{*}

$$\psi^\mu(z) = \sum_{n \in \mathbb{Z} + \frac{1}{2}} z^{-n - \frac{1}{2}} \psi_n^\mu, \quad (5.14)$$

is equivalent to $\{\psi_n^\mu, \psi_m^\nu\} = \delta_{n+m}^{\mu\nu}$. The free fermion stress energy tensor is $T_\psi = \frac{1}{2}(\partial\psi)\psi$, and from (5.13) we have

$$T_\psi(z) \cdot \psi^\mu(w) = \frac{1}{2} \frac{\psi^\mu(w)}{(z-w)^2} + \frac{\partial\psi^\mu(w)}{(z-w)} + \text{regular terms} , \quad (5.15)$$

so that $h = \frac{1}{2}$. Furthermore, the operator product of T_ψ with itself tells that a collection of $2n$ fermions has $c = n$. From these basic building blocks one can construct other conformal fields by considering operator products. For instance, the first regular term in the expansion (5.13) is given by

$$J^{\mu\nu}(z) = : \psi^\mu \psi^\nu : , \quad \mu, \nu = 1, \dots, 2n. \quad (5.16)$$

One finds that the modes $J_m^{\mu\nu} = \frac{1}{2\pi i} \oint dz z^m J^{\mu\nu}(z)$ realize a $\widehat{SO}(2n)$ Kac-Moody algebra; in particular, the zero modes $J_0^{\mu\nu}$ are the usual $SO(2n)$ generators, under which ψ_m^μ transform as vectors. As $J^{\mu\nu}(z)$ is given above in terms of fermions, one calls this a *fermionic* construction of the Kac-Moody algebra. Later we will discuss a *bosonic* version.

^{*} We take ψ here to be single valued on the complex plane. A more detailed discussion of fermion boundary conditions will be given in the next section

A very important operator in superconformal theories, i.e., in fermionic strings, is the superpartner of the stress energy tensor, which has in the simplest case the form

$$T_F(z) = i\psi^\mu \partial X_\mu(z) = \sum_m z^{-m-\frac{3}{2}} G_m . \quad (5.17)$$

It has weight $h = \frac{3}{2}$, and the corresponding equ. (5.4) together with (5.5) and

$$T_F(z) \cdot T_F(w) = \frac{\frac{1}{3}c}{(z-w)^3} + \frac{2T(w)}{(z-w)} + \text{regular terms} \quad (5.18)$$

constitute the $N = 1$ superconformal algebra. Equivalently, the mode components satisfy the super Virasoro algebra. More precisely, since we are dealing here only with analytical fields, we describe a $(0,1)$ (*right-moving*) supersymmetry. The zero mode $G_0 = \frac{1}{2\pi i} \oint dz z^{\frac{1}{2}} T_F(z)$ is precisely the two-dimensional supercharge; from (5.18) follows

$$\frac{1}{2}\{G_0, G_0\} = L_0 - \frac{c}{24}. \quad (5.19)$$

T_F indeed transforms fermions into bosons and vice versa:

$$\begin{aligned} T_F(z) \cdot \partial X^\mu(w) &= -\frac{i}{(z-w)^2} \psi^\mu \\ T_F(z) \cdot \psi^\mu(w) &= \frac{i}{(z-w)} \partial X^\mu \end{aligned} \quad (5.20)$$

5.2 STATES

The reason why we considered *conformal fields* is that there is a one to one correspondence between conformal fields ϕ_h of weight h and *highest weight states*[†] $|h\rangle = \phi_h(0)|0\rangle$ of the Virasoro algebra, defined by

$$\begin{aligned} L_0 |h\rangle &= h |h\rangle , \\ L_{+n} |h\rangle &= 0. \end{aligned} \quad (5.21)$$

Thus, h basically is the *mass* of the state $|h\rangle$. Such highest weight states characterize representations of the Virasoro algebra. The other states in a given representation (the *descendants*, which complete the *Verma module*) are obtained by acting with products of L_{-n} to $|h\rangle$; the corresponding operators (e.g., derivatives of ϕ_h) are however not conformal fields, as they create higher poles in (5.4).

Physical states in a string theory are generated by certain combinations of conformal fields called *vertex operators*. They describe the emission of states from the world-sheet,

[†] $\phi_h(0)$ on the plane corresponds to an incoming state at $t = -\infty$ on the world-sheet cylinder.

i.e., the in- or out-coming asymptotic states. For open strings, they have the typical form

$$V(z) = \phi_h(z) e^{ikX}(z) . \quad (5.22)$$

In order to produce conformally invariant correlation functions $\langle V(z_1) \dots V(z_k) \rangle$, these objects must have conformal weight one. Thus, only if the on-shell condition $\frac{1}{2}k^2 = 1 - h$ is satisfied (plus, in general, additional on-shell polarization conditions to prevent higher poles in (5.4)) $V(z)$ is a good vertex operator and $|\phi, k\rangle = V(0)|0\rangle$ a highest weight state. For instance, the tachyon vertex is given by (5.22) with $\phi(z) \equiv 1$, ($k^2 = -m^2 = 2$), and the vector emission vertex in the bosonic string has $\phi(z) \equiv \xi^\mu \partial X_\mu(z)$ with $k^2 = \xi^\mu k_\mu = 0$.

In superconformal theories, matters are a little bit more subtle. Let us consider for a while a fermionic string theory involving a collection of world-sheet fermions ψ^μ in addition to the bosons, where μ is considered a space-time index: $\mu = 1, \dots, 2n$. In addition to states created by purely bosonic operators (i.e., by ∂X and exponentials of X), there exist also highest weight states of the form (see (5.14))

$$(\psi^{\mu_1} \psi^{\mu_2} \dots \psi^{\mu_k})(0)|0\rangle = \psi_{-\frac{1}{2}}^{\mu_1} \psi_{-\frac{1}{2}}^{\mu_2} \dots \psi_{-\frac{1}{2}}^{\mu_k} |0\rangle \quad (5.23)$$

corresponding to antisymmetric tensors of rank k ($h = \frac{1}{2}k$). Taking $\phi_h = \psi^{\mu_1} \psi^{\mu_2} \dots \psi^{\mu_k}$ in (5.22) yields (roughly) vertex operators describing the emission of antisymmetric tensor states; in particular,

$$V = \xi^\mu \psi_\mu e^{ikX} \quad (5.24)$$

is related to the vector boson emission vertex. A quite useful object is the G -parity operator

$$G_{NS} = (-1)^F, \quad (5.25)$$

(F is the two-dimensional fermion number) which is $+1$ (-1) on even (odd) antisymmetric tensors (5.23).

The crucial difference to the bosonic case is that the Hilbert space (i.e., the representation space of the super Virasoro algebra) splits into two sectors, called *Neveu-Schwarz* (NS) and *Ramond* (R). This is due to the fact that there are two *spin structures* on the cylinder: in the NS -sector the fermions ψ have antiperiodic, while in the R -sector they have periodic boundary conditions. Performing a conformal transformation $z = e^w$ we see that (because of the factor $e^{\frac{1}{2}w} = z^{\frac{1}{2}}$ in (5.2)) fermions in the NS -sector are periodic (single-valued), and those in the R -sector antiperiodic (double-valued) along cycles enclosing the origin of the complex plane. Thus in the NS -sector the mode numbers n of ψ_n are half-integral, and in the R -sector integral. In particular, the zero modes in the R -sector obey

$$\{\psi_0^\mu, \psi_0^\nu\} = \delta^{\mu\nu}, \quad (5.26)$$

so they represent space-time ($SO(2n)$ -) γ -matrices. Accordingly the representation space on which ψ^μ act must be spinorial. Thus, denoting by $|\alpha\rangle$ (α = spinor index) the

ground state in the R -sector, a generic R -state (corresponding to (5.23)) looks like

$$(\psi^{\mu_1} \psi^{\mu_2} \dots \psi^{\mu_k})(0) |\alpha\rangle. \quad (5.27)$$

Clearly, while states in the NS -sector (5.23) correspond to space-time antisymmetric tensors, R -states (5.27) lead to space-time spinors. Like in the NS -sector, one can define a grading defined by the operator

$$G_R = \gamma_{2n+1} \otimes (-1)^F \quad (5.28)$$

with eigenvalues ± 1 (counting space-time helicity).

How can we create states in the R -sector via vertex operators ? If we view $\psi(z)$ as an analytic function on the plane, we can describe the double-valuedness around the origin (in the R -sector) as due to a square-root branch cut emanating from there: it flips the boundary conditions between periodic and antiperiodic. This cut creates stress energy (that is, it breaks translation invariance on the plane), and thus corresponds to an operator insertion at $z = 0$. This operator has to be an ordinary (super-)conformal field $S^\alpha(z)$, such as to generate a superconformal highest weight state:

$$|\alpha\rangle = S^\alpha(0) |0\rangle \quad (5.29)$$

S^α is called a *spin* field; it has a very complicated description in terms of the fundamental field ψ^μ . Basically it is the vertex operator for the emission of space-time spinors. In an unitary theory $G_0^2 \geq 0$, hence it follows from (5.19) that the ground state in the R -sector $|\alpha\rangle$ has weight $h = \frac{c}{24}$:

$$L_0 |\alpha\rangle = \frac{c}{24} |\alpha\rangle. \quad (5.30)$$

We thus encounter a problem: for $|\alpha\rangle$ to be a highest weight state and S^α a conformal field, we need $h = 1$. Furthermore, the property that S^α produces a branch cut with ψ is imposed by requiring that the operator product expansion of $\psi(z)$ with $S^\alpha(w)$ contains half-integral powers in $(z - w)$:

$$\psi^\mu(z) \cdot S^\alpha(w) = \frac{1}{(z - w)^{\frac{1}{2}}} (\gamma^\mu)_\beta^\alpha S^\beta + \text{regular terms} . \quad (5.31)$$

Thus naively the theory is *non-local* and as such ill-defined. We will discuss later how these difficulties can be resolved.

The above-explained construction is a special case of what one calls *twisting*: the R -sector is obtained from the NS -sector by a Z_2 -twist. One can of course consider more general, e.g., Z_N -twists, acting in particular also on the bosons ∂X ; this is precisely what happens in the case of *orbifolds*.

5.3 BASICS OF BOSONIZATION

Bosonization, a feature very special to two-dimensions, simply amounts to expressing fermionic field operators like ψ in terms of bosonic ones. The crucial point is that two conformal field theories are physically equivalent, *i.e.*, indistinguishable, if all correlation functions are identical. Now, (the singular parts of) all correlation functions are characterized by all possible operator product expansions (5.3) between the various fields. Thus bosonizing a theory essentially amounts to expressing any given two-dimensional fermionic operator by a certain bosonic expression in such a way that it reproduces all local operator product expansions. In particular, the realization of certain symmetries (*e.g.*, generated by $J^{\mu\nu}$ or T_F) depends only on operator products involving the symmetry currents and not on the way how these are represented.

To bosonize ψ^μ , $\mu = 1, \dots, 2n$, we first switch to a complex basis by defining n pairs

$$\begin{aligned} \frac{1}{\sqrt{2}}(\psi^1 + i\psi^2) &\rightarrow \psi^1, \\ \frac{1}{\sqrt{2}}(\psi^1 - i\psi^2) &\rightarrow \psi^{-1}, \\ \frac{1}{\sqrt{2}}(\psi^3 + i\psi^4) &\rightarrow \psi^2, \\ \frac{1}{\sqrt{2}}(\psi^3 - i\psi^4) &\rightarrow \psi^{-2}, \end{aligned} \quad (5.32)$$

and so on. Then (5.13) takes the form ($j, k = 1, \dots, n$):

$$\begin{aligned} \psi^k(z) \cdot \psi^{-j}(w) &= \frac{\delta_{kj}}{(z-w)} + J^{k,-j}(w) \\ &+ \text{other terms containing descendant fields.} \end{aligned} \quad (5.33)$$

Using (5.12) it is easy to see that we can express our $2n$ fermions by n bosons H_k in a way such that (5.33) is reproduced:

$$\psi^k = e^{iH_k} c_k, \quad \psi^{-k} = e^{-iH_k} c_{-k}. \quad (5.34)$$

(c_k are certain cocycle-phase generating operators which ensure that fermions with different indices really anticommute; in the following, we often will not write such factors explicitly.) Both conformal systems ($\{\psi^\mu\}$, $\{H_k\}$) have central charge $c = n$; in fact, it is only the total value of c that matters.

One can easily bosonize $J^{\mu\nu}$ by just reading it off from (5.33). Switching to the Cartan-Weyl-basis, we have for the off-diagonal ($k \neq j$) currents

$$\begin{aligned} J^{\pm k, \pm j}(z) &\equiv : \psi^{\pm k} \psi^{\pm j} : (z) \\ &= : e^{\pm iH_k} e^{\pm iH_j} : (z) c_{\pm k} c_{\pm j}, \end{aligned} \quad (5.35)$$

and for the diagonal currents, which span the Cartan subalgebra:

$$\begin{aligned} J^{k, -k}(z) &= : e^{iH_k} e^{-iH_k} : (z) \\ &= \lim_{w \rightarrow z} \left(e^{iH_k}(z) e^{-iH_k}(w) - \frac{1}{(z-w)} \right) \\ &= i\partial H_k(z). \end{aligned} \quad (5.36)$$

It is easy to check that these currents represent a bosonic realization of the $\widehat{SO(2n)}$ -

Kac-Moody algebra. Note that the price to pay for bosonization is non-covariance of the formalism: The off-diagonal currents are realized by solitons, i.e., circle-valued bosons, while the diagonal, abelian currents take the canonical form. (5.35) and (5.36) are only a particular example of the more general *Frenkel-Kac* construction (For a detailed discussion we refer to [27].); the latter works for any (simply-laced) Lie algebra, if the root vectors $(\pm e_k \pm e_j)$ of $SO(2n)$ are replaced by the length-squared two root vectors of the algebra in question.

In the last section we have given further examples of operators in the fermionic sector. For instance, in bosonic language the operator corresponding to $\psi^\mu \psi^\nu \dots \psi^\rho$ (5.23) is (a linear combination of operators) of the form

$$\phi_\lambda(z) = e^{i\lambda \cdot H}(z) , \quad (5.37)$$

where λ is an n -dimensional vector of the form $(\pm 1, 0, \pm 1, \dots)$, depending on the precise tensor structure. The G -parity operator (5.25) takes the form

$$G = e^{i\pi(\sum_{k=1}^n \lambda_k)} , \quad (5.38)$$

which is ± 1 depending on if $\sum \lambda_i$ is even or odd. Operators describing (anti-)commuting objects are related to vectors λ of even (odd) length-squares; one can easily see this from equ. (5.12) upon exchanging z and w .

It is clear that the collection of all operators of the form $e^{i\lambda H}$ is related to a *lattice*. This follows simply from the fact that in any operator product (5.12) the vectors add to give a new one: $\phi_\lambda(z) \cdot \phi_{\lambda'}(w) = \phi_{\lambda+\lambda'}(z-w)^{\lambda\lambda'} + \dots$. Consistency, that is, closure of the operator product algebra (5.3), then requires that the vertex operator $\phi_{\lambda+\lambda'}$ exists and generates a state $|\lambda + \lambda'\rangle$ of the theory. The lattice thus is formed by the set of vectors λ obtained by all possible additions of some given basic vectors. (Of course, there exist also vertex operators containing bosons ∂H ; however, these play only a trivial role in the following).

What we are after is a classification scheme for vertex operators or states, in order to characterize the spectrum of a given theory. Given the above lattice, operator products preserve a natural Z_2 -grading, that is, a split into vectors of even and odd length-squares; equivalently, a split into eigenvalues ± 1 of G_{NS} . More specifically, the lattice characterized above is precisely the weight lattice of D_n generated by the conjugacy classes (0) and (v). In particular, the non-zero roots of D_n $(\pm e_i \pm e_j)$ which appear in the bosonized currents (5.35) belong to the neutral class (0), while the fermions $\psi^{\pm i}$ belong to (v); the operator product expansion (5.13) of two fermions, which contains $J^{\mu\nu}$ in its regular terms, clearly reflects the rule $(v) + (v) = (0)$.

In this framework it is very easy to describe states in the R -sector. In the R -sector, the fields ψ^μ are twisted, as explained in the previous section. In bosonic language such a twist corresponds to a *shift* of the lattice, a fact that is made plausible by the identity

$$\left(\frac{1}{\prod (1 + q^{2n})} \right) \sum_{n \in \mathbb{Z}} q^{(n+\theta)^2} = q^{\theta^2} \prod_{n \geq 0} (1 + q^{2n+2\theta+1})(1 + q^{2n-2\theta+1}) , \quad (5.39)$$

which is a mathematical manifestation of bosonization: the RHS is the partition function of two fermions with twist θ , while the LHS represents the partition function of one

compact boson taking values on a (one-dimensional) integer lattice shifted by θ . More precisely, the sum at the LHS gives the contribution of all solitonic states created by $\{\phi_\lambda\}$, $\lambda = n + \theta$, $n \in \mathbf{Z}$, and the product is the contribution of all possible combinations of derivatives of ∂H . The case at hand corresponds to n copies of (5.39), i.e., to $2n$ fermions and to an integer lattice shifted by $\theta_1 = \dots = \theta_n = \pm \frac{1}{2} \bmod 2$. In other words, the R -sector is described by the spinor conjugacy classes (s) and (c) of D_n . To which particular spinor class a given state $|\lambda\rangle$ belongs is determined by the eigenvalue of G_R (5.28) which is, in bosonic language, up to a phase, identical to G_{NS} given in (5.38).

Every given conjugacy class can of course be represented by a particular vector belonging to that class; all other vectors in the same class are obtained by adding the root lattice, i.e., all vectors in the neutral class (0). Thus all states (up to derivatives) in the R -sector are of the form

$$(e^{i\lambda \cdot H} S^\alpha)(0) |0\rangle, \quad \lambda \in (0) \text{ of } D_n, \quad (5.40)$$

and we can think of the ground states of the R -sector $|\alpha\rangle = S^\alpha(0) |0\rangle$ as being generated by shift vectors of minimal lengths in the spinor conjugacy classes of D_n . Hence we find the following bosonic expression for the spin fields:

$$S^\alpha(z) = e^{i\lambda_s \cdot H}(z), \quad \lambda_s = (\pm \frac{1}{2}, \pm \frac{1}{2}, \dots, \pm \frac{1}{2}). \quad (5.41)$$

In fact, there are two spin fields, S^α and $S^{\dot{\alpha}}$, with opposite $SO(2n)$ helicity belonging to (s) and (c) of D_n ; they are distinguished by an even or odd number of $-$ signs in λ_s , respectively. The number 2^{n-1} of possible sign choices for λ_s in each case is precisely the dimension of the two spinor representations of $SO(2n)$: $\alpha, \dot{\alpha} = 1, \dots, 2^{n-1}$. The conformal weights of S^α , $S^{\dot{\alpha}}$ are $h_s = \frac{1}{2}\lambda_s^2 = \frac{1}{8}n$; since in a supersymmetric theory $n_f = n_b = 2n$, and $c_b = 2c_f = 2n$, we find $h_s = \frac{c}{24}$ in accordance with (5.30).

Many properties of states in the various conjugacy classes can be determined solely in terms of addition and multiplication rules of (0), (v), (s) and (c), in connection with the basic formula (5.12) for operator products of exponentials. For instance, the twisted boundary conditions for fermions in the R -sector follow from the branch cut arising from $(z - w)^{\lambda \cdot \lambda'}$ in (5.12) by the multiplication rule $(v) \cdot (s) \in \mathbf{Z} + \frac{1}{2}$.^{*} (Anti-)commutation properties are also easily determined from (5.12) upon exchanging z and w ; one finds that while all states in (0) commute and those in (v) anticommute, states in the spinor conjugacy classes (for generic n) have no definite (anti-)commutation properties.

The above discussion dealt only with periodic and antiperiodic fermionic boundary conditions, i.e., with Z_2 -twisted fermions. We have just seen that these can be described, in bosonic language, in terms of the weight lattice D_n of $SO(2n)$. It is obvious that one can easily generalize these concepts to other lattices (for instance, the weight lattices A_n of $SU(n + 1)$). (One can, of course, consider also lattices which are products of weight lattices of several Lie groups, and/or lattices which are not at all related to Lie groups.) One then describes fermions with more complicated boundary conditions, and other objects, which need not be either fermionic or bosonic. Like the spin fields,

^{*} This is nicely illustrated by (5.31), which also reflects the fusion rule $(v) + (s) = (c)$.

such operators are quite difficult to describe in terms of the fermionic variables ψ^μ . In bosonic language, however, all operators have the same (exponentiated) form, and ψ^μ are no more fundamental than S^α . Clearly, the bosonic lattice formulation provides the most simple and elegant description of such objects.

5.4 COMPACTIFICATION REVISITED

We have seen that certain conformal fields (like ψ^μ , $J^{\mu\nu}$ and spin fields) can easily be described in terms of free compact bosons. In fact, it is sometimes possible to interpret these as (torus-)compactified higher dimensional coordinates.

Consider, for example, torus compactification (as discussed in chapter 4) of the 26-dimensional closed bosonic string on an 8-dimensional even lorentzian self-dual lattice $(E_8)_L \times (E_8)_R$. This is the same as $(D_8)_L \times (D_8)_R$ with conjugacy classes $(0,0)$, $(0,s)$, $(s,0)$ and (s,s) . Using the compactified coordinates $X^I(z)$, $X^I(\bar{z})$ in the left- and right-moving sectors, it is easy to discuss the conformal field theory of this model in terms of operators like the currents of $(E_8)_L \times (E_8)_R$.

The currents belonging to (0) of the subgroup $D_8 \sim SO(16)$ are precisely given by (5.35) and (5.36); the currents (5.36) belonging to the Cartan subalgebra correspond to Kaluza-Klein states, the off-diagonal ones to winding states, i.e., solitons. The remaining currents belonging to (s) , i.e., to $E_8/SO(16)$ are realized by spin fields (5.41) of $SO(16)$. This is possible since a spinor weight of D_8 has length-squared two, so that it can represent a root vector (we say that the spinor conjugacy class of D_8 extends D_8 to E_8). Now, what is relevant in the two dimensional sense are the properties of the conformal fields of the theory, and not if they can be interpreted in terms of compactified higher dimensions. According to the fermion-boson equivalence in two dimensions, in order to describe the same theory we could also start with a fermionic version of the above objects. The $SO(16)$ currents are then represented by bilinears as in (5.16), however, the currents represented by the spin fields have a very complicated, transcendental dependence on the fermions [46]). In this formulation, there is no trace of higher dimensions left! Apparently the concept of compactification is quite delicate in the string context.

The above example represents a compactified theory in the very sense. There exist however more general conformal field theories, which do not admit a compactification interpretation (we will encounter such theories in the next chapter). One should therefore not confuse compact bosons (equivalent to fermions) with compactified space-time coordinates (which are a special case).

For closed strings, which are described by a combination of a right-moving (z -dependent) and a left-moving ($\bar{z}$ -dependent) conformal field theory, we require vertex operators to have conformal weight $(\bar{h}, h) = (1, 1)$. Consequently, these consist of a left- and a right-moving piece together with the exponentiated momentum,

$$V(\bar{z}, z) = \phi_h(z) \phi_{\bar{h}}^l(\bar{z}) e^{ik \cdot X}(\bar{z}, z), \quad (5.42)$$

with the condition $\bar{h} = h$. ϕ_h and $\phi_{\bar{h}}^l$, belonging to independent left- and right-moving sectors, are a priori not related to each other. For instance, for the compactified bosonic

string above,

$$V_\mu^{ab}(\bar{z}, z) = J^{ab}(\bar{z})\partial X_\mu(z)e^{ikX} \quad (5.43)$$

($\mu = 1, \dots, 18$) describes a gauge vector boson belonging to the adjoint representation of the left-moving E_8 with generators J^{ab} , whereas

$$V_\mu^{a'b'}(\bar{z}, z) = J^{a'b'}(z)\bar{\partial} X_\mu(\bar{z})e^{ikX} \quad (5.44)$$

describes the emission of a vector boson of the right-moving E_8 . On the other hand,

$$V_{\mu\nu}(\bar{z}, z) = \bar{\partial} X_\nu(\bar{z})\partial X_\mu(z)e^{ikX} \quad (5.45)$$

describes the emission of gravitons, antisymmetric tensor fields and dilatons in $d = 18$.

In the next section we will apply the above ideas to a bosonic description of heterotic strings, which in general do not admit a compactification interpretation.

6. NSR Model, Superstring and Heterotic String

6.1 THE COVARIANT NSR MODEL IN BOSONIC LATTICE FORMULATION

We will first introduce the covariant ten-dimensional *NSR* model [47] (the *spinning string*) in bosonic language, which will allow a straightforward generalization to lower dimensions in the next chapter.

The covariant *NSR* model can be described by a specific conformal field theory whose main variables are ten bosons X^μ and ten fermions ψ^μ , which play the role of space-time coordinates. Thus, the basic conformal operators are $\partial X^\mu(z)$ and $\psi^\mu(z)$, whose properties have been discussed already in the foregoing chapter. Since these fields depend *analytically* on z , we deal only with right-moving fields. What we are after is however closed, in particular, heterotic string theory. Thus we will eventually have to pair the right-moving *NSR* model with an a priori independent left-moving ($\bar{z}$ -dependent) conformal field theory. This will be explained further below.

On a flat background, and in superconformal gauge, the action takes the free form

$$S_{NSR} = -\frac{1}{2\pi} \int d^2z \left(\partial X^\mu \bar{\partial} X_\mu - \psi^\mu \bar{\partial} \psi_\mu + \mathcal{L}_{ghost} \right). \quad (6.1)$$

Let us denote by c and b the (anticommuting) ghost and antighost fields for fixing conformal invariance, and by γ and β the (commuting) ghost and antighost fields for fixing the fermionic part of superconformal invariance. Then the ghost lagrangian is given by

$$\frac{1}{2}\mathcal{L}_{ghost} = b\bar{\partial}c + \beta\bar{\partial}\gamma. \quad (6.2)$$

The conformal field theory of the X^μ and ψ^μ system was developed in the foregoing chapter; we indicated already, that this part of the theory by itself suffers from severe deficiencies. For example, ψ^μ as a two-dimensional spinor anticommutes with itself

(cf.(5.13)). Since it carries however a space-time vector index (it appears in emission vertices for space-time tensors), one would rather like to have it generating space-time bosons. Moreover, in order to describe the R -sector, we had to introduce the spin field S^α . This field has conformal weight $h = \frac{n}{8} = \frac{5}{8}$ and not $h = 1$, what would be needed for a vertex operator describing massless space-time fermions. Inserted at some point on the complex plane, S^α creates a branch cut (5.31) in order to change the boundary conditions of ψ^μ (in bosonic formulation, this is related to the rule $(v) \cdot (s) \in \mathbf{Z} + \frac{1}{2}$ for the conjugacy classes of D_5 , which is the Lie algebra of the Wick-rotated Lorentz group in ten dimensions). But the very presence of this cut makes the theory non-local. Another problem we encounter is that S^α does not anticommute with itself as it should as a space-time fermion. This fact is reflected by the rule $(s) \cdot (s) \in 2n + \frac{5}{4}, n \in \mathbf{Z}$ in D_5 , which also reveals the presence of even worse cuts.

It is important to notice that many of these and other unpleasant features trace back to properties of the various conjugacy classes of the lattice D_5 , if one uses the bosonic description of the ψ^μ . We thus expect that a resolution lies in the modification of this lattice.

The only new ingredient of the NSR model (6.1) compared to the conformal field theory of X^μ and ψ^μ as described in the last chapter is the ghost system. It is in fact a crucial ingredient, which can cure all above-mentioned diseases. To be more precise, let us first display some essential properties of the ghosts: $b(z) \cdot c(w) = c(w) \cdot b(z) = \frac{1}{(z-w)}$, $\beta(z) \cdot \gamma(w) = -\gamma(w) \cdot \beta(z) = -\frac{1}{(z-w)}$. Their conformal weights are

$$\begin{aligned} c : \quad & h = -1, \\ b : \quad & h = 2, \\ \gamma : \quad & h = -\frac{1}{2}, \\ \beta : \quad & h = \frac{3}{2}, \end{aligned} \tag{6.3}$$

The contribution to the full energy momentum tensor is $T_{gh} = c\partial b + 2\partial c b - \frac{1}{2}\gamma\partial\beta - \frac{3}{2}\partial\gamma\beta$. Substituting this into (5.5), we learn that the contribution to the central charge of the b, c system is $c_{bc} = -26$, and similarly $c_{\beta\gamma} = 11$; thus altogether we have $c_{ghost} = -15$ and this cancels precisely the contribution of the ten X^μ plus ten ψ^μ . So the conformal anomaly disappears, which seems to be crucial for the consistency of the theory. This does of course *not* imply that the string theory can only be consistent in ten dimensions. Rather, what counts here is only the *total* value of $c_{matter} = 15$, and not any particular decomposition in terms of contributions of X^μ and ψ^μ . Allowing for internal degrees of freedom, which carry 'missing' central charge, will allow us later to construct string theories in lower dimensions than ten.

As we intend to discuss only the spectrum of the theory and not other aspects (like correlation functions), we will explain only aspects of the ghost system relevant here. The role of the b, c system is then quite simple: For our purposes, all what the b, c system does is to provide the correct vacuum energy. In sect. 5.2 we described how the states of the theory are created by acting with conformal fields on some vacuum. We did however not specify the vacuum itself. The correct vacuum (of the bosonic string) is simply $|0\rangle = c(0)|0\rangle_{SL(2,C)}$, where $|0\rangle_{SL(2,C)}$ is the true $SL(2, C)$ invariant vacuum of

the theory, which is annihilated by $L_n, n \geq -1$. (Another way of describing the same situation is to redefine the vertex operators of the theory $V \rightarrow cV$; these objects then act on the $SL(2, C)$ invariant vacuum state.) Since c has weight $h = -1$, the level of $|0\rangle$ is -1 ; it represents the tachyon state of the bosonic string. Massless states, therefore, are build upon $|0\rangle$ using conformal fields of weight $h = 1$. This gives us back the mass formula of the bosonic string (4.10) ($\mathcal{N}$ counts the number of derivatives in a vertex operator).

In fermionic strings, we have in addition the commuting ghosts β, γ . Their role is much more involved. As we cannot discuss all issues in great detail, we will just give recipes for handling these ghosts (details can be found e.g. in [39]). It is very convenient to bosonize also these fields:

$$\beta = e^{-i\phi} \partial \xi, \quad \gamma = e^{i\phi} \eta. \quad (6.4)$$

Here, ϕ is a chiral boson with opposite norm, $\phi(z) \cdot \phi(w) = \log(z - w)$, and ξ and η are auxiliary fermions, which will play no role in the following. They are needed above since one cannot bosonize the commuting fields β, γ solely in terms of an anticommuting exponential of one boson.

One important feature of the ϕ -ghost is that the conformal field $e^{iq\phi}$ has a peculiar conformal weight:

$$\begin{aligned} T_{\beta\gamma}(z) \cdot e^{iq\phi}(w) &= -\frac{1}{2}q(q+2) \frac{e^{iq\phi}(w)}{(z-w)^2} + \dots \\ \rightarrow h &= -\frac{1}{2}q(q+2). \end{aligned} \quad (6.5)$$

The linear term $(-q)$ originates in an anomaly of the ghost number current $j = \beta\gamma$, and the negative sign is due to the negative metric of ϕ , which also implies

$$e^{iq\phi}(z) \cdot e^{ip\phi}(w) = (z-w)^{-qp} e^{i(q+p)\phi}(w) + \dots \quad (6.6)$$

Similar to the b, c system, also in the β, γ system the vacuum gets modified: it is characterized by a certain *bose ghost sea charge* q :

$$|0\rangle_{\beta\gamma} \equiv |q\rangle = e^{iq\phi}(0) |0\rangle. \quad (6.7)$$

To specify the possible values of q , note first that the $\beta\gamma$ ghosts are related to gauge degrees of the (non-dynamical) world-sheet gravitino, hence should have the same periodicity properties. Since that couples to the supercurrent $T_F = i\psi\partial X$, it follows that because of world-sheet supersymmetry the $\beta\gamma$ ghosts should have the same boundary conditions as the NSR fermions ψ^μ . This implies that we need *different* vacua in the NS - and R -sectors of ψ^μ : in the NS -sector $q \in \mathbf{Z}$ while in the R -sector $q \in \mathbf{Z} + \frac{1}{2}$. There exists now still an ∞ -fold ambiguity in what values of q we actually choose. In fact, all values differing by integers are equivalent. Thus, for several reasons, let us choose as ‘canonical’ ghost vacua

$$\begin{aligned} |0\rangle_{\beta\gamma}^{NS} &= |-1\rangle, \\ |0\rangle_{\beta\gamma}^R &= \left| -\frac{1}{2} \right\rangle. \end{aligned} \quad (6.8)$$

Later we will comment on other choices.

Similar to the b, c system, it is sometimes more convenient (in particular, for computing correlation functions) to redefine the vertex operators so that they act on a simple vacuum. Let us take here $|0\rangle = c(0)|0\rangle_{SL(2,C)}$ with level -1 . Then

$$V_{NS} e^{-i\phi}(z) \quad \text{and} \quad V_R e^{-i\frac{1}{2}\phi}(z) \quad (6.9)$$

are the vertex operators we will consider from now on. By (6.5), the mass formula reads

$$\begin{aligned} \frac{1}{8}m^2 &= h - 1, \\ h &= \frac{1}{2}\lambda^2 + \mathcal{N} - 1 + \begin{cases} \frac{1}{2} & NS\text{-sector} \\ \frac{3}{8} & R\text{-sector} \end{cases} \end{aligned} \quad (6.10)$$

which displays two different normal ordering constants in the NS - and R -sectors. This formula already cures some of our problems: the fermion emission vertices for both chiralities now become

$$V_F^{spinor(s)} = u_\alpha S^\alpha e^{-i\frac{1}{2}\phi} e^{ikX}, \quad V_F^{spinor(c)} = u_{\dot{\alpha}} S^{\dot{\alpha}} e^{-i\frac{1}{2}\phi} e^{ikX}, \quad (6.11)$$

and therefore (if $k^2 = 0$) have weight $h = 1$ (as opposed to $h = \frac{5}{8}$ without ghosts) what is needed for good vertex operators. Similarly, the vector boson emission vertex

$$V_B^{vector} = \xi_\mu \psi^\mu e^{-i\phi} e^{ikX}, \quad (6.12)$$

now also has $h = 1$ (as opposed to $h = \frac{1}{2}$) for vanishing momentum. V_F^{spinor} and V_B^{vector} therefore describe massless particles in ten dimensions. We can, of course, also build other operators like $V_B^{tachyon} = e^{-i\phi} e^{ikX}$ or $V_B^{tensor} = \xi_{\mu\nu} \psi^\mu \psi^\nu e^{-i\phi} e^{ikX}$, with mass $-\frac{1}{2}$ and $+\frac{1}{2}$, respectively. The ground states in the NS - and R -sectors are $|V_B^{tachyon}\rangle$ and $|V_F^{spinor}\rangle$, respectively.

We mentioned above that our other problems could possibly be solved by modifying the lattice D_5 related to the bosonized fermions. Contemplating on (6.9), the clever reader recognizes that part of these problems is solved already. To make this more clear, let us use a trick, invented by [15]: just consider the ghost charge q as part of an enlarged covariant lattice. That is, associate to any of the above (and all other possible) bosonized vertex operators (i.e., states) a six-dimensional vector $w = (\lambda | q)$, e.g.,

$$\begin{aligned} w^{spinor} &= (\pm\frac{1}{2}, \pm\frac{1}{2}, \pm\frac{1}{2}, \pm\frac{1}{2}, \pm\frac{1}{2} | -\frac{1}{2}) \\ w^{vector} &= (0, \dots, \pm 1, \dots, 0, \dots | -1) \\ w^{tachyon} &= (0, 0, 0, 0, 0 | -1) \end{aligned} \quad (6.13)$$

and $w^{tensor} = (0, \dots, \pm 1, \dots, \pm 1, \dots | -1)$, etc. In order to efficiently characterize the theory, let us investigate the nature of this six dimensional lattice Γ_6 . It looks rather like an extension $D_5 \rightarrow D_6$, but we have to take the switched sign in the ghost metric into account. The natural inner product on this lattice is defined by the power of $(z - w)$ in

operator product expansions, because that determines the various (anti-)commutation properties of the fields and measures locality. From (5.12) and (6.6) we have

$$V_\lambda e^{iq\phi}(z) \cdot V_{\lambda'} e^{iq'\phi}(w) = (z-w)^{\lambda \cdot \lambda' - qq'} V_{\lambda+\lambda'} e^{i(q+q')\phi}(w) + \dots \quad (6.14)$$

Therefore the natural inner product has *lorentzian* metric $(++++|-)$:

$$\langle w, w' \rangle = \lambda \cdot \lambda' - qq' \quad (6.15)$$

The same inner product appears also in the mass formula for arbitrary ghost charges:

$$\frac{1}{8}m^2 = \frac{1}{2} \langle w, w \rangle - q + \mathcal{N} - 1 \quad (6.16)$$

Accordingly, we denote the lorentzian lattice by $\Gamma_{5,1} = D_{5,1}$, with the comma indicating the sign flip in the metric.

As the ϕ -charge adds in (6.14), is it clear that for closure of the operator product algebra one needs also vertex operators with different ghost charges than the canonical ones (6.9). However, as mentioned above, vertex operators with ghost charges differing by integral units can represent (or generate) the same physical state. That is, there is a copy of every canonical vertex operator in every ghost sector; the different copies are also called *pictures*. There exists in fact an operation which maps a given vertex operator with ghost charge q into one with $q+1$:

$$V_{q+1}(z) =: \mathcal{P}_{+1} V_q(z) \quad (6.17)$$

The *picture changing operator* carries one unit of ghost charge, and is given by

$$\mathcal{P}_{+1}(z) = e^{i\phi} T_F(z) = ie^{i\phi} \psi \cdot \partial X(z) + \dots \quad (6.18)$$

Thus, apart from the ghost part, picture changing is more or less the same as performing a two dimensional supersymmetry transformation^{*}.

For instance, picture changing the vector emission vertex V_B^{vector} in the (-1) -picture (6.12) leads to its copy in the (0) -picture:

$$:\mathcal{P}_{+1} V_{B(-1)}^{vector}: (z) = V_{B(0)}^{vector}(z) = \xi \cdot [\partial X + i\psi(k \cdot \psi)] e^{ikX}(z) \quad (6.19)$$

which (apart from the ψ -dependent piece) is the same as the vector emission vertex of the bosonic string (5.43). One easily checks that the mass formula (6.16) gives the same answer for both $V_{B(-1)}^{vector}$ and $V_{B(0)}^{vector}$. This already follows from the fact that the conformal weight h of $\mathcal{P}_{+1}(z)$ vanishes. In general, $\mathcal{P}_{+1}$ always acts in a way that compensates the effect of changing the q -charge. Thus the mass spectrum of the theory is bounded from below, a subtle feature which is not at all manifest in the mass formula (6.16).

* In fact, the ghost part is necessary in order to have a well-defined local action of T_F on states in the R -sector; remember the branch cut of ψ in T_F . Roughly speaking, one may regard $e^{i\phi}$ as a bosonized anticommuting coordinate, playing the role of ϵ in $[\epsilon Q^{SU SY}, \dots]$.

Upon bosonization, we can represent $\mathcal{P}_{+1}$ by the vector $w^{PC} = (0 \dots, \pm 1, 0 \dots | +1) \in D_{5,1}$. Picture changing then acts on the lattice(-part of the theory) simply as shifting by w^{PC} . In order for this operation to be well-defined and local, w^{PC} has to have integral inner product with all other lattice vectors. Although trivially satisfied in the ten dimensional *NSR* model, this condition will prove quite non-trivial for the lower dimensional theories we will consider below, as for these theories the supercurrent T_F (i.e., w^{PC}) can take a different form.

The structure of (6.19) is more complicated than that of (6.12) (and even more complicated in higher pictures). In general, the canonical ghost sectors (6.9) have the virtue of being simplest in the sense that the vertex operators are related to lattice vectors in the most simple way. In the other ghost sectors, there is no manifest relation between lattice vectors and physical vertex operators; they are in general given by more complicated linear combinations. For our purposes, however, these complications play no role, as features like locality or (anti-)commutativity do not depend on them. That is, all relevant information about state classification is carried by the lattice vectors (oscillators, i.e., derivatives, also play no role). For our purposes, we can therefore represent the spectrum of vertex operator in all ghost sectors by the lattice $D_{5,1}$.

This certainly does *not* mean that the partition function of the physical states is given by the partition function of the lattice! There is a priori no one-to-one relation between lattice vectors and physical states[†]. There exists however a brute force method to count the physical degrees of freedom correctly: let us split every lattice vector $w = (\lambda | q) \in D_{5,1}$ into a four dimensional 'light-cone' (describing the Wick-rotated transverse Lorentz group $SO(8)$) and a two dimensional 'ghost' part, according to the decomposition

$$D_{5,1} = D_4^{light-cone} \times D_{1,1}^{ghost} \quad (6.20)$$

$$w = (u, x), \quad u \in D_4^{light-cone}, \quad x \in D_{1,1}^{ghost}.$$

Then, by definition, the physical states are those associated with

$$x = x_0 = \begin{cases} (0 | -1) & \text{NS-sector} \\ (-\frac{1}{2} | -\frac{1}{2}) & \text{R-sector} \end{cases} \quad (6.21)$$

One easily checks that this *physical state selection rule* is just a projection to light-cone states, in the canonical pictures (the astute reader also notices that for the above choice of x_0 above the mass formula (6.16) turns precisely into the light-cone mass formula). From the standpoint of conformal field theory, this separation of vectors into 'physical' and 'unphysical' ones is quite artificial (and ambiguous), as vertex operators are equally good in arbitrary pictures. The above selection rule is nothing but a help to identify and count the physical states of the theory.

[†] For example, w^{vector} (6.13) associated with $V_{B(-1)}^{vector}$ given in (6.12) describes ten degrees of freedom (which is the correct count), while $w^{vector} = (\dots \pm 1, \dots \pm 1, \dots | 0)$ associated with the picture changed, equivalent vertex operator $V_{B(0)}^{vector}$ (6.19) has 40 components.

Having now everything reduced to the properties of the lattice $D_{5,1}$, let us split it (i.e., the set of vertex operators) into cosets or conjugacy classes, much like we did for the case of D_n in section 5.3. In particular, as it is obvious from (6.13), V_B^{vector} belongs to (0) of $D_{5,1}$, $V_B^{tachyon}$ and V_B^{tensor} to (v) and V_F^{spinor} to (s) or (c), respectively. Any other vertex operator of the theory belongs to one of these four classes, and their lattice parts can be obtained by adding 'root' vectors belonging to the neutral class (0) to the vectors (6.13); these are simply ground state (lowest mass) representatives of the conjugacy classes (note in particular that the action of the picture changing operator $\mathcal{P}_{+1}$ belonging to (0) does not change the class of a vertex operator).

Then, like for D_n , we can represent all (anti-)commutation relations, fusion rules and possible nonlocalities of the NSR model in terms of addition and multiplication rules of the conjugacy classes of $D_{5,1}$ (The dot denotes the lorentzian inner product):

$$\begin{aligned}
 (0) + (0) &= (0), & (0) \cdot (0) &\in 2\mathbf{Z} \\
 (0) + (v) &= (v), & (0) \cdot (v) &\in \mathbf{Z} \\
 (0) + (s) &= (s), & (0) \cdot (s) &\in \mathbf{Z} \\
 (0) + (c) &= (c), & (0) \cdot (c) &\in \mathbf{Z} \\
 (v) + (v) &= (0), & (v) \cdot (v) &\in \mathbf{Z} \\
 (v) + (s) &= (c), & (v) \cdot (s) &\in \mathbf{Z} + \frac{1}{2} \\
 (v) + (c) &= (s), & (v) \cdot (c) &\in \mathbf{Z} + \frac{1}{2} \\
 (s) + (s) &= (0), & (s) \cdot (s) &\in \mathbf{Z} \\
 (c) + (c) &= (0), & (c) \cdot (c) &\in \mathbf{Z} \\
 (s) + (c) &= (v), & (s) \cdot (c) &\in \mathbf{Z} + \frac{1}{2}
 \end{aligned} \tag{6.22}$$

Note that all length(-squares) are integral, but not even; $D_{5,1}$ is therefore an *odd* lattice. (6.22) tells us that operators in (0) (even lengths) commute, those in (s), (c) (odd lengths) anticommute with each other. This is precisely what we like to have, since (0) contains (odd-rank antisymmetric) tensors, and (s), (c) spinors in space-time. Furthermore, $V_B^{vector} \in (0)$ and $V_F^{spinor} \in (s)$ are now local with respect to each other: the branch cut in $\psi(z) \cdot S(w)$ is cancelled by that of $e^{-i\phi}(z) \cdot e^{-i\phi/2}(w)$.

However, we are not yet completely done: the (even-rank antisymmetric) space-time tensors belonging to (v) anticommute and thus violate spin-statistics, and they are also non-local with respect to the spinors (as can be inferred from the $\frac{1}{2}$'s in (6.22)). Finally, the class (v) contains the tachyonic ground state of the NS -sector —an unpleasant feature. Now, what we have constructed so far is the spinning string, which is known to be inconsistent. One has to do little more to really obtain the superstring.

6.2 SUPERSTRING AND HETEROTIC STRING

The remaining diseases of the spinning string are simply eliminated by restricting to a subset of the possible vertex operators, characterized by (0) and (s) (or (c)). One easily concludes from (6.22) that all operator products are local and close within this subset. The spin-statistic relation is satisfied as (v) is absent now: tensors and spinors are related to even and odd vectors on $D_{5,1}$. In particular, as massless states we keep $|V_B^{vector}\rangle$ and $|V_F^{spinor}\rangle$ which form a supersymmetry multiplet. Actually one can prove (we will do this explicitly below) that the whole spectrum is supersymmetric. Accordingly, one calls this theory the superstring.

Keeping only states in (0) and (s) can also be expressed as projecting onto states $|w\rangle$ with 'positive G -parity' $G = 1$, where

$$G|w\rangle \equiv (-1)^{F+F^\phi}|w\rangle = e^{2\pi i \langle w^{spinor(s)}, w \rangle} |w\rangle \quad (6.23)$$

generalizes (5.38) by including the ϕ -ghost charge F^ϕ . This is called *GSO-projection* [48].

The *GSO*-projected lattice is smaller in the sense that it contains less conjugacy classes than the lattice we had before. Nevertheless we still will call it $D_{5,1}$, as according to usual notation a lattice is named after its root sublattice (0). (Remember that in order to characterize a lattice completely, one has to provide in addition a list of the conjugacy classes actually present.)

In fact, the projected lattice $D_{5,1}$ is *self-dual* as one can easily check. This self-duality is actually necessary for the consistency of the superstring: it ensures modular invariance of the partition function, which contains a lattice factor

$$\mathcal{L}_{5,1}(\tau) = \sum_{\substack{w=(u,x) \in D_{5,1} \\ x=x_0}} e^{i\pi\tau u^2} e^{2\pi i q} . \quad (6.24)$$

The sum is only over the states satisfying the selection rule (6.21), and the phase counts space-time fermions and bosons with opposite signs. The lattice function $\mathcal{L}_{5,1}$ differs in further ways from lattice partition functions (4.18) we encountered earlier. In particular, the lattice is odd rather than even, and in writing the exponent above we used the peculiar mass formula (6.16). It was shown in [16] that all these features conspire in precisely the right way to make the theory modular invariant. In the next section we will present an independent proof, which furthermore extends to higher loops.

Knowing this, we could have started already with a self-dual lattice from the very beginning and would never had encountered the problems of the spinning string; locality is guaranteed by the fact that on a self-dual lattice all inner products are integral. This will be our philosophy for constructing further string theories.

We have now everything prepared for generalizing the above to heterotic strings. We simply have to combine the above-explained, right-moving *NSR* model (not necessarily the superstring!) with a left-moving bosonic string. As the conformal anomaly $c_{bc} = -26$

for the bosonic string, we need in addition to the ten space-time coordinates X^μ 16 compact bosons $X^I(\bar{z})$ (or equivalently, 32 fermions):

$$S_{het} = -\frac{1}{2\pi} \int d^2z \left(\partial X^\mu \bar{\partial} X_\mu + \psi^\mu \bar{\partial} \psi_\mu + 2b\bar{\partial}c + 2\beta\bar{\partial}\gamma + \partial X^I \partial X_I + 2\bar{b}\partial\bar{c} \right). \quad (6.25)$$

($I = 1 \dots 16$). What is the momentum lattice of the compact bosons like? It is convenient to combine this 16-dimensional lattice Γ_{16} with that related to the right-moving *NSR* model:

$$\Gamma_{16;5,1} \equiv (\Gamma_{16})_L \times (D_{5,1})_R. \quad (6.26)$$

We cannot strongly enough emphasize that this does *not* necessarily represent a direct product decomposition! As (6.26) denotes only the root sublattices, there may still exist non-trivial correlations between the conjugacy classes of Γ_{16} and $D_{5,1}$. In order to characterize (6.26), let us consider (in analogy to (6.14)) the operator product of two *closed string* vertex operators (5.42):

$$V_w(\bar{z}, z) \cdot V_{w'}(\bar{w}, w) = (\bar{z} - \bar{w})^{w_L \cdot w'_L} (z - w)^{\lambda \cdot \lambda' - qq'} V_{w+w'}(w) + \dots,$$

$$V_w(\bar{z}, z) = e^{iw_L^I X^I}(\bar{z}) e^{i\lambda \cdot H} e^{iq\phi}(z), \quad w = (w_L; w_R), \quad w_R = (\lambda | q). \quad (6.27)$$

Upon rewriting the z - and w -dependent part using $(\bar{z} - \bar{w}) = \frac{|z-w|}{(z-w)}$, the condition for locality, i.e., the absence of branch cuts becomes

$$\langle w, w' \rangle \in \mathbf{Z}, \quad (6.28)$$

where the inner product has metric $[(-)^{16}; (+)^5 | (-)]$. Thus, $\Gamma_{16;5,1}$ has to be an integral, lorentzian lattice with respect to this metric. Note however that the two sets of $(-)$ -signs have completely different origins: the $(-)$ -sign in the left-moving sector is the same as that arising in the lattice of a compactified closed bosonic string, as discussed in chapter 4: w_L and w_R play the role of p_L and p_R in sect.4.2, respectively.

The full partition function of such a theory has the form

$$\text{Tr} \left[e^{i\pi\tau H'_R} e^{-i\pi\bar{\tau} H'_L} e^{2\pi i F^\phi} \right] = \frac{1}{\eta^{12} \bar{\eta}^{24}} \sum_{\substack{w=(w_L; w_R) \in \Gamma_{16;5,1} \\ w_R=(\lambda|q)=(u, x_0)}} e^{-i\pi\bar{\tau} w_L^2} e^{i\pi\tau u^2} e^{2\pi i q} \quad (6.29)$$

Modular invariance, as we will see, forces again this lattice to be (odd) self-dual. This does certainly not imply that Γ_{16} and $D_{5,1}$ should separately be self-dual lattices. That would correspond only to the simplest cases: taking in the right-moving sector the odd self-dual lattice $D_{5,1}$ corresponding to the superstring, requires Γ_{16} to be even self-dual. As explained in chapter 3, there exist only two such lattices, namely the root lattice of $E_8 \times E_8$ and the weight lattice of $\frac{Spin(32)}{\mathbb{Z}_2}$. These choices lead to the well-known supersymmetric string theories with gauge groups $E_8 \times E_8$ and $SO(32)$ (the vertex operators for the emission of the corresponding gauge bosons are accordingly given by $V^{ab}(\bar{z}, z) = J^{ab}(\bar{z}) V_{B(-1)}^{vector}$).

Allowing for a non-trivial correlation between conjugacy classes, the two factors in (6.26) need not be separately self-dual; only the combined lattice needs to be. This leads to further possibilities of constructing theories like the non-supersymmetric $SO(16) \times SO(16)$ -string [49] [50]: it is given by a lattice $\Gamma_{16;5,1} = (D_8 D_8)_L (D_{5,1})_R$ equipped with a certain list of correlated conjugacy classes. In this list, *all* conjugacy classes of $D_{5,1}$ of the spinning string do occur; however, the problems of the spinning string do not show up due to the particular, non-trivial interrelation with the left-moving sector. In particular, all branch cuts are cancelled by virtue of (6.28).

The above construction generalizes straightforwardly to the torus-compactified heterotic strings as introduced by [4]. In reducing the number of dimensions, the lattice gets larger as we include also the $2n$ torus-compactified left- and right-moving space-time coordinates. More specifically, in $d = 10 - 2n$ dimensions we need to have an odd self-dual lorentzian lattice

$$\Gamma_{16+2n;5+2n,1} = \Gamma_{16+2n;2n} \times D_{5,1} . \quad (6.30)$$

Here, $\Gamma_{16+2n;2n}$ is an even, lorentzian self-dual lattice as described by [4]. There exists a continuous variety of such lattices: for example, four dimensional theories are characterized by 132 parameters. These are related to background values of the various fields, precisely as in the case of the torus-compactified bosonic string described in chapter 4. As $D_{5,1}$ in (6.30) is odd self-dual, it represents the torus-compactified fermionic degrees of the superstring. Therefore this kind of theories is non-chiral.

It is now clear how to construct more general, possibly chiral theories: we simply have to look out for odd self-dual lattices (6.30), where the two factors are correlated in a non-trivial way. These theories then will not, in general, be describable as compactifications in the very sense. What we have to make sure of, of course, is that we correctly assign even and odd vectors to space-time tensors and spinors; in lower dimensional theories, there are then also additional constraints related to world-sheet supersymmetry in the internal sector (this will be explained below in chapters 7 and 8). The structure of the covariant lattice is more general than that of (6.30):

$$\Gamma_{16+2n;5+2n,1} = (\Gamma_{16+2n})_L \times \left(\Gamma_{3n}^{int} D_{5-n,1}^{spacetime} \right)_R \quad (6.31)$$

Having found this simple structure, we can forget now all subtleties related to the superconformal ghost system.

What remains to be proven is the connection between (multiloop) modular invariance of the partition function and self-duality of the lattice (6.31). Before this will be attacked, let us first explain and switch to an alternative lattice formulation which is easier to deal with.

6.3 THE EVEN LATTICE MAP

Although the *covariant lattices* introduced in the previous section specify the fermionic string theory completely, they are not very pleasant to deal with, because of their rather cumbersome metric, and because some of their vectors have odd norm. To recapitulate, what we will be interested in for string theories in $d = 10$ dimensions are odd self-dual lattices of the form $\Gamma_{16;5,1}$, where the six right-moving components belong to one of the four conjugacy classes of $D_{5,1}$. Furthermore the space-time spinors (i.e. spinor conjugacy classes of $D_{5,1}$) have to correspond to the vectors of odd length.

Now we will introduce a map of any such lattice to an even self-dual lattice. This map will have the useful property that it preserves modular invariance at all loop orders. Essentially, what we are going to do is to associate with any heterotic string theory (or type-II theory) a bosonic string theory, by mapping the right (or left) moving superconformal system to a bosonic one. This map works in general only in this direction.

This idea originates from [51], where it was used to argue that the fermionic strings are “vacua” of the bosonic string. Although this remains an attractive idea, this is not our interest here. For us the lattice map will merely be a useful tool for proving certain properties of fermionic strings. The form of the lattice map that we will use appeared first in [52], and in [44] it was shown that it preserves one-loop modular invariance. This was extended to higher loops in [53].

Basically, the idea is very simple. If we decompose one of the lattices of interest with respect to the $D_{5,1}$ conjugacy classes, we see that it consists of four sets of vectors

$$(\Delta_0, 0) + (\Delta_v, v) + (\Delta_s, s) + (\Delta_c, c), \quad (6.32)$$

where the second entry denotes the $D_{5,1}$ conjugacy classes, and the Δ ’s are simply the 16-dimensional vectors associated with them on the right lattice. Now define a new lattice $\Gamma_{16;8}$ by writing down a decomposition identical to (6.32), but with the second entry interpreted as a conjugacy class of D_8 instead of $D_{5,1}$. That is, we map

$$D_{5,1} = D_4^{light-cone} D_{1,1}^{ghost} \longleftrightarrow D_4^{light-cone} D_4^{ghost} = D_8. \quad (6.33)$$

Observe that the old and the new lattice have the same number of conjugacy classes, so that if the old lattice was unimodular, so is the new one. Now compute the list of norms and inner products of the conjugacy classes of D_8 . In comparison with (6.22) one finds that the only change is in the norm of the spinors, which change from odd into even length. But the spinors of $D_{5,1}$ were precisely the odd vectors on $\Gamma_{16;5,1}$. Hence the map preserves unimodularity and all mutual inner products, but it maps all the odd vectors into even ones. The result is a lattice $\Gamma_{16;8}$ which is even self-dual with respect to the metric $[(-)^{16}; (+)^8]$. Thus, by use of this one-to-one map, we are able to find an alternative *even lattice formulation* of heterotic string theories.

If we extend the lattice with an E_8 factor on the right, we get a $\Gamma_{16;16}$ lattice of that type, on which the bosonic string can be compactified (this can also be achieved by using D_{16} in the above construction, but that is less useful for discussing the spectrum). The

bosonic string compactified on the corresponding torus is all-loop modular invariant. Is this related to the modular invariance of the fermionic string from which it came?

To answer this question, one has to write down the multiloop partition function for the fermionic string, which has not been done yet. However, one usually assumes that as far as its modular transformation properties are concerned, it has the form (including only right-movers)

$$(Det D_{1/2})_\alpha^5 (Det D_{3/2})_\alpha^{-1},$$

where $D_{1/2}$ denotes the Dirac determinant, and represents the contribution of two Majorana Weyl fermions and $D_{3/2}$ is the Rarita-Schwinger determinant (which only has ghost contributions). The subscript α denotes the spin-structure. In general, this combination of determinants is multiplied with some α dependent factors from the right-movers.

The point is now, that the modular transformation properties of this object are known explicitly [54]. Using them, we can show that the following two spin-structure dependent quantities have identical modular transformations at any loop order

$$\begin{aligned} Y_\alpha &= \left(\frac{Det_{1/2}}{Det_{3/2}} \right)_\alpha \\ X_\alpha &= (Det_{1/2})_\alpha^4 \sum_\beta (Det_{1/2})_\beta^8 \end{aligned} \tag{6.34}$$

Thus if one replaces in a fermionic string partition function Y_α by X_α , the new one will be modular invariant if and only if the old one was.

This replacement is precisely what the lattice map achieves. To see that one has to make a simple basis transformation from spin-structure basis to conjugacy class basis. We will not explain this in detail here, but refer to [53]. Details aside, comparing Y_α and X_α the reader will undoubtedly recognize that the first represents the $D_{1,1}$ ghost and longitudinal part of the covariant lattice (cf. (6.20)), while the second represents D_4 conjugacy classes, multiplied with the E_8 partition function.

Let us apply the above to the superstring. As discussed in sect.6.2, the superstring is characterized by a $D_{5,1}$ -lattice with conjugacy classes (0) and (s). By use of (6.33), this can be mapped onto D_8 with the same conjugacy classes, which is the same as E_8 ! Thus, (odd) self-duality of $D_{5,1}^{superstring}$ can be inferred from the (even) self-duality of E_8 . This proves not only multiloop modular invariance of the superstring, but it also shows that (6.23) is the only projection leading to a modular invariant theory, as E_8 is the unique even self-dual lattice in 8 dimensions.

The beauty of the above reasoning lies in the fact that the ghost system is determined by the $N = 1$ conformal system, i.e. by a *world-sheet* property, that is the same for any fermionic string. Hence the replacement of Y by X , or the lattice map, is equally useful for any fermionic string theory, and in any space-time dimension.

So we might as well directly go to four-dimensional strings. As indicated at the end of the previous section, four-dimensional strings are specified by lattices of the form given in the left-hand side of (6.30), but without the restriction that it contains a $D_{5,1}$ root lattice. Since we are in four dimensions, it would be foolish to insist on

ten-dimensional Lorentz-invariance. Instead, we only need to require that $D_{2,1}$ roots be present (the Wick-rotated four-dimensional Lorentz-group is $SO(4)$, with Lie algebra D_2). Thus in $10 - 2n$ dimensions we would be looking for lattices $\Gamma_{16+2n;5+2n,1}$ of the form (6.31), with the requirement that the last $6 - n$ components belong to conjugacy classes of $D_{5-n,1}^{spacetime}$. We will again require that the lattice is odd self-dual, with odd lengths for spinors.

Using precisely the same lattice map, we replace now the $D_{5-n,1}^{spacetime}$ conjugacy classes by $D_{8-n}^{spacetime}$ conjugacy classes, that is

$$\begin{aligned} \Gamma_{16+2n;5+2n,1} &= (\Gamma_{16+2n})_L \left(\Gamma_{3n}^{int} D_{5-n,1}^{spacetime} \right)_R \\ &\longleftrightarrow (\Gamma_{16+2n})_L \left(\Gamma_{3n}^{int} D_{8-n}^{spacetime} \right)_R = \Gamma_{16+2n;8+2n} . \end{aligned} \quad (6.35)$$

Using the above we conclude then that the following statements are equivalent:

- The covariant lattice $\Gamma_{16+2n;5+2n,1}$ is odd self-dual (with odd norms for the spinors, as discussed above).
- The lattice $\Gamma_{16+2n;8+2n} \times E_8$ obtained by means of the lattice map is even self-dual.
- The bosonic string, compactified on the latter lattice is multi-loop modular invariant.
- The fermionic string defined by the odd self-dual covariant lattice is multi-loop modular invariant.

This is all that needs to be said about modular invariance. What corresponds to complicated spin-structure summations in fermionic constructions, is encoded in the simple statement that the lattice must be self-dual.

To some extent, we can drop the covariant lattice altogether, and work with even self-dual lattices. This is unfortunately not possible if one wants to study scattering amplitudes. For this purpose one needs the complete machinery of conformal field theory, and we have not succeeded to cast that into even lattice form. However, the even lattice formulation is extremely useful for discussing the spectrum.

Notice the map (6.35) above is a map between *lattices* (even with different dimensions) and *not* a map between individual vectors. The physical light-cone spectrum is obtained by applying a set of truncation rules to the bosonic string spectrum, which are designed to mimic exactly the corresponding procedure for the covariant, odd lattice. To get these rules, consider the physical state selection rule (6.21). It tell us to keep only states with special components in the $D_{1,1}$ sublattice, belonging to conjugacy classes (v) and (s) of that lattice. Since $D_{1,1}$ is mapped to D_4 by the lattice map, we take the identical conjugacy classes of D_4 to identify the physical states. Thus one physical state is represented by an entire conjugacy class of D_4 . One can choose a representative of the D_4 conjugacy class that makes all this even simpler. Namely, if one chooses some weight vector of 8_s and 8_v to represent the spinor and vector conjugacy class, then the mass of the light-cone states is correctly given by the compactified bosonic string mass formula (4.10), in which one should include the contribution from the D_4^{ghost} components. In

other words, if we make an (arbitrary) choice for the 8_v and 8_s weights, the physical state selection rule (6.21) for covariant lattices is replaced by

$$x = x_0 = \begin{cases} (0, 0, 0, 1) & NS\text{-sector} \\ (\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}) & R\text{-sector} \end{cases} \quad (6.36)$$

in the even lattice formulation ($w = (u, x) \in D_{8-n}^{spacetime}$, $u \in D_{4-n}^{light-cone}$, $x \in D_4^{ghost}$). The physical light-cone spectra are identical.

The above rules were first discovered in [51] and [52], and were a completely *ad hoc* prescription for getting the fermionic string from the bosonic one. Now we see that they do in fact follow from the physical state projection rule explained in sect. 6.1.

7. World-sheet supersymmetry

Up to now we have freely replaced world-sheet fermion by bosons. One has to be a little bit more careful about this, because one may endanger world-sheet supersymmetry. In the *NSR* sector of heterotic strings world-sheet supersymmetry is a manifest (local) $N = 1$ supersymmetry, which transforms the fermions ψ^μ into the bosons X^μ and vice-versa. The transformation is

$$\begin{aligned} \delta_\epsilon \psi^\mu &= i\epsilon \partial X^\mu \\ \delta_\epsilon X^\mu &= -i\epsilon \psi^\mu \end{aligned} \quad (7.1)$$

where ϵ is an infinitesimal grassmannian parameter. This transformation is generated by the operator

$$\mathcal{O} = \oint \frac{dz}{2\pi i} i\epsilon \psi^\mu \partial X_\mu = \oint \frac{dz}{2\pi i} \epsilon T_F, \quad (7.2)$$

in other words, the supersymmetry transformation of any conformal field $\Psi(z)$ is obtained by acting with $\mathcal{O}$

$$\delta_\epsilon \Psi = \mathcal{O} \Psi.$$

One may check that on ψ^μ and X^μ , $\mathcal{O}$ does indeed produce (7.1).

Replacing ψ^μ by bosonic fields does not in itself destroy world-sheet supersymmetry. In two dimensions one can realize supersymmetry entirely in terms of bosons. Indeed, by simply replacing ψ^μ in (7.2) by its bosonized form (5.32) one obtains a generator of world-sheet supersymmetry which is expressed entirely in terms of d bosons X^μ (where d is the number of space-time dimensions) and $\frac{d}{2}$ bosons H_i replacing the fermions.

In constructing four-dimensional strings we do however something more drastic than just bosonization. We allow the bosons related to the internal world-sheet fermions to mix freely with the internal bosons (we will use the word “internal” to indicate those components of ψ and X that do not have space-time indices). Since our philosophy is to construct new string theories directly in four dimensions, we start always with a completely bosonic formulation, i.e. in practice we never bosonize anything. Therefore it is not automatically guaranteed that the result respects world-sheet supersymmetry.

Why do we need world-sheet supersymmetry? One could answer this question by stating that world-sheet supersymmetry is an essential part of superconformal field theory, which is the foundation of fermionic string theory, but there is a more explicit way of seeing what goes wrong if one does not have it.

One thing that goes wrong is Lorentz-invariance. World-sheet supersymmetry is used to remove unphysical degrees of freedom from the covariant action of fermionic strings, so as to write them entirely in terms of the physical, light-cone degrees of freedom. Although one can always write down such a light-cone action, it is important that it came from a covariant action. This makes it possible to write down a Lorentz-algebra which does not just generate the transverse rotations of the light-cone states, but extends to the full Lorentz-algebra. This extension involves the familiar generators L^{i-} , whose commutators have to be checked to see if the algebra closes. In supersymmetric strings one of the terms appearing in the expression for L^{i-} is $\oint \psi^i T_F$.

One of the things that can go wrong with Lorentz invariance is that $[L^{i-}, L^{j-}]$ simply does not yield the answer required by the algebra of the Lorentz group. This situation occurs in theories with a conformal anomaly, and can be avoided by choosing the field content so that the conformal anomaly cancels. This is one of the oldest ways of obtaining the critical dimension.

It might appear that nothing prevents us from simply using the Lorentz algebra of ten-dimensional strings, regarding 6 of the eight 8 transverse Lorentz indices as internal. This algebra closes, and it is indeed a valid choice in torus compactifications. However, closure of the algebra is not enough. In addition, one has to require that L^{i-} takes physical states into physical states. Otherwise one faces a problem which is even worse than the conformal anomaly, namely the states of the theory won't even fit into representations of the Lorentz group. (In the presence of a conformal anomaly, the Lorentz algebra does not close, but for the bosonic string and the *NSR* formulation of the superstring it is still possible to assemble the massive states into representations of $SO(d-1)$, the little group of massive particles)

This is indeed precisely what happens, and one sees this most easily in the chiral sector of chiral string theories. At first sight, chirality and string theory are almost contradictory, because one would think that chiral states can always be excited into massive states. Massive chiral states would however violate Lorentz invariance, so a really clever mechanism is needed to ensure that all massive states combine into non-chiral pairs, while the massless ones do not. This is achieved by world-sheet supersymmetry. Indeed, this restriction on massive chiral states turned out to be powerful enough to determine all the conditions of world-sheet supersymmetry for the class of theories considered in [6].

A second way in which world-sheet supersymmetry enters is through the picture changing operator. This operator has the form $\mathcal{P}_{+1} = e^{i\phi} T_F$, where ϕ is the bosonized ghost of section (6.1). It is essential for the calculation of scattering amplitudes, and it too should take physical states into physical states, and have a well-defined action.

The problem of finding a bosonic realization of world-sheet supersymmetry consists thus of two parts

1. Find a purely bosonic expression for T_F that satisfies the supersymmetry algebra.
2. Make sure that it acts properly on all states.

The algebra that the supercurrent T_F must satisfy can be encoded in operator products, and reads (with finite terms omitted)

$$\begin{aligned} T(z)T_F(w) &= \frac{3/2}{(z-w)^2}T_F(w) + \frac{1}{(z-w)^2}\partial_w T_F(w) \\ T_F(z)T_F(w) &= \frac{10}{(z-w)^3} + \frac{2}{(z-w)}T(w) \end{aligned} \quad (7.3)$$

The first of these equations states that it must have conformal weight $\frac{3}{2}$. The second one is just (5.18), with $c = 15$, the conformal anomaly of 10 bosons and 10 Majorana-Weyl fermions. The form of the supercurrent is further restricted because we want it to be Lorentz invariant in d dimensions. This means that it must have the form

$$T_F = \psi^\mu \partial X_\mu + T_F^{int},$$

where the first term is just (5.17), and the second is some expression of conformal weight $\frac{3}{2}$, constructed out of $3n$ bosons. Here n is defined by $d = 10 - 2n$. The $3n$ bosons consist of $2n$ internal X 's and n bosons H_i representing the ψ 's as in (5.34). From now on, let us denote *all* compact bosons generically by X . Subtracting the space-time part of the supercurrent in (7.3), we see that T_F^{int} has to satisfy it with “10” replaced by $10 - d$, and $T(w)$ by $T(w)^{int}$, the energy momentum tensor of $3n$ bosons.

The first equation is quite easy to satisfy. The general solution has the form

$$T_F^{int} = \sum_{\vec{l}} \vec{B}(\vec{l}) \cdot \partial \vec{X} e^{i\vec{l} \cdot \vec{X}} + \sum_{\vec{t}} A(\vec{t}) e^{i\vec{t} \cdot \vec{X}}, \quad (7.4)$$

where $\vec{l}^2 = 1, \vec{t}^2 = 3$, and $\vec{l} \cdot \vec{B} = 0$. One can verify this by substituting (7.4) into the first equation of (7.3), using the standard energy momentum tensor for $3n$ bosons (see (5.11)).

The second equation in (7.3) imposes more complicated conditions on A and $\vec{B}$, and we do not know its general solution. However, as we will show in a moment, we do know many special solutions. First however we want to address the second point mentioned above, the action on T_F on the states in the theory.

If an operator of the form $\exp(i\vec{w}' \cdot \vec{X})$ acts on a state $|\vec{w}\rangle$ with lattice momentum $\vec{w}$, then it produces a state with momentum $\vec{w} + \vec{w}'$. Earlier in this section we have identified two operators whose action should take physical states into physical states, namely L^{i-} and $\mathcal{P}_{+1}$. If we write the right-moving part of the vectors of the covariant lattice in the form

$$w_R = (\text{internal components}, \text{weight of } D_{d/2} \mid \text{ghost charge}),$$

according to the decomposition

$$\Gamma_{16+2n;5+2n,1} = (\Gamma_{16+2n})_L \times \left(\Gamma_{3n}^{int} D_{5-n,1}^{spacetime} \right)_R, \quad D_{5-n,1}^{spacetime} = D_{d/2}^{lorentz} \Gamma_1^{ghost},$$

then we see that the internal part of the supercurrent contains exponentials of the above

form, with $\vec{w}'$ equal to

$$\begin{aligned} w^{L^{i-}} &= (\vec{l}, \vec{v} \mid 0) \text{ or } (\vec{t}, \vec{v} \mid 0) && \text{for } L^{i-} \\ w^{PC} &= (\vec{l}, 0 \mid -1) \text{ or } (\vec{t}, 0 \mid -1) && \text{for } \mathcal{P}_{+1} \end{aligned}$$

($\vec{v}$ denotes a vector weight of $D_{d/2}$.) If adding a vector $\vec{w}'$ to a lattice vector should give another lattice vector, then $\vec{w}'$ should lie on the lattice itself. For the picture changing operator one should furthermore require that its action on all states is *local*, i.e. does not introduce any branch cuts. In other words, if $e^{i\vec{w}'\cdot\vec{X}}$ is the vertex operator for some physical state with lattice momentum $\vec{w}$, then we should require that the operator product

$$e^{i\vec{w}'\cdot\vec{X}} e^{i\vec{w}\cdot\vec{X}} = (z - w)^{\vec{w}'\cdot\vec{w}} e^{i(\vec{w}'+\vec{w})\cdot\vec{X}} + \dots$$

does not contain any fractional powers of $(z - w)$. This implies that $\vec{w}'$ must have integral dot product with all lattice vectors, so that it must lie on the *dual* of the lattice. Fortunately our lattices are self-dual, so that these requirements on $\vec{w}'$ are compatible.

The requirements one gets from L^{i-} and $\mathcal{P}_{+1}$ are also compatible with each other, because in covariant lattices the $D_{d/2}$ components and the ghost charge belong to the same conjugacy class of $D_{d/2,1}$. What we conclude from this is that for every vector $\vec{l}$ and $\vec{t}$ appearing in T_F^{int} the lattice must contain vectors of the form $w_R = (\vec{l}, (v))$ and $(\vec{t}, (v))$, where (v) denotes generically the vector conjugacy class of $D_{d/2,1}$. This describes of course only the right-moving part of the lattice. The left components w_L of these vectors, which we will call *constraint vectors* in the following, must vanish.

There is yet another way to arrive at this conclusion, which we mention here because it uses more directly the requirement of supersymmetry, and can also be used to identify other world-sheet supersymmetries on the lattice. What these conditions are imposing on the theory is that the supersymmetry transformation not only respect the *local* properties of the fields, but also, on Riemann surfaces with non-trivial topology, their *global* properties. In (7.1) that would amount to requiring that both sides of the equation have the same periodicity. In the superstring, for which one only has (7.1), this is simply achieved by choosing the boundary conditions for ϵ equal to those of the *NSR* fermions on any Riemann surface with given spin structure. In bosonic language choosing a spin-structure is replaced by choosing a *soliton sector*, which simply means to consider vacuum with a certain lattice momentum vector $\vec{w}$. States in this sector which are created by vertex operators of the form $\exp(i\vec{w}'\cdot\vec{X})$ pick up a phase $\exp(2\pi i\vec{w}'\cdot\vec{w})$ if one takes σ_1 to $\sigma_1 + \pi$. When one considers one term of the supercurrent acting on it, then such a state is mapped into a state which picks up a different phase when moved around a closed loop. This phase change is $\exp(i\vec{p}\cdot\vec{w} + 2\pi i\phi(\vec{w}))$, where we have denoted the phase of ϵ in the sector $\vec{w}$ as $\phi(\vec{w})$. The vector $\vec{p}$ appearing here represents just one term in the supercurrent (i.e. it is one of the vectors $\vec{l}$ or $\vec{t}$ used above). If there were just one term, one can always adapt $\phi(\vec{w})$ to cancel the phase, so that supersymmetry respects all boundary conditions. If the supercurrent has a second term with a vector $\vec{p}'$, then this cancellation of the phase by $\phi(\vec{w})$ can work if and only if $\vec{p} - \vec{p}'$ has integer dot-product with any $\vec{w}$ on the lattice. But then it must lie on the lattice itself.

This argument tells us thus that the *difference* of any two vectors appearing in the supercurrent must lie on the lattice. This includes also the space-time part of the supercurrent. For that part the vector $\vec{p}$ belongs to the vector conjugacy class of $D_{d/2}$, so that we arrive at the same conclusions mentioned earlier.

There are two simple ways of obtaining supercurrents that satisfy the second equation of (7.3). The first is to construct one out of free fermions, and bosonizing it. Such a free fermion supercurrent has the form

$$T_F(z) = \frac{1}{6} f_{abc} \psi^a \psi^b \psi^c, \quad (7.5)$$

where the ψ 's are $6n$ fermions related to the $3n$ internal bosons by bosonization. If f_{abc} are the structure constants of a semi-simple Lie algebra, then one can check that (7.3) is satisfied [55]. The generic choice for this Lie algebra (which has to be of dimension $6n$) is $SU(2)^{2n}$. In dimensions below five there are additional solutions, which are of some interest for type-II strings, but not for chiral heterotic strings. To obtain a supercurrent of the form (7.4) one simply bosonizes the $6n$ fermions into $3n$ bosons using (5.32). There is some freedom in doing that, because one may associate the fermions in different ways with the $6n$ $SU(2)$ generators, but there is one especially simple way which yields only norm 3 vectors in the supercurrent. This is to take

$$T_F = N \sum_{i=1}^n [\psi_i^1 \psi_i^2 \psi_i^3 + \psi_i^{-1} \psi_i^{-2} \psi_i^{-3}]$$

where the notation is as in (5.32). We leave it as an exercise to determine the normalization constant N . The vectors $\vec{t}$ one gets from this supercurrent are, for $n = 1$,

$$\vec{t} = (\pm 1, \pm 1, \pm 1), \quad (7.6)$$

For larger n one just adds sets of 8 such vectors in orthogonal three-dimensional subspaces, so that altogether one has $8n$ vectors. This supercurrent was first used by Kawai et. al., and lead to their “triplet constraint”.

Although one might think that all supercurrents that satisfy (7.3) can be written as a trilinear expression of free fermions, this is not the case. A counterexample, found in [56] and [57] has vectors $\vec{t}$ of the form (for $n=1$)

$$\begin{aligned} \vec{t}_1 &= (\sqrt{3}, 0, 0) \\ \vec{t}_2 &= (0, \sqrt{3}, 0) \\ \vec{t}_3 &= (0, 0, \sqrt{3}) \end{aligned} \quad (7.7)$$

This pattern can be repeated n times if $n > 1$. By bosonizing the free fermionic supercurrent, one can only get vectors of the form $(\pm 1, \pm 1, \pm 1, 0, \dots, 0)$, i.e., there must exist a basis in which three entries are ± 1 , and the others zero. For one norm 3 vectors one can always achieve that, but not necessarily simultaneously for several vectors. In particular it is not possible for (7.7).

The mistaken impression that any supercurrent can be written in terms of free fermion trilinears, is based on the proof of [58] that the Fock space of any torus-compactified theory can always be written in terms of a fermionic representation. This may involve either fermions with Thirring interactions (to which the above does not apply, since one would not have a free fermion supercurrent), or even free fermions with complicated boundary conditions. Now of course a bosonic theory described in this fermionic way will still have world-sheet supersymmetry, since it is equivalent. But one cannot conclude from this argument that there must always exist a representation of the supercurrent of the trilinear form (7.5).

An illustrative example is a string theory on an E_8 lattice. The Fock space can be described either by means of fermionic oscillators with certain GSO projections, or with bosonic oscillators and lattice momenta. The Frenkel-Kac generators of the D_8 subalgebra of E_8 can also be described both bosonically and fermionically, the latter being fermion bilinears. But whereas the bosonic description extends trivially to the remaining spinor roots of $E_8/SO(16)$, the fermionic one does not. To describe these generators one would need a *transcendental* expression in terms of the fermions rather than a bilinear one (see the discussion in section (5.4)). We believe that the same applies to the supercurrent, with “bilinear” replaced by “trilinear”.

Finally we mention a second method for generating supercurrents, which owes its inspiration to asymmetric orbifolds. In this case one starts by taking the ten-dimensional form also in the internal space, i.e.

$$T_F^{int} = \sum_{I=1}^{2n} \psi^I \partial X_I.$$

The fermions can be bosonized trivially. Although the rest of the supercurrent is already bosonic, we can try to re-express the factors ∂X as sums of exponentials of the form $\exp(i\vec{r}\cdot\vec{X})$, where $\vec{r}^2 = 2$. These expressions are constrained by the fact that they must satisfy the same operator products as the ∂X ’s. An equivalent way of saying this is that they must be the generators of a $U(1)^{2n}$ algebra. One can find such subalgebras by rotating the ∂X ’s into the root system of a Lie algebra. One way of finding such solutions is to use the Weyl rotations of a Lie algebra, and apply the conjugation that maps twists into shifts. We will not discuss this in more detail here; for details we refer to [59]. The solutions already given above can be obtained in this way (this has been shown in [57]), as well as many others, listed in [59].

8. The Spectrum

To discuss the spectrum of covariant lattice theories we will use the even lattice formulation. It may be useful to summarize the requirements that we have derived in the foregoing chapters. To specify a covariant lattice theory in $d = 10 - 2n$ dimensions one must provide

- A lattice $\Gamma_{16+2n;8+2n}$ with left- and right dimensions as indicated
- A supercurrent that satisfies the $N = 1$ supersymmetry algebra, and is consistent with the lattice.

The supercurrent has the general form

$$T_F = \psi^\mu \partial X_\mu + T_F^{int}$$

where the internal part appears only below 10 dimensions. The fermions ψ^μ can be bosonized, and the momenta of the resulting bosons occupy $\frac{d}{2} = 5 - n$ of the $8 + 2n$ dimensions of the right part of the lattice. The internal part of the supercurrent is entirely bosonic, and has the form

$$T_F^{int} = \sum_{\vec{t}} A(\vec{t}) e^{i\vec{t} \cdot \vec{X}} + \sum_{\vec{l}} \vec{B}(\vec{l}) \cdot \partial \vec{X} e^{i\vec{l} \cdot \vec{X}},$$

where $\vec{t}^2 = 3$, $\vec{l}^2 = 1$ and $\vec{l} \cdot \vec{B} = 0$, and the coefficients satisfy the conditions discussed in the previous chapter. The momenta of the bosons $\vec{X}$ occupy $3n$ dimensions of the right part of the lattice. The remaining 3 components of the right lattice are related, in a somewhat indirect way, to the β, γ ghost system, as explained in chapter 6.

The lattice must satisfy the following requirements.

1. It must be even and self-dual with respect to a Lorentzian metric of the form $\text{diag}((-)^{16+2n}; (+)^{8+2n})$
2. The last $8 - n$ components of any vector must belong to one of the four conjugacy classes $(0), (v), (s)$ or (c) of the Lie algebra $D_{8-n}^{spacetime}$.
3. It must contain all vectors of the form $\vec{w}_L = 0, \vec{w}_R = (\vec{l}, (v))$ and $(\vec{t}, (v))$ for all $\vec{l}$ and $\vec{t}$ appearing in the supercurrent.

The last condition applies obviously only to dimensions less than 10.

As explained before, not all the states that one obtains from such a lattice are physical. The physical states are obtained by making a decomposition of D_{8-n} to its subalgebra $D_{4-n} \times D_4$. One keeps only states of which the last four components are fundamental vector or spinor weights of D_4 , and counts them with multiplicity one.

This decomposition of $D_{8-n}^{spacetime} \rightarrow D_{4-n}^{light-cone} \times D_4^{ghost}$ is very simple. The branching rules are as follows:

$$\begin{aligned} (0) &\rightarrow (0, 0) + (v, v) \\ (v) &\rightarrow (v, 0) + (0, v) \\ (s) &\rightarrow (c, c) + (s, s) \\ (c) &\rightarrow (s, c) + (c, s) \end{aligned}$$

We have written them in such a way that the second term is the one that survives the

physical state projection (6.36), because it has as its second ($= D_4$) entry (v) or (s). Of course we could have chosen the other spinor weight of D_4 as the one that gives physical states, but this merely amounts to a chirality flip of the entire theory.

Considering now the first entry, we can read off the spin of the particles in each conjugacy class of D_{8-n} . The first entry corresponds to D_{4-n} , which is the transverse (Wick-rotated) Lorentz group in $10 - 2n$ dimensions. We see that the following general rule holds for D_{8-n} conjugacy classes

$$\begin{aligned} (0) &\rightarrow \text{odd rank tensors} \\ (v) &\rightarrow \text{even rank tensors} \\ (s) &\rightarrow \text{spinors of chirality } + \\ (c) &\rightarrow \text{spinors of chirality } - \end{aligned} \tag{8.1}$$

The assignment of positive or negative chirality to the spinors is a matter of convention.

This list gives us the Lorentz representations of the pure lattice states in light-cone gauge. These are states that have a certain lattice momentum, but no bosonic oscillator excitations. The complete Fock space of physical states is obtained as follows

$$(\alpha\text{-oscillators})(\tilde{\alpha}\text{-oscillators})|\vec{w}_L; \vec{w}_R > |_{m_L=m_R}.$$

Here α denotes all 24 left-moving oscillators, of which $d - 2$ have a vector index of the transverse Lorentz group and will be denoted α^i . The remaining, “internal” ones will be denoted $\alpha^I, I = 1, \dots, 26 - d$.

For the $\tilde{\alpha}$ oscillators things are a bit more complicated. One has $d - 2 = 8 - 2n$ right-moving oscillators $\tilde{\alpha}^i$ in the vector representation of the transverse Lorentz group, which are simply the right-moving modes of the uncompactified coordinates. Then there are the oscillators belonging to the lattice degrees of freedom. Four of the $8 + 2n$ right-moving lattice dimensions are associated with ghosts and longitudinal modes, and the corresponding four bosonic oscillators should not be included; the remaining $2n + 4$ oscillators ($3n$ internal ones and $4 - n$ in the Cartan subalgebra of $D_{4-n}^{\text{light-cone}}$) should be allowed to act on the pure lattice states. They do not alter the Lorentz representation of the state on which they act. Obviously the $d - 2$ left and right-moving bosonic oscillators α^i and $\tilde{\alpha}^i$ do change the Lorentz representation. Thus, as we already pointed out, (8.1) is valid only for pure lattice states.

The last fact one has to know about the spectrum is the mass formula. This is extremely simple: It is just the mass formula of the compactified bosonic string:

$$\begin{aligned} \frac{1}{8}m_L^2 &= \frac{1}{2}w_L^2 + \mathcal{N} - 1 \\ \frac{1}{8}m_R^2 &= \frac{1}{2}w_R^2 + \tilde{\mathcal{N}} - 1 \end{aligned} \tag{8.2}$$

Here $(\vec{w}_L; \vec{w}_R)$ denotes a vector on $\Gamma_{16+2n; 8+2n}$. Notice that the entire vector contributes to the mass formula, including the D_4 components. Since every physical state necessarily is a fundamental vector or spinor of this D_4 , the lowest possible value of $\frac{1}{8}m_R^2$ is $-\frac{1}{2}$.

It follows that if we are only interested in massless states and tachyons, then we can ignore the right-moving oscillators, since they would increase the mass by too much. In the following we will use $M_R(M_L)$ to denote $\frac{1}{8}m_R^2$ ($\frac{1}{8}m_L^2$), and we will refer to this quantity simply as “mass”. Thus we will quote masses in units in which the bosonic string tachyon has mass -1.

8.1 TEN-DIMENSIONAL STRINGS

Now we have all the ingredients necessary to determine the spectrum. Since the ten-dimensional case is somewhat simpler, we will consider it first. The lowest states one can have in the four conjugacy classes of D_8 are discussed below.

The conjugacy class (0)

The shortest vectors in this conjugacy class are $\vec{w}_R = 0$ and $\vec{w}_R$ equal to a root of D_8 . The first cannot satisfy the physical state criterion, so that the physical state with the lowest value of m_R is obtained by taking $\vec{w}_R$ to be a root. Its mass is equal to zero, and according to (8.1) it yields a Lorentz vector. To satisfy the condition $M_L = M_R$ we have to pair it with a massless state from the left sector. There are two ways to achieve that: either by taking $\vec{w}_L^2 = 2$ or by taking $\vec{w}_L = 0$, and acting with one of the 24 bosonic oscillators. Eight of these oscillators are in the vector representation of the transverse Lorentz group $SO(8)$. Its tensor product with the right-moving sector yields the $SO(8)$ representations $(35) \oplus (28) \oplus (1)$, corresponding respectively to the graviton, the antisymmetric tensor B_{ij} , and a dilaton. The remaining 16 bosonic oscillators yield upon multiplication with the right sector 16 vector bosons. These turn out to be gauge bosons of a $U(1)^{16}$ gauge group. If there are any vectors $\vec{w}_L$ of norm 2 that combine on the right lattice with the (0) conjugacy class of D_8 , then they will give rise to additional gauge bosons, extending $U(1)^{16}$ to a non-abelian gauge group of rank 16.

Note that if the lattice contains vectors of the form $(\vec{r}; (0))$, with $\vec{r}^2 = 2$, then it contains in particular vectors of the form $(\vec{r}; 0)$. We will call such vectors, and also vectors of the form $(0; \vec{r})$ roots of the lorentzian lattice. Notice that all the physical states we have this discussed so far owe their existence to roots: the graviton and the 16 commuting gauge bosons to the roots of D_8 , and the remaining gauge bosons to roots of the left lattice. The graviton (and also the 16 abelian gauge bosons) are inevitable, since the D_8 roots are always present.

It can be shown that the roots of the left (right) part of the lattice must always form a root system of some simply laced Lie algebra. In the subspace spanned by these root systems, all lattice vectors must have components that have integer dot products with the root lattice of this Lie algebra. Hence in this subspace they must be weights.

If the roots of the left lattice span an l -dimensional subspace of the 16 left-moving dimensions, then the gauge bosons belong to a non-abelian rank l group, times an abelian group $U(1)^{16-l}$. We will denote this entire group as G_L . All other states in the theory belong to certain representations of this group.

The conjugacy class (v)

The lowest state that can appear here is the vector representation of D_8 . If it appears, there is a potential problem, because for this lattice vector $M_R = -\frac{1}{2}$. Because

the lattice is even, we know that $\vec{w}_L^2 = 1 \bmod 2$. Should there exist a lattice vector with both $\vec{w}_R^2 = 1$ and $\vec{w}_L^2 = 1$ then the theory would have a tachyon. It might happen that the lowest value of $\vec{w}_L^2$ is in fact 3 rather than 1, and indeed there is one example of that, the $O(16) \times O(16)$ string (see the section 9.1 for a discussion). Furthermore there are lattices on which the conjugacy class (v) is simply not present. This is the case for the supersymmetric strings (see below).

The conjugacy classes (s) and (c)

The lowest possible states are the D_8 spinors, which have $M_R = 0$. As was the case for the conjugacy class (0) , they can become physical either by pairing with a left-moving oscillator excitation or with a norm two vector. If the first happens, then in particular it can combine with an oscillator α_i . The left-right tensor product contains then a vector-spinor, usually called a gravitino.

If a gravitino is present, then the lattice state excited by the left-moving oscillators must have $\vec{w}_L = 0$, combined with a D_8 spinor. This lattice vector is a root of the *right* lattice, and by the arguments given above, it follows that the right lattice must have a larger root system than that of D_8 . It does not require a lot of thought to realize what it must be, namely E_8 . The presence of a gravitino implies that one must obtain supergravity in space-time. Again we see that a local symmetry of the theory is related to a root of the lattice.

Once one realizes that the spectrum has a gravitino, one would clearly expect it to be supersymmetric. For example, one would expect to find gaugino's, the superpartners of the gauge bosons. These are easy to identify: one can excite vectors with right-moving components in the (s) conjugacy class of D_8 in precisely the same way as those in the (0) conjugacy class discussed above. A little bit of thought will make it clear that these are the only massless spinors one can get; in ten-dimensional superstrings all massless fermions are either singlets or adjoints of the gauge group.

Of course one can also have non-supersymmetric strings. In that case, massless fermions can only arise from vectors of the form $(\vec{w}_L; (s))$, with $\vec{w}_L^2 = 2$. It should be emphasized that $\vec{w}_L$ cannot be equal to a root of the right lattice. If it were, one could subtract $(\vec{w}_L; 0)$ from the spinor, and obtain a gravitino by exciting the result with a bosonic oscillator. Thus in non-supersymmetric string theories in ten dimensions one cannot have spinors in the adjoint of the gauge group.

Up to now we have not distinguished the (s) and the (c) conjugacy class of D_8 . They give spinors with opposite space-time chirality. Whether the theory is chiral depends on the representations (i.e. the weights) with which the two conjugacy classes combine on the left lattice. If these are always the same, the theory is not chiral. Otherwise it is.

8.2 FOUR-DIMENSIONAL STRINGS

After this rather lengthy discussion of ten-dimensional strings, let us now see what changes if one goes to lower dimensions. The main difference is that now there can be additional contributions to M_R from the components of $\vec{w}_R$ which extend beyond the space-time lattice D_{8-n} . For definiteness (and for phenomenological reasons) we will consider here four dimensions. The discussion of 6 and 8 dimensions is not very different, the main difficulty being the mod-8 properties of spinors in different dimensions.

In four dimensions, the lattice is $\Gamma_{22;14}$, and the right part contains a D_5 space-time lattice. Again we will consider each of the conjugacy classes of D_5 .

The conjugacy class (0)

Here the discussion can be very brief, because there are almost no changes with respect to the ten-dimensional case. Roots of D_5 again give zero mass gravitons and gauge bosons, this time of course of a rank 22 gauge group. The 9 “internal” components of the right lattice play no role in the massless spectrum, as they would excite the mass above zero.

The conjugacy class (v)

As in ten dimensions, the lowest possible state here is a tachyon of mass $-\frac{1}{2}$. It originates from any lattice vectors of the form $(\vec{w}_L; 0, (v))$, $\vec{w}_L^2 = 1$, where the middle component refers to the nine internal dimensions of the lattice, according to the decomposition $\Gamma_{22;14} = (\Gamma_{22})_L \times \left(\Gamma_9^{int} D_5^{spacetime} \right)_R$. Clearly we have now the possibility to excite these nine components. In other words, the lattice may contain vectors of the form

$$(\vec{w}_L; \vec{u}_R, (v)), \quad \vec{w}_L^2 = 1 + \vec{u}_R^2.$$

This state would have mass $M_L = M_R = \frac{1}{2}\vec{u}_R^2 - \frac{1}{2}$. The possibilities that one has for $\vec{u}_R$ are severely restricted by the constraint vectors of the supercurrent. Since the lattice is self-dual, they must have integer dot-product with all the vectors $\vec{t}$ and $\vec{l}$ appearing in (7.4). For example, for the supercurrent (7.6) it is easy to see that the norm of $\vec{u}_R$ must be a multiple of $\frac{1}{4}$, so that in addition to the tachyon at mass $-\frac{1}{2}$, one may (depending on the lattice one chooses, of course) find tachyons of mass $-\frac{1}{4}$.

None of this is very important though, because one would normally not consider such tachyonic theories. More interesting is the possibility of having a vector $\vec{u}_R$ of norm 1. (Of course it is possible that the lattice has such vectors, without containing tachyonic vectors) In that case one obtains a state with $M_L = M_R = 0$, which is a massless scalar in a representation of G_L with weight vector $\vec{w}_L$, of norm 2.

This raises of course the possibility that $\vec{w}_L$ is actually a root of the left lattice. One can then go through an interesting chain of arguments. If $\vec{w}_L$ is a root (or, in other words, if one has a massless scalar in the adjoint representation of G_L), then one can subtract a root from $(\vec{w}_L; \vec{u}_R, (v))$, to obtain $(0; \vec{u}_R, (v))$. This is a root of the right part of the lattice, which has non-vanishing dot-product with the roots of D_5 . Again it is not hard to see that the lattice must contain a larger rootsystem than that of D_5 , of which the D_5 roots form a subset. This means that D_5 must be a *regular subalgebra*

of some larger Lie algebra. There are only a few possibilities: This algebra must be E_6, E_7, E_8 , or $D_{5+n}, n \geq 1$. Since the decomposition of the larger algebra must yield a vector representation of D_5 , it can only be D_{5+n} . We will see in a moment that this destroys any possibility of getting chiral fermions out of such a theory, but let us first consider another consequence of this extension of D_5 .

Of course we can excite the state corresponding to the extra roots by a right-moving oscillator α_i . This gives a massless space-time vector, and it should therefore be a gauge boson. That is indeed what it turns out to be. We will not go into the details of its interactions here, but we will just mention that its gauge symmetries originate from the right lattice, and involve the supercurrent in a non-trivial way. In fact, the gauge group turns out to be a non-regular subgroup of the Frenkel-Kac group of the right lattice. This group is of little relevance in heterotic strings, since it inhibits chirality, but it is more important in type-II strings; see [20], [21].

In chiral heterotic strings there is thus never a gauge group corresponding to the right-movers (this seems to have been proved more generally by Friedan and Shenker). As a consequence, the vectors $\vec{u}_R$ on the internal part of the right lattice only affect the *multiplicities* of the particles in the spectrum, not the representation they are in. There is no direct way of “measuring” $\vec{u}_R$, because there are no gauge symmetries coupling to it. Nevertheless these vectors do affect the theory in less direct ways, because the Yukawa couplings and Higgs self interactions depend on them.

The existence of this seemingly redundant nine-dimensional internal space is noteworthy for phenomenological reasons, because it makes it possible to have particles in the theory which have identical properties, but differ only in this part of the lattice. We need not point out that this might help explain the existence of fermion generations. Furthermore there can be interesting continuous and/or discrete *global* symmetries originating from this internal space.

The spinor conjugacy classes

Since a spinor of D_5 has norm $\frac{5}{4}$, it would appear that one could get tachyonic spinors of mass $-\frac{3}{8}$. The existence of tachyons in non-singlet representation of the Lorentz group signals an instability of the vacuum which would cause a breakdown of the Lorentz group. One might expect that world-sheet supersymmetry should prevent that. This does indeed appear to be the case. Observe that the D_5 spinor has half-integral dot-product with the constraint vectors of world sheet supersymmetry, because those vectors have (v) components in D_5 . If a spinor were not to have any components in the internal dimensions, this would be in disagreement with the fact that the lattice is self-dual. Therefore, any D_5 spinor must appear in a vector of the form $(\vec{w}_L; \vec{u}_R, (s))$, and $\vec{u}_R$ must have half-integral dot-product with all the vectors $\vec{l}$ or $\vec{t}$ in the supercurrent. In all cases we have examined it turns out that to satisfy this condition, $\vec{u}_R$ must have norm greater than $\frac{3}{4}$. Furthermore there seems always to exist a vector $\vec{u}_R$ which saturates this lower limit. The reader may find it instructive to verify this for the two supercurrents (7.6) and (7.7) which we discussed in the previous section. Although it ought to be possible to prove this without explicitly writing down the supercurrent, for the moment we will accept as a fact that fermionic tachyons do not appear in the spectrum.

results as the classification of supersymmetries can be obtained from the classic work of Cartan!

It is not difficult to extend this to other dimensions, and for the dimensions above four one gets then^{*}

$d = 4$	D_5	$\subset$	E_6	$N = 1$
			E_7	$N = 2$
			E_8	$N = 4$
$d = 6$	D_6	$\subset$	E_7	$N = 1$
			E_8	$N = 2$
$d = 8$	D_7	$\subset$	E_8	$N = 1$
$d = 10$	D_8	$\subset$	E_8	$N = 1$

(Some people seem to use different conventions for counting the number of supersymmetries, in particular in six dimensions. To compare your favorite convention with ours use the torus compactifications of the $N = 1$ theory in 10 dimensions; these correspond to E_8 in the table). For even more intriguing results in two dimensions we refer to [6].

It seems that everything that’s known about supersymmetry and supermultiplets can be found back in the representation theory of the exceptional groups. For example, consider $E_6 \supset D_5$ in four dimensions. The smallest representations of E_6 are the singlet (1), the (27) and the (78), which is the adjoint. Their D_5 content is

$$\begin{aligned} (1) &\rightarrow (1) \\ (27) &\rightarrow (10) \oplus (16) \oplus (1) \\ (78) &\rightarrow (45) \oplus (16) \oplus (\overline{16}) \oplus (1) \end{aligned}$$

According to our rules, the D_5 singlets contain no physical states, while the (10) yields a scalar and the (16) a fermion. Thus the (27) contains the scalar multiplet. (On the lattice the representation (27), which has weights of norm $\frac{4}{3}$, must appear in combination with an internal space vector $\vec{u}_R$). In a similar way one can see that (78) contains the vector multiplet. Note that to count the spinors we should consider only the (16), because the $(\overline{16})$ is just a CPT conjugate. Note also that for every (27) the lattice necessarily contains a CPT conjugate $(\overline{27})$.

In fact, one can push this further, as it is possible to relate the “unphysical” left-over singlet components to the auxiliary fields of $N = 1$ supergravity. Vertex operators for such auxiliary fields have recently been investigated [60]. It is conceivable that the above procedure works also for extended supergravities and/or higher dimensions, as well as for massive multiplets with arbitrary high spins, by using the representation theory of the appropriate exceptional group. We encourage the industrious student to work out the details.

^{*} This relation between space-time supersymmetry and exceptional groups is in fact completely general, and applies for instance also to Calabi-Yau compactified string theories. The reason is that space-time supersymmetry constrains the vertex operator algebra so that it can always be mapped to an exceptional algebra.

We have not said much about massive states, but something needs to be said about them in supersymmetric theories. One would expect that they too belong to supermultiplets. This can in fact be proved by using the triality properties in a somewhat unexpected way. These properties are known to be related to supersymmetry in ten dimensions, and to clarify the relation between the Neveu-Schwarz-Ramond and the Green-Schwarz formalism. In those arguments, one uses triality rotations in the transverse Lorentz group. Clearly that is of no use below ten dimensions, since one has a different Lorentz group. Instead we will show that one can use the D_4 that in our formulation describes the ghosts and longitudinal modes for this purpose. This part of the lattice is dimension independent, so that what follows holds in arbitrary dimensions.

The D_4 factor plays a role in distinguishing fermions from bosons, because the former have as their D_4 components weights of the spinor representation (8_s), whereas the latter have weights of the vector representation (8_v). These vectors can be mapped into each other by an $SO(4)$ rotation which maps all the entire D_4 weight lattice onto itself. This is called an automorphism of the lattice. If such an automorphism maps weights of a given representation into weights of *the same* representation it is called an *inner* automorphism. Clearly, if one has a Lie algebra lattice (i.e. a sublattice of the complete weight lattice, containing at least the root lattice), then all inner automorphisms are of course symmetries of it. There are also automorphisms that map weights of *different* representations to each other. Such automorphisms are called *outer*. In particular, D_4 has an outer automorphism which cyclically permutes (8_s), (8_c) and (8_v), known as triality. Clearly an arbitrary D_4 lattice (take for example $(0) + (s)$) is not invariant under such an automorphism.

It is a special property of the embedding of D_4 in the exceptional groups that this outer automorphism becomes an inner automorphism of the exceptional group. This means in particular that for every (8_v) of D_4 contained in some representation of an exceptional group, there is also an (8_s) and (8_c). For the spectrum that statement translates into having a fermion for every boson, and hence into supersymmetry at all levels (at least as far as multiplicities are concerned).

9. Model building

There are many ways of finding solutions to the conditions listed in the beginning of the previous section, but unfortunately none is systematic enough to achieve a complete classification. Only in ten dimensions the complete list of solutions is known (by this we mean the complete list of solutions that can be obtained with the covariant lattice construction, and which therefore have a rank 16 gauge group. In ten dimensions one knows just one additional solution, which has an E_8 gauge group and is tachyonic [49], [61]). We will consider this case first.

About the massless fermions one can say roughly the same as about the ten-dimensional case. The right-moving lattice vectors describing them must have norm 2. Because the lattice is lorentzian even, they appear in combination with even norm vectors $\vec{w}_L$ on the left. Two possibilities are of interest: $\vec{w}_L^2 = 0$ and $\vec{w}_L^2 = 2$.

In the first case one can excite the right-moving Fock space by means of bosonic oscillators α_i to obtain a gravitino. One can also excite it with an internal oscillator α_I or add a root of the left lattice (if it has any), to obtain a gaugino. The appearance of the gravitino tells us that the theory has a certain number of supersymmetries (more about this below). Again we see that the existence of local-space time symmetries is linked to the existence of certain roots of the right lattice.

In the second case, we may assume that $\vec{w}_L$ is not a root of the left lattice, for if were we could subtract $\vec{w}_L$ and obtain the case already discussed. If $\vec{w}_L$ is not a root, it is some vector weight of norm 2 of G_L , and one gets a massless spinor in the corresponding representation. If G_L has $U(1)$ factors part of $\vec{w}_L$ may correspond to $U(1)$ -charges, and the remainder is a weight of the non-abelian part of G_L . In any case the only representations of the non-abelian part of G_L that one can get are those with weights of norm less than or equal to two. Fortunately the representations of the standard model satisfy this bound.

Nothing we have said so far addresses the question of chirality. The situation is a bit different here in comparison with ten dimensions, and this has simply to do with the difference in CPT transformations of spinors in these dimensions. One can easily understand that there is difference by changing the sign of a spinor in the conjugacy class (s) of D_8 or D_5 : in the first case one gets a spinor in the same conjugacy class, whereas in the second case one obtains a spinor in (c). What does this have to with CPT? In four dimensions, a CPT transformation send a left-handed spinor in some representation (R) of some gauge group into a right-handed anti-spinor in the complex conjugate representation ($\bar{R}$). Remarkably, there is an automorphism of every lattice that achieves precisely the same, namely multiplying all lattice vectors with -1. As discussed above, this flips the chirality of the D_5 spinor conjugacy classes, and hence the space-time chirality, while at the same time changing the signs of all vectors on the left lattice. As is well-known, changing the sign of a weight changes it to a weight of the complex conjugate representation.

Thus first of all CPT is a symmetry of any theory we construct. Secondly, to get the usual counting of four-dimensional fermions, we should consider only one D_5 chirality, (s), say, and regard states from the other conjugacy class as anti-particles. Thus if the lattice contains three vectors $(\vec{w}; \vec{u}_R^l, (s)), l = 1, \dots, 3$, where $\vec{w}$ is a weight of a complex representation (R), we have three generations of fermions in the representation (R). The lattice will automatically also contain $(-\vec{w}; -\vec{u}_R^l, (c))$, but these are just the CPT conjugates of the three generations, and are not to be counted separately.

With this understanding, it is obvious how to count chiral fermions. If there are N generations of fermions in (R) and $\bar{N}$ in the complex conjugate ($\bar{R}$), then there are $N - \bar{N}$ chiral generations. This way of counting them assumes implicitly that pairs of opposite chirality will become massive, because there is no symmetry that would forbid such a mass term. We know however of no mechanism in string perturbation

theory that would actually achieve this. Most likely, the massless spectrum is not altered to all orders in perturbation theory, and even the non-chiral fermions will then simply stay massless. It is however quite clear that non-perturbative effects have a role to play in obtaining low-energy phenomenology, and in particular they have to break supersymmetry and/or the weak interaction gauge symmetry. Since we know nothing about non-perturbative effects, we can take the path of maximal optimism, and assume that they will do whatever we need them to do.

There does not exist a simple criterion for determining *a priori* which lattices can give chiral fermions, and which cannot. There is however a simple necessary condition, namely that the D_5 root system should not be part of the root system of a larger D_n group. If that were the case, all other D_5 representations originate from D_n representations. For the spinors that means that they always come in pairs of opposite chirality, because a D_n ($n > 5$) spinor (s) contains both (s) and (c) of D_5 . These spinors belong to the same representation of any gauge group, and hence they are chiral conjugates (More precisely, if one applies a CPT conjugation to the spinors in the (c) conjugacy class of D_5 , one can easily see that in the notation used above one would have $N = \bar{N}$). This loss of chirality is a well-known phenomenon in torus compactifications of higher dimensional chiral theories, and that is in fact precisely what is happening here: If the D_5 lattice extends to D_n one can regard the theory as a torus compactification of a higher dimensional one.

There are two situations in which such an extension of D_5 is forced on us. First of all, it occurs when we want to have gauge bosons belonging to G_R , as discussed above. Secondly, if the supercurrent contains any vectors $\vec{l}$ of length 1, the corresponding constraint vectors are roots, extending D_5 . Thus to get chiral fermions, one needs $\vec{B}(\vec{l}) = 0$.

This concludes the discussion of the spectrum of covariant lattice theories. We emphasize once more a feature of this discussion which we think is rather elegant, and is unique to the even lattice formulation: All local symmetries (gravity, gauge symmetries of G_L and G_R and supergravity) are related to roots on the lattice. The only massless states not originating from roots are the scalars and the fermions (other than gravitino's and gaugino's).

8.3 SPACE-TIME SUPERSYMMETRY

Space-time supersymmetry is a subject which merits a separate discussion. As we have seen, gravitino's originate from roots on the lattice which are spinors of D_5 . For reasons that should by now be familiar, this implies that D_5 is regular subalgebra of a larger Lie algebra. It is now quite clear that there is a restriction on the number of supersymmetries one can have, because there are not that many Lie algebras that extend D_5 by spinor roots; the only ones are E_6, E_7 and E_8 . By simply counting the number of gravitino's one gets, we deduce that these three possibilities correspond respectively to $N = 1, N = 2$ and $N = 4$ supergravity. Similarly, for type-II theories one can get this from both the left and the right-moving sector, yielding $N = 1, 2, 3, 4, 5, 6$ and 8 as the possible supersymmetries. We find it very intriguing that such relatively recent

9.1 TEN-DIMENSIONAL STRINGS

To obtain ten-dimensional strings we are looking for lattices $\Gamma_{16;8}$, where the right-moving components are D_8 weights. To find them, we can make use of a trick explained in section (3.4). Given a lorentzian even self-dual lattice of the required form, we can make an even self-dual euclidean lattice out of it by simply changing the sign of the last eight components of the metric. This seemingly absurd operation works only because the right lattice factor is known to be D_8 ; certainly changing the sign of the metric on an arbitrary lorentzian even self-dual does not preserve the self-duality. The result of this map is an even self-dual Euclidean lattice of dimension 24. But such lattices have been classified by Niemeier, so that classifying all ten-dimensional strings is reduced to merely finding the right mathematical literature (which in some libraries is still no small obstacle).

Not every Niemeier lattice yields a good string theory. To go back to a lorentzian lattice one has to be able to find a D_8 subalgebra. Obviously this must be a regular subalgebra, because we want to have a D_8 sublattice, not just a subalgebra. This is not difficult, and yields the following solutions

$$\begin{aligned}
 E_8 \times E_8 \times E_8 &\rightarrow E_8 \times E_8 \\
 D_{16} \times E_8 &\rightarrow SO(32) \\
 D_8 \times D_8 \times D_8 &\rightarrow SO(16) \times SO(16) \\
 D_{24} &\rightarrow SO(32) \\
 D_{12} \times D_{12} &\rightarrow SO(8) \times SO(24) \\
 D_{16} \times E_8 &\rightarrow SO(16) \times E_8 \\
 D_{10} \times E_7 \times E_7 &\rightarrow SO(4) \times E_7 \times E_7 \\
 D_9 \times A_{15} &\rightarrow SU(16) \times U(1)
 \end{aligned}$$

Here we have indicated on the left the Niemeier lattice, and on the right the gauge group it produces. Notice that there is one Niemeier lattice, $D_{16} \times E_8$, that produces two different string theories, because there are two ways of embedding D_8 .

The first two theories are supersymmetric, because $D_8 \subset E_8$. The last five are tachyonic, because D_8 is embedded in a D_n factor with $n > 8$. When the Niemeier lattice is mapped to $\Gamma_{16;8}$, this embedding leaves a trace in the form of lattice vectors $((v), (v))$, belonging to a $(D_{n-8})_L \times (D_8)_R$ sublattice originating from the D_n factor. Such vectors lead to tachyons. Finally, the third theory is the rather well-known $SO(16) \times SO(16)$ string, which does not have supersymmetry, and also no tachyons. When it was found it was the first of its kind, but in four dimensions such models are very common.

It is a useful exercise to determine the spectrum of some of these models, using the conjugacy classes of the Niemeier lattices. The answers can be found in [62].

9.2 FOUR-DIMENSIONAL STRINGS

In constructing lattices for four-dimensional strings we face a new technical difficulty, namely the constraint vectors for the supercurrent. Consider for example the choice (7.6). The first thing one should observe is that the form of this supercurrent forces all vectors in the nine internal dimensions to have as their components integers or half-integers. This is true because by addition of constraint vectors one gets, among others, vectors of the form

$$(0, \dots, 0, 2, 0, \dots, 0)$$

in the nine-dimensional internal space, and zero elsewhere. The structure of the lattice in this nine-dimensional space can then be summarized in a nice way by saying that all its vectors belong to conjugacy classes of $(D_1)^9$, where D_1 is the lattice introduced at the end of section (3.3).

Now the entire right lattice is of the form $D_5 \times (D_1)^9$, and it is not unreasonable to assume that also the left lattice can be written as a product of D_n factors. Since even under this assumption the lattices will turn out to be too numerous to be classified, whether one misses any solutions in this way is an academic question (our guess is that one does not). Any D_n factor on the left can always be decomposed into a product of D_1 factors, so one could try to enumerate all the even self-dual lattices based on

$$[(D_1)^{22}]_L [(D_1)^9 D_5]_R \quad (9.1)$$

conjugacy classes. One can imagine a *gedanken* computer program that does this. It would start with a lattice containing the conjugacy classes of the constraint vectors

$$\begin{aligned} &[(0)^{22}, (v), (v), (v), (0), (0), (0), (0), (0), (0), (v)] \\ &[(0)^{22}, (0), (0), (0), (v), (v), (v), (0), (0), (0), (v)] \\ &[(0)^{22}, (0), (0), (0), (0), (0), (0), (v), (v), (v), (v)], \end{aligned}$$

and then it would go through the list of conjugacy classes of (9.1) to find one of even lorentzian length, and with even inner product with all the vectors that are already on the the lattice. This procedure can be repeated to find a next conjugacy class, and when no new conjugacy class can be found, the lattice that one has is self-dual by definition.

Although writing this program is quite trivial, one is discouraged from running it because the total number of conjugacy classes of (9.1) is gigantic: 4^{32} . As explained in section (3.3), this means that a self-dual lattice based on (9.1) contains 2^{32} conjugacy classes, so that even checking them for tachyons, say, would cost an enormous amount of time.

Obviously this approach is quite useless in practice, but it does give us some insight in the extent of the problem. This can be made more precise by comparing it to a similar combinatorial problem, that of constructing all Euclidean even self-dual lattices of dimension 32. One could do that in an analogous way, namely by starting with $(D_1)^{32}$ (in 24 dimensions this approach would indeed give all 24 Niemeier lattices). As a combinatorial problem this is not too different from the one we were facing, because the

number of conjugacy classes is the same^{*}. But for this problem we know a (conservative) lower limit of the number of solutions, namely 8×10^7 (In [6] we gave an often misquoted rigorous upper limit on the number of theories, namely the number of even self-dual lattices of dimension 88. This number is finite, but larger than 10^{1500} . This was only intended to show that the number of solutions is finite, and does not in any way represent the actual number of solutions). This shows that we can expect to find a very large number of four-dimensional string theories for this choice of supercurrent.

A different, and more effective method for constructing lattices (for this supercurrent) was found by the authors of [63], who observed that they can all be obtained from *odd* self-dual lattices of dimension 22. By using techniques of the kind described in section (3.4), one can glue to this lattice a right-moving sector which contains the supercurrent constraint vectors. However from one odd self-dual lattice one gets many covariant lattices $\Gamma_{22,14}$, and a complete classification along these lines appears quite difficult. Of course the situation is improved considerably by imposing additional requirements, such as space-time supersymmetry.

Knowing the extent of the problem makes model building a somewhat arbitrary exercise. In fact “model building” is not even the correct terminology; “model finding” would be more appropriate. The objective is to *find* in the gigantic set of solutions one that corresponds to the standard model. It does not help to be just close to the standard model, since in string theory (ideally at least) one cannot adjust the model by adding a few higgses and tune a few parameters or by doing some of the other things model builders like to do.

This is a rather extreme view of the present situation, and in practice one can of course learn a lot from semi-realistic examples, and we will give some below. They are only intended as an illustration of the construction, and should not be confused with phenomenology. In [6] we produced many examples by modifying some of the Niemeier lattices. Here we will follow a perhaps more pedagogical route, the shift vector method. Any lattice of the form (9.1) can be transformed into any other one by adding a suitable set of shift vectors constructed out of D_1 “weights”. In many cases these D_1 conjugacy classes contain roots, which form the root system of some simply laced Lie algebra. It is then convenient to write the the lattice in terms of the corresponding Lie algebra lattice to reduce the list of conjugacy classes. The simplest roots appear in conjugacy classes of the form $(0, \dots, 0, v, v, 0, \dots, 0)$, which combine two D_1 factors into a D_2 . However, it is certainly not true that from (9.1) one can only get D_n gauge groups. For example, the Niemeier lattices, which can have A_n or E_n factors, can be written as D_1^{24} .

A useful starting point is the lattice $(D_{22})_L \times (D_{14})_R$. To the root lattice one has to add four conjugacy classes to obtain a unimodular lattice, and if one furthermore requires it to be even, there is not much choice: The conjugacy classes must be $(0, 0) + (v, v) + (s, s) + (c, c)$ (this is just an Englert-Neveu lattice with the left factor stretched by 8 dimensions). The phenomenology is extremely simple: The first term gives a graviton (plus $B_{\mu\nu}$ and a dilaton), and the gauge bosons of $SO(44)$ and the second

^{*} The fact that the constraint vectors must be on the lattice reduces the number of solutions, but the fact that not all 32 factors are on equal footing in (9.1) leads to an enhancement.

one gives a tachyon in the vector representation of $SO(44)$. The spinors are all massive and non-chiral, so that this example is certainly not of much interest. Note that the D_{14} lattice on the right contains the constraint vectors of world-sheet supersymmetry. In fact there are many possible sets of constraint vectors that one could choose. If one adds shift vectors, one has to make sure that they have integral dot product with one set of constraint vectors, so that these vectors are on the invariant sublattice Λ_0 defined in section (3.4).

Let us now make the model supersymmetric. This is easily achieved by adding a gravitino as a shift vector. To do so, split D_{14} into $D_6 \times D_8$ and add the conjugacy class $(0, 0, s)$ of $(D_{22})_L \times (D_6 \times D_8)_R$, where we embed the D_5 space-time lattice into D_8 (in the following we adopt the convention of embedding the space-time lattice in the last factor). Notice that two shift vectors in the same conjugacy class have the same effect, so that we can specify the shift vector by giving its conjugacy class with respect to the sublattice.

Using the shift vector method explained in section (3.4) we can easily determine the conjugacy classes that appear. Since we have three D_n factors, there are in fact $2^3 = 8$ of them, and they are

$$(0, 0, 0) + (v, v, 0) + (s, s, s) + (c, c, s) + (0, 0, s) + (v, v, s) + (s, s, 0) + (c, c, 0)$$

What we have now obtained is a ten-dimensional heterotic string, compactified on a lattice $\Gamma_{22;6} = D_{22} \times D_6$. The D_8 factor of the covariant lattice has been embedded in E_8 because of the addition of the gravitino. This example has in fact more supersymmetry than we bargained for, namely $N = 4$. It is of course not chiral.

This illustrates several general principles. First of all, any theory can be made supersymmetric by simply adding as a shift vector a lattice vector which produces a gravitino in the spectrum. More generally, one can enforce the appearance of any state one likes by simply adding the corresponding lattice vector as a shift vector. One may however get unwanted states in addition. Another noteworthy aspect is the use of spinor weights to break D_n factors into smaller ones.

There are two obvious obstacles to chirality in our example. The first is that the D_5 root system is part of a larger D_n root system (that of D_8 in this case), which always inhibits chirality as explained in the previous section. The second is that the $SO(44)$ gauge group simply doesn't have any complex representations, which is necessary for chirality in four dimensions. The only $SO(N)$ groups with complex representations are $SO(4k + 2)$, $k \in \mathbb{Z}$. Their chiral representations are the spinors, and the fundamental spinor representations have weight length $\frac{2k+1}{4}$. Clearly only the fundamental spinors of $SO(2)$, $SO(6)$, $SO(10)$ and $SO(14)$ can occur for massless fermions, since all others would have to high a mass. Furthermore a fermion can not be chiral due to spinors of $SO(6)$, because the $SO(6)$ spinor representations are anomalous, and modular invariant string theory is anomaly free. (Note that $SO(2)$ anomalies can be cancelled by the four-dimensional analog of the Green-Schwarz mechanism [18], [64].)

These chiral representations can only appear in combinations with total weight length equal to two. This still leaves many possibilities, and one of the simplest is the

$SO(10) \times SO(6)$ representation $(16, 4) + (16, \bar{4})$. The necessity to cancel the $SO(6)$ anomaly implies that the number of generations is a multiple of eight in this case.

Breaking $SO(44)$ into such small groups cannot be achieved by adding just one spinor shift vector, but two turn out to be sufficient. First decompose

$$(D_{22})_L(D_6 D_8)_R \rightarrow (D_{14} D_8)_L(D_4 D_2 D_2 D_6)_R$$

by adding a shift vector $(0, s; 0, s, 0, c)$. This shift vector projects out half of the gravitino's, so that we get an $N = 2$ model with an $SO(16) \times SO(28)$ gauge group. It is still not chiral, but we can now break the lattice root system further down to

$$(D_7 D_7 D_5 D_3)_L(D_2 D_2 D_2 D_1 D_1 D_1 D_5)_R$$

by adding the shift vector $(s, 0, 0, s; s, 0, c, s, 0, 0, c)$. This again removes one gravitino, so that we have $N = 1$ supersymmetry. The remaining lattice vector giving a gravitino is $(0, 0, 0, 0; 0, 0, 0, s, s, s, s)$. The constraint vectors appear in the conjugacy classes

$$\begin{aligned} &(0, 0, 0, 0; v, 0, v, v, 0, 0, v) \\ &(0, 0, 0, 0; 0, v, v, 0, v, 0, v) \\ &(0, 0, 0, 0; v, v, 0, 0, 0, v, v) \end{aligned}$$

To check if these vectors are on the lattice, it suffices to verify that they have integer dot product with the shift vectors we used.

Working out the spectrum takes some effort, since the lattice has 11 factors, and hence 2^{11} conjugacy classes. Most of them contribute only to massive states, and with some ingenuity one can determine the massless states by hand. However, this kind of labor is easy to computerize, which makes it much easier to experiment with other lattice constructions. The model we just constructed turns out to be chiral, and its complete fermion spectrum consists of the $SO(14) \times SO(14) \times SO(10) \times SO(6)$ representations

$$\begin{aligned} &2(1, 1, 16, 4) + 2(1, 1, 16, \bar{4}) \\ &+(14, 14, 1, 1) + (1, 1, 10, 6) \\ &+(14, 1, 10, 1) + (1, 14, 10, 1) \\ &+(14, 1, 1, 6) + (1, 14, 1, 6) \end{aligned}$$

Only the fermions in the first line are chiral, and give 16 generations of (16) 's of $SO(10)$. These fermions originate from the lattice vectors $(0, 0, s, s, 0, 0, c, 0, 0, s, c)$ and $(0, 0, s, c, 0, 0, s, 0, 0, s, c)$. Notice that the $(D_2)^3(D_1)^3$ factor of the right lattice only determines the multiplicities of the states in the theory, not their gauge quantum numbers. Since the model is supersymmetric, there are scalars in the same representations as the fermions.

This model is a well-known one, and appears in most of the papers on four-dimensional strings because it is one of the simplest to obtain. The construction given here closely resembles the one used in the fermionic constructions, and we have presented it in this way to allow the reader to make a comparison. Indeed, there is a one-to-one correspondence between the class of models based on (9.1) and fermionic models with “complex” boundary conditions, and (anti)-periodic boundary conditions.

One can continue from here, and add more shift vectors to slice the lattice into still smaller D_n chunks. It should be clear from the above that the reason for the rather large number of generations is the fact that their multiplicity is due to a spinor of $SO(6)$ and one of $SO(4)$. Thus the key to lowering the number of generations is to break these factors. This works indeed as expected, but we don't think it would be illuminating to show the gory details of such models. The model given above is just the tip of a gigantic iceberg. If one splits the lattice into more and more factors, there are more and more possibilities for doing this. Clearly the bulk of the models correspond to lattices which are split as much as possible. Unfortunately this “base of the iceberg” is the hardest part to explore, because of the huge number of conjugacy classes one has to deal with.

To conclude this chapter we will briefly discuss models for another choice of the supercurrent, namely (7.7). Just as our previous choice enforced a $(D_1)^9$ structure on the right lattice, the choice (7.7) restricts it as well. World-sheet supersymmetry requires that the lattice contain nine vectors of the form $(0, \dots, 0, \sqrt{3}, 0, \dots, 0, (v))$, where the last entry refers to D_5 , and the others to the remaining nine dimensions of the right lattice. Any other lattice vector in the conjugacy classes (0) or (v) of D_5 must have entries that are a multiple of $\frac{1}{3}\sqrt{3}$ in these nine dimensions; if it is a spinor of D_5 its entries in these nine dimensions must be an odd multiple of $\frac{1}{6}\sqrt{3}$. Vectors of the form $(0, \dots, 0, 2\sqrt{3}, 0, \dots, 0, (0))$ must always be on the lattice, because they are equal to twice the constraint vectors.

This suggests to define a one-dimensional “root”-lattice

$$\mathcal{U} = \{2k\sqrt{3} \mid k \in \mathbb{Z}\}$$

and its dual, or “weight”-lattice

$$\mathcal{U}^* = \{\frac{k}{6}\sqrt{3} \mid k \in \mathbb{Z}\}.$$

This lattice $\mathcal{U}^*$ contains $\mathcal{U}$ as a sublattice, and has 12 cosets (which, by analogy with lie-algebra lattices we shall call “conjugacy classes”) with respect to it. What we have just learned can then be rephrased as follows: the right lattice must be of the form $\mathcal{U}^9 \times D_5$. This notation must be interpreted in the usual sense, i.e. in addition to the root-lattice a certain set of conjugacy classes can appear.

The volume of the unit cell of the root-lattice $\mathcal{U}^9 \times D_5$ is $2(2\sqrt{3})^9$. To have a chance to get a self-dual lattice, we have to combine it a root-lattice on the left which completes this volume to an integer. This is easy: for example A_2, E_6 have a root lattice of volume $\sqrt{3}$, and $\mathcal{U}$ itself has volume $2\sqrt{3}$. Valid choices for the left lattice are for example $\mathcal{U} \times D_{21}$, or $A_2 \times D_{20}$, or $(A_2)^4 \times E_6 \times E_8$, or $\mathcal{U}^{21} \times D_1$. These are only a few possibilities from an obviously long list, and it should be clear that now one can once again start to construct models as we did above. This class is known to contain the Z_3 asymmetric orbifolds [8], [57]. We leave it as an exercise to find their conjugacy classes in this basis (the answer is essentially in [8], although in a different basis). They can then be used as a starting point for finding new theories by the shift vector method, yielding yet another, presumably very large set of theories. And this is still nowhere near the complete set, because there are many other supercurrents one can choose.

10. Final remarks

Despite the considerable length of this paper, there are still many things that we would have liked to discuss in some detail. Here we will merely list some of them with some brief comments and speculations.

- *Interactions* were not discussed in great detail, because this is not necessary with our construction. Since the entire method is based on conformal field theory, the formalism for calculating correlation functions on any Riemann surfaces is already built in. We don't expect these theories to be less consistent than the ten-dimensional ones. Explicit calculations of tree and one-loop diagrams have been done in [65], [64].
- *Overlaps* with other approaches are only partly understood. By construction, covariant lattice theories in four dimensions have gauge groups of rank 22. To the best of our knowledge other constructions yielding rank 22 gauge groups can be re-expressed in terms of covariant lattices. This has been shown for fermionic construction with complex fermion boundary conditions [5], and for asymmetric orbifolds with twists in the Weyl-group of a Lie-algebra [66],[59]. Furthermore the set of covariant lattice theories seems larger than that of fermionic constructions (of rank 22) with a trilinear supercurrent. The restriction to rank 22 can be overcome by considering
- *Twisted Covariant Lattices*, a new class of four-dimensional string theories, obtained by applying orbifold twists to covariant lattices. Since covariant lattices include Narain lattices, twisted covariant lattices include asymmetric orbifolds. The question is, does this class include anything more? Since one is twisting a much larger class of lattices, there is certainly reason to speculate that it might. The most interesting twists would be those that map bosonized world-sheet fermions into world-sheet scalars, something that fits quite naturally within our philosophy, and does not seem to be included in orbifold constructions. Obviously such twists will be severely restricted by modular invariance and world-sheet supersymmetry, but as far as we can tell there is no reason why these obstacles should be insurmountable.
- *The selection of the vacuum* has not been discussed, because we know of no principle that might distinguish one supersymmetric four-dimensional string from any other. Non-perturbative effects are usually invoked as a *deus ex machina* that will somehow achieve this, but this remains wishful thinking at present. A related question is whether four-dimensional strings are
- *Different vacua or different theories?* Probably regarding them as different vacua of the same, generic heterotic string theory is the best way of thinking about them, since after all different four-dimensional strings are nothing but the same conformal field theory with different boundary conditions. It is however not clear to us exactly in what sense one can regard all these theories as ground states of some space-time theory. Thinking about this question, one is automatically led to the question whether there are

- *Continuous parameters* connecting the theories considered here to other ones. One set of obvious candidates, namely Lorentz-rotations of the lattice, does not yield continuous parameters because in chiral theories they would ruin the constraints imposed on the right lattice by (among others) world-sheet supersymmetry. There could however very well be continuous parameters in the direction of lower rank theories.

Finally, there are many phenomenological questions that we wished we could answer. It would be very interesting to have an example that can approach the standards set in [67] for Calabi-Yau compactification, but the gigantic number of possibilities doesn't make the prospect of searching for one look very inviting. Probably the time is not yet ripe for genuine predictions from string theory, and it might be better to try to gain more insight into the vacuum structure first. In view of the extent of that problem, patience will be required.

ACKNOWLEDGEMENTS

We have benefited from discussions with many people, and are grateful for their help in improving our understanding of the subjects discussed here. We can't name them all without taking the risk of forgetting some, but we make an exception for Dieter Lüst, Bengt Nilsson and Nick Warner, whom we thank for the pleasant collaborations.

REFERENCES

- [1]. D. Gross, J. Harvey, E. Martinec and R. Rohm, Nucl. Phys. B256 (1985) 253; Nucl. Phys. B267 (1986) 75.
- [2]. P. Candelas, G. Horowitz, A. Strominger and E. Witten, Nucl. Phys. B258 (1985) 46.
- [3]. L. Dixon, J. Harvey, C. Vafa and E. Witten, Nucl. Phys. B261 (1985) 620; Nucl. Phys. B274 (1986) 285; M. Mueller and E. Witten, Phys. Lett. 182B (1986) 28; L. Ibañez, H. Nilles and F. Quevedo, Phys. Lett. 187B (1987) 25.
- [4]. K. Narain, Phys. Lett. 169B (1986) 41.
- [5]. H. Kawai, D. Lewellen and S. Tye, Nucl. Phys. B288 (1987) 1.
- [6]. W. Lerche, D. Lüst and A.N. Schellekens, Nucl. Phys. B287 (1987) 477.
- [7]. I. Antoniadis, C. Bachas and C. Kounnas, Nucl. Phys. B289 (1987) 87.
- [8]. K. Narain, M. Sarmadi and C. Vafa, Nucl. Phys. B288 (1987) 551.
- [9]. D. Gepner, Nucl. Phys. B287 (1987) 111; *Exactly Solvable String Compactifications on Manifolds of $SU(N)$ Holonomy*, Princeton preprint PUPT-1066.
- [10]. K. Bardacki, E. Rabinovici and B. Säring, *String models with $c < 1$ components*, CERN-TH.4760/87 (1987).
- [11]. C.S. Lam, *GSO Projections of modular invariant aperiodic strings*, Princeton preprint IASSNS-HEP-87/12 (1987); A. Chamseddine and J.-P. Derendinger, *Twisted superstrings in four dimensions* Zürich preprint ETH PT 87-2.

- [12]. J. Distler and B.R. Greene, *Aspects of (2,0) Compactifications*, Harvard preprint HUTP-87/A065
- [13]. J.H. Schwarz, *Review of Recent Developments in Superstring Theory*, CALTECH preprint CALT-68-1432; L. Dixon, *Some World-Sheet Properties of Superstring Compactifications on Orbifolds and Otherwise*, Princeton preprint PUPT-1074.
- [14]. C.A. Lütken, *A Large Class of Calabi-Yau Vacua*, CERN-TH.4900/87.
- [15]. A. Cohn, D. Friedan, Z. Qiu and S. Shenker, Nucl. Phys. B278 (1986) 577.
- [16]. W. Lerche and D. Lüst, Phys. Lett. 187B (1987) 45.
- [17]. M. B. Green, J. H. Schwarz and E. Witten, *Superstring Theory*, Cambridge University Press, (1987); J. H. Schwarz, Phys. Rep. 89 (1982) 223.
- [18]. M. Green and J. Schwarz, Phys. Lett. 149B (194) 117.
- [19]. A.N. Schellekens and N.P. Warner, Phys. Lett. 177B (1986) 317; Phys. Lett. 181B (1986) 339; Nucl. Phys. B287 (1987) 87.
- [20]. L. Castellani, R. d'Auria, F. Gliozzi and G. Sciuto, Phys. Lett. 168B (1986) 47; R. Bluhm, L. Dolan and P. Goddard, Nucl. Phys. B289 (1987) 364.
- [21]. H. Kawai, D. Lewellen and S. Tye, Phys. Lett. 191B (1987) 63; W. Lerche, B.E.W. Nilsson and A.N. Schellekens, Nucl. Phys. B294 (1987) 136; L. Dixon, V. Kaplunowsky and C. Vafa, Nucl. Phys. B294 (1987) 43.
- [22]. A. Polyakov, Phys. Lett. 103B (1981), 207; 211.
- [23]. L. Alvarez-Gaumé, G. Moore and C. Vafa, Comm. Math. Phys. 106 (1986) 1.
- [24]. J. Conway and M. Sloane, Journal of Number Theory 15 (1982) 83; Europ. J. Combinatorics 3 (1982) 219.
- [25]. H. Niemeier, Journal of Number Theory 5 (1973) 142.
- [26]. J.-P. Serre, *A Course in Arithmetic*, Springer Verlag, New York, (1973).
- [27]. P. Goddard and D. Olive, in *Vertex Operators in Mathematics and Physics*, Springer Verlag, (1984); Int. J. Mod. Phys. A (1986) 303.
- [28]. R. Slansky, Phys. Rep. 79 (1981) 1.
- [29]. F. Englert and A. Neveu, Phys. Lett. 163B (1985) 349.
- [30]. J. Leech, Canadian Journal of Mathematics 19 (1967) 251.
- [31]. A.N. Schellekens, contribution to *Perspectives in string theory*, Copenhagen (1987), CERN-preprint, to appear.
- [32]. C. Siegel, Ann. Math. 36 (1935) 527.
- [33]. K. Narain, Sarmadi, E. Witten Nucl. Phys. B279 (1986) 93.
- [34]. J. Polchinsky Comm. Math. Phys. 104 (1986) 37.
- [35]. L. Alvarez-Gaumé, *Supersymmetry and Index theory*, Bonn lectures (1984).
- [36]. P. Ginsparg and C. Vafa Nucl. Phys. B289 (1987) 414.
- [37]. Th. Ohmrodorf, Nucl. Phys. B281 (1987) 470.
- [38]. A. Belavin, A. Polyakov and A. Zamolodchikov, Nucl. Phys. B241 (1984) 333.

- [39]. D. Friedan, E. Martinec and S. Shenker, Nucl. Phys. B271 (1986) 93.
- [40]. D. Friedan, in *Unified String Theories* (Proc. Santa Barbara workshop 1985), M. Green and D. Gross, eds. , World Scientific (1986).
- [41]. S. Shenker in *Unified String Theories* (Proc. Santa Barbara workshop 1985), M. Green and D. Gross, eds. , World Scientific (1986).
- [42]. E. Martinec, University of Chicago preprint EFI 87-45, Lectures given at the 1987 spring school on Superstrings, Trieste.
- [43]. M. Peskin, SLAC-PUB 4251, lectures presented at the 3th Advanced Study Institute in elementary particle physics, Santa Cruz (1986).
- [44]. H. Nicolai and A.N. Schellekens, in *Proceedings of the 5th Adriatic Meeting on Superstrings, Anomalies and Unification*, Dubrovnik 1986.
- [45]. A. Kosteletzky, O. Lechtenfeld, W. Lerche, S. Samuel and S. Watamura, Nucl. Phys. B238 (1987) 173.
- [46]. P. Goddard, D. Olive and A. Schwimmer Phys. Lett. 157B (1985) 393.
- [47]. P. Ramond, Phys. Rev D3 (1971) 2415; A. Neveu and J. H. Schwarz Nucl. Phys. B31 (1971).
- [48]. F. Gliozzi, J. Scherk and D. Olive, Nucl. Phys. B122 (1977) 253
- [49]. L. Dixon and J. Harvey, Nucl. Phys. B274 (1986) 93.
- [50]. L. Alvarez-Gaumé, P. Ginsparg, G. Moore and C. Vafa, Phys. Lett. 171B (1986) 155.
- [51]. A. Casher, F. Englert, H. Nicolai and A. Taormina, Phys. Lett. 162B (1985) 121.
- [52]. F. Englert, H. Nicolai and A.N. Schellekens, Nucl. Phys. B274 (1986) 315.
- [53]. A.N. Schellekens, *Multiloop Modular Invariance of the Covariant Lattice Construction of Fermionic Strings*, CERN-TH/4834-87, to appear in Phys. Lett. B.
- [54]. E. Verlinde and H. Verlinde, Nucl. Phys. B288 (1987) 357.
- [55]. P. di Vecchia, V. G. Knizhnik, J. L. Petersen and P. Rossi, Nucl. Phys. B253 (1985) 701; P. Goddard and D. Olive, Nucl. Phys. B257 (1985) 245; I. Antoniadis, C. Bachas, C. Kounnas and P. Windey, Phys. Lett. 171B (1986) 51;
- [56]. W. Lerche, B.E.W. Nilsson and A.N. Schellekens, Nucl. Phys. B294 (1987) 136.
- [57]. H. Kawai, D. Lewellen and S. Tye, *Construction of four-dimensional fermionic strings with a modified supercurrent*, Cornell preprint CLNS 87/69.
- [58]. J. Bagger, D. Nemeschansky, N. Seiberg and S. Yankielowicz, Nucl. Phys. B289 (1987) 349.
- [59]. A.N. Schellekens and N.P. Warner, *Weyl groups, Supercurrents and Covariant Lattices*, CERN-TH.4916/87 (1987).
- [60]. J. Atick, L. Dixon, and A. Sen Nucl. Phys. B292 (1987) 109; M. Dine, I. Ichinose and N. Seiberg, Nucl. Phys. B293 (1987) 253; S. Cecotti, S. Ferrara and L. Girardello, *Gravitational Supertvertices in N=1, 4D-Superstrings*, CERN-TH.4802/87 (1987).

- [61]. H. Kawai, D. Lewellen and S. Tye, Phys. Rev. D34 (1986) 3794; D. Benett, M. Brene L. Mizrachi and H. Nielsen, Phys. Lett. 187B (1986) 179.
- [62]. W. Lerche, D. Lüst and A.N. Schellekens, Phys. Lett. 181B (1986) 71.
- [63]. J. Balog, P. Forgacs, Z. Horvath and P. Vecsernyés, Phys. Lett. 197B (1987) 395.
- [64]. W. Lerche, B.E.W. Nilsson, and A.N. Schellekens Nucl. Phys. B289 (1987) 609;
W. Lerche, B.E.W. Nilsson, A.N. Schellekens and N.P. Warner, *Anomaly cancelling terms from the elliptic genus*, CERN-TH.4765/87 (1987).
- [65]. D. Lüst, S. Theisen and G. Zoupanos, *Four-dimensional heterotic strings and conformal field theory* preprint MPI PAE PTh 83-87 (1987).
- [66]. D. Lüst and S. Theisen, *Four dimensional strings - Orbifolds and Covariant Lattices* preprint MPI-PAE-PTh83/87 (1987).
- [67]. B. Green, K.H. Kirklin, P.J. Miron and C.G. Ross Nucl. Phys. B278 (1986) 667.

THE COVARIANT LATTICE CONSTRUCTION OF FOUR-DIMENSIONAL STRINGS^{*}

W. Lerche[†]

and

A. N. Schellekens

CERN, 1211 Geneva 23, Switzerland

ABSTRACT

An elementary introduction is given to the covariant lattice construction of lower dimensional heterotic strings, and to the concepts on which it is based. This includes a review of properties of lattices, torus compactification and conformal field theory. Some more advanced topics, such as the bosonization of the superghost system, are also reviewed.

^{*} Based on lectures by the second author at the Summer School on Strings and Superstrings, Poiana Braşov, Romania, September 1987.

[†] Supported by the Max-Planck-Gesellschaft, Munich, FRG.
Address after 1st September 1988: California Institute of Technology, Pasadena, CA 91125, USA.

THE COVARIANT LATTICE CONSTRUCTION OF FOUR-DIMENSIONAL STRINGS^{*}

W. Lerche[†]

and

A. N. Schellekens

CERN, 1211 Geneva 23, Switzerland

ABSTRACT

An elementary introduction is given to the covariant lattice construction of lower dimensional heterotic strings, and to the concepts on which it is based. This includes a review of properties of lattices, torus compactification and conformal field theory. Some more advanced topics, such as the bosonization of the superghost system, are also reviewed.

^{*} Based on lectures by the second author at the Summer School on Strings and Superstrings, Poiana Braşov, Romania, September 1987.

[†] Supported by the Max-Planck-Gesellschaft, Munich, FRG.

Address after 1st September 1988: California Institute of Technology, Pasadena, CA 91125, USA.

THE COVARIANT LATTICE CONSTRUCTION OF FOUR-DIMENSIONAL STRINGS

W. LERCHE AND A.N. SCHELLEKENS

CERN, 1211 Geneva 23, Switzerland

1. Introduction

The main goal of string theory is an ambitious one: to obtain a consistent theory of all interactions, including gravity. Just a few years ago it seemed that gravity was all it could describe, but that has changed drastically with the discovery of the heterotic string [1]. Although at first it seemed that the theory would fail its first confrontation with experiment by predicting the wrong space-time dimension, that too changed very quickly after its birth. Indeed, what once may have seemed an insurmountable obstacle is now a gaping hole: not only can we construct string theories in four dimensions, but it is also so easy that we can make them in embarrassing abundance. Meanwhile the miracles that started the excitement have survived, although we are now so used to them that we tend to forget them: chiral string theories in four dimensions exist and their spectra have the same general features (gauge groups, representations and family replication) as that of the standard model. If all this is realized in a class of theories that holds the promise of offering a finite and consistent theory of quantum gravity, then there is certainly reason for excitement. The dream of having a *unique* theory of everything must be shelved for the moment, but fortunately uniqueness is the aspect that we can most easily live without. On the other hand, giving up consistency would be fatal.

The floodgates of four-dimensional string construction were first opened in [2]. The initial trickle of Calabi-Yau compactifications quickly grew into a raging torrent, which broadened into orbifold compactifications [3]. The theories that are now usually called four-dimensional strings are mathematically not very different from these early ideas, and it doesn't make much sense to try to make a distinction. If there is a difference, then it is one of philosophy rather than physics. Whereas initially one tried to compactify ten-dimensional strings, nowadays the emphasis is more on constructing string theories directly in four dimensions. We believe that the idea of compactification is too limited, and too much linked to notions of space-time that may be unsuitable for string theory, to be of lasting value. But all this is just a matter of interpretation, and does not influence the facts, such as the spectrum and the S-matrix.

The change of philosophy from compactification to four-dimensional strings started with the seminal paper of Narain [4], and gained momentum due to the work of Kawai. et. al. Soon thereafter many ways of constructing heterotic strings in four dimensions were discovered [5-8], the oldest being fermionic constructions, the covariant lattice construction, and asymmetric orbifolds, which are now understood to be close relatives.

Several other ideas have surfaced more recently [9-12]. For recent reviews of aspects of string vacua see e.g. [13], [14].

These lectures are devoted to a detailed description of the covariant lattice construction and its underlying principles. The construction is based on the bosonic formulation of the fermionic string and its ghost system [15]. The possibility of using this formalism for the construction of new modular invariant string theories was first discussed in [16], and it was used for four-dimensional strings in [6]. A related idea is the “charge lattice” of [5], which these authors obtained by bosonizing their fermionic construction. The result is a light-cone version of covariant lattices.

Although we tried to keep these notes as elementary as possible, some basic knowledge of string theory is assumed, and may be gotten from one of the many reviews of the subject, some of which are listed under [17]. In the next chapter we begin with an introduction to the principles of string construction. This section serves also a guide to the rest of the paper.

2. Basic principles of string construction

The most fundamental quantity in any string theory is a function $X^\mu(\sigma^0, \sigma^1)$ that specifies the space-time position of every point of the string (denoted by σ^1) for any value of the proper-time, σ^0 . Here μ labels the space-time coordinates. The space-time dimension is d , and our interest will ultimately be in the value $d = 4$. Throughout this paper we will only consider *closed* strings, which means that $X^\mu(\sigma^0, 0) = X^\mu(\sigma^0, \pi)$ (the choice of the σ_1 -interval is a matter of convention). When it propagates through space-time the string sweeps out a two-dimensional surface parametrized by σ^0 and σ^1 . This surface is called the *world-sheet*, in analogy to world-lines for point particles. For non-interacting closed strings it has the shape of a tube. If strings interact they do so by splitting and joining, and as a result they sweep out world-sheets that look like complicated pieces of plumbing.

The modern way of describing strings is to regard them as two-dimensional field theories. One regards $X^\mu(\sigma_0, \sigma_1)$ as a set of d scalar fields, which are functions of a two-dimensional coordinate σ^α , $\alpha = 0, 1$. Once one realizes that one is dealing with two-dimensional field theories, one is more or less naturally led to the questions: Which conditions should be imposed on such a field theory for it to describe a consistent string theory? And which two-dimensional field theories satisfy these conditions?

One has to be careful in formulating these conditions, because many claims about uniqueness of string theory were based on implicitly or explicitly imposed conditions which were unnecessarily strong. So we should begin by carefully listing and examining the conditions. Our list of what we believe to be necessary conditions for d -dimensional strings is:

1. The presence of d unconstrained scalar fields X^μ
2. Reparametrization invariance
3. Conformal invariance
4. Modular invariance

5. World sheet supersymmetry and superconformal invariance (for fermionic strings).

These are all conditions imposed on the world-sheet field theory. Remarkably, if one satisfies them one is guaranteed not to be confronted with most of the disasters that can occur in field theory in d -dimensional space-time. Reparametrization invariance means that the description of string propagation does not depend on how one chooses the coordinates σ_0 and σ_1 on the world-sheet. This has a simple interpretation in two-dimensional field theory language: it's just two-dimensional general coordinate invariance. The first two points are so basic to string theory that one would not even want to consider what would go wrong if they were not satisfied. For the last three conditions (which we will discuss in more detail in a moment) one can identify some desirable space-time properties that follow from them.

Conformal invariance guarantees Lorentz-invariance and absence of ghosts, general coordinate invariance in space-time and gauge invariance (if the theory has massless vector bosons, as is usually the case). Although it has been the subject of debate whether one can make sense of string theory without it, it seems clear that one gets into serious technical complications if one abandons it. The attempts to abandon it were motivated by the fact that conformal invariance restricts, naively at least, the space-time dimension to 26 for bosonic strings, and to 10 for fermionic ones. Since there are now other ways to get to lower dimensions, it seems highly preferable to keep it.

One of the nice consequences of modular invariance is that it ensures that chiral fermion representations coming from string theory are anomaly free. This is highly non-trivial, especially in higher dimensions, and the discovery of a rather spectacular cancellation in 10 dimensions [18] was the starting point of the current wave of interest in the subject. The precise relation with modular invariance was only understood much later [19]. Anomalies are certainly not the only problems one can expect in the absence of modular invariance, but it is the easiest one to identify.

World-sheet supersymmetry is essential for fermionic strings, and one of the problems one has in its absence is a violation of Lorentz invariance. We will discuss this in chapter 7. Here we will continue with a more detailed look at conformal and modular invariance.

2.1 CONFORMAL INVARIANCE

A conformal transformation is a coordinate dependent rescaling of the metric on the world-sheet. It is this invariance which makes string theory so simple in comparison with, for example, membranes. The standard two-dimensional action for free bosons,

$$S = \int d^2\sigma \sqrt{-g} g^{\alpha\beta} \partial_\alpha X^\mu \partial_\beta X_\mu,$$

where $g_{\alpha\beta}$ is the world-sheet metric and g its determinant, is not only general coordinate invariant, but has an additional invariance under $g_{\alpha\beta} \rightarrow \Lambda(\sigma^\alpha) g_{\alpha\beta}$. This invariance makes it possible to choose a gauge (light-cone gauge) in which one gets an action which depends only on the $d - 2$ “transverse” (i.e. transforming in the vector representation of the little group $SO(d - 2)$ of a massless particle) degrees of freedom X^i , and which

is non-interacting and manifestly ghost-free. In the absence of conformal invariance there is an extra mode of the metric, the Liouville mode, which enters into the action, and which makes life considerably more complicated. Thus we would like to maintain conformal invariance even if other terms are added to the action.

Although conformal invariance is a symmetry of the classical action, it is afflicted by an anomaly in the quantized theory. There are several contributions to this anomaly. In units in which a scalar contributes 1, the relevant ones are $\frac{1}{2}$ for a Majorana-Weyl fermion, -26 for the ghosts of general coordinate invariance, and +11 for the ghosts of local supersymmetry. Notice that though the graviton has no physical components in two dimensions, it still contributes via its ghosts. There are two obvious ways of cancelling these anomalies :

$$N = 0 : \quad 26 \text{ scalars} + \text{gravity ghosts}$$

$$N = 1 : \quad 10 \text{ scalars and Majorana Weyl fermions} + \text{gravity ghosts} + \text{superghosts}$$

Since there is always general coordinate invariance on the world sheet, one always has the corresponding ghost. In these solutions one can replace a scalar by two fermions or vice versa, but that amounts just to two-dimensional bose-fermi equivalence and does not really bring anything new. We have made our choice in such a way that supersymmetry is manifest, but that is not necessary and we will in fact choose a purely bosonic realization in our construction. One can also try to have more supersymmetries, but the resulting theories turn out to be relevant only in two space-time dimensions or less.

At this point we should make a trivial but crucial remark. The traditional interpretation of these solutions is to associate all the scalars with space-time coordinates. In that case the $N = 0$ system lives in 26 dimensions, and the $N = 1$ system in $d = 10$. This is not necessary however. One can equally well associate just a subset of the scalars with space-time coordinates, and regard the others as some sort of “internal” degrees of freedom. Then the numbers 26 and 10 are “maximal” rather than “critical” dimensions. In some cases, and in particular for bosonic strings, this means nothing more than that some dimensions are “compactified”, but for fermionic strings there are more interesting possibilities, as we will see.

A special feature of conformally invariant closed strings is that the modes of vibration of the string can be unambiguously separated into *left-moving* ones and *right-moving* ones. Since the world-sheet theory is non-interacting, and because there are no endpoints of the string that modes can bounce back from, a left-moving mode will move to the left forever. This left/right-mover separation goes so far that one can actually choose *different* conformal systems for them. This leads in an obvious way to three kinds of closed string theory:

Bosonic :	$N = 0$ left-movers;	$N = 0$ right-movers
Heterotic :	$N = 0$ left-movers;	$N = 1$ right-movers
Type II :	$N = 1$ left-movers;	$N = 1$ right-movers

This also specifies our convention for left- and right-movers in the heterotic string.

We will mainly consider heterotic strings in this paper, although everything we say is valid for type II strings as well. The latter seem however phenomenologically less promising at the moment [20], [21]. The heterotic string forces us to make a previous condition more precise. Since it has 26 bosons on the left and only 10 on the right, it is not *a priori* obvious in which dimensions it can exist. The precise requirement on the presence of a set of scalars X^μ is however that both its left and right-moving components are present in the theory. One cannot associate an unpaired left-moving mode with a flat space-time dimension (this would be in conflict with modular invariance). Thus the maximal dimension for the heterotic string is 10. Psychologically it was perhaps this heterotic string asymmetry that lead to a re-evaluation of the role of world-sheet scalars, making four-dimensional strings possible.

2.2 MODULAR INVARIANCE

So far we have imposed conditions to ensure that string theory is well-defined locally on the world-sheet. In the perturbation expansion of string theory one does however encounter world-sheets with non-trivial topology, where the *global* properties of the theory become important. The simplest of these diagrams is the torus, roughly corresponding to a one-loop diagram in field theory. Topologically the description of the multiloop diagrams in the expansion of closed (oriented) strings is very simple: they are all equivalent to multiple tori, and there is just one topology for each order in the expansion. These loops may have external lines (tubes) attached to them. Such surfaces are known as *Riemann surfaces*.

By "modular invariance" we mean absence of global anomalies on any of these surfaces. We also mean *factorization*, i.e. the amplitude for a multiloop diagram should equal the product of its subdiagrams in a limit where it is pulled apart into these subdiagrams. Although one should impose modular invariance as a condition on diagrams with and without external lines, it is sufficient to examine the ones without external lines, because all other diagrams can always be obtained from them as a subdiagram of some higher loop surface.

The exact conditions for modular invariance beyond one loop are not known for fermionic strings, since one does not know enough about the dependence on the "supermoduli", but reasonable *ansätze* exist, and it is under these assumptions that one usually checks modular invariance. For pedagogical reasons we will not say much about higher loop modular invariance except that it is a property of the class of theories we will consider (under the aforementioned assumptions, which have to be made for any fermionic string theory).

One-loop modular invariance is however rather easy to understand. A one-loop diagram is a torus, and it can be represented by points in the complex plane modulo a two-dimensional lattice. This corresponds to a choice of coordinates and a flat, constant metric on the torus. To regard the complex plane modulo a lattice simply means that points that differ by a lattice vector are identified with each other. A two dimensional lattice has two basis vectors, chosen so that all other vectors are integral linear combinations of them. By rotations and scale transformations one can always align one of the basis vectors with the real axis, and scale it to have length 1. The second one can

always be put into the positive upper half plane, where it can be specified by a complex number τ , with $Im(\tau) > 0$. We have now used all the transformations at our disposal to bring the metric in the simplest possible form. The remaining free parameter of the torus, τ , parametrizes metrics that cannot be mapped into each other by general coordinate transformations or Weyl transformations.

Not all these metrics are inequivalent however. Although they cannot be related by *local* transformations, some can be related by *global* ones. For example, if one defines a torus by $\tau' = \tau + 1$ one makes the same identification in the complex plane, but the metric would look rather different, i.e. have a different but equivalent parametrization. On a real-life, donut-shaped torus this transformation corresponds to cutting the torus along a cycle which one cannot be continuously deformed to a point, rotating one edge by 2π , and glueing the edges back together. A torus has two independent cycles of this type, and hence there is a second transformation of τ which leads to equivalent metrics, namely $\tau \rightarrow -\frac{1}{\tau}$. Together these transformations generate the group of *modular transformations*. Although one can write down an explicit expression for all its elements, one needs in practice usually only its two generators,

$$\begin{aligned}\tau &\rightarrow \tau + 1 \\ \tau &\rightarrow -\frac{1}{\tau}\end{aligned}\tag{2.1}$$

To calculate loop-diagrams à la Polyakov [22] one performs a path-integral over all fields in the two-dimensional action (for a given metric on the world-sheet), and furthermore one integrates over all inequivalent metrics. The expression for a one loop diagram is (after some gauge fixing) of the form

$$\int \frac{d^2\tau}{(Im\tau)^2} \mathcal{F}(\tau),$$

where $\mathcal{F}$ is some function which depends on the theory, and the other factor is the invariant integration volume. The function $\mathcal{F}$ (or rather a slightly modified function, defined more precisely in section (4.3)) is called the *partition function*. The issue of modular invariance centers on the boundaries of the τ -integral. If one were to integrate over the entire positive half plane, one would encounter the same metric an infinite number of times. One would like to restrict the integration to a *fundamental domain*, which is defined as a region in the upper half plane which contains every metric just once (the canonical choice is $-1 \leq Re\tau < 1, |\tau| \geq 1$), but this can only be done in a meaningful way if $\mathcal{F}(\tau)$ does not depend on the region one chooses, since otherwise the result would be ambiguous. Thus $\mathcal{F}$ should be invariant under (2.1). This is what is meant by modular invariance.

For an n -loop surface the problem is very similar. Instead of a single complex parameter τ one gets now a complex $n \times n$ matrix Ω , the *period matrix*. For discussions of modular transformations for higher genus Riemann surfaces we refer to [23].

2.3 FOUR DIMENSIONAL STRINGS: THE BASIC IDEA

There is a fairly logical path towards four-dimensional string constructions of the type we are going to consider, and it might be useful to have some idea about this before starting with the more technical aspects. Terms not explained below will be discussed in detail in the remaining chapters.

Consider first the ten-dimensional heterotic string. Its construction has been explained in much detail and with great clarity in [1], which is obligatory literature for anyone interested in string constructions. Here we concentrate on just one aspect, namely the role of the 16 left-moving bosons that are not paired with right-movers into space-time coordinates. Instead one regards them as compactified space-time coordinates (at least in the mathematical sense of the word, because as a physical concept this is somewhat inadequate (and unnecessary)). The center-of-mass coordinate of the string in these 16 dimensions is put in a periodic space (a torus) with periodicities specified by a 16 dimensional lattice. Modular invariance restricts the choice of this lattice (and in particular does not allow an infinite radius torus, which would correspond to flat space). One finds that it must be even and *self-dual*, and such 16-dimensional lattices have been classified, the only possibilities being $E_8 \times E_8$ or a certain subset of the $SO(32)$ weight lattice.

Narain took this construction one logical step further. Instead of insisting on 10-dimensional Lorentz-invariance he required only Lorentz-invariance in $d < 10$ dimensions (we will choose $d = 4$ here, for definiteness and other reasons), and consequently paired only 4 left- and right-moving bosons into space-time coordinates. The remaining ones, 22 left-movers and 6 right-movers are “compactified” (the physical meaning of this terminology has become less and less clear throughout the historical development, but the mathematical description is unambiguous) on a torus of total dimension $22+6$. The condition for modular invariance is that the lattice defining this torus must be even and self-dual with respect to a *lorentzian metric*. There is a continuous infinity of such lattices!

A problem with this approach is that it does not lead to chiral theories. This is related to the fact that the 10 right-moving Majorana-Weyl fermions of the heterotic string have remained untouched. Since they are responsible for the existence of space-time fermions, what happens is that all ten-dimensional space-time fermions which are chiral are trivially compactified on a torus, and it is a well-known fact that this leads to non-chiral theories in lower dimensions.

The next logical step in the development avoids this problem by making the Majorana-Weyl fermions participate in the process. To do so, they have to be on equal footing with the other fields, which can be achieved by bosonizing them. Actually we are not going to bosonize anything, but we will just start with 15 bosons (rather than with 10 bosons and 10 fermions) to cancel the conformal anomaly. The five bosons replacing the 10 fermions are included in the same torus, which now has dimension $22+11$. There is a small technical complication however: to get a lattice description of the theory one has to bosonize also the ghosts of local supersymmetry, and enlarge the lattice to include the charge of the resulting boson.

Now one can check the conditions for modular invariance, and the result is again that the torus has to correspond to a self-dual lattice, only this time it is not even but odd. This is all that needs to be said about modular invariance in these theories. It can be shown that (within the usual assumptions) this self-duality guarantees not just one-loop, but also higher-loop modular invariance.

Before we can make all this more precise we have to explain the corner stones of constructions of this type, and we will do so in the next three chapters on lattices, torus compactification and conformal field theory. These chapters are more or less independent, and readers sufficiently familiar with any of these subjects may want to skip some of them. Chapter 6 discusses the Neveu-Schwarz-Ramond model and its bosonic form, which is the heart of our construction. In particular we have to explain in some detail the properties of its bosonized ghost system. It is unfortunately not possible to explain this without the sophisticated techniques of conformal field theory, and although we have tried to supply the necessary information in chapter 5, we are aware of the fact that it may not be adequate to readers unfamiliar with it. Fortunately one finally arrives at a formulation of the theory where most of this can be forgotten again, at least as long as one is only interested in model building and spectrum calculations. For this reason we have formulated chapters 8 and 9 in such a way that they are self-contained, so that even readers who didn't fully grasp the ghost system should be able to apply the rules of model building. Meanwhile chapter 7 contains the discussion of world-sheet supersymmetry promised above.

3. Lattices

In this section we collect most of the known facts about self-dual lattices. In the mathematics literature, most work on this subject appears in journals on Number Theory, which are unfortunately not always easily available in physics libraries. Some of the relevant papers are listed under Ref. [24-26], and readers interested in proofs not provided below should consult these papers. We also recommend the reviews by Goddard and Olive [27].

3.1 DEFINITIONS

A lattice Λ is defined as a set of points in a vector space V of dimension N , of the form

$$\Lambda = \left\{ \sum_{i=1}^N n_i \vec{e}_i \mid n_i \in \mathbb{Z} \right\}, \quad (3.1)$$

where $\vec{e}_i$ ($i = 1, \dots, N$) is a set of basis vectors of V . The vector spaces we will consider are R^N , with a euclidean inner product, and $R^{p,q}$, ($p + q = N$), with a “lorentzian” inner product. In terms of the components v^a of vectors $\vec{v} \in V$ we define these inner products as $\vec{v} \cdot \vec{w} = v^a w^a d_{ab}$, with $d_{ab} = \delta_{ab}$ for R^N , and $d_{ab} = \epsilon(a) \delta_{ab}$, $\epsilon(a) = 1$ for $1 \leq a \leq p$, $\epsilon(a) = -1$ otherwise, for $R^{p,q}$. In the following we will denote these inner products as $(+)^N$ and $(+)^p(-)^q$ respectively, and we will refer to d_{ab} as the *signature*.

The *unit cell* of a lattice is the set of points $\{\vec{x} = \sum_i x_i \vec{e}_i | 0 \leq x_i < 1\}$. It contains precisely one lattice point (i.e. $\vec{x} = 0$), and every vector of V has precisely one “image” in the unit cell, obtained by adding a lattice vector to it. The *volume* $vol(\Lambda)$ of the unit cell of Λ is obtained by defining a matrix

$$g_{ij} = \vec{e}_i \cdot \vec{e}_j, \quad (3.2)$$

in terms of which

$$vol(\Lambda) = \sqrt{|\det g|}. \quad (3.3)$$

The matrix g_{ij} is called the *metric* of the lattice. It contains all information regarding the angles of the basis vectors and their lengths. Notice that if we change the signature from lorentzian to euclidean (by changing $(+)^p(-)^q$ to $(+)^{p+q}$) then $\det g$ changes at most by a sign, so that the volume of the unit cell remains the same.

The *dual* Λ^* of Λ is defined as

$$\Lambda^* = \{\vec{w} \in V | \vec{w} \cdot \vec{v} \in \mathbb{Z}, \forall \vec{v} \in \Lambda\}. \quad (3.4)$$

A basis for Λ^* is the set of vectors $\vec{e}_i^*$, with $\vec{e}_i^* \cdot \vec{e}_j = \delta_{ij}$. It is easy to see that this is indeed a basis for Λ^* , i.e.

$$\Lambda^* = \left\{ \sum_{i=1}^N m_i \vec{e}_i^* | m_i \in \mathbb{Z} \right\}. \quad (3.5)$$

To calculate the volume of Λ^* we define

$$g_{ij}^* = \vec{e}_i^* \cdot \vec{e}_j^*, \quad (3.6)$$

which is the metric of the dual lattice. Using the definition of the basis vectors one can easily show that $\sum_i e_i^{*a} e_i^b = d^{ab}$, and that therefore g^* is the inverse of g . Thus we have

$$vol(\Lambda) = (vol(\Lambda^*))^{-1}. \quad (3.7)$$

A lattice Λ is called

- *unimodular* if $vol(\Lambda) = 1$
- *integral* if $\vec{v} \cdot \vec{w} \in \mathbb{Z}, \forall \vec{v}, \vec{w} \in \Lambda$
- *even* if Λ is integral, and $\vec{v}^2 = 0 \pmod{2}, \forall \vec{v} \in \Lambda$
- *odd* if Λ is integral, but not even
- *self-dual* if $\Lambda = \Lambda^*$

Obviously Λ is integral if and only if $\Lambda \subset \Lambda^*$.

3.2 COSETS

If Λ_s is a sublattice of Λ , then we can decompose Λ in cosets with respect to Λ_s . To do so, choose a set of coset representatives $\vec{\beta}_\alpha$, so that

$$\begin{aligned}\vec{\beta}_\alpha &\in \Lambda, \vec{\beta}_\alpha \notin \Lambda_s \\ \vec{\beta}_\alpha - \vec{\beta}_{\alpha'} &\notin \Lambda_s \text{ if } \alpha \neq \alpha'.\end{aligned}\tag{3.8}$$

If we extend this set with the vector $\vec{\beta} = 0$ to represent Λ_s itself, we get

$$\Lambda = \sum_{\alpha} (\vec{\beta}_\alpha + \Lambda_s),\tag{3.9}$$

i.e. every vector in Λ can be written as $\vec{\beta}_\alpha + \vec{v}_s, \vec{v}_s \in \Lambda_s$. Every coset $\vec{\beta}_\alpha + \Lambda_s$ has precisely one point in the unit cell of Λ_s . Therefore, if there are N_s cosets (including Λ_s), we find

$$\text{vol}(\Lambda) = \frac{1}{N_s} \text{vol}(\Lambda_s).\tag{3.10}$$

This is a fairly trivial, but very useful observation. A direct consequence is that a lattice is self-dual if and only if it is both integral and unimodular.

3.3 LIE ALGEBRA LATTICES

It is often convenient to express a self-dual lattice in terms of Lie algebra lattices. In the following when we speak of a Lie algebra lattice we mean any sublattice of the weight lattice of a semi-simple Lie algebra (for a discussion of Lie algebras, roots and weights we refer to [28]). In particular we need the *simply laced* lattices, belonging to semi-simple Lie algebras which are products of D_n, A_n and E_6, E_7, E_8 . They have the property that all their roots have the same length. If one normalizes them so that they have all norm 2, the weight lattice Λ_W is the dual of the root lattice Λ_R , which is itself a sublattice of the weight lattice (the norm of a vector is its squared length)

$$\Lambda_W = \Lambda_R^* ; \Lambda_W \supset \Lambda_R\tag{3.11}$$

Therefore we can decompose Λ_W into cosets with respect to Λ_R . In the case of Lie algebra lattices these cosets are called *conjugacy classes*. In general cosets form a group under addition, with the sublattice Λ_s as the unit element. In this case this group is called the *center* of the Lie algebra.

We will make extensive use of D_n lattices, for which we denote the conjugacy classes as follows

$$\begin{aligned}(0) &= \Lambda_R = \{(k_1, \dots, k_n), k_i \in \mathbb{Z}, \sum_{i=1}^n k_i = 0 \bmod 2\} \\ (v) &= \Lambda_R + (1, 0, 0, \dots, 0) \\ (s) &= \Lambda_R + (\tfrac{1}{2}, \tfrac{1}{2}, \dots, \tfrac{1}{2}) \\ (c) &= \Lambda_R + (-\tfrac{1}{2}, \tfrac{1}{2}, \dots, \tfrac{1}{2})\end{aligned}\tag{3.12}$$

One may check that the center is isomorphic to Z_4 if n is odd, and to $Z_2 \times Z_2$ if n is even.

The irreducible representations of the semi-simple Lie algebras are characterized by their weight vectors, which for a given representation all belong to the same conjugacy class of the lattice. Hence we can specify conjugacy classes also by giving representations that belong to them. In the case of D_n , (0) contains all even rank tensor products of the vector representation (which includes the singlet and the adjoint), (v) contains all odd-rank tensors, and (s) and (c) contain spinors of opposite chirality.

A useful property of conjugacy classes is that their mutual inner products modulo integers depend only on the class, and not on the individual vectors. If $\vec{\beta}_1, \vec{\beta}_2 \in \Lambda_W$, and $\vec{\lambda}_1, \vec{\lambda}_2 \in \Lambda_R$, we have

$$(\vec{\beta}_1 + \vec{\lambda}_1) \cdot (\vec{\beta}_2 + \vec{\lambda}_2) = \vec{\beta}_1 \cdot \vec{\beta}_2 \bmod 1,$$

because $\vec{\beta}_i \cdot \vec{\lambda}_j \in Z$, and because the root lattice of simply laced algebras is integral (this follows from the fact that it is contained in its own dual, see (3.11)). In fact Λ_R is an even lattice, because its basis vectors are roots, which have norm 2. Using this property one can show by a similar argument that the norms of two vectors in the same conjugacy class are always equal modulo 2. These properties are clearly of great help in constructing even integral lattices, because one only needs to check the lengths and inner products of conjugacy class representatives. For this reason we have collected some information concerning the conjugacy classes of the (simple) simply laced algebras in the following table.

Lie algebra	conjugacy class	smallest representation	weight norm (mod 2)
D_n	(0)	singlet	0
	(v)	vector	1
	(s)	spinor	$\frac{n}{4}$
	(c)	conjugate spinor	$\frac{n}{4}$
A_n	(p)	rank p tensor ($0 \leq p < n$)	$\frac{p(n+1-p)}{n+1}$
E_6	(0)	singlet	0
	(1)	(27)	$\frac{4}{3}$
	(2)	($\overline{27}$)	$\frac{4}{3}$
E_7	(0)	singlet	0
	(1)	(56)	$\frac{3}{2}$
E_8	(0)	singlet	0

The volume of the unit cells of Λ_R and Λ_W are easy to obtain. Since they are each

other's dual

$$\text{vol}(\Lambda_R) = \text{vol}(\Lambda_W)^{-1}$$

If we denote the order of the center (i.e. the number of conjugacy classes) by N_c , we learn from (3.10) that

$$\text{vol}(\Lambda_W) = \frac{1}{N_c} \text{vol}(\Lambda_R),$$

so that

$$\begin{aligned} \text{vol}(\Lambda_R) &= \sqrt{N_c} \\ \text{vol}(\Lambda_W) &= \frac{1}{\sqrt{N_c}} \end{aligned}$$

All these results are valid for simple as well as semi-simple Lie algebras. The order of the center of a semi-simple Lie algebra is simply the product of the order of the centers of its simple factors. The same is true for the volume of the unit cell.

To specify a Lie algebra lattice completely, we have to supply two pieces of information: the semi-simple Lie algebra, and a list of the conjugacy classes that appear. For a given semi-simple Lie algebra there are in general many possible choices of conjugacy classes. This choice is restricted only by the requirement that the list must close under addition, which in particular means that the root lattice must always be present. The possible sets of conjugacy classes are thus in one-to-one correspondence with the subgroups of the center. Instead of giving a full list of conjugacy classes, one can give a shorter list of generators, which is a subset of conjugacy classes from which all others can be obtained by addition.

Since the volume of the unit cell of the root lattice is $\sqrt{N_c}$, it follows immediately that a Lie algebra lattice is unimodular if and only if it contains precisely $\sqrt{N_c}$ conjugacy classes. This is only possible if we choose a Lie algebra for which N_c is a perfect square. The lattice will be integral if and only if all conjugacy class generators have integral norm and integral mutual inner product. It will be even if and only if they have even norm and integral inner products.

Using the table one can now easily construct many examples of euclidean self-dual lattices. The reader is invited to check that

- D_n with conjugacy classes (0) and (v) is odd and self-dual (this lattice is also known as Z^n).
- D_n with (0) and (s) is odd self-dual for $n = 4 \bmod 8$. and even self-dual for $n = 0 \bmod 8$
- A_{n^2-1} with (0) and (n) is odd self-dual for n even and even self-dual for n odd.

In general we call vectors of norm 2 on euclidean lattices *roots*. The roots include in any case those of the Lie algebras explicitly used in the construction of the lattice (i.e. those in the conjugacy class (0)), but in some cases there may be additional norm 2 vectors in the other conjugacy classes. In that case we can always write the lattice in a different way, using larger Lie algebras and fewer conjugacy classes, although this is not always convenient. For example, the lattices D_8 with (0) + (s) and A_8 with (0) + (3)

defined above are both equal (up to rotations) to E_8 , which has only one conjugacy class, namely (0) (and which is therefore automatically self-dual!).

A large class of even lorentzian self-dual lattices in $R^{p,p}$ can be constructed as follows. Take a semi-simple Lie algebra of rank p . Consider the set of all vectors in $R^{p,p}$ of the form $(\vec{v}_1, \vec{v}_2)$, so that $\vec{v}_1$ and $\vec{v}_2$ belong to the same conjugacy class of the Lie algebra. With respect to the lorentzian metric any such vector has integral inner product with any other such vector, any furthermore they all have even length. It is also easy to see that the resulting lorentzian lattice is unimodular, so that it is a self-dual lattice. Lattices of this type were first described by Englert and Neveu [29].

To conclude this discussion of Lie algebra lattices, we want to introduce a minor, but useful extension of this set, namely the lattice D_1 . Although it is not part of Cartan's classification of simple Lie algebra's, it is easy enough to define by simply extending the definition of D_n to $n = 1$. The resulting one-dimensional root lattice consists simply of the even integers. The shortest non-trivial vector has norm four, so there are no roots on this root lattice. The vector conjugacy class consists of the odd integers, and the two spinor conjugacy classes are obtained by shifting the root lattice by $+\frac{1}{2}$ and $-\frac{1}{2}$.

3.4 LATTICE ENGINEERING

There are several tricks for making new self-dual lattices out of already known ones. These tricks are all variations on the same basic theme: one adds new conjugacy classes and removes old ones, in such a way that their total number stays the same (i.e. the lattice remains unimodular), and that all inner products remain integral.

A very general method is the following. Let Λ be a self-dual lattice. Choose a vector $\vec{s}$ so that $k\vec{s} \notin \Lambda, 1 \leq k \leq p-1$, but $p\vec{s} \in \Lambda$. For convenience we will assume that p is prime for the moment. Now we can define a lattice Λ_0 as the sublattice of Λ which has integer inner product with $\vec{s}$. Since $\vec{s} \notin \Lambda$ there must be a vector $\vec{t} \in \Lambda$ such that $\vec{s} \cdot \vec{t}$ is not integer. But then $p\vec{t}$ belongs to Λ_0 , and is the smallest multiple of $\vec{t}$ in Λ_0 . Hence we can decompose Λ in cosets

$$\Lambda = \sum_{k=0}^{p-1} (k\vec{t} + \Lambda_0)$$

Consider now the new lattice

$$\Lambda' = \sum_{k=0}^{p-1} (k\vec{s} + \Lambda_0)$$

clearly Λ' is unimodular. If furthermore $\vec{s}$ has integral norm, it is self-dual. If Λ is even and $\vec{s}$ has even length, Λ' is even self-dual.

Another useful trick is the following. Consider a Lie algebra lattice containing, among others, a D_n factor. Now replace this D_n factor by D_m , where $m = n + 8l, l \in \mathbb{Z}$. Since the unit cells of D_n and D_m have the same volume for any n and m , the new lattice is unimodular. Furthermore one can easily check that if $n = m \bmod 8$ all inner products and norms of the four conjugacy classes remain unchanged modulo two. Hence the new lattice will be (even) self-dual if the original lattice was (even) self-dual.

This transformation is of course meaningless if it results in $m = 0$. But one can give a meaning to it if m becomes negative. Consider for example a euclidean even self-dual lattice $\Lambda_p \times D_n$ with inner product with signature $(+)^{p+n}$. For $m < 0$ (with m defined as above) we can consider the lorentzian lattice $\Lambda_p \times D_m$ with signature $(+)^p(-)^{-m}$. This lattice is unimodular for the reasons explained above. Furthermore the lorentzian inner products of the new lattice are equal to the euclidean ones of the original one modulo two. Thus we see that shifting n by multiples of eight to negative values has the effect of changing a euclidean self-dual lattice to a lorentzian one.

3.5 LATTICE CLASSIFICATION THEOREMS

A lot is known about the possibilities for constructing self-dual lattices. One should keep in mind here that euclidean lattices can only be classified up to rotations, because any rotation respects the self-duality conditions. Likewise, lorentzian lattices can only be classified modulo Lorentz transformations. The known results are as follows:

1. *Lorentzian, odd.* All such lattices are Lorentz transformations of the lattice $Z^{p,q}$, which is the set of points in $R^{p,q}$ defined as $\{(n_1, \dots, n_p, m_1, \dots, m_q), n_i, m_j \in \mathbb{Z}\}$.

2. *Lorentzian, even.* Lattices of this type exist only for $|p - q| = 0 \bmod 8$. For $p = q + 8n$ they are Lorentz transformations of the lattice $nE_8 \times qP^{1,1}$. The E_8 lattice is most easily described as D_8 with conjugacy classes (0) and (s) , while $P^{1,1}$ is a lorentzian lattice of signature $(+)(-)$, which may be described as $D_1 \times D_1$ with conjugacy classes $(0,0) + (v,v) + (s,s) + (c,c)$ (this is a lattice of the Englert-Neveu type, discussed above).

3. *Euclidean, odd.* A class of such lattices which exists in any dimension is Z^n . In general, if any odd self-dual lattice has vectors of length one, they can only have mutual inner products $\pm 1, 0$. Thus they form an orthonormal basis for a subspace of R^n , which must be part of the basis of the lattice. Any other basis vector of the lattice must be orthogonal to this subspace. Thus if there are $2p$ length one vectors, the lattice is of the form $Z^p \otimes \Lambda^{n-p}$, where the second factor does not contain any vectors of length one.

An odd euclidean self-dual lattice Λ can be related to an even self-dual one of higher dimension in the following way [24]. Consider the sublattice Δ_0 of even norm vectors. Obviously Δ_0 has only one non-trivial coset in Λ , because the sum of two odd norm vectors is even. Thus

$$\Lambda = \Delta_0 + \Delta_v,$$

where Δ_v denotes the odd-norm vectors. Since $\text{vol}(\Delta_0)/\text{vol}(\Delta_0^*) = 4$, the dual of Δ_0 can be decomposed into four cosets,

$$\Delta_0^* = \Delta_0 + \Delta_v + \Delta_s + \Delta_c$$

These four cosets generate an abelian group of order four, which can only be Z_4 or $Z_2 \times Z_2$. This allows us to determine the mutual inner products of the four cosets and their norms modulo 2. If the group is Z_4 , one finds $\Delta_c = -\Delta_s$, and $2\Delta_s \subset \Delta_v$, so that the norm of all vectors in Δ_s and Δ_c is equal to $\frac{k}{4} \bmod 2, k$ odd. In the latter case $2\Delta_s = 2\Delta_c \subset \Delta_0$, so that the norm of all vectors in Δ_s and Δ_c is half-integer. One can then use $\Delta_s + \Delta_v = \Delta_c$, plus the fact that vectors in Δ_s cannot have integer inner

products with those in Δ_v , to show that the norm of vectors in Δ_s and Δ_c must be equal modulo 2. Thus for both possible abelian groups the vectors in Δ_s and Δ_c have, modulo two, norms equal to each other and to $\frac{k}{4}$, $0 \leq k < 7$. Consider then the lattice obtained by adding a D_{8-k} factor, so that the entire lattice consists of the vectors

$$(\Delta_0, 0) + (\Delta_v, v) + (\Delta_s, s) + (\Delta_c, c)$$

This lattice is unimodular, integral and even, so that it is even self-dual. So all odd self-dual lattice can be obtained from the even self-dual ones by inverting this procedure to remove D_n factors. This leaves us with the classification problem of

4. *Euclidean, even lattices.* These exist only in dimensions which are a multiple of 8, and are known explicitly for 8, 16 and 24 dimensions. In 8 dimensions, the only possibility is E_8 ; in 16 dimensions one can have $E_8 \otimes E_8$ and D_{16} with conjugacy classes (0) and (s). These two lattices cannot be rotated into each other. In 24 dimensions there are 24 even self-dual lattices, which are usually called the *Niemeyer lattices* [25], after the German mathematician who first classified them. A list of their conjugacy classes can be found in the first paper of Ref. [24].

One of the 24 lattices is the Leech-lattice [30], which differs from the other 23 because it contains no vectors of norm 2. Nevertheless, we can in a certain sense regard it as a Lie algebra lattice, which is a product of 24 D_1 factors, on which one has 2^{24} conjugacy classes. A list of these classes (or rather 12 classes that generate them) will appear in [31].

There is a rather amusing way of estimating the number of lattices in $8n$ dimensions. Every lattice has a discrete automorphism group. This is a discrete subgroup of $O(8n)$ which takes the lattice into itself. It consists of at least two elements, the identity and the reflection through the origin. Denote the total number of elements of this group for a lattice Λ as $g(\Lambda)$. Then the following identity can be proved [26], [32].

$$\sum \frac{1}{g(\Lambda)} = \frac{B_{4n}}{8n} \prod_{j=1}^{4n-1} \frac{B_{2j}}{4j},$$

where B_{2j} are the Bernoulli numbers. This formula is known as the Minkowski-Siegel mass formula. The sum is over all lattices in $8n$ dimensions, and adds up to a number that is less than half the total number of lattices in this dimension. Thus the right hand side, multiplied by 2, is a lower limit of the number of lattices. Calculating it for the lowest dimensions, one finds

dimension	lower limit	actual number
8	2.8×10^{-9}	1
16	4.9×10^{-18}	2
24	15.8×10^{-15}	24
32	8.0×10^7	?

This explains why the lattices of dimension 32 have not yet been classified. The explosive growth of the number of lattices that one observes here is related to the reason why the number of chiral four dimensional string theories is so much larger than the number of ten-dimensional ones, as we will see.

4. Torus Compactification of the Bosonic String

Armed with all relevant facts about lattices, we can now attack the problem of the bosonic string, propagating through a 26 dimensional space-time of which some dimensions have been "compactified". Of course our ultimate interest is in fermionic strings, but since we will use a bosonic approach to fermionic strings, a thorough understanding of the compactified bosonic string is essential. The presentation given here is essentially the one of [4], [33], although these authors compactified the heterotic string rather than the bosonic string. To warm up, and to introduce some notations, we will begin with a very brief reminder of the basics, the quantization of the uncompactified string.

4.1 THE UNCOMPACTIFIED CLOSED BOSONIC STRING

The propagation of a string through d dimensional space-time is governed by the action

$$S = -\frac{1}{4\pi\alpha'} \int d\sigma_0 \int_0^\pi d\sigma_1 \partial_\alpha X^\mu \partial^\alpha X^\nu \eta_{\mu\nu}, \quad (4.1)$$

where σ_0 and σ_1 are the world-sheet coordinates, indicating respectively the time direction and the direction along the string (one often denotes these variables as τ and σ , but we won't do this to prevent confusion with the usual notation for the one-loop modular parameter). In (4.1) $\eta_{\mu\nu}$ is a flat metric and $X^\mu(\sigma_0, \sigma_1)$ is a map from the world-sheet to space-time, which specifies the position of every point of the string at any time. The parameter α' (the "Regge slope parameter") is usually set to the value $\frac{1}{2}$, but we will refrain from that for a while. We shall only consider closed strings, for which one has the boundary condition

$$X^\mu(\sigma_0, 0) = X^\mu(\sigma_0, \pi). \quad (4.2)$$

From the action one can in the usual way derive the operators H and P which generate translations in the σ_0 and σ_1 directions on the world-sheet:

$$H = \frac{1}{4\pi\alpha'} \int_0^\pi d\sigma_1 [(\partial_0 X^\mu)(\partial_0 X_\mu) + (\partial_1 X^\mu)(\partial_1 X_\mu)] \quad (4.3)$$

$$P = \frac{1}{2\pi\alpha'} \int_0^\pi d\sigma_1 (\partial_0 X^\mu \partial_1 X_\mu) \quad (4.4)$$

The quantization of this system is simplest in light-cone gauge, in which the dynamical variables are $X^i, i = 1, \dots, d-2$. Their canonical momenta Π_i are easily obtained from the action

$$\Pi_i = \frac{1}{2\pi\alpha'} \partial_0 X_i.$$

The solutions of the equations of motion have the form

$$X^i(\sigma_0, \sigma_1) = q^i + 2\alpha' p^i \sigma_0 + \frac{i}{2} \sqrt{2\alpha'} \sum_{n \neq 0} \frac{1}{n} (\alpha_n^i e^{-2in(\sigma_0 + \sigma_1)} + \tilde{\alpha}_n^i e^{-2in(\sigma_0 - \sigma_1)}). \quad (4.5)$$

Using the canonical commutator

$$[X^i(\sigma_0, \sigma_1), \Pi^j(\sigma_0, \sigma'_1)] = i\delta^{ij}\delta(\sigma_1 - \sigma'_1) \quad (4.6)$$

one finds the commutation relations for the quantized modes

$$\begin{aligned} [q^i, p^j] &= i\delta^{ij} \\ [\alpha_m^i, \alpha_n^i] &= m\delta^{ij}\delta_{n+m,0} \\ [\tilde{\alpha}_m^i, \tilde{\alpha}_n^i] &= m\delta^{ij}\delta_{n+m,0} \\ [\alpha_m^i, \tilde{\alpha}_n^i] &= 0 \end{aligned} \quad (4.7)$$

The spectrum is obtained by requiring that both H and P vanish. This leads to the conditions

$$2p_+p_- - p_i^2 \quad (= p^2 = m^2) = \frac{2}{\alpha'}(\mathcal{N} + \tilde{\mathcal{N}} - 2), \quad (4.8)$$

where the -2 is due to the zero mode fluctuations of the oscillators, and we have defined the number operators

$$\mathcal{N} = \sum_{n=1}^{\infty} \alpha_{-n}^i \alpha_n^i$$

and similarly for $\tilde{\mathcal{N}}$. From the vanishing of P we get

$$\mathcal{N} = \tilde{\mathcal{N}}.$$

In the quantized theory this is imposed as a condition on the physical spectrum.

It is convenient to define an operator H' , which is equal to H , but without the zero-mode terms (i.e. the operators p^i and q^i). It is also useful to define the linear combinations

$$\begin{aligned} H_L &= \frac{1}{2}(H' - P) = 2(\tilde{\mathcal{N}} - 1) \\ H_R &= \frac{1}{2}(H' + P) = 2(\mathcal{N} - 1) \end{aligned}$$

which are the generators of world-sheet translations in the $\sigma_0 - \sigma_1$ and $\sigma_0 + \sigma_1$ directions respectively. Their eigenstates are called left- and right-moving modes respectively. Denoting the eigenvalues of H'_L and H'_R on a state $|\psi\rangle$ as $\frac{1}{2}\alpha'm_L^2$ and $\frac{1}{2}\alpha'm_R^2$ we can write the result obtained above as^{*}

$$\begin{aligned} H_L |\psi\rangle &= \frac{1}{2}\alpha'm_L^2 |\psi\rangle \\ H_R |\psi\rangle &= \frac{1}{2}\alpha'm_R^2 |\psi\rangle \\ m^2 &= \frac{1}{2}(m_L^2 + m_R^2) \end{aligned}$$

with the condition

$$m_L^2 = m_R^2.$$

^{*} For comparison with [6] note that we have reversed the assignment of $\tilde{\alpha}$ and $\tilde{\tau}$ here from right to left-movers here. This is more convenient in heterotic strings, where we will choose the NSR system to be right-moving, just as in [6].

4.2 THE COMPACTIFIED CLOSED BOSONIC STRING

Now we turn to the case of a torus compactified space-time. A torus in N dimensions is simply R^N modulo a lattice Λ . This means that points in R^N which differ by a vector on Λ are identified. If a string propagates in a space-time with this topology it is not necessary anymore that it closes into the same space-time point, as in (4.2). We should allow for the possibility that the points $\sigma_1 = 0$ and $\sigma_1 = \pi$ of the string are identified points in R^N :

$$X^i(\pi) = X^i(0) + 2\pi L^i,$$

where $L^i \in \Lambda$ and the factor 2π is put in for convenience. Physically, this corresponds to a string winding around the torus a certain number of times. To accommodate this possibility one modifies the string field X^i in the following way

$$X^i(\sigma_0, \sigma_1) = q^i + 2\alpha' p^i \sigma_0 + 2L^i \sigma_1 + \frac{i}{2} \sqrt{2\alpha'} \sum_{n \neq 0} \frac{1}{n} (\alpha_n^i e^{-2in(\sigma_0 + \sigma_1)} + \tilde{\alpha}_n^i e^{-2in(\sigma_0 - \sigma_1)}), \quad (4.9)$$

where L^i is an operator whose eigenvalues are the lattice vectors.

Having done this, we go now through the same exercise as for the uncompactified string. It is easy to verify that nothing changes in the canonical commutators (4.7) except for the trivial addition that L^i should commute with all other operators. The mass formula replacing (4.8) acquires on the right hand side an extra term $\frac{1}{\alpha'^2} L^2$. Furthermore, to interpret the left hand side as a mass in $d - N$ dimensions we have to move the internal components of p^i to the right, to get

$$p^2 = m^2 = \sum_{I=1}^N p_I^2 + \frac{1}{\alpha'^2} L^2 + \frac{2}{\alpha'} (\mathcal{N} + \tilde{\mathcal{N}} - 2).$$

Here we have chosen the first N components as the compactified ones. The momenta $p_I, I = 1, \dots, N$ are canonical momenta of a periodic coordinate, and therefore (for reasons very familiar from solid state physics) they are quantized. Indeed, if we write their eigenfunctions of momentum k_I as $e^{ik_I x_I}$, then we see that these functions are periodic on the boundaries of the unit cell of Λ if and only if $\vec{k} \cdot \vec{l} \in \mathbb{Z}$ for every $\vec{l} \in \Lambda$. This means that the eigenvalues of k_I must lie on the dual lattice Λ^* of Λ .

The condition obtained from the vanishing of P reads now

$$\mathcal{N} - \tilde{\mathcal{N}} + p_I L_I = 0$$

It is again useful to define left and right hamiltonians H' exactly as before, but with the understanding that only the space-time momenta are removed from H to get H' , not the internal ones. Then we get

$$\begin{aligned} \frac{1}{4} \alpha' m_R^2 &= \frac{1}{2} p_R^2 + \mathcal{N} - 1 \\ \frac{1}{4} \alpha' m_L^2 &= \frac{1}{2} p_L^2 + \tilde{\mathcal{N}} - 1, \end{aligned} \quad (4.10)$$

where as before left and right masses must be equal to each other and to m^2 . The left

and right momenta are defined as

$$\begin{aligned} p_R^I &= \frac{1}{\sqrt{2}}(\sqrt{\alpha'} p^I + \frac{1}{\sqrt{\alpha'}} L^I) \\ p_L^I &= \frac{1}{\sqrt{2}}(\sqrt{\alpha'} p^I - \frac{1}{\sqrt{\alpha'}} L^I) \end{aligned}$$

Before discussing the meaning of these left and right momenta, we want to mention a simple generalization, obtained by adding a background antisymmetric tensor field. This background field is coupled to X^μ via an additional term in the action

$$S_B = \frac{1}{2\pi\alpha'} \int d\sigma_0 \int_0^\pi d\sigma_1 \epsilon^{\alpha\beta} B_{\mu\nu} \partial_\alpha X^\mu \partial_\beta X^\nu$$

We take $B_{\mu\nu}$ to be a constant background field, which has non-vanishing components only in the compactified dimensions. The extra term is then a total derivative, which does not affect H or P . So how can it make any difference? It can, because there is a change in the canonical momentum operator, which now becomes

$$\begin{aligned} \Pi_\mu &= \frac{1}{2\pi\alpha'} (\partial_0 + B_{\mu\nu} \partial_1) X^\nu \\ &= \frac{1}{\pi} (p_\mu + \frac{1}{\alpha'} B_{\mu\nu} L^\nu + \text{oscillators}) \end{aligned}$$

For the oscillator contributions, everything remains the same. Although $\partial_1 X^\nu$ contains oscillator terms, there is a cancellation between left- and right-mover contributions to $[\Pi, X]$, so that the canonical commutators of the oscillators remain unchanged. The difference occurs for the zero modes, where p_μ is replaced as the generator of translations by $\tilde{p}_\mu = p_\mu + \frac{1}{\alpha'} B_{\mu\nu} L^\nu$. So it is $\tilde{p}$ rather than p which has its eigenvalues quantized on the dual of Λ . Although the mass formula remains the same as a function of p , it makes now more sense to express it in terms of $\tilde{p}$. The result is the same as (4.10) above, but with

$$\begin{aligned} p_R^I &= \frac{1}{\sqrt{2}}(\sqrt{\alpha'} \tilde{p}^I + \frac{1}{\sqrt{\alpha'}} (L^I - B^{IJ} L_J)) \\ p_L^I &= \frac{1}{\sqrt{2}}(\sqrt{\alpha'} \tilde{p}^I - \frac{1}{\sqrt{\alpha'}} (L^I + B^{IJ} L_J)) \end{aligned} \tag{4.11}$$

where $\tilde{p} \in \Lambda^*$ and $L \in \Lambda$. In the following we will omit the tilde on p . It is clear that the dependence on α' is not very significant. We can absorb it by changing the length of all vectors on Λ by $\sqrt{\alpha'}$ (and hence those on Λ^* by $\frac{1}{\sqrt{\alpha'}}$). This amounts to a rescaling of the size of the torus. From now on we will use this freedom to set $\alpha' = \frac{1}{2}$.

To interpret the result it is convenient to express the left and right momenta in terms of the basis vectors e_a^I of the lattice:

$$\begin{aligned} L^I &= e_a^I n_a \\ p^I &= e_a^I m_a = e_a^I g_{ab}^* m_b \end{aligned}$$

Then we get

$$\begin{aligned} p_R^I &= e_a^I \left(\frac{1}{2} g_{ab}^* m_b - g_{ab}^* B_{bc} n_c + n_a \right) \\ p_L^I &= e_a^I \left(\frac{1}{2} g_{ab}^* m_b + g_{ab}^* B_{bc} n_c - n_a \right) \end{aligned}$$

(upper and lower indices have no special significance here.) One can regard g (or its inverse g^*) as a constant background field, just as B . This becomes evident if one performs this change of basis directly in the action (4.1) : the effect is to change the flat metric δ_{IJ} (in the compactified dimensions) to g_{IJ} .

It is now instructive to calculate the inner product of the $2N$ dimensional vectors^{*} $P = (p_L; p_R)$ and $P' = (p'_L; p'_R)$ (the latter is expressed in terms of n' and m') with respect to the lorentzian signature $[(+)^N; (-)^N]$. The result is

$$\vec{P} \cdot \vec{P}' = nm' + mn'$$

From this expression we can learn many things. First of all, the lattice $\Gamma_{N;N}$ consisting of the vectors P defined above is an integral and even lorentzian lattice. Secondly, the dot-products do not depend on the background fields g and B . Therefore they must be parameters of a Lorentz transformation. Finally, using this background independence, we can make the choice $e_a^I = \delta_a^I$, $g_{IJ} = \delta_{IJ}$, $B_{IJ} = 0$. In this special case the lattice is $(D_1 \times D_1)^N$ with conjugacy classes $(0, 0) + (v, v) + (s, c) + (c, s)$ in each factor, which is self-dual. Because Lorentz transformations preserve self-duality, we conclude that $\Gamma_{N;N}$ is self-dual for any value of the background fields.

Thus compactification of the bosonic string on a torus R^N/Λ , with an additional background field B_{IJ} , can be described in terms of a new lattice $\Gamma_{N;N}$ which for any Λ and B is lorentzian self-dual. It should be clear that the lorentzian signature is a mathematical device for characterizing the lattice; it has nothing to do with the metric of space-time and does not lead to ghosts.

The number of parameters of g and B is $\frac{1}{2}N(N+1) + \frac{1}{2}N(N-1) = N^2$. One can verify that they are precisely the N^2 parameters of the coset $\frac{SO(N,N)}{SO(N) \times SO(N)}$, which is the set of Lorentz transformations of the lattice modulo the rotations of the left and right sector. These rotations are a symmetry of the theory, because all masses depend on p_L and p_R only via their norms. The genuine lorentz transformations mixing p_L and p_R on the other hand are not symmetries. Although the self-duality condition is invariant under such Lorentz transformations, the spectrum is not. So we have an N^2 parameter family of different string theories.

As seems usual in string theory, these results were originally discovered in the opposite order: First Narain [4] constructed the lorentzian lattices, and afterwards Narain, Sarmadi and Witten [33] gave the background field interpretation from which we started. We still have not discussed the relevance of the self-dual lattice which emerges from this construction. To do so, we have to consider the partition function of the compactified theory.

* We separate left- and right-moving sectors by a semicolon in the following.

4.3 PARTITION FUNCTIONS

Consider first the uncompactified bosonic string at one loop. The amplitude is an expression of the form

$$A_{1-loop} = \int \frac{\mathcal{D}X \mathcal{D}g}{V_{Diff} V_C} \exp(-S_E(g, X)), \quad (4.12)$$

where V_{Diff} and V_C denote the volume of the set of local diffeomorphism and of the conformal transformations respectively, and S_E is the bosonic string action (4.1), continued to euclidean space. What is meant by this expression is that one should integrate over all values of X and the world-sheet metric on the torus that are not equivalent by reparametrizations and conformal transformations. This is the analog of the process of gauge-fixing in gauge theories, and this problem can be tackled with very similar techniques. We will not (need to) explain this procedure here, but simply state the result (the reader interested in the details is referred to [34]):

$$A_{1-loop} = V \int \frac{d\tau}{(Im\tau)^2} (Im\tau)^{\frac{2-d}{2}} \int \mathcal{D}'X \exp[-S_E(g(\tau), X)] \quad (4.13)$$

The factor V denotes the volume of space-time, which of course is infinite. This infinity is easy to understand: since the action vanishes for constant X , one of the infinitely many integrals in $\mathcal{D}X$ is simply an integral of a constant over all of space-time. This infinite factor is nothing to worry about, and is simply to be regarded as a momentum conservation delta function with argument zero, which appears because we are considering a diagram without external lines. However, we have to make sure that we factor it out correctly, which can be achieved by putting the system in a box in order to regulate the infinity. After taking out the zero mode integral, we denote the remaining integral by $\mathcal{D}'X$. After gauge fixing, the components of the world-sheet metric are constants (i.e. they are independent of the world-sheet coordinates), which depend on the modular parameter τ . The remaining X integral is therefore a simple gaussian, and the result may be written as

$$\int \mathcal{D}'X e^{-S_E} = \frac{1}{\det'(-\Delta)}, \quad (4.14)$$

where Δ is the laplacian on the torus, $\Delta = g_{\alpha\beta} \partial_\alpha \partial_\beta$, and the prime on the determinant indicates that the zero-mode should be omitted. For every component of X^μ that is factored out one is left with a factor $\frac{1}{\sqrt{Im\tau}}$, which is nothing but the normalization of the zero-mode of the laplacian on the skewed torus (which has volume $Im\tau$). This does not quite account for all of the factors of $Im\tau$ in (4.13), but the extra one is a remnant of the gauge fixing procedure. The thing to remember is that every uncompactified X^i contributes a factor $(Im\tau)^{-\frac{1}{2}}$ to the integrand.

The factor (4.14) of (4.13) is called the (one-loop) *partition function*, and will be denoted as $\mathcal{F}(\tau)$. It is possible to evaluate it directly by regularization of the infinite product of eigenvalues of Δ (see the appendix of [34]), but at one loop there is a more intuitive way. Let us first recall a result for a one-dimensional bosonic system, specified

by a coordinate $X(t)$ and an action $S(X(t))$. One can prove (if the action satisfies certain conditions) the following relation between the path integral for a euclidean action S_E and the trace of the exponential of the hamiltonian H of the system

$$\int_{PBC} \mathcal{D}X e^{-S_E} = \text{Tr} e^{-\beta H}.$$

Here “PBC” means that $X(t)$ must satisfy periodic boundary conditions on the interval $[0, \beta]$. For a derivation of this relation see e.g. [35]. The result is easily generalized to X ’s having more components, or even an infinite number of components. The latter case is relevant for us, because the scalar field $X(\sigma_0, \sigma_1)$ appearing in string theory can be regarded as having an infinite number of components labeled by σ_1 (times a finite number of components because of the space-time index, which we omitted here). To use this relation, one should think of $Im\tau$ as β . There is a small complication if $Re\tau \neq 0$, because then the periodicity of X in $Im\tau$ depends on $Re\tau$. This can be taken into account by including a σ_1 shift operator in the trace, so that states at $\sigma_0 = 0$ are not identified with the same states at $\sigma_0 = Im\tau$, but with states of a string which is rotated in the σ_1 direction by an amount $Re\tau$. The generator of infinitesimal σ_1 translations is of course P , so that one gets

$$\begin{aligned} \mathcal{F}(\tau) &= \int_{T(\tau)} \mathcal{D}'X e^{-S_E} = \text{Tr} e^{\pi Im\tau H'} e^{i\pi Re\tau P} \\ &= \text{Tr} e^{i\pi\tau H_R} e^{-i\pi\bar{\tau} H_L} \end{aligned} \tag{4.15}$$

Here we have rescaled the variables σ_0 and σ_1 by factors of π . The integral is over all two-dimensional fields X that are periodic on a torus T specified by a modular parameter τ .

This way of writing the partition function admits a simple interpretation. Expanding it in terms of $q = e^{i\pi\tau}$ one gets a power series of the form $\mathcal{F}(\tau) = \sum d_{nm} q^n \bar{q}^m$. Obviously, d_{nm} is simply the number of states of left mass m and right mass n . (At this point we should add a word of warning. The states appearing in the spectrum are only those for which $n = m$. Notice however that these are not the only states contributing to the one-loop path integral. The reader who thinks that states with $n \neq m$ are projected out by the $Re\tau$ integral should realize that the lower boundary of the standard modular domain is not a straight line.)

It is usually a simple combinatorial exercise to read off the partition function from the expressions for H_L and H_R . Every term in the hamiltonian yields a factor in the partition function, so that one can derive the partition function term-by-term. Take for example a single bosonic oscillator α_n . It contributes to the right Hamiltonian via a term $2\alpha_{-n}\alpha_n$, and when it acts it increases the mass of a state by $2n$. Thus it contributes to the partition function a factor

$$(1 + q^{2n} + q^{4n} + q^{6n} + \dots) = \frac{1}{(1 - q^{2n})},$$

where on the right the term q^{2kn} corresponds to the contribution of $(\alpha_{-n})^k$. The partition function of the bosonic string contains such a factor for every oscillator.

For the uncompactified string we leave out the zero modes in H_L and H_R as discussed before. The remaining trace, i.e. the contribution from the bosonic oscillators to the partition function, is equal to

$$\mathcal{F}_B(\tau) = \text{Tr} q^{H'_R} \bar{q}^{H'_L} = |\eta(\tau)|^{-48} \quad (4.16)$$

where η is the famous Dedekind η -function

$$\eta(\tau) = q^{\frac{1}{12}} \prod_{n=1}^{\infty} (1 - q^{2n})$$

(as demonstrated in these equations, we will freely mix τ 's and q 's in the following.)

Since the oscillator contribution to the mass formula is the same for compactified and uncompactified coordinates, the partition function for the compactified string contains the same factor. The only difference is that the zero modes of the compactified coordinates are now included in the determinant as well as the hamiltonian. So first of all we have to multiply the uncompactified string result by $\sqrt{Im\tau}$ for every compactified coordinate to remove the zero-mode normalizations in (4.12). Secondly, we have to take into account the contribution to the trace due to the extra momentum terms in the mass formula (4.10). Since this is simply an extra term in the mass formulas, it produces an extra factor in the partition function, which together with the $\sqrt{Im\tau}$ factors is called the *correction factor*, because it gives the correction in the bosonic partition function for the fact that part of the zero-modes are compactified. From the mass formula (4.10) we get for the for a lattice $\Gamma_{N;N}$

$$C(\Gamma_{N;N}) = \mathcal{L}_{N;N}(\tau) (Im\tau)^{N/2}, \quad (4.17)$$

where $\mathcal{L}_{N;N}$ is the *lattice partition function*:

$$\mathcal{L}_{N;N} = \sum_{(p_L; p_R)} e^{i\pi\tau p_R^2} e^{-i\pi\bar{\tau} p_L^2}. \quad (4.18)$$

Here the sum is over all the vectors on the lattice.

4.4 MODULAR INVARIANCE

We can now discuss the modular transformations of these functions. As explained in chapter 2, we have two transformations to check, namely $\tau \rightarrow \tau + 1$ and $\tau \rightarrow -\frac{1}{\tau}$. The first one leaves the partition function invariant if and only if it is an even function of q . This is clearly the case for both compactified and uncompactified strings, as a consequence of the fact that there are 24 transverse dimensions, and because $\Gamma_{N;N}$ is even. The second modular transformation is a bit harder to check, and requires a trick, which goes by the name of “Poisson resummation formula”.

Define instead of (4.18) the function

$$P(\tau, \vec{z}) = \sum_{(p_L; p_R)} e^{i\pi\tau(\vec{p}_R + \vec{z}_R)^2} e^{-i\pi\bar{\tau}(\vec{p}_L + \vec{z}_L)^2}, \quad (4.19)$$

where $\vec{z} = (\vec{z}_L, \vec{z}_R)$ is a $2N$ vector, which is not restricted to lie on the lattice. Since we sum over all vectors on the lattice, the function P is obviously invariant if we shift $\vec{z}$ by a lattice vector. In other words, it is periodic with the periodicity of the lattice, and hence we can Fourier transform it. Normally the Fourier analysis of lattice functions would involve the dual lattice, but because our lattice is self-dual, we don't have to bother with that. The expression for P in terms of its Fourier transform $\tilde{P}$ is

$$P(\tau, \vec{z}) = \sum_{\vec{q}} e^{-2\pi i \vec{z} \cdot \vec{q}} \tilde{P}(\tau, \vec{q}) \quad (4.20)$$

The inverse transformation is of course

$$\tilde{P}(\tau, \vec{q}) = \int_{\text{unit cell}} d^{2N}z e^{-2\pi i \vec{z} \cdot \vec{q}} P(\tau, \vec{z}),$$

i.e. we have a multiple integral over all components of $\vec{z}$ with boundaries given by the unit cell of the lattice. The function P contains a sum over all lattice vectors. We can take the exponential multiplying it inside the sum, and furthermore we can shift the variables $\vec{z}$ in this exponential factor by $\vec{p}$. This shift introduces only a factor $\exp(2\pi i \vec{q} \cdot \vec{p})$, which is equal to one because both $\vec{q}$ and $\vec{p}$ are lattice vectors. The result is thus

$$\tilde{P}(\tau, \vec{q}) = \sum_{\vec{p}} \int_{\text{unit cell}} d^{2N}z e^{-2\pi i \vec{q} \cdot (\vec{z} + \vec{p})} e^{i\pi\tau(\vec{p}_R + \vec{z}_R)^2} e^{-i\pi\bar{\tau}(\vec{p}_L + \vec{z}_L)^2}$$

The sum combined with the integral gives as an integral over *all* unit cells of the lattice, i.e. over the entire space. Thus we get

$$\tilde{P}(\tau, \vec{q}) = \int_{R^{N:N}} d^{2N}x e^{-2\pi i \vec{q} \cdot \vec{x}} e^{i\pi\tau(x_R)^2} e^{-i\pi\bar{\tau}(x_L)^2},$$

where we have introduced a new integration variable $\vec{x}$ representing, in each cell, $\vec{p} + \vec{z}$. This integral is a simple gaussian, which yields

$$\tilde{P}(\tau, \vec{q}) = |\tau|^{-N} e^{-\frac{i\pi}{\tau}(q_R)^2} e^{\frac{i\pi}{\bar{\tau}}(q_L)^2}$$

Notice that we have made use here of the fact that the lattice is unimodular, so that the usual determinant factors that one expects from gaussian integrals are equal to one. Substituting this result into (4.20) and putting $\vec{z}$ equal to zero again we get the identity

$$P(-\frac{1}{\tau}) = |\tau|^N P(\tau) \quad (4.21)$$

One can assemble all these ingredients into the one-loop partition function of the compactified bosonic string. To check the modular invariance of the complete answer one

has to verify that the factors $|\tau|$ (which are called *modular weight factors*) cancel. They do, because there are also contribution from the transformation of $Im\tau$ and $\eta(\tau)$. The transformation of $Im\tau$ is straightforward: $Im(-\frac{1}{\tau}) = |\tau|^{-2}Im\tau$, which can be used to show that the correction factor (4.17) is invariant. To check the invariance of the uncompactified bosonic string, one needs furthermore

$$\eta(-\frac{1}{\tau}) = \sqrt{-i\tau} \eta(\tau).$$

The reader who checks this will notice that again $d = 26$ in (4.13) is essential.

It is useful to know that all of this extends in several ways. First of all there is an extension to lattices with different left and right dimensions, which appear in compactifications of heterotic strings, and also in the covariant lattice construction to be discussed in the following. The derivations are completely analogous, and the only difference is that in (4.21) one gets a modular weight factor which has different powers of τ and $\bar{\tau}$. Since these lattice functions appear in the partition functions with different powers of $\eta(\tau)$ and $\eta(\bar{\tau})$, that is no problem.

The extra argument $\vec{z}$ which we added to P in the derivation given above is of interest in its own right. It defines a more general *character valued* partition function, which has been used to prove anomaly anomaly cancellation. We refer to [19] for an extensive discussion of these issues.

Finally we want to point out that there is a straightforward extension of the correction factor to higher loops. The η -functions are replaced by the determinant of the laplacian on higher genus Riemann surfaces. The modular properties of this object are such that the uncompactified bosonic string is modular invariant for any genus. For the correction factor we can write down an explicit expression (see e.g. [36])

$$C_g(\Gamma) = \sum_{p_i \in \Gamma} \exp(i\pi p_R^i \Omega_{ij} p_R^j - i\pi p_L^i \bar{\Omega}_{ij} p_L^j) \det(Im\Omega)^{\frac{N}{2}}.$$

Although we will not do very much with this object in these lectures, it is good to know that it can be shown, by very similar techniques, to be modular invariant [37]. Since the construction of fermionic strings which follows is heavily based on self-dual lattices, it is very easy to prove that they are also multi-loop modular invariant.

5. Conformal Field Theory and Bosonization

5.1 GENERALITIES

In this section we recall elementary concepts of conformal field theory [38] [39] which will be relevant in our context. The list of more detailed reviews contains [40–45]. We emphasize that conformal field theory is *a priori* a more general notion than string theory. Rather, strings moving in a specific d -dimensional space-time represent particular examples of conformal field theories. We will of course restrict the discussion to that types of (free) conformal field theories we need for our particular construction of heterotic strings.

The basic objects are *conformal* or *primary fields* $\phi(z, \bar{z})$, which are defined to transform as follows under coordinate reparametrizations,

$$\phi(z, \bar{z}) = \left(\frac{\partial w}{\partial z} \right)^h \left(\frac{\partial \bar{w}}{\partial \bar{z}} \right)^{\bar{h}} \phi(w, \bar{w}), \quad (5.1)$$

such that $\phi(z, \bar{z})(dz)^h(d\bar{z})^{\bar{h}}$ is conformally invariant. The complex variable parametrizes either the world-sheet cylinder continued to euclidean space (in which case we use instead the variable $w = 2\sigma_0 - 2i\sigma_1$) or the conformally equivalent punctured complex plane (with $z = e^w$). Fields on the cylinder and on the plane are related by

$$\phi_{cyl}(z, \bar{z}) = z^h \bar{z}^{\bar{h}} \phi_{plane}(w, \bar{w}). \quad (5.2)$$

Any conformal field is thus characterized by a *conformal weight* $(h, \bar{h})$; the scaling weight is then given by $h + \bar{h}$ and the two-dimensional spin by $h - \bar{h}$. We will call fields with half-integer spin two-dimensional fermions, those with integer spin bosons (there exist however also objects which are neither bosons nor fermions). If $\bar{h} = 0$ ($h = 0$), a field is called *chiral* or *right-(left-)moving*; it depends then only (anti-)analytically on z . In the following, we will mainly be concerned with chiral fields. All relevant (local) properties of a given conformal field are characterized by operator product expansions with arbitrary other fields:

$$\phi_i(z) \cdot \phi_j(w) = \sum_k (z - w)^{-h_i - h_j + h_k} \alpha_{ijk} \phi_k(w). \quad (5.3)$$

The non-zero coefficients α_{ijk} define the *fusion rules*. Unless all powers in $(z - w)$ in all possible operator products are integral ($h_k - h_j - h_i \in \mathbf{Z}$), the theory is ill-defined (*non-local*), that is, the correlation functions cannot be unambiguously integrated because of branch cuts. As we will see, imposing locality is a quite strong constraint.

The conformal weight h of any field ϕ_h is determined by the operator product with the stress-energy tensor $T = T_{zz}$, as it generates (5.1):

$$T(z) \cdot \phi_h(w) = h \frac{\phi_h(w)}{(z - w)^2} + \frac{\partial_w \phi_h(w)}{(z - w)} + \text{regular terms}. \quad (5.4)$$

Conformal fields have the special property that they obey (5.4) without creating higher poles in $(z - w)$. For example, the fields (basically derivatives of ϕ_h) appearing in the

non-leading and regular terms in (5.4) are not conformal fields. Quite generally, taking the operator product (5.3) of any two conformal fields creates a series of other operators, of which however only a part are conformal fields.

$T(z)$ itself has weight $h = 2$,

$$T(z) \cdot T(w) = \frac{\frac{1}{2}c}{(z-w)^3} + 2 \frac{T(w)}{(z-w)^2} + \frac{\partial_w T(w)}{(z-w)} + \text{regular terms} , \quad (5.5)$$

but for $c \neq 0$ it is not a conformal field due to the 'anomalous' higher pole. To understand the meaning of the parameter c , let us expand $T(z)$ in modes

$$T(z) = \sum_n z^{-n-2} L_n \quad (5.6)$$

according to the general rule

$$\phi_h(z) = \sum_n z^{-n-h} \phi_{h,n} , \quad \phi_{h,n} = \frac{1}{2\pi i} \oint dz z^{n+h-1} \phi(z) \quad (5.7)$$

for expansions on the plane. The values of n must be such that the integral is well-defined. For example if $h \in \mathbb{Z}$ and $\phi(z)$ is single valued around the origin, then n takes all integral values. Contour integrating (5.5) leads now to the following equivalent expression:

$$[L_n, L_m] = \frac{c}{12} n(n^2 - 1) \delta_{n,-m} + (n-m) L_{n+m}. \quad (5.8)$$

Thus, c represents a central charge (anomaly) in the Virasoro algebra, and the zero mode L_0 is precisely one-half the Hamiltonian of the right-moving (analytic) sector of the theory (of course, the full theory contains in addition an isomorphic Virasoro algebra in the left-moving sector).

Quite generally, as above, the information encoded in operator product expansions is equivalent to the collection of all (anti-)commutation relations among the modes; operator products are just an economic way for writing these.

One of the most important fields is the derivative of a free boson, ∂X , which has $h = 1$. An example of X is the string functional described in sect. 2.1, mapped to the complex plane (called the *Fubini-Veneziano field*). X is itself not a conformal field, a fact which is related to its highly non-local operator product with itself (basically the two-dimensional propagator on the complex plane):

$$X(z) \cdot X(w) = -\log(z-w). \quad (5.9)$$

In contrast, ∂X is a local field:

$$\partial X(z) \cdot \partial X(w) = -\frac{1}{(z-w)^2} + \text{regular terms} . \quad (5.10)$$

For free bosons, the corresponding stress energy tensor is

$$T(z) = -\frac{1}{2} : \partial X \partial X : = -\frac{1}{2} \lim_{w \rightarrow z} \left(\partial X(z) \cdot \partial X(w) + \frac{1}{(z-w)^2} \right), \quad (5.11)$$

where the right hand side indicates what the normal ordering dots mean (similarly,

in the following, products of operators taken at the same point are always implicitly understood to be normal ordered, ie., all singularities are subtracted). Inserting (5.11) into (5.5) one learns that in this case $c = 1$.

Another important type of conformal field is given by $\phi_\lambda = :e^{i\lambda X}:$. From $T(z)\phi_\lambda(w) = \frac{1}{2}\lambda^2 \frac{1}{(z-w)^2}\phi_\lambda(w) + \dots$ follows $h = \frac{1}{2}\lambda^2$. Furthermore, we have from (5.9)

$$\begin{aligned} :e^{i\lambda \cdot X}: (z) \cdot :e^{i\lambda' \cdot X}: (w) &= (z-w)^{\lambda \cdot \lambda'} :e^{i\lambda \cdot X(z) + i\lambda' \cdot X(w)}: \\ &= (z-w)^{\lambda \cdot \lambda'} :e^{i(\lambda + \lambda') \cdot X(w)} (1 + i(z-w)\lambda \cdot \partial X(w) + \dots): \end{aligned} \quad (5.12)$$

Nothing in the above formally changes if we allow $\lambda \rightarrow \lambda^\mu$ and $X \rightarrow X^\mu$ to become vectors (for instance, the space-time coordinates of a string). For $2n$ bosons, $\mu = 1, \dots, 2n$, $T(z) = -\frac{1}{2}\partial X^\mu \partial X_\mu$ and accordingly $c = 2n$.

In *fermionic* string theories, there is a supersymmetry between $2n$ bosons X^μ and $2n$ fermions ψ^μ . The latter obey

$$\psi^\mu(z) \cdot \psi^\nu(w) = \frac{\delta^{\mu\nu}}{(z-w)} + \text{regular terms} , \quad (5.13)$$

which, defining ^{*}

$$\psi^\mu(z) = \sum_{n \in \mathbb{Z} + \frac{1}{2}} z^{-n - \frac{1}{2}} \psi_n^\mu, \quad (5.14)$$

is equivalent to $\{\psi_n^\mu, \psi_m^\nu\} = \delta_{n+m}^{\mu\nu}$. The free fermion stress energy tensor is $T_\psi = \frac{1}{2}(\partial\psi)\psi$, and from (5.13) we have

$$T_\psi(z) \cdot \psi^\mu(w) = \frac{1}{2} \frac{\psi^\mu(w)}{(z-w)^2} + \frac{\partial\psi^\mu(w)}{(z-w)} + \text{regular terms} , \quad (5.15)$$

so that $h = \frac{1}{2}$. Furthermore, the operator product of T_ψ with itself tells that a collection of $2n$ fermions has $c = n$. From these basic building blocks one can construct other conformal fields by considering operator products. For instance, the first regular term in the expansion (5.13) is given by

$$J^{\mu\nu}(z) = : \psi^\mu \psi^\nu : , \quad \mu, \nu = 1, \dots, 2n. \quad (5.16)$$

One finds that the modes $J_m^{\mu\nu} = \frac{1}{2\pi i} \oint dz z^m J^{\mu\nu}(z)$ realize a $\widehat{SO(2n)}$ Kac-Moody algebra; in particular, the zero modes $J_0^{\mu\nu}$ are the usual $SO(2n)$ generators, under which ψ_m^μ transform as vectors. As $J^{\mu\nu}(z)$ is given above in terms of fermions, one calls this a *fermionic* construction of the Kac-Moody algebra. Later we will discuss a *bosonic* version.

* We take ψ here to be single valued on the complex plane. A more detailed discussion of fermion boundary conditions will be given in the next section

A very important operator in superconformal theories, i.e., in fermionic strings, is the superpartner of the stress energy tensor, which has in the simplest case the form

$$T_F(z) = i\psi^\mu \partial X_\mu(z) = \sum_m z^{-m-\frac{3}{2}} G_m . \quad (5.17)$$

It has weight $h = \frac{3}{2}$, and the corresponding equ. (5.4) together with (5.5) and

$$T_F(z) \cdot T_F(w) = \frac{\frac{1}{3}c}{(z-w)^3} + \frac{2T(w)}{(z-w)} + \text{regular terms} \quad (5.18)$$

constitute the $N = 1$ superconformal algebra. Equivalently, the mode components satisfy the super Virasoro algebra. More precisely, since we are dealing here only with analytical fields, we describe a $(0,1)$ (right-moving) supersymmetry. The zero mode $G_0 = \frac{1}{2\pi i} \oint dz z^{\frac{1}{2}} T_F(z)$ is precisely the two-dimensional supercharge; from (5.18) follows

$$\frac{1}{2}\{G_0, G_0\} = L_0 - \frac{c}{24}. \quad (5.19)$$

T_F indeed transforms fermions into bosons and vice versa:

$$\begin{aligned} T_F(z) \cdot \partial X^\mu(w) &= -\frac{i}{(z-w)^2} \psi^\mu \\ T_F(z) \cdot \psi^\mu(w) &= \frac{i}{(z-w)} \partial X^\mu \end{aligned} \quad (5.20)$$

5.2 STATES

The reason why we considered conformal fields is that there is a one to one correspondence between conformal fields ϕ_h of weight h and *highest weight states*[†] $|h\rangle = \phi_h(0)|0\rangle$ of the Virasoro algebra, defined by

$$\begin{aligned} L_0 |h\rangle &= h |h\rangle, \\ L_{+n} |h\rangle &= 0. \end{aligned} \quad (5.21)$$

Thus, h basically is the mass of the state $|h\rangle$. Such highest weight states characterize representations of the Virasoro algebra. The other states in a given representation (the *descendants*, which complete the *Verma module*) are obtained by acting with products of L_{-n} to $|h\rangle$; the corresponding operators (e.g., derivatives of ϕ_h) are however not conformal fields, as they create higher poles in (5.4).

Physical states in a string theory are generated by certain combinations of conformal fields called *vertex operators*. They describe the emission of states from the world-sheet,

[†] $\phi_h(0)$ on the plane corresponds to an incoming state at $t = -\infty$ on the world-sheet cylinder.

i.e., the in- or out-coming asymptotic states. For open strings, they have the typical form

$$V(z) = \phi_h(z) e^{ikX}(z) . \quad (5.22)$$

In order to produce conformally invariant correlation functions $\langle V(z_1) \dots V(z_k) \rangle$, these objects must have conformal weight one. Thus, only if the on-shell condition $\frac{1}{2}k^2 = 1 - h$ is satisfied (plus, in general, additional on-shell polarization conditions to prevent higher poles in (5.4)) $V(z)$ is a good vertex operator and $|\phi, k\rangle = V(0)|0\rangle$ a highest weight state. For instance, the tachyon vertex is given by (5.22) with $\phi(z) \equiv 1$, ($k^2 = -m^2 = 2$), and the vector emission vertex in the bosonic string has $\phi(z) \equiv \xi^\mu \partial X_\mu(z)$ with $k^2 = \xi^\mu k_\mu = 0$.

In superconformal theories, matters are a little bit more subtle. Let us consider for a while a fermionic string theory involving a collection of world-sheet fermions ψ^μ in addition to the bosons, where μ is considered a space-time index: $\mu = 1, \dots, 2n$. In addition to states created by purely bosonic operators (i.e., by ∂X and exponentials of X), there exist also highest weight states of the form (see (5.14))

$$(\psi^{\mu_1} \psi^{\mu_2} \dots \psi^{\mu_k})(0)|0\rangle = \psi_{-\frac{1}{2}}^{\mu_1} \psi_{-\frac{1}{2}}^{\mu_2} \dots \psi_{-\frac{1}{2}}^{\mu_k} |0\rangle \quad (5.23)$$

corresponding to antisymmetric tensors of rank k ($h = \frac{1}{2}k$). Taking $\phi_h = \psi^{\mu_1} \psi^{\mu_2} \dots \psi^{\mu_k}$ in (5.22) yields (roughly) vertex operators describing the emission of antisymmetric tensor states; in particular,

$$V = \xi^\mu \psi_\mu e^{ikX} \quad (5.24)$$

is related to the vector boson emission vertex. A quite useful object is the G -parity operator

$$G_{NS} = (-1)^F, \quad (5.25)$$

(F is the two-dimensional fermion number) which is $+1$ (-1) on even (odd) antisymmetric tensors (5.23).

The crucial difference to the bosonic case is that the Hilbert space (i.e., the representation space of the super Virasoro algebra) splits into two sectors, called *Neveu-Schwarz* (NS) and *Ramond* (R). This is due to the fact that there are two *spin structures* on the cylinder: in the NS -sector the fermions ψ have antiperiodic, while in the R -sector they have periodic boundary conditions. Performing a conformal transformation $z = e^w$ we see that (because of the factor $e^{\frac{1}{2}w} = z^{\frac{1}{2}}$ in (5.2)) fermions in the NS -sector are periodic (single-valued), and those in the R -sector antiperiodic (double-valued) along cycles enclosing the origin of the complex plane. Thus in the NS -sector the mode numbers n of ψ_n are half-integral, and in the R -sector integral. In particular, the zero modes in the R -sector obey

$$\{\psi_0^\mu, \psi_0^\nu\} = \delta^{\mu\nu}, \quad (5.26)$$

so they represent space-time ($SO(2n)$ -) γ -matrices. Accordingly the representation space on which ψ^μ act must be spinorial. Thus, denoting by $|\alpha\rangle$ (α = spinor index) the

ground state in the R -sector, a generic R -state (corresponding to (5.23)) looks like

$$(\psi^{\mu_1} \psi^{\mu_2} \dots \psi^{\mu_k})(0) |\alpha\rangle . \quad (5.27)$$

Clearly, while states in the NS -sector (5.23) correspond to space-time antisymmetric tensors, R -states (5.27) lead to space-time spinors. Like in the NS -sector, one can define a grading defined by the operator

$$G_R = \gamma_{2n+1} \otimes (-1)^F \quad (5.28)$$

with eigenvalues ± 1 (counting space-time helicity).

How can we create states in the R -sector via vertex operators ? If we view $\psi(z)$ as an analytic function on the plane, we can describe the double-valuedness around the origin (in the R -sector) as due to a square-root branch cut emanating from there: it flips the boundary conditions between periodic and antiperiodic. This cut creates stress energy (that is, it breaks translation invariance on the plane), and thus corresponds to an operator insertion at $z = 0$. This operator has to be an ordinary (super-)conformal field $S^\alpha(z)$, such as to generate a superconformal highest weight state:

$$|\alpha\rangle = S^\alpha(0) |0\rangle \quad (5.29)$$

S^α is called a *spin* field; it has a very complicated description in terms of the fundamental field ψ^μ . Basically it is the vertex operator for the emission of space-time spinors. In an unitary theory $G_0^2 \geq 0$, hence it follows from (5.19) that the ground state in the R -sector $|\alpha\rangle$ has weight $h = \frac{c}{24}$:

$$L_0 |\alpha\rangle = \frac{c}{24} |\alpha\rangle . \quad (5.30)$$

We thus encounter a problem: for $|\alpha\rangle$ to be a highest weight state and S^α a conformal field, we need $h = 1$. Furthermore, the property that S^α produces a branch cut with ψ is imposed by requiring that the operator product expansion of $\psi(z)$ with $S^\alpha(w)$ contains half-integral powers in $(z - w)$:

$$\psi^\mu(z) \cdot S^\alpha(w) = \frac{1}{(z - w)^{\frac{1}{2}}} (\gamma^\mu)_\beta^\alpha S^\beta + \text{regular terms} . \quad (5.31)$$

Thus naively the theory is *non-local* and as such ill-defined. We will discuss later how these difficulties can be resolved.

The above-explained construction is a special case of what one calls *twisting*: the R -sector is obtained from the NS -sector by a Z_2 -twist. One can of course consider more general, e.g., Z_N -twists, acting in particular also on the bosons ∂X ; this is precisely what happens in the case of *orbifolds*.

5.3 BASICS OF BOSONIZATION

Bosonization, a feature very special to two-dimensions, simply amounts to expressing fermionic field operators like ψ in terms of bosonic ones. The crucial point is that two conformal field theories are physically equivalent, i.e., indistinguishable, if all correlation functions are identical. Now, (the singular parts of) all correlation functions are characterized by all possible operator product expansions (5.3) between the various fields. Thus bosonizing a theory essentially amounts to expressing any given two-dimensional fermionic operator by a certain bosonic expression in such a way that it reproduces all local operator product expansions. In particular, the realization of certain symmetries (e.g., generated by $J^{\mu\nu}$ or T_F) depends only on operator products involving the symmetry currents and not on the way how these are represented.

To bosonize ψ^μ , $\mu = 1, \dots, 2n$, we first switch to a complex basis by defining n pairs

$$\begin{aligned} \frac{1}{\sqrt{2}}(\psi^1 + i\psi^2) &\rightarrow \psi^1, \\ \frac{1}{\sqrt{2}}(\psi^1 - i\psi^2) &\rightarrow \psi^{-1}, \\ \frac{1}{\sqrt{2}}(\psi^3 + i\psi^4) &\rightarrow \psi^2, \\ \frac{1}{\sqrt{2}}(\psi^3 - i\psi^4) &\rightarrow \psi^{-2}, \end{aligned} \quad (5.32)$$

and so on. Then (5.13) takes the form ($j, k = 1, \dots, n$):

$$\begin{aligned} \psi^k(z) \cdot \psi^{-j}(w) &= \frac{\delta_{kj}}{(z-w)} + J^{k,-j}(w) \\ &+ \text{other terms containing descendant fields.} \end{aligned} \quad (5.33)$$

Using (5.12) it is easy to see that we can express our $2n$ fermions by n bosons H_k in a way such that (5.33) is reproduced:

$$\psi^k = e^{iH_k} c_k, \quad \psi^{-k} = e^{-iH_k} c_{-k}. \quad (5.34)$$

(c_k are certain cocycle-phase generating operators which ensure that fermions with different indices really anticommute; in the following, we often will not write such factors explicitly.) Both conformal systems ($\{\psi^\mu\}$, $\{H_k\}$) have central charge $c = n$; in fact, it is only the total value of c that matters.

One can easily bosonize $J^{\mu\nu}$ by just reading it off from (5.33). Switching to the Cartan-Weyl-basis, we have for the off-diagonal ($k \neq j$) currents

$$\begin{aligned} J^{\pm k, \pm j}(z) &\equiv : \psi^{\pm k} \psi^{\pm j} : (z) \\ &= : e^{\pm iH_k \pm iH_j} : (z) c_{\pm k} c_{\pm j}, \end{aligned} \quad (5.35)$$

and for the diagonal currents, which span the Cartan subalgebra:

$$\begin{aligned} J^{k, -k}(z) &= : e^{iH_k} e^{-iH_k} : (z) \\ &= \lim_{w \rightarrow z} \left(e^{iH_k}(z) e^{-iH_k}(w) - \frac{1}{(z-w)} \right) \\ &= i\partial H_k(z). \end{aligned} \quad (5.36)$$

It is easy to check that these currents represent a bosonic realization of the $\widehat{SO}(2n)$ -

Kac-Moody algebra. Note that the price to pay for bosonization is non-covariance of the formalism: The off-diagonal currents are realized by solitons, i.e., circle-valued bosons, while the diagonal, abelian currents take the canonical form. (5.35) and (5.36) are only a particular example of the more general *Frenkel-Kac* construction (For a detailed discussion we refer to [27].); the latter works for any (simply-laced) Lie algebra, if the root vectors $(\pm e_k \pm e_j)$ of $SO(2n)$ are replaced by the length-squared two root vectors of the algebra in question.

In the last section we have given further examples of operators in the fermionic sector. For instance, in bosonic language the operator corresponding to $\psi^\mu \psi^\nu \dots \psi^\rho$ (5.23) is (a linear combination of operators) of the form

$$\phi_\lambda(z) = e^{i\lambda \cdot H}(z) , \quad (5.37)$$

where λ is an n -dimensional vector of the form $(\pm 1, 0, \pm 1, \dots)$, depending on the precise tensor structure. The G -parity operator (5.25) takes the form

$$G = e^{i\pi(\sum_{k=1}^n \lambda_k)} , \quad (5.38)$$

which is ± 1 depending on if $\sum \lambda_i$ is even or odd. Operators describing (anti-)commuting objects are related to vectors λ of even (odd) length-squares; one can easily see this from equ. (5.12) upon exchanging z and w .

It is clear that the collection of all operators of the form $e^{i\lambda H}$ is related to a *lattice*. This follows simply from the fact that in any operator product (5.12) the vectors add to give a new one: $\phi_\lambda(z) \cdot \phi_{\lambda'}(w) = \phi_{\lambda+\lambda'}(z-w)^{\lambda\lambda'} + \dots$. Consistency, that is, closure of the operator product algebra (5.3), then requires that the vertex operator $\phi_{\lambda+\lambda'}$ exists and generates a state $|\lambda + \lambda'\rangle$ of the theory. The lattice thus is formed by the set of vectors λ obtained by all possible additions of some given basic vectors. (Of course, there exist also vertex operators containing bosons ∂H ; however, these play only a trivial role in the following).

What we are after is a classification scheme for vertex operators or states, in order to characterize the spectrum of a given theory. Given the above lattice, operator products preserve a natural Z_2 -grading, that is, a split into vectors of even and odd length-squares; equivalently, a split into eigenvalues ± 1 of G_{NS} . More specifically, the lattice characterized above is precisely the weight lattice of D_n generated by the conjugacy classes (0) and (v). In particular, the non-zero roots of D_n ($\pm e_i \pm e_j$) which appear in the bosonized currents (5.35) belong to the neutral class (0), while the fermions $\psi^{\pm i}$ belong to (v); the operator product expansion (5.13) of two fermions, which contains $J^{\mu\nu}$ in its regular terms, clearly reflects the rule $(v) + (v) = (0)$.

In this framework it is very easy to describe states in the R -sector. In the R -sector, the fields ψ^μ are twisted, as explained in the previous section. In bosonic language such a twist corresponds to a *shift* of the lattice, a fact that is made plausible by the identity

$$\left(\frac{1}{\prod (1 + q^{2n})} \right) \sum_{n \in \mathbb{Z}} q^{(n+\theta)^2} = q^{\theta^2} \prod_{n \geq 0} (1 + q^{2n+2\theta+1})(1 + q^{2n-2\theta+1}) , \quad (5.39)$$

which is a mathematical manifestation of bosonization: the RHS is the partition function of two fermions with twist θ , while the LHS represents the partition function of one

compact boson taking values on a (one-dimensional) integer lattice shifted by θ . More precisely, the sum at the LHS gives the contribution of all solitonic states created by $\{\phi_\lambda\}$, $\lambda = n + \theta, n \in \mathbf{Z}$, and the product is the contribution of all possible combinations of derivatives of ∂H . The case at hand corresponds to n copies of (5.39), i.e., to $2n$ fermions and to an integer lattice shifted by $\theta_1 = \dots = \theta_n = \pm \frac{1}{2} \bmod 2$. In other words, the R -sector is described by the spinor conjugacy classes (s) and (c) of D_n . To which particular spinor class a given state $|\lambda\rangle$ belongs is determined by the eigenvalue of G_R (5.28) which is, in bosonic language, up to a phase, identical to G_{NS} given in (5.38).

Every given conjugacy class can of course be represented by a particular vector belonging to that class; all other vectors in the same class are obtained by adding the root lattice, i.e., all vectors in the neutral class (0) . Thus all states (up to derivatives) in the R -sector are of the form

$$(e^{i\lambda \cdot H} S^\alpha)(0) |0\rangle, \quad \lambda \in (0) \text{ of } D_n, \quad (5.40)$$

and we can think of the ground states of the R -sector $|\alpha\rangle = S^\alpha(0) |0\rangle$ as being generated by shift vectors of minimal lengths in the spinor conjugacy classes of D_n . Hence we find the following bosonic expression for the spin fields:

$$S^\alpha(z) = e^{i\lambda_s \cdot H}(z), \quad \lambda_s = (\pm \frac{1}{2}, \pm \frac{1}{2}, \dots, \pm \frac{1}{2}). \quad (5.41)$$

In fact, there are two spin fields, S^α and $S^{\dot{\alpha}}$, with opposite $SO(2n)$ helicity belonging to (s) and (c) of D_n ; they are distinguished by an even or odd number of $-$ signs in λ_s , respectively. The number 2^{n-1} of possible sign choices for λ_s in each case is precisely the dimension of the two spinor representations of $SO(2n)$: $\alpha, \dot{\alpha} = 1, \dots, 2^{n-1}$. The conformal weights of $S^\alpha, S^{\dot{\alpha}}$ are $h_s = \frac{1}{2} \lambda_s^2 = \frac{1}{8} n$; since in a supersymmetric theory $n_f = n_b = 2n$, and $c_b = 2c_f = 2n$, we find $h_s = \frac{c}{24}$ in accordance with (5.30).

Many properties of states in the various conjugacy classes can be determined solely in terms of addition and multiplication rules of $(0), (v), (s)$ and (c) , in connection with the basic formula (5.12) for operator products of exponentials. For instance, the twisted boundary conditions for fermions in the R -sector follow from the branch cut arising from $(z - w)^{\lambda \cdot \lambda'}$ in (5.12) by the multiplication rule $(v) \cdot (s) \in \mathbf{Z} + \frac{1}{2}$.^{*} (Anti-)commutation properties are also easily determined from (5.12) upon exchanging z and w ; one finds that while all states in (0) commute and those in (v) anticommute, states in the spinor conjugacy classes (for generic n) have no definite (anti-)commutation properties.

The above discussion dealt only with periodic and antiperiodic fermionic boundary conditions, i.e., with Z_2 -twisted fermions. We have just seen that these can be described, in bosonic language, in terms of the weight lattice D_n of $SO(2n)$. It is obvious that one can easily generalize these concepts to other lattices (for instance, the weight lattices A_n of $SU(n+1)$). (One can, of course, consider also lattices which are products of weight lattices of several Lie groups, and/or lattices which are not at all related to Lie groups.) One then describes fermions with more complicated boundary conditions, and other objects, which need not be either fermionic or bosonic. Like the spin fields,

^{*} This is nicely illustrated by (5.31), which also reflects the fusion rule $(v) + (s) = (c)$.

such operators are quite difficult to describe in terms of the fermionic variables ψ^μ . In bosonic language, however, all operators have the same (exponentiated) form, and ψ^μ are no more fundamental than S^α . Clearly, the bosonic lattice formulation provides the most simple and elegant description of such objects.

5.4 COMPACTIFICATION REVISITED

We have seen that certain conformal fields (like ψ^μ , $J^{\mu\nu}$ and spin fields) can easily be described in terms of free compact bosons. In fact, it is sometimes possible to interpret these as (torus-)compactified higher dimensional coordinates.

Consider, for example, torus compactification (as discussed in chapter 4) of the 26-dimensional closed bosonic string on an 8-dimensional even lorentzian self-dual lattice $(E_8)_L \times (E_8)_R$. This is the same as $(D_8)_L \times (D_8)_R$ with conjugacy classes $(0,0)$, $(0,s)$, $(s,0)$ and (s,s) . Using the compactified coordinates $X^I(z)$, $X^I(\bar{z})$ in the left- and right-moving sectors, it is easy to discuss the conformal field theory of this model in terms of operators like the currents of $(E_8)_L \times (E_8)_R$.

The currents belonging to (0) of the subgroup $D_8 \sim SO(16)$ are precisely given by (5.35) and (5.36); the currents (5.36) belonging to the Cartan subalgebra correspond to Kaluza-Klein states, the off-diagonal ones to winding states, i.e., solitons. The remaining currents belonging to (s) , i.e., to $E_8/SO(16)$ are realized by spin fields (5.41) of $SO(16)$. This is possible since a spinor weight of D_8 has length-squared two, so that it can represent a root vector (we say that the spinor conjugacy class of D_8 extends D_8 to E_8). Now, what is relevant in the two dimensional sense are the properties of the conformal fields of the theory, and not if they can be interpreted in terms of compactified higher dimensions. According to the fermion-boson equivalence in two dimensions, in order to describe the same theory we could also start with a fermionic version of the above objects. The $SO(16)$ currents are then represented by bilinears as in (5.16), however, the currents represented by the spin fields have a very complicated, transcendental dependence on the fermions [46]). In this formulation, there is no trace of higher dimensions left! Apparently the concept of compactification is quite delicate in the string context.

The above example represents a compactified theory in the very sense. There exist however more general conformal field theories, which do not admit a compactification interpretation (we will encounter such theories in the next chapter). One should therefore not confuse compact bosons (equivalent to fermions) with compactified space-time coordinates (which are a special case).

For closed strings, which are described by a combination of a right-moving (z -dependent) and a left-moving ($\bar{z}$ -dependent) conformal field theory, we require vertex operators to have conformal weight $(\bar{h}, h) = (1, 1)$. Consequently, these consist of a left- and a right-moving piece together with the exponentiated momentum,

$$V(\bar{z}, z) = \phi_h(z) \phi_{\bar{h}}^l(\bar{z}) e^{ik \cdot X}(\bar{z}, z), \quad (5.42)$$

with the condition $\bar{h} = h$. ϕ_h and $\phi_{\bar{h}}^l$, belonging to independent left- and right-moving sectors, are a priori not related to each other. For instance, for the compactified bosonic

string above,

$$V_{\mu}^{ab}(\bar{z}, z) = J^{ab}(\bar{z}) \partial X_{\mu}(z) e^{ikX} \quad (5.43)$$

($\mu = 1, \dots, 18$) describes a gauge vector boson belonging to the adjoint representation of the left-moving E_8 with generators J^{ab} , whereas

$$V_{\mu}^{a'b'}(\bar{z}, z) = J^{a'b'}(z) \bar{\partial} X_{\mu}(\bar{z}) e^{ikX} \quad (5.44)$$

describes the emission of a vector boson of the right-moving E_8 . On the other hand,

$$V_{\mu\nu}(\bar{z}, z) = \bar{\partial} X_{\nu}(\bar{z}) \partial X_{\mu}(z) e^{ikX} \quad (5.45)$$

describes the emission of gravitons, antisymmetric tensor fields and dilatons in $d = 18$.

In the next section we will apply the above ideas to a bosonic description of heterotic strings, which in general do not admit a compactification interpretation.

6. NSR Model, Superstring and Heterotic String

6.1 THE COVARIANT NSR MODEL IN BOSONIC LATTICE FORMULATION

We will first introduce the covariant ten-dimensional *NSR* model [47] (the *spinning string*) in bosonic language, which will allow a straightforward generalization to lower dimensions in the next chapter.

The covariant *NSR* model can be described by a specific conformal field theory whose main variables are ten bosons X^{μ} and ten fermions ψ^{μ} , which play the role of space-time coordinates. Thus, the basic conformal operators are $\partial X^{\mu}(z)$ and $\psi^{\mu}(z)$, whose properties have been discussed already in the foregoing chapter. Since these fields depend *analytically* on z , we deal only with right-moving fields. What we are after is however closed, in particular, heterotic string theory. Thus we will eventually have to pair the right-moving *NSR* model with an a priori independent left-moving ($\bar{z}$ -dependent) conformal field theory. This will be explained further below.

On a flat background, and in superconformal gauge, the action takes the free form

$$S_{NSR} = -\frac{1}{2\pi} \int d^2z \left(\partial X^{\mu} \bar{\partial} X_{\mu} - \psi^{\mu} \bar{\partial} \psi_{\mu} + \mathcal{L}_{ghost} \right). \quad (6.1)$$

Let us denote by c and b the (anticommuting) ghost and antighost fields for fixing conformal invariance, and by γ and β the (commuting) ghost and antighost fields for fixing the fermionic part of superconformal invariance. Then the ghost lagrangian is given by

$$\frac{1}{2} \mathcal{L}_{ghost} = b \bar{\partial} c + \beta \bar{\partial} \gamma. \quad (6.2)$$

The conformal field theory of the X^{μ} and ψ^{μ} system was developed in the foregoing chapter; we indicated already, that this part of the theory by itself suffers from severe deficiencies. For example, ψ^{μ} as a two-dimensional spinor anticommutes with itself

(cf.(5.13)). Since it carries however a space-time vector index (it appears in emission vertices for space-time tensors), one would rather like to have it generating space-time bosons. Moreover, in order to describe the R -sector, we had to introduce the spin field S^α . This field has conformal weight $h = \frac{n}{8} = \frac{5}{8}$ and not $h = 1$, what would be needed for a vertex operator describing massless space-time fermions. Inserted at some point on the complex plane, S^α creates a branch cut (5.31) in order to change the boundary conditions of ψ^μ (in bosonic formulation, this is related to the rule $(v) \cdot (s) \in \mathbf{Z} + \frac{1}{2}$ for the conjugacy classes of D_5 , which is the Lie algebra of the Wick-rotated Lorentz group in ten dimensions). But the very presence of this cut makes the theory non-local. Another problem we encounter is that S^α does not anticommute with itself as it should as a space-time fermion. This fact is reflected by the rule $(s) \cdot (s) \in 2n + \frac{5}{4}, n \in \mathbf{Z}$ in D_5 , which also reveals the presence of even worse cuts.

It is important to notice that many of these and other unpleasant features trace back to properties of the various conjugacy classes of the lattice D_5 , if one uses the bosonic description of the ψ^μ . We thus expect that a resolution lies in the modification of this lattice.

The only new ingredient of the NSR model (6.1) compared to the conformal field theory of X^μ and ψ^μ as described in the last chapter is the ghost system. It is in fact a crucial ingredient, which can cure all above-mentioned diseases. To be more precise, let us first display some essential properties of the ghosts: $b(z) \cdot c(w) = c(w) \cdot b(z) = \frac{1}{(z-w)}$, $\beta(z) \cdot \gamma(w) = -\gamma(w) \cdot \beta(z) = -\frac{1}{(z-w)}$. Their conformal weights are

$$\begin{aligned} c : \quad & h = -1, \\ b : \quad & h = 2, \\ \gamma : \quad & h = -\frac{1}{2}, \\ \beta : \quad & h = \frac{3}{2}, \end{aligned} \tag{6.3}$$

The contribution to the full energy momentum tensor is $T_{gh} = c\partial b + 2\partial cb - \frac{1}{2}\gamma\partial\beta - \frac{3}{2}\partial\gamma\beta$. Substituting this into (5.5), we learn that the contribution to the central charge of the b, c system is $c_{bc} = -26$, and similarly $c_{\beta\gamma} = 11$; thus altogether we have $c_{ghost} = -15$ and this cancels precisely the contribution of the ten X^μ plus ten ψ^μ . So the conformal anomaly disappears, which seems to be crucial for the consistency of the theory. This does of course not imply that the string theory can only be consistent in ten dimensions. Rather, what counts here is only the *total* value of $c_{matter} = 15$, and not any particular decomposition in terms of contributions of X^μ and ψ^μ . Allowing for internal degrees of freedom, which carry 'missing' central charge, will allow us later to construct string theories in lower dimensions than ten.

As we intend to discuss only the spectrum of the theory and not other aspects (like correlation functions), we will explain only aspects of the ghost system relevant here. The role of the b, c system is then quite simple: For our purposes, all what the b, c system does is to provide the correct vacuum energy. In sect. 5.2 we described how the states of the theory are created by acting with conformal fields on some vacuum. We did however not specify the vacuum itself. The correct vacuum (of the bosonic string) is simply $|0\rangle = c(0)|0\rangle_{SL(2,C)}$, where $|0\rangle_{SL(2,C)}$ is the true $SL(2, C)$ invariant vacuum of

the theory, which is annihilated by $L_n, n \geq -1$. (Another way of describing the same situation is to redefine the vertex operators of the theory $V \rightarrow cV$; these objects then act on the $SL(2, C)$ invariant vacuum state.) Since c has weight $h = -1$, the level of $|0\rangle$ is -1 ; it represents the tachyon state of the bosonic string. Massless states, therefore, are build upon $|0\rangle$ using conformal fields of weight $h = 1$. This gives us back the mass formula of the bosonic string (4.10) ($\mathcal{N}$ counts the number of derivatives in a vertex operator).

In fermionic strings, we have in addition the commuting ghosts β, γ . Their role is much more involved. As we cannot discuss all issues in great detail, we will just give recipes for handling these ghosts (details can be found e.g. in [39]). It is very convenient to bosonize also these fields:

$$\beta = e^{-i\phi} \partial \xi, \quad \gamma = e^{i\phi} \eta. \quad (6.4)$$

Here, ϕ is a chiral boson with opposite norm, $\phi(z) \cdot \phi(w) = \log(z - w)$, and ξ and η are auxiliary fermions, which will play no role in the following. They are needed above since one cannot bosonize the commuting fields β, γ solely in terms of an anticommuting exponential of one boson.

One important feature of the ϕ -ghost is that the conformal field $e^{iq\phi}$ has a peculiar conformal weight:

$$\begin{aligned} T_{\beta\gamma}(z) \cdot e^{iq\phi}(w) &= -\frac{1}{2}q(q+2) \frac{e^{iq\phi}(w)}{(z-w)^2} + \dots \\ \rightarrow h &= -\frac{1}{2}q(q+2). \end{aligned} \quad (6.5)$$

The linear term $(-q)$ originates in an anomaly of the ghost number current $j = \beta\gamma$, and the negative sign is due to the negative metric of ϕ , which also implies

$$e^{iq\phi}(z) \cdot e^{ip\phi}(w) = (z-w)^{-qp} e^{i(q+p)\phi}(w) + \dots \quad (6.6)$$

Similar to the b, c system, also in the β, γ system the vacuum gets modified: it is characterized by a certain *bose ghost sea charge* q :

$$|0\rangle_{\beta\gamma} \equiv |q\rangle = e^{iq\phi}(0) |0\rangle. \quad (6.7)$$

To specify the possible values of q , note first that the $\beta\gamma$ ghosts are related to gauge degrees of the (non-dynamical) world-sheet gravitino, hence should have the same periodicity properties. Since that couples to the supercurrent $T_F = i\psi\partial X$, it follows that because of world-sheet supersymmetry the $\beta\gamma$ ghosts should have the same boundary conditions as the NSR fermions ψ^μ . This implies that we need *different* vacua in the NS - and R -sectors of ψ^μ : in the NS -sector $q \in \mathbf{Z}$ while in the R -sector $q \in \mathbf{Z} + \frac{1}{2}$. There exists now still an ∞ -fold ambiguity in what values of q we actually choose. In fact, all values differing by integers are equivalent. Thus, for several reasons, let us choose as ‘canonical’ ghost vacua

$$\begin{aligned} |0\rangle_{\beta\gamma}^{NS} &= |-1\rangle, \\ |0\rangle_{\beta\gamma}^R &= |-\frac{1}{2}\rangle. \end{aligned} \quad (6.8)$$

Later we will comment on other choices.

Similar to the b, c system, it is sometimes more convenient (in particular, for computing correlation functions) to redefine the vertex operators so that they act on a simple vacuum. Let us take here $|0\rangle = c(0)|0\rangle_{SL(2,C)}$ with level -1 . Then

$$V_{NS} e^{-i\phi}(z) \quad \text{and} \quad V_R e^{-i\frac{1}{2}\phi}(z) \quad (6.9)$$

are the vertex operators we will consider from now on. By (6.5), the mass formula reads

$$\begin{aligned} \frac{1}{8}m^2 &= h - 1, \\ h &= \frac{1}{2}\lambda^2 + \mathcal{N} - 1 + \begin{cases} \frac{1}{2} & NS\text{-sector} \\ \frac{3}{8} & R\text{-sector} \end{cases} \end{aligned} \quad (6.10)$$

which displays two different normal ordering constants in the NS - and R -sectors. This formula already cures some of our problems: the fermion emission vertices for both chiralities now become

$$V_F^{spinor(s)} = u_\alpha S^\alpha e^{-i\frac{1}{2}\phi} e^{ikX}, \quad V_F^{spinor(c)} = u_{\dot{\alpha}} S^{\dot{\alpha}} e^{-i\frac{1}{2}\phi} e^{ikX}, \quad (6.11)$$

and therefore (if $k^2 = 0$) have weight $h = 1$ (as opposed to $h = \frac{5}{8}$ without ghosts) what is needed for good vertex operators. Similarly, the vector boson emission vertex

$$V_B^{vector} = \xi_\mu \psi^\mu e^{-i\phi} e^{ikX}, \quad (6.12)$$

now also has $h = 1$ (as opposed to $h = \frac{1}{2}$) for vanishing momentum. V_F^{spinor} and V_B^{vector} therefore describe massless particles in ten dimensions. We can, of course, also build other operators like $V_B^{tachyon} = e^{-i\phi} e^{ikX}$ or $V_B^{tensor} = \xi_{\mu\nu} \psi^\mu \psi^\nu e^{-i\phi} e^{ikX}$, with mass $-\frac{1}{2}$ and $+\frac{1}{2}$, respectively. The ground states in the NS - and R -sectors are $|V_B^{tachyon}\rangle$ and $|V_F^{spinor}\rangle$, respectively.

We mentioned above that our other problems could possibly be solved by modifying the lattice D_5 related to the bosonized fermions. Contemplating on (6.9), the clever reader recognizes that part of these problems is solved already. To make this more clear, let us use a trick, invented by [15]: just consider the ghost charge q as part of an enlarged covariant lattice. That is, associate to any of the above (and all other possible) bosonized vertex operators (i.e., states) a six-dimensional vector $w = (\lambda | q)$, e.g.,

$$\begin{aligned} w^{spinor} &= (\pm\frac{1}{2}, \pm\frac{1}{2}, \pm\frac{1}{2}, \pm\frac{1}{2}, \pm\frac{1}{2} | -\frac{1}{2}) \\ w^{vector} &= (0, \dots, \pm 1, \dots, 0, \dots | -1) \\ w^{tachyon} &= (0, 0, 0, 0, 0 | -1) \end{aligned} \quad (6.13)$$

and $w^{tensor} = (0, \dots, \pm 1, \dots, \pm 1, \dots | -1)$, etc. In order to efficiently characterize the theory, let us investigate the nature of this six dimensional lattice Γ_6 . It looks rather like an extension $D_5 \rightarrow D_6$, but we have to take the switched sign in the ghost metric into account. The natural inner product on this lattice is defined by the power of $(z - w)$ in

operator product expansions, because that determines the various (anti-)commutation properties of the fields and measures locality. From (5.12) and (6.6) we have

$$V_\lambda e^{iq\phi}(z) \cdot V_{\lambda'} e^{iq'\phi}(w) = (z-w)^{\lambda \cdot \lambda' - qq'} V_{\lambda+\lambda'} e^{i(q+q')\phi}(w) + \dots \quad (6.14)$$

Therefore the natural inner product has *lorentzian* metric $(++++|-)$:

$$\langle w, w' \rangle = \lambda \cdot \lambda' - qq' \quad (6.15)$$

The same inner product appears also in the mass formula for arbitrary ghost charges:

$$\frac{1}{8}m^2 = \frac{1}{2} \langle w, w \rangle - q + \mathcal{N} - 1 \quad (6.16)$$

Accordingly, we denote the lorentzian lattice by $\Gamma_{5,1} = D_{5,1}$, with the comma indicating the sign flip in the metric.

As the ϕ -charge adds in (6.14), is it clear that for closure of the operator product algebra one needs also vertex operators with different ghost charges than the canonical ones (6.9). However, as mentioned above, vertex operators with ghost charges differing by integral units can represent (or generate) the same physical state. That is, there is a copy of every canonical vertex operator in every ghost sector; the different copies are also called *pictures*. There exists in fact an operation which maps a given vertex operator with ghost charge q into one with $q+1$:

$$V_{q+1}(z) =: \mathcal{P}_{+1} V_q(z) : \quad (6.17)$$

The *picture changing operator* carries one unit of ghost charge, and is given by

$$\mathcal{P}_{+1}(z) = e^{i\phi} T_F(z) = ie^{i\phi} \psi \cdot \partial X(z) + \dots \quad (6.18)$$

Thus, apart from the ghost part, picture changing is more or less the same as performing a two dimensional supersymmetry transformation^{*}.

For instance, picture changing the vector emission vertex V_B^{vector} in the (-1) -picture (6.12) leads to its copy in the (0) -picture:

$$:\mathcal{P}_{+1} V_{B(-1)}^{vector}: (z) = V_{B(0)}^{vector}(z) = \xi \cdot [\partial X + i\psi(k \cdot \psi)] e^{ikX}(z) \quad (6.19)$$

which (apart from the ψ -dependent piece) is the same as the vector emission vertex of the bosonic string (5.43). One easily checks that the mass formula (6.16) gives the same answer for both $V_{B(-1)}^{vector}$ and $V_{B(0)}^{vector}$. This already follows from the fact that the conformal weight h of $\mathcal{P}_{+1}(z)$ vanishes. In general, $\mathcal{P}_{+1}$ always acts in a way that compensates the effect of changing the q -charge. Thus the mass spectrum of the theory is bounded from below, a subtle feature which is not at all manifest in the mass formula (6.16).

* In fact, the ghost part is necessary in order to have a well-defined local action of T_F on states in the R -sector; remember the branch cut of ψ in T_F . Roughly speaking, one may regard $e^{i\phi}$ as a bosonized anticommuting coordinate, playing the role of ϵ in $[\epsilon Q^{SUSY}, \dots]$.

Upon bosonization, we can represent $\mathcal{P}_{+1}$ by the vector $w^{PC} = (0 \dots, \pm 1, 0 \dots | +1) \in D_{5,1}$. Picture changing then acts on the lattice(-part of the theory) simply as shifting by w^{PC} . In order for this operation to be well-defined and local, w^{PC} has to have integral inner product with all other lattice vectors. Although trivially satisfied in the ten dimensional *NSR* model, this condition will prove quite non-trivial for the lower dimensional theories we will consider below, as for these theories the supercurrent T_F (i.e., w^{PC}) can take a different form.

The structure of (6.19) is more complicated than that of (6.12) (and even more complicated in higher pictures). In general, the canonical ghost sectors (6.9) have the virtue of being simplest in the sense that the vertex operators are related to lattice vectors in the most simple way. In the other ghost sectors, there is no manifest relation between lattice vectors and physical vertex operators; they are in general given by more complicated linear combinations. For our purposes, however, these complications play no role, as features like locality or (anti-)commutativity do not depend on them. That is, all relevant information about state classification is carried by the lattice vectors (oscillators, i.e., derivatives, also play no role). For our purposes, we can therefore represent the spectrum of vertex operator in all ghost sectors by the lattice $D_{5,1}$.

This certainly does not mean that the partition function of the physical states is given by the partition function of the lattice! There is a priori no one-to-one relation between lattice vectors and physical states[†]. There exists however a brute force method to count the physical degrees of freedom correctly: let us split every lattice vector $w = (\lambda | q) \in D_{5,1}$ into a four dimensional 'light-cone' (describing the Wick-rotated transverse Lorentz group $SO(8)$) and a two dimensional 'ghost' part, according to the decomposition

$$D_{5,1} = D_4^{light-cone} \times D_{1,1}^{ghost} \quad (6.20)$$

$$w = (u, x), \quad u \in D_4^{light-cone}, \quad x \in D_{1,1}^{ghost}.$$

Then, by definition, the physical states are those associated with

$$x = x_0 = \begin{cases} (0 | -1) & NS\text{-sector} \\ (-\frac{1}{2} | -\frac{1}{2}) & R\text{-sector} . \end{cases} \quad (6.21)$$

One easily checks that this *physical state selection rule* is just a projection to light-cone states, in the canonical pictures (the astute reader also notices that for the above choice of x_0 above the mass formula (6.16) turns precisely into the light-cone mass formula). From the standpoint of conformal field theory, this separation of vectors into 'physical' and 'unphysical' ones is quite artificial (and ambiguous), as vertex operators are equally good in arbitrary pictures. The above selection rule is nothing but a help to identify and count the physical states of the theory.

[†] For example, w^{vector} (6.13) associated with $V_{B(-1)}^{vector}$ given in (6.12) describes ten degrees of freedom (which is the correct count), while $w^{vector} = (\dots \pm 1, \dots \pm 1, \dots | 0)$ associated with the picture changed, equivalent vertex operator $V_{B(0)}^{vector}$ (6.19) has 40 components.

Having now everything reduced to the properties of the lattice $D_{5,1}$, let us split it (i.e., the set of vertex operators) into cosets or conjugacy classes, much like we did for the case of D_n in section 5.3. In particular, as it is obvious from (6.13), V_B^{vector} belongs to (0) of $D_{5,1}$, $V_B^{tachyon}$ and V_B^{tensor} to (v) and V_F^{spinor} to (s) or (c), respectively. Any other vertex operator of the theory belongs to one of these four classes, and their lattice parts can be obtained by adding 'root' vectors belonging to the neutral class (0) to the vectors (6.13); these are simply ground state (lowest mass) representatives of the conjugacy classes (note in particular that the action of the picture changing operator $\mathcal{P}_{+1}$ belonging to (0) does not change the class of a vertex operator).

Then, like for D_n , we can represent all (anti-)commutation relations, fusion rules and possible nonlocalities of the *NSR* model in terms of addition and multiplication rules of the conjugacy classes of $D_{5,1}$ (The dot denotes the lorentzian inner product):

$$\begin{aligned}
 (0) + (0) &= (0), & (0) \cdot (0) &\in 2\mathbf{Z} \\
 (0) + (v) &= (v), & (0) \cdot (v) &\in \mathbf{Z} \\
 (0) + (s) &= (s), & (0) \cdot (s) &\in \mathbf{Z} \\
 (0) + (c) &= (c), & (0) \cdot (c) &\in \mathbf{Z} \\
 (v) + (v) &= (0), & (v) \cdot (v) &\in \mathbf{Z} \\
 (v) + (s) &= (c), & (v) \cdot (s) &\in \mathbf{Z} + \frac{1}{2} \\
 (v) + (c) &= (s), & (v) \cdot (c) &\in \mathbf{Z} + \frac{1}{2} \\
 (s) + (s) &= (0), & (s) \cdot (s) &\in \mathbf{Z} \\
 (c) + (c) &= (0), & (c) \cdot (c) &\in \mathbf{Z} \\
 (s) + (c) &= (v), & (s) \cdot (c) &\in \mathbf{Z} + \frac{1}{2}
 \end{aligned} \tag{6.22}$$

Note that all length(-squares) are integral, but not even; $D_{5,1}$ is therefore an *odd* lattice. (6.22) tells us that operators in (0) (even lengths) commute, those in (s), (c) (odd lengths) anticommute with each other. This is precisely what we like to have, since (0) contains (odd-rank antisymmetric) tensors, and (s), (c) spinors in space-time. Furthermore, $V_B^{vector} \in (0)$ and $V_F^{spinor} \in (s)$ are now local with respect to each other: the branch cut in $\psi(z) \cdot S(w)$ is cancelled by that of $e^{-i\phi}(z) \cdot e^{-i\phi/2}(w)$.

However, we are not yet completely done: the (even-rank antisymmetric) space-time tensors belonging to (v) anticommute and thus violate spin-statistics, and they are also non-local with respect to the spinors (as can be inferred from the $\frac{1}{2}$'s in (6.22)). Finally, the class (v) contains the tachyonic ground state of the *NS*-sector —an unpleasant feature. Now, what we have constructed so far is the spinning string, which is known to be inconsistent. One has to do little more to really obtain the superstring.

6.2 SUPERSTRING AND HETEROTIC STRING

The remaining diseases of the spinning string are simply eliminated by restricting to a subset of the possible vertex operators, characterized by (0) and (s) (or (c)). One easily concludes from (6.22) that all operator products are local and close within this subset. The spin-statistic relation is satisfied as (v) is absent now: tensors and spinors are related to even and odd vectors on $D_{5,1}$. In particular, as massless states we keep $|V_B^{vector}\rangle$ and $|V_F^{spinor}\rangle$ which form a supersymmetry multiplet. Actually one can prove (we will do this explicitly below) that the whole spectrum is supersymmetric. Accordingly, one calls this theory the superstring.

Keeping only states in (0) and (s) can also be expressed as projecting onto states $|w\rangle$ with 'positive G -parity' $G = 1$, where

$$G|w\rangle \equiv (-1)^{F+F^\phi}|w\rangle = e^{2\pi i \langle w^{spinor(s)}, w \rangle} |w\rangle \quad (6.23)$$

generalizes (5.38) by including the ϕ -ghost charge F^ϕ . This is called *GSO-projection* [48].

The *GSO*-projected lattice is smaller in the sense that it contains less conjugacy classes than the lattice we had before. Nevertheless we still will call it $D_{5,1}$, as according to usual notation a lattice is named after its root sublattice (0). (Remember that in order to characterize a lattice completely, one has to provide in addition a list of the conjugacy classes actually present.)

In fact, the projected lattice $D_{5,1}$ is *self-dual* as one can easily check. This self-duality is actually necessary for the consistency of the superstring: it ensures modular invariance of the partition function, which contains a lattice factor

$$\mathcal{L}_{5,1}(\tau) = \sum_{\substack{w=(u,x) \in D_{5,1} \\ x=x_0}} e^{i\pi\tau u^2} e^{2\pi i q} . \quad (6.24)$$

The sum is only over the states satisfying the selection rule (6.21), and the phase counts space-time fermions and bosons with opposite signs. The lattice function $\mathcal{L}_{5,1}$ differs in further ways from lattice partition functions (4.18) we encountered earlier. In particular, the lattice is odd rather than even, and in writing the exponent above we used the peculiar mass formula (6.16). It was shown in [16] that all these features conspire in precisely the right way to make the theory modular invariant. In the next section we will present an independent proof, which furthermore extends to higher loops.

Knowing this, we could have started already with a self-dual lattice from the very beginning and would never had encountered the problems of the spinning string; locality is guaranteed by the fact that on a self-dual lattice all inner products are integral. This will be our philosophy for constructing further string theories.

We have now everything prepared for generalizing the above to heterotic strings. We simply have to combine the above-explained, right-moving *NSR* model (not necessarily the superstring!) with a left-moving bosonic string. As the conformal anomaly $c_{bc} = -26$

for the bosonic string, we need in addition to the ten space-time coordinates X^μ 16 compact bosons $X^I(\bar{z})$ (or equivalently, 32 fermions):

$$S_{het} = -\frac{1}{2\pi} \int d^2z \left(\partial X^\mu \bar{\partial} X_\mu + \psi^\mu \bar{\partial} \psi_\mu + 2b\bar{\partial}c + 2\beta\bar{\partial}\gamma + \partial X^I \partial X_I + 2\bar{b}\partial\bar{c} \right). \quad (6.25)$$

($I = 1 \dots 16$). What is the momentum lattice of the compact bosons like? It is convenient to combine this 16-dimensional lattice Γ_{16} with that related to the right-moving *NSR* model:

$$\Gamma_{16;5,1} \equiv (\Gamma_{16})_L \times (D_{5,1})_R. \quad (6.26)$$

We cannot strongly enough emphasize that this does not necessarily represent a direct product decomposition! As (6.26) denotes only the root sublattices, there may still exist non-trivial correlations between the conjugacy classes of Γ_{16} and $D_{5,1}$. In order to characterize (6.26), let us consider (in analogy to (6.14)) the operator product of two closed string vertex operators (5.42):

$$V_w(\bar{z}, z) \cdot V_{w'}(\bar{w}, w) = (\bar{z} - \bar{w})^{w_L \cdot w'_L} (z - w)^{\lambda \cdot \lambda' - qq'} V_{w+w'}(w) + \dots,$$

$$V_w(\bar{z}, z) = e^{i w_L^I X^I}(\bar{z}) e^{i \lambda \cdot H} e^{i q \phi}(z), \quad w = (w_L; w_R), \quad w_R = (\lambda | q). \quad (6.27)$$

Upon rewriting the z - and w -dependent part using $(\bar{z} - \bar{w}) = \frac{|z-w|}{(z-w)}$, the condition for locality, i.e., the absence of branch cuts becomes

$$\langle w, w' \rangle \in \mathbf{Z}, \quad (6.28)$$

where the inner product has metric $[(-)^{16}; (+)^5 | (-)]$. Thus, $\Gamma_{16;5,1}$ has to be an integral, lorentzian lattice with respect to this metric. Note however that the two sets of $(-)$ -signs have completely different origins: the $(-)$ -sign in the left-moving sector is the same as that arising in the lattice of a compactified closed bosonic string, as discussed in chapter 4: w_L and w_R play the role of p_L and p_R in sect.4.2, respectively.

The full partition function of such a theory has the form

$$\text{Tr} \left[e^{i\pi\tau H'_R} e^{-i\pi\bar{\tau} H'_L} e^{2\pi i F^\phi} \right] = \frac{1}{\eta^{12} \bar{\eta}^{24}} \sum_{\substack{w=(w_L; w_R) \in \Gamma_{16;5,1} \\ w_R=(\lambda|q)=(u, x_u)}} e^{-i\pi\bar{\tau} w_L^2} e^{i\pi\tau u^2} e^{2\pi i q} \quad (6.29)$$

Modular invariance, as we will see, forces again this lattice to be (odd) self-dual. This does certainly not imply that Γ_{16} and $D_{5,1}$ should separately be self-dual lattices. That would correspond only to the simplest cases: taking in the right-moving sector the odd self-dual lattice $D_{5,1}$ corresponding to the superstring, requires Γ_{16} to be even self-dual. As explained in chapter 3, there exist only two such lattices, namely the root lattice of $E_8 \times E_8$ and the weight lattice of $\frac{Spin(32)}{Z_2}$. These choices lead to the well-known supersymmetric string theories with gauge groups $E_8 \times E_8$ and $SO(32)$ (the vertex operators for the emission of the corresponding gauge bosons are accordingly given by $V^{ab}(\bar{z}, z) = J^{ab}(\bar{z}) V_{B(-1)}^{vector}$).

Allowing for a non-trivial correlation between conjugacy classes, the two factors in (6.26) need not be separately self-dual; only the combined lattice needs to be. This leads to further possibilities of constructing theories like the non-supersymmetric $SO(16) \times SO(16)$ -string [49] [50]: it is given by a lattice $\Gamma_{16;5,1} = (D_8 D_8)_L (D_{5,1})_R$ equipped with a certain list of correlated conjugacy classes. In this list, *all* conjugacy classes of $D_{5,1}$ of the spinning string do occur; however, the problems of the spinning string do not show up due to the particular, non-trivial interrelation with the left-moving sector. In particular, all branch cuts are cancelled by virtue of (6.28).

The above construction generalizes straightforwardly to the torus-compactified heterotic strings as introduced by [4]. In reducing the number of dimensions, the lattice gets larger as we include also the $2n$ torus-compactified left- and right-moving space-time coordinates. More specifically, in $d = 10 - 2n$ dimensions we need to have an odd self-dual lorentzian lattice

$$\Gamma_{16+2n;5+2n,1} = \Gamma_{16+2n;2n} \times D_{5,1} . \quad (6.30)$$

Here, $\Gamma_{16+2n;2n}$ is an even, lorentzian self-dual lattice as described by [4]. There exists a continuous variety of such lattices: for example, four dimensional theories are characterized by 132 parameters. These are related to background values of the various fields, precisely as in the case of the torus-compactified bosonic string described in chapter 4. As $D_{5,1}$ in (6.30) is odd self-dual, it represents the torus-compactified fermionic degrees of the superstring. Therefore this kind of theories is non-chiral.

It is now clear how to construct more general, possibly chiral theories: we simply have to look out for odd self-dual lattices (6.30), where the two factors are correlated in a non-trivial way. These theories then will not, in general, be describable as compactifications in the very sense. What we have to make sure of, of course, is that we correctly assign even and odd vectors to space-time tensors and spinors; in lower dimensional theories, there are then also additional constraints related to world-sheet supersymmetry in the internal sector (this will be explained below in chapters 7 and 8). The structure of the covariant lattice is more general than that of (6.30):

$$\Gamma_{16+2n;5+2n,1} = (\Gamma_{16+2n})_L \times \left(\Gamma_{3n}^{int} D_{5-n,1}^{spacetime} \right)_R \quad (6.31)$$

Having found this simple structure, we can forget now all subtleties related to the superconformal ghost system.

What remains to be proven is the connection between (multiloop) modular invariance of the partition function and self-duality of the lattice (6.31). Before this will be attacked, let us first explain and switch to an alternative lattice formulation which is easier to deal with.

6.3 THE EVEN LATTICE MAP

Although the *covariant lattices* introduced in the previous section specify the fermionic string theory completely, they are not very pleasant to deal with, because of their rather cumbersome metric, and because some of their vectors have odd norm. To recapitulate, what we will be interested in for string theories in $d = 10$ dimensions are odd self-dual lattices of the form $\Gamma_{16;5,1}$, where the six right-moving components belong to one of the four conjugacy classes of $D_{5,1}$. Furthermore the space-time spinors (i.e. spinor conjugacy classes of $D_{5,1}$) have to correspond to the vectors of odd length.

Now we will introduce a map of any such lattice to an even self-dual lattice. This map will have the useful property that it preserves modular invariance at all loop orders. Essentially, what we are going to do is to associate with any heterotic string theory (or type-II theory) a bosonic string theory, by mapping the right (or left) moving superconformal system to a bosonic one. This map works in general only in this direction.

This idea originates from [51], where it was used to argue that the fermionic strings are “vacua” of the bosonic string. Although this remains an attractive idea, this is not our interest here. For us the lattice map will merely be a useful tool for proving certain properties of fermionic strings. The form of the lattice map that we will use appeared first in [52], and in [44] it was shown that it preserves one-loop modular invariance. This was extended to higher loops in [53].

Basically, the idea is very simple. If we decompose one of the lattices of interest with respect to the $D_{5,1}$ conjugacy classes, we see that it consists of four sets of vectors

$$(\Delta_0, 0) + (\Delta_v, v) + (\Delta_s, s) + (\Delta_c, c), \quad (6.32)$$

where the second entry denotes the $D_{5,1}$ conjugacy classes, and the Δ ’s are simply the 16-dimensional vectors associated with them on the right lattice. Now define a new lattice $\Gamma_{16;8}$ by writing down a decomposition identical to (6.32), but with the second entry interpreted as a conjugacy class of D_8 instead of $D_{5,1}$. That is, we map

$$D_{5,1} = D_4^{light-cone} D_{1,1}^{ghost} \longleftrightarrow D_4^{light-cone} D_4^{ghost} = D_8. \quad (6.33)$$

Observe that the old and the new lattice have the same number of conjugacy classes, so that if the old lattice was unimodular, so is the new one. Now compute the list of norms and inner products of the conjugacy classes of D_8 . In comparison with (6.22) one finds that the only change is in the norm of the spinors, which change from odd into even length. But the spinors of $D_{5,1}$ were precisely the odd vectors on $\Gamma_{16;5,1}$. Hence the map preserves unimodularity and all mutual inner products, but it maps all the odd vectors into even ones. The result is a lattice $\Gamma_{16;8}$ which is even self-dual with respect to the metric $[(-)^{16}; (+)^8]$. Thus, by use of this one-to-one map, we are able to find an alternative even lattice formulation of heterotic string theories.

If we extend the lattice with an E_8 factor on the right, we get a $\Gamma_{16;16}$ lattice of that type, on which the bosonic string can be compactified (this can also be achieved by using D_{16} in the above construction, but that is less useful for discussing the spectrum). The

bosonic string compactified on the corresponding torus is all-loop modular invariant. Is this related to the modular invariance of the fermionic string from which it came?

To answer this question, one has to write down the multiloop partition function for the fermionic string, which has not been done yet. However, one usually assumes that as far as its modular transformation properties are concerned, it has the form (including only right-movers)

$$(Det D_{1/2})_\alpha^5 (Det D_{3/2})_\alpha^{-1},$$

where $D_{1/2}$ denotes the Dirac determinant, and represents the contribution of two Majorana Weyl fermions and $D_{3/2}$ is the Rarita-Schwinger determinant (which only has ghost contributions). The subscript α denotes the spin-structure. In general, this combination of determinants is multiplied with some α dependent factors from the right-movers.

The point is now, that the modular transformation properties of this object are known explicitly [54]. Using them, we can show that the following two spin-structure dependent quantities have identical modular transformations at any loop order

$$\begin{aligned} Y_\alpha &= \left(\frac{Det_{1/2}}{Det_{3/2}} \right)_\alpha \\ X_\alpha &= (Det_{1/2})_\alpha^4 \sum_\beta (Det_{1/2})_\beta^8 \end{aligned} \tag{6.34}$$

Thus if one replaces in a fermionic string partition function Y_α by X_α , the new one will be modular invariant if and only if the old one was.

This replacement is precisely what the lattice map achieves. To see that one has to make a simple basis transformation from spin-structure basis to conjugacy class basis. We will not explain this in detail here, but refer to [53]. Details aside, comparing Y_α and X_α the reader will undoubtedly recognize that the first represents the $D_{1,1}$ ghost and longitudinal part of the covariant lattice (cf. (6.20)), while the second represents D_4 conjugacy classes, multiplied with the E_8 partition function.

Let us apply the above to the superstring. As discussed in sect.6.2, the superstring is characterized by a $D_{5,1}$ -lattice with conjugacy classes (0) and (s). By use of (6.33), this can be mapped onto D_8 with the same conjugacy classes, which is the same as E_8 ! Thus, (odd) self-duality of $D_{5,1}^{superstring}$ can be inferred from the (even) self-duality of E_8 . This proves not only multiloop modular invariance of the superstring, but it also shows that (6.23) is the only projection leading to a modular invariant theory, as E_8 is the unique even self-dual lattice in 8 dimensions.

The beauty of the above reasoning lies in the fact that the ghost system is determined by the $N = 1$ conformal system, i.e. by a *world-sheet* property, that is the same for any fermionic string. Hence the replacement of Y by X , or the lattice map, is equally useful for any fermionic string theory, and in any space-time dimension.

So we might as well directly go to four-dimensional strings. As indicated at the end of the previous section, four-dimensional strings are specified by lattices of the form given in the left-hand side of (6.30), but without the restriction that it contains a $D_{5,1}$ root lattice. Since we are in four dimensions, it would be foolish to insist on

ten-dimensional Lorentz-invariance. Instead, we only need to require that $D_{2,1}$ roots be present (the Wick-rotated four-dimensional Lorentz-group is $SO(4)$, with Lie algebra D_2). Thus in $10 - 2n$ dimensions we would be looking for lattices $\Gamma_{16+2n;5+2n,1}$ of the form (6.31), with the requirement that the last $6 - n$ components belong to conjugacy classes of $D_{5-n,1}^{spacetime}$. We will again require that the lattice is odd self-dual, with odd lengths for spinors.

Using precisely the same lattice map, we replace now the $D_{5-n,1}^{spacetime}$ conjugacy classes by $D_{8-n}^{spacetime}$ conjugacy classes, that is

$$\begin{aligned} \Gamma_{16+2n;5+2n,1} &= (\Gamma_{16+2n})_L \left(\Gamma_{3n}^{int} D_{5-n,1}^{spacetime} \right)_R \\ &\longleftrightarrow (\Gamma_{16+2n})_L \left(\Gamma_{3n}^{int} D_{8-n}^{spacetime} \right)_R = \Gamma_{16+2n;8+2n} . \end{aligned} \quad (6.35)$$

Using the above we conclude then that the following statements are equivalent:

- The covariant lattice $\Gamma_{16+2n;5+2n,1}$ is odd self-dual (with odd norms for the spinors, as discussed above).
- The lattice $\Gamma_{16+2n;8+2n} \times E_8$ obtained by means of the lattice map is even self-dual.
- The bosonic string, compactified on the latter lattice is multi-loop modular invariant.
- The fermionic string defined by the odd self-dual covariant lattice is multi-loop modular invariant.

This is all that needs to be said about modular invariance. What corresponds to complicated spin-structure summations in fermionic constructions, is encoded in the simple statement that the lattice must be self-dual.

To some extent, we can drop the covariant lattice altogether, and work with even self-dual lattices. This is unfortunately not possible if one wants to study scattering amplitudes. For this purpose one needs the complete machinery of conformal field theory, and we have not succeeded to cast that into even lattice form. However, the even lattice formulation is extremely useful for discussing the spectrum.

Notice the map (6.35) above is a map between *lattices* (even with different dimensions) and *not* a map between individual vectors. The physical light-cone spectrum is obtained by applying a set of truncation rules to the bosonic string spectrum, which are designed to mimic exactly the corresponding procedure for the covariant, odd lattice. To get these rules, consider the physical state selection rule (6.21). It tell us to keep only states with special components in the $D_{1,1}$ sublattice, belonging to conjugacy classes (v) and (s) of that lattice. Since $D_{1,1}$ is mapped to D_4 by the lattice map, we take the identical conjugacy classes of D_4 to identify the physical states. Thus one physical state is represented by an entire conjugacy class of D_4 . One can choose a representative of the D_4 conjugacy class that makes all this even simpler. Namely, if one chooses some weight vector of 8_s and 8_v to represent the spinor and vector conjugacy class, then the mass of the light-cone states is correctly given by the compactified bosonic string mass formula (4.10), in which one should include the contribution from the D_4^{ghost} components. In

other words, if we make an (arbitrary) choice for the 8_v and 8_s weights, the physical state selection rule (6.21) for covariant lattices is replaced by

$$x = x_0 = \begin{cases} (0, 0, 0, 1) & NS\text{-sector} \\ (\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}) & R\text{-sector} \end{cases} \quad (6.36)$$

in the even lattice formulation ($w = (u, x) \in D_{8-n}^{spacetime}$, $u \in D_{4-n}^{light-cone}$, $x \in D_4^{ghost}$). The physical light-cone spectra are identical.

The above rules were first discovered in [51] and [52], and were a completely *ad hoc* prescription for getting the fermionic string from the bosonic one. Now we see that they do in fact follow from the physical state projection rule explained in sect. 6.1.

7. World-sheet supersymmetry

Up to now we have freely replaced world-sheet fermion by bosons. One has to be a little bit more careful about this, because one may endanger world-sheet supersymmetry. In the *NSR* sector of heterotic strings world-sheet supersymmetry is a manifest (local) $N = 1$ supersymmetry, which transforms the fermions ψ^μ into the bosons X^μ and vice-versa. The transformation is

$$\begin{aligned} \delta_\epsilon \psi^\mu &= i\epsilon \partial X^\mu \\ \delta_\epsilon X^\mu &= -i\epsilon \psi^\mu \end{aligned} \quad (7.1)$$

where ϵ is an infinitesimal grassmannian parameter. This transformation is generated by the operator

$$\mathcal{O} = \oint \frac{dz}{2\pi i} i\epsilon \psi^\mu \partial X_\mu = \oint \frac{dz}{2\pi i} \epsilon T_F, \quad (7.2)$$

in other words, the supersymmetry transformation of any conformal field $\Psi(z)$ is obtained by acting with $\mathcal{O}$

$$\delta_\epsilon \Psi = \mathcal{O} \Psi.$$

One may check that on ψ^μ and X^μ , $\mathcal{O}$ does indeed produce (7.1).

Replacing ψ^μ by bosonic fields does not in itself destroy world-sheet supersymmetry. In two dimensions one can realize supersymmetry entirely in terms of bosons. Indeed, by simply replacing ψ^μ in (7.2) by its bosonized form (5.32) one obtains a generator of world-sheet supersymmetry which is expressed entirely in terms of d bosons X^μ (where d is the number of space-time dimensions) and $\frac{d}{2}$ bosons H_i replacing the fermions.

In constructing four-dimensional strings we do however something more drastic than just bosonization. We allow the bosons related to the internal world-sheet fermions to mix freely with the internal bosons (we will use the word “internal” to indicate those components of ψ and X that do not have space-time indices). Since our philosophy is to construct new string theories directly in four dimensions, we start always with a completely bosonic formulation, *i.e.* in practice we never bosonize anything. Therefore it is not automatically guaranteed that the result respects world-sheet supersymmetry.

Why do we need world-sheet supersymmetry? One could answer this question by stating that world-sheet supersymmetry is an essential part of superconformal field theory, which is the foundation of fermionic string theory, but there is a more explicit way of seeing what goes wrong if one does not have it.

One thing that goes wrong is Lorentz-invariance. World-sheet supersymmetry is used to remove unphysical degrees of freedom from the covariant action of fermionic strings, so as to write them entirely in terms of the physical, light-cone degrees of freedom. Although one can always write down such a light-cone action, it is important that it came from a covariant action. This makes it possible to write down a Lorentz-algebra which does not just generate the transverse rotations of the light-cone states, but extends to the full Lorentz-algebra. This extension involves the familiar generators L^{i-} , whose commutators have to be checked to see if the algebra closes. In supersymmetric strings one of the terms appearing in the expression for L^{i-} is $\oint \psi^i T_F$.

One of the things that can go wrong with Lorentz invariance is that $[L^{i-}, L^{j-}]$ simply does not yield the answer required by the algebra of the Lorentz group. This situation occurs in theories with a conformal anomaly, and can be avoided by choosing the field content so that the conformal anomaly cancels. This is one of the oldest ways of obtaining the critical dimension.

It might appear that nothing prevents us from simply using the Lorentz algebra of ten-dimensional strings, regarding 6 of the eight 8 transverse Lorentz indices as internal. This algebra closes, and it is indeed a valid choice in torus compactifications. However, closure of the algebra is not enough. In addition, one has to require that L^{i-} takes physical states into physical states. Otherwise one faces a problem which is even worse than the conformal anomaly, namely the states of the theory won't even fit into representations of the Lorentz group. (In the presence of a conformal anomaly, the Lorentz algebra does not close, but for the bosonic string and the *NSR* formulation of the superstring it is still possible to assemble the massive states into representations of $SO(d-1)$, the little group of massive particles)

This is indeed precisely what happens, and one sees this most easily in the chiral sector of chiral string theories. At first sight, chirality and string theory are almost contradictory, because one would think that chiral states can always be excited into massive states. Massive chiral states would however violate Lorentz invariance, so a really clever mechanism is needed to ensure that all massive states combine into non-chiral pairs, while the massless ones do not. This is achieved by world-sheet supersymmetry. Indeed, this restriction on massive chiral states turned out to be powerful enough to determine all the conditions of world-sheet supersymmetry for the class of theories considered in [6].

A second way in which world-sheet supersymmetry enters is through the picture changing operator. This operator has the form $\mathcal{P}_{+1} = e^{i\phi} T_F$, where ϕ is the bosonized ghost of section (6.1). It is essential for the calculation of scattering amplitudes, and it too should take physical states into physical states, and have a well-defined action.

The problem of finding a bosonic realization of world-sheet supersymmetry consists thus of two parts

1. Find a purely bosonic expression for T_F that satisfies the supersymmetry algebra.
2. Make sure that it acts properly on all states.

The algebra that the supercurrent T_F must satisfy can be encoded in operator products, and reads (with finite terms omitted)

$$\begin{aligned} T(z)T_F(w) &= \frac{3/2}{(z-w)^2}T_F(w) + \frac{1}{(z-w)^2}\partial_w T_F(w) \\ T_F(z)T_F(w) &= \frac{10}{(z-w)^3} + \frac{2}{(z-w)}T(w) \end{aligned} \quad (7.3)$$

The first of these equations states that it must have conformal weight $\frac{3}{2}$. The second one is just (5.18), with $c = 15$, the conformal anomaly of 10 bosons and 10 Majorana-Weyl fermions. The form of the supercurrent is further restricted because we want it to be Lorentz invariant in d dimensions. This means that it must have the form

$$T_F = \psi^\mu \partial X_\mu + T_F^{int},$$

where the first term is just (5.17), and the second is some expression of conformal weight $\frac{3}{2}$, constructed out of $3n$ bosons. Here n is defined by $d = 10 - 2n$. The $3n$ bosons consist of $2n$ internal X 's and n bosons H_i representing the ψ 's as in (5.34). From now on, let us denote all compact bosons generically by X . Subtracting the space-time part of the supercurrent in (7.3), we see that T_F^{int} has to satisfy it with “10” replaced by $10 - d$, and $T(w)$ by $T(w)^{int}$, the energy momentum tensor of $3n$ bosons.

The first equation is quite easy to satisfy. The general solution has the form

$$T_F^{int} = \sum_{\vec{l}} \vec{B}(\vec{l}) \cdot \partial \vec{X} e^{i\vec{l} \cdot \vec{X}} + \sum_{\vec{t}} A(\vec{t}) e^{i\vec{t} \cdot \vec{X}}, \quad (7.4)$$

where $\vec{l}^2 = 1, \vec{t}^2 = 3$, and $\vec{l} \cdot \vec{B} = 0$. One can verify this by substituting (7.4) into the first equation of (7.3), using the standard energy momentum tensor for $3n$ bosons (see (5.11)).

The second equation in (7.3) imposes more complicated conditions on A and $\vec{B}$, and we do not know its general solution. However, as we will show in a moment, we do know many special solutions. First however we want to address the second point mentioned above, the action on T_F on the states in the theory.

If an operator of the form $\exp(i\vec{w}' \cdot \vec{X})$ acts on a state $|\vec{w}\rangle$ with lattice momentum $\vec{w}$, then it produces a state with momentum $\vec{w} + \vec{w}'$. Earlier in this section we have identified two operators whose action should take physical states into physical states, namely L^{i-} and $\mathcal{P}_{+1}$. If we write the right-moving part of the vectors of the covariant lattice in the form

$$w_R = (\text{internal components, weight of } D_{d/2} \mid \text{ghost charge}),$$

according to the decomposition

$$\Gamma_{16+2n;5+2n,1} = (\Gamma_{16+2n})_L \times \left(\Gamma_{3n}^{int} D_{5-n,1}^{spacetime} \right)_R, \quad D_{5-n,1}^{spacetime} = D_{d/2}^{lorentz} \Gamma_1^{ghost},$$

then we see that the internal part of the supercurrent contains exponentials of the above

form, with $\vec{w}'$ equal to

$$\begin{aligned} w^{L^-} &= (\vec{l}, \vec{v} \mid 0) \text{ or } (\vec{t}, \vec{v} \mid 0) && \text{for } L^- \\ w^{PC} &= (\vec{l}, 0 \mid -1) \text{ or } (\vec{t}, 0 \mid -1) && \text{for } \mathcal{P}_{+1} \end{aligned}$$

($\vec{v}$ denotes a vector weight of $D_{d/2}$.) If adding a vector $\vec{w}'$ to a lattice vector should give another lattice vector, then $\vec{w}'$ should lie on the lattice itself. For the picture changing operator one should furthermore require that its action on all states is *local*, i.e. does not introduce any branch cuts. In other words, if $e^{i\vec{w} \cdot \vec{X}}$ is the vertex operator for some physical state with lattice momentum $\vec{w}$, then we should require that the operator product

$$e^{i\vec{w}' \cdot \vec{X}} e^{i\vec{w} \cdot \vec{X}} = (z - w)^{\vec{w}' \cdot \vec{w}} e^{i(\vec{w}' + \vec{w}) \cdot \vec{X}} + \dots$$

does not contain any fractional powers of $(z - w)$. This implies that $\vec{w}'$ must have integral dot product with all lattice vectors, so that it must lie on the *dual* of the lattice. Fortunately our lattices are self-dual, so that these requirements on $\vec{w}'$ are compatible.

The requirements one gets from L^- and $\mathcal{P}_{+1}$ are also compatible with each other, because in covariant lattices the $D_{d/2}$ components and the ghost charge belong to the same conjugacy class of $D_{d/2,1}$. What we conclude from this is that for every vector $\vec{l}$ and $\vec{t}$ appearing in T_F^{int} the lattice must contain vectors of the form $w_R = (\vec{l}, (v))$ and $(\vec{t}, (v))$, where (v) denotes generically the vector conjugacy class of $D_{d/2,1}$. This describes of course only the right-moving part of the lattice. The left components w_L of these vectors, which we will call *constraint vectors* in the following, must vanish.

There is yet another way to arrive at this conclusion, which we mention here because it uses more directly the requirement of supersymmetry, and can also be used to identify other world-sheet supersymmetries on the lattice. What these conditions are imposing on the theory is that the supersymmetry transformation not only respect the *local* properties of the fields, but also, on Riemann surfaces with non-trivial topology, their *global* properties. In (7.1) that would amount to requiring that both sides of the equation have the same periodicity. In the superstring, for which one only has (7.1), this is simply achieved by choosing the boundary conditions for ϵ equal to those of the *NSR* fermions on any Riemann surface with given spin structure. In bosonic language choosing a spin-structure is replaced by choosing a *soliton sector*, which simply means to consider vacuum with a certain lattice momentum vector $\vec{w}$. States in this sector which are created by vertex operators of the form $\exp(i\vec{w}' \cdot \vec{X})$ pick up a phase $\exp(2\pi i \vec{w}' \cdot \vec{w})$ if one takes σ_1 to $\sigma_1 + \pi$. When one considers one term of the supercurrent acting on it, then such a state is mapped into a state which picks up a different phase when moved around a closed loop. This phase change is $\exp(i\vec{p} \cdot \vec{w} + 2\pi i \phi(\vec{w}))$, where we have denoted the phase of ϵ in the sector $\vec{w}$ as $\phi(\vec{w})$. The vector $\vec{p}$ appearing here represents just one term in the supercurrent (i.e. it is one of the vectors $\vec{l}$ or $\vec{t}$ used above). If there were just one term, one can always adapt $\phi(\vec{w})$ to cancel the phase, so that supersymmetry respects all boundary conditions. If the supercurrent has a second term with a vector $\vec{p}'$, then this cancellation of the phase by $\phi(\vec{w})$ can work if and only if $\vec{p} - \vec{p}'$ has integer dot-product with any $\vec{w}$ on the lattice. But then it must lie on the lattice itself.

This argument tells us thus that the *difference* of any two vectors appearing in the supercurrent must lie on the lattice. This includes also the space-time part of the supercurrent. For that part the vector $\vec{p}$ belongs to the vector conjugacy class of $D_{d/2}$, so that we arrive at the same conclusions mentioned earlier.

There are two simple ways of obtaining supercurrents that satisfy the second equation of (7.3). The first is to construct one out of free fermions, and bosonizing it. Such a free fermion supercurrent has the form

$$T_F(z) = \frac{1}{6} f_{abc} \psi^a \psi^b \psi^c, \quad (7.5)$$

where the ψ 's are $6n$ fermions related to the $3n$ internal bosons by bosonization. If f_{abc} are the structure constants of a semi-simple Lie algebra, then one can check that (7.3) is satisfied [55]. The generic choice for this Lie algebra (which has to be of dimension $6n$) is $SU(2)^{2n}$. In dimensions below five there are additional solutions, which are of some interest for type-II strings, but not for chiral heterotic strings. To obtain a supercurrent of the form (7.4) one simply bosonizes the $6n$ fermions into $3n$ bosons using (5.32). There is some freedom in doing that, because one may associate the fermions in different ways with the $6n$ $SU(2)$ generators, but there is one especially simple way which yields only norm 3 vectors in the supercurrent. This is to take

$$T_F = N \sum_{i=1}^n [\psi_i^1 \psi_i^2 \psi_i^3 + \psi_i^{-1} \psi_i^{-2} \psi_i^{-3}]$$

where the notation is as in (5.32). We leave it as an exercise to determine the normalization constant N . The vectors $\vec{t}$ one gets from this supercurrent are, for $n = 1$,

$$\vec{t} = (\pm 1, \pm 1, \pm 1), \quad (7.6)$$

For larger n one just adds sets of 8 such vectors in orthogonal three-dimensional subspaces, so that altogether one has $8n$ vectors. This supercurrent was first used by Kawai et. al., and lead to their “triplet constraint”.

Although one might think that all supercurrents that satisfy (7.3) can be written as a trilinear expression of free fermions, this is not the case. A counterexample, found in [56] and [57] has vectors $\vec{t}$ of the form (for $n=1$)

$$\begin{aligned} \vec{t}_1 &= (\sqrt{3}, 0, 0) \\ \vec{t}_2 &= (0, \sqrt{3}, 0) \\ \vec{t}_3 &= (0, 0, \sqrt{3}) \end{aligned} \quad (7.7)$$

This pattern can be repeated n times if $n > 1$. By bosonizing the free fermionic supercurrent, one can only get vectors of the form $(\pm 1, \pm 1, \pm 1, 0, \dots, 0)$, i.e., there must exist a basis in which three entries are ± 1 , and the others zero. For one norm 3 vectors one can always achieve that, but not necessarily simultaneously for several vectors. In particular it is not possible for (7.7).

The mistaken impression that any supercurrent can be written in terms of free fermion trilinears, is based on the proof of [58] that the Fock space of any torus-compactified theory can always be written in terms of a fermionic representation. This may involve either fermions with Thirring interactions (to which the above does not apply, since one would not have a *free* fermion supercurrent), or even free fermions with complicated boundary conditions. Now of course a bosonic theory described in this fermionic way will still have world-sheet supersymmetry, since it is equivalent. But one cannot conclude from this argument that there must always exist a representation of the supercurrent of the trilinear form (7.5).

An illustrative example is a string theory on an E_8 lattice. The Fock space can be described either by means of fermionic oscillators with certain GSO projections, or with bosonic oscillators and lattice momenta. The Frenkel-Kac generators of the D_8 subalgebra of E_8 can also be described both bosonically and fermionically, the latter being fermion bilinears. But whereas the bosonic description extends trivially to the remaining spinor roots of $E_8/SO(16)$, the fermionic one does not. To describe these generators one would need a *transcendental* expression in terms of the fermions rather than a bilinear one (see the discussion in section (5.4)). We believe that the same applies to the supercurrent, with “bilinear” replaced by “trilinear”.

Finally we mention a second method for generating supercurrents, which owes its inspiration to asymmetric orbifolds. In this case one starts by taking the ten-dimensional form also in the internal space, i.e.

$$T_F^{int} = \sum_{I=1}^{2n} \psi^I \partial X_I.$$

The fermions can be bosonized trivially. Although the rest of the supercurrent is already bosonic, we can try to re-express the factors ∂X as sums of exponentials of the form $\exp(i\vec{r} \cdot \vec{X})$, where $\vec{r}^2 = 2$. These expressions are constrained by the fact that they must satisfy the same operator products as the ∂X ’s. An equivalent way of saying this is that they must be the generators of a $U(1)^{2n}$ algebra. One can find such subalgebras by rotating the ∂X ’s into the root system of a Lie algebra. One way of finding such solutions is to use the Weyl rotations of a Lie algebra, and apply the conjugation that maps twists into shifts. We will not discuss this in more detail here; for details we refer to [59]. The solutions already given above can be obtained in this way (this has been shown in [57]), as well as many others, listed in [59].

8. The Spectrum

To discuss the spectrum of covariant lattice theories we will use the even lattice formulation. It may be useful to summarize the requirements that we have derived in the foregoing chapters. To specify a covariant lattice theory in $d = 10 - 2n$ dimensions one must provide

- A lattice $\Gamma_{16+2n;8+2n}$ with left- and right dimensions as indicated
- A supercurrent that satisfies the $N = 1$ supersymmetry algebra, and is consistent with the lattice.

The supercurrent has the general form

$$T_F = \psi^\mu \partial X_\mu + T_F^{int}$$

where the internal part appears only below 10 dimensions. The fermions ψ^μ can be bosonized, and the momenta of the resulting bosons occupy $\frac{d}{2} = 5 - n$ of the $8 + 2n$ dimensions of the right part of the lattice. The internal part of the supercurrent is entirely bosonic, and has the form

$$T_F^{int} = \sum_{\vec{t}} A(\vec{t}) e^{i\vec{t} \cdot \vec{X}} + \sum_{\vec{l}} \vec{B}(\vec{l}) \cdot \partial \vec{X} e^{i\vec{l} \cdot \vec{X}},$$

where $\vec{t}^2 = 3$, $\vec{l}^2 = 1$ and $\vec{l} \cdot \vec{B} = 0$, and the coefficients satisfy the conditions discussed in the previous chapter. The momenta of the bosons $\vec{X}$ occupy $3n$ dimensions of the right part of the lattice. The remaining 3 components of the right lattice are related, in a somewhat indirect way, to the β, γ ghost system, as explained in chapter 6.

The lattice must satisfy the following requirements.

1. It must be even and self-dual with respect to a Lorentzian metric of the form $\text{diag}((-)^{16+2n}; (+)^{8+2n})$
2. The last $8 - n$ components of any vector must belong to one of the four conjugacy classes $(0), (v), (s)$ or (c) of the Lie algebra $D_{8-n}^{spacetime}$.
3. It must contain all vectors of the form $\vec{w}_L = 0, \vec{w}_R = (\vec{l}, (v))$ and $(\vec{t}, (v))$ for all $\vec{l}$ and $\vec{t}$ appearing in the supercurrent.

The last condition applies obviously only to dimensions less than 10.

As explained before, not all the states that one obtains from such a lattice are physical. The physical states are obtained by making a decomposition of D_{8-n} to its subalgebra $D_{4-n} \times D_4$. One keeps only states of which the last four components are fundamental vector or spinor weights of D_4 , and counts them with multiplicity one.

This decomposition of $D_{8-n}^{spacetime} \rightarrow D_{4-n}^{light-cone} \times D_4^{ghost}$ is very simple. The branching rules are as follows:

$$\begin{aligned} (0) &\rightarrow (0, 0) + (v, v) \\ (v) &\rightarrow (v, 0) + (0, v) \\ (s) &\rightarrow (c, c) + (s, s) \\ (c) &\rightarrow (s, c) + (c, s) \end{aligned}$$

We have written them in such a way that the second term is the one that survives the

physical state projection (6.36), because it has as its second ($= D_4$) entry (v) or (s). Of course we could have chosen the other spinor weight of D_4 as the one that gives physical states, but this merely amounts to a chirality flip of the entire theory.

Considering now the first entry, we can read off the spin of the particles in each conjugacy class of D_{8-n} . The first entry corresponds to D_{4-n} , which is the transverse (Wick-rotated) Lorentz group in $10 - 2n$ dimensions. We see that the following general rule holds for D_{8-n} conjugacy classes

$$\begin{aligned} (0) &\rightarrow \text{odd rank tensors} \\ (v) &\rightarrow \text{even rank tensors} \\ (s) &\rightarrow \text{spinors of chirality } + \\ (c) &\rightarrow \text{spinors of chirality } - \end{aligned} \tag{8.1}$$

The assignment of positive or negative chirality to the spinors is a matter of convention.

This list gives us the Lorentz representations of the pure lattice states in light-cone gauge. These are states that have a certain lattice momentum, but no bosonic oscillator excitations. The complete Fock space of physical states is obtained as follows

$$(\alpha\text{-oscillators})(\tilde{\alpha}\text{-oscillators})|\vec{w}_L; \vec{w}_R > |_{m_L=m_R}.$$

Here α denotes all 24 left-moving oscillators, of which $d - 2$ have a vector index of the transverse Lorentz group and will be denoted α^i . The remaining, “internal” ones will be denoted $\alpha^I, I = 1, \dots, 26 - d$.

For the $\tilde{\alpha}$ oscillators things are a bit more complicated. One has $d - 2 = 8 - 2n$ right-moving oscillators $\tilde{\alpha}^i$ in the vector representation of the transverse Lorentz group, which are simply the right-moving modes of the uncompactified coordinates. Then there are the oscillators belonging to the lattice degrees of freedom. Four of the $8 + 2n$ right-moving lattice dimensions are associated with ghosts and longitudinal modes, and the corresponding four bosonic oscillators should not be included; the remaining $2n + 4$ oscillators ($3n$ internal ones and $4 - n$ in the Cartan subalgebra of $D_{4-n}^{\text{light-cone}}$) should be allowed to act on the pure lattice states. They do not alter the Lorentz representation of the state on which they act. Obviously the $d - 2$ left and right-moving bosonic oscillators α^i and $\tilde{\alpha}^i$ do change the Lorentz representation. Thus, as we already pointed out, (8.1) is valid only for pure lattice states.

The last fact one has to know about the spectrum is the mass formula. This is extremely simple: It is just the mass formula of the compactified bosonic string:

$$\begin{aligned} \frac{1}{8}m_L^2 &= \frac{1}{2}w_L^2 + \mathcal{N} - 1 \\ \frac{1}{8}m_R^2 &= \frac{1}{2}w_R^2 + \tilde{\mathcal{N}} - 1 \end{aligned}, \tag{8.2}$$

Here $(\vec{w}_L; \vec{w}_R)$ denotes a vector on $\Gamma_{16+2n; 8+2n}$. Notice that the entire vector contributes to the mass formula, including the D_4 components. Since every physical state necessarily is a fundamental vector or spinor of this D_4 , the lowest possible value of $\frac{1}{8}m_R^2$ is $-\frac{1}{2}$.

It follows that if we are only interested in massless states and tachyons, then we can ignore the right-moving oscillators, since they would increase the mass by too much. In the following we will use $M_R(M_L)$ to denote $\frac{1}{8}m_R^2$ ($\frac{1}{8}m_L^2$), and we will refer to this quantity simply as “mass”. Thus we will quote masses in units in which the bosonic string tachyon has mass -1.

8.1 TEN-DIMENSIONAL STRINGS

Now we have all the ingredients necessary to determine the spectrum. Since the ten-dimensional case is somewhat simpler, we will consider it first. The lowest states one can have in the four conjugacy classes of D_8 are discussed below.

The conjugacy class (0)

The shortest vectors in this conjugacy class are $\vec{w}_R = 0$ and $\vec{w}_R$ equal to a root of D_8 . The first cannot satisfy the physical state criterion, so that the physical state with the lowest value of m_R is obtained by taking $\vec{w}_R$ to be a root. Its mass is equal to zero, and according to (8.1) it yields a Lorentz vector. To satisfy the condition $M_L = M_R$ we have to pair it with a massless state from the left sector. There are two ways to achieve that: either by taking $\vec{w}_L^2 = 2$ or by taking $\vec{w}_L = 0$, and acting with one of the 24 bosonic oscillators. Eight of these oscillators are in the vector representation of the transverse Lorentz group $SO(8)$. Its tensor product with the right-moving sector yields the $SO(8)$ representations $(35) \oplus (28) \oplus (1)$, corresponding respectively to the graviton, the antisymmetric tensor B_{ij} , and a dilaton. The remaining 16 bosonic oscillators yield upon multiplication with the right sector 16 vector bosons. These turn out to be gauge bosons of a $U(1)^{16}$ gauge group. If there are any vectors $\vec{w}_L$ of norm 2 that combine on the right lattice with the (0) conjugacy class of D_8 , then they will give rise to additional gauge bosons, extending $U(1)^{16}$ to a non-abelian gauge group of rank 16.

Note that if the lattice contains vectors of the form $(\vec{r}; (0))$, with $\vec{r}^2 = 2$, then it contains in particular vectors of the form $(\vec{r}; 0)$. We will call such vectors, and also vectors of the form $(0; \vec{r})$ roots of the lorentzian lattice. Notice that all the physical states we have this discussed so far owe their existence to roots: the graviton and the 16 commuting gauge bosons to the roots of D_8 , and the remaining gauge bosons to roots of the left lattice. The graviton (and also the 16 abelian gauge bosons) are inevitable, since the D_8 roots are always present.

It can be shown that the roots of the left (right) part of the lattice must always form a root system of some simply laced Lie algebra. In the subspace spanned by these root systems, all lattice vectors must have components that have integer dot products with the root lattice of this Lie algebra. Hence in this subspace they must be weights.

If the roots of the left lattice span an l -dimensional subspace of the 16 left-moving dimensions, then the gauge bosons belong to a non-abelian rank l group, times an abelian group $U(1)^{16-l}$. We will denote this entire group as G_L . All other states in the theory belong to certain representations of this group.

The conjugacy class (v)

The lowest state that can appear here is the vector representation of D_8 . If it appears, there is a potential problem, because for this lattice vector $M_R = -\frac{1}{2}$. Because

the lattice is even, we know that $\vec{w}_L^2 = 1 \bmod 2$. Should there exist a lattice vector with both $\vec{w}_R^2 = 1$ and $\vec{w}_L^2 = 1$ then the theory would have a tachyon. It might happen that the lowest value of $\vec{w}_L^2$ is in fact 3 rather than 1, and indeed there is one example of that, the $O(16) \times O(16)$ string (see the section 9.1 for a discussion). Furthermore there are lattices on which the conjugacy class (v) is simply not present. This is the case for the supersymmetric strings (see below).

The conjugacy classes (s) and (c)

The lowest possible states are the D_8 spinors, which have $M_R = 0$. As was the case for the conjugacy class (0) , they can become physical either by pairing with a left-moving oscillator excitation or with a norm two vector. If the first happens, then in particular it can combine with an oscillator α_i . The left-right tensor product contains then a vector-spinor, usually called a gravitino.

If a gravitino is present, then the lattice state excited by the left-moving oscillators must have $\vec{w}_L = 0$, combined with a D_8 spinor. This lattice vector is a root of the *right* lattice, and by the arguments given above, it follows that the right lattice must have a larger root system than that of D_8 . It does not require a lot of thought to realize what it must be, namely E_8 . The presence of a gravitino implies that one must obtain supergravity in space-time. Again we see that a local symmetry of the theory is related to a root of the lattice.

Once one realizes that the spectrum has a gravitino, one would clearly expect it to be supersymmetric. For example, one would expect to find gaugino's, the superpartners of the gauge bosons. These are easy to identify: one can excite vectors with right-moving components in the (s) conjugacy class of D_8 in precisely the same way as those in the (0) conjugacy class discussed above. A little bit of thought will make it clear that these are the only massless spinors one can get; in ten-dimensional superstrings all massless fermions are either singlets or adjoints of the gauge group.

Of course one can also have non-supersymmetric strings. In that case, massless fermions can only arise from vectors of the form $(\vec{w}_L; (s))$, with $\vec{w}_L^2 = 2$. It should be emphasized that $\vec{w}_L$ cannot be equal to a root of the right lattice. If it were, one could subtract $(\vec{w}_L; 0)$ from the spinor, and obtain a gravitino by exciting the result with a bosonic oscillator. Thus in non-supersymmetric string theories in ten dimensions one cannot have spinors in the adjoint of the gauge group.

Up to now we have not distinguished the (s) and the (c) conjugacy class of D_8 . They give spinors with opposite space-time chirality. Whether the theory is chiral depends on the representations (i.e. the weights) with which the two conjugacy classes combine on the left lattice. If these are always the same, the theory is not chiral. Otherwise it is.

8.2 FOUR-DIMENSIONAL STRINGS

After this rather lengthy discussion of ten-dimensional strings, let us now see what changes if one goes to lower dimensions. The main difference is that now there can be additional contributions to M_R from the components of $\vec{w}_R$ which extend beyond the space-time lattice D_{8-n} . For definiteness (and for phenomenological reasons) we will consider here four dimensions. The discussion of 6 and 8 dimensions is not very different, the main difficulty being the mod-8 properties of spinors in different dimensions.

In four dimensions, the lattice is $\Gamma_{22;14}$, and the right part contains a D_5 space-time lattice. Again we will consider each of the conjugacy classes of D_5 .

The conjugacy class (0)

Here the discussion can be very brief, because there are almost no changes with respect to the ten-dimensional case. Roots of D_5 again give zero mass gravitons and gauge bosons, this time of course of a rank 22 gauge group. The 9 “internal” components of the right lattice play no role in the massless spectrum, as they would excite the mass above zero.

The conjugacy class (v)

As in ten dimensions, the lowest possible state here is a tachyon of mass $-\frac{1}{2}$. It originates from any lattice vectors of the form $(\vec{w}_L; 0, (v))$, $\vec{w}_L^2 = 1$, where the middle component refers to the nine internal dimensions of the lattice, according to the decomposition $\Gamma_{22;14} = (\Gamma_{22})_L \times (\Gamma_9^{int} D_5^{spacetime})_R$. Clearly we have now the possibility to excite these nine components. In other words, the lattice may contain vectors of the form

$$(\vec{w}_L; \vec{u}_R, (v)), \quad \vec{w}_L^2 = 1 + \vec{u}_R^2.$$

This state would have mass $M_L = M_R = \frac{1}{2}\vec{u}_R^2 - \frac{1}{2}$. The possibilities that one has for $\vec{u}_R$ are severely restricted by the constraint vectors of the supercurrent. Since the lattice is self-dual, they must have integer dot-product with all the vectors $\vec{t}$ and $\vec{l}$ appearing in (7.4). For example, for the supercurrent (7.6) it is easy to see that the norm of $\vec{u}_R$ must be a multiple of $\frac{1}{4}$, so that in addition to the tachyon at mass $-\frac{1}{2}$, one may (depending on the lattice one chooses, of course) find tachyons of mass $-\frac{1}{4}$.

None of this is very important though, because one would normally not consider such tachyonic theories. More interesting is the possibility of having a vector $\vec{u}_R$ of norm 1. (Of course it is possible that the lattice has such vectors, without containing tachyonic vectors) In that case one obtains a state with $M_L = M_R = 0$, which is a massless scalar in a representation of G_L with weight vector $\vec{w}_L$, of norm 2.

This raises of course the possibility that $\vec{w}_L$ is actually a root of the left lattice. One can then go through an interesting chain of arguments. If $\vec{w}_L$ is a root (or, in other words, if one has a massless scalar in the adjoint representation of G_L), then one can subtract a root from $(\vec{w}_L; \vec{u}_R, (v))$, to obtain $(0; \vec{u}_R, (v))$. This is a root of the right part of the lattice, which has non-vanishing dot-product with the roots of D_5 . Again it is not hard to see that the lattice must contain a larger rootsystem than that of D_5 , of which the D_5 roots form a subset. This means that D_5 must be a *regular subalgebra*

of some larger Lie algebra. There are only a few possibilities: This algebra must be E_6, E_7, E_8 , or $D_{5+n}, n \geq 1$. Since the decomposition of the larger algebra must yield a vector representation of D_5 , it can only be D_{5+n} . We will see in a moment that this destroys any possibility of getting chiral fermions out of such a theory, but let us first consider another consequence of this extension of D_5 .

Of course we can excite the state corresponding to the extra roots by a right-moving oscillator α_i . This gives a massless space-time vector, and it should therefore be a gauge boson. That is indeed what it turns out to be. We will not go into the details of its interactions here, but we will just mention that its gauge symmetries originate from the right lattice, and involve the supercurrent in a non-trivial way. In fact, the gauge group turns out to be a non-regular subgroup of the Frenkel-Kac group of the right lattice. This group is of little relevance in heterotic strings, since it inhibits chirality, but it is more important in type-II strings; see [20], [21].

In chiral heterotic strings there is thus never a gauge group corresponding to the right-movers (this seems to have been proved more generally by Friedan and Shenker). As a consequence, the vectors $\vec{u}_R$ on the internal part of the right lattice only affect the *multiplicities* of the particles in the spectrum, not the representation they are in. There is no direct way of “measuring” $\vec{u}_R$, because there are no gauge symmetries coupling to it. Nevertheless these vectors do affect the theory in less direct ways, because the Yukawa couplings and Higgs self interactions depend on them.

The existence of this seemingly redundant nine-dimensional internal space is noteworthy for phenomenological reasons, because it makes it possible to have particles in the theory which have identical properties, but differ only in this part of the lattice. We need not point out that this might help explain the existence of fermion generations. Furthermore there can be interesting continuous and/or discrete *global* symmetries originating from this internal space.

The spinor conjugacy classes

Since a spinor of D_5 has norm $\frac{5}{4}$, it would appear that one could get tachyonic spinors of mass $-\frac{3}{8}$. The existence of tachyons in non-singlet representation of the Lorentz group signals an instability of the vacuum which would cause a breakdown of the Lorentz group. One might expect that world-sheet supersymmetry should prevent that. This does indeed appear to be the case. Observe that the D_5 spinor has half-integral dot-product with the constraint vectors of world sheet supersymmetry, because those vectors have (v) components in D_5 . If a spinor were not to have any components in the internal dimensions, this would be in disagreement with the fact that the lattice is self-dual. Therefore, any D_5 spinor must appear in a vector of the form $(\vec{w}_L; \vec{u}_R, (s))$, and $\vec{u}_R$ must have half-integral dot-product with all the vectors $\vec{l}$ or $\vec{\bar{l}}$ in the supercurrent. In all cases we have examined it turns out that to satisfy this condition, $\vec{u}_R$ must have norm greater than $\frac{3}{4}$. Furthermore there seems always to exist a vector $\vec{u}_R$ which saturates this lower limit. The reader may find it instructive to verify this for the two supercurrents (7.6) and (7.7) which we discussed in the previous section. Although it ought to be possible to prove this without explicitly writing down the supercurrent, for the moment we will accept as a fact that fermionic tachyons do not appear in the spectrum.

About the massless fermions one can say roughly the same as about the ten-dimensional case. The right-moving lattice vectors describing them must have norm 2. Because the lattice is lorentzian even, they appear in combination with even norm vectors $\vec{w}_L$ on the left. Two possibilities are of interest: $\vec{w}_L^2 = 0$ and $\vec{w}_L^2 = 2$.

In the first case one can excite the right-moving Fock space by means of bosonic oscillators α_i to obtain a gravitino. One can also excite it with an internal oscillator α_I or add a root of the left lattice (if it has any), to obtain a gaugino. The appearance of the gravitino tells us that the theory has a certain number of supersymmetries (more about this below). Again we see that the existence of local-space time symmetries is linked to the existence of certain roots of the right lattice.

In the second case, we may assume that $\vec{w}_L$ is not a root of the left lattice, for if were we could subtract $\vec{w}_L$ and obtain the case already discussed. If $\vec{w}_L$ is not a root, it is some vector weight of norm 2 of G_L , and one gets a massless spinor in the corresponding representation. If G_L has $U(1)$ factors part of $\vec{w}_L$ may correspond to $U(1)$ -charges, and the remainder is a weight of the non-abelian part of G_L . In any case the only representations of the non-abelian part of G_L that one can get are those with weights of norm less than or equal to two. Fortunately the representations of the standard model satisfy this bound.

Nothing we have said so far addresses the question of chirality. The situation is a bit different here in comparison with ten dimensions, and this has simply to do with the difference in CPT transformations of spinors in these dimensions. One can easily understand that there is difference by changing the sign of a spinor in the conjugacy class (s) of D_8 or D_5 : in the first case one gets a spinor in the same conjugacy class, whereas in the second case one obtains a spinor in (c). What does this have to with CPT? In four dimensions, a CPT transformation send a left-handed spinor in some representation (R) of some gauge group into a right-handed anti-spinor in the complex conjugate representation ($\bar{R}$). Remarkably, there is an automorphism of every lattice that achieves precisely the same, namely multiplying all lattice vectors with -1. As discussed above, this flips the chirality of the D_5 spinor conjugacy classes, and hence the space-time chirality, while at the same time changing the signs of all vectors on the left lattice. As is well-known, changing the sign of a weight changes it to a weight of the complex conjugate representation.

Thus first of all CPT is a symmetry of any theory we construct. Secondly, to get the usual counting of four-dimensional fermions, we should consider only one D_5 chirality, (s), say, and regard states from the other conjugacy class as anti-particles. Thus if the lattice contains three vectors $(\vec{w}; \vec{w}_R^l, (s)), l = 1, \dots, 3$, where $\vec{w}$ is a weight of a complex representation (R), we have three generations of fermions in the representation (R). The lattice will automatically also contain $(-\vec{w}; -\vec{w}_R^l, (c))$, but these are just the CPT conjugates of the three generations, and are not to be counted separately.

With this understanding, it is obvious how to count chiral fermions. If there are N generations of fermions in (R) and $\bar{N}$ in the complex conjugate ($\bar{R}$), then there are $N - \bar{N}$ chiral generations. This way of counting them assumes implicitly that pairs of opposite chirality will become massive, because there is no symmetry that would forbid such a mass term. We know however of no mechanism in string perturbation

theory that would actually achieve this. Most likely, the massless spectrum is not altered to all orders in perturbation theory, and even the non-chiral fermions will then simply stay massless. It is however quite clear that non-perturbative effects have a role to play in obtaining low-energy phenomenology, and in particular they have to break supersymmetry and/or the weak interaction gauge symmetry. Since we know nothing about non-perturbative effects, we can take the path of maximal optimism, and assume that they will do whatever we need them to do.

There does not exist a simple criterion for determining *a priori* which lattices can give chiral fermions, and which cannot. There is however a simple necessary condition, namely that the D_5 root system should not be part of the root system of a larger D_n group. If that were the case, all other D_5 representations originate from D_n representations. For the spinors that means that they always come in pairs of opposite chirality, because a D_n ($n > 5$) spinor (s) contains both (s) and (c) of D_5 . These spinors belong to the same representation of any gauge group, and hence they are chiral conjugates (More precisely, if one applies a CPT conjugation to the spinors in the (c) conjugacy class of D_5 , one can easily see that in the notation used above one would have $N = \bar{N}$). This loss of chirality is a well-known phenomenon in torus compactifications of higher dimensional chiral theories, and that is in fact precisely what is happening here: If the D_5 lattice extends to D_n one can regard the theory as a torus compactification of a higher dimensional one.

There are two situations in which such an extension of D_5 is forced on us. First of all, it occurs when we want to have gauge bosons belonging to G_R , as discussed above. Secondly, if the supercurrent contains any vectors $\vec{l}$ of length 1, the corresponding constraint vectors are roots, extending D_5 . Thus to get chiral fermions, one needs $\vec{B}(\vec{l}) = 0$.

This concludes the discussion of the spectrum of covariant lattice theories. We emphasize once more a feature of this discussion which we think is rather elegant, and is unique to the even lattice formulation: All local symmetries (gravity, gauge symmetries of G_L and G_R and supergravity) are related to roots on the lattice. The only massless states not originating from roots are the scalars and the fermions (other than gravitino's and gaugino's).

8.3 SPACE-TIME SUPERSYMMETRY

Space-time supersymmetry is a subject which merits a separate discussion. As we have seen, gravitino's originate from roots on the lattice which are spinors of D_5 . For reasons that should by now be familiar, this implies that D_5 is regular subalgebra of a larger Lie algebra. It is now quite clear that there is a restriction on the number of supersymmetries one can have, because there are not that many Lie algebras that extend D_5 by spinor roots; the only ones are E_6 , E_7 and E_8 . By simply counting the number of gravitino's one gets, we deduce that these three possibilities correspond respectively to $N = 1$, $N = 2$ and $N = 4$ supergravity. Similarly, for type-II theories one can get this from both the left and the right-moving sector, yielding $N = 1, 2, 3, 4, 5, 6$ and 8 as the possible supersymmetries. We find it very intriguing that such relatively recent

results as the classification of supersymmetries can be obtained from the classic work of Cartan!

It is not difficult to extend this to other dimensions, and for the dimensions above four one gets then ^{*}

$d = 4$	D_5	$\subset$	E_6	$N = 1$
			E_7	$N = 2$
			E_8	$N = 4$
$d = 6$	D_6	$\subset$	E_7	$N = 1$
			E_8	$N = 2$
$d = 8$	D_7	$\subset$	E_8	$N = 1$
$d = 10$	D_8	$\subset$	E_8	$N = 1$

(Some people seem to use different conventions for counting the number of supersymmetries, in particular in six dimensions. To compare your favorite convention with ours use the torus compactifications of the $N = 1$ theory in 10 dimensions; these correspond to E_8 in the table). For even more intriguing results in two dimensions we refer to [6].

It seems that everything that's known about supersymmetry and supermultiplets can be found back in the representation theory of the exceptional groups. For example, consider $E_6 \supset D_5$ in four dimensions. The smallest representations of E_6 are the singlet (1), the (27) and the (78), which is the adjoint. Their D_5 content is

$$\begin{aligned} (1) &\rightarrow (1) \\ (27) &\rightarrow (10) \oplus (16) \oplus (1) \\ (78) &\rightarrow (45) \oplus (16) \oplus (\overline{16}) \oplus (1) \end{aligned}$$

According to our rules, the D_5 singlets contain no physical states, while the (10) yields a scalar and the (16) a fermion. Thus the (27) contains the scalar multiplet. (On the lattice the representation (27), which has weights of norm $\frac{4}{3}$, must appear in combination with an internal space vector $\vec{u}_R$). In a similar way one can see that (78) contains the vector multiplet. Note that to count the spinors we should consider only the (16), because the ($\overline{16}$) is just a CPT conjugate. Note also that for every (27) the lattice necessarily contains a CPT conjugate ($\overline{27}$).

In fact, one can push this further, as it is possible to relate the “unphysical” left-over singlet components to the auxiliary fields of $N = 1$ supergravity. Vertex operators for such auxiliary fields have recently been investigated [60]. It is conceivable that the above procedure works also for extended supergravities and/or higher dimensions, as well as for massive multiplets with arbitrary high spins, by using the representation theory of the appropriate exceptional group. We encourage the industrious student to work out the details.

^{*} This relation between space-time supersymmetry and exceptional groups is in fact completely general, and applies for instance also to Calabi-Yau compactified string theories. The reason is that space-time supersymmetry constrains the vertex operator algebra so that it can always be mapped to an exceptional algebra.

We have not said much about massive states, but something needs to be said about them in supersymmetric theories. One would expect that they too belong to supermultiplets. This can in fact be proved by using the triality properties in a somewhat unexpected way. These properties are known to be related to supersymmetry in ten dimensions, and to clarify the relation between the Neveu-Schwarz-Ramond and the Green-Schwarz formalism. In those arguments, one uses triality rotations in the transverse Lorentz group. Clearly that is of no use below ten dimensions, since one has a different Lorentz group. Instead we will show that one can use the D_4 that in our formulation describes the ghosts and longitudinal modes for this purpose. This part of the lattice is dimension independent, so that what follows holds in arbitrary dimensions.

The D_4 factor plays a role in distinguishing fermions from bosons, because the former have as their D_4 components weights of the spinor representation (8_s), whereas the latter have weights of the vector representation (8_v). These vectors can be mapped into each other by an $SO(4)$ rotation which maps all the entire D_4 weight lattice onto itself. This is called an automorphism of the lattice. If such an automorphism maps weights of a given representation into weights of *the same* representation it is called an *inner* automorphism. Clearly, if one has a Lie algebra lattice (i.e. a sublattice of the complete weight lattice, containing at least the root lattice), then all inner automorphisms are of course symmetries of it. There are also automorphisms that map weights of *different* representations to each other. Such automorphism are called *outer*. In particular, D_4 has an outer automorphism which cyclically permutes (8_s), (8_c) and (8_v), known as triality. Clearly an arbitrary D_4 lattice (take for example $(0) + (s)$) is not invariant under such an automorphism.

It is a special property of the embedding of D_4 in the exceptional groups that this outer automorphism becomes an inner automorphism of the exceptional group. This means in particular that for every (8_v) of D_4 contained in some representation of an exceptional group, there is also an (8_s) and (8_c). For the spectrum that statement translates into having a fermion for every boson, and hence into supersymmetry at all levels (at least as far as multiplicities are concerned).

9. Model building

There are many ways of finding solutions to the conditions listed in the beginning of the previous section, but unfortunately none is systematic enough to achieve a complete classification. Only in ten dimensions the complete list of solutions is known (by this we mean the complete list of solutions that can be obtained with the covariant lattice construction, and which therefore have a rank 16 gauge group. In ten dimensions one knows just one additional solution, which has an E_8 gauge group and is tachyonic [49], [61]). We will consider this case first.

9.1 TEN-DIMENSIONAL STRINGS

To obtain ten-dimensional strings we are looking for lattices $\Gamma_{16;8}$, where the right-moving components are D_8 weights. To find them, we can make use of a trick explained in section (3.4). Given a lorentzian even self-dual lattice of the required form, we can make an even self-dual euclidean lattice out of it by simply changing the sign of the last eight components of the metric. This seemingly absurd operation works only because the right lattice factor is known to be D_8 ; certainly changing the sign of the metric on an arbitrary lorentzian even self-dual does not preserve the self-duality. The result of this map is an even self-dual Euclidean lattice of dimension 24. But such lattices have been classified by Niemeier, so that classifying all ten-dimensional strings is reduced to merely finding the right mathematical literature (which in some libraries is still no small obstacle).

Not every Niemeier lattice yields a good string theory. To go back to a lorentzian lattice one has to be able to find a D_8 subalgebra. Obviously this must be a regular subalgebra, because we want to have a D_8 sublattice, not just a subalgebra. This is not difficult, and yields the following solutions

$$\begin{aligned}
 E_8 \times E_8 \times E_8 &\rightarrow E_8 \times E_8 \\
 D_{16} \times E_8 &\rightarrow SO(32) \\
 D_8 \times D_8 \times D_8 &\rightarrow SO(16) \times SO(16) \\
 D_{24} &\rightarrow SO(32) \\
 D_{12} \times D_{12} &\rightarrow SO(8) \times SO(24) \\
 D_{16} \times E_8 &\rightarrow SO(16) \times E_8 \\
 D_{10} \times E_7 \times E_7 &\rightarrow SO(4) \times E_7 \times E_7 \\
 D_9 \times A_{15} &\rightarrow SU(16) \times U(1)
 \end{aligned}$$

Here we have indicated on the left the Niemeier lattice, and on the right the gauge group it produces. Notice that there is one Niemeier lattice, $D_{16} \times E_8$, that produces two different string theories, because there are two ways of embedding D_8 .

The first two theories are supersymmetric, because $D_8 \subset E_8$. The last five are tachyonic, because D_8 is embedded in a D_n factor with $n > 8$. When the Niemeier lattice is mapped to $\Gamma_{16;8}$, this embedding leaves a trace in the form of lattice vectors $((v), (v))$, belonging to a $(D_{n-8})_L \times (D_8)_R$ sublattice originating from the D_n factor. Such vectors lead to tachyons. Finally, the third theory is the rather well-known $SO(16) \times SO(16)$ string, which does not have supersymmetry, and also no tachyons. When it was found it was the first of its kind, but in four dimensions such models are very common.

It is a useful exercise to determine the spectrum of some of these models, using the conjugacy classes of the Niemeier lattices. The answers can be found in [62].

9.2 FOUR-DIMENSIONAL STRINGS

In constructing lattices for four-dimensional strings we face a new technical difficulty, namely the constraint vectors for the supercurrent. Consider for example the choice (7.6). The first thing one should observe is that the form of this supercurrent forces all vectors in the nine internal dimensions to have as their components integers or half-integers. This is true because by addition of constraint vectors one gets, among others, vectors of the form

$$(0, \dots, 0, 2, 0, \dots, 0)$$

in the nine-dimensional internal space, and zero elsewhere. The structure of the lattice in this nine-dimensional space can then be summarized in a nice way by saying that all its vectors belong to conjugacy classes of $(D_1)^9$, where D_1 is the lattice introduced at the end of section (3.3).

Now the entire right lattice is of the form $D_5 \times (D_1)^9$, and it is not unreasonable to assume that also the left lattice can be written as a product of D_n factors. Since even under this assumption the lattices will turn out to be too numerous to be classified, whether one misses any solutions in this way is an academic question (our guess is that one does not). Any D_n factor on the left can always be decomposed into a product of D_1 factors, so one could try to enumerate all the even self-dual lattices based on

$$[(D_1)^{22}]_L [(D_1)^9 D_5]_R \tag{9.1}$$

conjugacy classes. One can imagine a *gedanken* computer program that does this. It would start with a lattice containing the conjugacy classes of the constraint vectors

$$\begin{aligned} &[(0)^{22}, (v), (v), (v), (0), (0), (0), (0), (0), (0), (v)] \\ &[(0)^{22}, (0), (0), (0), (v), (v), (v), (0), (0), (0), (v)] \\ &[(0)^{22}, (0), (0), (0), (0), (0), (0), (v), (v), (v), (v)], \end{aligned}$$

and then it would go through the list of conjugacy classes of (9.1) to find one of even lorentzian length, and with even inner product with all the vectors that are already on the the lattice. This procedure can be repeated to find a next conjugacy class, and when no new conjugacy class can be found, the lattice that one has is self-dual by definition.

Although writing this program is quite trivial, one is discouraged from running it because the total number of conjugacy classes of (9.1) is gigantic: 4^{32} . As explained in section (3.3), this means that a self-dual lattice based on (9.1) contains 2^{32} conjugacy classes, so that even checking them for tachyons, say, would cost an enormous amount of time.

Obviously this approach is quite useless in practice, but it does give us some insight in the extent of the problem. This can be made more precise by comparing it to a similar combinatorial problem, that of constructing all Euclidean even self-dual lattices of dimension 32. One could do that in an analogous way, namely by starting with $(D_1)^{32}$ (in 24 dimensions this approach would indeed give all 24 Niemeier lattices). As a combinatorial problem this is not too different from the one we were facing, because the

number of conjugacy classes is the same^{*}. But for this problem we know a (conservative) lower limit of the number of solutions, namely 8×10^7 (In [6] we gave an often misquoted rigorous upper limit on the number of theories, namely the number of even self-dual lattices of dimension 88. This number is finite, but larger than 10^{1500} . This was only intended to show that the number of solutions is finite, and does not in any way represent the actual number of solutions). This shows that we can expect to find a very large number of four-dimensional string theories for this choice of supercurrent.

A different, and more effective method for constructing lattices (for this supercurrent) was found by the authors of [63], who observed that they can all be obtained from odd self-dual lattices of dimension 22. By using techniques of the kind described in section (3.4), one can glue to this lattice a right-moving sector which contains the supercurrent constraint vectors. However from one odd self-dual lattice one gets many covariant lattices $\Gamma_{22;14}$, and a complete classification along these lines appears quite difficult. Of course the situation is improved considerably by imposing additional requirements, such as space-time supersymmetry.

Knowing the extent of the problem makes model building a somewhat arbitrary exercise. In fact “model building” is not even the correct terminology; “model finding” would be more appropriate. The objective is to *find* in the gigantic set of solutions one that corresponds to the standard model. It does not help to be just close to the standard model, since in string theory (ideally at least) one cannot adjust the model by adding a few higgses and tune a few parameters or by doing some of the other things model builders like to do.

This is a rather extreme view of the present situation, and in practice one can of course learn a lot from semi-realistic examples, and we will give some below. They are only intended as an illustration of the construction, and should not be confused with phenomenology. In [6] we produced many examples by modifying some of the Niemeier lattices. Here we will follow a perhaps more pedagogical route, the shift vector method. Any lattice of the form (9.1) can be transformed into any other one by adding a suitable set of shift vectors constructed out of D_1 “weights”. In many cases these D_1 conjugacy classes contain roots, which form the root system of some simply laced Lie algebra. It is then convenient to write the the lattice in terms of the corresponding Lie algebra lattice to reduce the list of conjugacy classes. The simplest roots appear in conjugacy classes of the form $(0, \dots, 0, v, v, 0, \dots, 0)$, which combine two D_1 factors into a D_2 . However, it is certainly not true that from (9.1) one can only get D_n gauge groups. For example, the Niemeier lattices, which can have A_n or E_n factors, can be written as D_1^{24} .

A useful starting point is the lattice $(D_{22})_L \times (D_{14})_R$. To the root lattice one has to add four conjugacy classes to obtain a unimodular lattice, and if one furthermore requires it to be even, there is not much choice: The conjugacy classes must be $(0, 0) + (v, v) + (s, s) + (c, c)$ (this is just an Englert-Neveu lattice with the left factor stretched by 8 dimensions). The phenomenology is extremely simple: The first term gives a graviton (plus $B_{\mu\nu}$ and a dilaton), and the gauge bosons of $SO(44)$ and the second

^{*} The fact that the constraint vectors must be on the lattice reduces the number of solutions, but the fact that not all 32 factors are on equal footing in (9.1) leads to an enhancement.

one gives a tachyon in the vector representation of $SO(44)$. The spinors are all massive and non-chiral, so that this example is certainly not of much interest. Note that the D_{14} lattice on the right contains the constraint vectors of world-sheet supersymmetry. In fact there are many possible sets of constraint vectors that one could choose. If one adds shift vectors, one has to make sure that they have integral dot product with one set of constraint vectors, so that these vectors are on the invariant sublattice Λ_0 defined in section (3.4).

Let us now make the model supersymmetric. This is easily achieved by adding a gravitino as a shift vector. To do so, split D_{14} into $D_6 \times D_8$ and add the conjugacy class $(0, 0, s)$ of $(D_{22})_L \times (D_6 \times D_8)_R$, where we embed the D_5 space-time lattice into D_8 (in the following we adopt the convention of embedding the space-time lattice in the last factor). Notice that two shift vectors in the same conjugacy class have the same effect, so that we can specify the shift vector by giving its conjugacy class with respect to the sublattice.

Using the shift vector method explained in section (3.4) we can easily determine the conjugacy classes that appear. Since we have three D_n factors, there are in fact $2^3 = 8$ of them, and they are

$$(0, 0, 0) + (v, v, 0) + (s, s, s) + (c, c, s) + (0, 0, s) + (v, v, s) + (s, s, 0) + (c, c, 0)$$

What we have now obtained is a ten-dimensional heterotic string, compactified on a lattice $\Gamma_{22;6} = D_{22} \times D_6$. The D_8 factor of the covariant lattice has been embedded in E_8 because of the addition of the gravitino. This example has in fact more supersymmetry than we bargained for, namely $N = 4$. It is of course not chiral.

This illustrates several general principles. First of all, any theory can be made supersymmetric by simply adding as a shift vector a lattice vector which produces a gravitino in the spectrum. More generally, one can enforce the appearance of any state one likes by simply adding the corresponding lattice vector as a shift vector. One may however get unwanted states in addition. Another noteworthy aspect is the use of spinor weights to break D_n factors into smaller ones.

There are two obvious obstacles to chirality in our example. The first is that the D_5 root system is part of a larger D_n root system (that of D_8 in this case), which always inhibits chirality as explained in the previous section. The second is that the $SO(44)$ gauge group simply doesn't have any complex representations, which is necessary for chirality in four dimensions. The only $SO(N)$ groups with complex representations are $SO(4k+2)$, $k \in \mathbb{Z}$. Their chiral representations are the spinors, and the fundamental spinor representations have weight length $\frac{2k+1}{4}$. Clearly only the fundamental spinors of $SO(2)$, $SO(6)$, $SO(10)$ and $SO(14)$ can occur for massless fermions, since all others would have to high a mass. Furthermore a fermion can not be chiral due to spinors of $SO(6)$, because the $SO(6)$ spinor representations are anomalous, and modular invariant string theory is anomaly free. (Note that $SO(2)$ anomalies can be cancelled by the four-dimensional analog of the Green-Schwarz mechanism [18], [64].)

These chiral representations can only appear in combinations with total weight length equal to two. This still leaves many possibilities, and one of the simplest is the

$SO(10) \times SO(6)$ representation $(16, 4) + (16, \bar{4})$. The necessity to cancel the $SO(6)$ anomaly implies that the number of generations is a multiple of eight in this case.

Breaking $SO(44)$ into such small groups cannot be achieved by adding just one spinor shift vector, but two turn out to be sufficient. First decompose

$$(D_{22})_L(D_6 D_8)_R \rightarrow (D_{14} D_8)_L(D_4 D_2 D_2 D_6)_R$$

by adding a shift vector $(0, s; 0, s, 0, c)$. This shift vector projects out half of the gravitino's, so that we get an $N = 2$ model with an $SO(16) \times SO(28)$ gauge group. It is still not chiral, but we can now break the lattice root system further down to

$$(D_7 D_7 D_5 D_3)_L(D_2 D_2 D_2 D_1 D_1 D_1 D_5)_R$$

by adding the shift vector $(s, 0, 0, s; s, 0, c, s, 0, 0, c)$. This again removes one gravitino, so that we have $N = 1$ supersymmetry. The remaining lattice vector giving a gravitino is $(0, 0, 0, 0; 0, 0, 0, s, s, s, s)$. The constraint vectors appear in the conjugacy classes

$$\begin{aligned} &(0, 0, 0, 0; v, 0, v, v, 0, 0, v) \\ &(0, 0, 0, 0; 0, v, v, 0, v, 0, v) \\ &(0, 0, 0, 0; v, v, 0, 0, 0, v, v) \end{aligned}$$

To check if these vectors are on the lattice, it suffices to verify that they have integer dot product with the shift vectors we used.

Working out the spectrum takes some effort, since the lattice has 11 factors, and hence 2^{11} conjugacy classes. Most of them contribute only to massive states, and with some ingenuity one can determine the massless states by hand. However, this kind of labor is easy to computerize, which makes it much easier to experiment with other lattice constructions. The model we just constructed turns out to be chiral, and its complete fermion spectrum consists of the $SO(14) \times SO(14) \times SO(10) \times SO(6)$ representations

$$\begin{aligned} &2(1, 1, 16, 4) + 2(1, 1, 16, \bar{4}) \\ &+(14, 14, 1, 1) + (1, 1, 10, 6) \\ &+(14, 1, 10, 1) + (1, 14, 10, 1) \\ &+(14, 1, 1, 6) + (1, 14, 1, 6) \end{aligned}$$

Only the fermions in the first line are chiral, and give 16 generations of (16) 's of $SO(10)$. These fermions originate from the lattice vectors $(0, 0, s, s, 0, 0, c, 0, 0, s, c)$ and $(0, 0, s, c, 0, 0, s, 0, 0, s, c)$. Notice that the $(D_2)^3(D_1)^3$ factor of the right lattice only determines the multiplicities of the states in the theory, not their gauge quantum numbers. Since the model is supersymmetric, there are scalars in the same representations as the fermions.

This model is a well-known one, and appears in most of the papers on four-dimensional strings because it is one of the simplest to obtain. The construction given here closely resembles the one used in the fermionic constructions, and we have presented it in this way to allow the reader to make a comparison. Indeed, there is a one-to-one correspondence between the class of models based on (9.1) and fermionic models with “complex” boundary conditions, and (anti)-periodic boundary conditions.

One can continue from here, and add more shift vectors to slice the lattice into still smaller D_n chunks. It should be clear from the above that the reason for the rather large number of generations is the fact that their multiplicity is due to a spinor of $SO(6)$ and one of $SO(4)$. Thus the key to lowering the number of generations is to break these factors. This works indeed as expected, but we don't think it would be illuminating to show the gory details of such models. The model given above is just the tip of a gigantic iceberg. If one splits the lattice into more and more factors, there are more and more possibilities for doing this. Clearly the bulk of the models correspond to lattices which are split as much as possible. Unfortunately this “base of the iceberg” is the hardest part to explore, because of the huge number of conjugacy classes one has to deal with.

To conclude this chapter we will briefly discuss models for another choice of the supercurrent, namely (7.7). Just as our previous choice enforced a $(D_1)^9$ structure on the right lattice, the choice (7.7) restricts it as well. World-sheet supersymmetry requires that the lattice contain nine vectors of the form $(0, \dots, 0, \sqrt{3}, 0, \dots, 0, (v))$, where the last entry refers to D_5 , and the others to the remaining nine dimensions of the right lattice. Any other lattice vector in the conjugacy classes (0) or (v) of D_5 must have entries that are a multiple of $\frac{1}{3}\sqrt{3}$ in these nine dimensions; if it is a spinor of D_5 its entries in these nine dimensions must be an odd multiple of $\frac{1}{6}\sqrt{3}$. Vectors of the form $(0, \dots, 0, 2\sqrt{3}, 0, \dots, 0, (0))$ must always be on the lattice, because they are equal to twice the constraint vectors.

This suggests to define a one-dimensional “root”-lattice

$$\mathcal{U} = \{2k\sqrt{3} \mid k \in \mathbb{Z}\}$$

and its dual, or “weight”-lattice

$$\mathcal{U}^* = \{\frac{k}{6}\sqrt{3} \mid k \in \mathbb{Z}\}.$$

This lattice $\mathcal{U}^*$ contains $\mathcal{U}$ as a sublattice, and has 12 cosets (which, by analogy with lie-algebra lattices we shall call “conjugacy classes”) with respect to it. What we have just learned can then be rephrased as follows: the right lattice must be of the form $\mathcal{U}^9 \times D_5$. This notation must be interpreted in the usual sense, i.e. in addition to the root-lattice a certain set of conjugacy classes can appear.

The volume of the unit cell of the root-lattice $\mathcal{U}^9 \times D_5$ is $2(2\sqrt{3})^9$. To have a chance to get a self-dual lattice, we have to combine it a root-lattice on the left which completes this volume to an integer. This is easy: for example A_2, E_6 have a root lattice of volume $\sqrt{3}$, and $\mathcal{U}$ itself has volume $2\sqrt{3}$. Valid choices for the left lattice are for example $\mathcal{U} \times D_{21}$, or $A_2 \times D_{20}$, or $(A_2)^4 \times E_6 \times E_8$, or $\mathcal{U}^{21} \times D_1$. These are only a few possibilities from an obviously long list, and it should be clear that now one can once again start to construct models as we did above. This class is known to contain the Z_3 asymmetric orbifolds [8], [57]. We leave it as an exercise to find their conjugacy classes in this basis (the answer is essentially in [8], although in a different basis). They can then be used as a starting point for finding new theories by the shift vector method, yielding yet another, presumably very large set of theories. And this is still nowhere near the complete set, because there are many other supercurrents one can choose.

10. Final remarks

Despite the considerable length of this paper, there are still many things that we would have liked to discuss in some detail. Here we will merely list some of them with some brief comments and speculations.

- *Interactions* were not discussed in great detail, because this is not necessary with our construction. Since the entire method is based on conformal field theory, the formalism for calculating correlation functions on any Riemann surfaces is already built in. We don't expect these theories to be less consistent than the ten-dimensional ones. Explicit calculations of tree and one-loop diagrams have been done in [65], [64].
- *Overlaps* with other approaches are only partly understood. By construction, covariant lattice theories in four dimensions have gauge groups of rank 22. To the best of our knowledge other constructions yielding rank 22 gauge groups can be re-expressed in terms of covariant lattices. This has been shown for fermionic construction with complex fermion boundary conditions [5], and for asymmetric orbifolds with twists in the Weyl-group of a Lie-algebra [66],[59]. Furthermore the set of covariant lattice theories seems larger than that of fermionic constructions (of rank 22) with a trilinear supercurrent. The restriction to rank 22 can be overcome by considering
- *Twisted Covariant Lattices*, a new class of four-dimensional string theories, obtained by applying orbifold twists to covariant lattices. Since covariant lattices include Narain lattices, twisted covariant lattices include asymmetric orbifolds. The question is, does this class include anything more? Since one is twisting a much larger class of lattices, there is certainly reason to speculate that it might. The most interesting twists would be those that map bosonized world-sheet fermions into world-sheet scalars, something that fits quite naturally within our philosophy, and does not seem to be included in orbifold constructions. Obviously such twists will be severely restricted by modular invariance and world-sheet supersymmetry, but as far as we can tell there is no reason why these obstacles should be insurmountable.
- *The selection of the vacuum* has not been discussed, because we know of no principle that might distinguish one supersymmetric four-dimensional string from any other. Non-perturbative effects are usually invoked as a *deus ex machina* that will somehow achieve this, but this remains wishful thinking at present. A related question is whether four-dimensional strings are
- *Different vacua or different theories?* Probably regarding them as different vacua of the same, generic heterotic string theory is the best way of thinking about them, since after all different four-dimensional strings are nothing but the same conformal field theory with different boundary conditions. It is however not clear to us exactly in what sense one can regard all these theories as ground states of some space-time theory. Thinking about this question, one is automatically led to the question whether there are

- *Continuous parameters* connecting the theories considered here to other ones. One set of obvious candidates, namely Lorentz-rotations of the lattice, does not yield continuous parameters because in chiral theories they would ruin the constraints imposed on the right lattice by (among others) world-sheet supersymmetry. There could however very well be continuous parameters in the direction of lower rank theories.

Finally, there are many phenomenological questions that we wished we could answer. It would be very interesting to have an example that can approach the standards set in [67] for Calabi-Yau compactification, but the gigantic number of possibilities doesn't make the prospect of searching for one look very inviting. Probably the time is not yet ripe for genuine predictions from string theory, and it might be better to try to gain more insight into the vacuum structure first. In view of the extent of that problem, patience will be required.

ACKNOWLEDGEMENTS

We have benefited from discussions with many people, and are grateful for their help in improving our understanding of the subjects discussed here. We can't name them all without taking the risk of forgetting some, but we make an exception for Dieter Lüst, Bengt Nilsson and Nick Warner, whom we thank for the pleasant collaborations.

REFERENCES

- [1]. D. Gross, J. Harvey, E. Martinec and R. Rohm, Nucl. Phys. B256 (1985) 253; Nucl. Phys. B267 (1986) 75.
- [2]. P. Candelas, G. Horowitz, A. Strominger and E. Witten, Nucl. Phys. B258 (1985) 46.
- [3]. L. Dixon, J. Harvey, C. Vafa and E. Witten, Nucl. Phys. B261 (1985) 620; Nucl. Phys. B274 (1986) 285; M. Mueller and E. Witten, Phys. Lett. 182B (1986) 28; L. Ibañez, H. Nilles and F. Quevedo, Phys. Lett. 187B (1987) 25.
- [4]. K. Narain, Phys. Lett. 169B (1986) 41.
- [5]. H. Kawai, D. Lewellen and S. Tye, Nucl. Phys. B288 (1987) 1.
- [6]. W. Lerche, D. Lüst and A.N. Schellekens, Nucl. Phys. B287 (1987) 477.
- [7]. I. Antoniadis, C. Bachas and C. Kounnas, Nucl. Phys. B289 (1987) 87.
- [8]. K. Narain, M. Sarmadi and C. Vafa, Nucl. Phys. B288 (1987) 551.
- [9]. D. Gepner, Nucl. Phys. B287 (1987) 111; *Exactly Solvable String Compactifications on Manifolds of $SU(N)$ Holonomy*, Princeton preprint PUPT-1066.
- [10]. K. Bardacki, E. Rabinovici and B. Sähring, *String models with $c < 1$ components*, CERN-TH.4760/87 (1987).
- [11]. C.S. Lam, *GSO Projections of modular invariant aperiodic strings*, Princeton preprint IASSNS-HEP-87/12 (1987); A. Chamseddine and J.-P. Derendinger, *Twisted superstrings in four dimensions* Zürich preprint ETH PT 87-2.

- [12]. J. Distler and B.R. Greene, *Aspects of (2,0) Compactifications*, Harvard preprint HUTP-87/A065
- [13]. J.H. Schwarz, *Review of Recent Developments in Superstring Theory*, CALTECH preprint CALT-68-1432; L. Dixon, *Some World-Sheet Properties of Superstring Compactifications on Orbifolds and Otherwise*, Princeton preprint PUPT-1074.
- [14]. C.A. Lütken, *A Large Class of Calabi-Yau Vacua*, CERN-TH.4900/87.
- [15]. A. Cohn, D. Friedan, Z. Qiu and S. Shenker, Nucl. Phys. B278 (1986) 577.
- [16]. W. Lerche and D. Lüst, Phys. Lett. 187B (1987) 45.
- [17]. M. B. Green, J. H. Schwarz and E. Witten, *Superstring Theory*, Cambridge University Press, (1987); J. H. Schwarz, Phys. Rep. 89 (1982) 223.
- [18]. M. Green and J. Schwarz, Phys. Lett. 149B (194) 117.
- [19]. A.N. Schellekens and N.P. Warner, Phys. Lett. 177B (1986) 317; Phys. Lett. 181B (1986) 339; Nucl. Phys. B287 (1987) 87.
- [20]. L. Castellani, R. d'Auria, F. Gliozzi and G. Sciuto, Phys. Lett. 168B (1986) 47; R. Blum, L. Dolan and P. Goddard, Nucl. Phys. B289 (1987) 364.
- [21]. H. Kawai, D. Lewellen and S. Tye, Phys. Lett. 191B (1987) 63; W. Lerche, B.E.W. Nilsson and A.N. Schellekens, Nucl. Phys. B294 (1987) 136; L. Dixon, V. Kaplunowsky and C. Vafa, Nucl. Phys. B294 (1987) 43.
- [22]. A. Polyakov, Phys. Lett. 103B (1981), 207; 211.
- [23]. L. Alvarez-Gaumé, G. Moore and C. Vafa, Comm. Math. Phys. 106 (1986) 1.
- [24]. J. Conway and M. Sloane, Journal of Number Theory 15 (1982) 83; Europ. J. Combinatorics 3 (1982) 219.
- [25]. H. Niemeier, Journal of Number Theory 5 (1973) 142.
- [26]. J.-P. Serre, *A Course in Arithmetic*, Springer Verlag, New York, (1973).
- [27]. P. Goddard and D. Olive, in *Vertex Operators in Mathematics and Physics*, Springer Verlag, (1984); Int. J. Mod. Phys. A (1986) 303.
- [28]. R. Slansky, Phys. Rep. 79 (1981) 1.
- [29]. F. Englert and A. Neveu, Phys. Lett. 163B (1985) 349.
- [30]. J. Leech, Canadian Journal of Mathematics 19 (1967) 251.
- [31]. A.N. Schellekens, contribution to *Perspectives in string theory*, Copenhagen (1987), CERN-preprint, to appear.
- [32]. C. Siegel, Ann. Math. 36 (1935) 527.
- [33]. K. Narain, Sarmadi, E. Witten Nucl. Phys. B279 (1986) 93.
- [34]. J. Polchinsky Comm. Math. Phys. 104 (1986) 37.
- [35]. L. Alvarez-Gaumé, *Supersymmetry and Index theory*, Bonn lectures (1984).
- [36]. P. Ginsparg and C. Vafa Nucl. Phys. B289 (1987) 414.
- [37]. Th. Ohrndorf, Nucl. Phys. B281 (1987) 470.
- [38]. A. Belavin, A. Polyakov and A. Zamolodchikov, Nucl. Phys. B241 (1984) 333.

- [39]. D. Friedan, E. Martinec and S. Shenker, Nucl. Phys. B271 (1986) 93.
- [40]. D. Friedan, in *Unified String Theories* (Proc. Santa Barbara workshop 1985), M. Green and D. Gross, eds. , World Scientific (1986).
- [41]. S. Shenker in *Unified String Theories* (Proc. Santa Barbara workshop 1985), M. Green and D. Gross, eds. , World Scientific (1986).
- [42]. E. Martinec, University of Chicago preprint EFI 87-45, Lectures given at the 1987 spring school on Superstrings, Trieste.
- [43]. M. Peskin, SLAC-PUB 4251, lectures presented at the 3th Advanced Study Institute in elementary particle physics, Santa Cruz (1986).
- [44]. H. Nicolai and A.N. Schellekens, in *Proceedings of the 5th Adriatic Meeting on Superstrings, Anomalies and Unification*, Dubrovnik 1986.
- [45]. A. Kosteletzky, O. Lechtenfeld, W. Lerche, S. Samuel and S. Watamura, Nucl. Phys. B238 (1987) 173.
- [46]. P. Goddard, D. Olive and A. Schwimmer Phys. Lett. 157B (1985) 393.
- [47]. P. Ramond, Phys. Rev D3 (1971) 2415; A. Neveu and J. H. Schwarz Nucl. Phys. B31 (1971).
- [48]. F. Gliozzi, J. Scherk and D. Olive, Nucl. Phys. B122 (1977) 253
- [49]. L. Dixon and J. Harvey, Nucl. Phys. B274 (1986) 93.
- [50]. L. Alvarez-Gaumé, P. Ginsparg, G. Moore and C. Vafa, Phys. Lett. 171B (1986) 155.
- [51]. A. Casher, F. Englert, H. Nicolai and A. Taormina, Phys. Lett. 162B (1985) 121.
- [52]. F. Englert, H. Nicolai and A.N. Schellekens, Nucl. Phys. B274 (1986) 315.
- [53]. A.N. Schellekens, *Multiloop Modular Invariance of the Covariant Lattice Construction of Fermionic Strings*, CERN-TH/4834-87, to appear in Phys. Lett. B.
- [54]. E. Verlinde and H. Verlinde, Nucl. Phys. B288 (1987) 357.
- [55]. P. di Vecchia, V. G. Knizhnik, J. L. Petersen and P. Rossi, Nucl. Phys. B253 (1985) 701; P. Goddard and D. Olive, Nucl. Phys. B257 (1985) 245; I. Antoniadis, C. Bachas, C. Kounnas and P. Windey, Phys. Lett. 171B (1986) 51;
- [56]. W. Lerche, B.E.W. Nilsson and A.N. Schellekens, Nucl. Phys. B294 (1987) 136.
- [57]. H. Kawai, D. Lewellen and S. Tye, *Construction of four-dimensional fermionic strings with a modified supercurrent*, Cornell preprint CLNS 87/69.
- [58]. J. Bagger, D. Nemeschansky, N. Seiberg and S. Yankielowicz, Nucl. Phys. B289 (1987) 349.
- [59]. A.N. Schellekens and N.P. Warner, *Weyl groups, Supercurrents and Covariant Lattices*, CERN-TH.4916/87 (1987).
- [60]. J. Atick, L. Dixon, and A. Sen Nucl. Phys. B292 (1987) 109; M. Dine, I. Ichinose and N. Seiberg, Nucl. Phys. B293 (1987) 253; S. Cecotti, S. Ferrara and L. Girardello, *Gravitational Supervertices in $N=1$, 4D-Superstrings*, CERN-TH.4802/87 (1987).

- [61]. H. Kawai, D. Lewellen and S. Tye, Phys. Rev. D34 (1986) 3794; D. Benett, M. Brene L. Mizrachi and H. Nielsen, Phys. Lett. 187B (1986) 179.
- [62]. W. Lerche, D. Lüst and A.N. Schellekens, Phys. Lett. 181B (1986) 71.
- [63]. J. Balog, P. Forgacs, Z. Horvath and P. Vecsernyés, Phys. Lett. 197B (1987) 395.
- [64]. W. Lerche, B.E.W. Nilsson, and A.N. Schellekens Nucl. Phys. B289 (1987) 609; W. Lerche, B.E.W. Nilsson, A.N. Schellekens and N.P. Warner, *Anomaly cancelling terms from the elliptic genus*, CERN-TH.4765/87 (1987).
- [65]. D. Lüst, S. Theisen and G. Zoupanos, *Four-dimensional heterotic strings and conformal field theory* preprint MPI PAE PTh 83-87 (1987).
- [66]. D. Lüst and S. Theisen, *Four dimensional strings - Orbifolds and Covariant Lattices* preprint MPI-PAE-PTh83/87 (1987).
- [67]. B. Green, K.H. Kirklin, P.J. Miron and C.G. Ross Nucl. Phys. B278 (1986) 667.