COMMISSARIAT A L'ENERGIE ATOMIQUE
Centre d'Etudes Nucléaires de Saclay
91191 Gif-sur-Yvette Cedex

DPHPE 88-03
January 1988

Determination of the Neutral Current Chiral Coupling Constants
 u_L^2, u_R^2, d_L^2 and d_R^2
from a Neutrino and Antineutrino Deuterium Experiment

Amsterdam-Bergen-Bologna-Padova-Pisa-Saclay-Torino WA25 Collaboration

D.Allasia, C.Angelini, A.Baldini, M.Baldo-Ceolin, L.Bertanza, A.Bigi, F.Bobisut,
A.Borg, E.Calimani, P.Capiluppi, S.Ciampolillo, J.Derkaoui, M.L.Faccini-Turluer,
V.Flaminio, A.G.Frodesen, G.Giacomelli, B.Jongejans, C.Louedec, A.Margiotta-Neri,
A.Marzari-Chiesa, R.Pazzi, L.Riccati, A.Romero, A.Rossi, A.Sconza, P.Serra-Lugaresi,
A.Tenner, G.W. van Apeldoorn, P.van Dam, N.van Eijndhoven, D.Vignaud

DEPARTEMENT DE PHYSIQUE DES PARTICULES ELEMENTAIRES

Determination of the neutral current chiral coupling constants
 u_L^2 , u_R^2 , d_L^2 and d_R^2 from a neutrino and antineutrino deuterium experiment

Amsterdam-Bergen-Bologna-Padova-Pisa-Saclay-Torino WA25 Collaboration

D.Allasia⁷ , C.Angelini⁵ , A.Baldini⁵ , M.Baldo-Ceolin⁴ , L.Bertanza⁵ ,
 A.Bigi⁵ , F.Bobisut⁴ , A.Borg⁶ , E.Calimani⁴ , P.Capiluppi³ ,
 S.Ciampolillo⁴ , J.Derkaoui^{3*} , M.L.Faccini-Turluer⁶ , V.Flaminio⁵ ,
 A.G.Frodesen² , G.Giacomelli³ , B.Jongejans¹ , C.Louedec⁶ , A.Margiotta-Neri³ ,
 A.Marzari-Chiesa⁷ , R.Pazzi⁵ , L.Riccati⁷ , A.Romero⁷ , A.Rossi^{3**} ,
 A.Sconza⁴ , P.Serra-Lugaresi³ , A.Tenner¹ , G.W.van Apeldoorn¹ ,
 P.van Dam¹ , N.van Eijndhoven^{1***} , D.Vignaud⁶

1. NIKHEF-H , Amsterdam , The Netherlands
2. Institute of Physics, University of Bergen, Norway
3. Dipartimento di Fisica dell'Università di Bologna and INFN Sezione di Bologna, Italy
4. Dipartimento di Fisica dell'Università di Padova and INFN Sezione di Padova, Italy
5. Dipartimento di Fisica dell'Università di Pisa and INFN Sezione di Pisa, Italy
6. Département de Physique des Particules Élémentaires, CEN Saclay, France
7. Dipartimento di Fisica Sperimentale dell'Università di Torino and INFN Sezione di Torino, Italy

(*) Now at Faculté des Sciences , Université d'Oujda , Maroc

(**) And Dipartimento di Fisica dell'Università della Calabria , Italy

(***) Now at CERN

Abstract

The ratios of neutral current to charged current cross sections of ν and $\bar{\nu}$ interactions separately on proton and neutron targets have been measured. The Big European Bubble Chamber (BEBC), filled with deuterium and equipped with an external muon identifier (EMI) and an internal picket fence (IPF), was exposed to the CERN SPS (anti)neutrino wide band beam. The measured ratios are $R_{\nu p} = 0.405 \pm 0.024 \pm 0.021$, $R_{\nu n} = 0.243 \pm 0.013 \pm 0.016$, $R_{\bar{\nu} p} = 0.301 \pm 0.027 \pm 0.024$ and $R_{\bar{\nu} n} = 0.490 \pm 0.050 \pm 0.037$. (The first error is statistical and the second systematic). From combinations of these ratios the following neutral current chiral coupling constants have been determined: $u_L^2 = 0.099 \pm 0.018 \pm 0.008$, $d_L^2 = 0.202 \pm 0.020 \pm 0.019$, $u_R^2 = 0.020 \pm 0.016 \pm 0.009$ and $d_R^2 = 0.002 \pm 0.017 \pm 0.010$. These results agree with the predictions of the $SU(2) \times U(1)$ standard electroweak model. Assuming $\rho = 1$, the corresponding value of $\sin^2 \theta_w$ is found to be $0.247 \pm 0.018 \pm 0.023$, whereas a two-parameter fit to the data yields $\sin^2 \theta_w = 0.243 \pm 0.035 \pm 0.033$ and $\rho = 0.996 \pm 0.027 \pm 0.032$.

1. Introduction.

The GWS SU(2) $\times$ U(1) electroweak model is a serious candidate for a unified theory of weak and electromagnetic interactions [1]. In recent years a large effort has gone into a precise determination of its parameter $\sin^2\theta_w$ [2] : high accuracy is required to test the theory at the level of radiative corrections [3]. The electroweak model must also be tested in all possible sectors which involve neutral-current (NC) processes, i.e. processes mediated by the neutral intermediate vector boson (Z^0). In particular the (anti)neutrino-nucleon NC interactions are described by the left-handed and right-handed coupling constants of the various quarks to the Z^0 (see for example ref. [4]). The absence of flavour-changing neutral-current processes induces equality between strange (s) and charm (c) coupling constants and their corresponding non strange partners ($s_{L,R} = d_{L,R}$ and $c_{L,R} = u_{L,R}$). Thus one is left with four independent chiral coupling constants u_L^2 , d_L^2 , u_R^2 and d_R^2 . In the GWS model these coupling constants depend only on the single parameter $\sin^2\theta_w$.

Most of the experimental determinations of the complete set of coupling constants have been made by combining results from different experiments, a procedure which may introduce unknown systematic uncertainties. The simultaneous measurement of the four NC cross sections of neutrinos and antineutrinos on proton and neutron targets within the same experiment minimizes the systematic effects. Our collaboration, using the BEBC bubble chamber filled with deuterium, presented such a result a few years ago [5]. However the cuts applied to avoid contamination in the NC sample from neutral hadron induced events contributed significantly to the uncertainties. The experimental apparatus was improved by adding an internal picket fence, to get timing information and therefore discrimination between the events, as explained below. This paper describes the new set of experimental data and a new GWS-model-independent determination of the four chiral coupling constants.

The experimental procedure is described in section 2. The ratios R

$$R = \frac{NC}{CC} = \frac{\sigma(\bar{\nu} N \rightarrow \bar{\nu} X)}{\sigma(\bar{\nu} N \rightarrow \mu X)} \quad (1)$$

of neutral current to charged current (CC) $\bar{\nu}N$ processes (N being a proton, a neutron or a deuteron) averaged over the studied energy region are given in section 3. The determination of the chiral coupling constants from these ratios is made in

section 4 where the results are compared with the predictions of the electroweak model. The conclusions are given in section 5.

2. Experimental procedure.

a) Beam and detector.

The experiment was carried out by exposing the Big European Bubble Chamber (BEBC) [6] to the CERN SPS wide band neutrino and antineutrino beams produced by 400 GeV protons incident on a beryllium target. A magnetic horn system focused right sign π and K (π^+ and K^+ for the ν beam and π^- and K^- for the $\bar{\nu}$ beam) and defocused wrong sign π and K. In the antineutrino beam 25.8 % of the events produced in BEBC were induced by ν_μ and in the neutrino beam 3.3 % of these events were induced by $\bar{\nu}_\mu$. There is also a small contribution (4.8 % in the antineutrino beam and 2.3 % in the neutrino beam) of ν_e and $\bar{\nu}_e$ events. The ν_e and $\bar{\nu}_e$ come mainly from kaon decays. For the neutrino (antineutrino) interactions the average energy of the events is 60 GeV (48 GeV). More details concerning the beams can be found in [7].

The bubble chamber, filled with deuterium with a density of 0.140 g/cm^3 , was equipped with an external muon identifier (EMI) [8] consisting of two planes of proportional wire chambers, separated by a hadron absorber. The EMI surrounded the forward part of BEBC (Fig. 1) to identify muons emerging from charged current interactions. The internal picket fence (IPF) detector [9,10] was a system of proportional tube chambers surrounding the bubble chamber, giving 90% coverage in azimuth (see also Fig. 1). It was developed and integrated with the EMI system as an additional detector plane inside the BEBC shielding. Its basic structure was a two-layer vertical array of proportional tubes (2.20 m long and 7 mm internal diameter), separated from the BEBC liquid only by the pressure vessel (5 cm of stainless steel) and some insulation. The IPF was designed to provide timing information for the events measured in the bubble chamber and became operational in 1981. It is of crucial importance for the separation of neutral current events from the upstream produced hadronic background. Indeed during a wide band burst (3 ms of duration) several interactions may occur upstream of the chamber. To unambiguously identify an event on the basis of the time information, the 3 ms spill was divided into 250 ns intervals (time slots). With the IPF a specific

event which has at least one entering or one leaving track can be timed and assigned to a particular time slot. A trigger signal was derived from the IPF each time any of the proportional tubes recorded a hit. After a trigger occurred, a complete readout of the IPF system was performed and all hits occurring in that particular time slot were recorded. The use of this timing of events for separating NC from hadronic background is explained in section 2e.

b) Scanning and measurement.

The present study is based on 73000 ν pictures and 128000 $\bar{\nu}$ pictures obtained in the 1981 and 1983 exposures of BEBC to the wide band beam. A fiducial volume of 17.3 m^3 , corresponding to a mass of 2.42 t of deuterium, was used for the primary lepton interaction in the chamber. The fiducial volume was shaped in such a way that a minimum track length of 50 cm was guaranteed for the particles produced in the interactions. All pictures were scanned twice. Bare two and three prong events without any charged particle with a momentum above 4 GeV/c were not considered, because of the large neutral hadronic background expected in this sample. One prong events without a ν^0 or γ were singled out from the scanning procedure since their scanning efficiency was very poor. An estimate of the number of lost one prong events showed that their contribution is negligible in the present analysis.

Only events with a visible energy larger than 5 GeV (after subtraction of identified spectator protons) were considered for further analysis. The correction for scanning efficiency, which depends on the multiplicity of the event, was applied on an event-by-event basis. The overall efficiency was found to be 97.7 % and 98.2 % for the ν and $\bar{\nu}$ beam data respectively.

All events were measured on film-plane digitizers and passed through the standard CERN HYDRA programs for geometrical reconstruction and track extrapolation (THIRA) to the IPF and to the EMI. When necessary the momentum accuracy of the interacting hadrons was improved by measuring the interaction products and in case of large muon momenta the accuracy was improved by using the EMI+IPF spatial information. Leaving tracks produced by secondary interacting hadrons were also measured for low topology events to facilitate their association to a given time

slot. Strange neutral particles K^0 and Λ with visible decay products as well as electron pairs from γ 's were also measured.

The passing rate or processing efficiency reflects the efficiency of the measurement and of the reconstruction procedure. It was estimated for each multiplicity and separately for the two beams. The overall passing rates were found to be 91 % and 94 % for the ν and $\bar{\nu}$ beam data respectively. Corresponding corrections were also applied on an event by event basis (see section 3a).

c) EMI efficiency.

The selection of CC events and NC events, which is the basis of this analysis, relies primarily on the muon identification by the EMI. For this purpose all the leaving tracks of a given event were extrapolated towards the EMI using the THIRA program. Since practically all hadrons are stopped in the iron absorber a track was recognized as a muon if there was a hit in the outer plane of the EMI at the position predicted by the extrapolation in addition to a hit in the inner plane of the EMI or in the IPF. There are two types of EMI inefficiencies : the geometrical inefficiency, which depends on the muon momentum since low energy muons cannot reach the EMI, and the electronic inefficiency, independent of the muon momentum.

The geometrical efficiency was estimated by a Monte-Carlo method. It depends on the momentum, the transverse momentum and the charge of the muon. It drops rapidly for muons with a momentum below 4 GeV/c. This value was used to set a lower limit to select CC events. The CC events were weighted on an individual basis. The mean weight for right sign muons above 4 GeV/c is 1.02 and the mean weight for wrong sign muons above 4 GeV/c is 1.09.

When a muon traverses the EMI, a particular chamber may not give a signal due to electronic inefficiency (including dead regions, handling inefficiencies,...) and, consequently, the muon cannot be recognized. To estimate the electronic inefficiency of the inner plane, muon tracks recognized by the IPF and outer EMI hit associations were selected and a comparison of the predicted and associated hits in the inner plane was made. Assuming the electronic efficiency of the outer EMI plane to be the same as that of the inner EMI plane, the mean values for the

EMI electronic efficiency were found to be $(97.4 \pm 0.4) \%$ and $(97.0 \pm 0.4) \%$ for the neutrino and antineutrino beam data respectively. These values are the same but more accurate than those previously reported [5]. The electronic IPF efficiency was also evaluated, using the muon tracks identified by the two EMI planes. It was computed as the number of muons which have an identified hit in the IPF divided by the number of muons which have a predicted hit in the IPF and found to be $(94 \pm 1) \%$.

d) Proton-neutron separation.

Since BEBC was filled with deuterium, the target particle could either be a proton or a neutron. To distinguish between the two cases use was made of charge conservation. Odd (even) prong events correspond to interactions on proton (neutron). This simple scheme has to be modified for two reasons :

a) spectator protons. If, in the case of interactions on neutrons, the spectator proton has a momentum, due to Fermi motion, larger than 150 MeV/c (corresponding to a minimum visible track length in the chamber), the even prong becomes an odd prong event.

b) rescattering effects. Rescattering occurs when both nucleons are involved in the interaction. In (anti)neutrino interactions this may happen in two cases :
i) interaction of a secondary particle with the spectator nucleon. ii) coherent interaction of the incident (anti)neutrino with the two nucleons of the deuteron: the contribution of this process is small. In both cases the number of outgoing charged particles is odd irrespective of the primary nucleon charge. An estimate of the rescattering fraction f (i.e. the fraction of events where both nucleons were involved in the interaction) was made by a kinematical analysis of the (anti)neutrino data themselves [11], yielding :

$$f = 0.112 \pm 0.003(\text{stat}) \pm 0.010(\text{syst}) \quad (2)$$

The separation between proton and neutron events was done in a statistical way as follows. Since no genuine interaction on proton, nor events with rescattering, are expected to give protons in the backward hemisphere (if we except negligible effects due to Fermi motion), odd prong events with backward protons (BW) are classified as interactions on a neutron while the number of events with a forward going spectator is computed weighting the former with a factor :

$$\omega \approx [(1 - \beta \cos\theta) / (1 + \beta \cos\theta)]^2 \quad (3)$$

(β is the velocity of the backward spectator proton and θ is its production angle with respect to the beam direction in the laboratory system). This factor accounts for the effects due to Fermi motion of the target, namely: i) the flux variation due to the momentum and direction of the target neutrons in the laboratory system, giving the Möller flux factor $(1 + \beta \cos \theta)$ and ii) the (anti)neutrino neutron cross section variation due to the varying total energy of the reaction giving also a factor $\approx (1 + \beta \cos \theta)$. The number of events on neutron is then given by:

$$[(\text{even prongs}) + BW(1+\omega)](1-f)^{-1}$$

and the number of events on proton by:

$$[(\text{odd prongs}) - BW(1+\omega) - f(\text{events on neutron})]$$

e) Classification of events.

The measured events were analyzed by a computer program and by physicists in such a way that the identification of CC and NC events was optimal. There are however background events, contamination of NC events in the CC sample and vice-versa and inefficiencies in the procedure. Here a summary of the event classification algorithm is presented. More details are given elsewhere [12]. Corrections will be discussed in section 3. Our procedure is similar although not identical to the one used in the νp experiment [10].

The leaving tracks of a given event were extrapolated to the IPF and to the EMI. The extrapolation results were mapped onto the data from the IPF+EMI system, resulting in general in the assignment of a time slot to each event after hit association. This timing together with the observed IPF+EMI activities form the basis of the event classification. The procedure consists of:

- i) the identification of the time slot.
- ii) the identification of a muon.
- iii) the identification of signals (noise) in time with the event in the IPF tubes in a properly chosen cone upstream of the event (backward cone). (The vertex of the cone is the event vertex and its axis is defined as the direction opposite to the total visible momentum, see Fig. 2a). Such signals are due to background coming from interactions in the material in front of the chamber. The distribution of the hits not associated to any track in the upstream area within the event time slot is shown in Fig. 2b for all events with $E_{\text{vis}}^{\text{had}} > 5 \text{ GeV}$, the hatched area corresponding to CC events. From this distribution the cone size was taken to extend from -25° to $+25^\circ$.

iv) the search for a muon that traverses the experimental arrangement but that does not emerge from an event in BEBC (throughgoing muon); for this search we used also the information of the veto EMI counters.

The events were divided into 6 classes : i) CC events, i.e. events with one identified muon in one time slot and without noise in the backward cone. ii) DM (dimuon) events, i.e. events with two or more identified muons in one time slot. iii) NC events, i.e. events without any identified muon but assigned to a time slot by hits in the IPF and without any hit in the backward cone. iv) HA (hadron induced) events, i.e. events assigned to a time slot but for which there are hits in the backward cone, or an interaction in the material in front of BEBC signalled by a throughgoing muon and by the coincidence of signals from at least two of the electronic detectors (veto, IPF, inner EMI plane in the same time slot as the event). v) AS events, i.e. events associated to an upstream visible event in the same time slot. vi) DK (don't know) events, i.e. events for which no time slot could be identified or events for which no unique decision could be made.

Since it was verified that the number of hadrons incorrectly identified as muons is negligible for muons above 4 GeV/c , the CC sample is clean but not complete. The dimuon (DM) sample is added to the CC sample , the leading muon being chosen as the muon with the largest value of the momentum multiplied by the transverse momentum. The reliability of this selection was checked by a Monte-Carlo method. The NC sample contains true NC and a number of unidentified CC events which will be determined in section 3. The HA and AS samples are not considered further in this analysis. The DK sample (10% of the events) has to be investigated more carefully since it contains NC and HA events. The main source of DK events are low energy events for which there is no leaving track or for which the leaving tracks are not recognized by the IPF , due to its inefficiency. Less than 50% of the DK events (5% of the total sample) correspond to ambiguous time slots : an event is assigned to different classes in different time slots. The repartition of DK events between NC and HA was done assuming that the probability P for a NC or a HA to become a DK is the same if they have the same characteristics (energy, multiplicity, total transverse momentum, total hadronic energy, ...). If P depends on one of these kinematical variables (let's call it v) it should have the same value for a NC or a HA event with the same v . Several experimental distributions of the DK events were compared with the ones expected on the basis of the assumed hypothesis. It turned out that the energy was the most suitable variable to give

a reliable separation of DK between NC and HA. This analysis was done separately for each interaction. About 60% of the DK events (the exact percentage being slightly different for each interaction) were attributed to the NC sample. A systematic error of 10% on the NC coming from the DK events was estimated arising from this procedure.

3. Determination of the ratios.

To determine the ratios R of NC to CC processes we started from the two samples of data obtained with the neutrino beam and with the antineutrino beam. For each sample the events were separated between p and n targets as explained in section 2d. This defined four categories of interactions : νp , νn , $\bar{\nu} p$ and $\bar{\nu} n$.

a) Raw numbers.

All the selected events have a visible energy larger than 5 GeV , as stated before. For NC events this visible energy is purely hadronic. For CC events it contains also the muon energy. To compare in a similar way the two samples we apply on CC events a cut in visible hadronic energy at 5 GeV. The initial number of CC events is then CC events with a μ^- (for ν data) or μ^+ (for $\bar{\nu}$ data) , with a muon momentum p_μ larger than 4 GeV/c and with a visible hadronic energy larger than 5 GeV. All CC events which have p_μ smaller than 4 GeV/c and a visible energy larger than 5 GeV are placed in the NC sample. The initial number of NC events was formed by the NC events increased with the rejected CC events defined above. These numbers of events were then corrected for scanning efficiencies and passing rates and are given in the first line of Table 1.

b) Corrections.

All the corrections discussed below are given in Table 1.

The corrections due to the EMI+IPF inefficiencies were explained in section 2c. The CC events are weighted by an event dependent factor accounting for geometrical inefficiency and by a factor 1.03 accounting for electronic inefficiency. The correction for NC events decreases the raw number and was done

in a similar way starting from the number of CC events of both signs with $E_{\text{vis}} > 5 \text{ GeV}$ and $p_{\mu} > 4 \text{ GeV/c}$.

The following correction accounts for the NC events coming from the DK sample. About 60% of the DK events with $E_{\text{vis}} > 5 \text{ GeV}$, depending on the specific interaction, were assigned to the NC sample by the procedure explained in section 2e.

A further correction has to be applied to NC events due to low topology and low energy events (2 and 3 prong) which were rejected at the scanning level (no track above 4 GeV/c). The number of events lost was determined from experimental CC events by observing the hadron shower after removing the muon.

A Monte-Carlo computation was used to correct for : CC events of both helicities with $p_{\mu} < 4 \text{ GeV/c}$, and ν_e and $\bar{\nu}_e$ events. The simulation which was used consisted of two parts :

i) The simulation program NUBEAM [13] was used to generate the correct helicity ν_{μ} and $\bar{\nu}_{\mu}$ fluxes for the beam conditions of the present experiment. The right helicity fluxes were found to agree with the experimental results [14]. For the wrong helicity (ν_{μ} in $\bar{\nu}_{\mu}$ and $\bar{\nu}_{\mu}$ in ν_{μ}) NUBEAM did not reproduce correctly the data and the corresponding experimental distributions of CC events were used as input. NUBEAM was also used to provide ν_e and $\bar{\nu}_e$ flux distributions as a function of energy. The various spectra used are shown in Fig. 3.

ii) Events were then generated using the leptoproduction version of the Lund Monte-Carlo [15]. The number of events generated in each category (νp , νn , $\bar{\nu} p$ and $\bar{\nu} n$) was sufficiently large so that the statistical error was negligible. It was checked that Monte-Carlo distributions properly reproduced the main features of the observed events. The various samples of events were normalized in order to reproduce as well as possible the experimental data, in particular the contributions of wrong helicity and $\nu_e + \bar{\nu}_e$ events relative to the right helicity ones.

To correct for the $p_{\mu} < 4 \text{ GeV/c}$ cut we used the ratio of CC events with $p_{\mu} < 4 \text{ GeV/c}$ over CC events with $p_{\mu} > 4 \text{ GeV/c}$ predicted by the Monte-Carlo (A systematic error of 5% is estimated for this correction). The experimental number of CC events was increased in the same ratio. For NC events we subtracted CC events with p_{μ}

$< 4 \text{ GeV/c}$ and with $E_{\text{vis}} > 5 \text{ GeV}$ by normalizing the Monte-Carlo to the number of CC events with $p_{\mu} > 4 \text{ GeV/c}$.

The wrong helicity background consists of a) wrong helicity CC events with $p_{\mu} < 4 \text{ GeV/c}$ and b) wrong helicity NC events. The correction for a) was done as in the previous paragraph. The correction for b) was computed as follows. We calculated first the small $\bar{\nu}_{\mu}$ contribution in the neutrino beam, normalizing to the experimentally found $\bar{\nu}_{\mu}$ CC events and using the NC/CC ratio obtained in our previous experiment. Since this correction is relatively small the final values for the number of NC events are not sensitive to this value. For the ν_{μ} contribution in the $\bar{\nu}_{\mu}$ beam which is larger, we used the experimental NC/CC value that we obtained for ν_{μ} .

The correction for ν_e and $\bar{\nu}_e$ background was computed using the Monte-Carlo. The corresponding systematic error was determined taking into account a 10% uncertainty on the flux.

c) Final numbers.

The corrected numbers of CC and NC events in each category are summarized in Table 1. From these numbers we evaluate the ratios :

$$\begin{aligned} R_{\nu p} &= 0.405 \pm 0.024 \pm 0.021 \\ R_{\nu n} &= 0.243 \pm 0.013 \pm 0.016 \\ R_{\bar{\nu} p} &= 0.301 \pm 0.027 \pm 0.024 \\ R_{\bar{\nu} n} &= 0.490 \pm 0.050 \pm 0.037 \end{aligned} \tag{4}$$

The first error is statistical and the second one systematic. The sources of systematic errors are : correction for $p_{\mu} < 4 \text{ GeV/c}$, uncertainty in the DK repartition, uncertainties in the wrong helicity background and $\nu_e + \bar{\nu}_e$ contamination, uncertainties in the electronic and geometric efficiencies, uncertainty in two and three prong correction and uncertainty in the rescattering fraction.

We computed also the ratios NC/CC for the sum p+n (isoscalar target) and for the difference p-n :

$$R_{\nu} = \frac{\text{NC}(\nu p) + \text{NC}(\nu n)}{\text{CC}(\nu p) + \text{CC}(\nu n)} = 0.297 \pm 0.010 \pm 0.017$$

$$\begin{aligned}
R_{\nu}^{-} &= \frac{NC(\bar{\nu}p) + NC(\bar{\nu}n)}{CC(\bar{\nu}p) + CC(\bar{\nu}n)} = 0.372 \pm 0.021 \pm 0.027 \\
r_{\nu} &= \frac{NC(\nu p) - NC(\nu n)}{CC(\nu p) - CC(\nu n)} = 0.082 \pm 0.034 \pm 0.018 \\
r_{\nu}^{-} &= \frac{NC(\bar{\nu}p) - NC(\bar{\nu}n)}{CC(\bar{\nu}p) - CC(\bar{\nu}n)} = 0.021 \pm 0.100 \pm 0.044
\end{aligned} \tag{5}$$

All these NC/CC ratios have a cut in visible hadronic energy $E_h^{\text{vis}} > 5$ GeV. There are no other cuts.

Our results are compatible with other published results. For recent compilations see for example ref. [16]. Ratios on isoscalar targets, R_{ν} and R_{ν}^{-} , were previously obtained with a better precision [2,17] but in experiments which could not simultaneously determine r_{ν} and r_{ν}^{-} . The values of the four ratios $R_{\nu p}$, $R_{\nu n}$, $R_{\nu p}^{-}$ and $R_{\nu n}^{-}$ update the values that we published in a previous experiment [5] and are statistically independent of the former determination. The present values are obtained without any cut on the transverse momentum of the total hadronic system with respect to the beam direction. The values of $R_{\nu p}$ and $R_{\nu p}^{-}$ agree within 1.5 standard deviations with those obtained in the BEBC hydrogen experiment [18] which has slightly different statistics. The values of $R_{\nu n}$ and $R_{\nu n}^{-}$ are the only ones obtained so far (except for one experiment [19] which also obtained $R_{\nu n}$ but with a larger error).

4. Determination of the chiral coupling constants.

In the framework of the quark parton model the differential cross sections $d^2\sigma/dx.dy$ for (anti)neutrino-nucleon CC and NC processes depend on the quark distributions, on y and E (see for example [20]). For NC processes they depend also on the chiral coupling constants. The ratios R of the integrated cross sections can then be expressed as follows :

$$R = a_1 u_L^2 + a_2 d_L^2 + a_3 u_R^2 + a_4 d_R^2 \tag{6}$$

The coefficients a_i were computed by properly integrating over x and y and over the energy spectrum, using the Lund Monte-Carlo. We used the Q^2 dependent

Gluck-Hoffmann-Reya parametrization [21] of the quark distribution functions. To determine the four coupling constants we used R_ν , R_ν^- , r_ν and r_ν^- . Their expressions as a function of the numerical values of a_1 (with $E_h^{\text{vis}} > 5$ GeV) are:

$$\begin{aligned} R_\nu &= 0.921 u_L^2 + 0.979 d_L^2 + 0.354 u_R^2 + 0.387 d_R^2 \\ R_\nu^- &= 0.968 u_L^2 + 1.072 d_L^2 + 2.634 u_R^2 + 2.787 d_R^2 \\ r_\nu &= -1.125 u_L^2 + 0.982 d_L^2 - 0.307 u_R^2 + 0.229 d_R^2 \\ r_\nu^- &= 1.039 u_L^2 - 0.763 d_L^2 + 3.974 u_R^2 - 3.359 d_R^2 \end{aligned} \quad (7)$$

The coupling constants are then easily obtained :

$$\begin{aligned} u_L^2 &= 0.099 \pm 0.018 \pm 0.008 \\ d_L^2 &= 0.202 \pm 0.020 \pm 0.019 \\ u_R^2 &= 0.020 \pm 0.016 \pm 0.009 \\ d_R^2 &= 0.002 \pm 0.017 \pm 0.010 \end{aligned} \quad (8)$$

The following combinations of the coupling constants may also be directly computed:

$$\begin{aligned} u_L^2 + d_L^2 &= 0.301 \pm 0.013 \pm 0.021 \\ u_R^2 + d_R^2 &= 0.022 \pm 0.012 \pm 0.014 \\ u_L^2 - d_L^2 &= -0.104 \pm 0.036 \pm 0.019 \\ u_R^2 - d_R^2 &= 0.018 \pm 0.031 \pm 0.014 \end{aligned} \quad (9)$$

This complete set of values is obtained using only the data coming from the present experiment. It updates the one that we obtained in a previous experiment [5]. It is seen that, as far as the systematic errors are concerned, the present results do not significantly improve those given in ref. [5]. However we note that in [5] the coupling constants were obtained after applying a cut on the transverse momentum of the hadronic system. This cut would reduce the systematic error as calculated in the present analysis but it would introduce an additional systematic error that is hard to assess and that was not taken into account. This is why we prefer the method that we used in the present paper and we consider (8) and (9) as our best values. These values agree with the published results or with values that can be easily derived from them [17,18,22] and with those obtained in recent analyses [16]. The left-handed couplings are significantly stronger than the right-handed ones. Our values for $u_L^2 + d_L^2$ and $u_R^2 + d_R^2$ are in close agreement with those obtained from large statistics experiments on isoscalar targets [17].

For completeness we calculated also the four coupling constants of the Hung-Sakurai parametrization [23] : α , β , γ and δ respectively characterize the strength of isovector vector , isovector axial , isoscalar vector and isoscalar axial couplings. They are related to the chiral coupling constants themselves and not to their square value (see for example [20b]). So we assume that the sign is the one given by the standard model ($u_L > 0$, $d_L < 0$, $u_R < 0$ and $d_R > 0$). Their values are :

$$\begin{aligned}\alpha &= 0.57 \pm 0.07 \pm 0.09 \\ \beta &= 0.95 \pm 0.06 \pm 0.06 \\ \gamma &= -0.23 \pm 0.16 \pm 0.08 \\ \delta &= -0.04 \pm 0.11 \pm 0.06\end{aligned}\tag{10}$$

In the electroweak standard model β is predicted to be 1 and δ to be 0. Our values agree with these predictions within less than one standard deviation.

The four chiral coupling constants quoted above have been determined independently of the standard electroweak model. In this model they are closely related to the basic parameters $\sin^2 \theta_w$ and ρ :

$$\begin{aligned}u_L^2 &= \rho^2 (1/2 - 2/3 \sin^2 \theta_w)^2 \\ d_L^2 &= \rho^2 (-1/2 + 1/3 \sin^2 \theta_w)^2 \\ u_R^2 &= \rho^2 (-2/3 \sin^2 \theta_w)^2 \\ d_R^2 &= \rho^2 (1/3 \sin^2 \theta_w)^2\end{aligned}\tag{11}$$

From these relations a first test of the standard model can be performed :

$$\begin{aligned}u_L^2 + d_L^2 &= \rho^2 (1/2 + 1/3 \sin^2 \theta_w (5/3 \sin^2 \theta_w - 3)) > 0 \\ u_R^2 + d_R^2 &= 5/9 \rho^2 \sin^4 \theta_w > 0 \\ u_L^2 - d_L^2 &= 1/3 \rho^2 \sin^2 \theta_w (\sin^2 \theta_w - 1) < 0 \\ u_R^2 - d_R^2 &= 1/3 \rho^2 \sin^4 \theta_w > 0\end{aligned}\tag{12}$$

The values quoted in (9) show that $u_L^2 - d_L^2$ is negative at 2.5σ . This is the first evidence obtained in a single experiment. The test that $u_R^2 - d_R^2$ is positive is not significant. In Fig. 4 we show the left-handed (u_L^2 versus d_L^2) and right-handed (u_R^2 versus d_R^2) coupling constants. The elliptic contours enclose the one standard deviation region obtained from our analysis using the statistical

errors only. The lines correspond to the predictions of the electroweak model as a function of $\sin^2\theta_w$.

A test of the model is also provided by the experimental determination of the parameters ρ and $\sin^2\theta_w$. Fixing $\rho=1$, as predicted by the minimal standard model, a fit of the data yields :

$$\sin^2\theta_w = 0.247 \pm 0.029 \quad (13)$$

whereas a two-parameter fit yields :

$$\sin^2\theta_w = 0.243 \pm 0.046 \quad (14)$$

$$\rho = 0.996 \pm 0.041$$

The errors include both the statistical and systematic uncertainties. The value of $\sin^2\theta_w$ determined in the single parameter fit (eq. (13)) is consistent with the world average value, see for instance ref. [16].

5. Conclusion.

From an experiment using BEBC filled with deuterium we have measured neutral and charged current reactions induced by neutrinos and antineutrinos on proton and neutron separately. The use of an internal picket fence introduced in the experimental apparatus allowed a better determination of the neutral hadron induced events than in previous experiments and reduced significantly the errors on the NC/CC ratios $R_{\nu p}$, $R_{\bar{\nu} n}$, $R_{\nu n}$ and $R_{\bar{\nu} p}$. From these ratios the four chiral coupling constants u_L^2 , d_L^2 , u_R^2 and d_R^2 were determined **independently** of the electroweak standard model. Their values were found in agreement with the predictions of the standard SU(2)xU(1) electroweak model and a single parameter fit of the data gave the value : $\sin^2\theta_w = 0.247 \pm 0.029$.

These results are in agreement with recent analyses of the neutral current data [16].

Acknowledgements : We are indebted to the CERN staff for running the SPS , the (anti)neutrino beam, BEBC and the EMI+IPF system. We thank the scanning and measuring staff of the various institutes of our collaboration for their careful work.

References

- [1] S.L.Glashow , Nucl. Phys. 22 (1961) 579
A.Salam and J.Ward , Phys. Lett. 13 (1964) 168
S.Weinberg , Phys. Rev. Lett. 19 (1967) 1264
A.Salam , in "Elementary Particle Theory" , edited by N.Svartholm
(Almqvist and Forlag , Stockholm , 1968)
- [2] H.Abramowicz et al. , Phys. Rev. Lett. 57 (1986) 298
J.V.Allaby et al. , Phys. Lett. B177 (1986) 446
- [3] M.Davier , in Proc. of the XXII Int. Conf. on High Energy Physics ,
Leipzig (1984) , A.Meyer and E.Wieczorek editors , vol.1 , p.238
M.Davier , "Why do we need precision measurements of ρ and $\sin^2\theta_w$?
LAL report 84/33 (1984)
I.Liede , M.Roos and S.Sakakibara , Preprint , April 1985
- [4] J.D.Bjorken , Proc. Summer Institute on Particle Physics ,
SLAC-198 (Stanford , 1976) , p.1
L.M.Sehgal , Phys. Lett. 71B (1977) 99
- [5] D.Allasia et al. , Phys. Lett. B133 (1983) 129
- [6] BEBC User's Handbook , CERN , 1977-1979
- [7] E.H.M. Heijne , "Muon flux measurements with silicon detectors in the
CERN neutrino beams" , CERN Report 83-06 (1983)
- [8] C.Brand et al. , Nucl. Instr. and Meth. file (1976) 485
R.Beuselinck et al. , Nucl. Instr. and Meth. 154 (1978) 445
- [9] H.Foeth , Nucl. Instr. and Meth. 176 (1980) 203
C.Brand et al. , CERN Report EF 81-13 (1981)
- [10] H.Foeth , E.Hoffmann , M.L.Retter and S.J.Towers ,
Nucl. Instr. and Methods A253 (1987) 245
- [11] A. Tenner , to be published
- [12] N. van Eijndhoven , "Event classification in (anti)neutrino experiments
using an IPF-EMI detector system" , NIKHEF-H/87-6 (1987)
- [13] C.Visser , NUBEAM , CERN HYDRA application library (1979)
- [14] D.Allasia et al. , Nucl. Phys. B239 (1984) 301
C.Visser , Thesis , Amsterdam (1986)
- [15] B.Andersson et al. , Phys. Rep. 97C (1983) 31
T.Sjostrand , Comp. Phys. Comm. 27 (1982) 243

- G. Ingelman , Lund Monte-Carlo for deep inelastic lepton nucleon scattering , version 4.3 (April 1983)
- [16] U. Amaldi , A. Bohm , L.S. Durkin , P. Langacker , A.K. Mann , W.J. Marciano ,
A. Sirlin and H.H. Williams , Phys. Rev. D36 (1987) 1385
G. Costa , J. Ellis , G.L. Fogli , D.V. Nanopoulos and F. Zwirner ,
Nucl. Phys. B297 (1988) 244
 - [17] H. Abramowicz et al. , Z. Phys. C28 (1985) 51
P.G. Reutens et al. , Phys. Lett. B152 (1985) 404
M. Jonker et al. , Phys. Lett. B99 (1981) 265 and Phys. Lett. B103 (1981) 469
 - [18] G.T. Jones et al. , Phys. Lett. B178 (1986) 329
 - [19] T. Kafka et al. , Phys. Rev. Lett. 48 (1982) 910
 - [20] a) J.E. Kim , P. Langacker , M. Levine and H.H. Williams ,
Rev. Mod. Phys. 53 (1981) 211
b) L.M. Sehgal , Prog. in Part. and Nucl. Phys. 14 (1985) 1
 - [21] M. Gluck , E. Hofmann and E. Reya , Z. Phys. C13 (1982) 119
 - [22] J. Blietschau et al. , Phys. Lett. 88B (1979) 381
 - [23] P.Q. Hung and J.J. Sakurai , Ann. Rev. Nucl. Part. Sci. 31 (1981) 375
J.J. Sakurai , Proc. of the Int. Conf. on Neutrino Physics , Aachen (1976) ,
ed. by H. Faissner , H. Reitler and P.M. Zerwas , p.515

Table 1a

	νp		νn	
	CC	NC	CC	NC
raw numbers	2581	1777	5124	2884
EMI geometrical efficiency	+61	-93	+149	-197
EMI electronic efficiency	+77	-119	+154	-234
DK		+358		+516
2 and 3 prong		+103		+21
$p_\mu < 4 \text{ GeV/c}$	+506	-533	+1047	-1136
wrong helicity		-60		-59
$\nu_e + \bar{\nu}_e$		-128		-222
corrected numbers	3225	1305	6474	1573

Number of events and number of events entering the correction in the various categories (neutrino beam)

Table 1b

	$\bar{\nu}_p$		$\bar{\nu}_n$	
	CC	NC	CC	NC
raw numbers	1755	1045	987	1199
EMI geometrical efficiency	+38	-112	+26	-131
EMI electronic efficiency	+53	-119	+30	-97
DK		+226		+282
2 and 3 prong		+119		+34
$p_\mu < 4 \text{ GeV}/c$	+179	-190	+164	-176
wrong helicity		-225		-314
$\nu_e + \bar{\nu}_e$		-135		-205
corrected numbers	2025	609	1207	592

Number of events and number of events entering the correction in the various categories (antineutrino beam)

Figure captions

Figure 1 : Schematic layout of BEBC with the IPF and the EMI.

Figure 2 : a) Drawing of a typical event (vertex A) showing the backward cone (dashed lines).

b) Unassociated hit distributions in the upstream IPF area for the selected event time slots. The hatched distributions correspond to CC events. The angle (in degrees) of the hit is defined as the projection in the horizontal plane of the angle between the line connecting the hit position and the event vertex and the direction of the total visible momentum.

Figure 3 : a) Energy distribution for events in the neutrino beam with

$$E_h^{\text{vis}} > 5 \text{ GeV.}$$

The full curve corresponds to ν_μ induced events, the dashed curve to $\bar{\nu}_\mu$ induced events (wide band beam background) and the dotted curve to $\nu_e + \bar{\nu}_e$ induced events.

b) Energy distribution for events in the antineutrino beam with

$$E_h^{\text{vis}} > 5 \text{ GeV.}$$

The full curve corresponds to $\bar{\nu}_\mu$ induced events, the dashed curve to ν_μ induced events and the dotted curve to $\nu_e + \bar{\nu}_e$ induced events.

Figure 4 : Left-handed (u_L^2 versus d_L^2) and right-handed coupling constants.

The elliptic, one standard deviation, contours correspond to our data. The lines correspond to the electroweak model predictions as a function of $\sin^2 \theta_w$.

Fig. 1

a)

b)

Fig. 2

Fig. 3

Fig. 4