GRAND UNIFICATION, SUPERSYMMETRY, SUPERSTRINGS:
AN INTRODUCTION TO PHYSICS BEYOND THE STANDARD MODEL

L.E. Ibáñez ^{*)}
CERN - Geneva

ABSTRACT

I present an elementary introduction to some topics beyond the standard model (SM). After a short review of some puzzling aspects of the SM, I discuss its grand unification extensions. I give an elementary introduction to $N = 1$ supersymmetry, discuss the construction of the SUSY standard model and its experimental implications. A very brief discussion on superstrings is given at the end of the lectures.

^{*)} On leave of absence from Departamento de Física Teórica C-XI, Universidad Autónoma de Madrid, 28049 Madrid, Spain.

Based on Lectures given at the
1988 ASI School, St. Croix (Virgin Islands)
July 1988

literature as well as to the good number of reviews existing about these topics¹⁻³. Some of these are given in the list of references. The latter is not intended to be complete and I apologize for omissions.

2. PUZZLES IN THE STANDARD MODEL

It is really impressive how the $SU(3) \times SU(2) \times U(1)$ Standard Model (SM) essentially describes all present high energy physics data (except for low P_T strong interactions in which non-perturbative effects play an important role). We are so much used to this model that we very often forget all the aspects of it we do not really understand. The ideal theory for a particle theorist is probably one in which there are essentially no free parameters (all measurable ones are predicted) and an overall beautiful symmetry (as simple as possible) governs all the aspects of the theory. The standard model is far from this situation. The number of parameters "external" to the model (not predicted by it) is large. We have three gauge coupling constants g_1, g_2, g_3 as well as the θ -parameter governing the CP-violating sector ($\theta_{F, \mu\nu}^{\mu\nu}$) of the QCD Lagrangian (the latter is experimentally close to zero, theorists do not know why). To these we have to add the nine masses of quarks and leptons plus the possible Majorana masses of neutrinos (experimentally very small). We further have the three Cabibbo-Kobayashi-Maskawa angles plus the KM CP violating phase. Finally, one has the couplings appearing in the Higgs sector of the model. Thus the standard model is a field theory with a number (~ 20) of free parameters that, when given appropriate (experimental) values, describe remarkably well all known high-energy physics data, no more, no less. It is obvious we should be ambitious and try to explain the values of the ~ 20 free parameters of the model. The efforts of physicists to understand them are often referred to as "beyond the standard model physics". All (but the gauge coupling constraints) are parameters in one way or another related to the Higgs-Yukawa sector of the theory, a sector poorly understood and which has not been tested experimentally. One must admit from the outset that the efforts made in the last 15 years have not been very successful from the point of view of explaining the values of the free parameters of the standard model. However, the particular values of those parameters are not the only points not understood in the SM, as we will discuss below. I believe that in the last 15 years, a lot of conceptual progress in understanding the origin of the SM and its connection with gravity has been achieved.

Let us review now different puzzling qualitative features of the SM. First, let us recall the quantum numbers of each of the three generations with respect to $SU(3)_C \times SU(2)_L \times U(1)_Y$:

$$\left[\begin{array}{l} Q_L \equiv (U, D)_L \rightarrow (3, 2, 1/6) \\ U_L^c \rightarrow (\bar{3}, 1, -2/3) \\ D_L^c \rightarrow (\bar{3}, 1, 1/3) \\ L \equiv (\nu, e^-)_L \rightarrow (1, 2, -1/2) \\ E_L^c \rightarrow (1, 1, 1) \end{array} \right] \quad (1)$$

Altogether each generation contains $3 \times 2 + 3 + 2 + 1 = 15$ Weyl spinors with the above quantum numbers. One of the most relevant facts of the SM is that quark-lepton generations are "chiral". From the mathematical point of view, this means that the group representation of each family [i.e., quantum numbers in (1)] is complex, e.g., there are no left-handed quarks with $(\bar{3}, 2, -1/6)$ quantum numbers. Another way of stating the chirality

1. INTRODUCTION

During the last ten years the particle physics theorists have been bombing their experimental colleagues with new theories, symmetries, particles and promising experiments to be done. This includes grand unification, proton decay, neutrino oscillations, neutron oscillations, neutrino masses, supersymmetry, supergravity, gluinos, squarks, photinos, axions (visible, invisible and in between), fifth forces, heavy leptons, new quarks, extra Z 's, strings, etc. None of these very bright ideas have, up to now, been checked experimentally, and there is a reasonable tendency towards discouragement. However, one should think that the present situation is quite difficult for theorists. The leitmotiv of all those new ideas is to go beyond the standard model and try to understand a considerable number of puzzles left unexplained by that theory. This seems to require information about energy scales for which the experimental data are very scarce (if at all existent). One must hope that present and future accelerators (and underground experiments) will eventually unravel some of the expected new physics. In these lectures, I give an elementary introduction to some of the most popular ideas about extensions of the standard model (SM). This includes grand unification, supersymmetry and strings but the emphasis is put on the second. The reason is that supersymmetry is a theory which can be checked experimentally at accelerators. Unfortunately, the same cannot be said for grand unification and strings which will have to be checked (if at all) indirectly. Since the audience of the lectures was of experimentalists, I tried to be as little technical as possible although this is sometimes unavoidable. In Section 2, I discuss some of the features of the SM which are not well understood. The grand unification idea is discussed in Section 3 in which a simplified introduction to the simplest GUTs is given. The longest part of the lectures, devoted to supersymmetry, is contained in Section 4. After describing the properties of supersymmetric interactions, I present in some detail the supersymmetric version of the SM. Plausible SUSY spectra and experimental signatures are also given. Finally, in Section 5 I give an oversimplified description of string theories, particularly as candidates for unified theories of all interactions. A few final comments are left for Section 6.

These lectures are not intended to be self-contained but just to give a flavour of the topics covered. The reader is referred to the original

property is to say that left- and right-handed quarks and leptons couple differently to the gauge interactions [right-handed fields do not couple to $SU(2)_L$, the charged weak interactions].

An important consequence of chirality is that quarks and leptons cannot have mass terms in the SM Lagrangian. This is obvious because the mass terms of a fermion f is given by

$$= \bar{f}_R^f f_L + h.c. \quad (2)$$

where $f_{L(R)} = ((1\tau_Y)/2)f$, and a term like (2) is not invariant under the $SU(2) \times U(1)$ symmetry (instead of being a singlet transforms like a doublet). In order that quarks and leptons get a mass, one must break $SU(2) \times U(1)$ spontaneously. If there is a scalar (Higgs) with Yukawa couplings to fermions like

$$h \bar{f}_R^f f_L + h.c., \quad (3)$$

quarks and leptons get effective masses $\sim \langle h \rangle$ once the Higgs gets a vacuum expectation value (v.e.v.) $\langle h \rangle \neq 0$.

Why did Nature choose fermions to be chiral? A very symmetric and reasonable (non-chiral) assignment for candidate quarks and leptons would have been:

$$\begin{aligned} Q &\rightarrow (3, 2, 1/6) + (\bar{3}, 2, -1/6) \\ L &\rightarrow (1, 2, -1/2) + (1, 2, 1/2) \end{aligned} \quad (4)$$

but Nature decided to choose the more asymmetric (chiral) assignments of (1). There is no answer to this question within the SM.

Let us look back again at the quantum numbers of (1). Why those funny hypercharge assignments? Since charge Q and the diagonal generator of $SU(2)$, T^3 are related to hypercharge by

$$Q = T^3 + Y, \quad (5)$$

an equivalent way of stating the same question is the following. Why is the charge, e.g., of the d-quark one-third of that of the electron (why are the charges of proton and positron the same)? This is called charge quantization. Apparently there is no reason why this should be so. In fact, one can convince oneself that charge quantization is naturally required by the quantum consistency of the theory, i.e., the absence of gauge "anomalies"⁴. Gauge theories with chiral fermions, like the SM, have potential dangers (breakdown of unitarity of the S-matrix) with the existence of linear divergences in "triangle graphs" of the type shown in Fig. 1. The dangerous graphs involving the gauge boson coupled to Y are shown in that figure. Let us suppose we know the $SU(3) \times SU(2)$ quantum numbers of the quarks and leptons in a family [as in (1)] but we did not know the hypercharge assignments. Consider giving them arbitrary hypercharge values, e.g.

$$Y(Q_L) = a, Y(U_L^c) = b, Y(D_L^c) = c, Y(L) = d, Y(E_L^c) = e. \quad (6)$$

Each of the graphs in Fig. 1 are proportional to

$$\begin{aligned} i) & 3a(1/2) + d(1/2) \\ ii) & 2a(1/2) + b(1/2) + c(1/2) \\ iii) & 6a^3 + 3b^3 + 3c^3 + 2d^3 + e^3 \\ iv) & 6a + 3b + 3c + 2d + e \end{aligned} \quad (7)$$

Fig. 1

For example, the first term in i) is the contribution from Q_L to the triangle loop [factor 3 for three colours, a factor a from Y -boson coupling, a factor $1/2 = \text{tr}(T^3)^2$ from the coupling of two W 's, etc.], whereas the second is the contribution from L . Now, anomaly freedom dictates that all i)-iv) should vanish. Since an overall normalization is irrelevant, we may choose, e.g., $Y(E_L^c) = e = 1$. Then the reader can easily verify that the only solution allowed is

$$a = 1/6, b = -2/3, c = 1/3, d = -1/2, e = 1, \quad (8)$$

i.e., the assignments given in (1) are dictated by the absence of anomalies and so is charge quantization. The reader has probably noticed that the fourth graph involves gravitons⁵. There are reasons to believe that one should consider gravitational anomalies as well as purely gauge anomalies and thus, Eq. (7.iv) has to be reinforced. Anyway, it seems that charge quantization is not that arbitrary, even in the context of the SM, and is just a consequence of quantum consistency (freedom of anomalies by family, which is not strictly necessary. One could imagine a situation in which the contribution of each generation was anomalous but the sum of the contributions of the three families vanished. Why did anomalies have to cancel family by family? Again, this is something which is not understood within the scheme of the SM.

One can wonder about other remarkable qualitative features of the SM fermion content. We have seen that the cancellation of anomalies in the SM occurs in a very intricate way, quark and lepton contributions cancel against each other. A more simple world could be imagined in which quarks would feel only the strong force and leptons the weak force, e.g., families transforming like

$$(3,1,0) + (\bar{3},1,0) + (2,1,1/2) + (2,1,1/2) + \text{etc.} \quad (9)$$

Again this kind of particle contents is not chiral but why did Nature have to choose the chiral one of (1)? Why a gauge group like $SU(3) \times SU(2) \times U(1)$ and no other, to start with? One could think of more symmetric groups like $SU(3) \times SU(3)$ or $SU(2)^4$, etc. There is another very well-known puzzle which is the existence of three copies (generations) of quarks and leptons with identical gauge properties. Why three and not one, two or 13?

Eventually, we would like to understand more quantitative matters like the values of quark-lepton masses and mixing angles. The quark-lepton masses in GeV are shown in Table 1. One can observe three peculiar properties of the spectrum. i) The quark-lepton masses are, in general, very small compared with the natural scale of the masses in the SM. Since all masses are given by v.e.v.'s of the Higgs field, one would guess a natural mass scale to be $\sim M_H \sim 80 \text{ GeV}$. This is not the case and the first generation is even six orders of magnitude lighter. ii) There is a hierarchy of masses amongst the generations (3^{rd} heavier than the second which is heavier than the first). iii) Quarks in each family are heavier than the corresponding lepton. To these three properties of the masses one has to add another important piece of information. The Cabibbo-Kobayashi-Maskawa mixing matrix is, to a good approximation, close to the unit matrix (the biggest off-diagonal element is the Cabibbo angle $\theta \sim 1/5$). Within the scheme of the SM both masses and mixings depend on the shape and size of the Yukawa coupling constants G_D, G_E, G_U appearing in the Lagrangian:

$$G_D^i \bar{q}_L^j q_L^j + G_E^i \bar{q}_L^j q_L^j + G_U^i \bar{q}_L^j q_L^j + \text{h.c.}, \quad (10)$$

where $i, j = 1, 2, 3$ run over the three generations. The values of those couplings are "external" to the model so that we do not have any prediction for them within the SM. Of course, one would like to understand at least the above-mentioned four qualitative features of masses and mixing angles. To achieve that, we necessarily have to go beyond the SM.

All the aspects of the SM described above are probably sufficient to believe that the "theory of everything" cannot be just the standard model. There are further theoretical arguments that point towards difficulties associated with the SM field theory. To start with, the $U(1)$ piece of the SM as well as the Yukawa and Higgs sectors are non-asymptotically free field theories. Then, one knows that at a very large scale the coupling constants will get too large (Landau singularities). Furthermore, there are reasons to believe⁶ that the purely Higgs sector of the theory would be trivial (S matrix = 1). All this indicates that one cannot extrapolate

Table 1

Lepton	$e \rightarrow 5 \times 10^{-4}$	$\mu \rightarrow 1 \times 10^{-1}$	$\tau \rightarrow 1.8$
D-quarks	$d \rightarrow 7 \times 10^{-3}$	$s \rightarrow 1.5 \times 10^{-1}$	$b \rightarrow 4.5$
U-quarks	$u \rightarrow 4 \times 10^{-3}$	$c \rightarrow 1.2$	$t \rightarrow > 45$

the SM to very large energies. On the other hand, one knows that anyhow, when the energies considered are of the order of the Planck mass $M_P \approx 10^{18} \text{ GeV}$, the effective gravitational constant $G \sim E^2/M_P^2 \sim 1$ and gravitational interactions cannot be neglected in any standard model computation.

There is another (aesthetical?) problem for the SM which appears even at scales much below the Planck scale and that is the "naturalness problem". The point is that it is hard to understand how and why the scale of weak interactions is so small compared with the Planck scale, $M_P \ll M_P$. We know that the weak scale M_W is of the order of the Higgs mass parameter we put in our Lagrangian, $M_H \sim |\mu_{\text{Higgs}}|$. We can fix $|\mu_{\text{Higgs}}|$ to be $\sim 100 \text{ GeV}$, as experimentally required. However, radiative loop corrections tend to destabilize that fixing and give large masses to the Higgs scalar. The point is that graphs like those in Fig. 2 give rise to quadratically divergent contributions to the Higgs mass. Of course, one can just choose a renormalization prescription, order by order, in which $|\mu_{\text{Higgs}}| \sim 100 \text{ GeV}$. However, this will require a ridiculous fine-tuning of the counterterms in the Lagrangian in order to get the desired result⁷. This problem can be stated as the violation of a naturalness criterion which goes as follows⁸. "A small parameter (like $|\mu_{\text{Higgs}}|$) is unnatural unless the symmetry is increased by setting it = 0". Setting $|\mu_{\text{Higgs}}| = 0$ does not increase the original symmetry of the SM. Notice that a similar naturalness problem does not exist for fermions. The point is that one can see that all loop contributions to mass renormalization for a fermion are always proportional to the original bare mass. Thus, if that mass is set to zero, it will continue being zero even after radiative corrections. What happens is that when we set the mass of a fermion to zero, a "chiral symmetry" is preserved by which we can rotate f_L and f_R by arbitrary phases (a mass term would violate this). This symmetry is preserved by all the Lagrangian and hence remains true after radiative corrections. Thus, a vanishing fermion mass can be "natural", the symmetry of the Lagrangian increases. This is not the case for a scalar; a mass term like $m^2 \phi^\dagger \phi$ cannot be forbidden by any known consistent symmetry of a Lagrangian.

Fig. 2

In the above sense, the SM is unnatural, the weak scale M_W is unnaturally low. In order to avoid this naturalness problem, many different possibilities have been considered which can be essentially classified into two: compositeness and supersymmetry. The most attractive idea put forward in the first is the possibility that Higgses were not elementary but composite of some new fermions ("techniquarks") bound together by a new strong force ("technicolour").⁹ Although the original idea is very nice, it gets very cumbersome when one tries to implement the details and, in particular, give masses to quarks and leptons. Unless one assumes very peculiar behaviour of the technicolour interaction, one gets wrong predictions for flavour changing neutral current processes (FCNC) (rates too high for reactions like $K^0 \rightarrow \mu^+ \mu^-$, $\mu \rightarrow e \gamma$, etc.).¹⁰ Other ideas involve strongly interacting W bosons,¹¹ or strongly interacting Higgses.¹² Finally, there are "preon" models in which all quarks and leptons are composite. In all these schemes no outstanding model has been found and one typically finds that the number of "preonic" particles one has to introduce, is often bigger than the observed number of quarks and leptons so that not much simplicity is achieved. Furthermore, all the dynamics of these models is unknown so that one cannot really give any model-independent prediction (apart from the expectation that if the confining scale of preons is reached at an accelerator, one would start producing many new particles).

From all the above, the clever reader has probably realized that I am not particularly fond of the compositeness scenarios considered up to now (except, maybe, technicolour). In these lectures, I will discuss the supersymmetry alternative in some detail. There is an additional motivation for that. I have been discussing above a number of puzzles of the SM, but the SM is not all physics. It is well known that eventually gravity has to be included in the scheme and a consistent quantum theory of all interactions including gravity. In this direction, it is known that supersymmetry (particularly superstrings) is quite promising. Furthermore, eventually one would like to answer seemingly more fundamental questions like: why does space-time exist? Why do we live in four dimensions? Why is one of the co-ordinates timelike? etc. To answer these questions, it seems one would need a consistent unified theory of all interactions, including gravity. The work done in the last 15 years (grand unification, supersymmetry, supergravity, superstrings) seems to be on the right track. In the next lecture, I will very briefly discuss these topics emphasizing the chances of an experimental verification. I must remark that none of these ideas has been checked experimentally and so must be considered at the moment just as attractive speculations.

3. THE GRAND UNIFICATION IDEA

Before discussing the supersymmetry solution to the naturalness problem, we are going to consider a beautiful generalization of the SM, grand unified theories.^{13,14} In a remarkable paper¹⁵, Georgi, Quinn and Weinberg pointed out the following. Consider the couplings of the SM model $\alpha_1, \alpha_2, \alpha_3$ ($\alpha_1 = g_1^2/4\pi$) as measured at accelerators. It is well known that the size of the coupling constants depend on the energies at which they are measured ("running coupling constants"). The value at any other scale μ is given by the one-loop renormalization group equations:

$$\frac{1}{\alpha_i(\mu)} = \frac{1}{\alpha_i(M_U)} - 8\pi b_i \log(\mu/M_U), \quad i = 1, 2, 3 \quad (11)$$

where b_i are the coefficients of the corresponding "beta functions" and $\alpha_i(M_U)$ are the measured values at a scale, say, M_U . Plugging the known values for $\alpha_1(M_U)$ and b_i in Eq. (11), one obtains a qualitative scale

Fig. 3

dependence as shown in Fig. 3. The gauge coupling constants seem to get together at an extremely large mass scale $\sim 10^{15}$ GeV, not far away from the Planck mass ($\sim 10^{18}$ GeV). Thus, the following idea seems very plausible: the couplings join because beyond a scale $\sim 10^{15}$ GeV, there is a "unified theory" of the three $SU(3) \times SU(2) \times U(1)$ interactions into a single one.^{13,14} This will have a simple group G gauge invariance such that $G \supset SU(3) \times SU(2) \times U(1)$ and there will be a unique gauge coupling constant, g . The first task is then to look for a simple group G which contains the SM. The Lie algebras of groups were classified a long time ago by Cartan. The associated groups correspond to the $SU(n)$, $SO(n)$ (n odd), $SO(n)$ (n even) and $Sp(n)$ ("symplectic") classical groups. There is also a finite number (5) of Lie groups which are called "exceptional" and denoted G_2, F_4, E_6, E_7, E_8 . The rank (number of commuting diagonal generators) of the SM model is four [two from $SU(3)$ + one from $SU(2)$ + one from $U(1)$] so that the candidate group needs to have a rank at least equal to four. Cartan's classification for simple algebras of rank > 4 is shown in Table 2. A further constraint one has to impose to get a chiral spectrum is that the group must admit "complex representations" (recall our statements about chirality in the previous section). As it turns out, only the $SU(N)$, $SO(2n+4)$ and E_6 groups have complex representations and then we are left only with the groups inside a square in Table 2. The group of minimal rank is $SU(5)$ and this is the simplest candidate. For rank 5, we have $SO(10)$ and $SU(6)$ and for rank 6 we have E_6 and $SU(7)$. The $SU(6)$ and $SU(7)$ groups do not have any particular advantage over their subgroup $SU(5)$ so that it seems that $SU(5)$, $SO(10)$ and E_6 are the most attractive groups for "grand unification". Let us discuss them briefly in turn:

1) $SU(5)$. This is the simplest possible grand unification scheme.¹³ It has $5^2 - 1 = 24$ gauge bosons which we can schematically put in a matrix:

$$24 = \begin{pmatrix} \begin{matrix} X_1^+ Y_1^- \\ X_2^+ Y_2^- \\ X_3^+ Y_3^- \end{matrix} & \begin{matrix} Z^0 W^- \\ W^+ \gamma \end{matrix} \\ \begin{matrix} X_1^- X_2^+ \\ X_1^- X_3^+ \end{matrix} & \begin{matrix} Z^0 W^- \\ W^+ \gamma \end{matrix} \\ \begin{matrix} X_2^- X_3^+ \\ Y_1^+ Y_2^+ \\ Y_1^+ Y_3^+ \end{matrix} & \begin{matrix} Z^0 W^- \\ W^+ \gamma \end{matrix} \end{pmatrix} \quad (12)$$

gluons

Table 2

RANK	CLASSICAL	EXCEPTIONAL
4	$SU(5)$	$SO(8)$ F_4
5	$SU(6)$	$SO(10)$
6	$SU(7)$	$SO(12)$ E_6
7	$SU(8)$	$SO(14)$ E_7
8	$SU(9)$	$SO(16)$ E_8
N	$SU(N+1)$	$SO(2N)$
	$SO(2N+1)$	$Sp(2N)$
	$SO(2N+4)$	$SO(2N+4)$

Apart from the known ¹² gauge bosons (8 gluons + W^+, W^0, Z^0), there are 12 new ones X_i^+, Y_i^+ ($i = 1, 2, 3$ colours) which are triplets of colour and $SU(2)$ doublets. They have fractional charges $\pm 4/3, \pm 1/3$ and couple both to quarks and leptons. Since the complete $SU(5)$ symmetry is only realized above the unification scale $M_X \sim 10^{15}$ GeV, that will be the order of magnitude of the masses of X_i^+ and Y_i^+ . Thus, below M_X the $SU(5)$ symmetry will be spontaneously broken by some Higgs mechanism down to $SU(3) \times SU(2) \times U(1)$, and there will be essentially no obvious trace of the $SU(5)$ symmetry.

The quarks and leptons of each family in $SU(5)$ sit in a five-plet and a ten-plet under $SU(5)$:

$$\underline{5} = \begin{bmatrix} d_f \\ d_f^c \\ e \\ \nu_e \end{bmatrix}_L; \quad \underline{10} = \begin{bmatrix} 0 & u_f^c & -u_f^c & u_1 & d_1 \\ 0 & u_f^c & u_2 & d_2 & d_3 \\ 0 & u_3 & d_3 & e^c & 0 \end{bmatrix}_L \quad (13)$$

where we have represented the ten-plet by an antisymmetric 5×5 matrix (this multiplet results from the antisymmetric product of two five-plets and thus has $5 \times (5-1)/2 = 10$ independent components). These assignments may

look bizarre but one can convince oneself that a $(10+5)$ representation is the simplest chiral, anomaly-free fermion content within $SU(5)$. Notice that the 15 states of each SM generation just fit. There is no place for more quarks or leptons in (13). This is a remarkable fact.

Since the new gauge bosons X_i^+ and Y_i^+ have both colour and $SU(2)$ quantum numbers, they can convert quarks into leptons and also quarks into antiquarks (i.e., they violate baryon and lepton numbers). Thus processes like the ones in Fig. 4 are possible.

Fig. 4

They give rise to proton decay with a lifetime

$$\tau_p \sim \frac{M_X^4}{m^5} \sim 10^{30} \text{ years} \quad (14)$$

where m is a mass scale of order of the mass of the proton.

A model like $SU(5)$ has two widely different symmetry breaking scales, $M_X \sim 10^{15}$ GeV and $M_W \sim 10^2$ GeV:

$$SU(5) \xrightarrow{M_X} SU(3)_C \times SU(2)_L \times U(1)_Y \xrightarrow{M_W} SU(3)_C \times U(1)_Q \quad (15)$$

The first breaking is assumed to be produced by a Higgs mechanism triggered by a v.e.v. of a scalar field transforming like the adjoint representation of $SU(5)$. This will be a field with $5^2-1 = 24$ components whose v.e.v. can be represented in matrix form as:

$$\langle \phi^a, b \rangle = \begin{bmatrix} v & & 0 \\ & v & \\ 0 & -3v/2 & -3v/2 \end{bmatrix}, \quad v \sim 10^{15} \text{ GeV} \quad (16)$$

There is an explicit $SU(3) \times SU(2)$ symmetry left unbroken in (16). The second breaking at the weak scale will proceed as in the usual SM, with an $SU(2)_L$ doublet of Higgses. Within the original $SU(5)$ structure that doublet is contained in a five-plet H^a ($a = 1-5$) of Higgses.

ii) $SO(10)^{16}$. This gauge group contains $SU(5)$:

$$SO(10) \supset SU(5) \times U(1) \quad (17)$$

and thus it is convenient to analyze it in terms of this subgroup. [Other interesting subgroups are $SU(4) \times SU(2)_L \times SU(2)_R$ and $SU(3)_C \times SU(2)_L \times SU(2)_R \times U(1)_{B-L}$.] $SO(10)$ has $10 \times (10-1)/2 = 45$ gauge bosons which in terms of the $SU(5)$ subgroup transform as

$$45 = \underline{24} + \underline{10} + \underline{1} \quad (18)$$

There are then 33 extra gauge bosons in this model. An attractive feature of SO(10) is that [unlike SU(5)] all particles within a family fit in the simplest SO(10) chiral multiplet which has dimension 16:

$$16 = (\underbrace{\nu, u_1, u_2, u_3; e, d_1, d_2, d_3}_{\text{LEFT}}; \underbrace{d_3^c, d_2^c, d_1^c, e^c; u_3^c, u_2^c, u_1^c, \nu^c}_{\text{RIGHT}})_L \quad (19)$$

and has an explicit left-right symmetry. This implies the existence in each generation of an extra particle (the 16th fermion which is not in the SM) which has the quantum numbers of a right-handed neutrino ν_L^c (it is a singlet under the standard model interactions but it has lepton number).

As happened with SU(5), the extra gauge bosons, in general, violate baryon and lepton numbers and hence proton is again unstable with a lifetime $\sim 10^{30}$ years.

Since the rank of SO(10) is five, there are more symmetry breaking possibilities. One in general needs at least two different Higgs fields to do the breaking of SO(10) down to the SM and this breaking may proceed in several stages. Some symmetry breaking possibilities are¹⁴:

$$\begin{aligned} \text{SO}(10) &\xrightarrow{\phi, \phi' \neq 0} \text{SU}(3)_C \times \text{SU}(2)_L \times \text{U}(1)_Y \\ \text{SO}(10) &\xrightarrow{\phi \neq 0} \text{SU}(3)_C \times \text{SU}(2)_L \times \text{SU}(2)_R \times \text{U}(1)_{B-L} \rightarrow \text{SU}(3)_C \times \text{SU}(2)_L \times \text{U}(1)_Y \quad \phi' \neq 0 \quad (20) \\ \text{SO}(10) &\xrightarrow{\phi \neq 0} \text{SU}(4) \times \text{SU}(2)_L \times \text{SU}(2)_R \rightarrow \text{SU}(3)_C \times \text{SU}(2)_L \times \text{U}(1)_Y, \text{ etc.} \end{aligned}$$

where the $\text{U}(1)_{B-L}$ in (20) corresponds to a gauge boson which couples to baryon-lepton number. The latter is gauged within SO(10). We must remark that the symmetry breaking in SO(10). We must remark that the symmetry breaking in SO(10) requires scalar fields with relatively high dimensionalities (ϕ has dimension 45 or 54 whereas ϕ' has dimension 16 or 126).

ii) E_6 . This gauge group¹⁸ contains SO(10)

$$E_6 \supset \text{SO}(10) \times \text{U}(1) \quad (21)$$

so that it is useful to analyze it in terms of that subgroup. [Other interesting subgroups include $\text{SU}(3)_C \times \text{SU}(3)_L \times \text{SU}(3)_R$.] E_6 has 78 gauge bosons which in terms of the SO(10) subgroup have the multiplet structure

$$78 = \underline{45} + \underline{16} + \underline{1\bar{6}} + \underline{1} \quad (22)$$

There are then plenty of extra gauge bosons which, as with SU(5) and SO(10), can mediate proton decay. A complete generation of quarks and leptons fit in the fundamental multiplet of E_6 which has dimension 27. However, in this case an E_6 generation contains many (12) extra yet unobserved quarks and leptons. In terms of SO(10), and E_6 generation contains:

$$27 = \underline{16} + \underline{10} + \underline{1} \quad (23)$$

i.e., apart from an SM generation, one has two neutrino-like objects (ν_L^c and N) and new quarks ($g+\bar{g}$) and leptons ($\lambda+\bar{\lambda}$). Here, $g(\lambda)$ has the $\text{SU}(3) \times \text{SU}(2) \times \text{U}(1)$ quantum numbers of a right-handed d-quark (left-handed lepton).

Again since the rank of E_6 is large (6), there are plenty of possibilities for breaking the symmetry down to the standard model. All the large symmetry breaking may be done with a single scalar field of quite a large multiplicity, 351. This finishes our short review on SU(5), SO(10) and E_6 grand unified theories (GUTs).

There are a number of properties and predictions which appear in most GUTs. They include the following:

a) Charge quantization. Since the charge generator is $Q = Y + T^3$ and both $\text{SU}(2)_L \times \text{U}(1)_Y$ are included amongst the generators of the unifying simple group G , Q has to be traceless, like any other G generator. This implies then the same condition $[\text{Tr}(Q) = 0]$ that we found from imposing absence of the mixed hypercharge-gravitational anomaly (Fig. 1.iv). Thus charge quantization is an automatic property in GUTs.

b) Glashow-Weinberg angle prediction. Since the three coupling constants g_1, g_2, g_3 are assumed to join at the "grand unification mass" M_X , one can compute what is the latter by finding at what scale, e.g., g_2 and g_3 join. That is done by using the renormalization group equation in (11). Once we know the actual value of M_X , the value of the third coupling constant (e.g., g_1) may be computed by running it from M_X down to low energies where experiments are performed. Instead of using $\alpha_1, \alpha_2, \alpha_3$, it is more convenient to use the fine structure constant $\alpha_e, \sin^2 \theta_W$ and α_3 . The above computations yield (in leading log approximation)

$$\log\left(\frac{M_X}{M_W}\right) = \left(\frac{1}{\alpha_e(M_W)} - \frac{8}{3\alpha_3(M_W)}\right) / 8\pi\left(\frac{1}{3}b_1 + b_2 - \frac{8}{3}b_3\right) \quad (24a)$$

$$\sin^2 \theta_W(M_W) = \frac{3}{8} - 5\pi\alpha_e(M_W) (b_1 - b_2) \log\left(\frac{M_X}{M_W}\right) \quad (24b)$$

Assuming that the GUT symmetry breaks directly to the SM (no intermediate symmetry breaking), one finds a value for $M_X = 10^{14-15}$ GeV, implying a value of the Weinberg angle $\sin^2 \theta_W = 0.20$, which may be considered a successful prediction (we will further comment on this below). Notice that in models [like SO(10) or E_6] in which an intermediate scale of symmetry breaking is possible, Eqs. (24) get modified and there is more flexibility to accommodate different values of $\sin^2 \theta_W$.

c) Quark-lepton mass relations. In GUTs, quarks and leptons live in the same multiplets and one expects to find relationships between the Yukawa couplings of quarks and leptons. Thus, e.g., in SU(5) one has the following couplings¹⁹ between the fermions in $\underline{5} = (d_1^c, L)$ and $\underline{10} = (Q_L, u_1^c, e^c)$:

$$G_{D,L}^{ij} \bar{\psi}_1^j \phi_{10,H} + G_{U,L}^{ij} \bar{\psi}_{10}^j \phi_{10,H}^* + \text{h.c.} \quad (25)$$

Notice that we only have two independent terms instead of the three in the standard model [Eq. (10)]. We then have equal Yukawa couplings for d-quarks and leptons ($G_L = G_D$) as long as SU(5) is a good symmetry. Once SU(5) is broken, there are radiative corrections which contribute differently to quarks and leptons (in particular gluon exchange). Computing these radiative effects and using the renormalization group, one gets¹⁹

$$\frac{m_D}{m_L} = \frac{h_D(m_D)}{h_L(m_L)} = \frac{h_D(M_X)}{h_L(M_X)} \times \left(\frac{\alpha_3(M_D)}{\alpha_3(M_X)}\right)^{4/7} \approx 3.0-4.0 \quad (26)$$

which does compare well with the ratios m_b/m_c and m_s/m_c , but is definitely wrong for the first generation. However, one may argue that the first generation has some additional sources for its mass coming, e.g., from different type of Yukawas which makes it different. In $SO(10)$ the Higgs Weinberg-Salam doublets are contained in the fundamental multiplet, a ten-plet. Since all quarks and leptons are in the same multiplet, a 16-plet, there is only one Yukawa term:

$$G^{\frac{1}{2}} \psi_{16} \psi_{16} H_{10} + \text{h.c.} \quad (27)$$

However, H_{10} contains two $SU(2)$ doublets, one (H) coupling to d-quarks and charged leptons and the other ($\bar{H}$) coupling to U-quarks and neutrinos. Then, one has $m_{U,L} \approx G\langle H \rangle$ but $m_{U,\nu} \approx G\langle \bar{H} \rangle$. This gives us the same prediction for d-quarks and leptons as $SU(5)$ but also an additional prediction $m_t/m_b \approx m_c/m_s$ [of course, the renormalization effects, as it happened in Eq. (26), have to be taken into account]. Such a prediction is still not ruled out experimentally. Unfortunately, if the model contains one single ten-plet of Higgses, the mass matrices for U and D-quarks are going also to have identical shape and may be diagonalized by the same unitary matrices. U_U and U_D acting on the left-handed quarks (U_L, D_L). Since the Cabibbo-Kobayashi-Maskawa matrix $U = U_U^\dagger U_D = 1$, there would be no mixing at all in the simplest $SO(10)$ model. On the other hand, this could be considered as a good feature of the model, since after all the CKM matrix is quite similar to the unit matrix. Again one could argue that some other effects (maybe loop contributions?) slightly modify the CKM matrix and gives non-vanishing off-diagonal entries. E_6 gives results similar to those of $SO(10)$.

d) Neutrino masses. When discussing quark-lepton mass relationships in $SO(10)$, we saw that one expects $m_\nu = m_\nu$ in the simplest model. This looks like a disaster since present limits for neutrino masses are in the range $20 \text{ m}_\nu \lesssim 15 \text{ eV}$, $m_{\nu_\mu} \lesssim 150 \text{ keV}$, $m_{\nu_\tau} \lesssim 150 \text{ MeV}$. However, the above $SO(10)$ prediction refers to the "Dirac ν -mass", which corresponds to a term in the Lagrangian:

$$m_\nu \bar{\nu}_R \nu_L + \text{h.c.} \quad (28)$$

and involves both ν_L and ν_R (or equivalently, ν_L^c). However, the right-handed neutrino ν_R may also have a "Majorana" mass, a term of the form

$$M \nu_L^c \nu_L^c + \text{h.c.} \quad (29)$$

which involves only the ν_R . In fact, the last term is allowed by the $SU(3) \times SU(2) \times U(1)$ symmetry [unlike the one in Eq. (28)] and hence M may be as large as the unification mass (unlike m_ν which is at most $\sim M_U$). The natural value for M is $\sim M_X$ and that is usually the case in realistic models (anyway, one usually has $M \gg M_U$). We thus have a neutrino mass matrix for each generation²¹

$$M \sim \begin{bmatrix} \nu_L & \nu_L^c \\ \nu_L & 0 \\ \nu_L^c & M \end{bmatrix} \begin{bmatrix} \nu_L \\ \nu_L^c \\ M \end{bmatrix} \quad (30)$$

which has eigenvalues $\sim m_\nu^2/M$ and $\sim M$. Thus there is an eigenstate (essentially ν_L^c) with a very large mass $\sim M$ and hence undetectable. The other eigenstate (mostly the ν_L) has a mass $\sim m_\nu^2/M$. Notice that for $m_\nu \sim m_\mu \sim 1 \text{ GeV}$ and $M \sim M_X \sim 10^{15} \text{ GeV}$, one expects neutrino masses

extremely small, $\sim 10^{-6} \text{ eV}$. If M is smaller [as happens with $SO(10)$ models with an intermediate $SU(2)_R$ symmetry], the values for neutrino masses may be close to the experimental limits.

A similar situation occurs for E_6 . In the simplest version of $SU(5)$, there is no room for a ν_L in the multiplet $10+5$ so that fermion cannot be Dirac neutrino masses. One may of course add a singlet fermion to the $SU(5)$ model to play the role of ν_L but that would be a bit artificial. An interesting effect which may appear if neutrinos have a mass is neutrino oscillations (e.g., $\nu_e \leftrightarrow \nu_\mu, \nu_\tau$ transitions). This may happen because, if the ν 's have a mass, there will be in general a mixing matrix for them similar to the CKM matrix. Then, a neutrino produced in a weak interaction process will not be, in general, a mass eigenstate and the proportion of ν_e, ν_μ and ν_τ in a produced ν -state will oscillate with time as it propagates.²² Experimental searches for both neutrino masses and oscillations have been unsuccessful up to now²².

e) Baryon number violation. As we mentioned before, all grand unified theories predict baryon number violating processes like proton decay due to the exchange of new heavy gauge bosons. The expected lifetime τ very much depends on details of each model. In particular, τ goes like the fourth power of M_X and e.g., a factor 1.8 in the value of M_X leads to a τ an order of magnitude larger. The minimal version of $SU(5)$ (with just a 5 of Higgses and the usual three generations) has been in fact already ruled out by proton-decay experiments²³. However, since the dependence of τ on M_X is so strong it is easy to slightly modify the model (by adding, e.g., some new scalar or fermion²⁴) in such a way which is not ruled out by present experimental limits. Another baryon-number violating process which may appear in GUTs is neutron-antineutron oscillations ($n \leftrightarrow \bar{n}$) or $n\text{-}\bar{n}$ annihilation in nuclei. This is a process whose appearing or not is more model-dependent in relation with the particular Higgs particle content of the model under consideration¹.

The above are general features appearing in almost any model. Can one consider that the grand unification idea has been a success? Probably the most remarkable fact of GUTs [particularly $SU(5)$ and $SO(10)$] is how well a quark-lepton generation fits into unified multiplets. Charge quantization is also automatic in GUT theories. Another attractive point is that GUTs give a (modest) reduction in the number of couplings since g_1, g_2, g_3 are related by their unification at M_X . This gives the prediction for $\sin^2 \theta_W$ which we mentioned above and can be considered as a semi-quantitative success. In fact, detailed computations of $\sin^2 \theta_W$ (including higher order effects) in the minimal $SU(5)$ model give a result $\sin^2 \theta_W = 0.214 \pm 0.003$ to be compared with the experimental world average $\sin^2 \theta_W^{\text{ex}} = 0.228 \pm 0.0044$, which seems to be about three standard deviations away²⁵. However, we repeat again that slight changes of the model can accommodate the experimental value. Thus, the prediction for the Weinberg angle may be considered as a semi-quantitative success. Concerning the predictions for quark-lepton mass ratios, the success (if any) is very limited since they certainly yield wrong results for the lightest generation. On the other hand, GUTs predict the observed experimental fact that quarks are heavier than their lepton partners within each family. This also points towards the existence of a "big desert" between the weak scale and a big scale sufficiently high so that the renormalization effects (logarithmic) have time to give a large contribution. Anyway, the grand unification idea, as sketched in the previous lines, is probably too naive. It completely ignores the gravitational interactions which are known to become important at a scale $\sim 10^{18} \text{ GeV}$, very close to the hypothetical grand unification mass. It sounds very unlikely that those two scales were not related and if they are, how can we ignore the effects of gravity? Since gravity is not an

interaction like the others, no consistent quantum gravity field theory is known, one expects that new physics will govern the combined theory of the four interactions. Nevertheless, the above-mentioned attractive features (successes?) of GUTs seem to indicate that they may be partially right. It is plausible that GUTs may be just good "classification symmetries", in the same way that the SU(3) and SU(4) global flavour symmetries of the strong interactions are. Then GUT symmetries may be useful tools to study unified theories including gravity. This is in fact what happens in string theories in which, in general, one does not get a GUT structure of the type described above but grand unification tools are very useful in searching for potentially realistic strings. In the meantime GUTs have stimulated the search of baryon and lepton number violations, existence of monopoles, etc. whose experimental check is of the utmost importance.

4. SUPERSYMMETRY²

Let us come back now to the naturality problem we were discussing in Section 2. The problem is that there is no symmetry forbidding the appearance of a large mass term $m^2(H^\dagger H)$ for the Higgs scalar, ruining the electroweak symmetry breaking. Supersymmetry²⁶ is an exception to this fact. It introduces a concept of "chirality" for scalars by assigning a fermionic bispinor of definite handedness to each complex scalar. This is the simplest version of supersymmetry ("N=1") in which any complex scalar has a fermionic partner. As we will discuss, this symmetry between bosons and fermions leads to cancellations which improve the ultraviolet behaviour of Green's functions. Thus, for example, the quadratic divergences appearing in the graphs of Fig. 2 are exactly cancelled by the diagrams in Fig. 5 in which "supersymmetric partners" (denoted by a hat) of the standard particles circulate in the loops. The cancellation comes about because the coupling constants in Figs. 2 and 5 are equal and because of the relative minus sign between fermionic and bosonic loops. We should nevertheless remark that supersymmetry ("SUSY") by itself does not solve the "hierarchy problem". An explanation why the Weinberg-Salam (WS) Higgs doublet is light compared to the Planck mass is still lacking. SUSY just tells us that if we find a reason why the WS doublet is light, its lightness will be protected from radiative corrections²⁷.

We will consider in what follows only $N = 1$ supersymmetric theories. Theories with N up to eight can be formulated. However, only $N = 1$ SUSY is chiral in the sense we discussed above, i.e., only $N = 1$ SUSY allows that left-handed and right-handed quarks and leptons transform differently under the gauge group. Let us now give a telegraphic introduction to the supersymmetry concept.

Fig. 5

4.1. Supersymmetric field theory

Supersymmetry is a symmetry between bosons and fermions. A dynamical system has supersymmetry if it is invariant under a "SUSY rotation" that changes fermions into bosons and vice versa. This requires having the same number of bosonic and fermionic degrees of freedom. The simplest such system one can think of in field theory is the set of fermion bispinor ψ_α ($\alpha = 1, 2$) and a complex scalar ($\phi = A+iB$). The free massless Lagrangian of these two fields is itself supersymmetric

$$L_0 = \partial_\mu \phi^\dagger \partial^\mu \phi + i \bar{\psi} \not{\partial} \psi \quad (31)$$

This system has a Noether current which is conserved once one applies the equations of motion. The conserved current is

$$J_\alpha^\mu = (\not{\partial} \phi \sigma^\mu \psi)_\alpha ; \quad \partial_\mu J_\alpha^\mu = 0 \quad (32)$$

where $\sigma^\mu = (\mathbb{1}, \vec{\sigma})$ and $\alpha = 1, 2$ is a bispinor index. The conservation of this current implies the conservation of a charge:

$$Q_\alpha \equiv \int dx^3 J_\alpha^0 \quad (33)$$

Something which is unusual about this charge operator, is that it is fermionic [see Eq. (32)] and it is not Lorentz scalar; it transforms as a bispinor.

Up to now the symmetry induced by the charge Q does not look very useful since it refers to a trivial Lagrangian like the one considered in Eq. (31). What makes SUSY interesting is that such a symmetry can be defined for interacting Lagrangians. Thus if one considers, e.g.,

$$L = L_0 - h^2 |\phi|^4 - h(\phi \psi \psi + \text{h.c.}) \quad (34)$$

one can check again that a conserved current exists

$$J_\alpha^\mu = j_\alpha^\mu + h \sigma_\alpha^\mu{}_{\beta\gamma} (\phi^*)^2 \bar{\psi}^\beta \quad (35)$$

The Lagrangian in Eq. (34) is the one of the "massless Wess-Zumino model". It involves a Yukawa as well as a $|\phi|^4$ interaction. In order to obtain a conserved current, it is crucial that the Yukawa coupling constant h is the "square root" of the $|\phi|^4$ coupling. This is an example of the relationships between coupling constants which appear in SUSY theories. We will show below how to write down Lagrangians which are automatically supersymmetric. Before doing that, let us consider again the conserved fermionic charge Q . If $\bar{Q}$ is the complex conjugate one can form the anti-commutator $\{Q, \bar{Q}\}$. Since both Q and $\bar{Q}$ are conserved, the anti-commutator will be conserved. The only conserved quantity one can think of is the four-momentum p^μ contracted with σ^μ in order to get the right spinor structure. In fact, one has

$$\{Q_\alpha, \bar{Q}_\beta\} = 2p^\mu \sigma_{\alpha\beta}^\mu \quad (36)$$

Thus we see that the generator Q is peculiar in yet another aspect. It is not an internal symmetry generator like, e.g., the ones of the standard model symmetry $SU(3) \times SU(2) \times U(1)$, it is related to the generator of space-time translations, p^μ . In a loose sense, Q is the "square-root" of the generator of translations.

In order to do physics with this symmetry, we have to realize supersymmetry on fields. This will allow us to look for quantum field theories with actions invariant under supersymmetry. In the same way that one realizes Poincaré symmetry on fields with definite spin and mass [scalars $\phi(x)$, fermions $\psi_\alpha(x)$, gauge bosons $A^\mu(x)$, graviton $g^{\mu\nu}(x)$, etc.], one realizes supersymmetry on SUSY multiplets ("chiral multiplet", "vector multiplet", etc.). In the same way that the scalar field is the simplest object in usual field theory, the "chiral multiplet" is the simplest supersymmetric object. A chiral multiplet has the following field content:

$$\begin{aligned} \phi &= \text{complex scalar} \\ \psi_\alpha &= \text{bispinor} \\ \bar{F} &= \text{complex scalar (auxiliary)} \end{aligned} \quad (37)$$

The field F is there for technical reasons but it disappears from the action once one uses the equations of motion, so that we will skip it in what follows (it is unphysical). Notice that we denote with the same symbol the whole multiplet and its scalar component. A supersymmetric transformation "rotates" the "components" ϕ and ψ into each other. The infinitesimal "angle" of such a "rotation" has to be peculiar. It has to be a "fermionic" parameter since it transforms fermions into bosons and vice versa

$$\phi \rightarrow \phi + \theta\psi \quad (38a)$$

$$\psi \rightarrow \psi + i\theta\bar{F} \quad (38b)$$

where θ_α ($\alpha = 1, 2$) is the fermionic (anticommuting) infinitesimal parameter of the SUSY transformation. From Eq. (38a) one observes that the θ parameters have mass dimension $= -1/2$ to match the different dimensions of ϕ ($=1$) and ψ ($=3/2$). In (38b) the "rotation" of ψ into $\bar{F}$ requires a derivative ∂ in order to get the correct dimensions. The chiral multiplet is the building block for the supersymmetric matter, i.e., quarks and leptons will be contained in chiral representations. Thus, e.g., for the left-handed electron, we will have the chiral multiplet

$$\phi_{e_L} = (\tilde{e}_L, e_L) \quad (39)$$

where $\tilde{e}_L$ denotes the scalar partner of the left-handed electron (the "electron"). Notice again that a chiral multiplet contains a fermion bispinor (Weyl spinor) with definite chirality and a complex scalar superpartner. The natural object to describe a SUSY multiplet is a "superfield". These superfields unify in a single mathematical object the different bosonic and fermionic components of a supersymmetric multiplet. This simplifies very much the writing of SUSY actions as well as perturbative calculations. However, due to the elementary nature of these lectures, I will not describe here the "superspace" approach to $N = 1$ SUSY. Furthermore, once SUSY is broken (as it eventually should), the superfield approach is not particularly useful for phenomenological computations.

How are the interactions between supersymmetric multiplets? There are two different types of interactions which can be written for a renormalizable $N = 1$ globally supersymmetric theory: a) "chiral interactions" which are interactions between chiral multiplets including Yukawa couplings, $\lambda\phi^4$ interactions and mass terms; b) "gauge interactions" which we will see also include new gauge Yukawa couplings and gauge ϕ^4 couplings.

Let us describe first the chiral interactions. Take a set of chiral multiplets $\phi_i = (\phi_i, \psi_i, F_i)$ where i denotes any index numerating the different chiral multiplets or a gauge group label. The chiral interactions amongst these multiplets are completely determined by a function of the scalars ϕ_i which is called superpotential $W(\phi_i)$. It is important that W does not depend on ψ_i (it is an analytic function). The most general form for $W(\phi_i)$ compatible with renormalizability is

$$W(\phi_i) = \lambda_{ijk} \phi_i \phi_j \phi_k + m_{ij} \phi_i \phi_j + f_i \phi_i \quad (40)$$

This superpotential gives rise to a fermionic Lagrangian and a scalar potential. The Lagrangian relevant to fermions may be obtained in the following manner:

a) Fermionic chiral Lagrangian:

$$L_F = \sum_{i,j} \frac{\partial W}{\partial \phi_i} \bar{\psi}_i \psi_j + \text{h.c.} \quad (41)$$

This gives rise to Yukawa couplings amongst the ψ 's and the ϕ 's and also to fermion masses. This is obvious from Eq. (40). The λ 's turn out to be the Yukawa coupling constants and the m 's the fermion masses. As an example, let us consider the superpotential $W_e(\tilde{e}_L, \tilde{e}_R, H)$ which gives rise to the usual Yukawa coupling of the Weinberg-Salam model for the electron. This is

$$W_e = h_e \tilde{e}_L \tilde{e}_R H \quad (42)$$

where $\tilde{e}_L$ ($\tilde{e}_R$) is the scalar partner of the left-handed (right-handed) electron and H is the Higgs field. The corresponding term in the Lagrangian from Eq. (41) will be

$$L_F = h_e (e_L \tilde{e}_R) H + h_e (e_R \tilde{e}_L) \tilde{H} + h_e (e_L \tilde{H}) \tilde{e}_R + \text{h.c.} \quad (43)$$

Fig. 6

where $\tilde{H}$ is the fermionic partner of the Higgs scalar. Thus we observe that the usual Yukawa coupling comes along with two new Yukawa couplings with (this is important) the same coupling constant h_e . These new couplings are required by supersymmetry. In fact, the three vertices are obtained one from another by changing two of the three particles into their SUSY partners.

The scalar chiral interactions are obtained from the superpotential W as follows.

b) Scalar potential (chiral part)

$$V(\phi_i) = \sum_i \left| \frac{\partial W}{\partial \phi_i} \right|^2 \quad (44)$$

where the sum runs over all the scalars in the theory. In the example of the electron Yukawa superpotential one gets

$$V_e = |h_e \tilde{e}_R H|^2 + |h_e \tilde{e}_L H|^2 + |h_e \tilde{e}_L \tilde{e}_R|^2 \quad (45)$$

Fig. 7

From formulae (43) and (45) one observes that the supersymmetrization of the usual Yukawa for the electron requires the introduction of five new couplings (Figs. 3b-c and 4a-c) but there is only one independent coupling constant h_e , as in the non-SUSY version.

Formulae (41) and (44) are completely general and the specific dynamics is given by the particular W that we are given. We will see later on what is the superpotential of the SUSY standard model. Notice that in a SUSY theory Yukawa couplings and ϕ^4 couplings come together and one cannot consider them separately.

The second type of renormalizable interactions present in $N = 1$ supersymmetric theories are SUSY gauge interactions. This involves usual gauge interactions plus new SUSY interactions as we will show below. The gauge bosons in a SUSY theory are contained in "vector multiplets", the next to simplest multiplets after the chiral ones. Vector multiplets have the following field content:

$$\begin{aligned} \lambda_a &= \text{bispinor ("gaugino")} \\ V(\lambda_a, V^\mu, D; c, \chi_a, N) &= \text{gauge boson} \\ D &= \text{real scalar (auxiliary)} \end{aligned} \quad (46)$$

The fields c , χ_a and N are unphysical fields which can be made to disappear in a convenient "gauge", the so-called "Wess-Zumino gauge". One can introduce a generalization of the usual concept of gauge transformations of fields and gauge bosons to chiral multiplets and vector multiplets. These SUSY gauge transformations allow for the possibility of "gauging away" the unphysical fields c , χ_a and N . Concerning the auxiliary field D , it has again dimension $= 2$, as it happens with the

chiral auxiliary field F . Thus D is also a non-propagating unphysical field and only the gauge boson V^μ and its partner the gauginos λ_a remain as physical fields. All the rest are unphysical fields needed for technical reasons.

What are the SUSY gauge interactions? As an example, let us consider $SU(n)$ gauge interactions of a vector multiplet V with chiral multiplets ϕ_i :

$$\begin{aligned} V^a &= (\lambda_a^\alpha, V_\mu^a, D^a) \quad a = 1, \dots, n^2 - 1 \\ \phi_i &= (\phi_i, \psi_i^\alpha; F_i) \end{aligned} \quad (47)$$

The vector multiplet transforms as an adjoint of $SU(n)$ and the ϕ_i as any $SU(n)$ multiplet. Recall that $N = 1$ SUSY partners always transform the same under the gauge group (SUSY commutes with gauge symmetry). The SUSY gauge interactions are the following. First, there are the usual gauge couplings of gauge bosons amongst themselves and also coupled to fermions and scalars. In diagrammatic form, the usual couplings are

Fig. 8

In order that the gauge interactions be invariant under supersymmetry, one has to add the following new SUSY gauge couplings:

Fig. 9

The coupling of Fig. 9a is just the obvious gauge coupling of gauge bosons to gauginos, where the $SU(N)$ factors are the same as in Fig. 8a, [T is the $SU(N)$ generator matrix]. The coupling of Fig. 9b is a sort of "gauge Yukawa" where the $SU(N)$ factors are as those in Figs. 8c,d. Finally, Fig. 9c represents a "gauge contribution to the scalar potential"

$$V_g(\phi_i) = g^2/2 |\phi_i^\dagger T_{ij}^a \phi_j|^2 \quad (48)$$

where T_{ij}^a are the $SU(N)$ generators written in the representation under which the ϕ_i transform.

These complete the list of interactions present in a supersymmetric gauge theory. The reader will get more acquainted with them later on when we apply them to the case of the supersymmetric standard model. Now we can write down the complete scalar potential for a supersymmetric theory with both chiral and gauge multiplets. From Eqs. (44) and (48), one gets

$$V(\phi_i) = \sum_i |\frac{\partial W}{\partial \phi_i}|^2 + \frac{g^2}{2} |\phi_i^\dagger T_{ij}^a \phi_j|^2 \quad (49)$$

This is sometimes written in the literature as

$$V(\phi_i) = \sum_i |\nabla_i W|^2 + \frac{1}{2} |D^a|^2 \quad (50)$$

This is done because one can show that the equations of motion of the complete Lagrangian give

$$F_i^* = -\frac{\partial W}{\partial \phi_i} \quad (51a)$$

$$D^a = -g\phi_i^\dagger T_{ij}^a \phi_j \quad (51b)$$

for the auxiliary fields F_i and D^a . Equations (51a,b) are precisely the ones which show that those fields are unphysical since they can be written in terms of other (physical) fields, i.e., they are redundant degrees of freedom. Notice the important fact that the scalar potential in (50) is positive definite, which is not the case in a generic non-supersymmetric theory.

4.1. Supersymmetry breaking

Now that we have learnt how to build interacting theories which are invariant under SUSY transformations, we have to face the fact that SUSY is not an exactly conserved symmetry of nature. Exact SUSY would require that particles and SUSY partners should be degenerate in mass and this has not been observed experimentally. Thus, we have to consider the breaking of supersymmetry²⁹. There are two interesting ways for breaking SUSY: spontaneously or explicitly. From what happens with other symmetries in physics, the spontaneous breaking of supersymmetry seems quite attractive. It consists in having a Lagrangian explicitly SUSY invariant but a vacuum state which is not

$$Q_\alpha |0\rangle \neq 0 \quad (52)$$

The "order parameter" for SUSY breaking is not the v.e.v. of a scalar but the energy of the vacuum state. To see this, recall that making use of the anticommutation relations in Eq. (36) for Q and $\bar{Q}$ one can write

$$P_0 \equiv H = \frac{1}{4} (\bar{Q}_1 Q_1 + Q_1 \bar{Q}_1 + \bar{Q}_2 Q_2 + Q_2 \bar{Q}_2) \quad (53)$$

and hence $\langle 0|H|0\rangle > 0$. Then Eq. (53) implies

$$\langle 0|H|0\rangle > 0 \rightarrow Q_\alpha |0\rangle \neq 0 \text{ (broken SUSY)} \quad (54a)$$

$$\langle 0|H|0\rangle = 0 \rightarrow Q_\alpha |0\rangle = 0 \text{ (conserved SUSY)} \quad (54b)$$

and the vacuum energy is the order parameter of supersymmetry. Notice that all this is consistent with the fact we discussed above that the scalar potential is positive definite

$$\langle 0|H|0\rangle = \langle 0|V|0\rangle = (\sum_i |F_i|^2 + \frac{1}{2} |D^a|^2)_{\text{at the minimum}} > 0 \quad (55)$$

We thus observe that in order to break SUSY spontaneously, one must have $\langle 0|F_i|0\rangle \neq 0$ or $\langle 0|D^a|0\rangle \neq 0$ (or both). In other words, one breaks SUSY by arranging that some of the auxiliary fields F or/and D get a v.e.v.

One can show the existence of a SUSY version of the Goldstone theorem which implies the existence of a massless particle with the transformation properties of the broken generators. Since the broken generator Q_α is a fermionic spin = 1/2 object, the associated Goldstone particle is a spin = 1/2 fermion, the "goldstino". In general, the latter is the SUSY partner of the non-vanishing auxiliary field F or D which has broken SUSY. Thus, the goldstino is a fermion of a chiral multiplet if we have broken SUSY through a $\langle F \rangle \neq 0$ or a "gaugino" if we broke SUSY through a $\langle D \rangle \neq 0$. In any case, if SUSY is broken, a massless goldstino must exist. We anticipate however that when one makes a "local" version of $N = 1$ SUSY ("N=1 supergravity"), there is a "SUSY Higgs" mechanism in which the goldstino is swallowed by the spin = 3/2 gravitino (SUSY partner of the graviton) which then acquires a mass. No massless particle is thus left.

As we stated above, there is a second interesting way of breaking SUSY³⁰. One may break SUSY "explicitly but softly". This consists in adding to the SUSY Lagrangian terms which are explicitly non-supersymmetric but are such that they do not induce quadratic divergences in loop calculations. As we mentioned before, this is required if we want to use SUSY to solve the hierarchy problem. This method for breaking SUSY looks quite ad hoc and unappealing at first sight. However, as we will see later on, it turns out to be most natural if we couple the SUSY standard model to $N = 1$ supergravity. I just briefly discuss this type of SUSY breaking here since we will use it later on. One can see that the SUSY-breaking terms one can add to a SUSY Lagrangian without inducing quadratic divergences in loop calculations are the following³⁰:

i) Scalar mass terms of the two different types

$$a) m^2 \phi^* \phi \quad (56a)$$

$$b) m^2 (\phi \phi + \text{h.c.}) \quad (56b)$$

where ϕ is the scalar in a chiral multiplet.

ii) Gaugino mass terms:

$$c) M(\lambda\lambda) + \text{h.c.} \quad (56c)$$

iii) Trilinear scalar couplings

$$d) A\phi^3 + \text{h.c.} \quad (56d)$$

where A is a dimensionless coupling constant.

Any of these terms may be added to the SUSY Lagrangian without generating quadratic divergences in loop calculations. However, e.g., other $\dim=3$ terms like explicit chiral multiplet fermion masses induce those divergences, i.e., they are not soft.

4iii. No renormalization theorems

The improved degree of divergence of SUSY theories is what makes them interesting in trying to solve the hierarchy problem. For the case of $N = 1$ supersymmetry these properties may be stated very easily. Let us have theory with chiral interactions determined by a superpotential W of a general form like Eq. (40). Then one has the following theorem: "the parameters in W (λ 's, m 's, f 's) do not suffer any renormalization from loop corrections". This is true both for finite and infinite loop corrections. This theorem implies that, e.g., if $m = 0$ at the tree level, m will be still vanishing in any order of perturbation theory. This is particularly useful for the Higgs doublet of the WS model: if it is massless to start with, it will still be massless even after radiative corrections. This is a great improvement on the situation concerning the gauge hierarchy problem. We only (!) have to explain now why the WS Higgs doublet was massless to start with. Let us remark that the fact that the parameters λ and m do not get loop corrections does not mean that they do not "run". They depend on the energy scale at which they are measured because there is still wave function renormalization of the fields since the kinetic energy term in the Lagrangian does receive (logarithmically) divergent loop corrections.

4iv. The supersymmetric standard model

With the tools described in the previous sections, we can now build a SUSY extension of the standard model. It will include, apart from the usual particles (gauge bosons, quarks, leptons and Higgs doublets), their supersymmetric partners. Eventually, we will have to consider two symmetry breakings: i) SUSY breaking and ii) $SU(2) \times U(1)$ symmetry breaking. It will turn out that in the simplest scenario, these two symmetry breakings are related. But before studying this double process of symmetry breaking, let us specify the (minimal) particle content of the SUSY- $SU(3) \times SU(2) \times U(1)$ theory³¹. As we anticipated, the gauge bosons and their fermionic partners ("gauginos") will be contained in vector multiplets whereas the quarks, leptons and Higgses and their supersymmetric partners ("s-quarks", "s-leptons", and "Higgsinos") will be contained in chiral multiplets. The particle content is shown in Table 3.

Table 3

VECTOR MULTIPLETS

$$J = 1 \quad J = 1/2$$

g (gluon)	+	$\tilde{g}$ (gluino)
$W^{\pm}, W^0$	+	$\tilde{W}^{\pm}, \tilde{W}^0$ (winos)
B	+	$\tilde{B}$ (bino)

CHIRAL MULTIPLETS

$$J = 1/2 \quad J = 0$$

q_L, q_R (quarks)	+	$\tilde{q}_L, \tilde{q}_R$ (squarks)
ℓ_L, ℓ_R (leptons)	+	$\tilde{\ell}_L, \tilde{\ell}_R$ (sleptons)
H_1, H_2 (Higgses)	+	H_1, H_2 (Higgsinos)

In the table, each particle and its SUSY partner transform in the same way under $SU(3) \times SU(2) \times U(1)$. Notice that one introduces two different WS doublets H_1 and H_2 . This is easily understood if we recall how one gives masses to quarks in the usual WS model:

$$\text{d-quarks: } h_d (\tilde{d}_R d_L) H + \text{h.c.} \quad (57a)$$

$$\text{u-quarks: } h_u (\tilde{u}_R u_L) H^* + \text{h.c.} \quad (57b)$$

in Dirac spinor notation. Once H acquires a non-vanishing v.e.v. $\langle H \rangle \neq 0$, the quarks get a mass. A SUSY version of Eqs. (57a,b) would require [see Eqs. (40)-(42)] superpotentials

$$W_d = h_d \tilde{d}_L^c \tilde{d}_L H \quad (58a)$$

$$W_u = h_u \tilde{u}_L^c \tilde{u}_L H^* \quad (58b)$$

However, W_u cannot be an admissible superpotential because it depends on H^* (superpotentials cannot depend on complex conjugate fields). To solve this problem, one introduces two doublets H_1 and H_2 with opposite hypercharge so that one uses H_2 in Eq. (58b) (instead of H^*) and H_1 in (58a). Notice that this reasoning also implies that one cannot use, e.g., the scalar partners of $(\tilde{\nu}_L, \tilde{e}_L)$ as Higgs particles since they would only give masses to d-quarks and leptons but not to u-quarks. [Adding just one Higgs doublet H_2 will not do because, although one could then give masses to u-quarks, the fermionic Higgsino $\tilde{H}_2$ would induce $U(1)$ anomalies.]

The supersymmetric $SU(3) \times SU(2) \times U(1)$ gauge interactions amongst the particles in the SUSY standard model are easy to obtain from the rules we gave in previous sections. The explicit Lagrangian may be found, e.g., in Ref. 32. Concerning the chiral interactions they are completely determined by the superpotential W . The superpotential of the (minimal content) SUSY standard model is

$$W = \sum_{\text{generations}} (h_u \tilde{q}_L^c \tilde{u}_L H_2 + h_d \tilde{q}_L^c \tilde{d}_L H_1 + h_L \tilde{\ell}_L^c \tilde{\nu}_L H_1) + \mu H_1 H_2 \quad (59)$$

where $\tilde{q}_L$ and $\tilde{\ell}_L$ denote the $SU(2)_L$ doublets $(\tilde{u}_L, \tilde{d}_L)$ and $(\tilde{\nu}_L, \tilde{e}_L)$ and $\tilde{u}_L^c, \tilde{d}_L^c, \tilde{\nu}_L^c$ are the scalar partners of the right-handed quarks and leptons. $SU(3)_C$ and $SU(2)_L$ index contractions are understood. The Yukawa coupling constants h_u, h_d and h_L are, in general, non-diagonal 3×3 matrices in flavour space leading to the usual masses and Cabibbo angles between quarks. The superpotential (59) is the most general compatible with baryon and lepton number conservation. Notice that it just includes the SUSY version of the usual Yukawas plus a possible mass term for Higgses.

4v. SUSY and grand unification

The SUSY standard model may be easily unified³³, e.g., inside $SU(5)$. In this case, the multiplet content would be as in Table 4 where each family of quarks and leptons is contained in a $\mathbf{5} + \mathbf{\bar{5}}_{10}$. One introduces Higgses Φ_{24} (and partners $\tilde{\Phi}_{24}$) in order to break $SU(5)$. The usual WS doublets are contained in $H_2 + H_2^*$.

Table 4

VECTOR MULTIPLT		CHIRAL MULTIPLTS	
$J = 1$	$J = 1/2$	$J = 1/2$	$J = 0$
$A_{24}^0 + \lambda_{24}$		$3(\tilde{F}_3 + \tilde{T}_{10}) + 3(\tilde{F}_5 + \tilde{T}_{10})$ $\tilde{\phi}_{24} + \phi_{24}$ $\tilde{H}_5 + \tilde{H}_5 + H_5 + H_5$	

Most of the features of usual GUTS remain true in the supersymmetric case. The existence of the SUSY partners of the usual particles cause the $SU(3) \times SU(2) \times U(1)$ gauge coupling constants to join at a larger mass scale³⁴. This is due to the fact that the coefficients of the β -functions b_1, b_2, b_3 in Eq. (24a) change in the supersymmetric case ($b_1 = 11, b_2 = 1, b_3 = -3$). One thus finds $M_X^{SS} \approx 3 \times 10^{16}$ GeV. The value of the coupling constant at the GUT scale also increases, $\alpha_{SS} \sim 1/24$ compared to $\alpha \sim 1/40$ in the non-SUSY case. Interestingly enough, the prediction for the Weinberg angle only changes slightly³⁴:

$$\sin^2 \theta_W = \frac{3}{8} - \frac{5}{4} \frac{\alpha_e(M_W)}{(4\pi)} 6 \log \left(\frac{M_X}{M_W} \right)^{SS} = 0.23 \quad (60)$$

compared to the ~ 0.20 result in the non-SUSY case (leading log). This result is, however, very sensitive to the particle content and changes sharply³⁵ if one adds some new particle to the standard model (like extra Higgs doublets³⁴). The prediction for the m_b/m_τ ratio is practically unchanged also³⁴:

$$m_b/m_\tau = \left(\frac{\alpha_3(m_b)}{\alpha_3(m_\tau)} \right)^{4/7} \left(\frac{\alpha_3(M_W)}{\alpha_{SS}} \right)^{8/9} \approx 3.0 \quad (61)$$

In all these computations we have assumed that SUSY is broken in an effective way around the weak scale so that the scheme of mass scales is as in Fig. 10. This has to be so in order to preserve the masslessness of the Higgs doublets down to the M_W scale where they are used to break the $SU(2) \times U(1)$ symmetry.

Fig. 10

The situation concerning baryon number violation in the SUSY case is more uncertain. There are three types of mechanisms potentially contributing to proton decay, as exemplified in Fig. 11.

Fig. 11

The first mechanism is the usual one already discussed for the non-SUSY case. This one is very much suppressed in general since the unification mass (and hence the masses of X and Y) is very large and the lifetime is proportional to the fourth power of M_X :

$$\tau_P^{SS} \sim \left(\frac{M_{SS}}{M_X} \right)^4 \tau_P^{NS} \sim 10^{38 \pm 2} \text{ years} \quad (62)$$

Mechanism b) is also present in the non-SUSY case and the rate depends on the unknown mass of the Higgs H_3 . The latter is the colour triplet of Higgs fields which completes the five-plet of Higgses along with the WS doublets H_2 (i.e. $H_5 = H_2 + H_3$). The decay is undetectable unless $m_{H_3} \sim 10^{11}$ GeV. Modes including strange particles would be dominant ($p \rightarrow K \pi^+ \nu$, etc.). Finally, the mechanism in Fig. 3 is genuinely supersymmetric³⁶. Unfortunately, unknown parameters (gaugino masses, squark-slepton masses, H_3 masses) are required to compute the graph. Reasonable choices of those parameters show that graphs like the one in Fig. 3 would be dominant in the SUSY case. However, due to the uncertainties mentioned, no precise prediction for the proton lifetime can be given. Again the decay modes involve predominantly strange particles.

Let us remark now a very important point concerning baryon stability. In the supersymmetric version of the SM, one can add to the Lagrangian terms compatible with $SU(3) \times SU(2) \times U(1)$ and Poincaré symmetries, but explicitly violating baryon³⁷ and/or lepton number. The following terms could in principle be added³⁷ to the superpotential in Eq. (59):

$$\begin{aligned} \text{a) } & (\tilde{u}_L^c \tilde{d}_L^c \tilde{d}_L^c) & \text{b) } & (\tilde{q}_L \chi_L \tilde{q}_L^c) \\ \text{c) } & m(\chi_L H_2) & \text{d) } & (\chi_L \chi_L \chi_R) \end{aligned} \quad (63)$$

The first violates baryon number and the second, third and fourth lepton number. If both terms a) and b) are allowed, graphs like the one in Fig. 12 will give rise to fast proton decay within a few minutes since the mass of the squark exchanged is necessarily of the order of $\sim M_W$. If only term a) exists, proton would not decay (it does not violate lepton number) but there can be neutron-antineutron oscillations, although one can compute that the rate of these oscillations would not be ruled out by experimental limits for reasonable choices of the parameters³⁸.

Fig. 12

The existence of possible renormalizable terms [like those in Eq. (63)] explicitly violating baryon and/or lepton number is a new feature of supersymmetric models. No such terms can be written in the minimal standard model. Of course, one can just forbid these terms in the superpotential and forget about it. This is perfectly reasonable since there is no principle within the context of SUSY-GUTs which tells us what terms should be in the superpotential and which terms should not. However, eventually, if we have a theory which is able to give us information about superpotential terms, it will have to explain why the superpotential of Eq. (59) does not contain the terms in Eq. (63). We will comment again on this problem when we talk about string models. For the moment, we will assume that none of the terms in Eq. (63) are allowed in the superpotential.

4vi. SUSY-breaking in the SUSY standard model

As is well known, no supersymmetric particles have been detected experimentally up to now (see below). pp and e^+e^- collider results imply $m_{\tilde{g}} \gtrsim 80$ GeV, $m_{\tilde{q}} \gtrsim 25$ GeV and similar limits apply for the rest of the charged SUSY particles. How can we obtain this in a natural manner? One can try to break SUSY spontaneously in the manner we discussed before. However, it turns out to be quite complicated (if not impossible) to break SUSY spontaneously without adding a large number of new ad hoc particles. There is, however, a very elegant mechanism which gives us what we want (i.e., masses for SUSY partners) in a very economical way. Furthermore, it is much more satisfactory theoretically since it is based on the extension of the scheme to include gravity ($N = 1$ supergravity). We first describe the effects of this type of SUSY breaking on the SM Lagrangian and then we will briefly discuss their origin.

Once this type of SUSY-breaking has taken place the standard supersymmetric Lagrangian gets new terms as follows:

- i) equal positive $(\text{mass})^2$ for all scalars (squarks, sleptons, Higgses) in the theory, $m^2|\phi_i|^2$;

- ii) new purely scalar interactions proportional to the superpotential, $\sim mW(\phi_i)$;

- iii) equal gaugino masses M at the GUT scale.

The mass parameter m is chosen to be $m \sim M_1$ and the gaugino mass M is an independent parameter also of order M_{GUT} . Applying this to the SUSY SM one obtains the following Lagrangian

$$\begin{aligned}
 L = & L(\text{SUSY}) + \delta L \text{ (SUSY-breaking)} \\
 \delta L = & m^2 \sum_{i=\tilde{q}, \tilde{\ell}, H_1, H_2} |\phi_i|^2 + AmW_3(q, \tilde{\ell}, H_1, H_2) + B\mu m H_1 H_2 + \text{h.c.} \\
 & + M_1 \lambda_1 + \text{h.c.}
 \end{aligned} \quad (64)$$

where W_3 is the trilinear part of the superpotential in Eq. (59), A and B are dimensionless constants $\sim$ unity and λ_1 are the gaugino fields. Thus, Eq. (64) gives us additional terms to the Lagrangian which can diagrammatically be represented as in Fig. 13. Interestingly enough, these terms correspond to the list of soft breaking terms which do not generate quadratic divergences given in Eqs. (56).

Fig. 13

What is the magnitude of m ? It has to be heavier than say ~ 20 GeV, in order to explain why we do not see $\tilde{q}$ or $\tilde{\ell}$'s. However, it cannot be much bigger than $M_1 \sim 10^2$ GeV since then the Higgs doublet would have a very large mass and it would not be possible to give a non-vanishing v.e.v. to H_1 and H_2 to break the $SU(2) \times U(1)$ symmetry. Notice that the Higgs $(\text{mass})^2$ is positive and at this level it is not easy to see how one can break the $SU(2) \times U(1)$ symmetry. We will discuss that below. Before doing that, let us say a few words about the origin of these SUSY-breaking couplings.

The general idea is that it makes no sense to have supersymmetric $SU(3) \times SU(2) \times U(1)$ interactions but to have non-supersymmetric gravitation. In fact, since the $N = 1$ SUSY algebra [Eq. (36)] involves the generator of translations p^μ , a local version of SUSY would contain general co-ordinate transformations, i.e., the invariance group of Einstein gravitation. Thus, a local version of SUSY is equivalent to a supersymmetric version of gravity, i.e., $N = 1$ supergravity⁴³. We thus have to consider the coupling of the SM to $N = 1$ supergravity. The spin $= 2$ graviton $g_{\mu\nu}$ has a supersymmetric spin $= 3/2$ SUSY partner ψ^μ , the gravitino. The gravitino is the "gauge particle" corresponding to local supersymmetry transformations. When supersymmetry is broken (i.e., any of the auxiliary fields $\langle F \rangle$ or $\langle D \rangle \neq 0$) the gravitino field "swallows" the massless "goldstino" and becomes massive much in the same way the $W^\pm$'s become massive by swallowing the usual Higgs field (Fig. 14).

Fig. 14

The second important ingredient in this mechanism of supersymmetry breaking is to assume it takes place in a "hidden sector" of the theory^{39-41,44}. This means to assume the existence of further singlet chiral multiplets (or maybe new gauge interactions) not coupling directly with the "observable sector" of (s)quarks, (s)leptons, Higgses, etc. The details of this "hidden world" are irrelevant and the only important point is the assumption that in this sector supersymmetry breaking takes place⁴⁵ at a scale $M_{SB} \sim \sqrt{M_{\text{Planck}}}$ in 10^{10} GeV. Then one can show that the gravitino gains a mass

$$m_{3/2} \sim \frac{|M_{SB}|^2}{M_{\text{Planck}}} \sim M_H. \quad (65)$$

Since the "observable world" is isolated from the hidden world, it does not feel that SUSY has been broken directly (i.e., there are no mass splittings between SUSY partners $\sim M_{SB}$). However, gravitational (and SUSY-gravitational) interactions transmit the SUSY-breaking from the "hidden" to the "observable" sector. Since gravitational interactions are suppressed by powers of $(1/M_{\text{Planck}})$, the effective SUSY breaking scale in the observable sector is only $m \sim M_{SB}^2/M_{\text{Planck}} \sim m_{3/2}^2$. Thus the parameter m in Eq. (64) is essentially given by the gravitino mass. That is why in Eq. (65) we chose $m_{3/2} \sim M_H$, in order not to give too large a mass to the scalars in the standard SUSY model. Let us remark that, since the "hidden matter" and the gravitino couple only gravitationally to the quark-lepton world, they are experimentally unobservable. They may, however, have cosmological importance and may also be candidates for the "dark matter" in the Universe. Summarizing this brief discussion on the mechanism of supersymmetry breaking, the important point is that the SUSY partners of the standard particles get the couplings in Eq. (64) with $m \sim m_{3/2}^2 \sim M_H$. For the purposes of low-energy physics the details of the $N = 1$ supergravity theory can be skipped.

4vii. $SU(2) \times U(1)$ symmetry breaking

Let us now discuss how the electroweak symmetry may be broken in the SUSY SM. We have to study the scalar potential for the Higgs fields H_1 and H_2 and check if there is a minimum for which $\langle H_1 \rangle, \langle H_2 \rangle \neq 0$. Let us start by writing down the scalar potential for H_1 and H_2

$$V(H_1, H_2) = \frac{1}{2} \left((H_1^\dagger \frac{a}{2} H_1 + H_2^\dagger \frac{a}{2} H_2)^2 g^2 + (|H_2|^2 - |H_1|^2)^2 g'^2 \right) + \mu_1^2 |H_1|^2 + \mu_2^2 |H_2|^2 - \mu_3^2 (H_1 H_2 + \text{h.c.}) \quad (66)$$

where g and g' are the WSG gauge couplings and the τ^a 's ($a=1,2,3$) are $SU(2)_L$ Pauli matrices. This is obtained from formulae (44), (48) and (64) after defining

$$\mu_1^2 \equiv m^2 + \mu^2; \quad \mu_2^2 \equiv m^2 + \mu^2; \quad \mu_3^2 \equiv B\mu \quad (67)$$

Equation (66) is the SUSY version of the celebrated "bottom of a bottle" Higgs potential which induces $SU(2) \times U(1)$ breaking in the usual SM. However, the potential in Eq. (66) is problematic because $\mu_1^2 = \mu_2^2 > 0$. A minimum with non-vanishing $\langle H_1 \rangle, \langle H_2 \rangle$ may only be obtained if there is a negative (mass)² eigenvalue in the Higgs mass matrix, i.e., if $\mu_1^2 - \mu_4^2 < 0$. However, if $\mu_1^2 = \mu_4^2 < \mu_3^2$, the scalar potential is unbounded below in the direction $\langle H_1 \rangle = \langle H_2 \rangle \rightarrow \infty$. There is something wrong though in the above analysis. μ_1^2 and μ_2^2 are equal only at the Planck scale. Below that point they will get renormalized differently if the scalars H_1 and H_2 couple with different strengths to quarks and leptons. This is true in general for all the soft breaking appearing in Eq. (64). Thus at the Planck scale we would have

$$m_q^2 = m_\chi^2 = \mu_1^2 - \mu^2 = \mu_2^2 - \mu^2 = m^2 \quad (68a)$$

$$M_{\tilde{g}} = M_{\tilde{W}} = M_{\tilde{B}} = M \quad (68b)$$

$$A_U = A_D = A_L = A \quad (68c)$$

but below that mass scale, all these parameters may evolve differently according to their corresponding renormalization group equations. We will not write down those equations which may be found in Refs. 50, 47 and 48, but we will give a general idea of how these parameters evolve with energy. Equations (68a-c) also assume that a grand unification occurs not very far from the Planck scale [notice, e.g., Eq. (68b) which implies that all the gauginos are unified in mass]. As an example, it turns out that gaugino masses M evolve with energy just like the $3 \times 2 \times 1$ coupling constants

$$M_1 = \frac{\alpha_1(Q^2)}{\alpha_1(M_X^2)} M. \quad (69)$$

This allows us to predict that $M_{\tilde{g}} \sim (\alpha_3/\alpha_e) M_{\tilde{W}}$ where $\tilde{g}$ and $\tilde{W}$ are the photino and gluino respectively. For the (masses)² of squarks and leptons, one gets an evolution with energy of the type shown in Fig. 15 (in that figure $m = M = 100$ GeV and $A = 1$ is taken). We thus find that $\tilde{q}$'s are usually heavier than $\tilde{\chi}$'s (one can see that they have similar masses in the limit $M \rightarrow 0$).

What happens with the Higgs scalar H_1 and H_2 ? We know that a negative (mass)² term would be welcome in order to break $SU(2) \times U(1)$. The miracle occurs and the Higgs masses may run in a natural way as shown in Fig. 16 and μ_2^2 gets negative whereas μ_1^2 remains positive. Notice that this is what we want since if μ_1^2 were also negative, the potential Eq. (66) would be unbounded below.

The mechanism in Fig. 16 works only if the top quark is heavy enough⁵¹. What has the t-quark to do with all this? It is simply that there is a negative contribution to μ_2^2 in its renormalization group equation from the graphs in Fig. 17.

The negative sign comes from the fact that supergravity breaking gives mass to squarks but not to quarks and then the diagrams 17a and 17b fail to cancel giving a net negative contribution⁵¹

$$\delta\mu_2^2 = -\frac{3}{8\pi^2} h_U^2 m^2 \text{Log}(M_X/Q) \quad (70)$$

In order for this negative contribution to overwhelm the original positive μ_2^2 , h_U has to be large enough. Detailed numerical computations show that in order to get the appropriate negative μ_2^2 to break $SU(2) \times U(1)$ one needs $h_U(M_X) \gtrsim 0.1$. It is obvious that the U-quark of a possible fourth quark-lepton generation would also do the job. We will specify below how heavy the top quark has to be.

Let us come back now to the $H_{1,2}$ scalar potential in Eq. (66). One can see that the potential is minimized for^{46,47}

$$v^2 \equiv v_1^2 + v_2^2 = \frac{2(\mu_1^2 - \mu_2^2 - (\mu_1^2 + \mu_2^2)\cos 2\theta)}{(g^2 + g'^2)\cos^2 \theta} \quad (71)$$

Fig. 15

Fig. 16

Fig. 17

where $v_{1,2} < H_0^{1,2}$ and $\sin 2\theta = 2\mu_3^2/(\mu_1^2 + \mu_2^2)$. Now we have to impose that the $SU(2) \times U(1)$ symmetry is broken at the appropriate scale, i.e., it must be $v^2 = 2M_W^2/g^2$, where M_W is the experimentally measured W -mass. This condition and Eq. (71) lead to the following constraint

$$\frac{\mu_1^2 - \omega \mu_2^2}{\omega^2 - 1} = \frac{1}{2} M_Z^2 \quad (72)$$

where $\omega = \tan \theta = v_2/v_1$ and M_Z is the Z^0 boson mass. The quantities μ_1^2, μ_2^2 and ω^2 in Eq. (72) should be evaluated at the M_W scale. An interesting limit is obtained in the case in which $\mu_3^2 \ll (\mu_1^2 + \mu_2^2)$. In that case $\cos 2\theta \rightarrow 1$ and $\omega \rightarrow \infty$ and the condition for $SU(2) \times U(1)$ breaking at the right scale reads

$$\mu_2^2(Q^2 = M_W^2) = -\frac{1}{2} M_Z^2 \quad (73)$$

which implies that μ_2^2 has to get a negative value, as indicated in Fig. 16. This implies that the top quark has to be heavy enough. However, Eq. (72) is more general than Eq. (73) and one can see that it is not strictly necessary to drive μ_2^2 negative in order to get $SU(2) \times U(1)$ breaking, although, as we will comment below, the former is the most natural possibility.

The $SU(2) \times U(1)$ breaking implies certain relationships between the parameters in the model in such a way that $\mu_1^2(M_W), \mu_2^2(M_W)$ and $\omega(M_W)$ can fulfill Eq. (72). These quantities depend on the free parameters of the model. Recalling Eqs. (59), (64) and (67), we see that those free parameters are

$$m, M, A, \mu_3^2, \mu \quad (74)$$

plus the Yukawa couplings, in particular h_t which is the only one that can be large enough to affect the renormalization group equations for μ_1^2, μ_2^2 and ω^2 . The condition of $SU(2) \times U(1)$ breaking implies the existence of constraints amongst the parameters in (74) and h_t (or, equivalently, the top quark mass $m_t \approx h_t(M_W)v_2$). One can study those constraints numerically. For example, in the limit $\mu_3^2, \mu \rightarrow 0$ (relevant in an interesting class of models) one finds numerical results as in Fig. 18.

Fig. 18

It shows the relationship between the parameters M (gaugino masses at M_X), m (scalar masses at M_X) and m_t for a certain value of the A parameter. One observes that one has always $m_t \gtrsim 60$ GeV and that the larger M is, the higher is the value of m_t required. Similar results are obtained for other values of A . At this point, it is interesting to recall that in the supersymmetric SM the top quark (or any possible new quark for that matter) has an upper bound $50.47, 48$ for its mass coming from the renormalization group equation for h_t . One can find the following numerical formula for the top quark mass in terms of its Yukawa coupling at the unification scale $h_t(0)$:

$$m_t = \left(\frac{3.6h_t(0)}{\sqrt{1+11h_t^2(0)}} \right) v^2 \quad (75)$$

As $h_t(0) \rightarrow \infty$ (as a formal limit), one has $m_t \sim v^2 \sim 200$ GeV. Thus one can say that in the minimal SUSY SM, one has the limits for the top quark mass

$$60 \text{ GeV} \lesssim m_t \lesssim 200 \text{ GeV}$$

for the class of models in which $\mu_3^2 \ll \mu_1^2, \mu_2^2$. For the case with $\mu_3^2 \sim \mu_1^2, \mu_2^2$, the lower bound disappears. However, in many SUSY GUTs or string models, one expects $\mu_3^2 \ll \mu_1^2, \mu_2^2$. This is because $\mu_3^2 = B\mu$ and a natural value for μ [see Eq. (59)] is zero. Notice that μ and $m = m_3/2$ are completely independent parameters which, on the other hand, have to be (for phenomenological reasons) of the same order of magnitude $m, \mu \sim M_{\text{GUT}}$. It is rather unnatural that two completely independent parameters turn out to have similar values without any obvious reason. It is most reasonable to assume that $\mu = 0$ to start with and a value for $\mu_3^2 \neq 0$ is generated radiatively once supersymmetry is broken. One can easily build models in which this is the case. In this large class of models, the lowest bound in Eq. (76) will then apply. Let us finally remark that one cannot have $\mu_3^2 = 0$ because then one can see, examining the scalar potential Eq. (66), that a massless pseudoscalar ("axion") would appear of the type introduced by Peccei and Quinn. That type of axions is ruled out experimentally.

4viii. The supersymmetric particle spectrum

a) Masses of squarks and sleptons

There are four different contributions (a,b,c,d) to the masses of squarks and sleptons. The "helicity" structure of these four terms is as follows ($\phi = \tilde{q}, \tilde{\ell}$)

$$\begin{aligned} \phi_L^* & \left[m_a^2 L^2 + m_b^2 L^2 + m_c^2 \right] & \phi_R^* & \left[m_a^2 R^2 + m_b^2 R^2 + m_c^2 \right] \\ \phi_L^* & \left[m_a^2 L^2 + m_b^2 L^2 + m_c^2 \right] & \phi_R^* & \left[m_a^2 R^2 + m_b^2 R^2 + m_c^2 \right] \end{aligned} \quad (77)$$

The leading contributions are a) and b). Contribution a) is the one coming from supergravity (Fig. 13a) after renormalization group equations for M_u to the M_u scale are taken into account. Contribution b) comes from the $1/2 E_a |D^2|^2$ term in the SUSY scalar potential (Fig. 9c) once the doublets $H_{1,2}$ acquire a v.e.v. Solving the relevant renormalization group equations for the masses of squarks and sleptons, one can express their masses in terms of the parameters of the model m, M and $\cos 2\theta$. We give here for completeness the sum $m_a^2 + m_b^2$ for all the squarks and sleptons of a generation^{4/52}

$$m_{0L}^2 = m^2 + 2M^2 \left(\frac{1}{3} \tilde{\alpha}_3 f_3 + \frac{1}{6} \tilde{\alpha}_2 f_2 + \frac{1}{6} \tilde{\alpha}_1 f_1 \right) + \cos 2\theta M_u^2 \left(-\frac{1}{3} + \frac{1}{3} \sin^2 \theta_u \right) \quad (78a)$$

$$m_{1L}^2 = m^2 + 2M^2 \left(\frac{4}{3} \tilde{\alpha}_3 f_3 + \frac{1}{6} \tilde{\alpha}_2 f_2 + \frac{1}{6} \tilde{\alpha}_1 f_1 \right) + \cos 2\theta M_u^2 \left(\frac{1}{3} - \frac{1}{3} \sin^2 \theta_u \right) \quad (78b)$$

$$m_{2L}^2 = m^2 + 2M^2 \left(\frac{4}{3} f_3 \tilde{\alpha}_3 + \frac{1}{6} \tilde{\alpha}_1 f_1 \right) - \cos 2\theta M_u^2 \left(\frac{1}{3} \sin^2 \theta_u \right) \quad (78c)$$

$$m_{0R}^2 = m^2 + 2M^2 \left(\frac{4}{3} \tilde{\alpha}_3 f_3 + \frac{1}{6} \tilde{\alpha}_1 f_1 \right) + \cos 2\theta M_u^2 \left(\frac{1}{3} \sin^2 \theta_u \right) \quad (78d)$$

$$m_{1R}^2 = m^2 + 2M^2 \left(\frac{4}{3} \tilde{\alpha}_2 f_2 + \frac{1}{6} \tilde{\alpha}_1 f_1 \right) + \cos 2\theta M_u^2 \left(\frac{1}{3} \sin^2 \theta_u \right) \quad (78e)$$

$$m_{2R}^2 = m^2 + 2M^2 \left(\frac{4}{3} \tilde{\alpha}_2 f_2 + \frac{1}{6} \tilde{\alpha}_1 f_1 \right) - \cos 2\theta \frac{M_u^2}{2} \quad (78f)$$

$$m_{\tilde{e}R}^2 = m^2 + 2M^2 \left(\tilde{\alpha}_1 f_1 \right) + \cos 2\theta M_u^2 \sin^2 \theta_u \quad (78g)$$

Here θ_w is the Weinberg angle, M_u is the Z^0 mass and m, M, θ are the free parameters in Eq. (74). One also has

$$\begin{aligned} \tilde{\alpha}_1 & \equiv \frac{\alpha_1(M_u)}{4\pi} ; & f_i & \equiv \frac{1}{\beta_1} \left(1 - \frac{1}{(1+\beta_1 t)^2} \right) \\ \beta_1 & \equiv \tilde{\alpha}_1 b_1 ; & t & \equiv \log(M_u^2/M_{\tilde{X}}^2) = 67 \end{aligned} \quad (79)$$

and $b_i = (-3, 1, 11)$ for $i = 3, 2, 1$ are the coefficients of the Callan-Symanzik $SU(3) \times SU(2) \times U(1)$ β -functions. From the above equations we see that, unless M is negligible, the squarks will be heavier than the sleptons since $\tilde{\alpha}_3 \gg \tilde{\alpha}_{2,1}$. We also see that (for non-vanishing $\cos 2\theta$) the scalar neutrino will be the lightest s -fermion⁵³. Contribution c) in Eq. (77) is just equal to the mass of the corresponding quark or lepton partner and thus is essentially negligible except for the quarks of the third generation. Finally, contribution d) mixes left and right squarks and sleptons and is proportional to m_{MA} , where m_i is the corresponding quark (lepton) mass. Again, it is negligible for the first two generations.

Notice that the $\tilde{q}$ and $\tilde{\ell}$ spectrum depends essentially only on three parameters m, M and $\cos 2\theta$. Since there are $7 \times 3 = 21$ different $\tilde{q}'$ s and $\tilde{\ell}'$ s, it is obvious that this scheme has strong predictive power. Furthermore, finding a scalar spectrum obeying Eqs. (78) would be a verification not only of supersymmetry but also of the unification concept since a common scalar mass m is assumed at the GUT scale. Notice also that m, M and $\cos 2\theta$ are not arbitrary since they have to be such that the $SU(2) \times U(1)$ symmetry is broken at the appropriate scale. This further restricts the pattern of $\tilde{q}$ and $\tilde{\ell}$ masses. We show below examples of consistent supersymmetric spectra (Tables 5 and 6).

TABLE 5

Some allowed spectra for $m_t \sim 50 \text{ GeV}$. Masses are in GeV

	a)	b)	c)
m	30	20	20
M	-25	1	-100
A	-1	-2	-4
μ_0	10	5	103
$\sin 2\theta(M_u)$	0.62	0.67	0.94
$\sin 2\theta(M_{\tilde{X}})$	0.99	0.99	0.98
m_t	52	50	55
$\tilde{q}$	73	20	269
$\tilde{\ell}_1$	43	39	230
$\tilde{\ell}_2$	116	64	303
$\tilde{\ell}_3$	38	21	79
$\tilde{g}$	75	3	301
$\tilde{\gamma}$	12	0.5	46
N^0	88	92	172
	92	85	143
	14	6	106
	75	74	105
$\tilde{W}_2$	84	81	166
$\tilde{W}_1$	89	82	216
H_a	99	92	219
H_b	4	0.4	17
H_c	45	27	201

TABLE 6

Some allowed spectra for $m_t \geq 60$ GeV. Masses are in GeV

	a)	b)	c)	d)	e)	f)	g)	h)	i)
m	20	20	40	40	20	30	60	30	120
M	-60	100	100	-120	-120	-60	-60	-25	-25
A	-3	2	2	-1	0	-1	-2	-1	3
μ_0	2	40	6	5	12	2	2	3	17
$\sin 2\theta(M_X)$	0.41	-0.40	-0.15	0.23	0.45	0.10	0.12	0.17	-0.28
$\sin 2\theta(M_W)$	0.88	-0.83	-0.38	0.30	0.48	0.79	0.52	0.96	-0.59
m_t	60	62	69	69	68	77	83	90	107
q	161	268	270	323	321	162	169	71	134
$\tilde{g}_1$	102	219	204	264	269	102	64	56	58
$\tilde{g}_2$	212	312	324	374	369	222	241	146	193
$\tilde{g}_3$	91	83	94	106	100	64	86	42	124
$\tilde{g}_4$	42	54	69	74	65	51	75	38	123
$\tilde{g}_5$	22	60	58	75	68	21	47	16	117
$\tilde{g}_6$	180	301	301	361	361	180	180	75	107
$\tilde{g}_7$	113	125	133	145	144	112	113	97	105
$\tilde{g}_8$	71	98	63	58	64	71	71	82	76
$\tilde{g}_9$	3	43	4	2	10	2	2	3	13
$\tilde{g}_{10}$	50	74	22	16	32	44	28	65	28
$\tilde{g}_{11}$	109	127	135	146	145	112	117	90	110
$\tilde{g}_{12}$	82	120	98	108	106	81	92	80	139
$\tilde{g}_{13}$	91	122	93	95	99	90	93	91	131
$\tilde{g}_{14}$	12	37	52	68	57	14	40	5	63
$\tilde{g}_{15}$	26	92	59	76	73	23	49	20	116

b) Masses of Higgs particles

The supersymmetric SM contains two doublets of Higgses $H_1 = (H_1^+, H_1^0)$, $H_2 = (H_2^+, H_2^0)$ with altogether eight degrees of freedom. Three of them are swallowed by the $W^\pm$ and Z^0 to get a mass. There remain as physical scalar fields one charged Higgs $H^\pm$ and three neutral Higgses that we will call H_a, b, c . Using the scalar potential Eq. (66), one arrives at the following expressions for their masses

$$m_{H^\pm}^2 = M_W^2 + \mu_1^2 + \mu_2^2 \quad (80a)$$

$$m_{H_c}^2 = \mu_1^2 + \mu_2^2 \quad (80b)$$

$$m_{H_{a,b}}^2 = \frac{1}{2} [m_{H_c}^2 + (m_{H_a}^2 + m_{H_b}^2)^2 - 4m_{H_a}^2 m_{H_b}^2 \cos 2\theta]^{1/2} \quad (80c)$$

Examining these formulae, one observes the following facts. First, the charged Higgs is very massive, $m_{H^\pm} \gg M_W$ [see Eq. (80a)]. Concerning the neutral Higgses, two of them are relatively massive (H_a and H_c) whereas the other (H_b) is bounded above by M_Z . In fact, one can derive the following inequalities

$$(m_{H_a}^2 + m_{H_c}^2)^{1/2} > m_{H_a} > M_Z \quad (81a)$$

$$\cos 2\theta M_Z > m_{H_b} > 0 \quad (81b)$$

$$m_{H_a} > m_{H_c} > m_{H_b} \quad (81c)$$

Typically $\cos 2\theta \approx 0.1-1.0$ so that the scalar H_b may be rather light. This is to be compared with the Higgs field in the usual SM for which there is no bound. Here the bound comes about because the self-couplings of the Higgses are fixed (by gauge invariance and supersymmetry) to be proportional to the gauge coupling constant. Notice how Eqs. (81a-c) are fulfilled by the examples of consistent spectra shown in Tables 5 and 6.

c) Masses of SUSY fermions

Here we call SUSY fermions the gauginos and Higgsinos. For the case of the gluino, the result is very simple

$$M_{\tilde{g}} = (\alpha_3/\alpha_{GUT})M \quad (82)$$

but the mass spectrum of the $\tilde{W}$, $\tilde{B}$ and Higgsinos is more complicated because once $SU(2)_C \times U(1)$ is broken they mix amongst themselves. This is due to the existence of the SUSY gauge couplings of Fig. 8b which gives rise to mass terms which mix the charged winos with the charged Higgsinos and $\tilde{W}^0$ and $\tilde{B}$ with the neutral Higgsinos. Apart from these, there are the usual Majorana mass for the gauginos (Fig. 13d) and the direct Higgsino mass μ coming from the superpotential Eq. (59). In this way, we have a mass matrix for the charged winos-Higgsinos⁵⁶

$$\begin{pmatrix} \tilde{W}^+ & \tilde{H}^+ \\ \tilde{W}^- & \begin{pmatrix} M_2 & g_2 v_1 \\ g_2 v_2 & \mu \end{pmatrix} \end{pmatrix} \quad (83)$$

where M_2 is the direct wino Majorana mass and $v_{1,2} = \langle H_{1,2} \rangle$. If both M_2 and μ are large (larger than M_W) the eigenvalues of this matrix will both be large and will correspond to the wino ($\tilde{W}^+, \tilde{W}^-$) and Higgsino ($\tilde{H}^+, \tilde{H}^-$) states. In the limit $M_2 \rightarrow \infty$ and $\mu \rightarrow 0$, there will be two eigenstates with masses $(M_2, g_2^2 v_1 v_2 / M_2)$, the latter being lighter than the $W^\pm$ boson. In the limit $M_2 \rightarrow 0$, $\mu \rightarrow 0$ there are two eigenstates with masses $(g_2^2 v_1, g_2^2 v_2)$. The latter would correspond to the Dirac fermions $\phi_1 = (\tilde{W}^+, \tilde{H}^-)$, $\phi_2 = (\tilde{W}^-, \tilde{H}^+)$ respectively. Notice that in this case one has $M_W^2 = 1/2(M_2^2 + \mu^2)$ and ϕ_1 has to be lighter than the W -boson. Thus, in the interesting class of models in which μ is negligible, there is a charged fermion $\lambda^\pm$ which will be lighter than the W -boson. This fermion has good chances to be the lightest charged supersymmetric particle in the SUSY standard model. Values of this mass for several consistent choices of the parameters are shown in Tables 5 and 6. Notice how in many cases the lightest chargino $\tilde{\chi}_1^\pm$ will be available in forthcoming accelerators.

The mass matrix for the neutral "fermioninos" is more complicated⁵⁶

$$\begin{pmatrix} \tilde{W}^0 & \tilde{B}^0 & \tilde{H}_1^0 & \tilde{H}_2^0 \\ \begin{pmatrix} M_2 & 0 \\ 0 & M_1 \end{pmatrix} & \begin{pmatrix} -g_2 v_1 / \sqrt{2} & g_2 v_1 / \sqrt{2} \\ g_1 v_1 / \sqrt{2} & 0 \end{pmatrix} & \begin{pmatrix} -g_1 v_2 / \sqrt{2} & \mu \end{pmatrix} \\ \begin{pmatrix} -g_2 v_1 / \sqrt{2} & g_1 v_1 / \sqrt{2} \\ g_2 v_2 / \sqrt{2} & -g_1 v_2 / \sqrt{2} \end{pmatrix} & \begin{pmatrix} \mu & 0 \end{pmatrix} \end{pmatrix} \quad (84)$$

where M_2 and M_1 are related through Eq. (69) implying $M_1 = (3\alpha_1/5\alpha_2)M_2$. There is a linear combination of $\tilde{W}^0$ and $\tilde{B}^0$ (the photino $\tilde{\gamma}$) which does not mix directly with the Higgsinos and is given by

$$\tilde{\gamma} \approx g_1 \tilde{W}^0 + g_2 \tilde{B}^0 \quad (85)$$

with mass

$$M_{\tilde{\gamma}} = \frac{g_1^2}{g_1^2 + g_2^2} M_2. \quad (86)$$

Notice that $M_{\tilde{\gamma}}/M_{\tilde{Z}} = g_3/\alpha_e \approx 6$. The photino is the lightest SUSY particle in many models [when I say "model", I refer to possible ranges of the free parameters in Eq. (74)], particularly when M is not too large. On the other hand, a neutral Higgsino is also very often the lightest SUSY particle when μ is very small. In general, the mass matrix (84) has to be diagonalized numerically case by case. Again Tables 5 and 6 show examples of consistent spectra. Notice how often a neutralino $\tilde{N}^0$ is lighter than the photino. In some cases also, the sneutrino may be the lightest supersymmetric particle³³. We should remark that some of the details given above rest on the assumption of a "minimal content" of the SUSY SM. For example, some of the features of particle spectra (particularly regarding neutralinos and neutral Higgses as well as the lower bound on the top quark) change if an extra singlet multiplet S^0 coupling to Higgses with a Yukawa as $(S^0 H_1 H_2)$ exists. This possibility seems to appear in some string-inspired models in which direct Higgsino mass terms $\mu H_1 H_2$ seem difficult to get. However, most of the general features survive also in this case and a v.e.v. for S^0 acts as an "effective" μ parameter.

4ix. Supersymmetry and experiment

One of the most attractive points of the supersymmetric version of the SM is that this theory can be directly checked experimentally (it is not clear if this is the case for grand unification of superstrings). If no supersymmetric particles are found below ~ 1 TeV, low-energy supersymmetry would be ruled out. In fact, this upper limit is clearly pessimistic. Using naturality arguments, one may estimate "reasonable" upper bounds on the masses of squarks, sleptons, gluinos and "fermioninos" as in Fig. 19 (Barbieri et al. in Ref. 58). Again, one does not expect those limits to be saturated and one would expect slepton masses ~ 100 GeV and heavier squark and gluinos ~ 200 – 300 GeV although, of course, these are matters open to bet. Unfortunately, the range of masses explored by present accelerators is rather small compared to the expected range of SUSY particles masses.

The effect of SUSY particles may also appear indirectly through radiative corrections to SM processes or through SUSY contributions to rare decays. In this connection, it is important to remark that in the SUSY SM discussed above, the masses of squarks and sleptons are practically (up to Yukawa couplings) independent of the generation. This degeneracy between $\tilde{q}$'s and $\tilde{\ell}$'s of different families is very much welcome, since otherwise there would be sizeable flavour-changing neutral currents (FCNC). There is a SUSY version⁵⁹ of the Glashow-Iliopoulos-Maiani (GIM) mechanism at work which suppresses FCNC to the appropriate limits if the $\tilde{q}$'s and $\tilde{\ell}$'s are almost degenerate. Let us remark however that if some rare decay rate or process incompatible with the SM were to be detected experimentally, most probably it could be "explained" by many other possible models apart from supersymmetry. A reliable check of supersymmetry will require the production of SUSY particles at accelerators.

Fig. 19

In order to look for sparticles at accelerators, one needs to know something about the possible signatures. An important point to note is that, with a superpotential like the one in Eq. (59), there is a discrete symmetry ("R-parity") in the theory under which sparticles are "odd" and usual particles are "even"

$$\begin{aligned} & \text{R-parity} \\ & q, \lambda, H, g, \dots \rightarrow q, \lambda, H, g, \dots \\ & \tilde{q}, \tilde{\chi}, \tilde{H}, \tilde{g}, \dots \rightarrow -\tilde{q}, -\tilde{\chi}, -\tilde{H}, -\tilde{g}, \dots \end{aligned} \quad (87)$$

This means that sparticles can only be produced in pairs. Using these facts and the SUSY interactions described in Figs 6, 8 and 9, one can find the expected decay modes of each sparticle. Typical decays are depicted in Fig. 20 under the assumption that the gauginos are lighter than the squarks and sleptons (this is not always the case but the intelligent reader should be able to find the decay modes in the opposite situation). Of course, the decays in Fig. 20 are not the only possibilities. Notice also that, in general, the fermioninos ($\tilde{W}^\pm, \tilde{H}^\pm, \tilde{Z}^0, \tilde{\gamma}, \tilde{N}^0$) are not mass eigen-

states and one has to consider the eigenstates of matrices (83) and (84) as the physical fields. Notice that in all the decays in Fig. 20 there is a photino amongst the final state particles. For some choices of the supersymmetry breaking parameters (particularly when M is not too large), the photino is the lightest supersymmetric particle. Then R-parity conservation would imply that it is absolutely stable and will appear at the end of any decay of a supersymmetric particle. On the other hand, the photino interacts very weakly with matter with cross-sections typical of weak interactions (this is because they require the exchange of squarks and sleptons). Then the photino will not be detected by a conventional detector and will give rise to large missing momentum when produced in the final state. This is a typical signature which also appears in the more general case in which the lightest supersymmetric particle (LSP) is a Higgsino or a scalar neutrino.

Fig. 20

In the last few years, there has been an important experimental effort looking for SUSY particles and a number of useful limits have been put. The e^+e^- machines⁶⁰ give lower bounds for the masses of charged sleptons $m_{\tilde{l}} \gtrsim 25$ GeV. For the selection the limit $m_{\tilde{e}} \gtrsim 32$ GeV may be put from the non-observation of the decay $W \rightarrow \tilde{e}\nu$ (UAL)⁶¹. A nice indirect limit on the mass of the $\tilde{e}$ comes from the search for events produced by the (hypothetical) process depicted in Fig. 21. The signature is a photon plus missing energy and there is a well-understood conventional background from the $e^+e^- \rightarrow \nu\bar{\nu}$ process. The cross-section from Fig. 21 is proportional to $(m_{\tilde{e}})^{-4}$ and thus one obtains a limit on $m_{\tilde{e}}$ depending on $m_{\tilde{\gamma}}$ (provided $m_{\tilde{\gamma}} \lesssim \sqrt{s}/2$). The results from ASP⁶² (there are also similar limits from MAC⁶³) are shown in Fig. 22.

Fig. 21

Fig. 22

One finds, e.g., $m_{\tilde{g}} \gtrsim 50$ GeV for $m_{\tilde{q}} \lesssim 5$ GeV. The best limits on the masses of squarks and gluino come from the $p\bar{p}$ colliders through 64 diagrams like those in Fig. 23. The decay of squarks and/or gluinos involve photinos giving rise to missing E_T events. There is a soft spectrum conventional background from the $W \rightarrow \nu\bar{\nu}$ and $q\bar{q} \rightarrow Z^0(\nu\bar{\nu})g$ processes amongst others. The present UAL data⁶⁵ exclude an area in the $m_{\tilde{q}}-m_{\tilde{g}}$ plane as shown in Fig. 24. The tevatron will soon improve these limits, at least up to the ~ 80 GeV range. Concerning the "charginos" $\tilde{W}^\pm$, $\tilde{H}^\pm$, e^+e^- machines put a limit $m_{\tilde{W}^\pm} \gtrsim 22$ GeV and a slightly better (though more model-dependent) limit may be obtained from the non-observation in UA2⁶⁶ of decays like $\tilde{W}^\pm \rightarrow C^+\tilde{\gamma}$ ($C^+ = \tilde{W}^\pm - \tilde{H}^\pm$ physical linear combination). The limits on neutral particles like the three neutral physical Higgses, neutralinos (physical linear combinations of Higgsinos, photino and zeno) and sneutrinos are much weaker and the best (although very model-dependent) come from cosmological considerations. Due to the presence of too much dark matter, one gets from the non-closure of the Universe, a lower bound $m_{\tilde{W}^\pm} \gtrsim 5$ GeV for a stable photino and $m_{\tilde{W}^\pm} \gtrsim 5$ GeV for the lightest supersymmetric particle (LSP) being a Higgsino⁶⁷. I should remark at this point that many experimental limits on the masses of SUSY particles (like squarks and gluinos) assume that the LSP is the photino. A complete reanalysis of all experimental data for the hypothesis of a Higgsino being the LSP is still to be done but a general feature is that present experimental limits are in general worse since then photinos decay into Higgsinos + hadrons (leptons) and the missing energy signature is less efficient (unless the photino is sufficiently stable to escape the detector).

Fig. 23

Fig. 24

A word of caution should be given concerning the conservation of "R-parity". If terms like those of Eq. (63) are present in the superpotential of the SUSY-standard model, R-parity is explicitly broken and SUSY-particles need no longer be produced in pairs^{68,69}. Thus, e.g., if term (63a) exists, then the photino would decay mostly into strange particles, $\tilde{\gamma} \rightarrow u_d^c s^c$. If a term like (63d) exists, one could produce sneutrinos in e^+e^- collisions, $e^+e^- \rightarrow \tilde{\nu}$, etc. In the presence of these R-parity violating terms in the Lagrangian much of the expected SUSY-signatures would change. Notice, however, that, as we remarked above, not all the terms in Eq. (36) can be present simultaneously since they violate baryon or lepton number and when both things are simultaneously violated the proton would decay too fast⁶⁸. Thus, although R-parity violation may exist, to make it compatible with the observed conservation of baryon and lepton numbers would require that some of the terms in Eq. (63) were present but others do not. This makes this possibility rather unattractive but still certainly possible.

Let us end this section on supersymmetry with a few comments. The main goal of the introduction of a SUSY version of the Standard Model is to avoid the naturalness problem. Scalars may be relatively light (compared with the Planck mass) in supersymmetric theories. Furthermore, since the self-coupling of Higgs particles (at least in the minimal model) is governed by the gauge coupling constant, the Higgs particles of the SUSY-Standard Model are much more constrained than in the non-supersymmetric case. Moreover the breaking of the electroweak symmetry $SU(2) \times U(1)$ occurs in a much more elegant way as a consequence of supersymmetry breaking. Radiative corrections render a negative (mass)² contribution to one of the Higgses which then triggers $SU(2) \times U(1)$ breaking. This is to be compared with the standard model procedure in which one puts by hand a negative (mass)² term in the scalar potential.

The natural extension of $N = 1$ supersymmetry to a local symmetry ($N = 1$ supergravity) provides also for the possibility of an elegant supersymmetry breaking mechanism. This is assumed to take place in a "hidden sector" of the theory which couples to the "observable" world only through gravitational interactions. The details of the structure of the

hidden world are not relevant as long as supersymmetry breaking takes place in that sector at a scale $\sim \sqrt{M_{\text{W}} M_{\text{P}}} \sim 10^{10}$ GeV. This type of SUSY-breaking allows for the construction of a SUSY version of the Standard Model obeying all experimental constraints. We remark that this is a non-trivial result since other approaches like technicolour or composite models do in general have problems with experimental constraints (e.g., flavour-changing neutral currents).

In spite of all these attractive features, supersymmetry does not provide us with any new clues concerning questions like fermion masses and mixings or why there is family replication. Furthermore, one of the ingredients is $N = 1$ supergravity, which is known to be non-renormalizable as a field theory. Thus again, $N = 1$ supergravity coupled to the Standard Model (or to a GUT for that matter) cannot be the whole story.

5. STRINGS

The introduction of strings³ is certainly a more daring step than the ones we have been discussing before. It implies a change in many concepts usually taken for granted. Thus the fundamental objects are no longer particles but extended one-dimensional objects ("strings") and the observed particles correspond to some "vibration modes" of the string. Space and time (the number of dimensions) are now dynamical variables instead of the "arena" in which physics takes place. Many points are still poorly understood in string theory (e.g., there is no consistent field theory of closed strings) although in the last few years there has been much progress. Of course, in these lectures I will not pretend to give an introduction to string theory but will just try to give a flavour of the topic for the non-experts. I will try to emphasize how the topics we have been discussing in previous sections could fit within the string context. The interested reader is referred to the abundant reviews on the subject³.

We are all familiar with the dynamics of a relativistic free particle. Its co-ordinates X^μ are a function of the proper time τ (Fig. 25) and the action is given by

$$S = m \int_{\tau_i}^{\tau_f} d\tau \sqrt{-\dot{X}^2(\tau)} = m \int_{\tau_i}^{\tau_f} ds \quad (88)$$

where $\tau_{i,f}$ are the initial and final times respectively and ds is the infinitesimal interval. We see from Eq. (88) that the action of the free particle is given by the length of the world-line which is described when it propagates. An open string is a one-dimensional extended object parametrized by a real parameter σ (Fig. 25b). When it propagates in space it sweeps a "world-sheet" parametrized by σ and a time-like variable τ . Then each point in the string is described by a co-ordinate $X^\mu(\sigma, \tau)$. In analogy with Eq. (88) one can show that the action of a free relativistic string is given by the area swept by the world sheet. The simplest expression for the string action is obtained as⁷⁰

b) open string

a) particle

$$S = \int d\sigma d\tau \sqrt{-g} g^{ab} \partial_a X^\mu \partial_b X_\mu \quad (89)$$

where g^{ab} is a two-dimensional metric on the world-sheet ($a, b = \sigma, \tau$) and $\partial_a = \partial/\partial\sigma$, $\partial_b = \partial/\partial\tau$ ($g = \det|g^{ab}|$). This metric behaves as an auxiliary field and may be eliminated from Eq. (89) by making use of the equations of motion. In fact the action is invariant under reparametrizations of the world-sheet $\sigma, \tau \rightarrow \sigma', \tau'$ ($S \sim$ area, independently of how we choose to parametrize the surface) and one can use this invariance to choose a flat metric in Eq. (89):

$$S = \int d\sigma d\tau \eta^{ab} \partial_a X^\mu \partial_b X_\mu \quad (90)$$

with $\eta = \text{diag}(1, -1)$. An important point to realize is that Eq. (90) looks like the action of a massless bosonic field in a space-time with two dimensions (σ, τ). Thus the movement of a string along with time may be represented by a two-dimensional massless field theory. These theories are rather remarkable due to the fact that they possess a huge amount of symmetry ("conformal invariance"). Conformal invariance¹ is the symmetry of conformal mappings of the two-dimensional plane. It is convenient to work with a Euclidean version ($\tau \rightarrow -i\tau$) of the world-sheet and to use the complex variables $z, \bar{z}$: ($z \equiv \tau + i\sigma$) instead of σ and τ . Then conformal invariance implies invariance of the action under transformations

$$z \rightarrow f(z) \quad \bar{z} \rightarrow g(\bar{z}) \quad (91)$$

where $f(g)$ are holomorphic (antiholomorphic) arbitrary functions. Co-ordinate transformations of this type generate the "conformal group". For more than two dimensions the conformal group is finite-dimensional. Two dimensions (like in our case σ, τ) are special and in this case the group is infinite-dimensional. This enormous symmetry makes it possible to get rid of unphysical negative norm states.

The energy momentum tensor plays a special rôle since it is the generator of local conformal transformations. In fact one can see that the four possible components T ($a, b = \sigma, \tau$) of this tensor only two $T_{zz} \equiv T$ (and the complex conjugate $\bar{T}_{\bar{z}\bar{z}} \equiv \bar{T}$) are non-vanishing. For the action (90) the corresponding energy-momentum component is:

$$T = -\frac{1}{2} (\partial_z X)^2; \quad (92)$$

where the dots indicate normal ordering product of the fields. Thus T is a "composite" operator. The behaviour of fields $\phi(z)$ under a conformal transformation are given by the operator product expansions⁷¹ (OPE) (we omit normal ordering dots from now on)

$$T(w) \phi(z)_{w \rightarrow z} \sim \frac{d_\phi}{(w-z)^2} \phi(z) + \frac{1}{(w-z)} \partial_z \phi(z) + (\text{non-singular}) \quad (93)$$

d_ϕ is the "conformal dimension" of the field $\phi(z)$. One finds that the conformal dimension of the field $(\partial_z X)$ is $d = 1$. Recalling the form of T in Eq. (92) one would expect T to have conformal dimension $d = 2$. However, when computing the relevant operator product expansion [using Wick's theorem and the form of the propagator of the field, $\langle X^\mu(z) X^\nu(w) \rangle = -\eta^{\mu\nu} \log(z-w)$] one finds³

$$T(z)T(w) \sim \frac{c/2}{w-z} + \frac{2}{(z-w)^2} T(w) + \frac{1}{(z-w)} \partial_w T + \dots \quad (94)$$

where in the present case $c = 1$. The last two terms are the ones one would expect for a field for conformal dimension $d = 2$. One can see that the first most singular term, which is not expected at the classical

level, appears as a quantum mechanical effect. The coefficient c in Eq. (94) is called "central charge". The behaviour in Eq. (94) occurs in general for any conformal field theory and not only for the simple case of Eqs. (90) and (92). The appearance of such a term leads to a violation of conformal invariance at the quantum level (an "anomaly"). In fact one can consider the Fourier components of $T(z)$ on the circle centered in the origin³

$$L_n = \int \frac{dz}{2\pi i} z^{n+1} T(z) \quad (95)$$

They generate the infinitesimal conformal transformations corresponding to $f(z) = z^{n+1}$. The infinite operators L_n ($n = 0, \pm\infty$) constitute the Virasoro operators associated to the conformal invariance. Using Eqs. (94) and (95) one can obtain the commutators

$$[L_n, L_m] = (n-m) L_{n+m} + \frac{c}{12} n(n+1) (n-1) \delta_{m+n} \quad (96)$$

leading to the Virasoro algebra. The last term corresponds to the first in Eq. (96) and one can see that it has the typical aspect of an "anomalous" quantum term violating a classical (conformal) invariance. To have a consistent conformal invariant theory (which we need in order not to have ghosts), the overall central charge c (the sum over all the central charges of each independent conformal field) has to vanish.

In the case of an action like that in Eq. (90) the central charge would be $c = D$, $D =$ number of dimensions, since $c = 1$ for a bosonic field $X(z)$, and thus the system would not be conformally invariant at the quantum level. In fact this is not completely correct. In analogy with what happens when quantizing a usual non-Abelian Yang-Mills theory in which in order to avoid "double counting" due to the "gauge volume" one introduces the Faddeev-Popov ghosts, one also has to add "ghosts" to the system of a bosonic string. These "ghosts" are associated to the "gauge volume" of reparametrizations of the world sheet (ghosts "b" and "c"). They are scalars which anticommute, as usually happens with ghosts. They have conformal dimensions $d(c) = -1$, $d(b) = 2$. The associated piece of the energy-momentum tensor is $T(z) = -2b\partial_z c - \partial_b c$. Again, using Wick's theorem and the propagator $\langle b(z)c(w) \rangle = \frac{1}{2}(z-w)^{-1}$ one gets the operator product expansion

$$T_{\text{ghosts}}(z) T_{\text{ghosts}}(w) \sim \frac{(-13)}{(z-w)^4} + \dots \quad (97)$$

so that the ghost system has central charge $c = -26$. Thus the central charge of a bosonic string in D dimensions is

$$c = (D - 26) \quad (98)$$

and the theory is consistent only in 26 space-time dimensions. This is the well-known result that the purely bosonic string⁷³ is only consistent in 26 dimensions. Let us remark, however, that what $c = 0$ implies is the necessity of having 26 $X^\mu(\sigma, \tau)$ ($\mu = 0, \dots, 25$) fields but not necessarily all of them need to have an interpretation as physical space-time dimensions. As we will see later on, by appropriately choosing boundary conditions for the fields $X^\mu(\sigma, \tau)$ one can make that only a subset of them (e.g., $4 = 3+1$) admits an interpretation as observable space-time dimensions.

What is the mass spectrum of the bosonic string? The equations of motion for an action as that in Eq. (90) are the ones of a two-dimensional Klein-Gordon field

Fig. 26

$$\left(\frac{\partial^2}{\partial \sigma^2} - \frac{\partial^2}{\partial \tau^2}\right) X^\mu(\sigma, \tau) = 0 \quad (99)$$

The freedom to make conformal transformations allows one to work in the "light-cone gauge" in which only the D-2 transverse $X^i(\sigma, \tau)$ ($i = 1, \dots, D-2$) fields appear as independent physical fields (this is reminiscent of what happens with the physical transverse degrees of freedom of a usual gauge boson like, e.g., the photon). Let us consider the case of a closed string as in Fig. 26. The world-sheet is now a cylinder. The general solution of Eq. (99), subject to the boundary condition $X^i(\sigma, \tau) = X^i(\sigma + \pi, \tau)$ (so that the string is really closed), is

$$X^i(\sigma, \tau) = x_0^i + p^i \tau + \frac{i}{2} \sum_{n \neq 0} \left\{ \frac{\alpha_n^i}{n} e^{-2in(\tau - \sigma)} + \frac{\tilde{\alpha}_n^i}{n} e^{-i2n(\tau + \sigma)} \right\} \quad (100)$$

[here σ and τ have again a Minkowskian metric as in Eq. (90)]. Here x_0^i and p^i are the centre-of-mass position and momentum and the two terms in brackets correspond to string vibrations going in one direction ("right-moving") and in the opposite ("left-moving"). In complex (Euclidean) notation they would correspond to holomorphic ($X^i(z)$) and antiholomorphic ($\tilde{X}^i(\bar{z})$) fields. Upon quantization one has

$$[\alpha_n, \alpha_m] = n\delta_{n+m}, 0; \quad [\tilde{\alpha}_n, \tilde{\alpha}_m] = n\delta_{n+m}, 0 \quad (101)$$

One can explicitly compute the mass operator $m^2 = p^\mu p_\mu$ and one obtains

$$\frac{m^2}{4} = \frac{m_0^2}{4} + \frac{m^2}{4} = (N-1) + (N-1) \quad (102)$$

where N , $\tilde{N}$ are the "number operators"

$$N = \sum_{n=1}^{\infty} \alpha_{-n}^i \alpha_n^i; \quad \tilde{N} = \sum_{n=1}^{\infty} \tilde{\alpha}_{-n}^i \tilde{\alpha}_n^i, \quad (i = 1-24) \quad (103)$$

for "left" and "right" vibrations.

A general state in the Hilbert space of the string is of the usual Fock space form:

$$\prod \alpha_{-n}^i \dots \alpha_{-m}^j |0\rangle \quad (104)$$

The eigenvalues of N and $\tilde{N}$ are just positive integers counting the number of oscillators acting on the vacuum. One can see that the fact that there is no privileged point (origin for measuring σ) in the closed string imposes the constraint $(N-\tilde{N})|\psi\rangle = 0$ for any string state $|\psi\rangle$. Now from Eq. (102) one sees that the lightest states are

$$\begin{aligned} N = \tilde{N} = 0 + m^2 = -8, \quad |0\rangle \\ N = \tilde{N} = 1 + m^2 = 0, \quad \alpha_{-1}^i \tilde{\alpha}_{-1}^j |0\rangle \end{aligned} \quad (105)$$

The vacuum state $|0\rangle$ is tachyonic. This is the well-known problem of the purely bosonic string. Then for $N = \tilde{N} = 1$ there are massless states with two space indices (i, j) . A symmetric combination of the indices leads to a state which can be identified with the graviton g_{ij} . Thus the bosonic string is a theory of gravity! This is a remarkable fact since we have put gravitation nowhere to start with. There are more massless fields, the trace $\alpha_{-1}^i \alpha_{-1}^i |0\rangle$ is a scalar field ("dilaton") and the antisymmetric combination of the two oscillators correspond to a two-index antisymmetric field.

The bosonic string we have been discussing does not seem very promising: it contains a tachyon and has no massless fermions (quark-lepton candidates) nor gauge bosons. It turns out that in order to extend the string concept to fermions one has to extend the simple Lagrangian in Eq. (90) in a supersymmetric way. Thus one introduces fermionic two-dimensional fields $\psi^\mu(z)$ and $\tilde{\psi}^\mu(\bar{z})$ ($\mu = 0, \dots, D-1$). The relevant action is

$$S = \int d\sigma d\bar{\tau} \left\{ \partial_z X^\mu \partial_{\bar{z}} X_\mu - \psi^\mu \partial_z \psi_\mu - \tilde{\psi}^\mu \partial_{\bar{z}} \tilde{\psi}_\mu \right\} \quad (106)$$

This action, as we discussed when we talked about supersymmetry, possesses an $N = 1$ SUSY in the two-dimensional world-sheet (not necessarily implying space-time supersymmetry in the D-dimensional space!). (X^μ, ψ^μ) form SUSY multiplets and the bosonic energy momentum tensor T_B has now a SUSY-partner T_F :

$$T_B = -\frac{1}{2} (\partial_z X^\mu)^2 + \frac{1}{2} \psi^\mu \partial_z \psi_\mu; \quad T_F = -\frac{1}{2} \psi^\mu \partial_z \psi_\mu \quad (107)$$

and similarly for $T_{\bar{B}}, T_{\bar{F}}$. As it happened with the purely bosonic case, one has to introduce "ghosts" upon quantization of fermions. One introduces "superconformal ghosts" γ (conformal dimension $d = -\frac{1}{2}$) and β ($d = 3/2$). One can see that the contribution of the (β, γ) system to the central charge is $c = 11$. Since the contribution to each $\psi(z)$ fermion to c is $\frac{1}{2}$, one gets an overall central charge for the "superstring" from Eq. (106):

$$c = D + \frac{1}{2} D - 26 + 11 = 3/2 \quad (D=10) \quad (108)$$

$$X \quad \psi \quad \beta, \gamma, \beta$$

Thus the superstring is truly consistent only in $D = 10$ space-time dimensions. In (107), T_F is the generator of local supersymmetry on the world-sheet. T_F generates also a local supersymmetry for antiholomorphic fields and one says that the superstring has $(1,1)$ local world-sheet supersymmetry, referring to the $N = 1$ SUSY generated by T_F and $T_{\bar{F}}$ on left- and right-movers. The superstring has two types of "fermionic" states depending on the boundary conditions of the ψ^μ fields. If $\psi^\mu(\sigma) = (-1)^F \psi^\mu(\sigma + \pi)$, one has the "Neveu-Schwarz" sector of the string and for $\psi^\mu(\sigma) = \psi^\mu(\sigma + \pi)$ one gets the "Ramond" sector. One can see that the string states in the first sector are bosons and the ones in the second are fermions. A remarkable point is that the theory can be shown to be inconsistent if it contains different numbers of fermions and bosons. This means that one must have a complete matching of bosonic and fermionic states, i.e., one must have a space-time supersymmetry $^{5/6}$ (and not only world-sheet SUSY). Thus the spectrum of the ten-dimensional superstring must be supersymmetric. Due to this fact this theory has no tachyons and the massless fields include a supersymmetric version of gravity, ten-dimensional $N = 2$ supergravity. We will not discuss here why we are forced to have a fermion-boson matching in the superstring. Let me only remark that when one considers a one-loop amplitude for the superstring, a property called "modular invariance" must be requested for the string

amplitudes³. Otherwise "global anomalies" to some extent similar to the ones we discussed for gauge theories appear. When one imposes modular invariance of one-loop amplitudes one can see that necessarily the spectrum of the fermionic string is supersymmetric (a projection killing states which are not supersymmetric appears)⁵. Let me also remark that this property of "modular invariance", which plays a prominent rôle in string theory, also makes that all one-loop amplitudes are ultra-violet finite both in the bosonic string and in the superstring.

Although the superstring described above has no tachyons and contains massless fermions, it still is not suitable to describe observed physics (apart from the fact that $D = 10$). There are no massless gauge bosons one could identify with gluons, W 's or the photon. However, one can construct new more promising types of supersymmetric strings, the heterotic strings. To do this one makes use of another remarkable property of closed strings: left-moving and right-moving states are completely independent and follow separate dynamics. This we have already observed above and it allows one to have different strings for left- and right-movers. The heterotic strings⁷⁷ are right-moving superstrings and left-moving bosonic strings:

$$\left[\begin{array}{c} R-M \\ \phi^\mu(z) \end{array} \right] \quad L-M \quad \left[\begin{array}{c} X^\mu(\bar{z}), \quad \mu = 0-25 \\ (109) \end{array} \right]$$

This sounds a bit puzzling since the superstring is ten-dimensional and the bosonic string is 26-dimensional. However, one can see that for such a theory to be consistent, 16 of the 26 dimensions of the bosonic string have to be curled up ("compactified") on a 16-dimensional torus of fixed radii, so that the actual number of dimensions of a heterotic string is 10. "Compactified" on a torus means that one identifies the co-ordinates X^i , $i = 10-25$ with their translations by some "lattice vectors" $\{e_i\}$:

$$X^i = X^i + R \sum_{i=1}^{16} n_i e_i \quad (110)$$

where R is the "radius of compactification". To have an idea consider the case of a two-dimensional torus as in Fig. 27a. It may be obtained from Fig. 27b by gluing the sides A (identification $X = X + e_1$) and the sides B (identification $X = X + e_2$). When we force the co-ordinates X^i ($i = 10-25$) to move only on a 16-dimensional torus, the momenta associated to these co-ordinates p_i are quantized. One can further show that the "modular invariance" property we mentioned above requires the p_i to be very specific types of vectors⁷⁸ belonging to the "root lattices" of the Lie algebras of $E_8 \times E_8$ and $SO(32)$ ⁷⁸. I will not explain how this works out since it is rather technical. Let me just comment on the spectrum of these heterotic strings. It turns out that the modulus squared of p_i must be an even number (in string units, $R^2 = \alpha'$, where α' is the string tension which we have taken $\alpha' = \frac{1}{2}$ all over). The mass formula for the heterotic strings has the form (e.g., Ramond R -M sector)

Fig. 27

$$m^2 = m_R^2 + m_L^2; \quad \left\{ \begin{array}{l} \frac{m_R^2}{4} = N \\ \frac{m_L^2}{4} = N + \frac{p^2}{2} - 1 \end{array} \right. \quad (111)$$

where N , $\tilde{N}$ are the number operators. Notice there is no tachyon in the right-handed (superstring) contribution. The condition $(m_R^2 - m_L^2)/4 > 0$ (analogous to the $N = \tilde{N}$ condition of the bosonic string) makes then that no tachyons may appear in the heterotic string. There are massless states, e.g., for $N = \tilde{N} = 0$ and $(p^i)^2 = 2$. One can see that the vectors of the $E_8 \times E_8$ root-lattices with $(p^i)^2 = 2$ are just the roots. As is well known, each charged generator of a group has associated a "root vector". This is just telling us that the set of generators of a group $E_8 \times E_8$ (gauge bosons) will be massless. Thus in the heterotic strings there are massless gauge bosons. Furthermore, as happened with the "superstring", there is space-time supersymmetry. Altogether one can see that the massless spectrum is $N = 1$ supergravity coupled to an $E_8 \times E_8$ [or $SO(32)$] super Yang-Mills theory.

The heterotic strings seem more promising than unification theories since they have massless gauge bosons and also charged fermions (the SUSY partners of the gauge bosons). However, the theories are formulated in ten dimensions instead of four. This is easy to mend since not necessarily all the co-ordinates in Eq. (109) must admit a space-time interpretation. Thus it could well happen that the six extra unwanted dimensions were also "curled up" (compactified) on some torus (or other space). As an example, one could compactify the six extra right and left co-ordinates in Eq. (109) $X^i(Z)$, $X^i(Z)$ ($i = 4-9$) in three copies of the 2-torus of Fig. 27

$$X^i = X^i + \sum_{a=1}^9 R_a n_a e^i_a \quad (112)$$

where e^i_a , e^i_b , e^i_c could be orthogonal unit vectors as in Fig. 27b. R_a , the radii of compactification, are arbitrary since there are both left and right co-ordinates. However, it turns out that in order to get the correct size for the gravitational ($\sim 1/M_P^2$) and gauge ($g \sim 1$) interactions all the radii R_a of compactification have to be $R_a \sim 1/M_P$ and hence the size of the extra six dimensions is tremendously small. This means that to "probe" those extra dimensions we would need energies of the order of the Planck mass, $M_P \sim 10^{18}$ GeV. Thus at ordinary energies the physical number of dimensions will be just four.

What we have just been discussing is called toroidal string compactification and this method gives us the simplest four-dimensional strings. However, it is not yet what we want. The reason is that the theories so obtained in four dimensions are not chiral. This happens because the fermions (spinors) of an $N = 1$, $D = 10$ theory from which we started split into four spinors in four dimensions. This means that if we had one supersymmetry ($N = 1$) in $D = 10$ we will have four supersymmetries ($N = 4$) in $D = 4$. But we already mentioned that in four dimensions only $N = 1$ supersymmetry can be chiral. Furthermore, the groups $E_8 \times E_8$ or $SO(32)$ are not amongst the interesting unification groups we discussed in the first lecture.

One way to solve this problem is to compactify the six extra dimensions on an "orbifold"⁷⁹. Orbifolds of the type we are discussing are relatives of tori⁸⁰. You start from a torus and not only identify points which differ by some lattice vector but you identify points which are related by some discrete symmetry P of the torus. Consider as an example the two-torus in Fig. 27b. Now we are going to identify all the points on the torus which can be obtained from each other as a reflection with respect to the origin in Fig. 27b. Thus, e.g., the points E and E' in Fig. 28 are identified since they can be obtained from each other after a reflection with respect to the origin (plus a shift by $\vec{e}_1 + \vec{e}_2$). In this way all the points in the original torus are identified with points in the shaded area. One expresses this identification mathematically by saying that the above two-dimensional orbifold is the result of "dividing" the torus T^2 by an isometry P , $O_2 = T^2/P$. Again one can obtain a four-dimensional string by compactifying the six unwanted dimensions, e.g., on a six-dimensional orbifold with three copies of O_2 , i.e., $O_6 \equiv O_2 \times O_2 \times O_2$. Of course there are many more tori and orbifolds one can construct. The important point is that only the fields of the compactified string which are invariant under the symmetries P will remain as physical fields in the string spectrum. Thus we may get rid of three of the four supersymmetries of toroidal compactifications by appropriately choosing a symmetry P which kills three of the four existing massless gravitini. There are plenty of possibilities to do so leading each one to different four-dimensional strings.

Fig. 28

We thus have four-dimensional strings with $N = 1$ supersymmetry and massless gauge bosons and fermions. Again this is not enough since the gauge groups we have [$E_6 \times E_8$ or $SO(32)$] are not adequate (recall our discussion in the first lecture). Thus we want to break the gauge group down to some more promising subgroup. A natural process is to do something similar to the "dividing by a symmetry P " of the orbifold. One can "mod" the gauge degrees of freedom of the heterotic string by some subgroup G of $E_6 \times E_8$ [or $SO(32)$]. Then only the string states invariant under the combined action on space and gauge degrees of freedom $P \times G$ remain as physical states. It turns out that "modular invariance" constraints strongly correlate the possible forms of P and G . The simplest solution is to have similar forms for the symmetries P acting on the six compactified co-ordinates and the symmetry G acting on the gauge degrees of freedom. This type of compactifications is called (2,2) or "standard embedding" compactifications. The case of the $E_6 \times E_8$ string is particularly interesting. By acting with an action G inside one of the E_8 's, a (2,2) compactification leads to a gauge group

$$E_6 \times E_8 \rightarrow E_6 \times E_8 \quad (113)$$

and E_6 is one of the interesting groups for unification discussed in the first lecture. We are thus left with a $D = 4$, $N = 1$ supersymmetric theory coupled to a gauge group $E_6 \times E_8$. In fact one can see that the E_6 and E_8 sectors of the theory are decoupled from each other (only interact gravitationally amongst themselves) so that the E_8 world is a "hidden world". It is remarkable how similar this situation is to the one we discussed for the supersymmetric standard model. Thus the existence of a "hidden sector" of the theory is a generic fact in four-dimensional strings. One can also see that the chiral matter obtained in this type of strings is just a number of E_6 generations (27's) and antighenerations (27's). The number of quark-lepton families will be the net number ($N_{27} - N_{\bar{27}}$) since pairs $27 \times \bar{27}$ can in general gain large masses and disappear from the physical low energy spectrum (although this may be problematic in practice).

There are two other known methods to construct four-dimensional (2,2) strings. One of them is the compactification of the six extra dimensions in a particular class of six-dimensional manifolds called "complete-intersection Calabi-Yau manifolds" (CICY)⁸¹. The problem with this method is that the conformal field theory associated to these four-dimensional strings is not known. This was, however, historically the first method attempted and much of the string "phenomenology" done up to now has a "Calabi-Yau inspiration". There is a third method⁸² which is also very interesting. It just constructs a conformal invariant theory with the usual constraints of modular invariance and overall vanishing central charge, $c = 0$, forgetting about the possible interpretations of the unwanted degrees of freedom as extra dimensions. Consider first the right-moving (superstring) sector of the heterotic string. The degrees of freedom we want to get rid of (or at least do not want to have as physical dimensions) are $X^I(Z)$, $I = 4-9$ and $\psi^I(Z)$ ($I = 4-9$). The central charge of this superconformal system is $c = 9 = 6 \times 1 + 6 \times \frac{1}{2}$. Thus the idea is that any $c = 9$ superconformal system leading to a modular invariant string is in principle acceptable (the overall central charge will vanish as required). The simplest superconformal models in the market are the so-called "minimal models". These models are labelled by an integer $K = 1, \dots, \infty$ and have central charges

$$c = \frac{3K}{K+2}, \quad K = 1, \dots, \infty \quad (114)$$

so that they have $1 < c < 3$. The idea⁸² now is to combine a set of models of this type so that the overall central charge is $c = 9$. For example, nine copies of the $K = 1$ model or six copies of the $K = 2$ model, etc. All this refers to the right-moving sector. There is then a standard way⁸² to build the left-moving sector of the heterotic string so that the result is a (2,2) modular invariant for dimensional string.

These three methods to construct (2,2) four-dimensional strings are not unrelated. There are string models which may be constructed in more than one way^{83,84}. The situation concerning the relationship among these constructions is summarized in Fig. 29. How do these models look compared to observed physics? A first selection criterion is the number of generations. A complete catalogue of all these (2,2) models is still to be made but a lot of classification work has already been done. The situation at present is as follows. Concerning orbifolds, models with an Abelian P symmetry ("Abelian orbifolds") have been studied in some detail^{85,86}. The number of net generations found is always a multiple of six. Concerning non-Abelian orbifolds, although a complete classification

Fig. 29

has not been done, it has been shown that no three-generation model⁷⁹ can be found [in the case of (2,2) compactifications!]. There is a unique complete-intersection Calabi-Yau model with net number of generations = 3 (nine generations and six antigenerations)⁸⁶. Its phenomenology has been studied in some detail⁸⁷ although the lack of a known conformal field theory did not allow for a string formulation of the model. Finally, the analysis of the $N = 2$ superconformal constructions mentioned above is still incomplete⁸⁸. Some of the models seem to be equivalently obtained from orbifolds⁸³. However, and most interestingly, some of the models seem to correspond to some known CICY_{82,84}. In particular, the three-generation model mentioned above has a formulation as an $N = 2$ superconformal construction⁸². This is interesting since one can in principle compute all the details of the model, including Yukawa couplings. Still, the possibility that new three-generation (2,2) models exist is open.

We have seen that amongst the relatively large number of (2,2) models known to exist up to now, only one has a net number of generations equal to three; it is not so easy to get three-generation models. However, this model has also six pairs of $(\overline{27}+27)$ matter fields and it is still not clear whether they get a large mass and decouple from low-energy physics. This should be the case since otherwise the gauge coupling constants of the standard model would get too large before unification. Moreover, preliminary field-theoretical studies seem to show that the breaking of E_6 down to the standard model while maintaining sufficient proton stability and other experimental constraints may perhaps be possible but in a rather contrived way. Are there any other types of four-dimensional strings which may lead to realistic physics? The answer is yes. The (2,2) four-dimensional strings are the simplest solution to the conditions of modular invariance but certainly not the only ones⁷⁹. In orbifold language, there are many other possible actions G on the gauge degrees of freedom which may be consistent. Their common feature is that the gauge group is no longer $E_6 \times E_8$ but many other possibilities [including $SO(10)$, $SU(5)$ or even the standard model group $SU(3) \times SU(2) \times U(1)$] are now available. These other constructions are called (2,0) models. Very little is known about (2,0) models obtained from CICY or $N = 2$ superconformal models. On the other hand, many such orbifold models have been constructed^{85,89,90}, some of them with promising phenomenology. Other techniques in order to get (2,0) models also exist⁹²⁻⁹⁴.

The simplest quasi-realistic four-dimensional strings constructed are based on a (2,0) version of the " Z_3 " orbifold⁹¹. It is called so because the orbifold P symmetry is an Abelian Z_3 group. I will not give the details here and refer the reader to the original literature^{86,91,85}.

This class of models has three generations of quarks and leptons plus additional Higgs doublets and singlets. One of the most promising models has gauge group^{95,96}

$$SU(3)_C \times SU(2)_L \times U(1)_Y \times [U(1)_L \times U(1)' \times SO(10)'] \quad (115)$$

where the $SO(10)' \times U(1)'$ is "hidden" from the standard model physics (it only interacts with usual particles through gravitational interactions). The extra $U(1)_L$ gauge boson may or may not be there depending on the version of the model. The particle content is three generations of quarks and leptons plus a number of $SU(2)_L$ doublets and singlets. Preliminary studies of the Yukawa couplings have also been done. A different formalism ("four-dimensional fermionic strings")⁹³ has been used to obtain other (2,0) strings leading to a grand unified "flipped" $SU(5) \times U(1)_3$ model⁹⁷.

The examples of quasi-realistic four-dimensional strings found are encouraging but we are still far away from a situation in which (a) a given model is consistent with absolutely all the known phenomenological constraints and (b) any new prediction can be given. A point which is encouraging is that it is not so easy to find three-generation four-dimensional strings. If one insists on mimicking more and more details of the observed physics of quarks and leptons, more and more string candidates are ruled out. At the moment I do not think that anybody can seriously claim that a completely realistic string exists. Points like sufficient stability of the proton are hard to fulfill. Once the gauge symmetry is broken down to the standard model there is in general no symmetry forbidding the B - and L -violating operators we discussed in Eq. (63). One can "cook" the models so that these kinds of couplings do not appear but there does not seem to exist any symmetry principle based on string physics which may prefer a conservation of baryon or lepton number. In fact, strings do not like continuous global symmetries, they prefer gauge symmetries. A possible solution to this baryon stability problem of four-dimensional strings is the conservation of an unbroken extra gauge symmetry forbidding the dangerous couplings of Eq. (63). That is precisely what happens if an unbroken $SU(2)_L \times SU(2)_R \times U(1)_{B-L}$ symmetry remains close to the weak scale. This may be obtained in some four-dimensional string models⁹⁸.

In spite of the difficulties mentioned, four-dimensional strings are a big improvement over previous schemes. First of all, superstrings are believed to be finite theories. They are really unified theories, including for the first time a consistent theory of quantum gravity. Furthermore they are highly constrained theories. Thus, given a four-dimensional string, the particle content and gauge interactions are completely fixed. As an example, no exotic massless representations (e.g., colour sextet quarks or similar objects) are in general allowed. Also no adjoint Higgs fields ever appear in the massless sector of a string. Yukawa couplings are also determined by the corresponding conformal field theory associated to the model. This makes string models very easy to rule out. Let us remark, however, that all string models have some neutral scalar fields ("moduli" and "marginal operators") whose (unknown) vacuum expectation values govern the precise numerical values of quantities like Yukawa couplings. Then one may expect from a given model a prediction on what couplings are allowed and which ones are forbidden or qualitative features of the quark-lepton spectrum but no precise prediction on quark and lepton masses is to be expected as long as we do not find a dynamical reason which could explain why some four-dimensional strings are dynamically selected.

How do the topics of the previous lectures, grand unification and low-energy supersymmetry, fit in the string context? As we mentioned already, some of the grand unification ideas are also present in four-dimensional strings. Thus there are bigger gauge groups [Eg in $D = 10$ or maybe even some GUT like $SU(5)$, E_6 or $SO(10)$ in some models] although the way in which they appear is more sophisticated. Some models (but not all) maintain the standard predictions for $\sin^2\theta_W$. In general quark-lepton mass relations disappear and the Yukawa couplings are given by correlation functions between the conformal fields ("vertices") associated to each particle. Concerning low-energy SUSY, it fits quite naturally inside the four-dimensional string context. $N = 1$ supersymmetry in $D = 4$ guarantees the absence of tachyons in the string. Thus this provides us with another argument (independent of the use for solving the "naturalness" problem) in favour of the existence of a space-time supersymmetry in four dimensions. Furthermore, the existence of a "hidden sector" in which SUSY-breaking may occur appears automatically in string models. A natural possibility for the SUSY-breaking mechanism is a dynamical SUSY-breaking triggered by condensation of the gauginos⁹⁹ of the hidden gauge sector (i.e., $\langle\lambda\lambda\rangle \neq 0$). The size of this breaking can be hierarchically small compared to the Planck mass and may explain why the weak scale is so small compared to the Planck mass. Thus one would expect as the low-energy limit of a four-dimensional string something similar to the supersymmetric standard model described in the previous lecture. Of course the possibility exists of having some extra particles like new gauge interactions or heavy leptons, etc. The experimental detection of these hypothetical particles would be very important. Meanwhile, unfortunately, there is no serious string argument which predicts the existence of these new states.

6. FINAL COMMENTS

Four-dimensional strings seem to provide us with the first really unified theories of all interactions including gravity. Unfortunately strings are theories which reveal themselves at energies of the order of the Planck mass, $\sim 10^{18}$ GeV and we can only guess at their existence (if they are there) indirectly. Since direct experimental tests of strings seem out of question the risk exists of getting very little progress (if any) in the complete construction of the theory. Past experience tells us that experimental input has played a key rôle in finding the correct field theory of interactions amongst elementary particles. However, string theories are so far superior to the previously considered unifying schemes that one just cannot throw them away in the dustbin. An effort must be made both in understanding the theory better and trying to make contact with observed physics.

Fortunately enough, the prospects for low-energy SUSY to be checked experimentally are much brighter since new particles within the reach of present or future accelerators are predicted. Certainly, if SUSY particles are not found at supercollider energies, low-energy supersymmetry would be ruled out. If SUSY particles are found the detailed study of their properties would give us precious information about the SUSY breaking process and, maybe, about Planck mass physics.

ACKNOWLEDGEMENTS

I warmly thank Tom Ferbel for the nice atmosphere created at the School. Special thanks go to Captain Guido Martinelli whose skills as navigator were clearly demonstrated in our attempted trip to Buck Island. His providential command saved us from disaster. I also thank C. Albajar and P. Paus for providing me with some experimental information.

REFERENCES

1. For a review, see:
P. Langacker, Physics Reports 72C:185 (1981);
R. Slansky, Physics Reports 79:1 (1981);
G.G. Ross, "Grand Unified Theories", Benjamin Inc. (1984).
J. Mess and J. Bagger, "Supersymmetry and Supergravity", Princeton University Press, Princeton (1983);
P. West, "Introduction to Supersymmetry and Supergravity", World Scientific (1986).
For the phenomenology of supersymmetry, see:
H.P. Nilles, Physics Reports C10:1 (1984);
H. Haber and G. Kane, Physics Reports C117:75 (1985);
R. Barbieri and S. Ferrara, Surveys in High Energy Physics 4:33 (1983).
M.B. Green, J.H. Schwarz and E. Witten, Superstring Theory, Vols. I and II, Cambridge University Press (1986);
D. Gross, in Proceedings of the 1986 ASI School (Virgin Islands). Plenum Press (1987).
J. Schwarz, "Introduction to Superstrings", CALT-68-1290 (1985);
M. Peskin, "Introduction to String and Superstring Theory", SLAC-PUB-4251 (1987).
S. Adler, Phys.Rev. Vol. 177:2426 (1969);
J.S. Bell and R. Jackiw, Nuovo Cimento 60A:47 (1969).
L. Alvarez-Gaumé and E. Witten, Nucl.Phys. B234:269 (1983).
See, e.g.:
M. Luscher and P. Weisz, Nucl.Phys. B295:65 (1988);
A. Hasenfratz et al., Phys.Lett. 199:531 (1987);
M. Grzadowski and M. Lindner, Phys.Lett. 178B:81 (1986).
G. 't Hooft, in "Recent Developments in Gauge Theories", ed. by G. 't Hooft et al., Plenum Press, New York (1980), p. 135.
S. Weinberg, Phys.Rev. D13:974 (1976) and D19:1277 (1979);
L. Susskind, Phys.Rev. D20:2619 (1979);
E. Farhi and L. Susskind, Physics Reports 74C:277 (1981).
S. Dimopoulos and J. Ellis, Nucl.Phys. B182:505 (1981);
See, however:
S. Dimopoulos, H. Georgi and S. Raby, Phys.Lett. 127B:101 (1983);
B. Holdom, Phys.Rev. D24:1441 (1981);
T. Appelquist and L.C.R. Wigewardhana, Phys.Rev. D35:774 (1987).
L. Abbot and E. Farhi, Nucl.Phys. B189:547 (1981);
M. Claudson, E. Farhi and R. Jaffe, Phys.Rev. D34:873 (1986).
M. Chanowitz and M.K. Gaillard, Nucl.Phys. B261:379 (1985);
M. Chanowitz, M. Golden and H. Georgi, Phys.Rev. D36:149 (1987);
Phys.Rev.Lett. 57:2344 (1986).
H. Georgi and S.L. Glashow, Phys.Rev.Lett. 32:32 (1974).
J. Pati and A. Salam, Phys.Rev. D10:275 (1974).
H. Georgi, H. Quinn and S. Weinberg, Phys.Rev.Lett. 33:451 (1974).
H. Georgi, unpublished;
H. Fritzsch and P. Minkowski, Ann.Phys. 93:193 (1975);
H. Georgi and D.V. Nanopoulos, Nucl.Phys. B155:59 (1979).
H. Georgi and D.V. Nanopoulos, Nucl.Phys. B159:16 (1979);
F. Del Aguila and L.E. Ibañez, Nucl.Phys. B177:60 (1981).
F. Gusev and M. Serdaroglu, Nuovo Cimento 65A:337 (1981) and references therein;
Y. Achiman and B. Stech, Phys.Lett. 77B:389 (1978);
P. Ramond, in Proceedings of Sanibel Symposium (1979);
R. Barbieri and D.V. Nanopoulos, Phys.Lett. 91B:369 (1980).
A.J. Buras, J. Ellis, M.K. Gaillard and D.V. Nanopoulos, Nucl.Phys. B195:66 (1978).
See, e.g.: A. Savoy-Navarro, in Proceedings of ASI Summer School (1986), Plenum Press (1987).

21. M. Gell-Mann, P. Ramond and R. Slansky, unpublished (1979); T. Yanagida, Proceedings Workshop on the Unified Theory and the Baryon Number in the Universe, KEK, Japan (1979).
22. See, e.g.: G.G. Ross, Proceedings of the 1987 International Symposium on Lepton and Photon Interactions at Higher Energies, DESY, Hamburg (1987), CERN Preprint TH. 4881 (1987); See also: A. Savoy-Navarro in Ref. 20.
23. G. Blewitt et al., Phys.Rev.Lett. 54:22 (1985).
24. L.E. Ibañez, Nucl.Phys. B181:105 (1981).
25. W. Marciano and A. Sirlin, Phys.Rev. D22:2095 (1980) and Nucl.Phys. B189:442 (1981); C.H. Llewellyn Smith, G.G. Ross and J.F. Wheeler, Nucl.Phys. B177:263 (1981); G. Costa et al., CERN Preprint TH. 4675 (1987), LBL-23271 (1987); U. Amaldi et al., Phys.Rev. D36:1385 (1987).
26. Y.A. Gol'fand and E.P. Likhtman, JETP Lett. 13: (1971); D.V. Volkov and V.P. Akulov, JETP Lett. 16:438 (1972); J. Wess and B. Zumino, Nucl.Phys. B70:39 (1974).
27. M. Veltman, Acta Phys.Polon. B12:437 (1981); L. Maiani, Proceedings Summer School of Gif-Sur-Yvette (Paris 1980); E. Witten, Nucl.Phys. B188:513 (1981).
28. A. Salam and J. Strathdee, Nucl.Phys. B76:477 (1974); S. Ferrara, J. Wess and B. Zumino, Phys.Lett. 51B:239 (1974); S. Gates, M. Grisaru, M. Rocek and W. Siegel, "Superspace", Frontiers in Physics, Benjamin Inc. (1983).
29. P. Fayet and J. Iliopoulos, Phys.Lett. 51B:461 (1974); L. O'Raifeartaigh, Nucl.Phys. B96:331 (1975); P. Fayet, Phys.Lett. 58B:67 (1975).
30. L. Girardello and M. Grisaru, Nucl.Phys. B194:65 (1982); K. Harada and N. Sakai, Progr.Theor.Phys. 67 (1982) 1877.
31. P. Fayet, Phys.Lett. 69B:489 (1977); 84B:416 (1979); 78B:417 (1978); G. Farrar and P. Fayet, Phys.Lett. 79B:442 (1978); 89B:191 (1980).
32. H. Haber and G. Kane, Physics Reports C117:75 (1985).
33. S. Dimopoulos and H. Georgi, Nucl.Phys. B93:150 (1981); N. Sakai, Z.Phys. C11:153 (1982).
34. S. Dimopoulos, S. Raby and F. Wilczek, Phys.Rev. D24:1681 (1981); L.E. Ibañez and G.G. Ross, Phys.Lett. 105B:439 (1982); M.B. Einhorn and D.R.T. Jones, Nucl.Phys. B196:475 (1982).
35. L.E. Ibañez, Phys.Lett. 126B:196 (1983).
36. S. Weinberg, Phys.Rev. D26:287 (1982); N. Sakai and T. Yanagida, Nucl.Phys. B197:533 (1982); S. Dimopoulos, S. Raby and F. Wilczek, Phys.Lett. 112B:133 (1982); J. Ellis, D.V. Nanopoulos and S. Rudaz, Nucl.Phys. B202:43 (1982).
37. L. Hall and M. Suzuki, Nucl.Phys. B231:419 (1984); I.H. Lee, Nucl.Phys. B246:120 (1984); F. Zwirner, Phys.Lett. B132:103 (1983); R. Barbieri and A. Masiero, Nucl.Phys. B267:679 (1986); S. Dimopoulos and L. Hall, Phys.Lett. B196:135 (1987).
39. L.E. Ibañez, Phys.Lett. 118B:73 (1982); Nucl.Phys. B218:514 (1983).
40. R. Barbieri, S. Ferrara and C. Savoy, Phys.Lett. 119B:343 (1982).
41. P. Nath, R. Arnowitt and A. Chamseddine, Phys.Rev.Lett. 49:970 (1982); H.P. Millea, M. Srednicki and D. Wyler, Phys.Lett. 120:346 (1982).
43. R. Barbieri and S. Ferrara, Surveys in High Energy Physics 4:33 (1983); E. Cremmer, S. Ferrara, L. Girardello and A. Van Proeyen, Nucl.Phys. B212:413 (1983).
44. H.P. Millea, Phys.Lett. 115B:193 (1982).
45. S. Dimopoulos and S. Raby, Nucl.Phys. B219:479 (1983); M. Dine and W. Fischler, Nucl.Phys. B204:346 (1982); J. Polchinski and L. Susskind, Phys.Rev. D26:3661 (1982).

46. K. Inoue et al., Progr.Theor.Phys. 67:1859 (1982).
47. L.E. Ibañez and C. Lopez, Phys.Lett. 126B:54 (1983); Nucl.Phys. B233:511 (1984).
48. L. Alvarez-Gaumé, J. Polchinski and M. Wise, Nucl.Phys. B221:495 (1983).
49. J. Ellis, J. Hagelin, D.V. Nanopoulos and K. Tamvakis, Phys.Lett. 125B:275 (1983).
50. K. Inoue et al., Progr.Theor.Phys. 68:927 (1982).
51. L.E. Ibañez and G.G. Ross, Phys.Lett. 110B:227 (1982).
52. L.E. Ibañez, C. Lopez and C. Muñoz, Nucl.Phys. B256:218 (1985).
53. L.E. Ibañez, Phys.Lett. 137B:160 (1984); J. Hagelin, G. Kane and S. Raby, Nucl.Phys. B241:638 (1984).
54. C. Muñoz, Ph.D. Thesis, Universidad Autónoma de Madrid (1986), unpublished;
55. L.E. Ibañez and C. Muñoz, Paper in preparation (1989).
55. R. Flores and M. Sher, Ann.Phys. 148:95 (1983); H.P. Millea and M. Nussbaumer, Phys.Lett. 145B:73 (1984); P. Majumdar and P. Roy, Phys.Rev. D30:2432 (1984).
56. J. Ellis and G.G. Ross, Phys.Lett. 117B:397 (1982); A. Chamseddine, P. Nath and R. Arnowitt, Phys.Lett. 129B:445 (1983); P. Dias, S. Nandi and X. Tata, Phys.Lett. 129B:451 (1983).
57. S. Weinberg, Phys.Rev.Lett. 50:387 (1983); B. Grinstein, J. Polchinski and M. Wise, Phys.Lett. 130B:285 (1983).
58. R. Barbieri and G.F. Giudice, Nucl.Phys. B296:75 (1988).
59. J. Ellis and D.V. Nanopoulos, Phys.Lett. 110B:44 (1982); R. Barbieri and R. Gatto, Phys.Lett. 110B:211 (1982); T. Inami and G.S. Lim, Nucl.Phys. B207:593 (1982); M. Suzuki, Phys.Lett. 115B:46 (1982); M. Duncan, Nucl.Phys. B221:285 (1983); J. Donoghue, H.P. Millea and D. Wyler, Phys.Lett. 128B:55 (1983).
60. M. Davier, Proceedings 23rd International Conference on High-Energy Physics, Berkeley (1986), ed. S.G. Loken, World Scientific (1987), p. 25; S. Whitaker, *ibid.*, p. 602.
61. F. Pauss, CERN EP/88-158 (1988).
61. G. Arnison et al. (UAI Collaboration), Europhys.Lett. 1:327 (1986); C. Alajar et al. (UAI Collaboration), CERN preprint CERN-EP/88-168 (1988).
62. C. Hearty et al. (ASP Collaboration), Phys.Rev.Lett. 58:1711 (1987).
63. E. Fernandez et al. (MAC Collaboration), Phys.Rev.Lett. 54:1118 (1983).
64. G. Kane and J. Leveillé, Phys.Lett. 112B:227 (1982); P. Harrison and C.H. Llewellyn Smith, Nucl.Phys. B213:223 (1983); M.J. Herrero, L.E. Ibañez, C. Lopez and F.J. Yndurain, Phys.Lett. 132B:199 (1983); 145B:430 (1984); E. Reya and P. Roy, Phys.Lett. 141B:442 (1984); Phys.Rev. D32:645 (1985); J. Ellis and H. Kowalski, Phys.Lett. 142B:441 (1984); Nucl.Phys. B259:109 (1985).
65. C. Alajar et al. (UAI Collaboration), Phys.Lett. B198:261 (1987); See also: F. Pauss, CERN-EP/88-158 (1988) and references therein.
66. R. Ansari et al. (UAI Collaboration), Phys.Lett. B195:613 (1987).
67. H. Goldberg, Phys.Rev.Lett. 50:1419 (1983); J. Ellis, J. Hagelin, D.V. Nanopoulos and M. Srednicki, Phys.Lett. 127B:233 (1983); J. Ellis et al., Nucl.Phys. B238:453 (1984).
68. S. Dimopoulos and L. Hall, Phys.Lett. B207:216 (1988).
69. L. Hall, LBL Preprint LBL-26059 (1988).
70. A. Polyakov, Phys.Lett. 103B:207 (1981).
71. A. Belavin, A. Polyakov and A. Zamolodchikov, Nucl.Phys. B241:333 (1984).

72. D. Friedan, E. Martinec and S. Shenker, Nucl.Phys. B271:93 (1986).
73. V. Alessandrini, D. Amati, M. Le Bellac and D. Olive, Physics Reports 1C:170 (1971);
S. Mandelstam, Physics Reports 13C:259 (1974);
J. Scherk, Rev.Mod.Phys. 47:123 (1975).
74. P. Ramond, Phys.Rev. D3:2415 (1971);
A. Neveu and J. Schwarz, Nucl.Phys. B31:86 (1971); Phys.Rev. D4:1109 (1971).
75. F. Gliozzi, D. Olive and J. Scherk, Nucl.Phys. B122:253 (1977).
76. M. Green and J. Schwarz, Phys.Lett. 136B:367 (1984).
77. D. Gross, J. Harvey, E. Martinec and R. Rohm, Nucl.Phys. B256:468 (1985) and B267:75 (1986).
78. M.B. Green and J.H. Schwarz, Nucl.Phys. B255:93 (1985); Phys.Lett. 149B:117 (1984).
79. L. Dixon, J. Harvey, C. Vafa and E. Witten, Nucl.Phys. B261:651 (1985); Nucl.Phys. B274:285 (1986).
80. L.E. Ibañez, CERN Preprint TH. 4769 (1987), to be published in the Proceedings of the XVIIIth International GIFT Seminar on Theoretical Physics, World Scientific, Singapore (1988).
81. P. Candelas, G. Horowitz, A. Strominger and E. Witten, Nucl.Phys. B258:46 (1985);
P. Green, T. Hübsch and C.A. Lütken, CERN Preprint TH. 4933 (1987), to appear in Class.Quant.Gravity.
82. D. Gepner, Nucl.Phys. B296:757 (1988); Phys.Lett. 199B:380 (1987).
83. A. Font, L.E. Ibañez and F. Quevedo, CERN Preprint TH. 5217 (1988), to appear in Phys.Lett. B.
84. T. Eguchi, H. Ooguri, A. Taormina and S. Yang, Preprint UT-536 (1988);
K. Li and N. Warner, CERN Preprint TH. 5047 (1988);
B. Greene, C. Vafa and N. Warner, HUTP-88/A047 (1988).
85. L.E. Ibañez, J. Mas, H.P. Nilles and F. Quevedo, Nucl.Phys. B301:157 (1988).
86. S.T. Yau, in Proceedings of the Argonne Symposium on Anomalies, Geometry and Topology, World Scientific, Singapore (1985).
87. B. Greene, K. Kirklin, P. Miron and G. Ross, Nucl.Phys. B278:667 (1986); B279:574 (1986).
88. C.A. Lütken and G.G. Ross, CERN Preprint TH. 5058 (1988);
M. Lynker and R. Schimfirk, Texas Preprint UTG-17-88 (1988).
89. L.E. Ibañez, H.P. Nilles and F. Quevedo, Phys.Lett. B187:25 (1987).
90. L. Dixon, Ph.D. Thesis, Princeton (1985), unpublished.
91. L.E. Ibañez, J.E. Kim, H.P. Nilles and F. Quevedo, Phys.Lett. B191:282 (1987).
92. W. Lerche, A.N. Schellekens and N. Warner, CERN Preprint TH. 5155 (1988), to appear in Physics Reports.
93. I. Antoniadis, CERN Preprint TH. 5199 (1988), and references therein.
94. K. Narain, M. Sarmadi and C. Vafa, Nucl.Phys. B288:951 (1987).
95. A. Font, L.E. Ibañez, H.P. Nilles and F. Quevedo, Phys.Lett. B210:101 (1988).
96. J.A. Casas and C. Muñoz, Phys.Lett. 209B:214 (1988); Oxford-TP-32/88 (1988); CERN Preprint TH. 5052 (1988);
M. Quirós, IEM-Madrid Preprint (1988), and references therein.
97. I. Antoniadis, J. Ellis, J. Hagelin and D.V. Nanopoulos, CERN Preprint TH. 4931 (1987), 5005 (1988).
98. A. Font, L.E. Ibañez, F. Quevedo and A. Sierra, in preparation.
99. J.-P. Derendinger, L.E. Ibañez and H.P. Nilles, Phys.Lett. 155B:65 (1985);
M. Dine, R. Rohm, N. Seiberg and E. Witten, Phys.Lett. 156B:55 (1985).