CERN-TH 5351/89
IOA-223/89

LIGHT NEUTRINOS WITH LARGE MAGNETIC
MOMENTS AND THE SOLAR NEUTRINO
PROBLEM

G. K. Leontaris*

CERN-Geneva

and

K. Tamvakis

Theoretical Physics Division

The University of Ioannina

GR-451 10 Ioannina, Greece

ABSTRACT

We analyse the question of having a large enough neutrino magnetic moment in order to implement the VVO mechanism in the framework of an $SU(4) \times SU(2)_L \times SU(2)_R$ model, derivable from four dimensional superstrings. We find that the radiative correction to the seesaw type neutrino mass matrix can generate an appreciable magnetic moment without any seesaw type suppression.

*On leave from Theoretical Physics Division, The University of Ioannina, GR-45110 Ioannina, Greece

The long standing solar neutrino puzzle [1-3] has motivated a couple of years ago the suggestion of Voloshin, Vysotsky and Okun [4] that the deficit of solar neutrinos is a consequence of a magnetic moment type of interaction of the neutrinos with the magnetic field of the outer layers of the sun. A ν_e magnetic moment in the range

$$\mu_{\nu_e} = (10^{-11} - 10^{-10}) \mu_B$$

($\mu_B = e/2m_e$ is the Bohr magneton) is needed in order to achieve the required conversion into inert right-handed or higher-family neutrinos which are undetectable. Presently the most stringent upper bound on μ_{ν_e} is $0.7 \times 10^{-10} \mu_B$ [5] and the explanation of the solar neutrino deficit via the VVO mechanism is an allowed possibility.

The standard electroweak model with an additional Higgs isodoublet that gives a Dirac mass to ν_e through the coupling to a gauge singlet right-handed neutrino gives too small a value for μ_{ν_e} ($= 10^{-18} \mu_B$) [6] and people have looked into extensions of the standard model that can predict a μ_{ν_e} at the required value range, introducing exotic particles [7]. An important constraint that is present in all models of neutrino magnetic moment is the constraint imposed by the upper limit on the ν_e mass [8]. Large neutrino magnetic moments have to exist with small neutrino masses. A recent examination of this question in the framework of the standard seesaw model of neutrino masses has given a negative answer, i.e., too small a μ_{ν_e} [9].

In the present article we reexamine the question of the neutrino magnetic moment in relation to the neutrino mass in the framework of a $SU(4) \times SU(2)_L \times SU(2)_R$ model with $N=1$ supersymmetry which is derivable from four-dimensional fermionic superstrings. The model possesses a variety of attractive features such as naturally light doublets and fermion mass relations which have been analyzed elsewhere [10]. Our main motivation is provided by the promising fact that the neutrino mass in this model is in part provided by the coupling to gauge singlet fields that have no relation to the neutrino magnetic moment, thus, allowing a degree of independence among these two parameters. In what follows, first, we briefly present the content of the model and the superpotential and, then, proceed to derive the neutrino mass matrix and compute the relevant diagrams that contribute to the overall neutrino magnetic moment.

The chiral superfield content of the model is the following

$$F(4,2,1) = Q(3,2,1/6) + l(1,2,1/2)$$

$$F^c(4,1,2) = u^c(\bar{3},1,-2/3) + d^c(\bar{3},1,1/3) + e^c(1,1,1) + N^c(1,1,0)$$

$$H(4,1,2) = \bar{u}_H^c(3,1,2/3) + \bar{d}_H^c(3,1,-1/3) + \bar{e}_H^c(1,1,-1) + \bar{N}_H^c(1,1,0)$$

$$\bar{H}(\bar{4},1,2) = u_H^c(\bar{3},1,-2/3) + d_H^c(\bar{3},1,1/3) + e_H^c(1,1,1) + N_H^c(1,1,0)$$

$$h(1,2,2) = h(1,2,-1/2) + h^c(1,2,1/2)$$

$$D(6,1,1) = D(3,1,-1/3) + D^c(3,1,1/3)$$

$$\Phi(1,1,0)$$

in terms of $SU(4) \times SU(2)_L \times SU(2)_R$ and $SU(3)_c \times SU(2)_L \times U(1)_Y$ representations. The matter representations $F + F^c$ make up just the 16-representation of $SO(10)$, while $h + D$ is the 10-representation. In contrast, the GUT-Higgses $H + \bar{H}$ are only parts of the 16+16. The superpotential of the model will involve all existing parts of the $16 \times 16 \times (10)_H + 16 \times (\bar{16})_H \times 1 + (16)_H \times (16)_H \times (10)_H + (\bar{16})_H \times (\bar{16})_H \times (\bar{10})_H + (10)_H \times (10)_H \times 1 + 1 \times 1 \times 1$ couplings.

An additional Z_2 discrete symmetry $\bar{H} \leftrightarrow -\bar{H}$ rids the model of the unwanted piece of the $16 \times \bar{16}_H \times 1$ coupling[10]. Under the given representations and the imposed discrete symmetry this is the most general cubic super-potential. In terms of the fields it reads as follows

$$\begin{aligned} W = & \lambda_1^{ij} F_i F_j^c h + \lambda_2^{ij} F_i^c H \Phi_j + \lambda_3 H H D + \lambda_4 \bar{H} \bar{H} D + \lambda_5^i h h \Phi_i \\ & + \lambda_6^{ijk} \Phi_i \Phi_j \Phi_k + \lambda_7^{ij} F_i F_j D + \lambda_8^{ij} F_i^c F_j^c D + \lambda_9^i D D \Phi_i \end{aligned} \quad (1)$$

The indices $i, j = 1, 2, 3$ are generation indices. The cubic term for the three singlets ensures that they get no vacuum expectation value larger than the supersymmetry breaking. The introduction of an extra singlet Φ_0 coupled through a term $\lambda_6^{ij0} \Phi_i \Phi_j \Phi_0$ provides a mass for the Φ_i 's from the vacuum expectation value of Φ_0 . In order to guaranty a v.e.v. for Φ_0 the cubic Φ_0^3 term should be absent. This is only technically natural unless we complicate the imposed discrete symmetry.*

(*)For example, $Z_2 \times Z_2$ symmetry with $\Phi_1, \Phi_2 \rightarrow (-1, -1)$, $\Phi_3 \rightarrow (1, -1)$ and $\Phi_0 \rightarrow (-1, 1)$ leads to $W(\Phi) = (\Phi_1 + \Phi_2) \Phi_3 \Phi_0$ with supersymmetric minimum at $\Phi_1 = \Phi_2 = \Phi_3 = 0$ and $\Phi_0 \neq 0$.

A simpler possibility, which we adopt, is the introduction of a direct quadratic mass term for the Φ_i 's, $\mu_{ij}\Phi_i\Phi_j$. In what follows we shall assume that there is a mass μ_{ij} for the singlets Φ_i .

The symmetry breaking to $SU(3)_c \times SU(2)_L \times U(1)_Y$ occurs when the neutral scalars in H and $\bar{H}$ acquire non-zero vacuum expectation values

$$\begin{aligned} \langle H \rangle &= \langle \bar{N}_H^c \rangle = M \\ \langle \bar{H} \rangle &= \langle N_H^c \rangle = \bar{M} \end{aligned} \quad (2)$$

The coloured triplets in $\bar{H}$ and H combine with the coloured triplets in D through the couplings

$$\lambda_3 HHD + \lambda_4 \bar{H}\bar{H}D \longrightarrow \lambda_3 \langle \bar{N}_H^c \rangle d_H^c D^c + \lambda_4 \langle N_H^c \rangle d_H^c D \quad (3)$$

The neutrino mass-matrix turns out to be

$$\begin{pmatrix} 0 & m_u & 0 \\ m_u & 0 & \mathcal{M} \\ 0 & \mathcal{M} & \mu \end{pmatrix} \quad (4)$$

where the entries arise from the couplings

$$\lambda_1^{ij} \langle h_0 \rangle (u_i u_j^c + v_i N_j^c) = m_u^{ij} v_i N_j^c + \dots$$

$$\lambda_2^{ij} \langle \bar{N}_H^c \rangle N_i^c \Phi_j = \lambda_2^{ij} M N_i^c \Phi_j = \mathcal{M}^{ij} N_i^c \Phi_j$$

$$\mu_{ij} \Phi_i \Phi_j$$

Being agnostic about the value of the singlet mass-parameter μ , we arrive at the neutrino mass eigenvalues

$$m_u^2 \mu / \mathcal{M}^2 \quad \mu \pm \sqrt{\mu^2 + \mathcal{M}^2}$$

the light one corresponding dominantly to v . Even if μ were to be of order M_p we get tiny neutrino masses since M is forced to be superheavy from baryon decay.

In the case that μ is superheavy, the Higgs isodoublet mixing term $h h \Phi$ generates a tiny (order m_s^2/M) Higgs mixing through the diagrams of figure 1 which is not adequate to avoid an unwanted axion state. Then, an extra source of Higgs mixing is required. This could be an extra singlet with a small ($O(m_w)$) vacuum expectation value coupled as $h h \chi$.

This would be only technically natural. In the case that μ is small ($O(m_W)$), the radiatively generated Higgs mixing $\lambda_5^2 m_S / \pi^2$ is satisfactory for large enough values of λ_5 . This again is technically natural. In both cases we can assume that either way there exists a mixing term $m(h)hh$ for the Higgsinos with $m(\tilde{h}) \approx O(m_W)$.

The neutrino mass matrix is modified by radiative corrections. The νN^c entry is modified by contributions from the diagrams shown in figure 2. The correction is roughly (up to logarithmic terms), suppressing indices on generation space,

$$m_R \approx \lambda_1^2 m_e m(\tilde{h}) / 16\pi^2 m_S$$

and assuming $m(\tilde{e}) \approx m_S$.

In this model the masses of quarks and leptons are the same at tree-level but we assume that we have taken into account the radiative corrections for fermion masses.

The $\nu\Phi$ entries which are zero at the tree-level receive radiative contributions from the diagrams of figure 3 which are roughly of order (*)

$$m'_R \approx (\lambda_5 \lambda_6 m_u' m_S / 16\pi^2 \mathcal{M}) (\ln(\mathcal{M}^2 / m_\Phi^2) - 1)$$

with m_u' we denote the value of the coupling $\nu N^c \lambda_1 \langle h_0 \rangle$.

The correction to the Φ mass may also be relevant in the case that μ is not superheavy. It turns out roughly (figure (4)),

$$\mu_R \approx (\lambda_5^3 \lambda_6 \langle h \rangle^2 m_S / 16\pi^2 \mu^2)$$

No radiative corrections up to two loops at least appear for the ΦN^c vertex.

It is also worth noticing that, in the diagrams of figures 3 and 4 which are responsible for the m'_R and μ_R contributions, no charged particles circulate.

Thus finally the one-loop corrected neutrino mass-matrix takes the form

$$\begin{pmatrix} 0 & m_u + m_R & m'_R \\ m_u + m_R & 0 & \mathcal{M} \\ m'_R & \mathcal{M} & \mu + \mu_R \end{pmatrix} \quad (5)$$

(*) unless $m \gg \mathcal{M}$ in which case instead of $1/\mathcal{M}$ we have $\mathcal{M}/\mu^2$.

Again, there are two large eigenvalues $[\mu + \mu_R \pm ((\mu + \mu_R)^2 + 4\mathcal{M}^2)^{1/2}]$ and a small neutrino mass-eigenvalue

$$(m_u + m_R)^2 (\mu + \mu_R) / \mathcal{M}^2 - (m_u + m_R) m'_R / \mathcal{M}$$

Note that in the second term m'_R is already of order $1/\mathcal{M}$ so that the second term is

$$-(\lambda_5 \lambda_6 (m_u + m_R) m_u m_S / 8\pi^2 \mathcal{M}^2) \approx -(\lambda_5 \lambda_6 m_u^2 m_S / 8\pi^2 \mathcal{M}^2)$$

and its relevance depends on the relation between μ and $\lambda_5 \lambda_6 m_S / 8\pi^2$.

A neutrino magnetic moment interaction $\bar{\nu}_R \sigma_{\mu\nu} F^{\mu\nu} \nu_L$ is also present in this model generated by the same diagrams that give rise to the neutrino mass radiative contributions with a photon insertion in any of the lines representing charged particles. The fact is that to this order only figure 1 represents a diagram with charged internal lines and therefore only effective interaction terms of the type $\bar{\nu}_R \sigma_{\mu\nu} F^{\mu\nu} N_L^c$ and $\bar{N}_R^c \sigma_{\mu\nu} F^{\mu\nu} \nu_L$ will arise. Denoting with μ the contribution to the neutrino magnetic moment matrix corresponding to the νN^c entry and with μ', μ'' , the much smaller contributions to the $N^c \Phi$ and $\nu \Phi$ entries that might arise in higher loops ($\mu', \mu'' \ll \mu$), we have

$$\begin{aligned} L_{\text{int.}} &= (\bar{\nu}_R \bar{N}_R^c \bar{\Phi}_R) \sigma_{\mu\nu} F^{\mu\nu} \begin{pmatrix} 0 & \mu & \mu'' \\ \mu & 0 & \mu' \\ \mu'' & \mu' & 0 \end{pmatrix} \begin{pmatrix} \nu_L \\ N_L^c \\ \Phi_L \end{pmatrix} \\ &\approx (\bar{\nu}_R \bar{N}_R^c \bar{\Phi}_R) \sigma_{\mu\nu} F^{\mu\nu} \begin{pmatrix} 0 & \mu & 0 \\ \mu & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} \nu_L \\ N_L^c \\ \Phi_L \end{pmatrix} \end{aligned} \quad (6)$$

A rough estimate gives ($\mu_B = e/2m_e$)

$$\mu(\nu, N^c) \approx \mu_B (\lambda_1^2 m_e m(\tilde{h}) / 16\pi^2 m_S^2) (\ln(m_S^2 / m(\tilde{h})^2) - 1)$$

where we have suppressed family indices. Taking $m(h)$ as favourable as possible, i.e., $m(\tilde{h}) \approx m_S \approx m_W$ and the Yukawa couplings λ_1 to take the value, 0.7×10^{-3} , we still get

$$\mu \approx \mu_B \times 10^{-14}$$

which is smaller by three orders of magnitude from the required value. Nevertheless, the essential point of this analysis is that the value of the neutrino magnetic moment is not directly related to the neutrino mass value.

The simplest way out of these small values is to introduce a second Higgs isodoublet $h'(1,2,2) = h'(1,2,-1/2) + h'^c(1,2,1/2)$ with identical couplings to the standard fermions i.e. $\lambda_1' \bar{F} F^c h'$ which does not get a vacuum expectation value due to a direct mass $m(h') h' h'$ that overpowers any negative mass corrections due to radiative effects. $m(h')$ will be assumed to be near the electroweak scale. Then next to the diagram of figure 2 there is an analogue diagram with the lower Higgs internal lines replaced by h' . The dominant contribution is then

$$\mu \approx \mu_B (\lambda_1'^2 m_e m(\tilde{h}') / 16\pi^2 m_S^2) (\ln(m^2/m(\tilde{h}')^2) - 1)$$

The Yukawa couplings λ_1' have nothing to do with the lepton mass-matrix and could be assigned natural values $\lambda_1' \approx 10^{-2}$. Then, for $m'(\tilde{h}) \approx m_S \approx m_W$

$$\mu \approx \mu_B \times 10^{-11}$$

which is the desired order of magnitude.

We can conclude that the essential points concerning neutrino masses and magnetic moments are independent of the particular $SU(4) \times SU(2)_L \times SU_R(2)$ model we analysed and hold for a general class of models in which the seesaw mechanism is realized with additional neutral singlet fields. Thus, it is possible in this class of models to have light neutrinos and sizeable magnetic moments. Supersymmetry does not play a decorative role since it guarantees the smallness of radiative corrections and helps to obtain larger values for magnetic moments. The need for additional Higgs isodoublets that do not get vacuum expectation value is happily met by all models derived from superstrings[11].

References

- [1] R. Davis Jr et al., Phys. Rev. Lett. 20(1968)1205; J.K. Rowley, B.T. Cleveland and R. Davis, in "Solar neutrinos and neutrino astrophysics" A.I.P. Conf. Proc. No 126 (1985)p.1.
- [2] J.N. Bahcall et al., Rev. Mod. Phys. 54(1982)767
- [3] Kamiokande collaboration at the XXIV Int.Conf. on High Energy Physics, Munich, August 4-10 (1988).
- [4] M.I. Vysotsky and M.B. Voloshin, Yad. Fiz. 44(1986)43 (Sov.J. of Nucl. Phys. 44(1986)544); M.B. Voloshin, M.I. Vysotsky and B. Okun, Zh. Eksp. Teor. Fiz 91(1986) 745 (Sov. Phys. JETP 64(1986)446); A. Cisneros, Astrophys. Space. Sc. 10 (1971) 87.
- [5] P. Sutherland et al., Phys. Rev. D13(76) 2700; F. Reines, H. Gurr and H. Sobel, Phys. Rev. Lett. 37(1976)315.
- [6] S.T. Petcov, Yad. Fiz. 25, (1977) 641; B.W. Lee and R.E. Shrock, Phys.Rev. D16(77) 1444; K. Fujikawa and R.E. Schrock, Phys. Rev. Lett. 45 (1980) 963.
- [7] S.T. Petcov, Phys. Lett. 1158 (1982) 401; G.K. Leontaris, K. Tamvakis and J.D. Vergados, Phys.Lett. 162B(1985)153; G.K. Leontaris and K. Tamvakis, Phys. Lett. 191B (1987)421; M. Fukugita and T. Yanagida, Phys. Rev. Lett 58(1987)1807; G.K. Leontaris and C.E. Vayonakis, Phys. Lett. 206B (1988)271.
- [8] R. Barbieri and G. Fiorentini, Nucl. Phys. B304(1988)909; J. Liu, Un. of Michigan Preprint UM-TH-88-19 (1988); K.S. Babu and V.S. Mathur, Phys. Lett. 196B(1987)218; J.A. Grifols and S. Peris, Phys.Lett. 213B(1988)482; J. Pulido and R. Ralston, Phys. Rev. D38(1988)2864; M.J. Duncan, J.A. Grifols, A. Mendez and S.U. Sarkar, Phys. Lett. 191B (1987) 304. J. Maalampi, J.T. Peltoniemi and M. Roos, Univ. of Helsinki preprint KU-TFT 88-51(1988); J. Pulido, CFMC (Lisbon) preprint IFM-18/88 (1988); M.R. Valoshin, ITEP preprint 87-215(1987).
- [9] K. Enqvist, J. Maalampi and A. Masiero, Univ. of Helsinki preprint HU-TFT-89-1 (1989).
- [10] I. Antoniadis and G.K. Leontaris, Phys. Lett. 216B(1989)333.
- [11] I. Antoniadis, J. Ellis, J. Hagelin and D.V. Nanopoulos, Phys. Lett. 194B(1987)23; 208B(1988)207.

Figure Captions

Fig. 1. Higgs mixing.

Fig. 2. νN^c contribution to neutrino mass matrix and magnetic moment.

Fig. 3. $\nu \Phi$ contribution to neutrino mass matrix and magnetic moment.

Fig. 4. $\Phi \Phi$ radiative corrections.

FIGURE 1

FIGURE 2

FIGURE 3

FIGURE 4