EUROPEAN ORGANIZATION FOR NUCLEAR RESEARCH

Z032

BRUMMER Pete

SPS

01

CERN LIBRARIES, GENEVA

CERN/LEP-TH/89-34

CM-P00064396

Synchrotron-Radiation Limit of the Luminosity in TeV Linear Colliders

K. Hirata*, K. Oide *[†] and B. Zotter

Geneva, Switzerland

April 1989

Abstract

In linear colliders in the TeV region, the synchrotron radiation occurring in the final quadrupole magnet limits the smallest vertical r.m.s. beam size which can be reached at the interaction point. Due to the same effect, however, the distribution does not remain Gaussian, which helps the luminosity: it is much larger than what is expected from the r.m.s. beam size, but much smaller than the value without radiation in the quadrupole. The luminosity is still limited from above, but the maximum occurs at a much smaller value of the nominal beam size. A simple approximate expression for the luminosity is also proposed.

Submitted to Phys.Lett.B

*On leave of absence from KEK, National Laboratory for High Energy Physics, Tsukuba, Ibaraki 305, Japan.

[†]Stanford Linear Accelerator Center, Stanford University, Stanford, California 94309, U.S.A

1 Introduction

Recently, one of the authors[1] found that synchrotron radiation in the final quadrupole magnet limits the smallest possible r.m.s. beam size which can be reached at the interaction point (IP) of a linear collider. Its value depends essentially only on the normalized emittance. This fact seems to discourage efforts to achieve lower values of the beta-function at the IP (β^*). The present paper shows that such efforts may still be useful. For the final focus design, it is only important that the distribution at the IP be optimized such that the luminosity L becomes a maximum. It should be noted that this distribution may be very different from a Gaussian due to the strong effect of the radiation, even if it was close to it before entering the quadrupole magnet. For this reason we can not apply the usual formula for the luminosity L_G , which assumes a Gaussian distribution:

$$L \neq L_G \propto 1/\sigma, \quad (1.1)$$

where σ is the r.m.s. vertical beam size.

2 The Problem

As shown in Ref.[1], the perturbation Y of the vertical coordinate y at the IP due to the final quadrupole magnet can be written as $y = y_0 + Y$, where the stochastic part Y is approximated by

$$\begin{aligned} Y &= K y'_0 \int_0^L ds \frac{u(s)}{E_0} G(s) \\ G(s) &= \int_0^s ds' g^2(s'). \end{aligned} \quad (2.1)$$

Here y_0 and y'_0 are the nominal values (without radiation) of the vertical coordinates y and y' at the IP, K the (constant) focusing gradient of the lens, L its length, u a stochastic variable representing the energy of the emitted photon, E_0 the nominal energy of the particle, and

$$g(s) = \frac{\sin \sqrt{K} s}{\sqrt{K}} + l^* \cos \sqrt{K} s, \quad (2.2)$$

where l^* is the distance between the face of the magnet and the IP.

The distributions in y_0 and y'_0 are Gaussian with standard deviations

$$\sigma_0 = \sqrt{\varepsilon \beta^*}, \quad \sigma'_0 = \sqrt{\varepsilon / \beta^*}, \quad (2.3)$$

respectively. Here ε is the nominal emittance of the beam. In this connection, it is useful to define the effective beam size σ_{eff} by

$$\sigma_{eff} = \sigma_0 \frac{L_0}{L} = \sigma \frac{L_G}{L}, \quad (2.4)$$

where L_0 is the nominal luminosity without radiation in the last quadrupole. If we replace σ by σ_{eff} , we can calculate the real luminosity L with the usual formula for the Gaussian distribution.

In the approximation of Eq.(2.1)¹ Y is independent of y_0 . It is thus convenient to introduce the characteristic functions

$$\tilde{\rho}(k) = \langle e^{ik y} \rangle, \quad \tilde{\rho}_0(k) = \langle e^{ik y_0} \rangle, \quad \text{and} \quad \tilde{\rho}_Y(k) = \langle e^{ik Y} \rangle, \quad (2.5)$$

where $\langle \rangle$'s are averages over y , y_0 and Y , respectively. That is, they are Fourier transforms of the corresponding distribution functions. It can be shown that

$$\tilde{\rho}(k) = \tilde{\rho}_0(k) \tilde{\rho}_Y(k). \quad (2.6)$$

Since the luminosity satisfies

$$L \propto \int \tilde{\rho}(k) \tilde{\rho}(-k) dk, \quad (2.7)$$

we can express it as

$$L = L_0 \frac{\int dk \tilde{\rho}(k) \tilde{\rho}(-k)}{\int dk \tilde{\rho}_0(k) \tilde{\rho}_0(-k)} = L_0 D, \quad (2.8)$$

where D is the "luminosity enhancement factor": $D = L/L_0 = \sigma_0/\sigma_{eff}$. Since $\tilde{\rho}(-k) = \tilde{\rho}(k)^*$, we get

$$D = \frac{2\sigma_0}{\sqrt{\pi}} \int_0^\infty e^{-\sigma_0^2 k^2} |\tilde{\rho}_Y(k)|^2 dk. \quad (2.9)$$

Because $|\tilde{\rho}_Y(k)| < 1$, this formula clearly shows that, $D < 1$, i.e., *the luminosity is always reduced by radiation in the last quadrupole*: $L < L_0$. It will be shown later that furthermore $L > L_G$, so that

$$L_G < L < L_0. \quad (2.10)$$

3 Evaluation of the Luminosity

The main problem lies in evaluation of $|\tilde{\rho}_Y(k)|$. Note that Y in Eq.(2.1) is stochastic in several different ways. We will use the following notation for averaging: (i) $[]$: The first average is taken over all possible ways of the emission of photons. The position of the photon emission obeys the Poisson law, while the energy of the emitted photons obeys the classical radiation law[2]. (ii) $\langle \rangle$: The second average is taken over the angular distribution in y'_0 . It is implied that the first average $[]$ is already taken.

Following the derivation of the Campbell-Rice law[3], we may write²

$$[e^{ikY}] = \exp \int_0^L ds N(s) \{S(l) - 1\}. \quad (3.1)$$

¹Actually, Y depends weakly on y_0 . In Eq.(2.1), this has been neglected, which is justified when $\beta^* \ll l^*$.

²Note that Y , in Eq.(2.1), can be put into the form $Y = \int ds U(s)$. Let us divide the quadrupole magnet into many thin slices in such a way that the expectation value for the number of emitted photons $N(s)ds$ is much less than unity in each slice. In each slice, we

$$N(s) = \frac{5\alpha\gamma}{2\sqrt{3}} |Kg(s)y'_0|, \quad (3.2)$$

$$S(P) = \frac{3}{10} \frac{1}{P} \frac{1}{\sqrt{1+P^2}} \{(\sqrt{1+P^2} + P)^{5/3} - (\sqrt{1+P^2} - P)^{5/3}\}, \quad (3.3)$$

$$P = ky_0'^2 K^2 G(s) |g(s)| \frac{3}{2} \lambda \gamma^2. \quad (3.4)$$

Here, $N(s)$ is the expectation value of the number of emitted photons per unit length, α the fine structure constant, γ the relativistic Lorentz factor, $S(P)$ the Fourier transform[4] of the classical photon spectrum, and λ the Compton wave length of the electron.

The characteristic function $\tilde{\rho}_Y(k)$ can be obtained from

$$\tilde{\rho}_Y(k) = \langle [e^{ikY}] \rangle = \frac{1}{\sqrt{2\pi\sigma_0'}} \int dy'_0 \exp(-\frac{y_0'^2}{2\sigma_0'}) [e^{ikY}]. \quad (3.5)$$

This, with Eq.(2.6), gives us the complete description of $\tilde{\rho}(k)$. By expanding $\log \tilde{\rho}(k)$ into a Taylor series in k (cumulant expansion[3,5]), we obtain the "cumulants" C_n with respect to y . The second cumulant is the r.m.s. beam size,

$$C_2 \equiv \sigma^2 = \sigma_0^2 + A_1^2 F_1^2 \Sigma_6 + A_2 F_2 \Sigma_5. \quad (3.6)$$

The coefficients are defined[5] as

$$\Sigma_n \equiv \langle |y'_0|^n \rangle, \quad (3.7)$$

$$A_n = \frac{\sqrt{3}}{2\pi} \alpha \gamma \left(\frac{3}{2} \lambda \gamma^2\right)^n \int_0^\infty \xi^n d\xi \int_{\xi}^\infty K_{5/3}(\bar{\xi}) d\bar{\xi}. \quad (3.8)$$

The form factors $F_n = F_n(\sqrt{K}L, \sqrt{K}l^*)$ are defined by

$$F_n(x, y) = \int_0^x d\varphi |\sin \varphi + y \cos \varphi|^{n+1} \left[\int_0^\varphi (\sin \varphi' + y \cos \varphi')^2 d\varphi' \right]^n. \quad (3.9)$$

The function F in Ref.[1] is F_2 here. The second term of σ^2 was missing in Ref.[1]. It is negligible for the usual values of β^* , but becomes dominant for very small values of it.

The higher cumulants are useful parameters of the deviation from a Gaussian. When y is Gaussian, all C_n for $n > 2$ vanish. After some algebra, we get

$$\begin{aligned} C_4 = & A_1^4 F_1^4 (\Sigma_{12} - 3\Sigma_6^2) + 6A_1^2 F_1^2 A_2 F_2 (\Sigma_{11} - \Sigma_6 \Sigma_5) + 3A_2^2 F_2^2 (\Sigma_{10} - \Sigma_5^2) \\ & + 4A_1 F_1 A_3 F_3 \Sigma_{10} + A_4 F_4 \Sigma_9. \end{aligned} \quad (3.10)$$

In Table 1, we show the normalized 4-th cumulant and σ as functions of β^* , the parameters can ignore cases where more than two photons are emitted. We thus have

$$\begin{aligned} [e^{ikY}] &= \prod_i [e^{ikU_i}] \\ &= \prod_i (1 - N_i + N_i [e^{ikU_i}]_1) \\ &= \exp\left(\sum_i N_i \{[e^{ikU_i}]_1 - 1\}\right) \\ &= \exp\left(\int_0^L ds N(s) \{[e^{ikU(s)}]_1 - 1\}\right). \end{aligned}$$

Here, U_i and N_i are $U(s)ds$ and $N(s)ds$ at the i -th slice, respectively and $[]_1$ stands for the average over one photon process.

β^*	C_4/σ^4	N_{photon}	$\sigma_0(\text{nm})$	$\sigma(\text{nm})$	$\sigma_{eff}(\text{nm})$	$\sigma_{eff}(\text{nm})$
(μm)	exact	exact	exact	exact	Num	App
100	0.77	0.22	1.58	1.60	1.59	1.58
80	3.17	0.25	1.41	1.46	1.43	1.42
60	17.4	0.29	1.22	1.33	1.25	1.23
40	117	0.35	1.00	1.33	1.05	1.03
20	371	0.46	0.71	2.25	0.81	0.83
10	330	0.71	0.5	5.31	0.69	0.82
5	249	1.00	0.35	13.2	0.65	0.92
1	135	2.24	0.16	118	0.91	1.67
0_+	43.2	∞	0	∞	?	∞

Table 1: Beam size and related quantities which are exact, numerical (Num) and approximation (App) results.

being the same as those in Ref.[1]: $\varepsilon = 2.5 \times 10^{-14}m$, $\gamma = 10^6$, $L = 0.4m$, $l^* = 0.4m$ and $K = 8.5417m^{-2}$. It can be seen that the beam is far from a Gaussian when β^* is small. This is not in contradiction to the fact that the total number of photons, N_{photon} , becomes large for extremely small β^* as shown in the table. Since in that case σ and C_4 are dominated by the second term of Eq.(3.6) and the first term of Eq.(3.10), respectively, the ratio of C_4/σ^4 does not tend to zero, and hence the central limit theorem does not apply.

We could continue to calculate higher cumulants, however the expansion does not appear to be useful in describing $\tilde{\rho}_Y(k)$ since it converges badly for large value of k . The convergence radius is very small or even zero[5]. This could be related to the fact that $S(P)$ has branch points at $P = \pm i$.

To obtain $\tilde{\rho}_Y(k)$, and then D and σ_{eff} , we should average over y'_0 and k . Results of the numerical integration are shown in Fig.1, as well as in Table 1. They are in good agreement with two other independent simulations: real space multiparticle simulation[6] and frequency domain simulation[5]. In the former case, a large number of particle must be used to avoid excessive numerical fluctuations when β^* is small. In addition, the result for the luminosity is quite sensitive to the mesh size in y space.

In Fig.1, it is clearly shown that the r.m.s. beam size σ is not a good parameter to estimate the luminosity for small β^* . The reason is clear: due to the strong effect of radiation in the final quadrupole, the distribution becomes very different from a Gaussian. The beam is thinner than the Gaussian with the same r.m.s. width. The effective beam size, σ_{eff} , is still limited from below, but the minimum occurs at a much smaller value of β^* than that of σ , and the smallest value is only about half of the latter.

4 Approximation of the Luminosity

It seems quite difficult to obtain σ_{eff} analytically. Let us introduce an approximation. In P , we replace terms dependent on k , y'_0 and s by their averages with respect to the

Figure 1: Beam size as a function of β^* . Without radiation (dotted line), Gaussian approximation (full line), present approximation (dashed line), and numerical results (crosses).

integrations in Eqs.(2.9), (3.5) and (3.1). Thus,

$$P \rightarrow \bar{P} = \frac{1}{\sqrt{2}\sigma_0} \sigma_0'^2 \frac{3}{2} \lambda \gamma^2 K^2 < G(s)|g(s)| >, \quad (4.1)$$

where $< >$ is the average over s . It is then easy to perform integrations required:

$$D = \{w(iX)\}^2 \quad (4.2)$$

$$X = \frac{A_0 F_0}{\sqrt{2}} \sigma_0' \{1 - S(\bar{P})\}, \quad (4.3)$$

and where w is the complex error function. The approximate values for σ_{eff} thus obtained are also shown in Table 1 and Fig.1. The agreement is good enough for such a simple approximation. In the limit $\beta^* \rightarrow 0$, the model predicts that $D \rightarrow 0$ and $\sigma_{eff} \rightarrow \infty$.

For another set of parameters³, it is confirmed that the agreement is equally satisfactory as in the present case.

5 Summary

We have not considered the "disruption" of the beam due to collisions with the other beam at the IP[7]. Usually one assumes that the incoming beam is Gaussian in the transverse direction. However, the effect of the non-Gaussian distribution of the incoming beam on luminosity may be quite important. The enhancement factor should thus be corrected also for this effect, and the parameters should be optimized considering both the synchrotron radiation in the final quadrupole and the disruption due to the non-Gaussian distribution in the other beam.

A more detailed and extended paper will be published elsewhere.

Two of the authors (K.H and K.O) thank the members of the LEP theory group of CERN and the theory group of SLAC, respectively, for the hospitality extended to them.

References

- [1] K. Oide, Phys.Rev.Lett.**61** (1988) 1713.
- [2] For example, M. Sands, *The Physics of Electron Storage Rings*, SLAC-121 (1970).
- [3] S. O. Rice, Bell System Technical Journal **23** (1944) 282.
- [4] A. Erdélyi, ed., *Tables of integral transformations*, Vol.I, (Mcgraw-Hill, New York, 1954).
- [5] K.Hirata, *SOIL: Simulation of Oide Limit*, CERN/LEP-TH/89-03 (1989).
- [6] K.Oide 'FFSAD' (1988), unpublished. K.Hirata, in :Proceedings of the 2-nd Advanced ICFA Beam Dynamics Workshop, J.Hagel and E.Keil, eds. CERN 88-04 (1988), p.62.
- [7] R. Hollebeek, Nucl.Instrum.Meth.**184** (1981) 333.

³A possible set of parameters for a CERN linear collider: $\varepsilon = 5 \times 10^{-13}m$, $\gamma = 2 \times 10^6$, $L = 1.3m$, $l^* = 1.1878m$ and $K = 0.82m^{-2}$.