NULL SPINNING STRINGS

J. Gamboa[†]

*Instituto de Física, Univ. de São Paulo
CP 20516, São Paulo, CEP 01498, Brasil*

C. Ramírez⁺

*Inst. für Theor. Physik, Univ. Heidelberg
Philosophenweg 16, D 6900 Heidelberg, W. Germany*

M. Ruiz-Altba^{*}

*CERN, Theory Division
CH 1211 Genève 23*

ABSTRACT

We quantize in detail the closed string with vanishing tension, null closed string. For Weyl ordering of quantum operators, the spectrum is continuous and there is no critical dimension. For normal ordering, however, the usual critical dimension holds and the spectrum is **finite**: the field content is the same as in the field theory (infinite tension) limit of usual (closed) strings. The content of bosonic null strings is massless gravity. We carry through the local supersymmetrization of this system and find similar conclusions: normal ordered null spinning strings contain in their spectrum only the massless supergravity multiplet. However normal ordered bosonic null strings do not have a consistent quantum interpretation, whereas the null spinning strings do. It points to the likely origin of some spontaneous symmetry breakdown which generates a non-zero tension, and an infinite tower of massive states. We also consider the Hamiltonian formalism for quantum null (super)strings.

[†] 47551::JGamboa (hepnet); on leave from Centro de Estudios Científicos de Santiago, Casilla 16443, Santiago 9, Chile.

⁺ Q79 at dhdurzi.bitnet

^{*} RuizAltb at cernvm.cern.ch

I. Introduction

An approach to a better understanding of the beauty and subtleties of string theory is to simplify it by studying its zero tension limit [1, 2, 3, 4, 5, 6, 7] (see also [8, 9]) in complete analogy to massive and massless (super)particles [10, 11, 12].

The bosonic null string is a one dimensional extended object with zero tension, introduced and studied by Schild [1] several years ago. Classically, it describes a system of free massless point particles subject to kinematical constraints, that is their potential energy vanishes with the consequence that the system is unstable. In the proper time gauge, the kinematical constraints mean that the velocities of the point particles are perpendicular to the string and that they lie on a $D-2$ dimensional sphere; moreover there is a rather stringy constraint which sets the momentum flow along the string equal to zero. This last constraint remains unsolved also in the light cone gauge and it actually corresponds to the equality of the masses of the left and right sectors of the usual closed strings; it will play an important role in the quantization of null strings.

From the point of view of usual strings, the zero tension limit could be interpreted as corresponding to the very high energy region where the mass gap between two string modes is tiny and in the limit leads to either a continuous or massless spectrum. Null strings have the peculiarity that, although their action can be written in a reparametrization and Weyl invariant way, their physical solutions do not display the full two dimensional conformal invariance. In fact, this is a reflection of the singularity of the geometry involved.

Quantized strings describe consistent physical theories only at a certain critical dimension [13]. On the other hand, for point particles no such critical dimensions arise. For usual quantum spinning strings with non zero tension [14, 15], the situation is similar to that of bosonic strings: their consistency implies also a critical dimension. Moreover, they enjoy the non trivial feature that their spectrum is supersymmetric only at the critical dimension [16]. In contrast, for space time supersymmetric strings [17] the situation is in some sense opposite: supersymmetry lacks a geometric interpretation and the covariant quantization is difficult as first and second class constraints are inherently mixed, but the explicit supersymmetry of its spectrum makes it most suitable for an immediate physical interpretation.

The quantization of null bosonic strings was carried out by Lizzi *et al.* [2], with the result that there are no critical dimensions for the null string, that the mass spectrum is continuous and that the corresponding wave functions are plane waves. Thus their results were consistent with the view of the null strings as a system of free massless point particles. However it seems that they missed the fact that the momentum along the string

does not vanish for plane waves with the consequence that Lorentz covariance is broken. From this approach, it is not clear why there are no critical dimensions because, as long as consistency of the Hilbert space of the physical states is not involved, it depends only on the ordering prescription which can lead to very different results, as shown in [5]. More recently, Marquard and Scholl [18] have obtained the critical dimension considering the null tension limit perturbatively. The interest of their approach lies on the fact that it can be easily generalized for higher dimensional extended objects, allowing in this way for the computation of their critical dimensions.

A related issue concerns the quantum inconsistency [19, 20] of supermembranes [21]. It turns out that, at the classical level, the potential for a tensionful membrane has flat directions and hence, a spherical membrane will naturally stretch itself into a string where no energy is paid for an increase in length, that is it will tend to a null string. De Wit *et al.* have shown that quantum corrections to the classical potential destroy the flat directions and thus a quantum bosonic membrane has a discrete spectrum and will not degenerate into a null string. However, quantum supersymmetric membranes retain the flat directions and hence their spectrum is continuous - hence supermembranes are unstable: they degenerate into null supersymmetric strings. Let us advance that the spectrum for Weyl ordered null superstrings is also continuous, in agreement with the possible interpolation of null strings as degenerate (and favoured) configurations of quantum (super)membranes. Nevertheless, these null strings enjoy no critical dimension. Another application of null strings are the dust strings considered as classical sources of gravity [22].

In this article we consider the issue of the consistent quantization of null string theory. For bosonic null strings it turns out that they can be consistently quantized with Weyl operator ordering. The result is a continuous mass spectrum and no critical dimension, *i.e.*, consistent with Lizzi *et al.* although our wave functions have a more complicated structure essentially given by Bessel functions. Furthermore, we tackle the question of the local supersymmetrization of null strings in the lagrangian formalism which, as far as we know, has never been discussed in the literature. It turns up that the naive supersymmetrization works only after some nontrivial reparametrization and that, up to kinematical constraints, it is equivalent to a system of $N=2$ free massless spinning particles [23]. In this case quantization gives similar results: with Weyl ordering, the mass spectrum remains continuous, and there are no critical dimensions. With the normal order quantization is consistent only at the critical dimension $D=10$ and, of course after GSO projection, the spectrum is the massless spectrum of type II superstrings, *i.e.*, $N=2$ supergravity. However in this case the quantum Hilbert space is well defined, *i.e.*, it has only states with positive norm, thus reflecting the old well known fact that only supersymmetric strings can be consistently

quantized. Finally, we consider all these questions from the point of view of the Hamiltonian formalism and BRST quantization which confirm the preceding results. We show also that there is a generalization of the formalism from which we can obtain chiral bosons as well as null strings.

We shall review the Lagrangian approach for the well known bosonic null strings [1, 24, 6] in Section II. We shall then fix the light cone gauge, canonically quantize, and find the physical spectrum for the Weyl and normal ordered quantum null strings. (Sections III and IV), commenting on the origin of the differences with the authors of Ref. [2]. In Section V we establish the Lagrangian formulation of the null spinning string. In Section VI we discuss its light cone quantization and consider the issue of spacetime supersymmetry. We briefly consider interactions *à la* Mandelstam in Section VII, and then study in Section VIII the null string constraints from an algebraic point of view. General considerations allow us to relate null strings to chiral bosons. We supersymmetrize the null string in the Hamiltonian approach, and discover the null spinning string constraint algebra [4]. Section IX details the BRST [25] quantization of the null spinning string; we have minimized the overlap of this Section with our work in Ref. [5]. Finally, the conclusions yield some insight into the likely equivalence between our null spinning string and the recently formulated null superstring [7], as well as some comments on other approaches in the literature where the tensionless limit of strings seems to play a role. Appendix A reviews the basics of the superspace formulation for the spinning string [26]. In Appendix B we have collected some results concerning the null string wave functions constructed in Section III.

II. The Bosonic Null String

The formal problem of writing down an action for the tensionless string can be overcome exactly as in the massless limit of the free particle action. Suppose we have a generic action

$$S = T \int L(x) dx$$

where T is a parameter of non vanishing dimension. This action is equivalent to

$$S' = \frac{1}{2} \int \left(\frac{L^2}{z} + z T^2 \right) dx$$

where z is an auxiliary field, whose equation of motion inserted back into S' yields S . The limit $T \rightarrow 0$ of S is thus given by

$$S_0 = \frac{1}{2} \int \frac{L^2}{z} dx$$

The action for the null string follows from using for L the Nambu action [1, 3, 27, 28]:

$$S_0 = -\frac{1}{2} \int \frac{g \mu^2 \sigma}{z} \quad (2.1)$$

where $g = \det g_{mn}$, $g_{mn} = \partial_m X^\mu \partial_n X_\mu$, with $m, n = 0, 1, \mu = 0, 1, \dots, D-1$. In order to maintain conformal invariance, z must transform as an invariant density under world-sheet reparametrizations and its Weyl weight must be zero. The equation of motion of z gives $g = 0$, i.e., the worldsheet is a null surface from which actually came the name of null strings [1]. Computing c from S_0 the momentum conjugate to X^μ , we get:

$$P^\mu = \frac{1}{z} \left\{ X'^2 \dot{X}^\mu - \dot{X} \cdot X' \right\} \quad (2.2)$$

which satisfies the constraints:

$$\begin{aligned} P^\mu \cdot X' &= 0 \\ P^2 &= \frac{X'^2}{g} = 0 \end{aligned} \quad (2.3)$$

The dynamics of the system described by this action can be studied after the gauge fixing [24, 6]

$$g_{mn} = \begin{pmatrix} -\dot{X}^2 & 0 \\ 0 & X'^2 \end{pmatrix}$$

$$S_0 = \frac{1}{2} \int d\tau d\sigma \frac{1}{z} \dot{X}^2 X'^2$$

so that

$$\dot{X} \cdot X' = 0$$

and

The equations of motion are then

$$\begin{aligned}\dot{X}^2 \cdot X'^2 &= 0 \\ \partial_\tau \left(\frac{1}{2} X'^2 \dot{X}^\mu \right) + \partial_\sigma \left(\frac{1}{2} \dot{X}^2 X'^\mu \right) &= 0\end{aligned}$$

These equations have non-trivial solutions only if we take

$$\dot{X}(\sigma)^2 = 0$$

Observe that the singular geometry of the world-sheet does not allow the full conformal symmetry of the string, but only a subgroup of it.

We may further fix the gauge $z = X'^2$, so that the momentum is $P^\mu = \dot{X}^\mu$, and the action is simply

$$S_0 = \frac{1}{2} \int \dot{X}^2 \quad (2.4)$$

The fields $X^\mu(\tau, \sigma)$ satisfy then the equation of motion

$$\ddot{X}_\mu(\sigma) = 0 \quad (2.5)$$

subject to the first-class constraints

$$\dot{X}^2 = \dot{X} \cdot X' = 0 \quad (2.6)$$

The gauge just derived is the closest analogue to the conformal gauge for usual strings, and we shall call it Schild's (or null) gauge. The metric takes, then, the canonical null form $\begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$ instead of the canonical conformal $\begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$. The remaining gauge degrees of freedom are subject to $\xi^0 = 0$, where ξ^m are the infinitesimal reparametrizations of the world-sheet. This can be used, together with the constraints, to fix the light-cone gauge. We shall do that in the next section and subsequently quantize.

Physically, $X^\mu(\tau, \sigma)$ describes a set of free massless point-particles (one for every point σ along the string) whose velocity is perpendicular to the string [1, 2, 24]. The crucial differences with respect to the usual string can be traced to the difference in the constraints (2.6), which for the tensionful string read

$$\dot{X}^2 + T^2 X'^2 = \dot{X} \cdot X' = 0 \quad (2.7)$$

Having set $T = 0$, we have lost the ("nearest-neighbour") interaction—that is why the null string is made out of free particles instead of coupled harmonic oscillators. These particles, nevertheless, do not form plane waves: they still satisfy the inherently stringy $\dot{X} \cdot X' = 0$ constraint. Note that there are no boundary conditions for the open null string in Schild's

gauge [6]: no momentum flows along σ at any point, so in particular no momentum flows out at the end-points either. This fact has a consequence that for open strings functions of σ cannot be expanded in Fourier series so that their boundary values are well-defined, this is a problem particularly when we consider constraints because setting them to zero will not be equivalent to set to zero their Fourier expansion. The closed null string satisfies, of course, the customary periodicity. Therefore, unless explicitly stated otherwise, we shall deal throughout this article with closed null strings.

III. Light-Cone Quantization of the Bosonic Null String

The safest way to study the physical content of a gauge theory is to avoid the subtleties of covariant quantization by considering the theory with all the gauge degrees of freedom fixed. In general, this procedure is not manifestly covariant under the original global symmetries, which are now non linearly realized. The requirement that there be a realization of these symmetries in the quantized theory provides us with a consistency test for the theory. For usual strings with non zero tension, this amounts to requiring Lorentz covariance in the light cone gauge, which proves to hold only at the critical dimension. It also restricts the mass spectrum of the theory. From the previous Section, we write

$$X^\mu(\tau, \sigma) = P^\mu(\sigma)\tau + Y^\mu(\sigma) \quad (3.1)$$

where $P^\mu(\sigma)$ and $Y^\mu(\sigma)$ are subject to the constraints

$$P^2(\sigma) = P(\sigma) \cdot Y'(\sigma) = 0 \quad (3.2)$$

The light cone gauge is defined by the condition $X^+ = \frac{1}{\sqrt{2\pi}} p^+ \tau$, with p^+ a constant. The constraints can be solved in this gauge as

$$\begin{aligned} P^-(\sigma) &= \frac{\sqrt{2\pi}}{2p^+} \tilde{P}^2(\sigma) \\ Y'^- &= \frac{\sqrt{2\pi}}{p^+} \tilde{P}^2 \cdot \tilde{Y}'^i \end{aligned} \quad (3.3)$$

where the vector products on the right-hand side include only the transverse components p^I and Y'^I , $I = 1, \dots, d = D - 2$.

It is convenient to Fourier expand the independent (transverse) degrees of freedom. Periodicity of the bosonic co-ordinates in $\sigma \in [0, 2\pi]$ implies, rather trivially, that

$$\begin{aligned} P^\mu(\sigma) &= \frac{1}{\sqrt{2\pi}} \sum_{m \in \mathbb{Z}} p_m^\mu e^{im\sigma} \\ Y'^\mu(\sigma) &= \frac{1}{\sqrt{2\pi}} \sum_{m \in \mathbb{Z}} y_m^\mu e^{im\sigma} \end{aligned} \quad (3.4)$$

The zero mode of the second constraint in (3.2) is in the light cone gauge given by:

$$P_\sigma = \frac{1}{2\pi} \int_{-\pi}^{\pi} \tilde{P}^2 \cdot \tilde{Y}'^i d\sigma = i \sum_{m \in \mathbb{Z}} m \tilde{p}_{-m} \cdot \tilde{y}_m = 0 \quad (3.5)$$

that is, it remains unsolved and can be interpreted classically as the requirement that the momentum flowing along the string, in the σ direction, vanish. It reflects the symmetry of

the null string under rigid σ rotations. The condition $P_\sigma = 0$ is the (closed!) null-string analogue of the usual string constraint $L_0 = \bar{L}_0 = 0$.

The mass M of the string is given by the expression [2]

$$M^2 = - \left[\int d\sigma P(\sigma) \right]^2 = \sum_{n \neq 0} p_{-n}^\mu p_n^\mu \geq 0 \quad (3.6)$$

where we have used the constraint $P^2(\sigma) = 0$ and also the fact that, in the light-cone gauge, the only mode of P^+ which does not vanish is the zero mode (p^+).

As usual, the quantization prescription is

$$\begin{aligned} [y_0^-, p^+] &= -i \\ [Y'^I(\sigma), P'^J(\sigma')] &= i\delta^{IJ}\delta(\sigma - \sigma') \end{aligned} \quad (3.7)$$

In modes, the second commutator reads as

$$[y_n^I, p_m^J] = i\delta^{IJ}\delta_{n+m,0} \quad (3.8)$$

After quantization, the condition $P_\sigma = 0$ will be replaced by the requirement that physical states be eigenstates of P_σ with some eigenvalue, fixed by the definition of operator ordering. Thus we have to solve the eigenvalue problem of the mass operator and P_σ . Expand P_σ and M^2 as follows:

$$\begin{aligned} P_\sigma &= i \sum_{n>0} n (\tilde{p}_{-n} \cdot \tilde{y}_n - \tilde{p}_n \cdot \tilde{y}_{-n}) = \sum_{n>0} n S_n \\ M^2 &= 2 \sum_{n>0} \tilde{p}_{-n} \cdot \tilde{p}_n = 2 \sum_{n>0} T_n \end{aligned} \quad (3.9)$$

It turns out that the commutators $[M^2, P_\sigma]$, $[M^2, S_n]$, $[M^2, T_n]$, $[P_\sigma, S_n]$, $[P_\sigma, T_n]$, $[S_n, T_n]$, $[S_n, S_m]$, and $[T_n, T_m]$ vanish. Moreover,

$$\begin{aligned} [P_\sigma, y_m^I] &= m [S_m, y_m^I] = m p_m^I \\ [P_\sigma, p_m^I] &= m [S_m, p_m^I] = m p_m^I \\ [M^2, y_m^I] &= [T_m, y_m^I] = -2i p_m^I \\ [M^2, p_m^I] &= [T_m, p_m^I] = 0 \end{aligned} \quad (3.10)$$

where the commutators of M^2 hold for $m \neq 0$. Therefore if we take the mass operator plus the zero modes, i.e., $2p_0^-$, it will give us the Hamilton operator [2]. Thus, the problem of finding mass eigenstates which are also eigenstates of P_σ can be reduced to that of simultaneously diagonalizing S_n and T_n :

$$S_n \Psi_n = s_n \Psi_n \quad \text{and} \quad T_n \Psi_n = t_n \Psi_n \quad (3.11)$$

The wave function of a generic state will be written as a product of wave functions, a factor for each $n \geq 0$:

$$\Psi = \prod_{n=0}^{\infty} \Psi_n \quad (3.12)$$

The zero modes p_0^I and y_0^I are not contained in P_n and M^2 . To them corresponds a plane wave describing the motion of the string's centre of mass,

$$\Psi_{n=0} = e^{i\vec{k}_0 \cdot \vec{y}_0} \quad (3.13)$$

In what follows, we shall discuss the wave functions Ψ_n ($n > 0$). We will first quantize with the Weyl order which essentially amounts to realize the commutation relations in the usual way, i.e., representing the momentum by the derivative with respect to the co-ordinate:

$$\begin{aligned} y_n^I &= v_n^I, & y_{-n}^I &= (v_n^I)^*, & (n > 0) \\ p_{-n}^I &= -i \frac{\partial}{\partial v_n^I}, & p_n^I &= -i \frac{\partial}{\partial (v_n^I)^*}, & (n > 0) \end{aligned} \quad (3.14)$$

where $\bar{v}_n \in \mathbb{C}^d$ and $\bar{v}_n^*$ is its complex conjugate. For conciseness, we drop the level label n and restore it at the end, when we assemble results.

The wave functions depend on v^I and v_I^* . The scalar product of two wave functions Φ and Ψ

$$(\Phi, \Psi) = \int d^d v d^d v^* \Phi^*(\bar{v}, \bar{v}^*) \Psi(\bar{v}, \bar{v}^*) \quad (3.15)$$

is such that the operators

$$\begin{aligned} S &= v^I \partial_{v^I} - v_I^* \partial_{v_I^*} \\ T &= -\partial_{v^I} \partial_{v_I^*} \end{aligned} \quad (3.16)$$

are hermitian.

In order to construct the eigenstates of S , we start from wave functions Ψ_g depending only on the scalar $\mathbf{x} = \bar{v} \cdot \bar{v}^*$ and satisfying

$$S \Psi_g = 0 \quad (3.17)$$

The general solution to the eigenstate problem of S will be given by tensorial wave functions

$$\Psi_{j_1 \dots j_q}^{i_1 \dots i_q} = N \left(v^{i_1} \dots v^{i_q} v_{j_1}^* \dots v_{j_q}^* \right) \Psi_g(x) \quad (3.18)$$

satisfying tracelessness, i.e.,

$$\delta_{i_a}^{j_a} \Psi_{j_1 \dots j_q}^{i_1 \dots i_q} = 0 \quad (\forall a \in [1, q], \bar{a} \in [1, \bar{q}]) \quad (3.19)$$

The notation $(\dots)_0$ means precisely that only the traceless part of the tensor product of v 's and v^* 's is retained. Of course the traces are just states with a smaller number of v 's and v^* 's.

These wave functions are, by construction, eigenfunctions of S :

$$S \Psi_{j_1 \dots j_q}^{i_1 \dots i_q} = s \Psi_{j_1 \dots j_q}^{i_1 \dots i_q}; \quad s = q - \bar{q} \quad (3.20)$$

Furthermore, the Laplace operator T in Eq. (3.16) acts as

$$\begin{aligned} T \Psi_{j_1 \dots j_q}^{i_1 \dots i_q} &= \left(v^{i_1} \dots v^{i_q} v_{j_1}^* \dots v_{j_q}^* \right)_{j_1 \dots j_q} (T \Psi_g - h \Psi_g') \\ &= t \Psi_{j_1 \dots j_q}^{i_1 \dots i_q} \quad h = q + \bar{q} \end{aligned} \quad (3.21)$$

so that $\Psi_g(x)$ must satisfy the differential equation

$$x \Psi_g''(x) + (h + d) \Psi_g'(x) + t \Psi_g(x) = 0 \quad (3.22)$$

whose solution can be expressed in terms of Bessel functions (see Appendix B) as follows (for convenience of notation, we set $t = m^2$ and call $r = \sqrt{x}$ the length of $\bar{v}$):

$$\Psi_g(r) = N_m r^{1-h-d} J_{h+1-d}(2mr) \quad (3.23)$$

This function is bounded as $r \rightarrow \infty$ only for $m^2 \geq 0$, and thus the mass spectrum is positive. The normalization constant N_m depends on m in such a way that for $m = 0$ the wave function does not vanish (refer to Appendix B for details).

The wave functions (3.18), with Ψ_g given by (3.23), will depend on the space-time dimension $(d+2)$ and on the three quantum numbers m , $s = q - \bar{q}$ and $h = q + \bar{q}$, the first of which is real, the last two integral (note that both $q, \bar{q} \geq 0$ so that h is a non-negative integer). For fixed m , the degeneracy is infinite and contains all possible integer spins, and even for fixed s the degeneracy is infinite and does not depend on the value of s , the degenerate states being labelled by h . The consistency of the Hilbert space generated by these wave functions is ensured by the fact that the following orthogonality relationships hold (Appendix B):

$$\left(\Psi_{j_1 \dots j_q; m, i_1 \dots i_q}^{k_1 \dots k_q} \right) = \delta_{qq'} \delta_{\bar{q}\bar{q}'} \left[\delta_{(i_1 \dots i_q)(k_1 \dots k_q)} \delta_{(j_1 \dots j_q)(\bar{j}_1 \dots \bar{j}_q)} \right]_0 \delta(m - m') \quad (3.24)$$

where

$$\delta_{(k_1 \dots k_q)(i_1 \dots i_q)} = \delta_{(k_1 i_1)}^{(1)} \delta_{(k_2 i_2)}^{(2)} \dots \delta_{(k_q i_q)}^{(q)}$$

is a symmetrized product of Kronecker deltas. Thus we can build normalizable wave packets as superpositions of the wave functions above constructed. In general, the wave-functions

of the bosonic null string are infinite products of tensorial wave functions satisfying (3.11), with values of P_σ and M^2 given by

$$\begin{aligned} P_\sigma &= \sum_{n \geq 0} n(q_n \cdot \bar{q}_n) \\ M^2 &= \sum_{n \geq 0} m_n^2 \end{aligned} \quad (3.25)$$

where $q_n, \bar{q}_n$ are nonnegative integers and m_n^2 are nonnegative real numbers. The mass spectrum is, in principle, continuous, ranging from zero to infinity. Clearly, its degeneracy is huge and the spin content is independent of the possible value of P_σ , which has to be fixed by the requirement of Lorentz covariance.

It is perhaps worth pausing to consider the result that the mass spectrum for the null string is continuous. Since our theory had no parameter to start with, the only other foreseeable alternative was that the spectrum would have consisted only of massless states. From the point of view of the limit $\alpha' \rightarrow \infty$ of usual string theories, it is clear that all the states in the leading Regge trajectory collapse to zero mass. In fact, any state of finite mass collapses to zero mass. But since there is an infinity of daughter trajectories, for any real value of α' , however large, there are still infinitely many states with arbitrarily high mass. We believe this reasoning explains heuristically our rigorous result: the mass spectrum is continuous, not massless. In fact, this is a consequence of the ordering chosen so far, p 's to the right of y 's, which essentially agrees with the Weyl ordering used in Ref. [2]. We shall now see that, with normal ordering, only a finite number of massless states survive in the Hilbert space.

IV. Lorentz Covariance of the Light Cone Quantization

Up to now we have neglected the problems arising from ordering ambiguities. Now we must take them into account. There are two inequivalent orderings of interest. The first is the one already used in the preceding section and amounts to putting the momenta on the right of the co-ordinate modes. This order is obviously equivalent to the Weyl order, as can be seen from (3.9). Weyl ordering was used by *Lizzi et al.*[2] and, as already shown by them, it does not give rise to a critical dimension. Zheleukhin agrees with them as well [3]. It is rather straightforward to check that computations involving this ordering are actually equivalent to classical computations. With this order we have shown that, up to Lorentz covariance, the Hilbert space of the physical states (and the observables acting on them) is consistent. However, there is no compelling reason, like zero point energy, which would tell us *a priori* that this ordering is the good one. An alternative ordering prescription was used by us in Ref.[5], where we chose to put the positive indexed modes of P and Y to the right of the negative indexed ones (this order is often referred to as “normal”). As shown in Ref. [5], this order implies the existence of a critical dimension for null strings, precisely the same critical dimension as for their tensionful siblings. We now want to discuss these issues in some detail, in the framework of the required Lorentz covariance.

The generators for Lorentz transformations are

$$\begin{aligned} M^{\mu\nu} &= \frac{1}{2\pi} \int_{-\pi}^{\pi} : \{ P^\mu(\sigma) Y^\nu(\sigma) - P^\nu(\sigma) Y^\mu(\sigma) \} : d\sigma = \\ &= \sum_n : (p_n^\mu y_{-n}^\nu - p_n^\nu y_{-n}^\mu) : \end{aligned} \quad (4.1)$$

As usual, the only problematic commutator is $[M^{I-}, M^{J-}]$, where

$$M^{-I} = p_0^I y_0^- + \frac{i}{p^+} \sum_n \frac{1}{n} : p_n^I y_{-n}^- : - \frac{1}{p^+} \sum_n : p_n^I y_{-n}^- : \quad (4.2)$$

where we defined

$$\begin{aligned} p_n &= p^+ p_n^- = \frac{1}{2} \sum_m : \tilde{p}_{n-m} \cdot \tilde{p}_m : \\ y_n &= i \sum_m : \tilde{p}_{n-m} \cdot \tilde{y}_m := \begin{cases} i p_n^+ y_n^- & \text{if } n \neq 0 \\ P_n^- & \text{if } n=0 \end{cases} \end{aligned} \quad (4.3)$$

As for usual strings, they constitute a realization of the constraint algebra, ie:

$$\begin{aligned} [p_m, p_n] &= 0 \\ [y_m, p_n] &= (n-m) p_{m+n} \\ [y_m, y_n] &= (n-m) y_{m+n} + C(m) \delta_{m+n} \end{aligned} \quad (4.4)$$

where we have allowed for the most general central term $C(m) = \alpha_3 m^3 + \alpha_1 m$ compatible with the Jacobi identities.

A classical computation yields

$$[M^{IJ-}, M^{J-}] = -\frac{2}{p^+} \sum_{m>0} \frac{1}{m} p_{-m}^I p_m^J P_\sigma \quad (4.5)$$

Hence, even classically, solutions with non-vanishing P_σ break the Lorentz covariance. As already mentioned, calculations with the Weyl order give the same results as classical calculations and, therefore, in this case the central term in (4.4) vanishes and closure of the Lorentz algebra implies that P_σ must vanish when acting on physical states:

$$P_\sigma |\text{phys}\rangle = \sum_{n>0} n(q_n - \bar{q}_n) |\text{phys}\rangle = 0 \quad (4.6)$$

Therefore, in this case Lorentz covariance holds in all dimensions and there are no critical dimensions. Further, from [5], we know that the normal order, with all the positive-indexed modes on the right of the negative-indexed ones, leads to a theory with critical dimension $D=26$. From (3.10) we see that the normal order requires that P_σ take only negative values:

$$P_\sigma = i \sum_{n>0} n(\vec{p}_{-n} \cdot \vec{y}_n - \vec{y}_{-n} \cdot \vec{p}_n) \quad (4.7)$$

The Fock space is created out of a vacuum $|0\rangle$ annihilated by y_n^I and p_n^I for n positive. Thus

$$P_\sigma |0\rangle = y_n^I |0\rangle = p_n^I |0\rangle = 0 \quad , \quad (n > 0) \quad (4.8)$$

as well as $M^2 |0\rangle = 0$. In this case, the central term is given by [5]:

$$C(n) = \frac{d}{6}(n^3 - n) \quad (4.9)$$

Letting a denote the eigenvalue of P_σ on physical states, after a rather tedious computation we obtain for the Lorentz algebra

$$[M^{I-}, M^{J-}] = \frac{1}{p^+ 2} \sum_{m>0} \left[\left(\frac{d}{6} - 4 \right) m - \left(\frac{d}{6} + 2a \right) \frac{1}{m} \right] p_{-m}^I p_m^J - \frac{2}{p^+} \sum_{m>0} \frac{1}{m} p_{-m}^I p_m^J (P_\sigma - a) \quad (4.10)$$

Clearly, the r.h.s. vanishes if and only if $d = 24$ and $a = -2$. Physical states will be those eigenstates of P_σ with eigenvalue $P_\sigma = -2$ which are, at the same time, mass eigenstates. It is easy to see that the only states which fulfill these conditions are the massless states $p_{-2}^I |k\rangle$, $p_{-1}^I p_{-1}^J |k\rangle$, and $(p_{-1}^I y_{-1}^J - p_{-1}^J y_{-1}^I) |k\rangle$ i.e., with spin content 0, 1,

and 2 (dilaton, Kalb-Ramond, graviton), k is the momentum of the center of mass of the spin. Indeed, all the eigenstates of $P_\sigma = -2$ are $p_{-2}^I |k\rangle$, $p_{-1}^I p_{-1}^J |k\rangle$, $p_{-1}^I y_{-1}^J |k\rangle$, $y_{-1}^I y_{-1}^J |k\rangle$. Their mass matrix is given by

$$M^2 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 2 & 0 \end{pmatrix} \quad (4.11)$$

which has zero eigenvalues with the above mentioned eigenstates. The physical interpretation of this result is unclear because the first two of these eigenstates have zero norm whereas the antisymmetric one has negative norm. In order to clarify the order issue, we "diagonalize" M^2 and P_σ by the redefinitions:

$$\begin{aligned} a_m^I &= \frac{1}{\sqrt{2}} (p_m^I + i m y_m^I) \\ b_m^I &= \frac{1}{\sqrt{2}} (p_m^I - i m y_m^I) \end{aligned} \quad (4.12)$$

such that $y_n = A_n - B_n$ and $p_n = A_n + B_n + C_n$, with

$$\begin{aligned} A_n &= \frac{1}{2} \sum_m \vec{a}_{n-m} \cdot \vec{a}_m \\ B_n &= \frac{1}{2} \sum_m \vec{b}_{n-m} \cdot \vec{b}_m \\ C_n &= \frac{1}{2} \sum_m \vec{a}_{n-m} \cdot \vec{b}_m \end{aligned} \quad (4.13)$$

The following commutators hold:

$$[a_m^I, a_n^J] = -[b_m^I, b_n^J] = -m \delta^{IJ} \delta_{m+n,0} \quad (4.14)$$

and

$$\begin{aligned} [A_n, A_m] &= (n-m) A_{m+n} + \mathcal{A}(n) \delta_{m+n,0} \\ [B_n, B_m] &= (n-m) B_{m+n} + \mathcal{B}(n) \delta_{m+n,0} \\ [C_n, C_m] &= (n-m) (A_{m+n} - B_{m+n}) - [\mathcal{A}(n) + \mathcal{B}(n)] \delta_{m+n,0} \end{aligned} \quad (4.15)$$

the corresponding transformations are:

$$\begin{aligned} [A_n, a_m^I] &= m a_{m+n}^I, & [B_n, b_m^I] &= -m b_{m+n}^I \\ [C_n, a_m^I] &= m b_{m+n}^I, & [C_n, b_m^I] &= -m a_{m+n}^I \end{aligned} \quad (4.16)$$

Furthermore, $P_\sigma = A_0 - B_0$ and $M^2 = A_0 + B_0 + C_0$ are given by the zero modes. With normal ordering (positive modes of a and b to the right), we have:

$$\mathcal{A}(n) = \mathcal{B}(n) = \frac{d}{12}(n^3 - n)$$

and the corresponding central term in (4.4) will be:

$$\mathcal{C}(n) = \mathcal{A}(n) + \mathcal{B}(n) = \frac{d}{6}(n^3 - n) \quad (4.17)$$

However, from the commutators (4.14) we see that this quantization is actually not unitary. We can circumvent this problem by redefining $b_n^I \rightarrow d_n^I = ib_n^I$. Some signs change so that now, with the same ordering, the mass eigenstates

$$\left(a_m^I \pm d_m^I \right) |0\rangle \quad (4.18)$$

have zero norm because

$$\left(a_m^I \pm d_m^I \right)^\dagger = d_m^I \mp d_m^I \quad (4.19)$$

and so on. In fact, at this point we are back to the situation leading to the mass matrix (4.11). The remaining possibility is to order all operators with the positive modes of a_m^I to the right of the negative modes of b_m^I (or viceversa); in this case

$$\mathcal{A}(m) = -R(m)$$

and there is no contribution to the central term of (4.4). It can be easily verified that this order is in fact equivalent to the Weyl order, which yields trivially vanishing central terms, i.e., $\mathcal{A}(n) = \mathcal{B}(n) = 0$. Note that also for usual strings does Weyl order yield no anomalous central term and thus no critical dimensions.

Therefore, the issue of the orderings and critical dimensions of null strings resembles to some extent the situation of the usual strings, i.e., there are essentially two orderings: the normal order which gives a theory with critical dimension $D=26$ and the Weyl order which gives a theory without critical dimensions. However, opposite to the case of usual strings where the quantization with Weyl order is not consistent, in the case of null strings it is consistent.

Quantization with normal order leads to very nice results but, unfortunately, there are physical states have zero norm. In fact, this may not be as terrible as it appears at first sight. The normal ordered null string is the rigorously quantum theory which obtains in the limit $T \rightarrow 0$ of normal strings, and we have found that pure stringy consistency implies that there are zero norm “physical” states. From this we could conjecture that

there are states of the usual tensionfull bosonic string whose norm vanishes in the limit $T \rightarrow 0$.[†] This agrees in some amount with the conjectured behavior of usual strings at very high energies [29], and the most likely physical interpretation of this fact is that it is not meaningful to construct asymptotic in and out states with null strings, i.e., string theory at infinitely high energies is *not* a usual quantum field theory [30, 31, 32, 33, 29].

Null strings do not describe a system of harmonic oscillators as usual strings do, but rather a system of massless point particles related by purely kinematical constraints. In the light cone gauge, there is a reminiscence of these constraints in the fact that the net momentum flow along the string must take a definite value, zero classically and with hermitian ordering, -2 with normal ordering. It is very interesting that the normal ordered null string develops this sort of chirality (the net momentum along the string is non-zero, so left is distinguished from right). This intriguing spontaneous generation of chirality is probably the clue to the instability of null strings. Indeed, even if we have not written it down, the curious -2 for the P_σ eigenvalue must have energy units. Our theory had no dimensional parameters to start with, and a continuous mass spectrum does not require any mass parameter. But a discrete momentum eigenvalue does. In the end, since physical states are massless, there is still no mass scale whatsoever in the theory. So, one could conjecture that the normal ordered null string (a) is unstable, and (b) it generates a “vacuum expectation value” for “left right” asymmetry. It remains beyond the scope of this paper to study the dynamics of the flow to the stable chiral phase, presumably realized by usual (heterotic?) strings.

[†] We are particularly grateful to J. Labastida and A. Morozov for most enjoyable discussions on this point.

V. Lagrangian Formulation of the Null Spinning String

Let us now consider the supersymmetric null string, starting from the action for two-dimensional supergravity and taking the string tension to zero following the general procedure outlined in Section II. The superspace approach clarifies the gauge fixing process, and is best suited for the analysis of the spectrum. Our notation closely follows Howe's for the usual spinning string [26], which is briefly reviewed in the Appendix.

In complete analogy to the bosonic action (2.1), consider the action

$$S_0 = \frac{1}{2} \int d^2\sigma d^2\theta \frac{1}{Z} E^2 (p^\alpha X^\mu \mathcal{D}_\alpha X_\mu)^2 \quad (5.1)$$

which is invariant under local world sheet supersymmetric, superWeyl, and global space-time Poincaré transformations. The superfield Z transforms as an invariant density, whose components are

$$Z = z + \theta^\alpha \rho_\alpha - \frac{1}{2} \theta^2 b \quad (5.2)$$

In the WZ gauge we obtain an action with the structure of a σ model.

$$\begin{aligned} S_0 = - \int d^2\sigma d^2\theta \frac{c^2}{z} & \left[-\frac{1}{4} \psi^2 \chi_m \gamma^n \gamma^m \chi_n \right. \\ & + \left(-\psi^2 g^{mn} \eta^{\mu\nu} + \psi^\mu \gamma^m \gamma^n \psi^\nu \right) \partial_m X_\mu \partial_n X_\nu + i \psi^2 \psi^\mu \gamma^m \partial_m \psi_\mu \\ & + \left(-\psi^2 \eta^{\mu\nu} + \psi^\mu \psi^\nu \right) B_\mu B_\nu + 2i \psi^\mu \gamma^m \psi^\nu B_\mu \partial_m X_\nu \\ & \left. + \frac{1}{2} \psi^2 \rho \gamma^m \psi^\mu \partial_m X_\mu + \frac{1}{z} \psi^2 \rho \psi^\mu B_\mu - \frac{1}{4z} \left(b - \frac{1}{z} \rho^2 \right) \psi^2 \psi^2 \right] \end{aligned} \quad (5.3)$$

Observe that this Lagrangian contains as auxiliary fields (i.e., non propagating Lagrange multipliers) the zweibein e_m^α , the last component B^μ of X^μ , and all the components of Z : z , ρ_α and b . Moreover, there is a kinetic term for the gravitino χ_m which is actually pure gauge. Indeed, we may fix the gauge such that

$$\chi_m = \gamma_m \phi \quad (5.4)$$

where the spinor ϕ is the γ trace of χ_m :

$$\phi^\alpha = \frac{1}{2} (\gamma^m \chi_m)^\alpha \quad (5.5)$$

and therefore

$$\gamma^m \gamma_m \chi_m = 0 \quad (5.6)$$

We shall fully fix the conformal gauge, consisting of the above gauge for χ_m and the choice of a diagonal zweibein,

$$e_m^\alpha = f_m \delta_m^\alpha \quad (5.7)$$

It is, however, best to keep the off diagonal components of the zweibein, denoted as g_0 and g_1 , and set them to zero only *after* deriving the equations of motion. We need the equations of motion

$$\frac{\partial L_0}{\partial A_i} \Big|_{\text{conformal gauge}} = 0 \quad (5.8)$$

for the auxiliary fields $A_i = (e_m^\alpha, Z, B) = (f_0, f_1, g_0, g_1, z, \rho, b, B)$, in order to get the constraints.

Since we wish to exhibit all the constraints as primary constraints, it is most convenient to effect a redefinition of the fields before solving for the equations of motion. This field redefinition amounts to a rescaling of the Grassman variables $\theta_\alpha \rightarrow f_0^{-1/2} \theta_\alpha$ followed by a further rescaling of the field z . Explicitly, the redefinitions are

$$\begin{aligned} X^\mu & \longrightarrow X^\mu, & \psi^\mu & \longrightarrow \sqrt{f_0} \psi^\mu, & B^\mu & \longrightarrow f_0 B^\mu \\ z & \longrightarrow f_0 z, & \rho^\alpha & \longrightarrow \sqrt{f_0} \rho^\alpha, & b & \longrightarrow f_0 b \end{aligned} \quad (5.9)$$

In terms of the new fields, the Lagrangian becomes

$$\begin{aligned} L_0 = - \frac{c^2}{z} & \left[\left(-\psi^2 g^{mn} \eta^{\mu\nu} + \psi^\mu \gamma^m \gamma^n \psi^\nu \right) \partial_m X_\mu \partial_n X_\nu \right. \\ & + \frac{i}{f_0} \psi^2 \psi \gamma^m \partial_m \psi + \frac{1}{f_0^2} \left(-\psi^2 \eta^{\mu\nu} + \psi^\mu \psi^\nu \right) B_\mu B_\nu \\ & + i \left(\frac{2}{f_0} B \psi \gamma^m \psi + \frac{1}{z} \psi^2 \rho \gamma^m \psi \right) \partial_m X + \frac{1}{f_0 z} \psi^2 \rho \psi B \\ & \left. - \frac{1}{4z} \left(b - \frac{1}{z} \rho^2 \right) \psi^2 \psi^2 \right] + \mathcal{L}(\chi_m) \end{aligned} \quad (5.10)$$

The equation for b in the conformal gauge

$$\frac{f_0 f_1^2}{2 \gamma^2} \psi^2 \psi^2 = 0 \quad (5.11)$$

implies that

$$f_0 = 0 \quad (5.12)$$

because if we set f_1 or ψ^2 equal to zero and substitute back into the Lagrangian, the results are of no practical interest. Therefore, we set $f_0 = 0$ in the conformal gauge. This means that the metric is singular and thus the original fictitious conformal invariance is broken. The canonical form for the metric is $\begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$ instead of $\begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$. The equations of motion for the remaining auxiliary fields are then

$$\frac{\partial L_0}{\partial g_0} = \frac{f_1}{z} \left[\psi^2 \left(2 \dot{X} X' + i \psi \gamma^0 \psi' \right) + 2 \dot{X} \psi \psi X' - 2i B \psi \gamma^0 \psi X' \right] = 0 \quad (5.13)$$

$$\frac{\partial L_0}{\partial \eta^1} = -\frac{f_1}{z} \left[\psi^2 \psi \gamma^1 \dot{\psi} + 2B \psi \gamma^1 \dot{\psi} \dot{X} \right] = 0 \quad (5.14)$$

$$\begin{aligned} \frac{\partial L_0}{\partial f_0} = -\frac{f_1}{z} & \left[2\dot{X} \psi \psi X' + 2iB \psi \gamma^1 \dot{\psi} X' + \psi^2 \psi \gamma^1 \dot{\psi}' \right. \\ & \left. + \frac{f_1}{z} \psi^2 \left(\rho \psi B + i\rho \gamma^0 \dot{\psi} \dot{X} - \frac{1}{4} b \psi^2 \right) \right] = 0 \end{aligned} \quad (5.15)$$

$$\frac{\partial L_0}{\partial f_1} = \frac{2f_1}{z} \left[-\psi^2 \dot{X}^2 + \left(\dot{X} \psi \right)^2 - i\psi^2 \psi \gamma^0 \dot{\psi} + \psi^2 B^2 - (B\psi)^2 - 2iB \psi \gamma^0 \dot{\psi} \dot{X} \right] = 0 \quad (5.16)$$

$$\frac{\partial L_0}{\partial \rho_\alpha} = 0 \quad (5.17)$$

$$\frac{\partial L_0}{\partial B^\mu} = \frac{2f_1^2}{z} \left[\psi^2 B^\mu + (B\psi) \psi^\mu + i\psi^\mu \gamma^0 \dot{\psi} \dot{X} \right] = 0 \quad (5.18)$$

B^μ can be solved from this last equation if we multiply it by ψ_α^μ and use the fact that $\psi_\alpha \cdot \psi_\beta = \frac{1}{2} \epsilon_{\alpha\beta} \psi^2$. The result is

$$B^\mu = \frac{2i}{3} \frac{1}{\psi^2} \psi^\mu \gamma^0 \dot{\psi} \dot{X} \quad (5.19)$$

which, put back into the Lagrangian, yields

$$L_0 = \frac{f_1^2}{z} \psi^2 \left[\dot{X}^2 + i\psi \gamma^0 \dot{\psi} - \frac{2}{3} \frac{(\dot{X} \psi)^2}{\psi^2} \right] \quad (5.20)$$

so that the equation of motion for z gives once more Eq.(5.16).

Further substituting (5.19) into (5.14), we get

$$\psi \gamma^0 \dot{\psi} = 0 \quad (5.21)$$

which, together with (5.16), yields

$$\dot{X}^2 \psi^2 - \frac{2}{3} (\dot{X} \psi)^2 = 0 \quad (5.22)$$

We may now use the equation of motion for ψ

$$i \left(\gamma^0 \psi \right)_\alpha + \frac{4}{3\psi^2} \left[\frac{(\dot{X} \psi)}{\psi^2} \psi_\alpha^\mu - \dot{X} \psi_\alpha \dot{X}^\mu \right] = 0 \quad (5.23)$$

which, multiplied by ψ_β^μ and B^μ , gives respectively

$$\psi_\alpha \cdot \psi_\beta = 0 \quad (5.24)$$

and

$$\left(\dot{X} \psi \right)^2 \dot{X} \psi_\alpha = 0 \implies \left(\dot{X} \psi \right)^2 = 0 \quad (5.25)$$

which in virtue of Lorentz covariance means

$$\dot{X} \psi_\alpha = 0 \quad (5.26)$$

Therefore, the Lagrangian we find is

$$L_0 = -\frac{f_1^2}{z} \psi^2 \left(\dot{X}^2 + i\psi \gamma^0 \dot{\psi} \right) = -\frac{f_1^2}{z} \psi^2 \left(\dot{X}^2 + i\psi_\alpha \dot{\psi}_\alpha \right) \quad (5.27)$$

plus the constraints

$$\dot{X}^2 = \dot{X} \psi_\alpha = \dot{X} X' + \frac{i}{2} \psi_\alpha \psi_\alpha' = 0 \quad (5.28)$$

The null spinning string describes an $N = 2$ massless spinning superparticle [23] at every point of the string. The null spinning string is a line of such $N = 2$ massless spinning particles, subject to the constraints (5.28).

With a bit more work, the Lagrangian (5.27) can be cast in an even simpler form. Indeed, the gauge freedom remaining after fixing the gauge (5.7) is given by

$$\begin{aligned} \xi^1 + i\zeta \gamma^5 \phi &= 0 \\ f_1 \dot{\zeta}_\alpha - \frac{1}{2} f_1 \dot{\zeta}_\alpha &= 0 \end{aligned} \quad (5.29)$$

whereas ξ_0 and the Lorentz and superWeyl transformations are unrestricted, ζ_α is the supersymmetry parameter. Upon a short calculation, the transformation law for $f_1^2 \psi^2 / z$ can be elucidated :

$$\delta \left(\frac{f_1^2 \psi^2}{z} \right) = \xi^m \partial_m \left(\frac{f_1^2 \psi^2}{z} \right) + (\xi^\mu - \xi^0) \frac{f_1^2 \psi^2}{z} \quad (5.30)$$

whereby the gauge freedom in ξ^0 may be used to set

$$z = 2 f_1^2 \psi^2. \quad (5.31)$$

Moreover, superWeyl invariance can be used to set $\phi = 0$ and $f_1 = 1$, thus $\dot{\xi}^1 = \dot{\zeta}_\alpha = 0$. In this gauge, the Lagrangian reads (compare with (2.4))

$$L_0 = \frac{1}{2} \dot{X}^2 + \frac{i}{2} \psi_\alpha \dot{\psi}_\alpha \quad (5.32)$$

and the equations of motion are

$$\ddot{X}^\mu = \dot{\psi}_\alpha^\mu = 0, \quad (5.33)$$

whereas the (primary and first class) constraints are given by (5.28). The remaining gauge freedom is restricted by

$$\begin{aligned}\xi^0 &= \dot{\xi}^1 = \dot{\xi}_\alpha = 0 \\ \xi^\mu &= \dot{\xi}^0.\end{aligned}\quad (5.34)$$

We have succeeded in constructing an action for the null spinning string whose interpretation is rather simple. Just as the bosonic null string looked like a bunch of free massless bosons, the spinning null string contains, in addition, free massless fermions. Of the four constraints (5.28), two are fermionic and two are bosonic. One of these latter ones is exactly the transverse constraint of the bosonic case, whereas the longitudinal constraint has acquired a purely fermionic piece. We shall have the occasion to study in some depth these constraints and their algebra in Section VII below.

VI. Light Cone Quantization of the Null Spinning String

Canonical quantization of the Lagrangian (5.32) is straightforward after using the left-over gauge freedom (5.34) to fix a light cone gauge [34] in order to retain the physical degrees of freedom. The most general solution to (5.33) does not involve any oscillators,

$$\begin{aligned}X^\mu(\tau, \sigma) &= F^\mu(\sigma)\tau + Y^\mu(\sigma) \\ \psi_\alpha^\mu(\tau, \sigma) &= \psi_\alpha^\mu(\sigma)\end{aligned}\quad (6.1)$$

Since the fermionic variables have no τ dependence, they do not contribute to the momentum of the string. Hence the fermionic creation operators will increase and enrich the degeneracy of the spectrum, without contributing to the mass. On the other hand, fermions do contribute to the anomaly and hence both to the critical dimension and to the net momentum along the string, P_σ .

In the light-cone gauge ($X^+ = \frac{1}{\sqrt{2}\pi} p^+ \tau$, $\psi_\alpha^+ = 0$), the solution to the constraints (5.28) is

$$\begin{aligned}P^- &= \frac{\sqrt{2}\pi}{2p^+} \vec{P}^2 \\ Y^{--I} &= \frac{\sqrt{2}\pi}{p^+} \left(\vec{P} \cdot \vec{Y}^I + \frac{i}{2} \vec{\psi}_\alpha \cdot \vec{\psi}_\alpha^I \right) \\ \psi_\alpha^- &= \frac{\sqrt{2}\pi}{p^+} \vec{P} \cdot \vec{\psi}_\alpha\end{aligned}\quad (6.2)$$

where $\vec{u} \cdot \vec{v} = u^I v^I$, $I = 1, \dots, D-2$. From the second equation above, we find that the net momentum along the spinning null string is

$$P_\sigma = \frac{1}{2\pi} \int_{-\pi}^{\pi} \left(\vec{P} \cdot \vec{Y}^I + \frac{i}{2} \vec{\psi}_\alpha \cdot \vec{\psi}_\alpha^I \right) = P_\sigma^B + P_\sigma^F \quad (6.3)$$

We may now proceed to the canonical quantization and postulate the following (graded) brackets[†] for the independent degrees of freedom

$$\begin{aligned}\left[p^I(\sigma), y^J(\sigma') \right] &= -i\delta^{IJ} \delta(\sigma - \sigma') \\ \left[\psi_\alpha^I(\sigma), \psi_\beta^J(\sigma') \right] &= \delta_{\alpha\beta} \delta^{IJ} \delta(\sigma - \sigma')\end{aligned}\quad (6.4)$$

As for the bosonic case, we consider only closed null strings. The mode expansion for the bosonic variables is given by (3.4) above, whereas, in complete analogy to the usual string, the fermionic co-ordinates are double valued. We have two sectors, depending on

[†] Here we are actually using the Dirac bracket, since the fact that the momentum of a fermionic variable is the variable itself yields a second class constraint which we have already trivially resolved [35].

whether the fermionic co-ordinates satisfy periodic boundary conditions (Ramond sector [14])

$$\psi_\alpha^I(\sigma) = \frac{1}{\sqrt{2\pi}} \sum_{m \in \mathbb{Z}} \psi_{\alpha m}^I e^{im\sigma} \quad (6.5)$$

or antiperiodic ones (Neveu Schwarz sector [15])

$$\psi_\alpha^I(\sigma) = \frac{1}{\sqrt{2\pi}} \sum_{r \in \mathbb{Z} + \frac{1}{2}} \psi_{\alpha r}^I e^{ir\sigma} \quad (6.6)$$

The (mass)² operator is exactly the same as in the purely bosonic case, Eq. (3.9). The bosonic part P_σ^B of P_σ is also given by Eq. (3.9) above. The expansion for P_σ^F is

$$P_\sigma^F = \frac{1}{2} \sum_{\alpha=1,2} \sum_{a \in \{\frac{a}{2}, \frac{a+1}{2}\}} a \tilde{\psi}_\alpha^a \cdot \tilde{\psi}_{-\alpha}^a \quad (6.7)$$

where the index a takes integer or half odd values depending on the sector (Ramond, R, or Neveu Schwarz, N) in which $\Psi_\alpha(\sigma)$ is realized. From the commutators (6.4) we obtain the following non vanishing commutators for the string "modes":

$$\begin{aligned} [y_0^-, p^+] &= -i \\ [p_m^I, y_n^I] &= -i \delta^{IJ} \delta_{m+n} \\ [\psi_{\alpha a}^I, \psi_{\beta b}^I] &= \delta^{IJ} \delta_{\alpha\beta} \delta_{a+b} \end{aligned} \quad (6.8)$$

from which it is easy to establish that the fermionic operators ψ_m are ladder operators for P_σ , but do not change the mass:

$$\begin{aligned} [M^2, \psi_{\alpha a}^I] &= 0 \\ [P_\sigma, \psi_{\alpha a}^I] &= a \psi_{\alpha a}^I \end{aligned} \quad (6.9)$$

That is, the operators $\psi_{\alpha a}^I$ will increase ($a > 0$) or decrease ($a < 0$) the value of P_σ . The wave functions for the spinning string are thus obtained from the bosonic wave functions (3.18) by applying on them creation and annihilation fermionic operators.

Concretely, the commutation relations (6.8) can be realized in two ways, depending on whether $\psi_{a>0}$ or $\psi_{a<0}$ are taken as co-ordinate variables. This ambiguity amounts to defining the normal order by putting positive indexed modes to the right of negative-indexed ones, or vice-versa. Explicitly, we realize the commutation relations with the help of Grassman variables η

$$\psi_{\alpha a}^I = \eta_\alpha \quad , \quad \psi_{\alpha -a}^I = \frac{\partial}{\partial \eta_\alpha} \quad (6.10)$$

and the two orderings mentioned above correspond to imposing (6.10) on either positive or negative a for every value of α , separately.

Clearly, the large variety of possible ordering schemes corresponds to a large variety of quantum theories with the same classical limit. These theories differ among themselves, quite pointedly, in the value for the central extension of (the Virasoro subalgebra of) the null string superalgebra. This situation is actually entirely analogous to that in usual string theory, where $D = 26$ (or 10 in the supersymmetric case) arises only as a consequence of normal ordering. It is difficult to provide a criterion for choosing one ordering over another, and all the arbitrariness is only increased when we come to null strings, for which there is no potential and hence no zero point fluctuations in the ordinary sense.

Lorentz covariance for Null Spinning Strings

The generators of the Lorentz group are

$$M^{\mu\nu} = \frac{1}{2\pi} \int_{-\pi}^{\pi} d\sigma \left[P^\mu(\sigma) Y^\nu(\sigma) - P^\nu(\sigma) Y^\mu(\sigma) + i \tilde{\Psi}_\alpha^\mu(\sigma) \tilde{\Psi}_\alpha^\nu(\sigma) \right] \quad (6.11)$$

or, in modes, $M^{\mu\nu} = L^{\mu\nu} + K^{\mu\nu}$ with $L^{\mu\nu}$ given by (4.1) and

$$K^{\mu\nu} = i \sum_{\alpha=1,2} \sum_m \tilde{\psi}_{\alpha m}^\mu \tilde{\psi}_{\alpha -m}^\nu \quad (6.12)$$

In order to avoid confusions we shall work in a fixed sector (R, R, N, N, R or N, N), i.e., to each value of α corresponds a fixed sector R or N. All mode expressions are understood to be appropriately ordered, of course.

The realization of the constraints is now

$$\begin{aligned} y_n &= y_n^B - \sum_{\alpha=1,2} \sum_a \frac{a}{2} \tilde{\psi}_{\alpha, n+a}^+ \cdot \tilde{\psi}_{\alpha, -a}^- \\ p_n &= p_n^B \\ \phi_{\alpha a} &= p^+ \tilde{\psi}_{\alpha, a}^- = \sum_m \tilde{p}_m^+ \cdot \tilde{\psi}_{\alpha, n-a}^- \end{aligned} \quad (6.13)$$

where p_n^B and y_n^B are given by (4.3). We have

$$\begin{aligned} [y_n, y_m] &= (n-m)y_{m+n} + A(m)\delta_{m+n} \\ [y_n, \phi_{\alpha, a}] &= \left(m - \frac{n}{2}\right) \phi_{\alpha, a+n} \\ \{\phi_{\alpha, a}, \phi_{\beta, b}\} &= 2\eta_{a+b} \delta_{\alpha\beta} \end{aligned} \quad (6.14)$$

where $A(m) = A^B(m) + A^F(m)$ is the ordering prescription dependent central term. The value of A^F depends also on the sector, i.e., on whether the fermion modes are integer or half-odd.

Consider the contributions to A^F coming from one of the two fermionic variables i.e., for $\alpha = 1$ or 2. Each of the two fermionic terms of the first eq. in (6.13) constitute, on their own, a Virasoro algebra:

$$[u_{\alpha,n}, u_{\alpha,m}] = (m-n)u_{\alpha,m+n} + A_{\alpha}(n) \quad (6.15)$$

where

$$u_{\alpha,n} = \sum_a \frac{a}{2} \bar{\psi}_{\alpha,n+a} \cdot \psi_{\alpha,-a} \quad (6.16)$$

As in [5], we obtain the following central terms:

$$\begin{aligned} A_{\alpha}^R(n) &= \pm \frac{d}{24} (m^3 + 2m) \\ A_{\alpha}^N(n) &= \pm \frac{d}{24} (m^3 - m) \end{aligned} \quad (6.17)$$

where the plus (minus) sign corresponds to the order with the positive (negative) modes to the right. Therefore, as well as in the bosonic case, the total fermionic central term $A^F(m)$ depends on the relative ordering of the two fermionic variables and, essentially, we have two inequivalent orderings: the normal order where all the positive modes are on the right of the negative ones and the Weyl order which in fact amounts to taking for $\alpha = 1$ the normal order, and for $\alpha = 2$ the antinormal order (i.e., the negative modes on the right). For the Weyl order, there is no central term. For the normal order, for the total central term $A(m) = A^B(m) + A^F(m)$ we have the following results:

$$\begin{aligned} A^{B-R}(m) &= \frac{d}{4} m^3 \\ A^{N-N}(m) &= \frac{d}{4} (m^3 - m) \\ A^{R-N}(m) &= A^{N-R}(m) = \frac{d}{4} \left(m^3 - \frac{1}{2} m \right) \end{aligned} \quad (6.18)$$

We did not consider mixed orderings, e.g., normal for the bosonic sector and Weyl for the fermionic sector, because they lead to unphysical critical dimensions. Using this for the Lorentz algebra, we find that

$$[M^{I-}, M^{J-}] = \frac{1}{p+2} \sum_{m>0} \left(p_{-m}^I p_m^J - p_{-m}^J p_m^I \right) \left[\frac{A(m)}{m^2} - 2m - \frac{2}{m} P_{\sigma} \right] \quad (6.19)$$

Thus, if we insert the central terms (6.18) we will get the critical dimension $D = 10$ and the P_{σ} intercept values: $a_{R-R} = 0$, $a_{N-N} = \dots = 1$ and $a_{R-N} = a_{N-R} = -\frac{1}{2}$.

We may now establish what the physical content of the theory of spinning null strings will be: with Weyl ordering the mass spectrum is continuous from zero to infinity, and the wave functions are the bosonic wave functions of Section III, on which fermionic creation operators act. Thus at any given mass value one finds infinities of bosonic and fermionic degrees of freedom, such that both sectors contain all spin values. From the naive point of view of counting degrees of freedom, there is trivial space-time supersymmetry.

With normal order the situation is much more simple and interesting: just as for the bosonic string, the spectrum contains only massless states (the fermionic variables do not contribute to the mass):

$$\begin{aligned} N-N &: p_{-1}^I |0\rangle & \psi_{\alpha, -\frac{1}{2}}^I \psi_{\beta, -\frac{1}{2}}^J |0\rangle \\ R-R &: |AB\rangle \\ R-N &: \psi_{1, -\frac{1}{2}}^I |A\rangle \\ N-R &: \psi_{2, -\frac{1}{2}}^J |A\rangle \end{aligned} \quad (6.20)$$

where A and B indices are 8 dimensional spinor indices for the Ramond zero modes, realized as Γ matrices. In this case only the state $p_{-1}^I |0\rangle$ of the N sector has zero norm, all the other states generated by fermionic creation operators have positive norm. However this spectrum is not supersymmetric and the only way we know to achieve it is the GSO projection. In fact for null strings the GSO projection also has the consequence of ensuring the quantum consistency of the theory by eliminating the zero norm states. Thus, after the GSO projection the content of the null spinning strings will be given by the massless sector of the type II superstrings, i.e., :

$$\begin{aligned} N-N &: \psi_{1, -\frac{1}{2}}^I \psi_{2, -\frac{1}{2}}^J |0\rangle \\ R-R &: |AB\rangle \\ R-N &: \psi_{1, -\frac{1}{2}}^I |A\rangle \\ N-R &: \psi_{2, -\frac{1}{2}}^J |A\rangle \end{aligned} \quad (6.21)$$

Therefore the null spinning string spectrum is that of $N = 2$ (II-type) 10 dimensional supergravity. After this beautiful results, the reader is probably anxious to turn the page and find phenomenologically sound null string theories. Alas, we have not yet constructed a heterotic null string, and it ought to be quite tricky: there are no left- and right-movers in the null string.

The extension to extended supersymmetric constructions ($N > 1$) remains to be fully explored, although a comment is in order [5]. We have refrained from complexifying our

VII. Interactions for Null Strings

One would guess that the simplest way to introduce interactions among null strings would be to find reasonable vertex operators for a variety of states, and to compute the averages of some of these vertex operators weighted by the action. Unfortunately, however, the action for the null string is essentially the determinant of the induced world-sheet metric, which is zero on shell. We have not been able to produce an appealing vertex-operator formalism for null (super)strings, though that is hardly evidence that it does not exist. And yet, it might well be that there is no operator formalism for null string scattering. This situation would be completely consistent with the fact that physical states (in the normal quantization) have zero norm. For the case of Weyl order, we suspect the theory to be inherently and trivially free, so again the amplitudes might not be computable as for usual strings. We concentrate, for simplicity, in the bosonic case, and follow Mandelstam in the light cone gauge [36].

In this marvellous gauge, all states are physical and the world sheet can be represented by a rectangle, whose horizontal axis is τ and whose ordinate is σ . The height of this rectangle is p^+ , whereas the length is, when we visualize S matrix elements, tending to infinity. The strip may have horizontal cuts in it, representing splitting and joining of strings. Actually, since we are dealing with closed strings, the total number of edges at any given τ must be even and the closedness is enforced by identifying them pairwise.

The identification of in and out states at the left and right ends of the strip presents no particular problems due to the zero norm of these states, and such a scattering diagram can be computed by evaluating [36, 13] the determinant of the propagator on the strip, subject to the appropriate boundary conditions. For usual strings, the relevant operator is $\partial_\sigma^2 - \partial_\tau^2$. In the null case, the operator of interest is simply $-\partial_\tau^2$, whose determinant can be evaluated on the strip and is almost unaffected by the number and position of horizontal cuts (cuts in the τ direction). The only dependence on the cuts comes from (a) the coupling constant g associated to every point where a cut ends or begins, and (b) the total number of cuts at different heights (different values of σ). Point (a) is trivial, but point (b) deserves some explanation. The determinant of the operator $-\partial_\tau^2$ is simply the product of the eigenvalues of the eigenfunctions of this operator *on the strip*. These functions are of the form

$$f_n(\sigma, \tau) = e^{in\tau/T} h(\sigma) \quad (7.1)$$

where we have assumed $\tau \in [-T/2, T/2]$. Now, $h(\sigma)$ must be expanded in a basis. If there are no cuts whatsoever on the strip,

$$h(\sigma) = e^{2\pi i m \sigma / p^+} \quad (7.2)$$

real fermions (technically, we deal with $N = 1$ and not $N = 2$), in contrast to the massless spinning particle studied by Brink *et al.* [23]: once the operators ψ^I are complexified, their (anticommuting) zero mode can be realized either fermionically or bosonically, thus enlarging the spectrum. This allowed the authors of Ref. [23] to include the graviton in the spinning particle spectrum. For us, however, enlarged supersymmetry does not appear so crucial, since we already have massless bosonic states with two symmetric space time indices.

and

$$\det(-\partial_{\mu\nu}^2) = \prod_{m \in \mathbb{Z}} \prod_{n \in \mathbb{Z}} n^2 \quad (7.3)$$

The product of all integers squared can be evaluated using ζ function regularization, and it is some number. The problem comes from the second infinite product. By way of comparison, consider a light cone process where two null strings merge into one, i.e., a strip with one cut from the left end to somewhere in the middle. The basis for the $h(\sigma)$ functions is now given by only those functions (7.2) which have a node at $\sigma = p_1^+/p^+$, i.e., only those functions which are periodic in all three ranges $\sigma \in [0, p^+]$, $[0, p_1^+]$, and $[p_1^+, p^+]$. We have not yet studied in detail how to regularize these products nor have we found explicitly the dependence of the answer on the number of cuts, i.e., on the genus and number of external states. An obvious way to regularize the whole thing is to introduce a small tension to modify the null string propagator and make it into the usual one. We then expect Gross and Mende's results to presumably apply[37]. Of course it would be more appealing to show directly that the ratio of any diagram with cuts to that of the uncut strip is zero, and thus

$$S = 1 \quad (7.4)$$

This is completely consistent with all our intuition about null strings but we still do not have a proof of it. The interesting question of what is the physically relevant limit (does $g \rightarrow \infty$ when $T \rightarrow 0$? how fast? may we introduce a finite coupling for null strings?, etc.), and which are the consistent first quantized interactions for Weyl and normal ordered null strings, will be addressed elsewhere.

It is clear that the result $S = 1$ should follow rather trivially, if true, from the null string formalism. A relevant and useful check on this statement should come from the study of null string field theory. Below we give the explicit form of the BRST charge for null strings. We have not investigated whether the symmetries of the theory allow for the introduction of cubic (and/or quartic?, higher order?) interactions in the second quantization of null (spinning) strings.

VIII. Hamiltonian Constraints for the Null String

In this section we turn to the constrained Hamiltonian formalism and deal with the null string as a dynamical system defined by the first class constraints. For the null bosonic string, the constraints (2.6) are, in the Hamiltonian formulation [38, 39, 40],

$$\begin{aligned} \mathcal{H}_\perp &= P^2 \\ \mathcal{H}_\parallel &= P \cdot \partial_\sigma X \end{aligned} \quad (8.1)$$

where $X^\mu(\tau, \sigma)$ is the position of the string in space time, and $P^\mu(\tau, \sigma)$ is its conjugate momentum.

We shall supersymmetrize the above constraints and recover, from a different perspective, the spinning string discussed in previous sections. Before doing so, we find of great interest to squeeze everything we can from the constraints (8.1). In particular, let us be very general and assume that we know two things about a dynamical system with X and P as phase space co-ordinates. Firstly, the (first class) constraints (8.1) are satisfied. Secondly, the co-ordinates X and P depend on two internal smooth parameters but their dynamics does not. In other words, the second input is "world sheet covariance" or reparametrization invariance. The Hamiltonian is then a pure constraint,

$$\mathcal{H} = \lambda \mathcal{H}_\perp + \mu \mathcal{H}_\parallel \quad (8.2)$$

where the Lagrange multipliers λ and μ are arbitrary. Impose now a third assumption, concerning the formulation of this (very general) theory in the Lagrangian formulation: assume that the momentum is linear in the parameter derivatives of the space time co-ordinate:

$$P = \alpha \dot{X} + \beta X' \quad (8.3)$$

Substituting this Ansatz into the expression for the Lagrangian density $\mathcal{L}$, and then identifying $P = \partial \mathcal{L} / \partial \dot{X}$, a simple system of equations for α , β , μ and λ arises, and it has two different solutions $\alpha = 0$, $\alpha = \frac{1}{2\mu}$.

In the first case, $P = \beta X'$ is independent of $\dot{X}$. With the gauge choice $\beta = \mu\beta + \lambda = 1$, we find chiral bosons [41] with Hamiltonian $\mathcal{H} = P \cdot X'$.

In the second case, $\beta = -\frac{\mu}{2\lambda}$ and we find null strings in a general gauge in which the induced world sheet metric is $\begin{pmatrix} \mu^2 & \mu \\ \mu & 1 \end{pmatrix}$, of zero determinant indeed. Schild's gauge corresponds to the gauge choice $\mu = 0$.

This relationship between chiral bosons and null strings is interesting but, unfortunately, we have not found any way of exploiting it fruitfully: it is not possible to go from one system to the other smoothly.

Now we shall supersymmetrize, as advertised, the null bosonic string in the constrained Hamiltonian formalism. The equal time Poisson bracket is quite canonical, since the relationship between P and X does not involve any second class constraint:

$$[P^\mu(\sigma), X^\nu(\sigma)] = \eta^{\mu\nu} \delta(\sigma - \sigma') \quad (8.4)$$

It is easy to check that the constraints (8.4) satisfy the following Poisson bracket algebra*

$$\begin{aligned} [\mathcal{H}_\perp(\sigma), \mathcal{H}_\perp(\sigma')] &= 0 \\ [\mathcal{H}_\perp(\sigma), \mathcal{H}_1(\sigma')] &= (\mathcal{H}_\perp(\sigma) + \mathcal{H}_\perp(\sigma')) \partial_\sigma \delta(\sigma - \sigma') \\ [\mathcal{H}_1(\sigma), \mathcal{H}_1(\sigma')] &= (\mathcal{H}_1(\sigma) + \mathcal{H}_1(\sigma')) \partial_\sigma \delta(\sigma - \sigma') \end{aligned} \quad (8.5)$$

This is the (bosonic) null string constraint algebra.

For the usual string, $\mathcal{H}_\perp \pm 2T\mathcal{H}_1 = (P^\mu \pm T \partial_\sigma X^\mu)^2$ are perfect squares, so one can choose light cone co-ordinates and carry through the supersymmetrization in the left and right directions of the light cone [14, 15]. The supersymmetrization of the null string constraints (8.1) is not as straightforward, because no linear combination of the constraints is a perfect square and hence, the square root method is not immediately applicable. Taking after the super numerical variables proposed by Collins and Tucker [42], we propose to implement the supersymmetrization of the null string in the normal and tangential directions to its dynamical evolution. Accordingly, we define the supercurrents or fermionic constraints by

$$\mathcal{S}_a(\sigma) = P_\mu(\sigma) \Psi_a^\mu(\sigma) \quad (8.6)$$

where the subindex $a = \perp, 1$ denotes the normal and tangential directions. The anticommuting variables $\Psi_\perp^\mu$ and Ψ_1^μ transform as vectors in space time and as spinors on the world-sheet. They satisfy a Clifford algebra

$$[\Psi_a^\mu(\sigma), \Psi_b^\nu(\sigma')] = i \delta_{ab} \eta^{\mu\nu} \delta(\sigma - \sigma') \quad (8.7)$$

so that

$$[\mathcal{S}_a(\sigma), \mathcal{S}_b(\sigma')] = i \mathcal{H}_\perp \delta_{ab} \delta(\sigma - \sigma') \quad (8.8)$$

and, obviously,

$$\begin{aligned} [\mathcal{H}_\perp(\sigma), \mathcal{H}_\perp(\sigma')] &= 0 \\ [\mathcal{S}_a(\sigma), \mathcal{H}_\perp(\sigma')] &= 0 \end{aligned} \quad (8.9)$$

In order to close the superalgebra, $\mathcal{H}_1$ must be modified. The correct extension is given by [4, 5]

$$\mathcal{H}_1(\sigma) = P \cdot \partial_\sigma X(\sigma) - \frac{i}{2} \left(\Psi_\perp(\sigma) \cdot \frac{\partial}{\partial \sigma} \Psi_\perp(\sigma) + \Psi_1(\sigma) \cdot \frac{\partial}{\partial \sigma} \Psi_1(\sigma) \right) \quad (8.10)$$

* The bracket $[,]$ is graded according to the variables it acts upon. We use the bracket throughout, the transition to quantum brackets will take place below without a change in bracket notation.

so that

$$\begin{aligned} [\mathcal{S}_a(\sigma), \mathcal{H}_1(\sigma')] &= \frac{1}{2} (\mathcal{S}_a(\sigma) + 2\mathcal{S}_a(\sigma')) \partial_\sigma \delta(\sigma - \sigma') \\ [\mathcal{H}_\perp(\sigma), \mathcal{H}_1(\sigma')] &= (\mathcal{H}_\perp(\sigma) + \mathcal{H}_\perp(\sigma')) \partial_\sigma \delta(\sigma - \sigma') \\ [\mathcal{H}_1(\sigma), \mathcal{H}_1(\sigma')] &= (\mathcal{H}_1(\sigma) + \mathcal{H}_1(\sigma')) \partial_\sigma \delta(\sigma - \sigma') \end{aligned} \quad (8.11)$$

The supersymmetric null string is described by the superalgebra in Eqs. (8.8), (8.9) and (8.11). Closure of the superalgebra guarantees that $\mathcal{S}_a$ and $\mathcal{H}_a$ can be imposed consistently as first class constraints. The fermionic constraints $\mathcal{S}_a$ generate local $N = 1$ supersymmetry transformations $\delta A = [A, \epsilon^a \mathcal{S}_a]$ in the temporal ($a = \perp$) and spatial ($a = 1$) directions. Explicitly,

$$\begin{aligned} \delta \Psi_a^\mu(\sigma) &= \epsilon_a(\sigma) P^\mu(\sigma) \\ \delta X^\mu(\sigma) &= \sum_a \epsilon^a(\sigma) \Psi_a^\mu(\sigma) \\ \delta P^\mu(\sigma) &= 0 \end{aligned} \quad (8.12)$$

One could extend this construction to $N > 1$ supersymmetry as in the superparticle case [43] by adding several Clifford algebras, but we shall restrict ourselves in this article to the $N = 1$ case (see, however, Ref.[5]).

The total Hamiltonian for this general co-ordinate invariant system is

$$H = \int d\sigma \sum_a (\lambda^a \mathcal{H}_a + \bar{\lambda}^a \mathcal{S}_a) \quad (8.13)$$

where λ^a are bosonic Lagrange multipliers and $\bar{\lambda}^a$ are fermionic.

The boundary conditions for the null spinning string are derived by requiring the bosonic variables λ^a , $\mathcal{H}_a$ to satisfy the same boundary conditions as if no fermions were present. Let us restrict ourselves for the time being to closed strings, for which the bosonic variables are periodic in $\sigma \in [0, 2\pi]$, then the Ψ 's are the only variables whose boundary conditions must be determined, by requiring that the functionals $\int \lambda^a \mathcal{H}_a$ and $\int \bar{\lambda}^a \mathcal{S}_a$ be well defined, i.e., they contain no surface terms. The variation of the functional $\int \bar{\lambda}^1 \mathcal{H}_1$, with terms of the form $\int \Psi \partial_\sigma \Psi$, produces the boundary term

$$\lambda^1 \left(\Psi_\perp \delta \Psi_\perp^\mu + \Psi_1 \delta \Psi_1^\mu \right) \Big|_0^{2\pi}, \quad (8.14)$$

which implies $\Psi_a^\mu(2\pi) = \pm \Psi_a^\mu(0)$, leading to two different sectors for each fermionic variable. The corresponding boundary conditions on the fermionic Lagrange multipliers $\bar{\lambda}^a$ are also periodic or anti-periodic.

The Fourier expansion is accordingly

$$\begin{aligned}
X^\mu(\sigma) &= \frac{1}{\sqrt{2\pi}} \sum_{n \in \mathbb{Z}} x_n^\mu e^{in\sigma} \\
P^\mu(\sigma) &= \frac{1}{\sqrt{2\pi}} \sum_{n \in \mathbb{Z}} p_n^\mu e^{in\sigma} \\
\Psi_a^\mu(\sigma) &= \frac{1}{\sqrt{2\pi}} \sum_{n \in \{\frac{a}{2} + \frac{1}{2}\}} \psi_n^{a\mu} e^{in\sigma}
\end{aligned} \tag{8.15}$$

with $a = \pm, i$. The non-vanishing brackets are

$$\begin{aligned}
[p_m^\mu, x_n^\nu]_- &= -i\eta^{\mu\nu} \delta_{m+n,0} \\
[\psi_m^{a\mu}, \psi_n^{b\nu}]_+ &= \delta^{ab} \eta^{\mu\nu} \delta_{m+n,0}
\end{aligned} \tag{8.16}$$

Note that $p_{-n}^\mu = p_n^{\mu\dagger}$, and similarly for x_n^μ and $\psi_n^{a\mu}$.

The algebra of constraints for the null spinning string can be neatly expressed in terms of the above operator eigenmodes. Using the definition $G(\sigma) = \frac{1}{2\pi} \sum G_n e^{in\sigma}$ for all the constraints G , the constraint modes are

$$\begin{aligned}
\mathcal{H}_n^\perp &= \sum_{k \in \mathbb{Z}} p_{n-k} \cdot p_k \\
\mathcal{H}_n^i &= i \sum_{k \in \mathbb{Z}} k p_{n-k} \cdot x_k - \frac{1}{2} \sum_{k \in \{\frac{n}{2} + \frac{1}{2}\}} k \psi_{n-k}^a \cdot \psi_k^a \\
S_n^a &= \sum_{k \in \{\frac{n}{2} + \frac{1}{2}\}} p_{n-k} \cdot \psi_k^a
\end{aligned} \tag{8.17}$$

where in the last expression the sum runs over integers or half-odd numbers depending on the sector. Care must be exercised to (normal) order all operators. Note that the constraint modes are hermitian.

The algebra satisfied by these constraint modes (in any of the sectors) is given by

$$\begin{aligned}
[\mathcal{H}_m^\perp, \mathcal{H}_n^\perp] &= 0 \\
[\mathcal{H}_m^\perp, \mathcal{H}_n^i] &= (n-m)\mathcal{H}_{m+n}^\perp + (b_3 m^3 + b_1 m) \delta_{m+n,0} \\
[\mathcal{H}_m^i, \mathcal{H}_n^i] &= (n-m)\mathcal{H}_{m+n}^i + (a_3 m^3 + a_1 m) \delta_{m+n,0} \\
[\mathcal{H}_m^\perp, S_n^a] &= 0 \\
[\mathcal{H}_m^i, S_n^a] &= (n - \frac{m}{2}) S_{m+n}^a \\
[S_m^a, S_n^b] &= \delta^{ab} \left(\mathcal{H}_{m+n}^\perp - (2b_3 m^2 + \frac{b_1}{2}) \delta_{m+n,0} \right)
\end{aligned} \tag{8.18}$$

where the most general central terms allowed by the Jacobi identities are shown.

The null super Virasoro algebra (8.18) is interesting on its own, and the central extensions give a clue as to the quantum consistency of the theory. Using the mode expansions (8.17), it is easy to check that the parameters b and β of the above extensions are zero for the null spinning string, whereas a and α reveal the conformal anomaly of the system and will be dealt with at length below. To find a physical realization of the algebra (8.18) with a non-trivial central extension b remains an open problem. This is equally true, in fact, for the purely bosonic case, which looks like the usual (anomalous) Virasoro algebra plus the “tail” provided by the $\mathcal{H}_\perp$ modes, which behave as faithful representations of the $\mathcal{H}_\perp$ Virasoro algebra except that, tragically and crucially...they commute among themselves!

IX. BRST Quantization of the Null Spinning String

We quantize the null spinning string using the BRST invariance [25] in its path integral BFV formalism [44, 45], both because of its elegance [46] and because the formalism is most suited to the constrained Hamiltonian approach.

The Hamiltonian (8.13) is[†]

$$H = \sum_{\alpha=\perp,1} \int_0^\pi d\sigma \left(\lambda^\alpha \mathcal{H}_\alpha + \dot{\lambda}^\alpha \mathcal{S}_\alpha \right) \quad (9.1)$$

The primary first class constraints $\mathcal{H}_\alpha$ and $\mathcal{S}_\alpha$ involve the variables X^μ , P^μ and Ψ^μ . They span the classical algebra found above, which we display in one place for convenience:

$$\begin{aligned} [\mathcal{H}_\perp(\sigma), \mathcal{H}_\perp(\sigma')] &= 0 \\ [\mathcal{H}_\perp(\sigma), \mathcal{H}_1(\sigma)] &= \left(\mathcal{H}_\perp(\sigma) + \mathcal{H}_\perp(\sigma') \right) \partial_\sigma \delta(\sigma - \sigma') \\ [\mathcal{H}_1(\sigma), \mathcal{H}_1(\sigma')] &= \left(\mathcal{H}_1(\sigma) + \mathcal{H}_1(\sigma') \right) \partial_\sigma \delta(\sigma - \sigma') \\ [\mathcal{S}_\alpha(\sigma), \mathcal{H}_\perp(\sigma)] &= 0 \\ [\mathcal{S}_\alpha(\sigma), \mathcal{H}_1(\sigma)] &= \frac{1}{2} \left(\mathcal{S}_\alpha(\sigma) + 2\mathcal{S}_\alpha(\sigma') \right) \partial_\sigma \delta(\sigma - \sigma') \\ [\mathcal{S}_\alpha(\sigma), \mathcal{S}_b(\sigma')] &= i \mathcal{H}_\perp \delta_{ab} \delta(\sigma - \sigma') \end{aligned} \quad (9.2)$$

The phase space of the theory includes the Lagrange multipliers also as canonical variables. The untilded Lagrange multipliers λ^α are even, the tilded ones $\dot{\lambda}^\alpha$ are odd. Their conjugate momenta are $\pi_\alpha(\sigma)$ and $\tilde{\pi}_\alpha(\sigma)$, with non zero graded brackets

$$\left\{ \pi_\alpha(\sigma), \lambda^b(\sigma') \right\} = \left[\tilde{\pi}_\alpha(\sigma), \dot{\lambda}^b(\sigma') \right] = -\delta_\alpha^b \delta(\sigma - \sigma') \quad (9.3)$$

The first class constraints of the theory are the original $\mathcal{H}$'s and $\mathcal{S}$'s, and the π 's. Enlarge the phase space by introducing ghosts η^α and $\tilde{\eta}^\alpha$ for the original constraints $\mathcal{H}_\alpha$ and $\mathcal{S}_\alpha$, and ghosts N^α and $\tilde{N}^\alpha$ for the Lagrange momenta π_α and $\tilde{\pi}_\alpha$. Introduce also the corresponding conjugate ghost momenta $\mathcal{P}_\alpha$, $\tilde{\mathcal{P}}_\alpha$, Π_α and $\tilde{\Pi}_\alpha$, with non zero brackets

$$\left[\mathcal{P}_\alpha(\sigma), \eta^b(\sigma') \right] = \left[\tilde{\mathcal{P}}_\alpha(\sigma), \tilde{\eta}^b(\sigma') \right] = \left[\Pi_\alpha(\sigma), N^b(\sigma') \right] = \left[\tilde{\Pi}_\alpha(\sigma), \tilde{N}^b(\sigma') \right] = -\delta_\alpha^b \delta(\sigma - \sigma') \quad (9.4)$$

Note the untilded ghosts and their momenta are odd (anticommuting).

The classical BRST charge is a sum of two terms [47, 48],

$$\Omega = \Omega^0 + \Omega^1 \quad (9.5)$$

[†] The analysis could be carried through in terms of Fourier modes, in notation closer to the usual Virasoro modes – we prefer to display the full σ dependence for once in order to avoid misleading simplifications.

where Ω^0 is the sum of products (ghost) $\times$ (constraint)

$$\Omega^0 = \sum_{\alpha=\perp,1} \int d\sigma \left(\eta^\alpha \mathcal{H}_\alpha + \tilde{\eta}^\alpha \mathcal{S}_\alpha + N^\alpha \pi_\alpha + \tilde{N}^\alpha \tilde{\pi}_\alpha \right) \quad (9.6)$$

and Ω^1 takes care of the non-trivial constraint algebra,

$$\Omega^1 = -\frac{1}{2} (-)^{\epsilon_\alpha \epsilon_\beta} \eta^\alpha \tilde{\eta}^\beta C_{\alpha\beta}^a \mathcal{P}_a \quad (9.7)$$

The structure functions $C_{\alpha\beta}^a$ can be read off from the algebra $[G_a, G_b] = C_{ab}^\alpha G_\alpha$ with G_a any of the first class constraints (the $\mathcal{H}$'s, $\mathcal{S}$'s and π 's), and $\epsilon_b = 0(1)$ if the constraint G_b is bosonic (fermionic). In the case at hand, the non vanishing structure functions following from the classical superalgebra (9.2) are as follows:

$$\begin{aligned} C_{\perp 1}^1 &= C_{11}^1 = [\delta(\sigma - \sigma'') + \delta(\sigma' - \sigma'')] \partial_\sigma \delta(\sigma - \sigma') \\ C_{\alpha b}^1 &= i\delta(\sigma - \sigma') \delta(\sigma' - \sigma'') \delta_{\alpha b} \\ C_{\alpha 1}^b &= \left[\frac{1}{2} \delta(\sigma - \sigma'') + \delta(\sigma' - \sigma'') \right] \partial_\sigma \delta(\sigma - \sigma') \delta_\alpha^b \end{aligned} \quad (9.8)$$

Explicitly, then,

$$\begin{aligned} \Omega^1 &= - \int d\sigma d\sigma' \delta(\sigma - \sigma') \left\{ \eta^1(\sigma') \eta^1(\sigma) \mathcal{P}_1(\sigma) + \eta^1(\sigma) \eta^1(\sigma') [\mathcal{P}_\perp(\sigma) + \mathcal{P}_\perp(\sigma')] \right. \\ &\quad \left. + \sum_{\alpha=\perp,1} \eta^1(\sigma') \tilde{\eta}^\alpha(\sigma) \left[\frac{1}{2} \tilde{\mathcal{P}}_\alpha(\sigma) + \tilde{\mathcal{P}}_\alpha(\sigma') \right] \right\} \\ &\quad - \frac{1}{2} \sum_{\alpha=\perp,1} \int d\sigma \tilde{\eta}^\alpha(\sigma) \tilde{\eta}^\alpha(\sigma) \mathcal{P}_\perp(\sigma) \end{aligned} \quad (9.9)$$

Using the null spinning string constraint algebra and the commutation relations among all the fields, it is straightforward to check that the classical BRST charge is nilpotent,

$$[\Omega, \Omega] = 0 \quad (9.10)$$

It is somewhat easier to analyze the effect of the central extensions (resulting from an ordering prescription for quantum operators) in terms of Fourier modes, using the algebra (8.18). In obvious notation, one finds that

$$\begin{aligned} [\Omega, \Omega] &= \sum_{n \in \mathbb{Z}} \left[(b_3 n^3 + b_1 n) \eta_n^\perp \eta_{-n}^\perp + (a_3 n^3 + a_1 n) \eta_n^1 \eta_{-n}^1 \right] \\ &\quad + \sum_{\alpha=\perp,1} \sum_{n \in \mathbb{Z}} \left(2b_\alpha n^2 + \frac{b_1}{2} \right) \tilde{\eta}_n^\alpha \eta_{-n}^\alpha \end{aligned} \quad (9.11)$$

which proves nilpotency of the BRST charge Ω iff the anomalous central terms vanish.

For the quantum BRST charge, one has [5]

$$\begin{aligned} \Omega = & \eta_k^a \mathcal{H}_{-k}^a + \tilde{\eta}_k^a S_{-k}^a + (m-n) \eta_m^1 \mathcal{P}_{-m-n}^\perp + \frac{1}{2}(m-n) \eta_m^1 \eta_n^1 \mathcal{P}_{-m-n}^\perp \\ & + (m-n) \frac{n}{2} \tilde{\eta}_m^a \tilde{\eta}_n^1 \mathcal{P}_{-m-n}^\perp - \frac{1}{2} \tilde{\eta}_m^a \tilde{\eta}_n^a \mathcal{P}_{-m-n}^\perp + \alpha \eta_0^1 \end{aligned} \quad (9.12)$$

where all repeated indices are summed over, all operator products are assumed to be ordered according to some prescription, and α is an unknown constant, which takes into account that $\mathcal{H}_0^1$ presents ordering problems.

The single Virasoro subalgebra for the full system is generated by

$$L_m = [\Omega, \mathcal{T}_m^1] = L_m^{\text{matter}} + L_m^\perp + L_m^1 + L_m^{\frac{1}{2}} + L_m^{\frac{1}{2}} + L_m^1 - \alpha \delta_m \quad (9.13)$$

where

$$\begin{aligned} L_m^{\text{matter}} &= -\mathcal{H}_m^1 = L_m^x + L_m^{\psi_1} + L_m^{\psi_1} \\ L_m^x &= -ik p_{m-k} \cdot x_k \\ L_m^{\psi_1} &= \frac{k}{2} \psi_{m-k}^a \cdot \psi_k^a \\ L_m^a &= (k+m) \eta_k^a \mathcal{P}_{m-k}^a \\ L_m^{\frac{1}{2}} &= (k + \frac{m}{2}) \tilde{\eta}_k^a \tilde{\mathcal{P}}_{m-k}^a \end{aligned} \quad (9.14)$$

with no sum over a in the last three expressions. Each set of the above operators satisfy an anomalous Virasoro algebra,

$$[L_m^{\frac{1}{2}}, L_n^{\frac{1}{2}}] = (m-n) L_{m+n}^{\frac{1}{2}} + A_m^{\frac{1}{2}} \delta_{m+n} \quad (9.15)$$

In the case of Weyl ordering, $A_m = \alpha = 0$ and there is no critical dimension [2]. If we normal order all modes [5], then to compute the anomaly $A_m = \sum A_m^i$ we may use the identity

$$\langle 0 | L_m L_{-m} | 0 \rangle = A_m - 2m\alpha \quad (9.16)$$

We find that

$$\begin{aligned} A_m^x &= \frac{D}{6}(m^3 - m) \\ A_m^\perp &= A_m^1 = \frac{1}{6}(-13m^3 + m) \\ A_m^{\frac{1}{2}R} &= A_m^{\frac{1}{2}L} = \frac{1}{12}(11m^3 - 2m) \\ A_m^{\frac{1}{2}N} &= A_m^{\frac{1}{2}N} = \frac{1}{12}(11m^3 + m) \\ A_m^{\psi_1 R} &= A_m^{\psi_1 R} = \frac{D}{24}(m^3 + 2m) \\ A_m^{\psi_1 N} &= A_m^{\psi_1 N} = \frac{D}{24}(m^3 - m) \end{aligned} \quad (9.17)$$

Hence, quantum nilpotency of the BRST charge Ω implies in this case [5]

$$\left. \begin{aligned} \text{RR} : \quad D=10, \quad \alpha=0 \\ \text{NN} : \quad D=10, \quad \alpha=-1 \\ \text{RN} : \quad D=10, \quad \alpha=-1/2 \\ \text{NR} : \end{aligned} \right\} \quad (9.18)$$

Before concluding this section, we should like to derive from the BRST charge the action (or rather, an action) for null (supersymmetric) strings. Let us work with the σ dependence, rather than the modes.

In terms of an arbitrary odd and imaginary gauge fixing function Ψ , the effective Hamiltonian density is given by

$$\mathcal{H}_{\text{eff}} = -[\Psi, \Omega] \quad (9.19)$$

and the effective action is

$$S_{\text{eff}} = \int_{t_1}^{t_2} \mathcal{L} dt \quad (9.20)$$

where we choose the following BRST invariant boundary conditions [45]:[†]

$$\pi_a(t_i) = p_a(t_i) = \Pi_a(t_i) = \tilde{\pi}_a(t_i) = \tilde{p}_a(t_i) = \tilde{\Pi}_a(t_i) = 0 \quad (9.21)$$

The effective Lagrangian is

$$\mathcal{L} = -\mathcal{H}_{\text{eff}} + \int d\sigma \left[\dot{X} \cdot P + \sum_{a=1,1} \dot{\Psi}_a \cdot \Psi_a + \left(\dot{\lambda}^a \pi_a + \dot{\eta}^a \mathcal{P}_a + \dot{N}^a \Pi_a + [\text{ilded}] \right) \right] \quad (9.22)$$

so that the generating functional reads as

$$Z = \int \mathcal{D}X^\mu \mathcal{D}P^\mu \mathcal{D}\Psi_a^\mu (\mathcal{D}\eta^a \mathcal{D}\mathcal{P}_a \mathcal{D}\lambda^a \mathcal{D}\pi_a \mathcal{D}N^a \mathcal{D}\Pi_a) \times (\text{ilded}) e^{iS_{\text{eff}}} \quad (9.23)$$

Following Henneaux [45], we choose as a gauge fixing function something of the form

$$\Psi = \sum_a \int d\sigma (\mathcal{P}_a \lambda^a + \Pi_a \chi^a + [\text{ilded}]) \quad (9.24)$$

where χ^a and λ^a do not involve any ghosts. Even with this restriction, there is still an enormous variety of possible gauge fixings. The simplest one would appear to be merely

[†] Because of these boundary conditions, the ghosts Π are often referred to as $\tilde{\eta}$, and accordingly the momenta N are labelled as $\tilde{P}$. We shall not indulge in such notation.

$\chi^a = \tilde{\chi}^a = 0$, but in fact the most convenient gauge is (to our taste) Teitelboim's proper time gauge [49, 50, 51], generalized to the current situation:

$$\begin{aligned}\chi^\perp &= 0 \\ \chi^1 &= -\frac{1}{\epsilon}\lambda^1 \\ \tilde{\chi}^a &= -\frac{1}{\epsilon}\tilde{\lambda}^a\end{aligned}\quad (9.25)$$

where ϵ is an arbitrary parameter. A possible variation of this generalization is to take $\tilde{\chi}^\perp = 0$ and $\tilde{\chi}^1$ as above.

It is simple to find the variation of Ψ under a BRST transformation:

$$\begin{aligned}\mathcal{H}_{\text{eff}} &= \int d\sigma \left[\lambda^a \mathcal{H}_a + \tilde{\lambda}^a \mathcal{S}_a - \mathcal{P}_a N^a - \tilde{\mathcal{P}}_a \tilde{N}^a + \frac{1}{\epsilon} \lambda^1 \pi_1 + \frac{1}{\epsilon} \tilde{\lambda}^a \pi_a + \frac{1}{\epsilon} N^1 \Pi_1 + \frac{1}{\epsilon} \tilde{N}^a \tilde{\Pi}_a + \tilde{\lambda}^a \tilde{\eta}^a \mathcal{P}_\perp \right] \\ &+ \int d\sigma d\sigma' \delta'(\sigma - \sigma') \left[\lambda^1(\sigma) \tilde{\eta}^a(\sigma') \left(\mathcal{P}_a(\sigma) + \frac{1}{2} \mathcal{P}(\sigma') \right) - \tilde{\lambda}^a(\sigma) \eta^1(\sigma') \left(\frac{1}{2} \mathcal{P}_a(\sigma) + \mathcal{P}(\sigma') \right) \right] \\ &- \left(\eta^1(\sigma) \lambda^1(\sigma') - \eta^1(\sigma') \lambda^1(\sigma) \right) \mathcal{P}_1(\sigma) + \left(\lambda^1(\sigma) \eta^\perp(\sigma') + \lambda^\perp(\sigma) \eta^1(\sigma') \right) \left(\mathcal{P}_\perp(\sigma) + \mathcal{P}_\perp(\sigma') \right)\end{aligned}\quad (9.26)$$

We now perform a rescaling [52]

$$\begin{aligned}\pi_1 &\rightarrow \epsilon \pi_1 \\ N^1 &\rightarrow \epsilon N^1 \\ \tilde{\pi}_a &\rightarrow \epsilon \tilde{\pi}_a \\ \tilde{N}^a &\rightarrow \epsilon \tilde{N}^a\end{aligned}\quad (9.27)$$

in the generating functional (9.23), after which the parameter ϵ is taken to zero. The effective Lagrangian is then, in terms of the phase space variables,

$$\begin{aligned}\mathcal{L} &= \int d\sigma \left[\dot{X} \cdot P + \dot{\Psi}_a \cdot \Psi_a + \dot{\lambda}^\perp \pi_\perp + \tilde{\eta}^a \mathcal{P}_a + \tilde{\eta}^a \tilde{\mathcal{P}}_a + \dot{N}^\perp \Pi_\perp \right. \\ &\quad \left. - \lambda^\perp P^2 - \lambda^1 \left(P \cdot X' + \frac{i}{2} \Psi_a \cdot \Psi'_a \right) - \tilde{\lambda}^a P \cdot \Psi_a \right. \\ &\quad \left. - N^\perp \mathcal{P}_\perp - \lambda^1 \pi_1 - \tilde{\lambda}^a \tilde{\pi}_a + \Pi_1 N^1 + \tilde{\Pi}_a \tilde{N}^a + \tilde{\lambda}^a \tilde{\eta}^a \mathcal{P}_\perp \right] \\ &+ \int d\sigma d\sigma' \delta(\sigma - \sigma') \left[\lambda^1(\sigma) \tilde{\eta}^a(\sigma') \left(\mathcal{P}_a(\sigma) + \frac{1}{2} \mathcal{P}_a(\sigma') \right) \right. \\ &\quad \left. + \left(\lambda^1(\sigma) \eta^\perp(\sigma') + \lambda^\perp(\sigma) \eta^1(\sigma') \right) \left(\mathcal{P}_\perp(\sigma) + \mathcal{P}_\perp(\sigma') \right) \right. \\ &\quad \left. + \left(\eta^1(\sigma') \lambda^1(\sigma) - \eta^1(\sigma) \lambda^1(\sigma') \right) \mathcal{P}_1(\sigma) - \tilde{\lambda}^a(\sigma) \eta^1(\sigma') \left(\frac{1}{2} \tilde{\mathcal{P}}_a(\sigma) + \tilde{\mathcal{P}}_a(\sigma') \right) \right]\end{aligned}\quad (9.28)$$

The functional integral in the Lagrange momenta π_a , $\tilde{\pi}_a$ can now be performed, yielding a bunch of δ functions of the Lagrange multipliers (note that fermionic δ functions are well defined)

$$\delta(\dot{\lambda}^\perp) \delta(\dot{\lambda}^\perp) \delta(\lambda^1) \delta(\dot{\lambda}^1) \quad (9.29)$$

which allow us to eliminate the τ dependence of $\lambda^\perp$, and to eliminate $\dot{\lambda}^\perp$, λ^1 , and $\dot{\lambda}^1$ altogether. The ghost momenta Π_a , $\tilde{\Pi}_a$ associated to the Lagrange multipliers can be integrated out as well, yielding similarly

$$\delta(\dot{N}^\perp) \delta(\dot{N}^\perp) \delta(N^1) \delta(\dot{N}^1) \quad (9.30)$$

The Gaussian integration over the co-ordinate momenta P^μ can also be performed, taking care that the invariant world sheet measure is in this gauge $d\tau d\sigma \lambda_\perp$ and that the bosonic zero modes for $\lambda_\perp$ may take only positive values [53]. The resulting vacuum functional is

$$Z = \int \mathcal{D}X^\mu \mathcal{D}\Psi_a^\mu \mathcal{D}\lambda^\perp \mathcal{D}N^\perp \mathcal{D}\eta^a \mathcal{D}\tilde{\eta}^a \mathcal{D}\tilde{\pi}_a \mathcal{D}\tilde{\lambda}^\perp \delta(\dot{N}^\perp) \exp \left\{ i \int_{t_i}^{t_f} d\tau \int d\sigma \left[-\frac{1}{4\lambda^\perp} \dot{X}^2 + \dot{\Psi}_a \cdot \Psi_a + \tilde{\eta}^a \mathcal{P}_a + \tilde{\eta}^a \tilde{\mathcal{P}}_a - \left(\lambda^\perp \eta^\perp + \lambda^\perp \eta^1 \right) \mathcal{P}_\perp \right] \right\} \quad (9.31)$$

The ghost integrals involve an operator matrix whose determinant contains $\lambda^\perp$ and information about the topology (trivial in the free case) of the world sheet. Its evaluation (with ζ function regularization, for instance) requires a careful distinction between the Ramond and Neveu-Schwarz sectors and will be presented elsewhere. It can be written formally as

$$Z = \int \mathcal{D}X^\mu \mathcal{D}\Psi_a^\mu \mathcal{D}\lambda^\perp \delta(\dot{\lambda}^\perp) F[\lambda^\perp] \exp \left\{ i \int_{t_i}^{t_f} d\tau \int d\sigma \left[\frac{1}{4\lambda^\perp} \dot{X}^2 + \dot{\Psi}_a \cdot \Psi_a \right] \right\} \quad (9.32)$$

where we expect the functional $F[\lambda^\perp]$ to eliminate the negative norm states from the original degrees of freedom. Evidently, the action for the matter fields is reduced to that for free bosonic and fermionic co-ordinates, with the multiplier $\lambda^\perp$ playing the role of a conformal factor.

X. Conclusions

In this work, we have introduced and studied a system with more structure than the point particle, and yet simpler than the usual strings with non zero tension because the transverse constraint $\mathcal{H}_\perp$ is merely p^2 . Classically, from the viewpoint of a given point of the string, this means that the nearest neighbour interaction, the harmonic oscillator potential, is absent. Hence every point on the string is pretty much free, except of course that the longitudinal constraint $\mathcal{H}_\parallel$ enforces its velocity to be perpendicular to the string itself. In a theory of higher dimensional objects with a Nambu action, the limit of vanishing tension has even more dramatic consequences: the non linear interaction present in the transverse constraint $\mathcal{H}_\perp$ disappears, and thus quantization is possible [54, 55, 18].

It could appear as if the $T \rightarrow 0$ limit of string theories is, modulo technicalities, a simple limiting case. Naively, it corresponds to the slope of the Regge trajectories becoming infinite *i.e.*, projecting all the massive states onto the $M^2 = 0$ axis. The opposite limit, $T^{-1} \rightarrow 0$, has been used in the past to find the field theory limit of string theories, because it represents the Regge slope going to zero, hence the boosting to infinite mass of all the massive states. Our analysis above supports the idea that, once the Regge slope is fixed (infinite), the mass intercept remains at zero, and only the same finite (and small) number of massless states as were present in the field theory limit survive.

This most intriguing feature of the (normal ordered) null (super)string is closely related to the duality symmetry observed for closed strings and which relates the characteristic length of compactification R to its inverse $\frac{1}{R}$ [56] as well as for the temperature in string thermodynamics $T \rightarrow \frac{1}{T}$ [30].

Surprisingly, the quantization of the Schild action and its supersymmetric extension yield two very different theories depending on the operator ordering prescription adopted. Note that the only two possibilities for the spectrum of a theory without any mass scale are realized in these two scenarios: massless and continuous spectra. The massless case carries the bonus of a finite spectrum.

Hermitian or Weyl ordering yields (contrary to the usual string case) a consistent theory which, nevertheless, does not appear to be the limit of the usual string; it has no conformal anomaly, hence no critical dimension, however the fact that its mass spectrum is continuous points out that in the supersymmetric case it could be interpreted as a degenerated plane. Further, the infinitely enormous degeneracy of the continuous spectrum probably allows for space-time supersymmetry regardless of the GSO projection. An implementation of consistent interactions would perhaps help to clear this issue.

In the normal ordering the conformal anomaly shows up to yield a definite critical

dimension and a definite (non zero) intercept for the generator of rigid rotations along the string, P_σ . The closed null (super)string constraint algebra contains only one Virasoro subalgebra, and the appearance of a central term reflects itself on a shift in the momentum along the string P_σ , not the mass squared operator.

The independence of the string parametrization on the chosen origin for the space-like string co-ordinate σ is a natural condition for the quantum consistency of the Hilbert space, merely the zero-mode of the longitudinal constraint, and not something extra put in by hand. The quantum net non zero twist developed by some of the sectors as a consequence of the non zero “intercept”, plays essentially the same role as the intercept of the tensionful strings, namely it determines the spin content of the massless states, which is the same in both cases.

In this context, supersymmetry appears once more to be essential for the quantum consistency of strings, whereas the null bosonic strings suffer of zero norm states, null superstrings have a well defined Hilbert space where the GSO projection play a crucial role in order to eliminate consistently the zero norm states. In any case the study of the interactions of null strings is necessary in order to clarify their applicability to the study of the very high energy limit of string theories. Classically we would expect that there are only frontal interactions so that the structure of the quantum theory would be extremely simple as stressed, for instance, by Gross [29].

Another area worth investigating is the null string second quantized formalism [57], the starting point for which is the nilpotent BRST charge of Section IX. Note that in usual string field theory, one is led to contemplate interactions among strings with different values of $\alpha' = T^{-1}$ [58, 59]. Whether the formalism already developed allows us to consider strings with $\alpha' = \infty$ remains to be explored. Note that the BRST charge of the Heisenberg algebra [60] which yields massless particles in arbitrary representations of the Lorentz group [61] is not the same as ours, which has much more structure.

Allowing for some free thought, one might conjecture that null (super)strings are to usual strings as massless fields are to massive ones: their crucial purpose is to realize the gauge degrees of freedom. In strict analogy to the particle case, a symmetry of the null (supersymmetric) string, so far unidentified, could yield a finite non zero tension after its breaking.

Another open question concerns the thermodynamics of null (supersymmetric) strings themselves, not the thermodynamics of usual strings at high temperatures [30].[†] In principle, the limiting or Hagedorn temperature is zero, so one would expect that the

[†] We have found indeed the finite spectrum conjectured by the authors of Ref.[30].

interacting theory, if it exists, must be established previous to any relevant explorations in this direction. Nevertheless, the simple fact that the degeneracy of states is independent of the mass level (for the free Weyl-ordered null string) already allows us to explore the beast. The partition function for the null string looks like

$$Z = \int_0^\infty dm g(m) e^{-\beta m} = g \int_0^\infty dm e^{-\beta m} = g/\beta \quad (10.1)$$

where g is a big overall infinity. The free energy is continuous, the specific heat is constant, and $< \Delta E^2 > / < E >^2 = 1$. These and related issues are currently under investigation. We expect null superstrings to help significantly in the quest for understanding quantum gravity and many other stringy miracles.

Acknowledgements. We would like to thank L. Alvarez Gaumé, A.P. Balachandran, J. Barcelos Neto, J. van der Bij, M. Scholl, C. Gómez, R. Grimm, J. Labastida, A. Morozov, S. Ouvry, A.V. Ramallo, V.O. Rivelles, G. Sierra, J. Zanelli and A.A. Zheltukhin for interesting discussions and comments. Only after the completion of this work did we learn about the interesting work of Zheltukhin [3, 54, 9] on null strings and membranes. We have some points in common, on issues of interpretation we are in friendly disagreement. M.R. thanks J. Wess at the Institut für Theoretische Physik, Karlsruhe, and M. Schmidt at the Institut für Theoretische Physik, Heidelberg, for hospitality at some stage of this work. C.R. thanks the CERN Theory Division for hospitality at some other stage of this work. J.G.'s work was partially financed through a CAPES fellowship. C.R. is supported by the Deutsche Forschungsgemeinschaft.

Appendix A

We specify notations by reviewing spinning strings in (1, 1) superspace, following Howe [26]. The action is

$$S = T \int d^2\sigma d^2\theta E P^\alpha X^\mu P_\alpha X_\mu$$

where $E = \text{sdet}(E_M^\Lambda)$ is the invariant density and P_α are the fermionic covariant derivatives. Also, $M = (m, \alpha)$, with $\alpha = 1, 2$. The derivative of the superfield

$$X^\mu(\sigma, \theta) = X^\mu(\sigma) + \theta^\alpha \psi_\alpha^\mu(\sigma) - \frac{1}{2} \theta^2 B^\mu(\sigma)$$

is

$$\mathcal{D}_\alpha X^\mu = E_\alpha^M \partial_M X^\mu$$

and its transformation under infinitesimal reparametrizations is given by

$$\delta X^\mu = \xi^M(\sigma, \theta) \partial_M X^\mu(\sigma, \theta)$$

The geometrical quantities of superspace are the torsion and the curvature, which can be constrained so that only one auxiliary field survives in the supergravity multiplet. Furthermore, the reparametrization freedom can be restricted by allowing only transformations with parameters $\xi^M(\sigma)$ (Wess-Zumino gauge) so that the only gauge fields are the graviton e_m^α and the gravitino χ_m^α . Therefore in this WZ gauge the supergravity multiplet contains only $(e_m^\alpha, \chi_m^\alpha, A)$, with A the auxiliary field.

The superstring action is also invariant under superWeyl transformations

$$\begin{aligned} \delta E_M^\alpha &= \Lambda E_M^\alpha \\ \delta E_M^\alpha &= \frac{1}{2} \Lambda E_M^\alpha - \frac{i}{2} E_M^\alpha \gamma_m^{\alpha\beta} P_\beta \Lambda \end{aligned}$$

Due to this invariance, the superstring Lagrangian reduces in the WZ gauge to

$$L = c \left(1 - \frac{1}{8} \theta^2 \chi_m^\alpha \gamma^\mu \gamma^\nu \chi_n^\alpha \right) P^\alpha X \cdot P^\alpha X \quad (A.1)$$

where $c = \text{det}(e_m^\alpha)$ and P_α is almost the flat superspace covariant derivative,

$$\begin{aligned} \mathcal{D}_\alpha &= \partial_\alpha + i(\gamma^\mu \theta)_\alpha \partial_m \\ \gamma^m &= \gamma^\alpha \gamma^\mu \end{aligned}$$

Note that the auxiliary field A does not appear in the Lagrangian (A.1).

In components,

$$\begin{aligned} L = c \left[\psi^2 - 2\theta\psi B - 2i\theta\gamma^m \partial_m X \right. \\ \left. + \theta^2 \left(g^{mn} \partial_m X \partial_n X - i\psi\gamma^m \partial_m \psi + B^2 - \frac{1}{8} \psi^2 \chi_m^\alpha \gamma^\mu \gamma^\nu \chi_n^\alpha \right) \right] \end{aligned}$$

Appendix B

In Section III we have shown that the wave functions of the null string (in the Weyl ordering) are constructed on the solution to the differential equation

$$\Psi''(r) + \frac{1}{r}[2(h+d)-1]\Psi'(r) + 4m^2\Psi(r) = 0 \quad (\text{B.1})$$

obtained from (3.22) by the rescaling $x \rightarrow r = \sqrt{x}$. It is well known that equations of this type have solutions which can be expressed in terms of Bessel functions [62]. In general, there are solutions for any complex value of m^2 , but we are interested in *normalizable* wave-functions; i.e., m^2 has to be real because it is the eigenvalue of a hermitian operator (3.16). Thus the wave functions corresponding to different values of m^2 must be orthogonal. Since m^2 is a real and not discrete variable, normalizability must be understood in the larger sense that we can build normalizable (in the strict sense) wave packets. In other words, the orthonormality conditions for the wave functions $\Psi_{gm} = \Psi_m$ read

$$(\Psi_m, \Psi_{m'}) = \delta(m - m') \quad (\text{B.2})$$

In fact, if we take Bessel functions of the first kind, the corresponding solutions immediately fulfill this condition for $m, m' \neq 0$. The special case $m = 0$ will be dealt with separately below, although it will turn out that it can be obtained from Ψ_m as $m \rightarrow 0$.

For simplicity, consider first the case $h = 0$ (i.e., $q = \bar{q} = 0$). The solution to (B.1) is

$$\begin{aligned} \Psi_m(r) &= N_m r^{1-d} J_{d-1}(2mr) \\ &= N_m m^{d-1} \sum_{k=0}^{\infty} \frac{(-1)^k (mr)^{2k}}{k!(k+d-1)!} \end{aligned} \quad (\text{B.3})$$

We see that the tachyonic solutions (with $m^2 < 0$) are positive for all values of $r > 0$ and hence cannot possibly be normalizable. Thus the physical solutions have $m \geq 0$.

The scalar product of two wave functions is given by (3.15). In spherical co-ordinates, it reads

$$\begin{aligned} (\Psi_m, \Psi_{m'}) &= N_m N_{m'} \Omega_{2d} \int_0^\infty dr r^{2d-1} \Psi_m(r) \Psi_{m'}(r) \\ &= N_m N_{m'} \Omega_{2d} \int_0^\infty dr r J_{d-1}(2mr) J_{d-1}(2m'r) \end{aligned} \quad (\text{B.4})$$

where $\Omega_{2d} = \frac{2\pi^d}{(d-1)!}$ is the area of the unit sphere. The integral in (B.4) can be computed with the help of the standard result

$$\int x J_n(ax) J_n(bx) dx = \frac{x}{a^2 - b^2} [a J_{n+1}(ax) J_n(bx) - b J_n(ax) J_{n+1}(bx)] \quad (\text{B.5})$$

Making free use of distributions,

$$\begin{aligned} \int_0^\infty x J_n(ax) J_n(bx) dx &= \\ &= \lim_{\Lambda \rightarrow \infty} \frac{\Lambda}{a^2 - b^2} [a J_{n+1}(a\Lambda) J_n(b\Lambda) - b J_n(a\Lambda) J_{n+1}(b\Lambda)] \\ &= \frac{1}{\pi} \frac{1}{\sqrt{ab}} \lim_{\Lambda \rightarrow \infty} \left[\frac{\sin((a-b)\Lambda)}{a-b} - (-1)^n \frac{\cos((a+b)\Lambda)}{a+b} \right] \\ &= \frac{1}{a} \delta(a-b) - \frac{(-1)^n}{\sqrt{ab}} \delta'(a+b) \end{aligned} \quad (\text{B.6})$$

where we used $J_n(0) = 0$ for $n \in \mathbb{N}$ and the asymptotic form

$$J_n(x) \simeq \sqrt{\frac{2}{\pi x}} \cos \left[x - (2n+1)\frac{\pi}{4} \right] \quad \text{as } x \rightarrow \infty \quad (\text{B.7})$$

Thus, for $m, m' > 0$, we get

$$(\Psi_m, \Psi_{m'}) = \frac{\pi^d}{(d-1)!} \frac{N_m^2}{m} \delta(m - m') \quad (\text{B.8})$$

where the normalization constant is

$$N_m = \sqrt{\frac{(d-1)!}{\pi^d}} m \quad (\text{B.9})$$

which would mean that $\Psi_{m=0}(r) = 0$.

In fact, the massless $m = 0$ case of Equation (B.1) deserves special attention. The general solution is

$$\Psi_0 = Ar^{2(1-d)} + B \quad (\text{B.10})$$

where A and B are constants. We have

$$(\Psi_0, \Psi_m) = N_m \Omega_{2d} \int_0^\infty dr (Ar^{2-d} + Br^d) J_{d-1}(2mr) \quad (\text{B.11})$$

which can be computed as (B.4) above using

$$\begin{aligned} \int dx x^{n+1} J_n(x) &= x^{n+1} J_{n+1}(x) \\ \int dx x^{-n+1} J_n(x) &= -x^{-n+1} J_{n-1}(x) \end{aligned}$$

Thus for $m \neq 0$ we have

$$(\Psi_0, \Psi_m) = -AN_m \frac{\pi^d m^{\frac{d-3}{2}}}{(d-1)!(d-2)!} + B \sqrt{\frac{2}{\pi}} \frac{\Omega_{2d}}{2^{d+1}} \lim_{\Lambda \rightarrow \infty} \frac{N_m}{m^{3/4}} \Lambda^{d-1/2} \cos \left[m\Lambda - (2d+1)\frac{\pi}{4} \right] \quad (\text{B.12})$$

For $m \neq 0$, the second term is already zero so that the r.h.s. vanishes only if $A = 0$. Therefore, for $m = 0$, the solution is a constant

$$\Psi_0 = B \quad (\text{B.13})$$

Hence, if we take for $m \neq 0$ the wave function (B.3), with N_m given by (B.9), and for $m = 0$ we take (B.13), the wave functions fulfill (6.5). Actually, the wave functions Ψ_m do not have to be normalized to "1" as in (6.5); for arbitrary N_m the probability density of the mass eigenvalue m in the normalized wave packet

$$\Psi(r) = \int_0^\infty \rho(m) \Psi_m(r) dr \quad (\text{B.14})$$

would not be $\rho(m)$ but rather $\tilde{\rho}(m) = N_m \rho(m)$, such that $\tilde{\rho} \in L^2(\mathbb{R}^+)$. Hence, if we take $N_m = N/m^{d-1}$ with $N = (d-1)!B$, the limit of (B.3) for $m \rightarrow 0$ will be just (B.13).

The general wave functions with $q, \bar{q} \neq 0$ take the form (3.18):

$$\Psi_{j_1 \dots j_{d-m}}^{i_1 \dots i_q} = N \left(v^{i_1} \dots v^{i_q} r_{j_1}^* \dots r_{j_m}^* \right)_{||} \Psi_{m,h}(r) \quad (\text{B.15})$$

where $\Psi_{m,h}(r)$ is the solution of (B.1)

$$\Psi_{m,h}(r) = N_m r^{1-d-h} J_{d+h-1}(2mr) \quad (\text{B.16})$$

The scalar product of two such functions is

$$\begin{aligned} \left(\Psi_{j_1 \dots j_{d-m}}^{i_1 \dots i_q}, \Psi_{k_1 \dots k_q}^{j_1 \dots j_m} \right) &= N_m N_{m'} \int_0^\infty dr r^{1-h} J_{d+h-1}(2mr) J_{d+h'-1}(2m'r) \\ &\quad \times \int d\Omega \left(v_{i_1}^{i_1} \dots v_{i_q}^{i_q} v_{j_1}^{j_1} \dots v_{j_m}^{j_m} \right)_{||} \left(v_{k_1}^{k_1} \dots v_{k_q}^{k_q} v_{l_1}^{l_1} \dots v_{l_m}^{l_m} \right)_{||} \end{aligned} \quad (\text{B.17})$$

The integral is over a $2d$ dimensional Euclidean space with vector length squared

$$r^2 = \bar{r}r = \delta_{ij}^i r_i^i r_j^j \quad (\text{B.18})$$

so the solid angle integral in (B.17) yields

$$\begin{aligned} \int d\Omega \left(v_{i_1}^{i_1} \dots v_{i_q}^{i_q} v_{j_1}^{j_1} \dots v_{j_m}^{j_m} \right)_{||} \left(v_{k_1}^{k_1} \dots v_{k_q}^{k_q} v_{l_1}^{l_1} \dots v_{l_m}^{l_m} \right)_{||} \\ = c r^{2h} \delta_{qq'} \delta_{\bar{q}\bar{q}'} \left[\delta_{(i_1}^{(k_1} \delta_{i_2}^{k_2} \dots \delta_{i_q)}^{k_q)} \delta_{(j_1}^{(l_1} \delta_{j_2}^{l_2} \dots \delta_{j_m)}^{l_m)} \right] \end{aligned} \quad (\text{B.19})$$

Using (B.4) and (B.8), we get (3.24).

References

- [1] A. Schild, *Phys. Rev. D* **16** (1977) 1722.
- [2] F. Lizzi, B. Rai, G. Sparano and A. Srivastava, *Phys. Lett.* **182B** (1986) 326.
- [3] A.A. Zheltukhin, *Fisika Lett.* **46** (1987) 208; *Yad. Fiz.* **48** (1988) 487.
- [4] J. Gamboa and M. Ruiz Altaba, *La Guerra de Schild Supersimétrica*, in *Proceedings, Sexto Simposio Chileno de Física*, ed. M. Massman et al., Socialismo o Barbarie (1988) Santiago de Chile, p. 258.
- [5] J. Gamboa, C. Ramirez and M. Ruiz Altaba, *Quantum Null Superstrings*, preprint CERN-TH.5367/89 (April 1989), to appear in *Phys. Lett. B*.
- [6] R. Amorim and J. Barcelos Neto, *Z. Phys.* **C38** (1988) 643.
- [7] J. Barcelos Neto and M. Ruiz Altaba, *Superstrings with Zero Tension*, preprint CERN-TH.5294/89 (February 1989), to appear in *Phys. Lett. B*.
- [8] A.P. Balachandran, F. Lizzi and G. Sparano, *Nucl. Phys.* **B265** (1986) 608.
- [9] A.A. Zheltukhin, *Theor. Mat. Fiz.* **77** (1988) 377.
- [10] L. Brink and J.H. Schwarz, *Phys. Lett.* **100B** (1981) 310.
- [11] W. Siegel, *Phys. Lett.* **128B** (1983) 397.
- [12] S. Monaghan, *Phys. Lett.* **B178** (1986) 231.
- [13] For an in depth review, see M.B. Green, J.H. Schwarz and E. Witten, *Superstring Theory*, Cambridge University Press (1987) New York.
- [14] P. Ramond, *Phys. Rev. D* **3** (1971) 2415.
- [15] A. Neveu and J.H. Schwarz, *Nucl. Phys.* **B31** (1971) 80; *Phys. Rev. D* **4** (1971) 1109.
- [16] F. Gliozzi, J. Scherk and D. Olive, *Phys. Lett.* **65B** (1976) 282; *Nucl. Phys.* **B122** (1977) 253.
- [17] J.H. Schwarz and M.B. Green, *Nucl. Phys.* **B181** (1981) 502; *Phys. Lett.* **109B** (1982) 444; *Phys. Lett.* **136B** (1984) 367.
- [18] U. Marquard and M. Scholl, *Phys. Lett.* **B209** (1988) 434; Karlsruhe preprints KA-THEP-15 (March 1989), KA-THEP-4 (May 1989).
- [19] B. De Wit, J. Hoppe and H. Nicolai, *Nucl. Phys.* **B305** (1988) 545.
- [20] B. de Wit, M. Luescher, H. Nicolai, DESY preprint 88-162 (1988).
- [21] E. Bergshoeff, E. Sezgin and P.K. Townsend, *Phys. Lett.* **189B** (1987) 75; *Ann. Phys.* (N.Y.) **185** (1988) 330.
- [22] S. Deser and R. Jackiw, MIT preprint CTP 1673 (1989).
- [23] L. Brink, P. di Vecchia and P. Howe, *Nucl. Phys.* **B118** (1977) 76.
- [24] A. Karlhede and U. Lindström, *Class. Quantum Grav.* **3** (1986) L73.
- [25] C. Becchi, A. Roulet and R. Stora, *Phys. Lett.* **52B** (1974) 344; *Ann. Phys.* (N.Y.) **98** (1976) 287; I.V. Tyutin, Lebedev report FIAN 39 (1979) unpublished.
- [26] P. Howe, *J. Phys.* **A12** (1979) 393.
- [27] J. Gamboa and V. Tapia, *A generic action for strings*, Trieste preprint ICTP IC/87/265 (1987) unpublished.
- [28] J. Gamboa and M. Ruiz Altaba, *Phys. Lett.* **B205** (1988) 405.
- [29] D.J. Gross, *Phys. Rev. Lett.* **60** (1988) 1229.
- [30] J.J. Atick and E. Witten, *Nucl. Phys.* **B310** (1988) 291.
- [31] I. Klebanov and L. Susskind, *Nucl. Phys.* **B309** (1988) 175.
- [32] G. 't Hooft, Proc. Moscow Seminar on Quantum Gravity (USSR Academy of Sciences, May 1987).

- [33] G. Veneziano, *Europhysics Lett.* **2** (1986) 133; D. Amati, M. Ciafaloni and G. Veneziano, *Phys. Lett.* **216B** (1989) 41; *Int. J. Mod. Phys.* **3A** (1988) 1615.
- [34] P. Goddard, J. Goldstone, C. Rebbi and C.B. Thorn, *Nucl. Phys.* **B56** (1973) 109.
- [35] R. Casalbuoni, *Nuovo Cim.* **A33** (1976) 115.
- [36] S. Mandelstam, in *Strings and Unification*, eds. M.B. Green and D.J. Gross, World Scientific (Singapore) 1987.
- [37] D.J. Gross and P. Mende, *Phys. Lett.* **197B** (1987) 129; *Nucl. Phys.* **B303** (1988) 407.
- [38] P.A.M. Dirac, *Can. J. Math.* **2** (1950) 129; *Lectures on Quantum Mechanics*, Academic Press (1965) New York.
- [39] A. Hanson, T. Regge and C. Teitelboim, *Hamiltonian Constrained Systems*, Accademia dei Lincei (1976) Roma.
- [40] K. Sundermeyer, *Constrained Hamiltonian Systems*, Lecture Notes in Physics, Springer (1985) Heidelberg.
- [41] R. Floreani and R. Jackiw, *Phys. Rev. Lett.* **59** (1987) 1873.
- [42] P.A. Collins and R.W. Tucker, *Phys. Lett.* **64B** (1976) 207; *Nucl. Phys.* **B121** (1977) 307.
- [43] P. Howe, S. Penati, M. Pernici and P.K. Townsend, *Phys. Lett.* **B215** (1988) 555; CERN preprint TH.5305/89 (1989).
- [44] E.S. Fradkin and G.A. Vilkovisky, *Phys. Lett.* **55B** (1975) 224; I.A. Batalin and G.A. Vilkovisky, *Phys. Lett.* **69B** (1977) 309; E.S. Fradkin and T.E. Fradkina, *Phys. Lett.* **72B** (1978) 343.
- [45] M. Henneaux, *Phys. Reports* **126** (1985) 1. It includes very complete references.
- [46] C. Teitelboim, in *Proceedings of the La Plata Latin American School of Physics (ELAF 87)*, eds. L. Epele and C. Garcia Canal, World Scientific (1988) Singapore.
- [47] M. Kato and O.K. Ogawa, *Nucl. Phys.* **B212** (1983) 443.
- [48] S. Hwang, *Phys. Rev.* **D28** (1983) 2614.
- [49] C. Teitelboim, in *Superspace and Supergravity*, eds. S.W. Hawking and M. Roček, Cambridge University Press (1981), New York; *Phys. Lett.* **96B** (1980) 77; *Phys. Rev.* **D28** (1983) 297.
- [50] B.S. Skagerstam and A. Stern, *Nucl. Phys.* **B294** (1987) 636; T. Lee, *Ann. Phys. (N.Y.)* **183** (1988) 191.
- [51] J. Gamboa and V. Rivelles, *Spinning Self Dual Particles*, São Paulo preprint (1988); *BEST Formalism and the Quantization of Hamiltonian Systems with First Class Constraints*, to appear in *Proceedings, Conference honoring J. Douglas*, ed. T.M. Rassias, World Scientific (1989) Singapore.
- [52] E.S. Fradkin and G.A. Vilkovisky, *Quantization of relativistic systems with constraints: equivalence of canonical and covariant formalisms in quantum theory of gravitational field*, CERN preprint TH 2332 (1977) unpublished.
- [53] C. Teitelboim and M. Henneaux, *Ann. Phys. (N.Y.)* **143** (1982) 127.
- [54] I.A. Bandos and A.A. Zheltukhin, *Multidimensional Null Membranes: Absence of Critical Dimensions*, Khar'kov preprint (1989) to appear in *Yad. Fiz.*
- [55] J. Barcelos-Neto, J. Gamboa, C. Ramírez and M. Ruiz Altaba, in preparation.
- [56] See, e.g., K. Kikkawa and M. Yamasaki, *Phys. Lett.* **149B** (1987) 357; J. Laner, J. Mas and H.P. Nilles, München preprint MPI PAE/PTH 19/89 (1989).
- [57] P. Ramond and M. Ruiz Altaba, in *Proceedings of the II Mexican School of Particles and Fields*, eds. J.L. Lucio and A. Zepeda, World Scientific (1987) Singapore; and references therein.
- [58] H. Hata, K. Itoh, T. Kugo, H. Kunitomo and K. Ogawa, *Progr. Theor. Phys.* **78** (1987) 453.
- [59] A. Neveu and P. West, *Nucl. Phys.* **B293** (1987) 266.
- [60] S. Onyri and J. Stern, *Phys. Lett.* **B177** (1986) 335.
- [61] J. Labastida, *Phys. Rev. Lett.* **58** (1987) 531; *Massless Particles in Arbitrary Representations of the Lorentz Group*, IAS preprint IASSNS HEP 87/43 (1987).
- [62] S. Gradshteyn and I.M. Ryzhik, *Table of integrals, series and products*, Academic Press (1980) New York, p.971.