EUROPEAN ORGANIZATION FOR NUCLEAR RESEARCH

CERN-EP/89-93

3 August 1989

PHYSICS IN $p\bar{p}$ COLLISIONS

Felicitas PAUSS

CERN, Geneva, Switzerland

ABSTRACT

A review of experimental results obtained by UA1 and UA2 before the upgrading of the Proton-Antiproton Collider at CERN is presented. Topics covered in these lectures are the physics of the Intermediate Vector Bosons, heavy-flavour production, $B^0-\bar{B}^0$ mixing, the search for the top quark, and searches for physics beyond the Standard Model. We give preliminary results from the 1988/89 Collider run at CERN [UA1, upgraded UA2 and improved CERN $p\bar{p}$ Collider (ACOL)]. The physics potential of the full upgrading is discussed as well. Also, where available, results from the Fermilab Tevatron Collider are quoted. And finally, from the results obtained we infer the possible features of physics at future hadron colliders (LHC and SSC).

Lectures presented at the
XXVIII. Internationale Universitätswochen für Kernphysik
Schladming, Austria, 1989

1. INTRODUCTION

In a series of data-taking runs between 1982 and 1985, an integrated luminosity of close to 1 pb^{-1} was accumulated at the CERN $p\bar{p}$ Collider by each of the two major experiments, UA1 and UA2. The study of general event characteristics of hadron-hadron collisions at $\sqrt{s} = 630 \text{ GeV}$ has provided revealing information about the physics of the Standard Model (SM). A long list of important results covering a wide range of physics topics has come from the analyses of these data, e.g. the detailed measurements of jet production and fragmentation properties, the production and decay characteristics of the charged and neutral Intermediate Vector Bosons (IVBs), heavy-flavour production, B^0 - $\bar{B}^0$ mixing, etc. The main objective of all these studies is to know if the SM of electroweak and strong interactions is verified by the data, or if deviations are found which would hint at new physics.

Tests of the electroweak sector have been performed in measuring the SM parameters (Section 2). The strong sector (QCD) has been explored by studying jets, photons, the production of IVBs (Section 2), and the production of heavy flavour (Section 3). Within the minimal version of the SM there are three more elementary particles still waiting to be discovered, namely the ν_τ , the top (t) quark, and the neutral Higgs boson. Even though the ν_τ has not yet been discovered directly, there are indirect arguments for its existence, e.g. observation of the $W \rightarrow \tau\nu$ decay at the rate consistent with e- μ - τ universality of the weak charged-current coupling at $Q^2 = m_W^2$. The upper limit of 1.2×10^{-3} (90% CL) on $\text{BR}(B^0 \rightarrow \mu^+ \mu^- X)$ is strong indirect evidence that the t-quark exists. At the CERN $p\bar{p}$ Collider, the main sources of t-quarks are the decay $W \rightarrow t\bar{b}$, if kinematically allowed, and QCD production of $t\bar{t}$ pairs by gluon-gluon fusion or $q\bar{q}$ annihilation. At Fermilab, QCD $t\bar{t}$ production is the dominant production mechanism for all m_t . The search for the t-quark at CERN and Fermilab will be discussed in Section 4.

One of the most important discoveries still to be pursued is the experimental observation of the Higgs boson—which is believed to be the source of particle masses in the SM. The Higgs mass itself is, unfortunately, not predicted ($m_H \leq 1 \text{ TeV}/c^2$), and the only ‘safe’ experimental limit on the Higgs mass is $m_H > 15 \text{ MeV}/c^2$. There are two ways of searching for the Higgs at existing Colliders [1]: i) $p\bar{p} \rightarrow W^* \rightarrow WH + X$ (or $W \rightarrow W^*H$), i.e. the production of the Higgs with either a real or a virtual IVB; and ii) $p\bar{p} \rightarrow H + X$. In the first case the number of events expected is small, even for a very light Higgs. For example, for 1 pb^{-1} , one expects about four events ($H + \text{IVB}$) produced for $m_H < 1 \text{ GeV}/c^2$ at $\sqrt{s} = 630 \text{ GeV}$. A decay signature of $H \rightarrow e^+e^-$ ($\mu^+\mu^-$, $\pi^+\pi^-$, or K^+K^-) could lead to possible background-free detection of the Higgs, but, owing to the small branching ratios, a very high statistics W/Z data sample is required. At present $p\bar{p}$ Collider energies the dominant mechanism for Higgs production is through gluon-gluon fusion via a heavy quark loop. The Higgs decays into the heaviest fermion pair, i.e. into $b\bar{b}$, if $m_H > 10 \text{ GeV}/c^2$. Even though in this case the rates are large enough to observe the Higgs up to masses of about $50 \text{ GeV}/c^2$ (about $100 \text{ GeV}/c^2$) at $\sqrt{s} = 630 \text{ GeV}$ (1.8 TeV), the background from QCD jets is about a factor of 10^5 above a possible $H \rightarrow b\bar{b}$ signal. It therefore seems very unlikely that the Higgs will be discovered at existing hadron colliders.

There is a general consensus, however, that the SM is not expected to be our ultimate theory. Theorists have proposed numerous possible solutions to the well-known problems of the SM. For example, compositeness [2] could solve the Flavour problem, i.e. the understanding of the number of matter species. The Hierarchy problem—the understanding of the origin of the different particle masses—could be solved by supersymmetry [3]. It is expected that new physics should show up for $m \leq O(\text{TeV}/c^2)$. Unfortunately, the precise mass scale of this new physics is, as in the case of the Higgs, not known. Searches for physics beyond the SM can therefore provide limits on possible new mass scales of new physics (Section 5).

This review includes mostly data from the pre-ACOL era. These results have already been summarized and published in various conference proceedings. Therefore, instead of repeating details for the selected physics topics, the relevant results are stated and the interested reader is referred to the original publications.

The CERN $p\bar{p}$ Collider has been undergoing an important upgrading programme (addition of a new separate antiproton collector ring and six-bunch operation of the SPS). The performance reached during the first run of ACOL at the end of 1987 was far below design value ($L \approx 50 \text{ nb}^{-1}$ only). However, the following two runs (autumn 1988 and spring 1989) have been very successful, reaching an order of magnitude increase in luminosity. By end of June 1989, the total integrated luminosity delivered by the machine is 8.4 pb^{-1} , out of which 4.7 pb^{-1} (6.7 pb^{-1}) are recorded on tape by UA1 (UA2). Expectations from the present data samples will be discussed.

The electromagnetic (e.m.) calorimeters of the UA1 detector have been removed for the 1988 and 1989 runs in order to prepare for the installation of the new uranium-TMP calorimeter [4]. The muon detection capability has been improved by adding additional iron shielding in the forward region. The calorimetric measurements rely on the hadron calorimeters alone.

UA2 has been running since 1987 with the upgraded detector [5]. Full e.m. and hadronic calorimeter coverage has been achieved with the addition of new end-cap modules covering the angular regions $6^\circ < \theta < 40^\circ$ with respect to the beam directions. The electron identification is improved by a new central detector assembly consisting of a jet chamber, a silicon-pad detector, two layers of transition radiation detectors, and a scintillating-fibre detector.

The first physics run at $\sqrt{s} = 1.8 \text{ TeV}$ $p\bar{p}$ collisions at the Fermilab Tevatron Collider took place in 1987, resulting in a delivered luminosity of $\sim 50 \text{ nb}^{-1}$, out of which $\sim 30 \text{ nb}^{-1}$ have been collected by the Collider Detector at Fermilab (CDF) [6]. Published results from these data will be discussed in these lectures. The Tevatron started a long collider run in June 1988, which ended in June 1989, with excellent performance of the machine. The luminosity delivered is 9.5 pb^{-1} , with about 50% data-taking efficiency of the CDF detector (i.e. 4.7 pb^{-1}). Expectations from the present data sample will be discussed. The next Collider run at Fermilab is scheduled for the beginning of 1991.

Finally, some examples will be given of the possible features of physics at future hadron colliders (LHC and SSC), inferred from our present understanding of physics at hadron colliders.

2. PRODUCTION AND DECAY OF IVBs

2.1 IVB masses and Standard Model parameters

The data samples used in the UA1 analysis correspond to integrated luminosities of 136 nb^{-1} and 568 nb^{-1} at the Collider energies of $\sqrt{s} = 546 \text{ GeV}$ and 630 GeV , respectively. The UA2 results are based on 142 nb^{-1} and 768 nb^{-1} at the two energies. The full statistics of UA1 [7] and UA2 [8] result in a total of about 650 $W \rightarrow (e, \mu, \tau)\nu$ decays and about 100 $Z \rightarrow e^+e^-$ or $\mu^+\mu^-$ decays, observed in both experiments. The major fraction of the statistics is dominated by the $W \rightarrow e\nu$ and $Z \rightarrow e^+e^-$ channels.

The W and Z masses are the two parameters that can be directly measured by the Collider experiments. The most precise values for m_W and m_Z obtained from the electron channel are [9]

$$m_W = 82.7 \pm 1.0 \pm 2.7 \text{ GeV}/c^2 \quad (\text{UA1}),$$

$$m_W = 80.2 \pm 0.6 \pm 0.5 \pm 1.3 \text{ GeV}/c^2 \quad (\text{UA2}),$$

$$m_Z = 93.1 \pm 1.0 \pm 3.1 \text{ GeV}/c^2 \quad (\text{UA1}) ,$$

$$m_Z = 91.5 \pm 1.2 \pm 1.7 \text{ GeV}/c^2 \quad (\text{UA2}) .$$

The first errors quoted are statistical, followed by the systematic uncertainties. In the case of m_W , UA2 quotes the systematic uncertainties separately for transverse-mass determination and energy scale. The errors are already dominated by the energy-scale uncertainties of the e.m. calorimeters (about $\pm 3\%$ for UA1 and about $\pm 1.5\%$ for UA2).

From a data sample corresponding to 25.3 nb^{-1} integrated luminosity at $\sqrt{s} = 1.8 \text{ TeV}$, CDF obtained 22 $W \rightarrow e\nu$ events from an analysis with missing transverse energy $\cancel{E}_T$ larger than 25 GeV and with electron transverse energy larger than 15 GeV. A fit of the expected spectrum to the measured transverse mass distribution of the 22 events leads to [10]

$$m_W = 80.0 \pm 3.3 \pm 2.4 \text{ GeV}/c^2 \quad (\text{CDF}) ,$$

where the first error is statistical, and the second error, which is systematic, is dominated by the uncertainty in the absolute energy scale.

One of the important tests of the electroweak model is the need for radiative corrections. The relation between the radiative correction (Δr) and m_W and m_Z is given by

$$(1 - \Delta r) = (A^2/m_W^2)/[1 - (m_W^2/m_Z^2)] ,$$

where $A = (\pi\alpha/\sqrt{2}G_F)^{1/2} = 37.2810 \pm 0.0003 \text{ GeV}$. With this relation, we obtain [9]:

$$\Delta r = 0.036 \pm 0.100 \pm 0.067 \quad (\text{UA1}) ,$$

$$\Delta r = 0.068 \pm 0.087 \pm 0.030 \quad (\text{UA2}) .$$

A more precise value of Δr can be arrived at by using $\sin^2 \theta_w = 0.233 \pm 0.003 \pm 0.005$ from a world average of neutral-current neutrino experiments as additional input [11]:

$$\Delta r = 0.127 \pm 0.023 \pm 0.060 \quad (\text{UA1}) ,$$

$$\Delta r = 0.068 \pm 0.022 \pm 0.032 \quad (\text{UA2}) .$$

The experimental results are summarized in Fig. 1, where correlations between the uncertainties of the m_Z and m_Z - m_W measurements are shown. As can be seen from the error ellipse, the measurements of the IVB masses are not yet sensitive to the radiative corrections.

The one-loop radiative correction on the IVB masses has been calculated [12] to be

$$\Delta r = 0.0713 \pm 0.0013 ,$$

assuming that $m_t = 45 \text{ GeV}/c^2$ and $m_H = 100 \text{ GeV}/c^2$. The value of Δr decreases with increasing t-quark mass ($m_t \approx 250 \text{ GeV}/c^2$ leads to $\Delta r = 0$). Hence a precise measurement of Δr provides information on m_t . A global fit to all neutral-current data and to the W and the Z mass would correspond to $m_t < 180 \text{ GeV}/c^2$ (90% CL) for $m_H = 100 \text{ GeV}/c^2$ and assuming that only three fermion families exist [13].

2.2 IVB production cross-sections and e- μ - τ universality

Figures 2a,b show the UA1 and UA2 values for $\sigma \cdot \text{BP}(\Psi \rightarrow \ell \nu)$ and $\sigma \cdot \text{BR}(Z \rightarrow \ell^+ \ell^-)$ at $\sqrt{s} = 546$ GeV and at $\sqrt{s} = 630$ GeV. The experimental errors are dominated by systematic uncertainties in the luminosity in the case of $W \rightarrow e \nu$ and by statistics in the case of $Z \rightarrow e^+ e^-$ [9]. Also shown are the theoretical predictions and their error bands on the total W and Z production cross-sections taken from Ref. [14]. The leptonic branching ratios for $W \rightarrow \ell \nu$ and $Z \rightarrow \ell^+ \ell^-$ have been calculated for $m_t > (m_W - m_b)$. The theoretical predictions are affected by several uncertainties, e.g. i) various choices of structure functions and Q^2 -scale, ii) uncertainty in $\text{BR}(W \rightarrow \ell \nu)$ and $\text{BR}(Z \rightarrow \ell^+ \ell^-)$ due to the as yet unknown t-quark mass. The sensitivity of the experimental results to the t-quark mass and to the choice of structure functions [15] is illustrated in Figs. 3a,b.

The W production cross-section at the Tevatron energy of 1.8 TeV is expected to be about a factor of 3 larger compared with the CERN Collider energy of 0.63 TeV. CDF has published the measurement of $\sigma \cdot \text{BR}(W \rightarrow e \nu)$ obtained from a data sample of $25 \times 3 \text{ nb}^{-1}$ integrated luminosity. Using the 22 observed $W \rightarrow e \nu$ events and correcting for background and efficiency, CDF obtains [10]

$$\sigma \cdot \text{BR} = 2.6 \pm 0.6 \pm 0.5 \text{ nb}.$$

This result is shown together with the UA measurements in Fig. 4. The expected increase in W production is observed.

The UA1 data provide the first experimental verification of the universality of weak charged- and neutral-current coupling at $Q^2 = m_W^2$ (m_Z^2). Defining the weak charged-coupling constants by $(g_i/g_j)^2 = \Gamma(W \rightarrow \ell_i \nu_i)/\Gamma(W \rightarrow \ell_j \nu_j)$, and similarly for the weak neutral-coupling constants k, UA1 obtains [9]

$$g_\mu/g_e = 1.00 \pm 0.07 (\text{stat.}) \pm 0.04 (\text{syst.}),$$

$$g_\tau/g_e = 1.01 \pm 0.10 (\text{stat.}) \pm 0.06 (\text{syst.}),$$

$$k_\mu/k_e = 1.02 \pm 0.15 (\text{stat.}) \pm 0.04 (\text{syst.}).$$

2.3 IVB transverse-momentum distribution

In the framework of the QCD improved Drell-Yan model for quark-antiquark annihilation, the radiation of gluons from the incoming partons gives rise to a non-zero IVB transverse momentum and leads to the associated emission of high transverse-momentum jets. The normalized W transverse-momentum distributions obtained by UA1 and UA2 are shown in Fig. 5 [16]. The UA2 results are shown only for $p_T^W > 15$ GeV/c. The experimental data are compared with the theoretical predictions of Ref. [14]; also shown are the theoretical uncertainties indicated by the hatched band. The measurements agree well with the QCD prediction, except possibly at the very highest values of p_T^W . For $p_T^W > 60$ GeV/c, the theory predicts (0.8 ± 0.3) events for a total of three events observed (one event by UA2 and two events by UA1). The UA1 events have been discussed in detail in Ref. [17]. These events contain two high- p_T jets with an invariant mass compatible with the W mass, which makes their interpretation in terms of QCD correction to the W production somewhat unlikely. The analyses of the large new data samples at CERN and Fermilab are expected to clarify the question of a possible excess of high- p_T W's.

Figure 6 displays the inclusive p_T cross-sections for jets [18], direct photons [19], low-mass Drell-Yan $\mu^+ \mu^-$ pairs [20], and W production [17], as measured by UA1. The QCD predictions are

from Stirling [21], scaled by 1.5 for the jet cross-section, from Aurenche et al. [22] for direct photons, and the line describing the W cross-section is from Altarelli et al. [14]. For $p_T^W, p_T^\gamma \geq 50$ GeV/c, the effect of finite W and Z masses are reduced relative to photon production, and the absolute production cross-sections for W and γ become comparable. This is a direct manifestation of electroweak unification, the ratio of W and γ coupling to quarks being $\sin \theta_w$.

2.4 Measurement of α_s using W + jet data

Jets produced in association with the W provide a means of measuring α_s . To first approximation, the yield of W + 1-jet events is proportional to α_s . UA2 used the measured one-jet to zero-jet ratio by comparing this experimental value with the QCD-predicted value obtained from Monte Carlo simulation of W production [23]. Requiring that the experimental ratio R_{exp} be reproduced by the Monte Carlo ratio $R_{\text{MC}}(\alpha_s)$, the following value for α_s was obtained:

$$\alpha_s(m_W^2) = 0.13 \pm 0.03 (\text{stat.}) \pm 0.03 (\text{syst.1}) \pm 0.02 (\text{syst.2}),$$

where syst.1 represents the experimental systematic error and syst.2 is an estimate of the theoretical uncertainty. This measurement represents the first α_s determination that is (almost) free of K-factor effects in hadronic collisions. The result of this analysis is compared with other experimental determinations of α_s in Fig. 7 [23].

2.5 Number of light neutrino species

The measured ratio $R = \sigma_W \cdot \text{BR}(W \rightarrow \ell \nu) / \sigma_Z \cdot \text{BR}(Z \rightarrow \ell^+ \ell^-)$ depends on the t-quark mass and on the number of neutrino species via the branching ratios. The systematic measurement error largely cancels in the ratio R, and hence the available Z statistics limits the current precision on R. Combining all the available data of UA1 and UA2 (assuming lepton universality) results in [8, 24]

$$R = 8.4^{+1.2}_{-0.9},$$

$$R < 10.1 \quad \text{at 90\% CL.}$$

The theoretical prediction for R depends on: i) the ratio of the total production cross-section $R_0 = \sigma_W / \sigma_Z$ (an important source of uncertainty comes from the choice of structure functions; recent calculations of R_0 give values between 3.0 and 3.4 with a typical error of 0.15); ii) the partial width of the IVBs, which are precisely determined by the SM; iii) the total decay widths of W and Z, which depend on the t-quark mass and on the number of light neutrino species (N_ν).

The combined (UA1 and UA2) limit on N_ν as a function of m_t is shown in Fig. 8 [24], using the conservative theoretical input for $\sigma_W / \sigma_Z = 3.13$. Assuming that there is no W decay into fourth-generation leptons and that $m_\nu < m_Z/2$, one obtains

$$N_\nu < 5.7 \quad (90\% \text{ CL}).$$

If a new heavy charged lepton exists with a mass corresponding to the UA1 lower limit of 41 GeV/c², the limit obtained becomes less restrictive by one neutrino type. The data are not yet accurate enough to place an upper limit on the t-quark mass.

2.6 Outlook

Although the production cross-sections for W and Z increase with increasing $\sqrt{s}$, one expects that precision measurements in the W,Z sector will become more difficult at higher $\sqrt{s}$. For example, the

factor of ~ 3 increase of σ_W and σ_Z at the Tevatron compared with that at CERN has to be compared with about a factor of 10 increase in QCD two-jet production ($p_T^{\text{jet}} \approx 40 \text{ GeV}/c$), thus resulting in a worse signal-to-background ratio at the Tevatron. In addition, the mean p_T^W at the Tevatron is expected to be almost twice that at CERN, thus significantly broadening the Jacobian peak observed in $W \rightarrow e\nu$ decay, and affecting the precision measurement of m_W .

From an integrated luminosity collected during the 1988/89 runs of 4.7 pb^{-1} for UA1, 6.7 pb^{-1} for UA2, and 4.7 pb^{-1} for CDF, a total of 1600 (5000) reconstructed $W \rightarrow e\nu$ decays and 200 (500) reconstructed $Z \rightarrow e^+e^-$ in UA2 (CDF) can be expected. UA1 (CDF) will have collected about 500 (2500) $W \rightarrow \mu\nu$ and about 100 (200) $Z \rightarrow \mu^+\mu^-$ decays.

With the existing data sample and the Collider runs in ≥ 1990 , it should be possible to perform detailed tests of the SM, e.g. make a precise measurement of the W mass.

Using 10 pb^{-1} integrated luminosity UA2 estimated the expected error on the W and Z mass determination to be [25]

$$\begin{aligned}\delta m_W &\approx \pm 0.22 \text{ (stat.)} \pm 0.20 \text{ (syst. 1)} \pm 0.80 \text{ (syst. 2)} \text{ GeV}/c^2 \\ \delta m_Z &\approx \pm 0.25 \text{ (stat.)} \pm 0.92 \text{ (syst. 2)} \text{ GeV}/c^2\end{aligned}$$

where the errors are due to statistics (stat.), method (syst. 1) and calibration (syst. 2), assuming an uncertainty of $\pm 1\%$ on the absolute energy scale. This last error cancels in the measurement of $R = m_W/m_Z$, the expected precision on this ratio is

$$\delta(m_W/m_Z) \approx \pm 0.004.$$

The precise measurement of m_Z (to $\pm 50 \text{ MeV}/c^2$) expected from LEP allows to calibrate the mass scale of the hadron collider experiments. For UA2 this leads to an overall error on m_W (adding all contributions in quadrature) of

$$\delta m_W \approx \pm 0.37 \text{ GeV}/c^2.$$

UA1 estimated the error on m_W , assuming 50 pb^{-1} and an absolute energy scale uncertainty of $\sim 0.5\%$ obtainable with the uranium-TMP calorimeter, to be

$$\delta m_W \approx \pm 0.10 \text{ (stat.)} \pm 0.15 \text{ (syst.) GeV}/c^2,$$

again assuming a precise Z -mass value from LEP. This accuracy on m_W is unlikely to improve significantly until LEP operates above the threshold for W pair production.

At LHC and SSC energies, about 10^9 W 's and Z 's will be produced per year at a luminosity of $10^{33} \text{ cm}^{-2} \text{ s}^{-1}$. This has to be compared with $\sim 10^7$ Z 's produced in three years of running at LEP with $L = 1.7 \times 10^{31} \text{ cm}^{-2} \text{ s}^{-1}$, and $\sim 10^4$ W^+W^- at LEP 200. Is it possible to use the LHC/SSC to search for rare W and Z decays, either to probe new physics or to perform precision measurements of the SM? For the Z , the very unfavourable signal-to-background ratio at hadron colliders is to be contrasted with the complete purity of the signal at e^+e^- Z factories; thus, looking for rare decays will very likely be the domain of LEP. However, observing the conventional decays $Z \rightarrow e^+e^-$ and $\mu^+\mu^-$ at hadron colliders is of great importance for testing perturbative QCD.

The same argument holds for the standard W -decays. Note that the calculations depend on parton distributions at small $x \approx 0.002$ [compared with $x \approx 0.15$ (0.04) at CERN (Fermilab) energies].

With $\sim 10^9$ W events produced per year, a branching ratio of $BR \geq 10^{-6}$ – 10^{-7} can in principle be reached. Looking for hadronic decay modes seems rather discouraging owing to the overwhelming QCD-jet background. Leptonic decay modes are detectable, but it should be remembered that the W rapidity distribution is flat for $|y| \leq 4$ at SSC energies. For more details on possible detection of rare W-decays see Ref. 26.

It is clear that the sensitivity for observing interesting purely leptonic decay modes will be limited by large background — therefore rather challenging requirements for the detector performances are necessary.

3. HEAVY-FLAVOUR PRODUCTION

3.1 Inclusive b-quark production

At the CERN Collider, the predicted cross-sections for charm and beauty production are about $100 \mu\text{b}$ and $10 \mu\text{b}$, respectively, according to QCD estimates. These cross-sections need to be extracted from a large background of hadronic production due to $gg \rightarrow gg$. Furthermore, it is necessary to distinguish between charm and beauty production, which are expected to have comparable cross-sections at large p_T . Charm and beauty quarks can be tagged by the presence of one or more muons coming from the semileptonic decay of the heavy quarks. These muons are expected to be embedded in hadronic jets.

Inclusive muon production has been studied in UA1 for $p_T^\mu > 6 \text{ GeV}/c$ and within a pseudorapidity range $|\eta| < 1.5$ [27]. Figure 9 shows the p_T spectrum for muons after the decay background has been subtracted. The decays of charged π and K mesons amount to about 70% at $p_T^\mu = 6 \text{ GeV}/c$ and decrease to about 20% at $p_T^\mu = 20 \text{ GeV}/c$. The solid curve is the prediction for the sum of all expected contributions listed in Fig. 9. The agreement with the data shown in this figure is very good. In the interval of $10 < p_T^\mu < 15 \text{ GeV}/c$, heavy-flavour production is expected to be the dominant contribution. The harder fragmentation of the b-quark, compared with charm, leads on the average to a harder p_T spectrum of muons; thus $b\bar{b}$ is favoured over $c\bar{c}$ once a p_T cut is applied to the muons. An independent estimate of the beauty fraction is made by fitting the distribution of p_T^{rel} , the transverse momentum of the muons with respect to the nearest jet axis [27]. Figure 10 shows the p_T^{rel} distribution together with the fitted contributions from beauty, charm, and decay background, which for single-muon events results in

$$b\bar{b}/(b\bar{b} + c\bar{c}) = (76 \pm 12)\% .$$

The b-quark production cross-section has been evaluated using four different data sets [27]:

- J/ψ events: $p_T(J/\psi) > 5 \text{ GeV}/c$;
- low-mass dimuons: $2m_\mu < m_{\mu\mu} < 6 \text{ GeV}/c^2$; $p_T^{\mu 1} > 3 \text{ GeV}/c$; $p_T^{\mu 2} > 3 \text{ GeV}/c$;
- high-mass dimuons: $m_{\mu\mu} > 6 \text{ GeV}/c^2$; $p_T^{\mu 1} > 3 \text{ GeV}/c$; $p_T^{\mu 2} > 3 \text{ GeV}/c$;
- single muons: $p_T^\mu > 6 \text{ GeV}/c$.

The measured lepton cross-sections $\sigma(\mu)$ are related to the cross-sections for producing b-quarks, with quark transverse momentum above p_T^{min} , by the following expression:

$$\sigma(p_T^b > p_T^{\text{min}}) = \sigma(\mu)[\sigma_{\text{MC}}(p_T^b > p_T^{\text{min}})/\sigma_{\text{MC}}(\mu)] ,$$

where the cross-sections with subscript ‘MC’ are evaluated with the ISAJET Monte Carlo program [28]. The b-quark cross-section is then quoted for $p_T^b > p_T^{\text{min}}$, where p_T^{min} is defined in such a way that 90% of the muon events have $p_T^b > p_T^{\text{min}}$. Figure 11 shows the resulting inclusive b-quark cross-sections for six values of p_T^{min} , and for $|y^b| < 1.5$. The solid curve is the next-to-leading-order calculation of Nason et al. [29], and the dashed line corresponds to the full ISAJET prediction. The

level of agreement between the cross-section measurements made with quite different techniques (using all available μ data samples) and the theoretical predictions are very satisfactory.

The total cross-section for beauty production is obtained by normalizing the $O(\alpha_s^3)$ prediction to the data and extrapolating to $p_T^{\min} = 0$. Excluding the high-mass dimuon measurement as well as the measurement for $p_T^{\min} > 15$ GeV/c results in [27]

$$\sigma(p\bar{p} \rightarrow b \text{ or } \bar{b}, |y| < 1.5) = 14.7 \pm 4.7 \mu\text{b}.$$

Finally, extrapolating to all rapidities gives

$$\sigma(p\bar{p} \rightarrow b\bar{b}) = 10.2 \pm 3.3 \mu\text{b}.$$

This can be compared with a recent theoretical calculation [30], which gives

$$\begin{aligned} \sigma(p\bar{p} \rightarrow b\bar{b}) &= (12 \pm 7) \mu\text{b} & \text{for } m_b &= 5 \text{ GeV}/c^2, \\ \sigma(p\bar{p} \rightarrow b\bar{b}) &= (19 \pm 10) \mu\text{b} & \text{for } m_b &= 4.5 \text{ GeV}/c^2. \end{aligned}$$

Within the large experimental and theoretical errors, there is good agreement between the QCD prediction and the UA1 data.

3.2 B^0 - $\bar{B}^0$ oscillations

Evidence for B^0 - $\bar{B}^0$ oscillations has been reported by UA1 from the observation of an apparent excess of like-sign dimuon events [31]. Oscillation in the B_d^0 - $\bar{B}_d^0$ system has been observed by ARGUS and CLEO [32].

An experimentally accessible quantity that measures the degree of oscillations at the Collider is the fraction of wrong-sign decays:

$$\chi = P(B^0 \rightarrow \bar{B}^0 \rightarrow \mu^-) / P(B^0 \rightarrow \mu^\pm).$$

The experimental result obtained by UA1 yields

$$\chi = 0.121 \pm 0.047,$$

which is an average over all beauty states. Oscillations can occur for the neutral-meson states B_d^0 and B_s^0 . The corresponding mixing parameters χ_d and χ_s are related to χ via the expression

$$\chi = f_d \chi_d + f_s \chi_s,$$

where $f_{d(s)}$ is the fraction of beauty quarks hadronizing into $B_{d(s)}$ mesons. To obtain a limit on χ_d and χ_s from UA1 data, the values $f_d = 0.36$ and $f_s = 0.18$ have been used, based on the ratio of K^+/π^+ measured at the CERN Intersecting Storage Rings (ISR) [33]. Figure 12a shows the resulting ± 1 standard-deviation band obtained by UA1, compared with the weighted average of the ARGUS and CLEO results ($\chi_d = 0.162 \pm 0.039$). Figure 12b shows the corresponding combined 90% CL limit. The line in Fig. 12b indicates the upper limit on χ_d as a function of χ_s , which can be derived from existing limits on the Kobayashi-Maskawa (KM) matrix elements, assuming that we have a SM with three families. Combining this constraint with the experimental limits, only a small allowed region in the χ_s - χ_d plane remains, indicating almost maximal mixing in the B_s^0 channel [34].

4. SEARCH FOR HEAVY QUARKS (t , b')

In the framework of the SM, the t -quark should exist and its mass should be smaller than about $200 \text{ GeV}/c^2$ in order to be consistent with low-energy data. Thus, taking into account the lower limit on m_t from TRISTAN [35], the t -quark mass is expected to be in the range

$$28.0 < m_t \leq 180 \text{ GeV}/c^2.$$

An indirect lower mass limit of about $50 \text{ GeV}/c^2$ has also been obtained from analyses of $B^0-\bar{B}^0$ mixing [36].

The most promising channel for finding the t -quark at the CERN Collider is the decay $W \rightarrow t\bar{b}$, followed by the semileptonic decay of the t -quark. This channel dominates t -quark production at $\sqrt{s} = 630 \text{ GeV}$ if m_t is in the range $40\text{--}70 \text{ GeV}/c^2$. For m_t outside this range, direct $t\bar{t}$ production is the dominant process. The t -quark production cross-section for $t\bar{b}$ can be derived from the measurement $\sigma \cdot \text{BR}(W \rightarrow \ell\nu)$, taking into account the m_t dependence of the branching ratio for $W \rightarrow t\bar{b}$ and QCD corrections to the partial decay widths. The $t\bar{t}$ production cross-section can be estimated, using perturbative QCD, at the next-to-leading order and taking into account theoretical uncertainties. This leads, for example — assuming that $m_t = 40 \text{ GeV}/c^2$ — to $\sigma(t\bar{b}) = 1.41 \text{ nb}$ and $\sigma(t\bar{t}) = 0.64 \text{ nb}$ at $\sqrt{s} = 0.630 \text{ TeV}$. The production through the decay $Z \rightarrow t\bar{t}$ has a comparatively negligible cross-section and is not considered here.

UA1 has searched for heavy quarks using the distinctive signature of the semileptonic decay modes in the e and μ channels [37]. The experimental statistics available ($\sim 0.7 \text{ pb}^{-1}$) allowed a systematic investigation of the region $m_t \leq 45 \text{ GeV}/c^2$. A signal for the t -quark was searched for in the isolated $e/\mu + \geq 1\text{-jet}$ channel. The characteristic topology of a t -quark decaying semileptonically allows other sources of charged leptons to be discriminated, such as W , Z , Drell-Yan, J/Ψ , Υ , $b\bar{b}$, and $c\bar{c}$ — the main physics backgrounds to a new heavy-quark signal in the UA1 analysis.

Since no evidence for a t -quark signal was found in the data ($\sim 0.7 \text{ pb}^{-1}$), limits have been placed on the t -quark production cross-section in $p\bar{p}$ collisions. A lower limit on m_t is obtained by comparing this experimental limit with the theoretical prediction of the cross-section, which is a function of m_t . The contribution from $W \rightarrow t\bar{b}$ decays is fixed by the measured $\sigma \cdot \text{BR}(W \rightarrow e\nu)$, therefore the result is a limit on $\sigma(t\bar{t})$. UA1 has published lower limits of $44 \text{ GeV}/c^2$ and $32 \text{ GeV}/c^2$ on the masses of the t and b' (fourth-generation) quarks, respectively, at the 95% CL [37]. The results depend on the $O(\alpha_s)$ QCD prediction for the heavy-quark production cross-section. Using the $O(\alpha_s^3)$ calculation of Nason et al. [29] and the DFLM structure functions, Altarelli et al. [30] have revised these limits to

$$m_t > 41 \text{ GeV}/c^2 \quad (95\% \text{ CL}),$$

$$m_{b'} > 34 \text{ GeV}/c^2 \quad (95\% \text{ CL}),$$

as shown in Fig. 13. Since no contribution from $W \rightarrow t'\bar{b}'$ is included, the 95% CL upper limit on $\sigma(b'\bar{b}')$ is larger than the $\sigma(t\bar{t})$ limit.

To derive the b' -quark limit, it was assumed that the b' is lighter than the t - and t' -quarks, so the b' decays to a c -quark or a u -quark. Since there exists no lower bound on $V_{b'u}$ and $V_{b'c}$, these couplings could be very small, thus leading to long lifetimes of the b' . If the b' -quark has such a long lifetime, the quoted UA1 limit on $\sigma(b'\bar{b}')$ is no longer valid, because the selection of lepton candidates requires that the lepton track points back to the primary vertex of the $p\bar{p}$ interaction. However, such a long-lived b' provides an alternative signature: the b' , after travelling some

distance, decays and produces a secondary vertex of charged particles in the UA1 central drift chamber and a high-momentum jet in the calorimeter. This event topology has been used to search for long-lived heavy quarks. The result of this search is plotted in Fig. 14 and details of the analysis can be found in Ref. [38].

4.1 Outlook

One of the most important analysis topics of the new data of UA1, UA2, and CDF ($\sim 5 \text{ pb}^{-1}$, $\sim 7 \text{ pb}^{-1}$, and $\sim 5 \text{ pb}^{-1}$, respectively), is the search for the t -quark.

The $p\bar{p} \rightarrow t$ -quark production cross-section (from both $t\bar{b}$ and $t\bar{t}$) as a function of m_t at $\sqrt{s} = 0.63$ and 1.8 TeV is shown in Fig. 15. At CERN energies the $W \rightarrow t\bar{b}$ production dominates for $m_t \leq 70 \text{ GeV}/c^2$, which has the advantage of a simpler topology than for $t\bar{t}$ and a more precisely known production rate, since normalization to the measured $W \rightarrow e\nu$ cross-section is possible. At Fermilab, QCD $t\bar{t}$ production is the dominant production mechanism for all m_t . From Fig. 15 it is evident that the Fermilab Collider has a substantial advantage over the CERN Collider in t -production rate. Especially at large m_t (e.g. $m_t = 100 \text{ GeV}/c^2$), the $t\bar{t}$ cross-section increases by a factor of ~ 40 between $\sqrt{s} = 0.63$ and 1.8 TeV . If $m_t > m_W$, the interesting situation occurs that the t -quark decays into a real $W + b$ and the process $p\bar{p} \rightarrow t\bar{t} + X$ leads to final states with two real W 's and two b -quarks. The average p_T of each t -quark is $\sim m_t/2$. The W —because of its large mass—will carry most of the transverse momentum of the parent t -quark, and will therefore be produced at large p_T ($p_T^W \approx 45 \text{ GeV}/c$). The $p\bar{p} \rightarrow t\bar{t} + X \rightarrow W^+W^- b\bar{b} + X$ final state has therefore a very distinctive experimental signature: large transverse momentum p_T^t , highly unbalanced transverse momenta of the two t -quark decay products ($p_T^W \gg p_T^b$), and unbalanced W decay products, characteristic of large p_T^W production.

Preliminary results on the search for the t -quark using the new data samples are available from all three Collider experiments [39].

The new data of UA1 allow the t -quark to be searched for in the muon channel only, since the e.m. calorimeters have been removed during the 1988/89 data-taking period. Applying analysis techniques similar to those discussed in Section 4 to the 1988 data (1.3 pb^{-1}), UA1 does not observe a signal for t -quark production in the single-muon and dimuon data samples. Combining the additional data from 1988 ($\mu + \text{jet}$, $\mu\mu$) and data collected before 1988 ($\mu + \text{jet}$, $\mu\mu$, $e + \text{jet}$, and $e + \mu$), UA1 obtained a new, still preliminary, limit on m_t :

$$m_t > 56 \text{ GeV}/c^2 \quad (95\% \text{ CL}) ,$$

$$m_t > 60 \text{ GeV}/c^2 \quad (90\% \text{ CL}) ,$$

and for the b' -quark, the new UA1 lower limit on the b' mass is

$$m_{b'} > 38 \text{ GeV}/c^2 \quad (95\% \text{ CL}) .$$

Including the new data from the 1989 run, UA1 expects a sensitivity to t -quark masses approaching the kinematic limit of the W channel.

UA2 also presented preliminary results from the 1988 data [39] on the t -quark search in the e -channel (2.5 pb^{-1}). The selection requires missing transverse momentum $\not{p}_T > 15 \text{ GeV}$, $p_T^e > 11.5 \text{ GeV}$ and $15 < m_T(e\nu) < 50 \text{ GeV}/c^2$ for the t signal region. An increase in acceptance is expected if the $\not{p}_T$ is reduced to 10 GeV ; however, this requires more rejection power against background due to conversions and overlaps, i.e. tighter quality cuts on the electron candidate. So far the data are consistent with the expected background for both selections, where the dominant physics

contribution comes from $W + \text{jet}$ production. In 2.5 pb^{-1} of analysed data, no t -quark signal is observed in the mass range $30 < m_t < 60 \text{ GeV}/c^2$. The corresponding confidence limits on m_t are not yet quoted. UA2 expects a sensitivity to t -quark masses approaching the kinematic limit in the W -channel from the full data sample including the 1989 run.

A status report on the t -quark search by CDF has been presented [39]. The search strategy at the Fermilab Tevatron is different from that at the CERN $p\bar{p}$ Collider; this is because the $W \rightarrow t\bar{b}$ contribution is small and $t\bar{t}$ can be observed in many channels. Owing to the large cross-section and large integrated luminosity, CDF can afford to look for $e^\pm \mu^\mp$ events from $t\bar{t}$ decay. This channel provides a very clear signature, no background from $W + \text{jets}$, and negligible background from $b\bar{b}$ due to p_T and isolation cuts. In a data sample corresponding to 2.0 pb^{-1} , CDF searched for events with $E_T^\tau > 15 \text{ GeV}$ and $p_T^\tau > 12 \text{ GeV}/c$. The expectation from top in 2.0 pb^{-1} for this channel varies from 22 events ($m_t = 30 \text{ GeV}/c^2$) to 1.8 events ($m_t = 70 \text{ GeV}/c^2$). From zero events observed in the data, it was preliminarily concluded that $30 < m_t < 60 \text{ GeV}/c^2$ is unlikely. The corresponding confidence limits on m_t are not yet quoted. From the analysis of the full data sample ($\sim 5 \text{ pb}^{-1}$), CDF should have a sensitivity to t -quark masses up to $\sim 80 \text{ GeV}/c^2$, which could be extended to $\sim 120 \text{ GeV}/c^2$ with 20 pb^{-1} (expected by the end of 1991).

Heavy-quark production at the LHC/SSC will be dominated by pair-production through gg fusion [40]. The total production cross-section for heavy quark pairs as a function of m_Q is plotted in Fig. 16a at $\sqrt{s} = 20 \text{ TeV}$. The available centre-of-mass energy is an important parameter for heavy-quark production, e.g. for $m_Q = 0.5 \text{ TeV}/c^2$, the cross-section decreases by a factor of ~ 10 if $\sqrt{s}$ decreases from 20 to 10 TeV. Assuming that $Q \rightarrow qW$, the most promising decay mode to be detected is $Q\bar{Q} \rightarrow (W \rightarrow \ell\nu) + 4 \text{ jets}$. For one year of running at $L = 10^{33} \text{ cm}^{-2} \text{ s}^{-1}$, a mass reach of [40]

$$m_Q \approx 0.8 \text{ TeV}/c^2 \quad \text{at the LHC ,}$$

$$m_Q \approx 1.0 \text{ TeV}/c^2 \quad \text{at the SSC ,}$$

can be expected provided that the main background from $W + \text{jet}$ production is kept small. This can be achieved by cutting in a range of p_T^τ values, for which the signal exceeds the inclusive $W \rightarrow e\nu$ background, as can be seen in Fig. 16b.

5. SEARCHES FOR NEW PHYSICS

5.1 Search for heavy leptons and the number of neutrino species

An analysis of events with large missing transverse energy ($\vec{E}_T$) due to the production of one or more energetic neutrinos or weakly interacting particles, can extend our understanding of the SM, and is a sensitive way to search for new physics. A fourth generation of quarks and leptons would contain a charged heavy lepton L , together with its associated ν_L . Assuming that there is a universal strength of coupling to W and Z particles, and that $m_L < m_W$, the heavy lepton can be produced in W decays. One then looks for a process $p\bar{p} \rightarrow W + X$ (with $W \rightarrow L\nu$, and $L \rightarrow q\bar{q} + \nu$). The experimental signature would be events with one or more jets plus large $\vec{E}_T$. This decay pattern gives events that are qualitatively similar to $W \rightarrow \tau\nu$ ($\tau \rightarrow \text{hadrons} + \nu$), except for the phase-space factor and the typically larger invariant mass of the $q\bar{q}$ system. UA1 has used the $\vec{E}_T + \text{jet}$ data sample—which has led to the identification of $W \rightarrow \tau\nu$ decays—to place a limit on m_L [41]. Using the calculated heavy-lepton contribution and taking into account the existence of the additional neutrino (ν_L) coupling to the Z , it was found that

$$m_L > 41 \text{ GeV}/c^2 \quad (90\% \text{ CL}) .$$

Similarly, the rate of events from the process $p\bar{p} \rightarrow Z + X$ ($Z \rightarrow \nu\nu$) is proportional to the number of light-neutrino species coupling to the Z . If the Z is produced in association with an energetic jet (Z production at high p_T), such events are expected to appear as hadronic events with large values of E_T . The same data sample and method as were used for the heavy-lepton search have been employed by UA1 [41] to derive a limit on the number of light-neutrino species:

$$N_\nu \leq 10 \quad (90\% \text{ CL}) .$$

5.2 Searches for supersymmetric particles

As already mentioned, supersymmetric theories are attractive because they provide solutions for some of the fundamental problems of the SM. In supersymmetric theories, fermions and bosons are linked by a common symmetry, every particle having a supersymmetric partner, a ‘sparticle’. The coupling of these new particles is well defined and therefore production cross-sections can be calculated. Supersymmetric particles are pair-produced (e.g. $p\bar{p} \rightarrow \tilde{g}\tilde{g}$), they decay into lighter SUSY particles (e.g. $\tilde{g} \rightarrow q\bar{q}\tilde{\gamma}$), and the lightest SUSY particle (LSP) is expected to be stable and non-interacting, and thus will escape detection. In many models, the LSP is assumed to be the photino ($\tilde{\gamma}$).

At hadron colliders the dominant final states containing SUSY particles are $\tilde{g}\tilde{g}$, $\tilde{g}\tilde{q}$, and $\tilde{q}\tilde{q}$ [42]. In the case where $m_{\tilde{q}} > m_{\tilde{g}}$, the squark decays into $q\tilde{g}$ and the gluino decays into $q\bar{q}\tilde{\gamma}$; in the case where $m_{\tilde{q}} < m_{\tilde{g}}$, the decay modes are $\tilde{q} \rightarrow q\tilde{\gamma}$ and $\tilde{g} \rightarrow \tilde{q}\tilde{q}$ or $(q\bar{q})$. In both cases, the event topology consists of hadronic jets (from the fragmentation of outgoing q and $\bar{q}$) and E_T (from the outgoing $\tilde{\gamma}$ ’s). The number of jets depends upon the production sub-process and the $\tilde{q}$ and $\tilde{g}$ masses. When $m_{\tilde{q}} \ll m_{\tilde{g}}$, $\tilde{q}\tilde{q}$ pair-production dominates the production of strongly interacting sparticles, whereas $\tilde{g}\tilde{g}$ dominates if $m_{\tilde{q}} \gg m_{\tilde{g}}$.

UA1 has performed a search for gluinos and squarks [43] using the jet(s) + E_T signature (assuming a stable $\tilde{\gamma}$, which escapes detection). The domain of the $(m_{\tilde{g}}, m_{\tilde{q}})$ plane excluded by the UA1 analysis is shown in Fig. 17. In the case where $m_{\tilde{g}} \gg m_{\tilde{q}}$, the lower bound $m_{\tilde{q}} > 45 \text{ GeV}/c^2$ is obtained. Similarly, if $m_{\tilde{q}} \gg m_{\tilde{g}}$, the lower bound $m_{\tilde{g}} > 53 \text{ GeV}/c^2$ is found. For $m_{\tilde{g}} \approx m_{\tilde{q}}$, a more stringent limit of $\tilde{m} > 75 \text{ GeV}/c^2$ is obtained owing to the increase in the number of events expected from $\tilde{g}\tilde{q}$ production. The results are valid for photino masses up to $\sim 20 \text{ GeV}/c^2$. For $m_{\tilde{\gamma}} > 20 \text{ GeV}/c^2$, however, the squark and gluino mass limits decrease rapidly.

In the search for $\tilde{g}$ and $\tilde{q}$ by UA2 [44] two cases were considered: i) the photino is the LSP and is therefore stable, and ii) the photino decays into $\tilde{H}\gamma$, where $\tilde{H}$ escapes detection. In the first case, a somewhat lower mass limit for $\tilde{q}$ and $\tilde{g}$ has been obtained, compared with that from UA1. In the case of an unstable photino, the better granularity of the UA2 calorimeter allows an analysis of events containing two photons + jets in the final state. In this analysis, UA2 excludes gluino masses between 15 and 20 GeV/c^2 and squark masses between 9 and 46 GeV/c^2 . Again, assuming $m_{\tilde{g}} \approx m_{\tilde{q}}$, a limit of $\tilde{m} > 60 \text{ GeV}/c^2$ is obtained.

CDF has performed a search for gluinos and squarks using the jet(s) + E_T signature (assuming a stable $\tilde{\gamma}$, which escapes detection). The data sample used corresponds to 25.3 nb^{-1} of integrated luminosity [45]. The domain in the $(m_{\tilde{g}}, m_{\tilde{q}})$ plane excluded by the CDF analysis is shown in Fig. 18. The asymptotic mass limits are $m_{\tilde{g}} > 73 \text{ GeV}/c^2$ and $m_{\tilde{q}} > 74 \text{ GeV}/c^2$ at 90% CL. The lower-limit curve in Fig. 18 has been chosen where the acceptance dropped to 0.1%. The quoted results are valid for photino masses up to 30 GeV/c^2 in the high-mass region.

Supersymmetric leptons can be searched for in $W \rightarrow \tilde{\ell}\tilde{\nu}$ and $Z \rightarrow \tilde{\ell}^+\tilde{\ell}^-$ decays. In each case $\tilde{\ell} \rightarrow \ell\tilde{\gamma}$ is assumed to dominate, and one expects $\ell + E_T$ final states from W decays and $\ell^+\ell^- + E_T$ final states from Z decays. The former may be distinguished from the conventional $W \rightarrow \ell\nu$ decays

by its softer lepton spectrum and different angular distribution. The $Z \rightarrow \tilde{\ell}^+ \tilde{\ell}^-$ decay can be distinguished from conventional Drell-Yan background by the additional $\tilde{E}_T$ signature.

The Z data sample can also be used to extract a limit for the mass of the wino under the assumption that the decay $Z \rightarrow \tilde{W} \tilde{W}$ is kinematically possible. In this case one expects to observe again $\ell^+ \ell^-$ pairs in association with large $\tilde{E}_T$, under the assumption that $\tilde{W} \rightarrow e \tilde{\nu}$, where $\tilde{\nu}$ escapes detection (stable $\tilde{\nu}$ or $\tilde{\nu} \rightarrow \nu \tilde{\gamma}$).

Figures 19a,b show the 90% CL for $m_{\tilde{e}}$ and $m_{\tilde{W}}$ as a function of $m_{\tilde{\tau}}$ and $m_{\tilde{Z}}$, respectively, as obtained by UA2 [44]. In the $\tilde{W}$ case the UA2 result significantly improves the values reached in $e^+ e^-$ experiments so far. Using the $W \rightarrow e \nu$ data samples, UA1 [9] obtained a selectron mass limit of $\tilde{m} > 32 \text{ GeV}/c^2$, for $m_{\tilde{W}}$, $m_{\tilde{Z}} \gg m_{\tilde{e}}$, and $m_{\tilde{e}} = m_{\tilde{\tau}}$.

5.3 Searches for additional vector bosons

Additional vector bosons—presumably heavy—are predicted in many possible extensions of the SM [46]. These heavy particles (W', Z') would be detected in UA1 and UA2, depending on their mass, their coupling to quarks, and their branching ratio to leptons.

In the UA1 data sample, no $W \rightarrow e \nu$ and no $Z \rightarrow e^+ e^-$ candidates have been observed with electron-neutrino transverse mass or $e^+ e^-$ mass in excess of the expected distribution for Standard Model IVB decays [9]. These results have been used to set limits on the production and decay of more massive W(Z)-like objects. Normalizing the cross-section times branching ratio $(\sigma \cdot \text{BR}')_{\text{IVB}'}$ to those of the SM vector bosons results in limits on $m_{W'}$ and $m_{Z'}$ as a function of this ratio, as shown in Fig. 20. For standard coupling ($R = 1$), UA1 obtained the limits [9]

$$m_{W'} > 220 \text{ GeV}/c^2, \quad m_{Z'} > 173 \text{ GeV}/c^2 \quad (90\% \text{ CL}).$$

From the analysis of the final UA2 data sample, the general limits for $m_{W'}$ and $m_{Z'}$ as a function of $\lambda_q^2 \cdot \text{BR}_e$ (coupling times branching ratio) are derived, as shown in Fig. 21 [44]. For standard couplings, i.e. $\lambda_q^2 \cdot \text{BR}_e = 1$, UA2 obtains

$$m_{W'} > 209 \text{ GeV}/c^2, \quad m_{Z'} > 180 \text{ GeV}/c^2 \quad (90\% \text{ CL}).$$

Most models derived from superstring theories predict at least one extra Z' with $\lambda_q^2 \approx 0.2$ and BR_e varying from 0.3 to 1.0. For the resulting values of $\lambda_q^2 \cdot \text{BR}_e$ between 0.06 and 0.2, UA2 does not exclude any significant mass values.

5.4 Axigluon search

Axigluons are predicted by gauge theories based on the chiral colour group $SU(3)_L \times SU(3)_R$ [47]. In these theories, the $SU(3)_L \times SU(3)_R$ symmetry is broken down into the QCD $SU(3)_c$ symmetry at some scale that is comparable to the weak scale. A model-independent prediction of all chiral colour theories is the existence of a new massive colour octet of gauge bosons, the axigluons. The dominant decay channel of the axigluon is expected to be into two jets. The width of the axigluon depends on the number of particles into which it can decay. Within the model, the width is expected to be in the range $0.1 m_A < \Gamma < 0.4 m_A$.

A search for axigluons in the two-jet channel has been performed by UA1 [48]. Assuming that there is no interference between the normal QCD jet-production mechanism and the axigluon, the QCD and axigluon contributions have been added to obtain the total theoretical prediction in the presence of the axigluon. The data are fitted to the prediction using the maximum likelihood method. Figure 22 displays the limit, at 95% CL, on the production rate of axigluons in $p\bar{p}$ collisions at $\sqrt{s} = 630 \text{ GeV}$. It can be seen that the axigluon is excluded at the 95% CL for

$$150 \text{ GeV}/c^2 < m_A < 315 \text{ GeV}/c^2 .$$

This result has been generalized to obtain limits on the production rate of a heavy-state X decaying into two jets, for very small and very large widths, and also for the two spin cases (0 and 1). These limits are also shown in Fig. 22.

5.5 Quark substructure

As already mentioned, there are several theoretical motivations for compositeness. The proliferation of flavour may be a hint that quarks and leptons are composite. A possible manifestation of quark and lepton compositeness could be a new contact interaction having the new strong interaction scale Λ_c [49]. At the CERN Collider, the effect of a four-quark interaction term ($\sim 1/\Lambda_c$) would manifest itself in observed deviations from the QCD jet-production cross-section. Such deviations would be observable at scattering energies below Λ_c , and have been searched for by UA1, UA2 and CDF.

UA2 has concentrated on the overall normalization of the large- p_T jet cross-section [50]. Finite values of Λ_c would produce an excess of events at large p_T compared with the standard QCD prediction ($\Lambda_c = \infty$). Taking into account the theoretical and experimental uncertainties, UA2 obtained the limit

$$\Lambda_c > 370 \text{ GeV} \quad (95\% \text{ CL}) .$$

UA1 has used the angular distribution of high-mass jet pairs to set a limit on Λ_c [51]. Figure 23 shows the measured angular distribution, where χ is defined as $(1 + \cos \theta)$ over $(1 - \cos \theta)$, θ being the angle between the axis of the jet pair and the beam direction in the jet-jet rest frame [52]. A finite quark size would modify this angular distribution in such a way that more events are expected at small χ values. By varying the value of the parameter Λ_c in the fit to the measured angular distribution, and taking into account the systematic uncertainty on the jet-energy scale, UA1 obtained

$$\Lambda_c > 415 \text{ GeV} \quad (95\% \text{ CL}) .$$

To search for possible quark substructure, CDF has compared the measured inclusive jet cross-section at $\sqrt{s} = 1.8 \text{ TeV}$ with the predictions of leading-order QCD modified by the addition of a contact interaction with scale Λ_c [53]. Jet transverse energies up to 250 GeV have been observed, and the data with $E_T^{\text{jet}} < 130 \text{ GeV}$ were used to normalize the prediction. Taking into account Poisson fluctuations in the highest E_T -bins, the absence of jets with $E_T > 255 \text{ GeV}$, and uncertainties in structure functions and process-scales, CDF obtained the limit

$$\Lambda_c > 700 \text{ GeV} \quad (95\% \text{ CL}) .$$

5.6 Outlook

Searches for new physics have so far been fruitless at both CERN and Fermilab. However, in several cases, significant lower mass limits could be placed on hypothetical new particles.

What is the discovery potential of hadron colliders in the near and the distant future?

Searches for new particles using the new data samples of UA1, UA2, and CDF can be divided into two classes: a) where $\sqrt{s}$ does not play an important role, and b) where the production cross-section is strongly dependent on $\sqrt{s}$.

Examples for class (a) are

- fourth-generation of charged leptons searched for in W decays,
- scalar leptons ($\tilde{e}$, $\tilde{\mu}$, $\tilde{\nu}$) and $\tilde{W}$ searched for in W and Z decays.

One expects sensitivity up to the kinematic limit using present data. Figure 24 shows, as an example, the expected limits obtainable by UA2 for 10 pb^{-1} from the decay $Z \rightarrow \tilde{e}\tilde{e} \rightarrow ee\tilde{\gamma}\tilde{\gamma}$ [54].

Examples for class (b) are

- gluino and squark,
- additional vector bosons,
- compositeness scale Λ_c .

We have already seen the obvious benefit of having higher $\sqrt{s}$ in the search for gluinos and squarks. The mass reach at the Tevatron was about 20 GeV higher for about a factor of 30 less integrated luminosity. From the present data samples, one expects limits of the order of

$$\tilde{m} \approx 120 \text{ GeV}/c^2 \quad (\text{UA2}),$$

$$\tilde{m} \approx 200 \text{ GeV}/c^2 \quad (\text{CDF}).$$

Because of the absence of background in the decay of additional vector bosons (W' , Z') into leptons, hadron colliders make a large mass range accessible to experiments. Figure 25 shows the production cross-section for W' and Z' at $\sqrt{s} = 0.63$ and 1.8 TeV, assuming standard couplings to leptons and quarks, and $W' \rightarrow e\nu$ and $Z' \rightarrow ee$ decays. The corresponding sensitivity from the present data samples is expected to be

for $m_{W'}$: $\sim 300 \text{ GeV}$ (UA2) and $\sim 600 \text{ GeV}$ (CDF),

for $m_{Z'}$: $\sim 250 \text{ GeV}$ (UA2) and $\sim 520 \text{ GeV}$ (CDF).

An improvement in the limit on Λ_c of $\geq 650 \text{ GeV}$ from CERN data will be achievable. CDF expects to reach $\Lambda_c > 1.6 \text{ TeV}$ from present data samples, again demonstrating the clear benefit of higher $\sqrt{s}$.

Finally, we discuss the possible discovery potential of LHC/SSC with respect to two topics: supersymmetry and Z' .

The detection of supersymmetric particles has received a lot of attention in various physics studies dealing with future colliders [55]. Present limits on sparticles are not yet problematic for low-energy supersymmetry. Sparticle masses may still be too high to be produced at the CERN pp Collider, at the Tevatron, at LEP, and at the SLC. If this is the case, it will be necessary to await the next generation of high-energy colliders, probing parton-parton collisions at energies of the order of 1 TeV. However, we have to keep in mind that even if the sparticle masses are larger than a few hundred GeV, nearly all low-energy SUSY models imply the existence of at least a light Higgs scalar with mass less than $O(m_Z)$. Therefore, the non-observation of a neutral Higgs boson up to LEP 200 energies would put rather tight constraints on possible supersymmetric parameters [56].

Squarks and gluinos should be copiously produced at the LHC/SSC. For example, for a total integrated luminosity of $10^{40} \text{ cm}^{-2} = 10 \text{ fb}^{-1}$ (corresponding to one year of running at $L = 10^{33} \text{ cm}^{-2} \text{ s}^{-1}$), we expect $\sim 10^4$ gluino pairs to be produced at the SSC for $m_{\tilde{g}} = 1.5 \text{ TeV}$, or at the LHC for $m_{\tilde{g}} = 1 \text{ TeV}$. The sparticle signatures depend on the details of the assumed masses and decay patterns. The 'classical' signature of large E_T plus two or more high- E_T jets results from the assumption that $\tilde{q}$ and $\tilde{g}$ will decay directly to the LSP. This assumption has been used in the analysis of UA1, UA2 and CDF data. Although these signatures should remain valid at LHC and SSC energies, it has been emphasized that for large squark and gluino masses, direct decays to the LSP becomes less important [57]. Instead, the decays to heavier neutralinos and charginos become dominant. These cascade decays generally occur for both the produced gluinos and squarks,

and result in a large variety of possible final states, often containing W and Z bosons in a complex event topology.

Detailed studies of signal and backgrounds have been performed for the direct decays of sparticles into the LSP [58]. Figure 26 shows an example of the $\cancel{E}_T$ spectrum using one of the possible event selections at LHC energies. The resulting signal-to-background ratios are typically $\sim 10:1$ at large $\cancel{E}_T$. Taking into account uncertainties involved in these calculations, it was concluded that squark and gluino masses, up to $\sim 1 \text{ TeV}/c^2$ are potentially accessible for 10 fb^{-1} integrated luminosity. Extrapolating the LHC study to SSC energies, a discovery limit of $\sim 1.5\text{--}2 \text{ TeV}/c^2$ can be expected for the gluino and squark mass, because the signal cross-section increases much faster than the background cross-sections for higher $\sqrt{s}$. The analysis discussed above has been extended to study the feasibility of searching for $\tilde{g}$ and $\tilde{q}$ if the LHC runs at a luminosity of $\sim 5 \times 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$ [59]. It was concluded that, provided the experimental problems associated with running at high luminosity can be overcome, an extension of the discovery limit for $\tilde{g}$ and $\tilde{q}$ of masses $\sim 2 \text{ TeV}/c^2$ should be feasible at the LHC.

A study of gluino final states containing Z's from cascade decays was carried out for SSC energies [57]. It was concluded that it is possible to find event signatures that give promising results for signal-to-background ratios even for these complex decay modes.

Additional vector bosons are predicted in many possible extensions of the Standard Model. In order to study the discovery potential for an additional neutral gauge boson Z' at the LHC/SSC, a superstring-inspired model was used [60]. This model predicts one (or two) new Z' with masses in the range of $\leq 1 \text{ TeV}/c^2$. Figure 27 shows the predicted $\sigma \cdot \text{BR}(Z' \rightarrow \ell^+ \ell^-)$ for the minimal superstring model at LHC and SSC energies. Since no significant background is expected for the leptonic decay modes, it was concluded that a search can be made for Z' with masses up to 4 TeV (6 TeV) using the $Z' \rightarrow \ell^+ \ell^-$ decay channel at the LHC for 10 fb^{-1} (500 fb^{-1}) integrated luminosity. The corresponding discovery limit at the SSC is 7 TeV for 10 fb^{-1} integrated luminosity. This shows that the discovery potential for additional vector bosons is very high in hadron-hadron machines, and also demonstrates the benefit of having higher available centre-of-mass energies.

6. CONCLUSIONS

The data collected between 1982 and 1985 by the UA1 and UA2 experiments at the CERN $p\bar{p}$ Collider, corresponding to integrated luminosities of about 1 pb^{-1} , have provided many important physics results. Excellent agreement with the Standard Model has been observed. Searches for new physics have so far been fruitless; however, in several cases, significant lower mass limits could be placed on hypothetical new particles.

First results from CDF at the Fermilab Collider are in agreement with the expectations owing to the increase in $\sqrt{s}$, and no deviations from the Standard Model have been observed. The benefit of the larger $\sqrt{s}$ has been best demonstrated in searches for new physics ($m_{\tilde{g}}$, $m_{\tilde{q}}$, Λ_c).

From the data taken with ACOL and at Fermilab, it is expected that the Standard Model predictions can be tested more precisely, and that hopefully at least one of its missing elements—the top quark—will be found. The preliminary results on searches for the t-quark using a subsample of available data, lead to the conclusion that the t-quark must be very heavy compared with other known quarks and leptons, provided the Standard Model is correct and that no new physics influences the production and decay properties of the t-quark. The goal of discovering new physics still remains, but will in some cases be more easily accessible at the Tevatron and at LEP.

Acknowledgement

I wish to thank the Scientific Reports Editing and Text Processing Section at CERN for their competence and patience in preparing the final version of this paper.

REFERENCES

- [1] S.L. Glashow et al., Phys. Rev. **D18** (1978) 1724.
E. Eichten et al., Rev. Mod. Phys. **56** (1984) 579.
- [2] M.E. Peskin, Proc. Int. Symp. on Lepton and Photon Interactions at High Energies, Kyoto, 1985, eds. M. Konuma and C. Takahashi (Kyoto Univ., 1985), p. 714.
H. Harari, Proc. 5th Topical Workshop on Proton-Antiproton Collider Physics, Saint-Vincent, 1985, ed. M. Greco (World Scientific, Singapore, 1985), p. 429.
- [3] P. Fayet and S. Ferrara, Phys. Rep. **32** (1977) 249.
H.P. Nilles, Phys. Rep. **110** (1984) 1.
J. Ellis et al., Nucl. Phys. **B238** (1984) 453.
H.E. Haber and G.L. Kane, Phys. Rep. **117** (1985) 75.
- [4] J. Dowell (UA1 Collab.), Proc. 6th Topical Workshop on Proton-Antiproton Collider Physics, Aachen 1986, eds. K. Eggert et al. (World Scientific, Singapore, 1987), p. 419.
- [5] C. Booth (UA2 Collab.), *ibid.*, p. 381.
- [6] F. Abe et al., Nucl. Instrum. Methods Phys. Res., **A271** (1988) 387.
- [7] UA1 Collaboration papers:
G. Arnison et al., Phys. Lett. **122B** (1983) 103; **126B** (1983) 398; **129B** (1983) 273; **134B** (1984) 469; **147B** (1984) 241; **166B** (1986) 484; Europhys. Lett. **1** (1986) 327.
C. Albajar et al., Phys. Lett. **185B** (1987) 233.
- [8] UA2 Collaboration papers:
M. Banner et al., Phys. Lett. **122B** (1983) 476.
P. Bagnaia et al., Phys. Lett. **129B** (1983) 130; Z. Phys. **C24** (1984) 1.
J.A. Appel et al., Z. Phys. **C30** (1986) 1.
R. Ansari et al., Phys. Lett. **186B** (1987) 440.
- [9] C. Albajar et al. (UA1 Collab.), preprint CERN-EP/88-168 (1988), submitted to Z. Physik.
- [10] F. Abe et al. (CDF Collab.), Phys. Rev. Lett. **62** (1989) 1005.
- [11] H. Abramowicz et al., Phys. Rev. Lett. **57** (1986) 298.
J.V. Allaby et al., Z. Phys. **C36** (1987) 611.
D. Bogert et al., Phys. Rev. Lett. **55** (1985) 1969.
P. Reutens et al., Phys. Lett. **152B** (1985) 404.
- [12] F. Jegerlehner, Z. Phys. **C32** (1986) 425.
W.J. Marciano and A. Sirlin, Phys. Rev. **D22** (1980) 2695 and **D29** (1984) 945.
- [13] U. Amaldi et al., Phys. Rev. **D36** (1987) 1385.
J.V. Allaby et al., preprint CERN-EP/87-140 (1987).
G. Costa et al., Nucl. Phys. **B297** (1988) 244.
G. Gounaris and D. Schildknecht, preprint CERN-TH.4940 (1987).
- [14] G. Altarelli et al., Nucl. Phys. **B157** (1979) 461 and **B246** (1984) 12.
G. Altarelli et al., Z. Phys. **C27** (1985) 617.
- [15] R. Voss, Proc. Int. Symp. on Lepton and Photon Interactions at High Energies, Hamburg, 1987, eds. W. Bartel and R. Rückl (North-Holland Publ. Co., Amsterdam, 1988), p. 525.
- [16] P. Jenni, *ibid.*, p. 341.
- [17] C. Albajar et al. (UA1 Collab.), Phys. Lett. **193B** (1987) 389.
- [18] G. Arnison et al. (UA1 Collab.), Phys. Lett. **172B** (1986) 461.
- [19] C. Albajar et al. (UA1 Collab.), Phys. Lett. **209B** (1988) 385.
- [20] C. Albajar et al. (UA1 Collab.), Phys. Lett. **209B** (1988) 397.

- [21] W.J. Stirling, Proc. 6th Topical Workshop on Proton-Antiproton Collider Physics, Aachen, 1986, eds. K. Eggert, H. Faissner and E. Radermacher (World Scientific, Singapore, 1987), p. 301.
- [22] P. Aurenche et al., Nucl. Phys. **B297** (1988) 661; Phys. Lett. **140B** (1984) 87.
- [23] R. Ansari et al. (UA2 Collab.), Phys. Lett. **215B** (1988) 175.
- [24] C. Albajar et al. (UA1 Collab.), Phys. Lett. **198B** (1987) 271.
- [25] L. Di Lella, Proc. 7th Topical Workshop on Proton-Antiproton Collider Physics, Fermilab, 1988, eds. R. Raja et al. (World Scientific, Singapore, 1989), p. 849.
- [26] F.E. Paige, Proc. Workshop on Experiments, Detectors, and Experimental Areas for the Supercollider, Berkeley, USA, 1987, eds. R. Donaldson and M. Gilchriese (World Scientific, Singapore, 1989), p. 750.
- [27] C. Albajar et al. (UA1 Collab.), Z. Phys. **C37** (1988) 489; Phys. Lett. **186B** (1987) 237; **200B** (1988) 380; **213B** (1988) 405.
- [28] F.E. Paige and S.D. Protopopescu, Brookhaven National Laboratory report, BNL-38034 (1986).
- [29] P. Nason, S. Dawson and R.K. Ellis, Nucl. Phys. **B303** (1988) 607.
- [30] G. Altarelli et al., preprint CERN-TH.4978/88 (1988).
- [31] C. Albajar et al. (UA1 Collab.), Phys. Lett. **186B** (1987) 247.
- [32] H. Albrecht et al., Phys. Lett. **192B** (1987) 246.
H. Schröder, Proc. 24th Int. Conf. on High-Energy Physics, Munich, 1988, eds. R. Kotthaus and J.H. Kühn (Springer-Verlag, Berlin, Heidelberg, 1989), p. 73.
- [33] A. Breakstone et al., Phys. Lett. **135B** (1984) 510.
- [34] K. Eggert and H.G. Moser, Proc. 7th Topical Workshop on Proton-Antiproton Collider Physics, Fermilab, 1988, eds. R. Raja et al. (World Scientific, Singapore, 1989), p. 599.
- [35] T. Kamae, Proc. 24th Int. Conf. on High-Energy Physics, Munich 1988, eds. R. Kotthaus and J.H. Kühn (Springer-Verlag, Berlin, Heidelberg, 1989), p. 156.
- [36] J. Ellis, J.S. Hagelin and S. Rudaz, Phys. Lett. **192B** (1987) 201.
V. Barger, T. Hau and D.V. Nanopoulos, Phys. Lett. **194B** (1987) 312.
I.I. Bigi and A.I. Sanda, Phys. Lett. **194B** (1987) 307.
L.L. Chau and W.Y. Keung, UC Davis preprint, UCD-87-02 (1987).
H. Harari and Y. Nir, Stanford preprint SLAC-PUB-4341 (1987).
G. Altarelli and P.J. Franzini, preprint CERN-TH-4745/87 (1987).
H. Albrecht et al. (ARGUS Collab.), Phys. Lett. **192B** (1987) 245.
- [37] C. Albajar et al. (UA1 Collab.), Z. Phys. **C37** (1988) 504.
- [38] J. Kroll, Proc. 24th Int. Conf. on High-Energy Physics, Munich, 1988, eds. R. Kotthaus and J.H. Kühn (Springer-Verlag, Berlin, Heidelberg, 1989), p. 1416.
- [39] Seminar at CERN, Search for the top quark, 18 April 1989:
M. Della Negra (UA1),
L. Fayard (UA2),
G. Bellettini (CDF).
- [40] D. Froidevaux, Proc. Workshop on Physics at Future Accelerators, La Thuile, Italy and CERN, Geneva, Switzerland, ed. J. Mulvey, CERN 87-07 (1987), Vol. I, p. 61.
- [41] C. Albajar et al. (UA1 Collab.), Phys. Lett. **185B** (1987) 241.
- [42] S. Dawson et al., Phys. Rev. **D31** (1985) 1581.
- [43] C. Albajar et al. (UA1 Collab.), Phys. Lett. **198B** (1987) 261.
- [44] R. Ansari et al. (UA2 Collab.), Phys. Lett. **195B** (1987) 613.
- [45] F. Abe et al. (CDF Collab.), Phys. Rev. Lett. **62** (1989) 1825.

- [46] P. Langacker et al., Phys. Rev. **D30** (1984) 1470.
 E. Cohen et al., Phys. Lett. **165B** (1985) 76.
 M.B. Green, J.H. Schwarz and E. Witten, Superstring theory (Cambridge Univ. Press, Cambridge, 1987).
 F. del Aguila et al., Nucl. Phys. **B284** (1987) 530 and **B287** (1987) 419.
- [47] P.H. Frampton and S.L. Glashow, Phys. Lett. **190B** (1987) 157.
- [48] P.A. Sphicas, Proc. 7th Topical Workshop on Proton-Antiproton Collider Physics, Fermilab, 1988, eds. R. Raja et al. (World Scientific, Singapore, 1989), p. 239.
- [49] E. Eichten et al., Phys. Rev. Lett. **50** (1983) 811.
- [50] J.A. Appel et al. (UA2 Collab.), Phys. Lett. **160B** (1985) 349.
- [51] G. Arnison et al. (UA1 Collab.), Phys. Lett. **177B** (1986) 244.
- [52] J.C. Collins and D.E. Soper, Phys. Rev. **D16** (1977) 2219.
- [53] F. Abe et al. (CDF Collab.), Phys. Rev. Lett. **62** (1989) 613.
- [54] D. Froidevaux and P. Jenni, Proton-Antiproton Collider Physics, eds. G. Altarelli and L. Di Lella (World Scientific, Singapore, 1989), p. 323.
- [55] For a recent review, see
 F. Pauss, Proc. 24th Int. Conf. on High-Energy Physics, Munich 1988, eds. R. Kotthaus and J.H. Kühn (Springer-Verlag, Berlin, Heidelberg, 1989), p. 1275, and references therein.
- [56] R. Barbieri, Proc. Rencontres de Physique de la Vallée d'Aoste, La Thuile, Italy, 1987, ed. M. Greco (Editions Frontières, Gif-sur-Yvette, 1987), p. 523.
 J. Gunion et al., Univ. Calif. Davis report UCD-88-11 and LBL-25033 (1988).
 G.F. Giudice, Phys. Lett. **B208** (1988) 315.
- [57] R.M. Barnett et al., Proc. Workshop on Experiments, Detectors, and Experimental Areas for the Super collider, Berkeley, 1987, eds. R. Donaldson and H. Gilchriese (World Scientific, Singapore, 1989), p. 178.
 H. Baer and E. Berger, Phys. Rev. **D34** (1986) 1361.
 H. Baer et al., Phys. Rev. **D36** (1987) 96.
- [58] R. Batley, Proc. Workshop on Physics at Future Accelerators, La Thuile, Italy, and CERN, Geneva, Switzerland, ed. J. Mulvey, CERN 87-07 (1987), Vol. II, p. 109.
- [59] F. Pauss et al., The Feasibility of Experiments at High Luminosity at the Large Hadron Collider, ed. J. Mulvey, CERN 88-02 (1988), p. 79.
- [60] J. Ellis and F. Pauss, Proc. Workshop on Physics at Future Accelerators, La Thuile, Italy, and CERN, Geneva, Switzerland, ed. J. Mulvey, CERN 87-07 (1987), Vol. I, p. 80.
 J. Ellis, Proc. of the 1987 ICFA Seminar on Future Perspectives in High-Energy Physics, BNL, Upton, NY, USA, October 1987, ed. P.F. Dahl, BNL 52114, p. 117.

Figure captions

Fig. 1 Confidence contours (68%) in the ($m_Z - m_W$) versus m_Z plane, taking into account the statistical error (dashed line) and with statistical and systematic errors combined in quadrature (solid line). The shaded region is allowed by the average of recent low-energy measurements of $\sin^2 \theta_w$. The upper dashed-dotted curve (a) includes radiative corrections, the lower one (b) is without radiative corrections.

Fig. 2 Partial cross-sections versus c.m. energy: a) for $W \rightarrow \ell \nu$ production and b) for $Z \rightarrow \ell \ell$ production. The theoretical predictions and their error bands on the total W and Z production cross-sections are taken from Ref. [14], and the leptonic branching ratios for $W \rightarrow \ell \nu$ and $Z \rightarrow \ell \ell$ have been calculated under the assumption that $m_t > (m_W - m_b)$.

Fig. 3 Measurements and theoretical predictions for a) $W \rightarrow \ell \nu$ and b) $Z \rightarrow \ell \ell$ partial cross-sections as a function of m_t . The line labelled ‘1’ represents the predictions [14] used in Figs. 2a,b. The line labelled ‘2’ includes a 20% increase due to the BCDMS results on structure function parametrization [15]. The uncertainties are indicated by the error bars. The hatched band represents the measurements.

Fig. 4 Cross-section times branching ratio for $W \rightarrow e \nu$ versus E_{cm} measured by UA1, UA2 and CDF [10]. The prediction is from Ref. 14, adjusted for a W mass of $80 \text{ GeV}/c^2$. The dotted lines indicate the 1σ error limits for the theoretical curve.

Fig. 5 W transverse-momentum distributions from UA1 (full points) and UA2 (open points). Only data with $p_T^W > 15 \text{ GeV}/c$ are shown for UA2. The curves are QCD predictions as reported in Ref. [17], based on calculations of Ref. [14] and extrapolating for high p_T^W .

Fig. 6 Inclusive transverse-momentum distributions for W’s, jets, direct photons, and nearly on-shell photons from low-mass dimuons as measured in the UA1 experiment [9]. The curves are QCD predictions as described in the text.

Fig. 7 Recent experimental determinations of the strong coupling constant compared with second-order QCD predictions [23].

Fig. 8 The total number of light-neutrino species as a function of the t-quark mass, deduced from the combined UA1 and UA2 data.

Fig. 9 Inclusive transverse-momentum distribution for muons, after decay background subtraction. Predictions made with ISAJET are shown for comparison [27].

Fig. 10 Distribution of the transverse momentum relative to the nearby jet, for muons of $p_T > 10 \text{ GeV}/c$ in a sample of events with at least one jet of $E_T > 10 \text{ GeV}$ [27].

Fig. 11 Cross-section for b (or $\bar{b}$) quark production with transverse momentum above $p_T^{\min}$ and $|y| < 1.5$ [27]. The interpretation of the symbols is as follows: ($\blacksquare$) single muons, ($\blacktriangle$) low-mass dimuons, ($\circ$) high-mass dimuons, and ($\bullet$) J/ψ . Also shown are predictions from ISAJET, as well as from second- and third-order QCD calculations [29].

Fig. 12 Beauty-oscillation results in the χ_s - χ_d plane: a) one-standard-deviation bands, and b) combined 90% CL limit. The region below the line is allowed by the present constraints from the KM matrix, assuming three generations [34].

Fig. 13 The UA1 experimental upper limits (95% CL) on $t\bar{t}$ and $b'\bar{b}'$ production cross-sections in $p\bar{p}$ collisions at $\sqrt{s} = 0.63$ TeV. The shaded area is the predicted cross-section for heavy quarks, from Ref. [30].

Fig. 14 The region excluded at 95% CL by UA1 in the $m_{b'}-\tau_{b'}$ plane, for a long-lived b' -quark [38].

Fig. 15 The $p\bar{p} \rightarrow$ top production cross-section (from both $t\bar{t}$ and $t\bar{b}$) as a function of m_t at $\sqrt{s} = 0.63$ and 1.8 TeV.

Fig. 16 a) The total production cross-section for heavy-quark pairs as a function of the heavy-quark mass m_Q at $\sqrt{s} = 20$ TeV. Also shown is the cross-section for $pp \rightarrow Q\bar{Q} \rightarrow (W \rightarrow \ell\nu) + 4$ jets, assuming that $Q \rightarrow Wq$ [40].

b) The inclusive electron transverse momentum p_T spectrum from $W \rightarrow e\nu$ decays (dashed line) and for the process $pp \rightarrow Q\bar{Q} \rightarrow (W \rightarrow e\nu) + 4$ jets, for two different mass values of the heavy-quark Q (solid lines) [40].

Fig. 17 Limits on squark and gluino masses (90% CL) obtained by UA1. The arrows indicate the asymptotic values of the 90% CL contours as the squark or gluino mass becomes infinitely large [43].

Fig. 18 Limits on squark and gluino masses (90% CL) obtained by CDF. The arrows indicate the asymptotic values of the 90% CL contours, where the $\tilde{g}$ or $\tilde{q}$ mass is very large. The lower-limit curve corresponds to an CDF acceptance of 0.1% [45].

Fig. 19 Limits (90% CL) on a) $\tilde{e}$ and $\tilde{\gamma}$ masses and b) $\tilde{W}$ and $\tilde{\nu}$ masses obtained by UA2 [44]. The hatched area illustrates the systematic uncertainty. Also shown are limits from e^+e^- experiments.

Fig. 20 The W' (Z') lower mass limits (90% CL) shown as a function of the W' (Z') coupling strength obtained by UA1 [9].

Fig. 21 Limits (90% CL) on additional vector bosons W' and Z' depending on $\lambda_q^2 \cdot BR_e$ obtained by UA2 [44]: a) for the W' obtained from an analysis of single-electron candidates and b) for the Z' obtained from an analysis of electron pairs.

Fig. 22 Cross-section times branching ratio limits (95% CL) for the spin-0 and spin-1 particle X as a function of the mass m_X [48].

Fig. 23 Normalized angular distribution for very high mass jet-pairs as a function of χ . The solid curve is a QCD prediction ($\Lambda_c = \infty$); the dotted curve corresponds to $\Lambda_c = 300$ GeV, which is clearly excluded by the UA1 data [51].

Fig. 24 Expected limits (for 10 pb^{-1}) on $\tilde{e}$ and $\tilde{\gamma}$ at 90% CL from a search for $Z \rightarrow \tilde{e}\tilde{e} \rightarrow ee\tilde{\gamma}\tilde{\gamma}$ decays in UA2 [54].

Fig. 25 Production of W' and Z' at ACOL and Tevatron as a function of their mass, assuming standard couplings to leptons and quarks [54].

Fig. 26 Missing transverse energy distribution at $\sqrt{s} = 17$ TeV after possible selection cuts for squark (solid histogram) and gluino (dashed histogram) production and for the total background (points with error bars). A squark and gluino mass of $1 \text{ TeV}/c^2$ is assumed [58].

Fig. 27 Cross-section times maximal $\ell^+\ell^-$ branching ratio for Z' in a minimal rank-5 superstring-inspired model at $\sqrt{s} = 17$ and 40 TeV [60].

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6

Fig. 7

Fig. 9

Fig. 8

Fig. 10

Fig. 11

Fig. 12

Fig. 13

Fig. 14

Fig. 15

Fig. 16

Fig. 17

Fig. 18

Fig. 19

Fig. 20

Fig. 21

Fig. 22

Fig. 23

Fig. 24

Fig. 25

Fig. 27

Fig. 26