CERN-TH.5506/89

ROLE OF $\sin^2 \hat{\theta}_W(m_Z)$ AT THE Z^0 PEAK

A. Sirlin^{*)}
CERN - Geneva

ABSTRACT

The possible role of the $\overline{MS}$ parameter $\sin^2 \hat{\theta}_W(m_Z)$ in discussing physics at the Z^0 peak is outlined. A radiative correction $\Delta \hat{r}$ (the analogue of Δr), which links directly $\sin^2 \hat{\theta}_W(m_Z)$ to m_Z , is introduced and evaluated in the Standard Model. Assuming $60 \text{ GeV} < m_t < 200 \text{ GeV}$, $10 \text{ GeV} < m_H < 1 \text{ TeV}$, employing $\Delta \hat{r}$ and $m_Z = 91.12 \pm 0.16 \text{ GeV}$ leads to an accurate determination $\sin^2 \hat{\theta}_W(m_Z) = 0.2328 \pm 0.0012 \pm 0.0022$ (the first error is due to δm_Z and the second reflects the m_t, m_H uncertainty). It is interesting to note that this result is consistent with the simplest SUSY GUTs prediction. Accurate values for $\hat{\alpha}_1(m_Z)$ ($i = 1, 2$) are also given. Constraints due to gauge invariance in defining effective weak interaction angles are discussed.

*) Permanent address: Department of Physics, New York University, New York, NY 10003, USA.

In current discussions of physics at the Z^0 peak, the idea has been advanced of employing effective weak mixing angles that "absorb" to a considerable degree the top and Higgs mass dependence of the radiative corrections, as well as many unknown effects of new physics [1,2]. Such a quantity can be used, for instance, in "improved Born approximations" for the cross-sections and asymmetries, and can be correlated to the measured Z^0 mass with small theoretical uncertainty.

The first purpose of this note is to point out that there is an important parameter of the theory, to wit $\sin^2\hat{\theta}_W(m_Z)$, which is very well suited for this purpose. Here $\sin^2\hat{\theta}_W(m_Z)$ means the $\overline{MS}$ running constant, defined by modified minimal subtraction, evaluated at the m_Z scale. Apart from the fact that this is the natural parameter in GUTs predictions, it has long been emphasized that its use in electroweak calculations has an important simplifying property, namely that the electroweak form factor $\hat{\kappa}(q^2)$ multiplying $\sin^2\hat{\theta}_W(m_Z)$ [or $\sin^2\hat{\theta}_W(m_Z)$] is independent of the Higgs mass and only weakly dependent on the top mass [3,4]. The second purpose of this note is to show how $\sin^2\hat{\theta}_W(m_Z)$ can be introduced into the discussion of the Z^0 amplitude and derive a $\Delta\hat{\kappa}$ correction (the analogue of Δr [5]), which allows us to evaluate directly $\sin^2\hat{\theta}_W(m_Z)$ from m_Z . As we will see, $\Delta\hat{\kappa}$ has a simple mathematical structure. Assuming that $60 \text{ GeV} < m_t < 200 \text{ GeV}$ and $10 \text{ GeV} < m_H < 1 \text{ TeV}$, we then apply $\Delta\hat{\kappa}$ to determine this basic parameter with good accuracy in the Standard Model. Finally we discuss the m_t dependence of the residual corrections and comment briefly on another possible gauge invariant definition of an effective $\sin^2\hat{\theta}_W$.

A convenient framework to introduce $\sin^2\hat{\theta}_W(m_Z)$ is provided by the analysis of Ref. [6]. In that paper, it is shown that the $O(\alpha)$ contributions proportional to $\sin^2\hat{\theta}_W \equiv 1 - m_W^2/m_Z^2$ in the Z^0 mediated amplitude involve the combination

$$\sin^2\theta_W \left\{ 1 - \frac{c}{s} \frac{\hat{A}_{YZ}(q^2)}{q^2} - \frac{\alpha}{4\pi s^2} V(q^2) \right. \\ \left. - \text{Re} \left[\frac{c^2}{s^2} \left(\frac{A_{ZZ}(m_Z^2)}{m_Z^2} - \frac{A_{WW}(m_W^2)}{m_W^2} \right) + \frac{c}{s} \left(\frac{A_{YZ}(m_Z^2) + A_{YZ}(0)}{m_Z^2} \right) \right] \right\} \quad (1)$$

Here $\hat{A}_{YZ}(q^2)$ is the renormalized γZ self-energy, which in Ref. [6] satisfies $\text{Re}\hat{A}_{YZ}(m_Z^2) = 0$, $V(q^2)$ are finite vertex corrections, the A 's without carets are unrenormalized self-energies, and henceforth $s^2 = \sin^2\theta_W$, $\hat{s}^2 = \sin^2\hat{\theta}_W(m_Z)$, with an analogous meaning for c^2 , $\hat{c}^2$. As $\text{Re}\hat{A}_{YZ}(m_Z^2) = 0$, the important contribution for our discussion is the block of unrenormalized self-energies in Eq. (1), because it is here that the unknown Higgs and top dependences reside. The introduction of $\hat{s}^2$ follows then a strategy analogous to that developed in Ref. [4]. Defining

$$X \equiv \text{Re} \left(\frac{A_{ZZ}(m_Z^2)}{m_Z^2} - \frac{A_{WW}(m_W^2)}{m_W^2} \right) \quad (2)$$

one recognizes that $(c^2/s^2)X$ is the counterterm for $\sin^2\theta_W$, so that $\sin^2\hat{\theta}_W[1 - c^2/s^2 X] = \sin^2\theta_W^0$. Expressing (1) in terms of the bare parameter $\sin^2\theta_W^0$ removes $(c^2/s^2)X$ from the form factor, but leaves a divergent expression, namely $(c/s)(A_{YZ}(m_Z^2) + A_{YZ}(0))/m_Z^2$. [Note that in the original Eq. (1) the divergences of this term cancel against those of $(c^2/s^2)X$.] Next, one chooses the counterterms present in $\sin^2\theta_W^0$ to perform the $\overline{MS}$ renormalization, i.e., one subtracts the terms involving $(n-4)^{-1} + (\gamma - \ln(4\pi))/2$ in $(c/s)(A_{YZ}(m_Z^2) + A_{YZ}(0))/m_Z^2$ and sets the 't Hooft mass scale μ to the value m_Z . In terms of $\hat{s}^2$, Eq. (1) becomes

$$\hat{s}^2 \left\{ 1 - \frac{c}{s} \frac{\hat{A}_{YZ}(q^2)}{q^2} - \frac{\alpha}{4\pi \hat{s}^2} V(q^2) \right. \\ \left. - \frac{c}{s} \text{Re} \left(\frac{A_{YZ}(m_Z^2) + A_{YZ}(0)}{m_Z^2} \right) \frac{1}{\overline{MS}(\mu = m_Z)} \right\} \quad (3)$$

The term $(c^2/s^2)X$, which contains the Higgs as well as the dominant top dependence, has been absorbed in the $\overline{MS}$ redefinition of $\sin^2\theta_W^0$. There is a residual m_t dependence in the last term of (3) but, as we will see, it is very weak. This reasoning also shows how $\hat{s}^2$ and s^2 are related to $O(\alpha)$, namely

$$\hat{s}^2 = \left[1 - (c^2/s^2)X \frac{1}{\overline{MS}} \right] s^2 \quad (4)$$

where, and hereafter, the choice $\mu = m_Z$ is understood. Equation (4) allows us to pass from one definition to the other and has been employed in the past [7]. It is desirable, however, to have a direct link between $\hat{s}^2$ and m_Z . This can be achieved by introducing a new radiative correction $\Delta\hat{r}$ by means of the relation

$$\hat{s}^2 \hat{c}^2 (1 - \Delta\hat{r}) = s^2 c^2 (1 - \Delta r) \quad (5)$$

Inserting (4) into (5) leads to

$$\Delta\hat{r} = \Delta r - \left(\frac{\hat{c}^2}{s^2} - 1 \right) X \overline{MS} \quad (6)$$

The simplest way to combine the two terms is to note that Δr is μ -independent, so

that the right-hand side of (6) can also be written as $[\Delta r - X(c^2/s^2 - 1)]_{\overline{MS}}$. As the potentially large top dependence of Δr is contained precisely in a term of the form $(c^2/s^2)X$, we see that $\hat{\Delta}^2$ is obtained from Δr by erasing the big enhancement factor $c^2/s^2 = 3.3$ in this term, and performing the $\overline{MS}$ subtraction in what remains. Comparison with Eq. (34b) of Ref. [5] shows then that

$$\Delta \hat{r} = \left\{ \text{Re} \left(\frac{A_{ZZ}(m_Z^2)}{m_Z^2} - \frac{A_{WW}(0)}{m_W^2} \right) - 2 \delta e/e \right. \\ \left. + \frac{\alpha}{4\pi \hat{s}^2} \left[\left(\frac{Z}{2} - 6\hat{s}^2 \right) \frac{\ln \hat{c}^2}{\hat{s}^2} + 6 \right] \right\}_{\overline{MS}} \quad (7)$$

where $\delta e/e$ is the charge renormalization counterterm discussed in that work. It is worth noting that the structure of (7) is rather simple; in particular, the two terms involving $\text{Re} A_{WW}(m_W^2)/m_W^2$ in Δr have now cancelled! Using the results of Refs. [5] and [8], as well as $\text{Re} \Gamma_{\gamma\gamma}^{(5)}(m_Z^2) = -0.0288 \pm 0.0009$ [9], setting $m_Z = 91.12$ GeV [10, 11], $\hat{s}^2 = 0.233$ and calling $\tau \equiv m_t^2/m_Z^2$, $\xi \equiv m_H^2/m_Z^2$, we find

$$\Delta \hat{r} = 0.0701 + \frac{\alpha}{2\pi \hat{s}^2} \left\{ f_1(\tau) + f_2(\xi) \right\} \quad (8a)$$

$$f_1(\tau) = 0.1863 \left[\ln \tau + \frac{1}{3} + (3+D)(\Lambda(D)-1) \right] \\ - \frac{3}{2} \frac{\tau}{\tau^2} \left[\Lambda(D) - 1 + \frac{1}{4} \right] - \frac{8}{9} \hat{s}^2 \ln \tau \quad (8b)$$

$$f_2(\xi) = \frac{H(\xi)}{2\hat{c}^2} - \frac{3}{8} \frac{(\xi \ln \xi - \hat{c}^2 \ln \hat{c}^2)}{(\xi - \hat{c}^2)} \quad (8c)$$

where $H(\xi)$ is given analytically in Eqs. (A9) and (A10) of Ref. [8], $D \equiv 4\epsilon-1$ and, for $\tau > 1/4$, $\Lambda(D) = D \tan^{-1}(1/D^2)$. From (5) it follows that $\hat{s}^2$ depends on m_Z and $\hat{\Delta}^2$ in the same way as s^2 depends on m_Z and Δr . Therefore

$$\hat{s}^2 = \frac{1}{2} \left\{ 1 - \left[1 - \left(\frac{74.562 \text{ GeV}}{m_Z} \right)^2 \frac{1}{(1-\hat{\Delta}^2)} \right]^{1/2} \right\} \quad (9)$$

The radiative correction $\Delta \hat{r}$ given in Eqs. (8) and the values of $\hat{s}^2$ derived by inserting $m_Z = 91.12 \pm 0.16$ GeV (the weighted average of Refs. [10] and [11]) into Eq. (9) are shown in Table 1 as a function of m_t and m_H . From these values, one

finds

$$\sin^2 \theta_W(m_Z) = 0.2328 \pm 0.0012 \pm 0.0022 \quad (10)$$

where the first error is due to the experimental uncertainty in m_Z and the second arises from our ignorance of m_t and m_H under the assumption $60 \text{ GeV} < m_t < 200 \text{ GeV}$, $10 \text{ GeV} < m_H < 1 \text{ TeV}$. As the experimental error will decrease rapidly in the near future, we have here a determination of a fundamental parameter of the electroweak theory at the 1% level. The theoretical error will further decrease if the lower bound on m_t increases or, obviously, if m_t and/or m_H become known. It is interesting to note that (10) is consistent with the simplest SUSY GUTs prediction [7, 12]. In conjunction with $\hat{\Delta}^{-1}(m_Z) = 127.9 \pm 0.3$, where the error reflects hadronic uncertainties [12] and the central value was obtained by setting $m_W = 79.79 \text{ GeV}$, $m_Z = 91.12 \text{ GeV}$ and $m_t > m_Z$ in the analysis of Ref. [13], Eq. (10) leads to $\hat{\Delta}_2(m_Z) = \hat{\Delta}(m_Z)/\sin^2 \theta_W(m_Z) = (3.36 \pm 0.02 \pm 0.03) \cdot 10^{-2}$ and $\hat{\Delta}_1(m_Z) = (5/3) \hat{\Delta}(m_Z)/\cos^2 \theta_W(m_Z) = (1.699 \pm 0.003 \pm 0.006) \cdot 10^{-2}$ (again, the first errors quoted arise from δm_Z).

It is interesting to reflect on the complementary nature of $\sin^2 \theta_W$ and $\sin^2 \theta_W(m_Z)$. While the first can be obtained from ν or, of course, from m_t^2/m_Z^2 , with little or no hindrance from our ignorance of m_t , m_H , the same is true of the second if we use m_Z as input. Vice versa, if the inputs are exchanged, both parameters become sensitive to m_t and, to a lesser degree, m_H .

It is also worthwhile to note that the bosonic contribution to $\Delta \hat{r}$ is given by $(\alpha/2\pi \hat{s}^2)[2.905 + f_2(\xi)]$ and amounts to 0.97%, 1.22% and 1.57% for $m_H = 10 \text{ GeV}$, 100 GeV and 1 TeV respectively. Table 2 shows the top contribution $(\hat{c}/\hat{s}) \text{Re} A_{\gamma Z}^{(t)}(m_Z^2)/m_Z^2$ to the last term in Eq. (3). This quantity contains the residual m_t dependence in the resonance region after the $\overline{MS}$ renormalization. One sees that its contribution to the electroweak form factor $\hat{\kappa}$ multiplying $\hat{s}^2$ is quite small.

The reader may wonder whether it would be possible to absorb in a redefinition of $\sin^2 \theta_W$ the complete block of unrenormalized self-energies shown in Eq. (1), just by itself. Although this is, in fact, a procedure currently advocated as a practical way of analyzing physics in the resonance region, it has, in the opinion of this author, a serious conceptual drawback. The point is that in the bosonic sector only the X contribution to the block is gauge invariant. This is parenthetically the reason why the $\overline{MS}$ renormalization does not destroy in this case the gauge invariance of the residual $O(\alpha)$ corrections: it simply absorbs a gauge-invariant part! On the other hand, it is possible to absorb the complete fermionic contribution to the block because, at least at the one-loop level, it is gauge

Invariant. We have refrained from doing this because we would lose the identification of $\hat{g}^2$ with the $\overline{MS}$ coupling and, as we have seen, the residual m_t dependence is mild. An alternative gauge-invariant procedure is to absorb the complete block of unrenormalized self-energies in (1), together with a subset of vertex diagrams evaluated exactly on resonance. As the latter are not universal, it is then necessary to choose a particular case, such as the $Z\bar{e}e$ amplitude on resonance. This point will be discussed in greater detail in a separate communication.

ACKNOWLEDGEMENTS

The author is indebted to G. Altarelli, A. Blondel, J. Ellis and W. Hollik for motivating lectures and conversations, to G. Degraassi for very useful discussions and correspondence, and to the CERN Theory Group for its warm hospitality during the exciting month of August 1989. This research was supported in part by the National Science Foundation under Grant No. PHY-8715995.

m_t (GeV)	Δt	$\sin^2 \theta_W(m_Z)$
60	0.0673	0.2344
90	0.0661	0.2340
100	0.0656	0.2338
120	0.0643	0.2334
150	0.0619	0.2325
180	0.0589	0.2314
200	0.0567	0.2307

Table 1

Δt and $\sin^2 \theta_W(m_Z)$ as a function of m_t . These values correspond to $m_H = 100$ GeV. For $m_H = 1$ TeV, one must add +0.0035 to every value in the Δt column and either +0.0012 or +0.0013 to the entries in the $\hat{g}^2$ column. For $m_H = 10$ GeV, one adds -0.0025 to every value in the Δt column, -0.0009 to the first six entries in the $\hat{g}^2$ column and -0.0008 to the last.

m_t (GeV)	$\frac{\Delta \text{Re} A_Z^{(t)}}{s} \frac{(m_Z^2)^{\overline{MS}}}{m_Z^2}$
60	$-1.0 \cdot 10^{-3}$
100	$2.1 \cdot 10^{-6}$
150	$6.2 \cdot 10^{-4}$
200	$1.0 \cdot 10^{-3}$

Table 2

Top contribution to the electroweak form factor multiplying $\hat{g}^2$ in the resonance region [cf. Eq. (3)].

REFERENCES

- [1] G. Altarelli, talk delivered at the NATO Advanced Research Workshop on "Radiative Corrections: Results and Perspectives", University of Sussex, United Kingdom, July 1989.
- [2] A. Blondel, report CERN-EP/89-84, and references therein.
- [3] W.J. Marciano and A. Sirlin, Phys. Rev. Lett. 46 (1981) 163.
- [4] S. Sarantakos, W.J. Marciano and A. Sirlin, Nucl. Phys. B217 (1983) 84.
- [5] A. Sirlin, Phys. Rev. D22 (1980) 971.
- [6] A. Sirlin, "Considerations Concerning the Renormalization of the Electroweak Sector of the Standard Model", Universitat de Valencia (Burjassot) preprint (1989).
- [7] U. Amaldi et al., Phys. Rev. D36 (1987) 1385;
G. Costa et al., Nucl. Phys. B297 (1988) 244.
- [8] W.J. Marciano and A. Sirlin, Phys. Rev. D22 (1980) 2695.
- [9] H. Burkhardt et al., in "Polarization at LEP", eds. G. Alexander et al., CERN-88-06 (1988).
- [10] Mark II Collaboration, presented by G. Feldman, Proceedings of the XIV International Symposium on Lepton and Photon Interactions, Stanford (August 1989).

- [11] F. Abe et al., Phys. Rev. Lett. 63 (1989) 720.
- [12] W.J. Marciano, Brookhaven report BNL-41498 (1988).
- [13] W.J. Marciano and A. Sirlin, Phys. Rev. Lett. 56 (1986) 22.