Ref.TH.2505-CERN

IS THE LATEST ATTEMPT TO CONTRADICT THE TUBE MODEL CREDIBLE?

L. Bergström

Royal Institute of Technology, Stockholm

and

S. Fredriksson^{*)}

CERN -- Geneva

ABSTRACT

It is shown that recent data from pion and proton collisions with emulsion nuclei at 200 GeV/c are in perfect agreement with expectations from the collective tube model. The opposite conclusion in publications by Azimov et al. was built on a series of incorrect arguments.

^{*)} On leave of absence from the Department of Theoretical Physics, Royal Institute of Technology, Stockholm.

During the last year it has been claimed in papers from a big emulsion collaboration^{1),2)} that data are in conflict with "predictions" from the so-called Collective Tube Model (CTM) for high-energy multiple production off nuclei. In spite of our efforts to obtain a correct presentation, these articles have now been followed by an additional letter³⁾, in which further misinterpretations and misquotations occur. We feel, therefore, compelled to a public rebuttal and clarification.

We are at present performing a detailed analysis of all aspects of proton production and target fragmentation in high-energy collisions with nuclei from the CTM point of view. As we consider it urgent to prevent further spreading of trivial mistreatments of the CTM, we break out the part on emulsion data and present it in a shortened form in this letter. All details that we have to rush over will be carefully presented in an upcoming paper.

The CTM, reviewed in Refs. 4) and 5), is built on the postulate:

"A non-diffractive particle-nucleus hit is equivalent to a particle-tube hit, where the whole tube of nuclear matter, traversed by the projectile, collides like one single heavy hadron."

The problem is that we do not know in general how the detailed quantum numbers of the tube influence the reaction. Luckily, however, many important features of high-energy collisions are known from particle-particle experiments to be roughly independent of the target quantum numbers, typical examples being central production of pions and beam fragmentation. We therefore use the following working hypothesis of "universality"^{4),5)}:

"Whenever an inclusive cross-section is known to be a universal function of the invariants s , t and $x = 2p_L^*/\sqrt{s}$ (and eventually of the nature of the projectile), independent of the target quantum numbers, the tube influences the collision only by its mass, which reflects in the kinematics of the process."

By evaluating s , t and x for a target mass of im_N , corresponding to i nucleons in the tube, we easily get the CTM prediction for any "universal" cross-section $f^{hA \rightarrow cX} = \sigma_{inel}^{-1} E d\sigma/d^3p$ in the practical form:

$$f^{hA \rightarrow cX}(p_{beam}, y_{lab}, p_T) = \sum_{i=1}^A P(i, A) f^{hN \rightarrow cX}(ip_{beam}, y_{lab} + \ln i, p_T) \quad (1)$$

Here $f^{hN \rightarrow cX}$ is the same cross-section with a nucleon target. $P(i, A)$ is the probability to hit a tube with i nucleons in a target with A nucleons. Often one can safely omit the summation and use $i = \langle i \rangle = A^{1/3}$ in (1).

The papers by Azimov et al. contain around a dozen errors. We have tried to correct for all of them in our analysis, although a detailed list of all corrections would be far too long for this letter. We will only comment on (the few) mistakes that have direct influence on the conclusions about the CTM. All assumptions in Refs. 1)-3) that are "questionable", but difficult to prove incorrect, are left unchanged, to simplify the comparison between our analysis and an incorrect one.

The most serious mistake in Ref. 3) is the assumption that "universality" and hence Eq. (1) is valid for all cross-sections, i.e., that everything is independent of the structure of the target. This is obviously not true for the very important part of rapidity spectra that comes from target fragmentation. It is unfortunately too late to point out that the CTM papers quoted in Ref. 3) all contain explicit warnings against such an interpretation.

The approach we adopt is the following. Produced particles are assumed to come from either beam fragmentation, central production ("pionization"), or target fragmentation. The first two mechanisms are roughly target independent and hence give spectra that obey Eq. (1). Target fragments however come from the decay of the tube itself, or equivalently, from the fragmentation of the nucleons (quarks) in the tube. Intuition tells us that this mechanism, in any model lacking complicated intranuclear cascades, should give a spectrum that grows roughly proportionally to the size (No. of nucleons, No. of quarks) of the effective target for all rapidities. As expected, such a behaviour is in definite disagreement with (1). In order to make a correct test of the CTM, we therefore calculate, fully within the CTM, the fraction of target fragments in rapidity spectra, subtract them from data^{1),2)} and show that the remaining fraction obeys Eq. (1). The rest of this letter contains the details of our CTM calculation of tube fragment spectra, as well as a discussion of two-particle correlation data³⁾, which were also claimed to be in disagreement with the CTM. Our quantitative results are presented in Figs. 1-2. Fig. 2b should be directly compared with Fig. 2 of Ref. 3) and with the incorrect conclusion drawn from it.

In the CTM, target fragmentation is most naturally looked upon as a two-step process; the tube recoils in a collective manner and then decays. For the recoil we have shown⁶⁾ that the following assumption is in excellent agreement with data⁷⁾ on $\pi\text{Ne} \rightarrow \text{pX}$ at 10.5 GeV/c.

A.1) The tube recoils like one single nucleon, i.e., the x spectra of $h + \text{tube} \rightarrow \text{tube} + X$ and $h + N \rightarrow N + X$ are equal.

For the tube decay it is not enough this time to assume just a reasonable "softness"⁶⁾, because we need the angle distribution of secondaries as well as the

fraction of pions in order to reproduce the pseudo-rapidity (η) spectra of all charged particles. We therefore add:

A.2) The recoiling tube, having no transverse motion, decays in its rest system to nucleons and pions, both with relative momentum spread $\exp(-6E_{\text{rel}})$.

Still we need to know the total number of pions that fragment out from the tube. The most natural assumption for this is:

A.3) A nucleon "shakes off" pions in connection to the recoil and decay of the tube according only to the boost it gets in the collision, independently of the number of nucleons surrounding it in the tube.

Together with A.1) this gives a simple scaling law for the "non-universal" inclusive cross-section of any kind of target fragment t :

$$f^{hA \rightarrow tX}(p_{\text{beam}}, y_{\text{tub}}, p_T) = \sum_{i=1}^A i P(i, A) f^{hN \rightarrow tX}(p_{\text{beam}}/i, y_{\text{tub}}, p_T) \quad (2)$$

This is indeed very different from Eq. (1). The dependence on p_{beam}/i is weak⁶⁾, so the contribution from target fragments grows as expected like $A^{1/3}$. A more detailed parton-tube approach⁸⁾ for the backward region gives very similar results (but is not directly applicable to the region $\eta > 0$). The reason that we still need assumption A.2) is that we lack detailed knowledge of target fragmentation in hN collisions.

Before going into details, a few qualitative observations can be made. In the region $0 \leq \eta \leq 1$, where the bulk of tube fragments lies, we expect the η spectrum $\sigma_{\text{inel}}^{-1} d\sigma/d\eta$ of charged particles to be roughly proportional to $\langle i \rangle \approx A^{1/3}$. At smaller η values, the spectrum should grow slightly faster with A due to the velocity cut in data and other kinematic details. The ratios of hEm to hN spectra^{1), 2)} are indeed around 4 at $\eta = 0.5$ for both proton and pion beams ($\langle A \rangle \approx 64$ in emulsion). The target fragments are expected to show up as an extra peak relative spectra from nucleon targets. In pEm data at $p_{\text{beam}} \leq 200 \text{ GeV}/c$ ¹⁾ this peak is not visible, since the broad distribution of central pions overlaps too much. In πEm , however, the central peak lies at higher η values, just like in πp compared to pp data, so the separate peak of tube fragments is clearly visible^{1), 2)}. Hence we do not share the view in Refs. 1) and 2) that the observed "bimodality" reflects some hitherto unexplored physical phenomenon.

To perform the quantitative analysis, we first take the average number of charged pions fragmenting from a proton in a high-energy pp collision to be 0.8.

This number we get through the following rough estimates. Assume that a proton "kicks out" exactly k charged pions with probability c^k ($k \geq 1$). To reproduce the 40%⁹⁾ rate of protons that "redress" to neutrons, we need $c \approx 0.35$. Hence the number of charged pions per proton is $\sum k c^k \approx 0.8$. This is consistent with more detailed estimates¹⁰⁾ from the old two-component model.

Estimating tube fragment spectra is now straightforward. Compared to the previous calculation⁶⁾ of η spectra of protons, we integrate over tube decay, use the relation between η and η , take the velocity cut $v > 0.7c$ into account and finally add pion and proton contributions. The crucial $d\sigma/dx$ for recoils was taken, as in Ref. 6), from rather old data on $\pi p \rightarrow pX$ at 6 GeV/c¹¹⁾. More recent data on $\pi p \rightarrow pX$ at 8 and 16 GeV/c¹²⁾ and on $\pi p \rightarrow$ target charge + X at 100 GeV/c¹³⁾ are, however, in perfect agreement with this $d\sigma/dx$. We use the same recoil cross-section for a proton beam. Data on $pp \rightarrow pX$ are of no use there, because one cannot discriminate large momentum target recoils from retarded projectiles.

In Fig. 1 it is shown how the split-up in target fragments and "universal" particles works for the η spectrum of $\pi Em \rightarrow$ all charged + X²⁾. The analysis in Ref. 3) was, however, built on spectra taken at fixed n_g , the number of grey prongs. A quadratic relation between n_g and tube size i was assumed to be valid event by event. Such a relation might appear in Glauber-like models with the recoils cascading in the nucleus, but in the CTM it is not at all clear that a meaningful relation exists. Triggering on a fixed n_g , defined in an ad hoc way as the number of prongs with $0.3c \leq v \leq 0.7c$, might give any kind of biases in individual events. A detailed analysis of this would, in any model, mean that also complicated nuclear effects must be taken into account. For simplicity, we assume, however, that there is a meaningful average relation between n_g and $\langle i \rangle$, in the sense that events with a fixed n_g look the same on the average as completely unbiased events from a target of $\langle i \rangle$ nucleons. The only way to extract this relation without further assumptions is to look at probabilities. We match the measured¹⁾ distribution of n_g against the theoretical distribution of tube sizes in emulsion, so that the 27% events with $n_g = 0$ are assumed to come from the 27% smallest tubes, etc.

The main features of the "non-target" spectra that we extract by the described procedure are displayed in Fig. 2 as functions of both n_g and $\langle i \rangle$. Figure 2a shows the position in η of the peak. Figure 2b shows the half-width of the spectrum at half the height. Figure 2c shows the height of the peak. The curves in Fig. 2 prove that the behaviour of the "universal" part of η spectra is in perfect agreement with expectations from the CTM, as quantified in Eq. (1). With increasing tube size i and fixed p_{beam} the peak should move $\frac{1}{2} \ln i$ units to

the left, the half-width H should increase like $0.83 [\ln(p_{\text{beam}} i/2m_p)]^{\frac{1}{2}}$ and the whole spectrum should blow up like i^α/H with $\alpha \approx 0.285$ ³⁾. Here we use that $H \approx 0.83D$, where D is the dispersion, for which we take the same parametrization as in Ref. 3).

Also the arguments ³⁾ against the CTM built on studies of the two-particle correlator $R_2^{hA}(\eta_1, \eta_2)$ are too hasty, because a correct analysis shows that all differences between R_2^{hA} and R_2^{hN} are due to target fragmentation effects, which were neglected in Ref. 3). Data are presented for $R_2^{hEm}(\eta) = R_2^{hEm}(\eta, \eta)$ with both pion and proton beams. The η and n_g (or tube size) dependences are indeed easy to understand in the CTM. $R_2^{pp}(\eta)$ is symmetric in η_{cms} and negative in the beam and target fragmentation regions, because the probability to have two charged target (beam) fragments is very small in pp collisions. However, the average number of such particles exceeds two already with a tube size $i = 2$. Since these fragments in addition are spread out less in phase space than central hadrons, the correlator R_2^{hA} "jumps up" to a large value when going from hp to h tube already with a small tube. When the tube size is increased further, a new effect sets in that again successively reduces R_2^{hA} somewhat in parts of the target fragmentation region: The central bump moves towards lower η with increasing i in line with Eq. (1) and starts to overlap with tube fragments. The correlations are then partially destroyed, because a target fragment and a central hadron do not have a mutual short-range coupling. The expected trends are therefore that $R_2^{h\text{-tube}}$

- a) $\approx R_2^{hp}$ outside the target fragmentation region;
- b) $\gg R_2^{hp}$ in the target fragmentation region already for $i = 2$;
- c) decreases with increasing tube size $i > 2$ in the region where tube fragments overlap with other hadrons, i.e., at $0 \lesssim \eta \lesssim 3$ for pEm and at $1 \lesssim \eta \lesssim 4$ for πEm ;
- d) decreases with increasing η .

All this is in good qualitative agreement with data. A quantitative comparison with data is complicated and would require extra assumptions about the detailed correlations between tube fragments. Such information could hardly be extracted from πp and pp data.

In conclusion, we have shown that

- a) the influence of target fragments is extremely important in existing high energy emulsion data, and in fact explains the recently found "bimodality" of η spectra from πEm collisions;
- b) when CTM (or any other cascade free model) estimates of target fragment spectra are subtracted from data, the rest behaves the way a "universal" cross-section should do in the CTM, e.g., gets a width that grows with growing tube size;

- c) two-particle correlations are easily understood in the CTM if effects of tube decay are not neglected;
- d) it is wisest to consult the inventors of a model before publishing hasty "counterproofs".

After the completion of this manuscript we became aware of a recent experimental paper¹⁵⁾, where it is confirmed that the η spectrum in 200 GeV/c nA collisions grows like $A^{1/3}$ in the extreme backward region $\eta \approx -2$. The authors give strong support to the view presented here by showing that the backward spectrum is not "universal" in the meaning of Eq. (1). They however seem unable to understand¹⁶⁾ that the target fragmentation region of any collision is dominated by target fragments, and hence end up with the incorrect conclusion that the CTM "is in serious disagreement with data". In fact, their backward yields are trivially in agreement with any model that lacks slow hadron cascading.

Acknowledgements

We are grateful to B. Andersson, G. Bellini, A. Capella, A. Dar, E. Meroni, N. Nikolaev, D. Tow and R. Weiner for inspiring discussions about this analysis, as well as to Y. Afek, G. Bellini, G. Berlad, A. Dar, G. Eilam, M. Longo, I. Otterlund and F. Takagi for providing us with unpublished material of relevance to this work.

REFERENCES

- 1) S.A. Azimov et al., paper submitted to the VIII Int. Conf. on High Energy Physics and Nuclear Structure, Zurich 1977; the pEm data presented in the figures of this paper as " η spectra", with $\eta = -\ln \tan (\theta/2)$, are most likely λ spectra, where λ is the "cosmic ray variable" $-\ln \tan \theta$. Only by correcting for this can we get consistency with other emulsion data¹⁴⁾.
- 2) Z.A. Anzon et al., Nuclear Phys. B129 (1977) 205.
- 3) S.A. Azimov et al., Phys. Letters 73B (1978) 339.
- 4) Y. Afek et al., in Proceedings of the Topical meeting on multiparticle production on nuclei at very high energies (ed. G. Bellini, L. Bertocchi and P.G. Rancoita), Trieste 1976; and references therein.
- 5) S. Fredriksson, preprint TH.2423-CERN (1977), to be published in the Proceedings of the Triangle Seminar: Recent developments in high energy physics, Campione d'Italia 1977; and references therein.
- 6) L. Bergström and S. Fredriksson, Phys. Letters 68B (1977) 177.
- 7) W.M. Yeager et al., Phys. Rev. D16 (1977) 1294.
- 8) F. Takagi, Tohoku University preprint TU/77/176 (1977).
- 9) M. Antinucci et al., Nuovo Cimento Letters 6 (1973) 121.
- 10) R.G. Roberts, in Phenomenology of Particles at High Energies (ed. R.L. Crawford and R. Jennings), Academic Press, London (1974).
- 11) D.J. Crennell et al., Brookhaven National Laboratory report BNL-16417 (1971).
- 12) P. Bosetti et al., Nuclear Phys. B54 (1973) 141.
- 13) J. Whitmore et al., Phys. Rev. D16 (1977) 3137.
- 14) J. Hébert et al., Phys. Letters 48B (1974) 467; Phys. Rev. D15 (1977) 1867.
- 15) D.L. Burke et al., Univ. of Michigan preprint UM HE 78-6 (1978).
- 16) M.J. Longo, private communication.

FIGURE CAPTIONS

Fig. 1 : Tube model estimates of the composition of measured²⁾ η spectra of all charged particles with $v > 0.7c$ in 200 GeV/c π Em collisions.

Fig. 2 : Properties of the extracted target independent part of η spectra (total yields minus CTM estimates of target fragments), versus n_g and corresponding tube sizes $\langle i \rangle$. Curves show parameter free expectations from the CTM via Eq. (1). Horizontal error bars show uncertainties in the empirical relations between n_g and $\langle i \rangle$.

a) Peak position in η . Curves: $\eta_{\text{peak}}^{\text{hN}} - \frac{1}{2} \ln \langle i \rangle$.

b) Half-width of spectra. Curve: $0.83 [\ln (p_{\text{beam}} \langle i \rangle / 2m_N)]^{\frac{1}{2}}$.

c) Maximal height of spectra. Curves: $C^{\text{hN}} \langle i \rangle^{0.285} / H$, where C^{hN} is fitted to hN data and H is the half-width in b).

FIG.1

FIG.2