SAPIENZA
UNIVERSITÀ DI ROMA

Recent results on soft QCD with the ATLAS experiment

Simone Monzani

Università di Roma1

On behalf of the ATLAS collaboration

Montpellier, 29/06/15

Outline

- Bose-Einstein correlations ([ArXiv:1502.07947](#))
- Measurement of transverse polarization of Λ and $\bar{\Lambda}$ ([ArXiv:1412.1692](#))
- Measurement of distributions sensitive to the underlying events in inclusive Z-boson production ([Eur. Phys. J. C \(2014\) 74:3195](#))
- Measurement of the total cross section from elastic scattering in pp collisions ([Nuclear physics B \(2014\), 486-548](#))
- Conclusions

Bose-Einstein correlations

Bose-Einstein correlations

- Bose-Einstein correlations (BEC): correlations between two identical bosons (consequence of the symmetry of identical bosons wave function)
- BEC effect corresponds to an enhancement in two identical boson correlation function when the two particles are near in momentum space
- BEC sensitive probe of the space-time geometry of the hadronization region, allows the determination of the size and the shape of the source from which particles are emitted

$$C_2(p_1, p_2) = \frac{\rho(p_1, p_2)}{\rho_0(p_1, p_2)} = C_0 [1 + \Omega(\lambda, QR)] (1 + \varepsilon Q)$$

2 particles
correlation function

C_0 : normalization factor

ε : long-range momentum correlations

λ : strength parameter, 0 (1) for purely coherent (chaotic) sources

R : effective radius of the source size

$$R_2(Q) = \frac{C_2(Q)}{C_2^{MC}(Q)}$$

To account to resonances effects

$$Q^2 = -(p_1 - p_2)^2$$

MC constructed on unlike-sign charged-particle pairs

- 0.9 TeV
- 7 TeV
- 7 TeV (high multiplicity)

At least one vertex from at least two tracks (like-sign charged particles)
 with $p_T > 100 \text{ GeV}$ and $|\eta| < 2.5$

At least 120 tracks for high multiplicity events

Bose-Einstein correlations

Full phase-space results

λ and R determined by measurements

BEC parameters for fits of the two-particle double-ratio correlation function $R_2(Q)$

$\lambda = 0.74 \pm 0.11$, $R = 1.83 \pm 0.25$ at $\sqrt{s} = 0.9$ TeV for $n_{\text{ch}} \geq 2$

$\lambda = 0.71 \pm 0.07$, $R = 2.06 \pm 0.22$ at $\sqrt{s} = 7$ TeV for $n_{\text{ch}} \geq 2$,

$\lambda = 0.52 \pm 0.06$, $R = 2.36 \pm 0.30$ at $\sqrt{s} = 7$ TeV for $n_{\text{ch}} \geq 150$

Values close to the ones obtained by ALICE and CMS

Dependence of the BEC parameters λ and R on n_{ch} and k_T

BEC param.	Fit function	0.9 TeV	7 TeV	
			Minimum-bias events	High-multiplicity events
$R(n_{\text{ch}})$	$p_0 \sqrt[3]{n_{\text{ch}}}$ p_0	$p_0 = 0.64 \pm 0.07$ fm ($n_{\text{ch}} \leq 82$) —	$p_0 = 0.63 \pm 0.05$ fm ($n_{\text{ch}} < 55$) $p_0 = 2.28 \pm 0.32$ fm ($n_{\text{ch}} \geq 55$)	—
$\lambda(n_{\text{ch}})$	$p_0 e^{-p_1 n_{\text{ch}}}$	$p_0 = 1.06 \pm 0.10$ $p_1 = 0.011 \pm 0.004$	$p_0 = 0.96 \pm 0.07$ $p_1 = 0.0038 \pm 0.0008$	
$R(k_T)$	$p_0 e^{-p_1 k_T}$	$p_0 = 2.64 \pm 0.33$ fm $p_1 = 1.48 \pm 0.67$ GeV ⁻¹	$p_0 = 2.88 \pm 0.27$ fm $p_1 = 1.05 \pm 0.58$ GeV ⁻¹	$p_0 = 3.39 \pm 0.54$ fm $p_1 = 0.92 \pm 0.73$ GeV ⁻¹
$\lambda(k_T)$	$p_0 e^{-p_1 k_T}$	$p_0 = 1.20 \pm 0.18$ $p_1 = 2.00 \pm 0.35$ GeV ⁻¹	$p_0 = 1.12 \pm 0.10$ $p_1 = 1.54 \pm 0.26$ GeV ⁻¹	$p_0 = 0.75 \pm 0.10$ $p_1 = 0.91 \pm 0.45$ GeV ⁻¹

Bose-Einstein correlations

The saturation of R for high-multiplicity particles (≈ 240) is observed for the first time and can be predicted by Pomeron-based models (due to overlapping of colliding protons).

Multiplicity (n_{ch}) dependence

R independent to multiplicity for $n_{ch} > 50$

k_T dependence

$$k_T = \frac{|p_{T_1} + p_{T_2}|}{2}$$

Measurement of the transverse polarization of Λ and $\bar{\Lambda}$

Measurement of the transverse polarization of Λ and $\bar{\Lambda}$

P: polarization vector transverse to the momentum of Λ $0 < P < 1$

$\alpha = 0.642 \pm 0.013$
World average of the P-violating decay asymmetry for Λ

Probability distribution of θ^*

$$g(t; p) = \frac{1}{2} (1 + \alpha P t)$$

$$t = \cos \theta^*$$

From previous experiments:

- polarization measured only in experiment with fixed target up to 40 GeV in the cms system and t up to 62 GeV at ISR
- the magnitude of the Λ polarization increases with p_T until it saturates at about 1 GeV
- the magnitude of the Λ polarization decreases with decreasing $|x_F|$
- the Λ polarization does not depend strongly on the center of mass energy, tested up to 40 GeV

Measurement of the transverse polarization of Λ and $\bar{\Lambda}$

Probability distribution function modified by detector efficiency and resolution effects (t' cosine of the measured decay angle)

$$E(P) = \int_{-1}^1 g_{\text{det}}(t'; P) dt' = E(0) + [E(1) - E(0)]P$$

Method of moments

$$E_i^{\text{exp}}(P, E_{\text{bkg}}) = f_i^{\text{sig}} \{E_i^{\text{MC}}(0) + [E_i^{\text{MC}}(1) - E_i^{\text{MC}}(0)]P\} + (1 - f_i^{\text{sig}})E_{\text{bkg}}$$

MC weighting using t

Measurement of the transverse polarization of Λ and $\bar{\Lambda}$

$$P_{\Lambda} = -0.010 \pm 0.005(\text{stat}) \pm 0.004(\text{syst})$$

$$P_{\bar{\Lambda}} = 0.002 \pm 0.006(\text{stat}) \pm 0.004(\text{syst})$$

- Both consistent with 0 within the estimated uncertainty
- Comparison with others experiments (in agreement)

Fiducial Λ phase space

- $0.8 \text{ GeV} < p_T < 15 \text{ GeV}$
- $5 \cdot 10^{-5} < x_F < 0.01$
- $|\eta| < 2.5$
- $1100 \text{ MeV} < m_{p\pi} < 1127 \text{ MeV}$

Measurement of distributions sensitive to the underlying events in inclusive Z-boson production

Measurement of distributions sensitive to the underlying events in inclusive Z-boson production

Underlying events (UE)

- partons not participating in the hard-scattering process
- additional semi-hard scatters in pp collisions, multiple partons interactions (MPI)
- initial and final state gluon radiation contribute to UE

- Most active transverse region: trans-max
- Less active transverse region: trans-min

Flat distributions of UE activity, differences between the 2 transverse regions for hard scatter

↓
Trans-diff
(dominated by the hard scattering component)

Measurement of distributions sensitive to the underlying events in inclusive Z-boson production

Towards, trans-max/min show similar behaviour

POWHEG best agreement at low values, ALPGEN and SHERPA at higher

Trans-min region most sensitive to UE

Measurement of distributions sensitive to the underlying events in inclusive Z-boson production

ALPGEN and SHERPA best agreement at high p_T^Z

No model in agreement in the whole range (additional jets)

Comparison with other processes with a jet instead of a Z as leading particle

Leading jet more restricted as no jet with a p_T higher than the leading one can be considered

Measurement of the total cross section from elastic scattering in pp collisions

Measurement of the total cross section from elastic scattering in pp collisions

Nuclear physics B (2014), 486-548

- The total hadronic cross section is a fundamental parameter of strong interactions.
- Large distances are involved in the collision process and thus perturbation theory is not applicable
- Constraints through the Froissart-Martin bound or the optical theorem

- 2 tracking stations at 238 and 241 m from the interaction point (IP)
- $|\eta| > 8.5$

Measurement of the total cross section from elastic scattering in pp collisions

$$\frac{d\sigma}{dt} = \frac{1}{16\pi} \left| f_N(t) + f_C(t) e^{i\alpha\phi(t)} \right|^2 \quad \text{Differential elastic cross section}$$

f_C : Coulomb amplitude (function of $1/t$)
 f_N : purely strongly interacting amplitude
 (function of total cross section and $e^{-Bt/2}$)

$$-t = (\theta^* \times p)^2$$

θ^* : scattering angle at IP

$$B = 19.73 \pm 0.14(\text{stat.}) \pm 0.26(\text{syst.}) \text{ GeV}^{-2}$$

fit in the range

$$-0.01 \text{ GeV}^2 < t < 0.1 \text{ GeV}^2$$

where ALFA

acceptance > 10% and

single exponential

region

$$\sigma_{el} = 24.00 \pm 0.19 (\text{stat.}) \pm 0.57 (\text{syst.}) \text{ mb}$$

Parametrization integrated over the full t range

Hypothesis B is constant

Measurement of the total cross section from elastic scattering in pp collisions

$$\sigma_{tot}^2 = \frac{16\pi(\hbar c)^2}{1 + \rho^2} \left. \frac{d\sigma_{el}}{dt} \right|_{t \rightarrow 0}$$

Total cross section extrapolated at low t from the elastic cross section

$$\sigma_{tot} = 95.35 \pm 1.30 \text{ mb}$$

$$\sigma_{el} = 24.00 \pm 0.19 \text{ (stat.)} \pm 0.57 \text{ (syst.) mb}$$

2 σ far from model but other models have less increase

$$\sigma_{inel} = 71.34 \pm 0.36 \text{ (stat.)} \pm 0.83 \text{ (syst.) mb}$$

Less uncertainty than before
(no extrapolation to low diffractive masses outside the fiducial region)

Observed=inside the fiducial range

Conclusions

- Study of like-sign hadrons for Bose-Einstein correlation (BEC), clear signal BEC at low-Q, studied through the BEC parameters characterizing the correlation strength and the correlation source size
- Transverse polarization of Λ and Λ hyperons consistent with 0 at low x_F confirming the behaviour of previous experiments showing a decrease of polarization as function of x_F
- Measurement of UE is strongly affected by jets radiated from hard scatter, NLO/multileg generators give a better description of it, the Trans-min region is the least affected by extra-jets
- The total cross section is a fundamental parameter of strong interactions, it has been measured at 7 TeV center-of-mass energy confirming its rise proportional to $[\ln(s)]^2$, however the “asymptotic” energy dependence is yet to be determined
- All these studies repeated at 13 TeV will be important, since energy evolution is phenomenologically constrained

Back-up

Measurement of the transverse polarization of Λ and $\bar{\Lambda}$

From previous experiments:

- the magnitude of the Λ polarization increases with p_T until it saturates at about 1 GeV
- the magnitude of the Λ polarization decreases with decreasing $|x_F|$
- the Λ polarization does not depend strongly on the center of mass energy, tested up to 40 GeV

Fiducial Λ phase space

423498 Λ 378237 $\bar{\Lambda}$

- $0.8 \text{ GeV} < p_T < 15 \text{ GeV}$
- $5 \cdot 10^{-5} < x_F < 0.01$
- $|\eta| < 2.5$
- $1100 \text{ MeV} < m_{p\pi} < 1127 \text{ MeV}$

Selection criteria

- Vertex fit probability greater than 0.05
- Λ transverse decay distance significance and impact parameter significance

Background

K_0^s decays

$\gamma \rightarrow e^+ e^-$

$$\frac{L_{xy}}{\sigma_{L_{xy}}} > 15 \quad \frac{a_0}{\sigma_{a_0}} < 3$$

- reduction of physics background,
 $480 \text{ MeV} < m_{\pi\pi} < 515 \text{ MeV} \quad m_{ee} < 75 \text{ MeV}$

Measurement of distributions sensitive to the underlying events in inclusive Z-boson production

Events selection (Z decay)

- at least 2 primary vertex associated tracks with $p_T > 400$ MeV
- electrons and muons with $p_T > 20$ GeV, $|\eta| < 2.4$ and $z_0 \sin\theta < 10$ mm
- dileptons invariant mass of oppositely charged leptons between 66 and 116 MeV

Events selection (UE)

- $p_T > 0.5$ GeV, $|\eta| < 2.5$
- a minimum of 1 pixel and 6 in the semiconductor tracker
- 1 hit in the innermost pixel layer
- impact parameters $|d_0| < 1.5$ mm and $z_0 \sin\theta < 1.5$ mm
- for tracks with $p_T > 10$ GeV, goodness of fit greater than 0.01

Measurement of distributions sensitive to the underlying events in inclusive Z-boson production

Measurement of the total cross section from elastic scattering in pp collisions

Trigger conditions for elastic-scattering events, and 1 reconstructed track in all 4 detectors of the arm which fired the trigger.

60 μm from the edge of the detector