UAB-LEAE 90-04
March 90

Higher order corrections to the process

$$e^+e^- \rightarrow \nu\bar{\nu}\gamma$$

R. Miquel,
Laboratori de Física d'Altes Energies
Universitat Autònoma de Barcelona, E-08193 Bellaterra (Barcelona) Spain

C. Mana
Grupo de Altas Energias
C.I.E.M.A.T. - Madrid, Spain

and
M. Martinez
CERN, CH-1211 Geneva, Switzerland

March 23, 1990

Abstract

A calculation of the neutrino counting reaction $e^+e^- \rightarrow \nu\bar{\nu}\gamma$ is presented and implemented in a Monte Carlo generator of unweighted events. The non-photon corrections are included following the star-scheme. The photonic corrections have been exponentiated following the "inductive" and the "Structure Functions" approach. For a realistic experimental set-up, the total corrections around the Z^0 pole amount to $\sim 5\%$ with respect to the one-loop result. The final precision achieved is better than 1%.

1 Introduction

The radiative process $e^+e^- \rightarrow \nu\bar{\nu}\gamma$ has been proposed long time ago as a direct and clean way to determine the number of generations with massless neutrinos within the Standard Model framework. It will also be a potential source of background when searching for other neutral objects like $\tilde{\nu}$ and $\tilde{\gamma}$. The first calculations of this process were done at the Born level using the contact approximation for the W-exchange diagrams [1-2]. More recently, complete tree-level calculations have been performed and one-loop radiative corrections taken into account [3-4]. Since their effect is large, we have undertaken the task to include higher order corrections (more than one-loop) in order to keep the theoretical uncertainties below the 1% level which is of the order of the experimental accuracy one can achieve. This is not a trivial task because, since the radiated photons are the observed particles in the final state, the well known procedures of inclusive exponentiation, accurate to $\sim .1\%$, can not be directly used for our purpose. The results of this work, part of which can be extended to any process $e^+e^- \rightarrow f\bar{f}$ ($f \neq e$) involving neutral currents, are described in the present paper which is organized as follows. In section 2 we review the results for the lowest order cross-section. The following two sections deal with radiative corrections. As usual, they are divided in "photonic" corrections, described in section 3, and "non-photonic" corrections explained in section 4. Finally, the conclusions of the present work are given in section 5 together with some numerical examples.

2 The process $e^+e^- \rightarrow \nu\bar{\nu}\gamma$ in lowest order

The Feynman diagrams which describe the process $e^+e^- \rightarrow \nu\bar{\nu}\gamma$ in lowest order are depicted in fig. 1. A detailed description of the complete lowest order matrix element squared for different initial state helicity configurations can be found in [3] and will not be reproduced here. The results for the integrated total cross section ¹ when $E_\gamma \geq 1$ GeV and $|\cos\theta_\gamma| < \cos 15^\circ$ are shown in table 1 for different approximations as well as for the exact calculation.

$\sqrt{s}$ (GeV)	σ_Z	σ_{ct}	σ_{ZW}	σ_T
98	126.5	129.5	128.7	128.7
150	5.80	9.97	7.74	7.79

Table 1: Comparison between different approximations for the tree-level cross-section.

The first column shows the total cross-section when only the Z^0 diagrams (1.a and 1.b in fig. 1) are included. The second one is obtained neglecting the two W exchange diagram (1.e) and taking the limit $M_W \rightarrow \infty$ in the diagrams 1.c and 1.d. The third column shows the total cross-section when only the two W exchange diagrams is neglected and finally, the last column shows the result of the complete lowest order calculation. One can see that, as expected, the contribution of the diagram with two W exchange is negligible

¹The free parameters were set to $M_Z = 92$ GeV and $\sin^2\theta_W = .23$.

at LEP/SLC energies. Furthermore, the contact approximation (second column), which corresponds to the classical result of ref. [1] is good enough at LEP I/SLC energies but shows a larger deviation for higher center of mass energies. Last, one can see comparing the first and fourth columns that the cross-section is largely dominated by the Z^0 annihilation diagrams. The contribution of the W-exchange diagrams (difference between columns 1 and 4) is shown in fig. 2 as function of the center of mass energy. For energies between M_Z and 110 GeV it amounts to few per-cent of the total cross-section. Therefore, it is justified to compute radiative corrections only to the Z^0 annihilation diagrams. This will be the subject of the following sections.

3 Photonic corrections

We have divided the radiative corrections in two different gauge invariant groups: "photonic" corrections, that is, all the diagrams obtained from the lowest order ones by the attachment of photons in all possible ways, and "non-photonic" corrections which contain the details of the non-abelian structure of the theory.

As far as the photonic corrections concerns, they have no special theoretical interest but, as a common feature to all four-fermion neutral current processes around the Z^0 pole, they are the dominant ones. We have divided them into those containing virtual photons (fig. 3) and the ones where extra real photons are radiated (fig. 4).

At one-loop level, virtual photonic corrections are given by the interference of the diagrams shown in figure 3 with the ones contributing at lowest order. We have performed the calculation using the reduction techniques developed by Passarino and Veltman [5]. The results are in agreement with the ones shown in refs. [4] and [6]. The details of the calculation can be seen in [7].

The real photon corrections have been subdivided in "soft-photon" and "hard-photon" corrections. The first ones come from the diagrams of fig. 4 when the most externally attached photon has an energy smaller than certain cut-off $x_0 \cdot E_b$ being E_b the beam energy and x_0 a small number (typically around 10^{-2} to 10^{-3}). These corrections are computed analytically neglecting the soft-photon kinematical effects and combined with the virtual photon ones to give an infrared finite cross-section which depends on x_0 . This x_0 dependence is cancelled when the hard-photon corrections, that is, photons with energy larger than $x_0 \cdot E_b$ are included. These corrections have been computed without any approximation using the spinor technique and are explained in detail in [7-8]. When added together, the initial state radiative corrections are large in absolute value ($\sim -40\%$). This is a common feature to all four fermion processes mediated by neutral currents at energies close to the Z^0 pole. In general, the cross-section for a process $e^+e^- \rightarrow f\bar{f}$ ($f \neq e$) including one-loop initial state corrections can be expressed as

$$\sigma_1(s) = \sigma_0(s)(1 + \delta_1 + \beta \ln x_0) + \int_{x_0}^1 \beta \left(\frac{1}{x} - 1 + \frac{x}{2} \right) \sigma_0(s') dx \quad (1)$$

In the first term, which represents the virtual and soft-photon cross-section, x_0 is the soft-photon cut-off, β is defined as

$$\beta \equiv \frac{2\alpha}{\pi} \left(\ln \frac{s}{m_e^2} - 1 \right) \quad (2)$$

and δ_1 is the x_0 -independent correction given by

$$\delta = \frac{3}{4}\beta + \frac{\alpha}{\pi} \left(\frac{\pi^2}{3} - \frac{1}{2} \right) \quad (3)$$

In the second term, which corresponds to the hard photon contribution to the total cross-section, x is the photon energy normalized to the beam energy ($x = E_\gamma/E_b$) and $s' = s(1-x)$ is the effective center of mass energy squared. The function β factorizes the whole expression and can be regarded as an effective coupling constant for bremsstrahlung. At LEP/SLC energies it is large ($\beta \simeq .11$) but, since it has a smooth dependence with the center of mass energy, is of the same order as for the lower energies of PETRA/PEP and TRISTAN. However, since now we are close to a relatively narrow resonance, $\sigma_0(s)$ falls rapidly when far from the pole. Thus, the dominant part of the integral in (1) will have an effective upper limit of

$$x_M \sim \frac{\Gamma_Z/2}{E_b} \quad (4)$$

with Γ_Z the Z^0 width and E_b the beam energy. Therefore, the total cross-section will be approximately given by

$$\sigma_1(s) \simeq \sigma_0(s) \left(1 + \delta_1 + \beta \ln \frac{\Gamma_Z}{M_Z} \right) \quad (5)$$

Inserting standard values, one gets $\beta \ln \frac{\Gamma_Z}{M_Z} \sim -40\%$. It is clear that in the case of final state radiation or in the absence of resonances the restriction on x_M disappears and we shall have no large logarithmic terms². Therefore, in order to have the experimentally required accuracy, higher order corrections have to be considered.

We have implemented them in two different ways. The first one ("inductive exponentiation") is based on an explicit two-loop calculation of initial-state Q.E.D. corrections described in [9]. Analyzing in detail the various contributions, the second order cross-section can be expressed as:

$$\begin{aligned} \sigma_2(s) = & \sigma_0(s) \left[1 + \delta_1 + \delta_2 + \beta \ln x_0 + \delta_1 \beta \ln x_0 + \frac{1}{2} \beta^2 \ln^2 x_0 \right] \\ & + \int_{x_0}^1 \sigma_0(s') \left[\beta \left(\frac{1}{x} - 1 + \frac{x}{2} \right) (1 + \delta_1 + \beta \ln x) + \delta_2^H \right] dx \end{aligned} \quad (6)$$

where

$$\begin{aligned} \delta_2 & \equiv O(\alpha^2) \text{ correction independent of } x_0 \\ \delta_2^H & \equiv O(\alpha^2) \text{ non - leading hard correction} \\ s' & \equiv s(1-x) \end{aligned} \quad (7)$$

It is clear that the virtual and soft photon part shows an underlying structure of the type

$$\begin{aligned} \sigma_2^{VS}(s) & = \sigma_0(s) (1 + \delta_1 + \delta_2) \left(1 + \beta \ln x_0 + \frac{1}{2} \beta^2 \ln^2 x_0 + \dots \right) \\ & = \sigma_0(s) (1 + \delta_2 + \delta_2) x_0^\beta \end{aligned} \quad (8)$$

²This will not be the case if strong kinematic cuts are applied, like for instance small acolinearities in $e^+e^- \rightarrow \mu^+\mu^-$.

Similarly, for the hard-bremsstrahlung part one can induce an expression, consistent with the soft-photon one in the limit $x \rightarrow x_0$, of the form

$$\sigma_2(s)^H = \int_{x_0}^1 dx \left[\beta \left(\frac{1}{x} - 1 + \frac{x}{2} \right) x^\beta (1 + \delta_1 + \delta_2) + \delta_2^H \right] \sigma_0(s') \quad (9)$$

by resumming all the Sudakov-type logarithms. The sum of these two parts gives an expression of the total cross-section with the following features:

- The large infrared corrections, coming from the x_0 -dependent terms have been resummed to all orders
- The terms independent of x_0 are exact at the two-loop level. Thus, the precision will be of the size of δ_3 , that is, about one per mil.

This approach can be very easily implemented in any one-loop Monte Carlo generator. In our case, where σ_0 is the cross-section with one final state photon, this has been done with the following transformations:

- In the virtual and soft-photon part, we have substituted the infrared term $\beta \ln x_0$ in the correction by an overall multiplicative factor x_0^β .
- In the hard-photon part we have multiplied the whole expression by $x^\beta (1 + \delta_1)$. Certainly, this is not rigorous for hard-photon emission but, since terms of order β^2 are included correctly, the difference with respect to a rigorous treatment will be of order β^3 , that is, about few per mil.

Following this approach we are neglecting the effect of all photons with an energy smaller than $x_0 \cdot E_b$. However, this is not a problem in a practical case since, once we have an exponentiated expression, x_0 can be as small as we want. On the other hand, when the energy radiated is above this cut-off, we assume that it is carried away by one or two photons. This is an approximation since we are adding the contribution of an infinite number of radiated photons in at most two. However, for center of mass energies around the Z^0 pole, if the radiated energy is large enough it is carried away by one or at most two energetic photons.³ This effect is properly taken into account in the total cross-section but only approximately in the differential distributions being therefore accurate to few parts in 10^{-3} . The results of reference [11] confirm this statement.

The second approach is based on the Structure Functions formalism [12]. In this technique, the effect of initial state radiation is accounted for by convoluting the lowest order cross-section with the Structure Function for electrons and positrons in analogy with a Drell-Yan process in Q.C.D., that is

$$\sigma(s) = \int_{z_1^m}^1 dz_1 \int_{z_2^m}^1 dz_2 D_{e^+}(z_1, s) D_{e^-}(z_2, s) \sigma_0(s z_1 z_2) \quad (10)$$

³On the Z^0 pole, the probability to radiate an energy larger than 500 MeV in 3 or more photons is ≤ 2 per mil [10-11].

$\sqrt{s}$ (GeV)	92	94	100
σ^I (pb)	63.7 ± 0.2	175.5 ± 0.6	96.7 ± 0.4
σ^I (pb)	67.5 ± 0.3	184.5 ± 0.7	95.6 ± 0.5
σ^{SF} (pb)	67.6 ± 0.4	185.1 ± 1.3	94.8 ± 0.7

Table 2: Comparison between the one-loop cross-section and the exponentiated ones following the "inductive" and the "Structure Function" approaches.

where $D_e(z, s)$ is the probability to find an electron/positron with fractional energy

$$z = 1 - \sum_i \frac{E_{\gamma_i}}{\sqrt{s}/2} \quad (11)$$

at a momentum transfer squared $Q^2 = s$ and the lowest order cross-section is evaluated at the reduced center of mass energy $s' = sz_1z_2$. The functions $D_e(z, s)$ satisfy the Altareli-Parisi evolution equations and can be computed to the desired accuracy. In particular, it can be solved to all orders in the soft-photon limit (large z) leading to the well known Gribov-Lipatov solution [13]. The hard photon corrections up to order β^2 can be obtained by comparing a Taylor-expansion of the total cross-section in this limit with an explicit iterative second order solution of the evolution equations. Finally, the second order virtual corrections are added [12] by comparing the expansion with the complete second order calculation given in [9]. The result is:

$$\begin{aligned} D_e(z, s) = & \frac{1}{2}\beta(1-z)^{\frac{\beta}{2}-1}\Delta' - \frac{1}{4}(1+z)\beta \\ & + \frac{1}{32}\beta^2 \left[(1+z)[3\ln z - 4\ln(1-z)] - \frac{4}{1-z}\ln z - 5 - z \right] \end{aligned} \quad (12)$$

where $\Delta'^2 = \Delta + \frac{\beta^2\pi^2}{24}$ represents the virtual corrections up to order $O(\alpha^2)$ being

$$\begin{aligned} \Delta = & 1 + \left(\frac{\alpha}{\pi}\right) \left\{ \frac{3}{2}L + \frac{1}{3}\pi^2 - 2 \right\} \\ & + \left(\frac{\alpha}{\pi}\right)^2 \left\{ \left[\frac{9}{8} - 2\zeta(2) \right] L^2 + \left[-\frac{45}{16} + \frac{11}{2}\zeta(2) + 3\zeta(3) \right] L \right. \\ & \left. + \frac{6}{5}[\zeta(2)]^2 - \frac{9}{2}\zeta(3) - 6\zeta(2)\ln 2 + \frac{3}{8}\zeta(2) + \frac{57}{12} \right\} \end{aligned} \quad (13)$$

This is the expression we shall use to evaluate the total cross-section.

The Structure Functions approach justifies the previous "inductive" solution because, being more rigorous in the treatment of the infrared part, shows the same leading-log behaviour and, by construction, reproduces the same $O(\alpha^2)$ terms. A comparison between the two approaches is shown in table 2.

The first line (σ^I) correspond to the total cross-section at one-loop level. The second line (σ^I) to the one with the higher order corrections included in the inductive way and

the third one (σ^{SF}) to the Structure Functions approach. In all cases, the self-energy correction to the Z^0 propagator has been included. As one can see, there is a very good agreement (differences of the order of few per mil around the pole) in this range of energies between the two approaches used to include higher order effects. On the other hand, the corrections with respect to the one-loop calculations are sizeable (few per cent) especially at the pole and compensate the overestimation of the radiative effects by the first order calculation.

In order to obtain the differential cross-sections and an unweighted sample of events we have implemented the following Monte Carlo procedure:

- First, we generate the fractional energy of the electron (z_1) and the positron (z_2) in the collision according to an approximated distribution function

$$\tilde{D}_e(z, s) = \frac{\beta}{2}(1-z)^{\beta/2-1} \quad (14)$$

and compute the new center of mass energy $s' = sz_1z_2$ of the collision

- Then we generate the phase-space variables for the two neutrinos and the visible photon according to the differential tree level cross-section at the reduced center of mass energy s' and boost them to the lab reference frame.
- Next, we compute the weight of the event as

$$W = \frac{D_e(z_1, s) D_e(z_2, s)}{\tilde{D}_e(z_1, s) \tilde{D}_e(z_2, s)} \quad (15)$$

Computing the mean of those weights and multiplying by the integral of the approximant we will find the total cross-section and, with the help of a rejection algorithm, we will have a sample of unweighted events.

- Once we have an unweighted event, we compute the amount of energy taken by the non-visible photons: $x = 1 - z_1z_2$. If $x < x_0$, the event is accepted, otherwise, we throw away this event and generate an unweighted event according to the second order hard-photon differential cross-section weighted by $x^\beta(1 + \delta_1)$.

This second implementation should give more accurate differential cross-sections than the previous one because we have more kinematical information, that is, we know how the radiated energy is shared between the two initial state fermions. Furthermore, all the effects of collinear radiation are taken into account in a precise way. However, the transverse component of the radiation is not explicitly present since it has been integrated over. With our approach of taking the exact second order matrix element for the hard-photon radiation, we take it into account in a sensible way.

4 Non-photonic corrections

Among the "non-photonic" radiative corrections, the dominant ones at LEP/SLC energies are the corrections to the gauge boson propagators. They are universal in the sense that

$\sqrt{s}$ (GeV)	92	94	100
σ^{OL} (pb)	63.7 ± 0.2	175.5 ± 0.6	96.7 ± 0.4
σ^{QED} (pb)	67.5 ± 0.3	184.5 ± 0.7	95.6 ± 0.5
σ^* (pb)	66.4 ± 0.3	175.9 ± 0.8	93.3 ± 0.5

Table 3: Comparison between the one-loop corrected cross-section, the one-loop result with exponentiation and the exponentiated cross-section in the star-scheme.

do not depend on the specific process under consideration. To include them, we have used the star scheme [14] in which the gauge boson self-energies are reabsorbed, by means of the Dyson equations, in four finite universal running parameters:

$$\begin{aligned}
e_*^2(q^2) &= \frac{e_0^2}{1 - \Pi_{\gamma\gamma}(q^2)/q^2} \\
s_*^2(q^2) &= s_0^2 - s_0 c_0 \frac{\Pi_{Z\gamma}(q^2)}{q^2 - \Pi_{\gamma\gamma}} \\
G_{\mu\nu}^{-1}(q^2) &= G_{\mu 0}^{-1} - 4\sqrt{2} \left(\frac{s_0}{c_0} \Pi_{WW}(q^2) - \frac{s_0 c_0}{e_0} \Pi_{Z\gamma}(q^2) - \frac{s_0^2}{e_0^2} \Pi_{\gamma\gamma}(q^2) \right) \\
\rho_*^{-1}(q^2) &= \rho_0^{-1} - 4\sqrt{2} G_{\mu\nu}(q^2) (\Pi_{ZZ}(q^2) - \Pi_{WW}(q^2))
\end{aligned} \tag{16}$$

where Π_{ij} are the one-loop Dyson-resummed (i.e., leading-log terms included to all orders) gauge boson self-energy corrections and all quantities with subindex 0 denote "bare" quantities. These parameters are finite and renormalization scheme independent since they are functions of "bare" quantities only. For the rest, we shall assume that only Higgs-doublets are present ($\rho_0 = 1$) and thus we are left with only three independent running parameters ($e_*^2(q^2)$, $s_*^2(q^2)$, $G_{\mu\nu}(q^2)$) whose value has to be fixed at a given q^2 . Once this is done, all the relevant quantities like the neutral current matrix element and the Z^0 width can be rewritten in terms of the "starred" parameters. On doing so, all the leading-log terms of the self energies are automatically included and summed up to all orders. It is worth to note that since the "starred" parameters are universal, the running coupling constants which appear in the numerator of the neutral current matrix element cancel at the pole with those appearing in the Z^0 width giving a total cross-section which is insensitive to the self-energy corrections.

Finally, we have included the remaining $O(\alpha)$ and $O(\alpha_s)$ corrections to the width, which are known to be important on the Z^0 pole, in the way proposed in reference [15].

The effect of including "non-photon" radiative corrections is shown in table 3 for three different center of mass energies. The one-loop corrected cross-section is shown in the first line. The second one corresponds to the cross-section when only photonic corrections are included through the "inductive" approach (besides the self-energy corrections to the Z^0 boson). Finally the total cross-section including also the "non-photon" corrections within the "star" scheme is shown in the third line. It is clear from the table that Q.E.D. corrections are the dominant ones on the peak and therefore the total cross-section is larger than the one-loop corrected one. Slightly above the peak the two corrections are

about the same size and opposite signs so the global correction is almost zero. Finally, far above the peak, both corrections are negative and the total cross-section is smaller than the one at one-loop.

The differences in the total cross-section between the pure one-loop results and the ones obtained after applying all the corrections using the "inductive" and the "Structure Functions" approaches are shown in figure 5 for the previously referred set of cuts. The dotted line represents the one-loop result, the dashed line the one with the "Structure Functions" approach and the solid line represents the result of the "inductive" exponentiation. Their difference in the photonic energy distribution, $\cos(\theta)_\gamma$, and the photonic transverse momentum are shown respectively in figures 6,7 and 8 with the same convention for the lines that figure 5. In contrast to the angular distribution, which shows no significant differences, the reflected Z^0 peak in the photon energy distribution is enhanced with respect to the one-loop result. This affects as well the transverse momentum distribution.

5 Summary and conclusions

In the present paper we have discussed the effect of higher order (more than one-loop) corrections to the process $e^+e^- \rightarrow \nu\bar{\nu}\gamma$. They have been implemented in our previous Monte Carlo generator of unweighted events [3]. In particular, the soft-photon leading logarithmic terms have been included to all orders of perturbations following two different approaches, the "inductive" exponentiation and the Structure Function technique, both giving results which differ by few per mil around the pole. As far as non-photonic corrections concerns, by using the "star" scheme the leading logarithmic terms of the gauge-boson self-energies are resummed to all orders. We have also included the relevant corrections to the Z^0 width. Most of the topics and procedures discussed are valid for any general process $e^+e^- \rightarrow f\bar{f}$ ($f \neq e$). After these implementations the overall precision of our calculation, implemented in an unweighted Monte Carlo generator of events, can be conservatively estimated to be better than 1%.

Acknowledgements

We wish to thank Pascal Gay for his work in testing the Monte Carlo generator.

REFERENCES

- 1 K.J.F. Gaemers, R. Gastmans and F.M. Renard, Phys. Rev. D19 (1979) 1605.
- 2 G. Barbiellini, B. Richter and J.L. Siegrist, Phys. Lett. 106B (1981) 414.
- 3 F.A. Berends, G.J.H. Burgers, C. Mana, M. Martinez and W.L. van Neerven, Nucl. Phys. B301 (1988) 583.
- 4 M. Igarashi and N. Nakazawa, Nucl. Phys. B288 (1987) 301.
- 5 G. Passarino and M. Veltman, Nucl. Phys. B160 (1979) 151.
- 6 F.A. Berends, G.J.H. Burgers and W.L. van Neerven, Phys. Lett. 177B (1986) 191.
- 7 R. Miquel, Ph.D. Thesis, UAB-LFAE 89-02 (1989).
- 8 C. Mana, M. Martinez and F. Cornet, DESY prep. DESY 86-114 (1986).
- 9 F.A. Berends, G.J.H. Burgers and W.L. van Neerven, Phys. Lett. 185B (1987) 395.
- 10 M. Martinez, R. Miquel and C. Mana UAB prep. UAB-LFAE 89-06 (1989) (see also proceedings of the NATO Advanced Research Workshop held at the Univ. of Sussex, Brighton, 1989).
- 11 P. Colas, L. Mirabito and Z. Was, MPI-PAE/Exp. E1 (1989) 211.
- 12 E.A. Kuraev and V.S. Fadin, Sov. J. Nucl. Phys. 41/3 (1985) 466. O. Nicrosini and L. Trentadue, Phys. Lett. B196 (1987) 551.
- 13 V.N. Gribov and L.N. Lipatov, Sov. J. Nucl. Phys. 15 (1972) 438,675.
- 14 D.C. Kennedy and B.W. Lynn, SLAC-PUB-4039 (1988).
- 15 W. Wetzel, CERN Yellow Report, CERN 86-02 (Eds. J. Ellis and R. Peccei).

FIGURE CAPTIONS

- 1 Feynman diagrams of the process $e^+e^- \rightarrow \nu\bar{\nu}\gamma$ in the Born approximation.
- 2 Contribution of the W amplitudes in lowest order.
- 3 Feynman diagrams contributing to the photonic virtual corrections.
- 4 Feynman diagrams contributing to the real photon corrections.
- 5 Total cross-section in pb. as function of $\sqrt{s}$ in GeV for the process $e^+e^- \rightarrow \nu\bar{\nu}\gamma$ within the cuts specified in the text. The dotted line represents the one-loop result, the dashed line the one with the "Structure Functions" approach and the solid line represents the result of the "inductive" exponentiation.
- 6 Differential cross-section with respect to the photon energy normalized to the beam energy. The convention for the lines is the same as in figure 5.
- 7 Differential cross-section with respect to the photon $\cos\theta$. The convention for the lines is the same as in figure 5.
- 8 Differential cross-section with respect to the photon transverse momentum normalized to the beam energy. The convention for the lines is the same as in figure 5.

Fig.1.

Fig.3.

Fig.2.

Fig.4.

Fig.5.

Fig.6.

Fig.7.

Fig.8.