MEASUREMENT OF FREQUENCIES OF VARIOUS DECAY MODES
OF CHARMED PARTICLES D^0 , D^+ , D_s^+ AND Λ_c^+
INCLUDING THE OBSERVATION OF NEW CHANNELS

(ACCMOR Collaboration)

Amsterdam^{a)}-Bristol^{b)}-CERN^{c)}-Cracow^{d)}-Munich^{e)}-
Rutherford^{f)}-Valencia^{g)} Collaboration

S. Barlage^{e1}, H. Becker^{e2}, T. Böhringer^{c3}, M. Bosman^e, V. Castillo^{c4}, V. Chabaud^c,
C. Damerell^f, C. Daum^a, H. Dietl^e, A. Gillman^f, R. Gilmore^b, T. Gooch^b, L. Görlich^d,
P. Gras^g, Z. Hajduk^d, E. Higong^g, D.P. Kelsey^{c5}, R. Klanner^{e6}, S. Kwan^c, B. Lücking^e,
G. Lütjens^e, G. Lutz^e, J. Malos^b, W. Männer^e, E. Neugebauer^{e7}, H. Palka^d, M. Pepé^{f8},
J. Richardson^{f9}, K. Rybicki^d, H.J. Seebrunner^c, U. Stierlin^e, H.G. Tiecke^a,
G. Waltermann^e, S. Watts^h, P. Weilhammer^c, F. Wickens^f,
L.W. Wiggers^a, M. Witek^d and T. Zeludziejewicz^{d10}

ABSTRACT

In the CERN NA32 experiment a high-resolution silicon vertex detector and a purely topological approach have been used to investigate various decays of charmed particles. We observe ~ 620 fully reconstructed decays of D^0 in 12 channels and determine the branching ratios. For four D^0 decay modes involving a single (unseen) π^0 the small and narrow D^{*+} - D^0 mass difference is used to measure their branching ratios. We also observe ~ 280 fully reconstructed D^+ decays in 10 channels, ~ 90 D_s^+ decays in 11 channels as well as 160 Λ_c^+ decays in 7 channels and measure the corresponding branching ratios. In order to determine the branching fractions we use a sample of 584 events, each containing two decays consistent with associated production of charmed particles. We report measurements of the branching fractions for 20 decay channels of D^0 and 18 decay channels of D^+ . For the D_s^+ , we measure the branching fractions within a subset of 16 three- and five-prong decay channels. For the Λ_c^+ , we determine the branching fractions within a sample of 11 three- and five-prong decay channels, nearly all involving a proton.

Submitted to Zeitschrift für Physik C

a) NIKHEF-H, Amsterdam, NL	1) Now at KNMI, De Bilt, NL
b) Univ. of Bristol, Bristol, UK	2) Now at Gesamthochschule, Saarbrücken, FRG
c) CERN, Geneva, CH	3) Now at University of Lausanne, Lausanne, CH
d) Inst. of Nucl. Physics, Cracow, Poland	4) Now at Univ. of Valencia, Valencia, Spain
e) Max-Planck-Inst. f. Physik, Munich, FRG	5) Now at Rutherford Appleton Lab., Chilton, Didcot, UK
f) Rutherford Appleton Lab., Chilton, Didcot, UK	6) Now at DESY, Hamburg, FRG
g) IFIC, CSIC & Univ. of Valencia, Valencia, Spain	7) Now at Universität-GH Siegen, Siegen, FRG
h) Brunel Univ., Uxbridge, Mddx, UK	8) Now at CERN, Geneva, Switzerland
	9) Now at Univ. of Geneva, Geneva, CH
	10) Now at University of Melbourne, Melbourne, Australia

1. INTRODUCTION

In spite of great progress in the last decade, our knowledge of the decay modes of charmed particles is still far from being complete. Even for the best-known meson, D^0 , (throughout this paper a particle symbol stands for particle and antiparticle) the sum of all branching fractions^{*)} for hadronic decays listed in the last edition of the Review of Particle Properties (hereafter referred to as RPP [1]) amounts to only ~63%, far below ~85% expected from the branching fractions for the semileptonic decays. For D^+ , where we suffer from the difficulty of observation of one-prong decays, the corresponding numbers are ~33% and ~62%. For D_s^+ and Λ_c^+ , several branching ratios of Cabibbo-favoured decays have been determined; neither are Cabibbo-suppressed decays known nor have branching fractions been firmly established.

Especially difficult are measurements of branching fractions since they demand an absolute normalization. The MARK II collaboration [2] has introduced a novel method comparing the number of events in which a single D decay is identified ("single-tag" events) with the number of those in which a pair of Ds is reconstructed ("double-tag" events). This method is independent of the charm cross-section. The single-tag events determine the branching ratios while the double-tag events set the absolute scale of branching fractions. The results of MARK III dominate the field of branching fractions for the D^0 and D^+ mesons.

The LEBC-EHS collaboration [3] has obtained independent results using a topological normalization. However, their results suffer from low statistics; a branching fraction of ~0.5% corresponds to only one fully reconstructed event (i.e. a decay without unseen neutral products). The channels with unseen neutrals were statistically separated using the estimate of effective mass of neutral decay products. The estimate is defined as

$$m_o^2 = m_p^2 + m_{vis}^2 - 2 m_p \sqrt{m_{vis}^2 + (p_t^{vis})^2} \quad \{1\}$$

where m_{vis} is the effective mass of the charged decay products, m_p is the mass of the parent charmed particle and p_t^{vis} is the total visible transverse momentum with respect to the direction of the parent.

For this class of events, a single decay in the LEBC-EHS study represents a branching fraction of ~2%.

---

*) Branching fractions are understood as absolute frequencies (expressed in percentages) of decay modes. The ratio of two branching fractions, which we call a branching ratio, represents the relative frequency of two decay modes.

Preliminary results on branching ratios from our experiment were presented at the Munich Conference [4]. An analysis of D^0 decay modes based on our final sample was recently published [5]. In that study single-tag events (one fully reconstructed decay vertex) were used for the measurement of branching ratios as well as for the observation of new decay modes. Then the double-tag sample allowed the determination of branching fractions. The sample consisted of 84 events containing any decay of the D^0 (mostly involving unseen neutrals) and a fully reconstructed decay of $\bar{D}^0$, D^- , $\bar{D}_s^-$ or $\bar{\Lambda}_c^-$.

This paper describes a similar study for the four charmed particles; D^0 , D^+ , D_s^+ and Λ_c^+ . Now we use a larger sample of double-tag events than in ref. [5], namely all events consistent with associated charm production (two decay vertices, fully reconstructed or not). In the sample a single decay of D^0 or D^+ represents a branching fraction of 0.1% to 0.2%. The paper is organized as follows :

Section 2 describes the apparatus, the data processing and the Monte Carlo corrections. In section 3, branching ratios are measured for single-tag events involving decay modes of D^+ , D_s^+ and Λ_c^+ including either charged products only or charged products and a reconstructed V^0 . Additionally in section 4 we determine the branching ratios of four D^0 decay modes involving a single π^0 . Section 5 is devoted to the double-tag sample and to the determination of branching fractions. The results are discussed in section 6. Finally, conclusions are listed in section 7.

2. APPARATUS, DATA PROCESSING AND ACCEPTANCE CORRECTIONS

2.1 *The Apparatus*

The second phase of the NA32 experiment was performed at the CERN SPS using a negative 230 GeV/c beam (96% pions and 4% kaons). Charm decays were reconstructed with an improved silicon vertex detector and a large acceptance spectrometer (see Fig. 1). The latter consisted of two magnets, 48 planes of drift chambers arranged in four groups and three multicellular Cerenkov counters allowing π, K, p identification in the momentum range of (4-80) GeV/c. A two-level trigger, described in ref. [6], selected preferentially events with a (pK^-) or a (K^+K^-) pair, since this experiment was originally intended to study $\Lambda_c^+ \rightarrow pK^-\pi^+$ and $D_s^+ \rightarrow K^+K^-\pi^+$ decays; in fact a gain in sensitivity of about five has been achieved for these decays. Nevertheless we have also observed charm signals in many other channels. This is due to associated charm production often yielding two kaons of opposite charge, to pions misidentified as kaons at the trigger level and finally to kaons or protons from the primary vertex.

The vertex detector consisted of a beam telescope with seven silicon microstrip detectors (MSDs) and a vertex telescope of two charge-coupled devices (CCDs) at 10 and 20 mm down-

stream of a thin 2.5 mm Cu target followed by eight MSDs starting from 65 mm behind the target. The CCDs allow the measurements of space points with a precision of $\sim 5 \mu\text{m}$ in two orthogonal directions, while the two-track resolution is $\sim 40 \mu\text{m}$. The precise space-point information from the CCDs, very important for the observation of short-lived charmed baryons decaying in vacuum immediately behind the target, was also essential for our study of double-tag events. More details about the apparatus can be found in ref. [7].

The results discussed in this paper are based on the full sample of ~ 17 million triggers collected in 1985-86.

2.2 Data Processing

A very precise track measurement in our vertex detector allows a purely topological charm search, which is restricted neither to a limited number of decay modes nor to any mass window around m_P (see Equ. {1}), thus allowing the determination of charm branching fractions. The reconstruction is done in five steps (see ref. [7] for more details).

First, tracks are reconstructed in the drift chambers of the spectrometer.

In the second step, the beam track and all outgoing tracks are reconstructed in the beam and vertex telescopes, without using the drift chamber information. Then, tracks found in the drift chambers and in the vertex telescope are matched. Unmatched drift chamber tracks are used as a loose guidance to recover additional tracks out of complex clusters of signals in the vertex telescope. Particles are identified using the information of the Cerenkov counters. We obtain probabilities for each particle being a pion, a kaon or a proton. Finally, reconstruction of the primary vertex is performed. First all tracks are taken together. For each secondary track we calculate its impact parameter (the distance of closest approach in space to the primary vertex), its error and the probability that it belongs to the primary vertex. If for any track this probability is less than 1%, we remove the track with the smallest probability and repeat the procedure.

A loose selection of charm decays is made in the third step. We demand the primary vertex to be inside the Cu target and at least two tracks not originating from the primary vertex or one such track and a K^0 or Λ^0 reconstructed in the drift chambers. This step reduces the original sample by a factor of seven.

The fourth step, a search for secondary vertices, reduces the sample by a further factor of four. The tracks not belonging to the primary vertex are used to reconstruct one or more secondary vertices. We demand secondary vertices to be downstream of the Cu target, outside of the CCDs ($380 \mu\text{m}$ thick) and upstream of the first MSD. Tracks, which were originally attributed to the primary vertex are checked for compatibility with any of the secondary ver-

tices. In case of ambiguity, we keep all solutions ascribing a probability to each of them (topological probability). Then we try to identify the parent charmed particle in each decay vertex. In some cases this is not unique (e.g. $D_s^+ \rightarrow K^+K^-\pi^+$ can be confused with $D^+ \rightarrow K^-\pi^+\pi^+$ and/or $\Lambda_c^+ \rightarrow pK^-\pi^+$) and we calculate the probability of each identification from the Cerenkov information for the decay products. In most cases there is no ambiguity; otherwise we select the solution with the largest total (topological and identification) probability. In this way we remove the vast majority of reflections.

Events passing the fourth step and having two or more secondary vertices will be further analysed in section 5, which is devoted to the double-tag sample.

The fifth and last step is a search for fully reconstructed charm vertices. We demand that the total momentum vector of the decay originates from the primary vertex with a probability exceeding 1%. For decays involving a V^0 , the impact parameter of the total momentum vector has to be less than 80 μm . Finally, we check whether the particle identification in the secondary vertex corresponds to any decay mode of D^0 , D^+ , D_s^+ or Λ_c^+ (the Ξ_c^+ and Ξ_c^0 are described in refs. [8,9]). We consider all decay modes with prong number (number of charged secondaries) from two to six and without any π^0 . The results for D^0 have already been published [5]; the remaining single-tag events (D^+ , D_s^+ and Λ_c^+) will be discussed in section 3.

Concluding this section, we would like to stress that the effective mass of secondaries coming from the decay vertex has not been used in the above selection nor was it restricted to any particular decay modes. This enables us to study a larger number of decay modes than in any previous work. It also makes possible an analysis of double-tag events in a fixed-target experiment. This is due to the excellent spatial resolution of silicon detectors and to the purely topological approach.

2.3 Acceptance Corrections

Simulation of the geometrical acceptance of our apparatus, of the trigger and of the selection criteria requires a complex Monte Carlo program. This program generates an uncorrelated pair of charmed particles, $P_i\bar{P}_j$. For the simulation of double-tag events these particles decay into the observed channels, and we repeat the calculation of the acceptance A_{ij} for each combination of decays. For single-tag events P_i is the decay in question and $\bar{P}_j$ is always a mixture of $\bar{D}^0$ and D^- (with the empirical ratio of 2:1) decaying according to the branching fractions from ref. [1]. Each decay is generated from a phase-space distribution. The distributions of the Feynman variable x_F and of the transverse momentum of P_i or $\bar{P}_j$ [10] as well as the lifetime of Λ_c^+ [7] are taken from our experiment; the lifetimes of other particles come from ref. [1]. The generated pair of charmed particles is mixed with tracks from an event randomly chosen from an interaction-trigger sample recorded with our apparatus, thus faking a real event

(the sum of momenta of all tracks is required not to exceed 230 GeV/c). All tracks from such an event are subsequently traced through the apparatus and subjected to the same cuts as those made in our analysis. In particular we demand that

- i) the trigger conditions have to be satisfied by a pair of oppositely charged kaons/protons or by pions mistakenly identified as heavier hadrons in our apparatus; these particles may come either from charm decay or from the primary identification..
- ii) all decay tracks must be seen in the vertex telescope and in the first group of drift chambers.
- iii) any kaon/proton from the decay has to pass the second-magnet exit window and be seen in the Cerenkov counters and in the drift chambers downstream; the kaon decay probability and the kaon/proton interaction probability are also taken into account.
- iv) all the topological cuts used for vertex selection are applied.
- v) the V^0 decay products (if any) must be seen in the first group of drift chambers, one of them has to pass the second-magnet exit window and be seen in the drift chambers downstream.

The distributions of multiplicity, momenta, impact parameters and other characteristics for accepted events agree with those for real ones (for more details see ref. [10]). The values of $x_F > 0$ acceptance for single-tag events are given in Tables 1-4. It should be pointed out that the results on branching ratios and branching fractions are less sensitive to the details of the Monte Carlo program than e.g. the cross-section for charm production, because some of the systematic effects cancel in the ratios.

In addition to the acceptance corrections, the Monte Carlo program is also used for two purposes connected with our analysis of double-tag events.

If P_i decays into a channel involving neutral particles, we obtain the distributions of m_{vis} and m_0^2 for accepted events. This knowledge is essential for disentangling the various decay modes of charmed particles (see section 5.2).

If P_i decays semileptonically, we can determine the contamination of the corresponding hadronic channels (our identification procedure does not discriminate between pions and leptons). For accepted events we attribute the pion mass to electrons or muons, and then calculate m_{vis} and m_0^2 to find the hadronic channels being contaminated. Knowing the semileptonic branching fractions, we can thus calculate corrections for the hadronic channels (see section 5.4).

3. DETERMINATION OF BRANCHING RATIOS FOR D^+ , D_s^+ AND Λ_c^+

The single-tag events have been selected in section 2.2 without any restriction on m_{vis} . Now we look at the mass plots and search for charm signals. This is done for all possible decay modes with two to six charged secondaries and without any π^0 . Some channels with a single π^0 will be studied in the next section devoted to D^0 ; in this section we deal with charged charmed particles decaying into three or five charged secondaries. The decay modes involving an additional reconstructed V^0 (K^0 or Λ^0) are also included here. *In the following subsections we deal also with some channels for which no significant signal was seen. In these cases upper limits can be calculated. Moreover, where these channels have been studied in the double-tag sample, both measurements are combined. This may result in an improvement of the upper limit or in the confirmation of a signal.*

3.1 *Branching Ratios for the D^+ Meson*

The invariant mass distributions for ten decay modes of the D^+ are shown in Fig. 2; in some plots the D_s^+ is also visible. We fit the distributions by a Gaussian signal (or two signals) and a linear background. The results for signals are shown in Table 1. Channels studied for the first time (e.g. the five-body decay $D^+ \rightarrow K^-\pi^+\pi^+\pi^+\pi^-$) are marked with asterisks. The $(K^+K^-)_{nr}$ means that the effective mass of the kaon pair is outside the resonance (in this case the ϕ) region.

In Table 1 we also show the acceptance values. For channels involving a K^0 these include the relevant factors for the K_S^0 and its decay into $\pi^+\pi^-$ as well as the efficiency of our V^0 reconstruction package estimated to be $(83 \pm 7)\%$. The contribution of this error is shown as the systematic error in Table 1. The branching ratios are calculated with respect to the $D^+ \rightarrow K^-\pi^+\pi^+$ channel. It is seen in Table 1 that they are consistent with the results of other experiments.

3.2 *Branching Ratios for D_s^+*

In Fig. 3 we show the invariant mass distributions for those decay modes of the D_s^+ which have not been shown in Fig. 2. In Table 2 all results are summarized using $D_s^+ \rightarrow \phi\pi^+$ as the reference mode. Hereafter our branching ratios are corrected for the unobserved $\phi \rightarrow K^0\bar{K}^0$ and $K^{*0} \rightarrow K^0\pi^0$ decays whenever needed. A row in brackets always refers to a sub-channel of the mode described in the preceding row. In general, our branching ratios are consistent with those of other experiments. We observe a significant signal in the $D_s^+ \rightarrow K^-K^0\pi^+\pi^+$ channel which is dominated by $D_s^+ \rightarrow K^{*+}\bar{K}^{*0}$.

3.3 Branching Ratios for Λ_c^+

Figure 4 shows the invariant mass distributions for the Λ_c^+ decay modes. The branching ratios in Table 3 are calculated with respect to $\Lambda_c^+ \rightarrow pK^-\pi^+$. A study of this decay channel shows a considerable $p\bar{K}^{*0}$ component (see Fig. 5a) amounting to $(24 \pm 7)\%$. Contrary to the results quoted in ref. [1], we do not observe any significant $\Delta^{++}K^-$ component (see Fig. 5b). There is an indication of a signal in three Cabibbo-suppressed modes, i.e. $\Lambda_c^+ \rightarrow pf_0(975)$, $\Lambda_c^+ \rightarrow pK^+K^-$ (or $\Lambda_c^+ \rightarrow p\phi$) and $\Lambda_c^+ \rightarrow p\pi^+\pi^+\pi^-\pi^-$. It is interesting to note that the $f_0(975)$ has already been observed by the Tagged Photon Spectrometer (TPS) collaboration [11] in the $D_s^+ \rightarrow \pi^+\pi^+\pi^-$ channel. Our indication, if confirmed, could mean that also in the Λ_c^+ decays the $\bar{s}s$ final state is more likely than $\bar{u}u$ or $\bar{d}d$ just as in the Cabibbo-suppressed D^0 decays where the K^+K^- final state is much more likely than the $\pi^+\pi^-$ one. Here also our results are consistent with the Λ_c^+ branching ratios measured in other experiments.

4. BRANCHING RATIOS OF THE D^0 MESON

D^0 branching ratios have already been measured from fully reconstructed decays in our sample (see ref. [5] and Table 6). In this section we determine branching ratios for another class of D^0 decay modes, namely those with a single π^0 . Our apparatus does not register π^0 s. However, one can use the D^{*+} - D^0 mass difference to identify these events. For each charged pion from the primary vertex ("external" pion) we calculate m_{vis}^π , the effective mass of the pion with all charged particles originating from the D^0 vertex. Our Monte Carlo calculation shows that the spread of $\Delta m = m_{vis}^\pi - m_{vis}$ around the nominal value of 145.4 MeV is reasonably small (see Table 4), even for D^0 decay modes involving a π^0 . Thus a peaking of Δm distribution in this region is a signature of $D^{*+} \rightarrow D^0\pi^+$ decay. This is, however, valid only for sufficiently high m_{vis} ; we take $m_{vis} > 1300$ MeV. For $D^0 \rightarrow \pi^+\pi^-\pi^0$ and $D^0 \rightarrow \pi^+\pi^+\pi^-\pi^0$ this is automatically satisfied since we demand $m_{vis} > m_D - m_K$; otherwise a Cabibbo-favoured interpretation would be possible (see section 5.2). For $D^0 \rightarrow K^-\pi^+\pi^+\pi^-\pi^0$ there are very few events with $m_{vis} < 1300$ MeV, so it is only for a $D^0 \rightarrow K^-\pi^+\pi^0$ channel that the above cut considerably reduces the acceptance. For the last mode we also demand $m_0^2 < 0.25 \text{ GeV}^2$ (see equ. {1}) to separate this mode from the $D^0 \rightarrow K^-\pi^+\pi^0\pi^0$ (see section 5.2). All these cuts are used when calculating the acceptance quoted in Table 4.

For each secondary vertex fulfilling the above criteria, we calculate Δm for each pion from the primary vertex. For $D^0 \rightarrow K^-\pi^+\pi^0$ as well as for $D^0 \rightarrow K^-\pi^+\pi^+\pi^-\pi^0$, the charge of the "external" pion must be opposite to that of the kaon ("right charge" combination); the "wrong charge" combination, e.g. $(K^-\pi^+\pi^0)\pi^-$ can be used for the background estimation. The same is true for $D^0 \rightarrow K^-\pi^+\pi^+\pi^-$ which is used as a reference channel. For $D^0 \rightarrow \pi^+\pi^-\pi^0$ as well as for $D^0 \rightarrow \pi^+\pi^+\pi^-\pi^0$ both charges of pions must be taken and the background estimation can be based only on the combined Δm ("right" + "wrong charge") plot.

The Δm distributions are shown in Fig. 6 and the results are listed in Table 4. As expected, the D^* signal is less significant in the Cabibbo-suppressed channels $D^0 \rightarrow \pi^+\pi^-\pi^0$ and $D^0 \rightarrow \pi^+\pi^+\pi^-\pi^0$. The branching ratios determined in this section together with those published previously [5] for the D^0 will be used in section 5.5.

5. STUDY OF DOUBLE-TAG EVENTS

5.1 *Final Selection of Events*

Since only $\sim 12\%$ of decays in our double-tag sample are fully reconstructed and for the rest we cannot apply the powerful constraint of $m_{\text{vis}} = m_p$, the sample is further cleaned by the following cuts :

- i) in order to avoid a contamination by secondary interactions the position of each secondary vertex must be at least two standard deviations away from the Cu target edge and from the edges of both CCDs,
- ii) decays attributed to short-lived D^0 , D_s^+ and Λ_c^+ must occur upstream of the second CCD (the resulting loss of genuine events is only $\sim 2\%$ and we gain important CCD information),
- iii) $\pi^+\pi^-$ vertices are accepted only if $m_{\text{vis}} > 580$ MeV in order to remove all K^0 decays (the resulting loss of $D^0 \rightarrow \bar{K}^0\pi^+\pi^-$ etc. decays is taken into account in our acceptance corrections). The value of 580 MeV comes from a detailed inspection of the tail of the very high statistics K^0 mass distribution seen in our apparatus. Because of pion/kaon (proton/kaon) ambiguity some K^0 (Λ^0) decays imitate $D^0 \rightarrow K^-\pi^+\pi^0\pi^0$ events. They are also rejected from our sample. Similarly we remove $K^+ \rightarrow \pi^+\pi^+\pi^-$ decays.
- iv) the "visible" transverse momentum p_t^{vis} with respect to the direction of the parent charmed particle cannot exceed the maximum transverse momentum kinematically allowed for any particle in a decay channel under consideration,
- v) in addition to the "normal" tracks reconstructed both in the vertex telescope and in the drift chambers, we also find tracks seen in the vertex telescope (including CCDs) but not in the drift chambers and thus of unknown momentum. If such a track does not belong to the primary vertex but can be assigned to the secondary vertex, then the latter vertex is rejected. In this way we remove part of the background described in section 5.3.

After this selection we are left with 673 double-tag events.

5.2 Assignment of Vertices to Decay Modes

The small fraction of decay vertices in the double-tag sample corresponding to fully reconstructed charm decays forces us to identify most channels using the values of m_{vis} and m_0^2 (see equ. {1}). In case of ambiguity a Cabibbo-favoured interpretation is always preferred to a Cabibbo-suppressed one. A $\pi^+\pi^-$ vertex for example is assigned to the $D^0 \rightarrow \pi^+\pi^-\pi^0$ channel only if $m_{\text{vis}} > m_D - m_K$; otherwise the vertex is interpreted as $D^0 \rightarrow \bar{K}^0\pi^+\pi^-$ or $D^0 \rightarrow \bar{K}^0\pi^+\pi^-\pi^0$. Obviously, this overestimates the latter channel at the expense of the former. A corresponding correction is based on the Monte Carlo distribution of m_{vis} for $D^0 \rightarrow \pi^+\pi^-\pi^0$ decays passing our apparatus and selection criteria (see section 2.3). From the distribution we calculate the ratio r_i of $D^0 \rightarrow \pi^+\pi^-\pi^0$ decays with m_{vis} below $m_D - m_K$ to that above this value. Then the total weight w_i (see equ. {3} in section 5.4) of the $D^0 \rightarrow \pi^+\pi^-\pi^0$ channel is multiplied by $(1+r_i)$ and a value of $r_i w_i$ is subtracted from the channels involving a $\bar{K}^0$. We treat the $\pi^+\pi^+\pi^-$ vertices analogously. They are assigned to the decay $D_s^+ \rightarrow \pi^+\pi^+\pi^-\pi^0$ or to the channel $D^+ \rightarrow \pi^+\pi^+\pi^-\pi^0$ only if $m_{\text{vis}} > m_D - m_K$, and a similar comparison is made. However, contrary to the D^0 decays, there is an important complication resulting from the fact that the parent charmed particle is unknown. This is solved with the help of the m_0^2 parameter.

The m_0^2 parameter has been used by the LEBC-EHS collaboration [3] to discriminate statistically between channels with one or more neutral decay products. We follow and develop this method which is now applied to more channels as well as to the statistical identification of the parent charmed particle P (remember that its mass m_P enters explicitly equ. {1} for the m_0^2). If we want to discriminate between channel A and channel B, we produce Monte Carlo distributions of the m_0^2 for both channels and find a value of m_{cut}^2 such that the fraction s_i of channel A decays with $m_0^2 > m_{\text{cut}}^2$ is equal to that of channel B with $m_0^2 < m_{\text{cut}}^2$. Then we assign decays with $m_0^2 < m_{\text{cut}}^2$ to channel A and the remaining ones to channel B. Though the assignment is wrong for some decays (the fraction s_i represents the confusion level), it is statistically correct provided both channels are of similar abundance. The values of m_{cut}^2 and s_i used in our analysis are shown in Table 5. The values of m_{cut}^2 and $p = 1 - s_i$ of the LEBC-EHS collaboration for the various channels is very similar to ours. Two remarks should be made for the rows 5 -7 of Table 5, where we do not *a priori* know the parent particle. First, we calculate the m_0^2 assuming the parent particle to be D_s^+ both for real events and in our Monte Carlo. Secondly, m_0^2 is sensitive to the difference in m_P only for high values of m_{vis} . But, as explained above, it is only for these values that the separation is made.

As it has already been mentioned, the above separation is exactly valid only if channel A and channel B are of equal abundance. Since this is not necessarily the case, we have performed a more detailed analysis for the high-statistics decay modes. Namely, we have simulated the m_{vis} distributions for two or more channels with the same charged decay products but

with various neutral ones. Then we have fitted the observed distributions of m_{vis} and m_0^2 , the relative contributions of various channels being the free parameters. This was done for a cleaner subsample of events fulfilling additional selection criteria (see section 5.3). The results for $\pi^+\pi^-$ and $K^-\pi^+$ vertices confirm those obtained with the help of the m_{cut}^2 . For the latter channel the fit was consistent with no significant contribution of a Cabibbo-suppressed channel $D^0 \rightarrow K^-K^0\pi^+$ which has also not been observed in the single tags. The relevant upper limit is well above the value quoted in ref. [1]. Also for the $\pi^+\pi^+\pi^-$ vertices, we confirm the results obtained using the m_{cut}^2 parameter and obtain only an upper limit for the $\Lambda_c^+ \rightarrow \Lambda^0\pi^+\pi^+\pi^-$ channel seen in the single tag sample. It is only for the $K^-\pi^+\pi^+$ vertices that our fit demands three components, i.e. $D^+ \rightarrow K^-\pi^+\pi^+\pi^0$, $D^+ \rightarrow K^-K^0\pi^+\pi^+$ (Cabibbo-suppressed) and $D_s^+ \rightarrow K^-K^0\pi^+\pi^+$. For the fourth possible channel i.e. $D^+ \rightarrow K^-\pi^+\pi^+\pi^0\pi^0$, we obtain only an upper limit.

Unfortunately, even such a fit is not sensitive enough to discriminate between channels with two or three neutral decay products. However the latter are likely to be much weaker than the former (cf. section 5.5)

We have partially checked this procedure using single-tag events with an observed K^0 . For each event we have calculated m_0^2 ignoring the K^0 and we have counted events with $m_0^2 > m_{\text{cut}}^2$. The fraction t_i of such events, representing a direct measurement of the confusion level, is shown in the last column of Table 5. Within statistics the agreement is fair, especially considering the high background in the $D^+ \rightarrow \bar{K}^0\pi^+\pi^+\pi^-$ channel (see Fig. 2b).

After assigning all decay vertices we are left with 584 events consistent with a charm-anti-charm interpretation, 58 charm-charm (or anticharm-anticharm) pairs and 31 events in which one of the vertices cannot be assigned to any charm decay. Six of the last sample are easily explained by the loss of a single track going outside of our acceptance (e.g. a pK^- vertex is probably due to a $\Lambda_c^+ \rightarrow pK^-\pi^+$ decay, the pion being lost). The loss of such events is included in our acceptance corrections. For the remaining 25, usually four- and six-prong vertices with m_{vis} exceeding that of charmed particles we have no obvious explanation. Some of them can be due to mistakes of our reconstruction program erroneously assigning a track to a secondary vertex or even joining two vertices together and others could be decays of particles containing a bottom quark with a subsequent decay of a charmed decay product. Since the last conjecture involves neutral decay products in both vertices, one cannot make any definite statement.

Ignoring these events, we are left with 642 double-tag events in which all 1284 secondary vertices can be interpreted in terms of decays of D^0 , D^+ , D_s^+ or Λ_c^+ . Among these there are two events with three secondary vertices and at least 58 events in which both parent particles contain a charmed quark of the same sign. This is a clear demonstration of double associated charm production.

5.3 Corrections for Background

For the final analysis we use only the 584 events consistent with the production of two particles of opposite charm. The identification of the charmed quark in the parent particle is trivial for D^+ , D_s^+ and Λ_c^+ since the sign of the parent charge is identical to that of the charmed quark. It is also obvious for D^0 decays involving a charged kaon ("tagged D^0 "). However, for the "non-tagged" D^0 decays into none or two charged kaons (including all Cabibbo-suppressed channels) both signs of charmed quark are possible. In such a case we assume the sign opposite to that in the other secondary vertex. In this way, we overestimate the frequencies of the above modes by including some charm-charm pairs into the charm-anticharm sample. In order to calculate the corresponding correction we count the numbers of tagged D^0 s paired with another particle of definite charm (D^+ , D_s^+ , Λ_c^+ or another tagged D^0). There are 253 such decays in the charm-anticharm sample and 37 in the charm-charm one. Assuming that the same ratio holds for non-tagged D^0 (those with none or two charged kaons) we multiply the total weight w_i (see equ. {3}) for such channels by $253/(37+253) = 0.87$ to remove the charm-charm contamination.

Potentially, the most dangerous background is due to charm vertices with one or more charged decay products going outside our acceptance. This could for example shift a $D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-$ event into the $D^+ \rightarrow K^- \pi^+ \pi^+ \pi^0$ category. The resulting loss in the D^0 channel is included in our acceptance but the unwanted "gain" of D^+ is not and the corresponding calculation would need some nontrivial assumptions about the charm production characteristics. This background has been reduced by the cuts iv) and v) in section 5.1, nevertheless some background is still expected to remain and to distort our results. We expect this background to occur in two-prong D^0 decays and three-prong D^+ or D_s^+ decays. Four-prong D^0 should contain very little background because the five-prong D^+ and D_s^+ decay channels are fairly weak (not to mention the six-prong D^0). The Λ_c^+ decays are usually protected from the background by demanding a proton among the charged decay products. We expect the background to be relatively strong at high m_0^2 .

In order to determine this background, we proceed as follows. One constructs a plane defined by the beam direction and the direction of the parent charmed particle P and studies the projection $p_{\text{proj}}^{\text{vis}}$ of the total visible momentum onto this plane (see Fig. 7). If the angle θ of $p_{\text{proj}}^{\text{vis}}$ with respect to the direction of the parent particle is positive (i.e. directed away from the axis of our apparatus) the secondary particle needed to balance the transverse momentum goes into the apparatus and should be registered if it was a charged one. Therefore this category of decays (we denote their number in the i -th channel by N_i^+) is clean, and it was this subsample which was used in the fits to m_{vis} and m_0^2 distributions in section 5.2. However, if the angle θ is negative (this happens for N_i^- decays) the particle demanded by the transverse momentum

conservation could go outside our acceptance and thus be lost even if charged. If this is the case, then we expect $N_i^+ < N_i^-$. If there is no background, we get $N_i^+ \equiv N_i^-$. The case $N_i^+ > N_i^-$ can only be due to fluctuations. In fact, for all our channels N_i^+ never exceeds N_i^- by more than 1.2 standard deviations. For each decay channel, we introduce a correction factor f_i defined as

$$f_i = (2N_i^+) / (N_i^+ + N_i^-) \quad \{2\}$$

For a complete sample of fully reconstructed decays θ fluctuates around zero, $N^+ = 70$, $N^- = 76$ and

$$f = \frac{2 \times 70}{70 + 76} = 0.96 \pm 0.08 \cong 1$$

as expected. The values of f_i for other channels are given in Tables 6 - 9. It is seen that for some channels f_i is significantly lower than unity. This happens indeed mainly for two- and three-prong channels. As expected, the most significant deviation occurs for channels with the largest values of m_0^2 , i.e. for $D^0 \rightarrow \bar{K}^0 \pi^+ \pi^- \pi^0$ and $D^+ \rightarrow \bar{K}^0 \pi^+ \pi^+ \pi^- \pi^0$.

If f_i is smaller than unity by at least 1.2 standard deviations, we replace the total weight w_i (see equ. {4}) for a given channel by $f_i w_i$. An alternative solution is to consider only vertices with $\theta > 0$; it yields very similar results but with larger statistical errors.

5.4 Calculation of Branching Fractions, Discussion of Systematic Errors

For each decay channel, we calculate the total weight as

$$w_i = \sum_j (N_{ij} / A_{ij}) \quad \{3\}$$

where N_{ij} is the number of events in which one of the vertices belongs to the i -th channel and another to the j -th one.

A_{ij} is the acceptance for such a combination.

These weights w_i are then corrected as explained in section 5.3.

Then the branching fractions BF_i are calculated as

$$BF_i = \left[(1 - \mathcal{L}) \times w_i \right] / \sum_j w_j \quad \{4\}$$

where $\mathcal{L}$ is the fraction of zero- or one-prong channels which are not observed in our experiment. Thus we normalize the branching fractions to $(1 - \mathcal{L})$. For D^0 $\mathcal{L}_0 = (10.0 \pm 1.7)\%$ (average from the results quoted by Hitlin [18] so that $1 - \mathcal{L}_0 = (90.0 \pm 1.7)\%$. The resulting relative error of $\sim 2\%$ is much smaller than those in Table 6. For D^+ $\mathcal{L}_1 = (55 \pm 4)\%$ also from ref.[18]; thus $1 - \mathcal{L}_1 = (45 \pm 4)\%$. The resulting systematic error of 9% is still smaller than those in Table 7.

The situation is much worse for D_s^+ and Λ_c^+ since $\mathcal{L}_1$ is completely unknown. Therefore in Tables 8 and 9 we give the branching fractions within our subsample of events. For the D_s^+ meson this consists of three- and five-prong decays. For the Λ_c^+ it is the three- and five-prong decays involving a proton as well as the decay $\Lambda_c^+ \rightarrow \Lambda^0 \pi^+ \pi^+ \pi^-$ which form our subsample. This will be further discussed in section 6.3.

A similar hierarchy of quality of results is valid for the semileptonic decay channels. For the D^0 , the channels are fairly well known (see ref.[19] and references therein) and the corresponding corrections (cf. section 2.3) introduce reasonably small errors. For the D^+ the situation is worse, the errors are larger (see ref.[20] and references therein) but the correction is still possible. However, for the D_s^+ only the sum of all semileptonic branching fractions has been measured and found to be $(9 \pm 9 \pm 1.4)\%$ [12]. From the experimental near-equality of the lifetimes of D^0 and D_s^+ we expect this fraction to be $\sim 8\%$. Since we expect several semileptonic decay modes their individual branching fractions would be of the order of $(1-2)\%$ which could lead to considerable changes in our results. The problem is less serious for the Λ_c^+ since the inclusive semileptonic branching fraction is only $(4.5 \pm 1.7)\%$ [1].

5.5 Global Fits and Problem of Channels with Many Neutrals

For each charmed particle we perform a global fit using all branching ratios (with relatively small errors) and branching fractions (with large errors). The fit rearranges the branching fractions within their large errors so that the ratios would agree with the branching ratios determined in section 3 and section 4. The fit results are shown in Tables 6-9. The χ^2/NDF values are 25/15 for the D^0 , 12/9 for the D^+ , 8/9 for the D_s^+ and 5/6 for the Λ_c^+ . It should be stressed that only the values of $\mathcal{L}_0$ and $\mathcal{L}_1$ (corrections for lost low-multiplicity decays) as well as subtraction of semileptonic channels are based on the results of other studies; everything else comes from our experiment. The changes of $\mathcal{L}_0$ and $\mathcal{L}_1$ would scale our results up or down without changing their ratios. The changes of exclusive semileptonic branching fractions would only influence the contaminated channels keeping all the rest unchanged. The same is true for the uncertainty due to contamination by higher-multiplicity channels which will now be discussed in a more detailed way.

As described in section 5.2, the method of m_{cut}^2 cannot discriminate between channels with two or three neutral decay products. An inspection of D^0 decay modes shows that the most affected channels are probably $D^0 \rightarrow K^-\pi^+\pi^0\pi^0$ (contaminated by $D^0 \rightarrow K^-\pi^+\pi^0\pi^0\pi^0$) and $D^0 \rightarrow \bar{K}^0\pi^+\pi^-\pi^0$ (possibly containing $D^0 \rightarrow \bar{K}^0\pi^+\pi^-\pi^0\pi^0$). We can make a rough estimate of the strength of the above five-body channels since we measure the branching fractions for two similar channels, i.e. $D^0 \rightarrow K^-\pi^+\pi^+\pi^-\pi^0$ and $D^0 \rightarrow \bar{K}^0\pi^+\pi^+\pi^-\pi^-$, to be of the order of $\sim 2\%$. We do not know the resonant component (if any) of the five-body channels and, consequently, we cannot relate various charge states by a proper isospin Clebsch-Gordan coefficient. However, it seems reasonable to assume that all the branching fractions for channels involving a kaon and five pions will be also of the order of a few percent. In that case, the contamination will be nearly an order of magnitude lower than the branching fraction of the contaminated channel which is $\sim 15\%$.

We expect a still weaker contamination for the D^+ channels, since an additional π^0 in the Cabibbo-favoured channels would produce decays into a kaon and five pions which are certainly strongly suppressed by phase space and have not yet been firmly established. For the Cabibbo-suppressed channel there is a possible admixture of $D^+ \rightarrow \pi^+\pi^+\pi^-\pi^0\pi^0$ but this is small (cf. branching fraction for $D^+ \rightarrow \pi^+\pi^+\pi^+\pi^-\pi^-$), badly known and well within the error of the branching fraction for the former channel.

For the D_s^+ decay modes involving a single π^0 , we cannot determine the contamination by channels with two π^0 s because we are limited to low values of m_0^2 by the procedure described in section 5.2. A rough estimate can be obtained by the inspection of relevant five-prong channels. This would suggest the largest contamination of the $D_s^+ \rightarrow \phi\pi^+\pi^0$ channel to be that from $D_s^+ \rightarrow \phi\pi^+\pi^0\pi^0$ since the branching fraction for $D_s^+ \rightarrow \phi\pi^+\pi^+\pi^-$ is only 2.5 times smaller than that for $D_s^+ \rightarrow \phi\pi^+\pi^0$.

For the Λ_c^+ we expect little contamination of Cabibbo-favoured channels since this would need six-body modes which should be strongly suppressed by the phase space.

6. DISCUSSION OF RESULTS

6.1 *Comparison with Other Results*

Before making a detailed comparison of our branching fractions with those of other experiments, let us comment on our earlier study of D^0 [5]. This was based on a full sample of ~ 620 single-tag events and a subsample of 84 double-tag events, namely those in which a D^0 was associated with a fully reconstructed $\bar{D}^0$, D^- , D_s^- or $\bar{\Lambda}_c^-$. The results shown in Table 6 come from 681 D^0 s in the present double-tag sample. In addition, we now use four more

branching ratios (see section 4). The present results supersede those of ref.[5], while being entirely consistent with them.

Now let us compare our branching fractions for D^0 and D^+ with those listed in RPP [1]. An inspection of Tables 6 and 7 shows that our errors are comparable to those of ref.[1] and that the consistency is fair except for our branching fraction for the $D^0 \rightarrow \pi^+\pi^-\pi^0$ channel which is high. This can be at least partially explained by the admixture of decays with more π^0 s, but even then the value of $\sim 3\%$ seems to be too high for Cabibbo-suppressed channels.

Let us remark that the RPP [1] values are dominated by the branching fractions of the MARK III collaboration but also include the results of other experiments. This inclusion has reduced the branching fractions by (10-15)% with respect to the original results of MARK III. Our results confirm this reduction.

An additional check can be made by calculating some inclusive branching fractions from our results. In Table 10 they are compared to the values quoted in RPP. For $D^0 \rightarrow \bar{K}^0 (K^0) + \text{anything}$ one should increase our value by up to 10% to account for K^0 s from zero-prong decays. In the last row we have subtracted 0.8% due to the $D^+ \rightarrow K^+\bar{K}^0$ channel which is not accessible to us. In general there is a reasonable consistency.

For D_s^+ and Λ_c^+ no direct comparison of branching fractions is possible. However, we can consider some branching ratios which were not accessible in our single-tag sample, namely some D_s^+ (and D^+) decay channels involving a single π^0 . The branching ratios are calculated from Tables 7 and 8 (the uncertainty connected with the $\mathcal{L}_1$ cancels). In Table 11 they are compared with the results of the TPS [21] and NA14 [22] collaborations. Again the agreement is reasonable. Of particular interest is the ratio $D_s^+ \rightarrow \pi^+\pi^+\pi^-\pi^0 / D_s^+ \rightarrow \phi\pi^+$. Assuming that the former channel is completely dominated by $D_s^+ \rightarrow \eta'\pi^+$ mode, we obtain $D_s^+ \rightarrow \eta'\pi^+ / D_s^+ \rightarrow \phi\pi^+ < 5$ at 90% C.L. This value is to be compared with 5.0 ± 2.3 measured by MARK III [23] and $5.0 \pm 1.8 \pm 1.2$ reported by NA14 [22].

Considering also the fair agreement of our branching ratios with those of other experiments (see ref.[5] and section 2), we can conclude that there is an overall consistency between our results and the values reported by other groups. In some cases like $D^0 \rightarrow K^-\pi^+$ and $D^0 \rightarrow K^-\pi^+\pi^+\pi^-$, the errors of the branching fractions averaged from RPP and our results will be smaller than 10%.

6.2 Observation of New Decay Channels

The good consistency between our results and those of other experiments shows that the fairly complicated procedure described in section 5 leads to proper conclusions. This supports the significance of our branching fractions for the decay channels which were either unknown

or suggested, but without any knowledge of branching ratios (those labelled "seen" in RPP [1]). Before reviewing them, let us remark that the branching fractions measured for the first time in this work refer to many-body decays. The study of resonant substructures is difficult, especially for the double-tag events where we miss neutral decay products. Consequently most of our branching fractions cannot be compared to the predictions of Bauer, Stech and Wirbel [24] which refer only to two-body decays (see section 6.4).

We study eight new decay modes of the D^0 , jointly amounting to $\sim 7\%$. Two of them are Cabibbo-favoured decays into a kaon and four pions. The decay $D^0 \rightarrow \bar{K}^0 \pi^+ \pi^+ \pi^- \pi^-$ is suggested in the single-tag sample and is fairly significant in the double-tag events. The decay $D^0 \rightarrow K^- \pi^+ \pi^+ \pi^- \pi^0$ is clearly seen in both samples. Thus the D^0 decay into a kaon and four pions has been firmly established. The remaining new decay modes are Cabibbo-suppressed. The most significant is the mode $D^0 \rightarrow K^+ K^- \pi^+ \pi^-$ (with a sizable $D^0 \rightarrow \phi \pi^+ \pi^-$ component) clearly seen in the single-tag sample (cf. ref.[5]) and confirmed in the double-tag one. The mode $D^0 \rightarrow \pi^+ \pi^+ \pi^- \pi^- \pi^0$ is only indicated in the former (cf. section 4) but fairly significant in the latter. Our results on $D^0 \rightarrow K^+ K^- \bar{K}^0 \pi^0$ and $D^0 \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$ (a couple of events in the double-tag sample) can be treated as indications only. Finally, our $D^0 \rightarrow \pi^+ \pi^+ \pi^+ \pi^- \pi^- \pi^-$ can only be used to calculate an upper limit since the weak signal in the single-tag sample is not confirmed in the double-tag one.

A similar situation exists for the D^+ decays, for which the statistics amounts to only half of that for the D^0 . Here we observe a very significant decay into a kaon and four charged pions. This is a 4σ effect in the single-tag sample and a single event in the double-tag one. The remaining new significant decay modes are Cabibbo-suppressed ones and could be seen only in the latter, namely $D^+ \rightarrow \pi^+ \pi^+ \pi^- \pi^0$, $D^+ \rightarrow K^+ K^- \pi^+ \pi^0$ and $D^+ \rightarrow \phi \pi^+ \pi^0$. The combined (single + double tags) evidence for $D^+ \rightarrow \pi^+ \pi^+ \pi^+ \pi^- \pi^-$ is only a 2σ effect and clearly needs confirmation. The same is true for $D^+ \rightarrow K^- K^0 \pi^+ \pi^+$, which might be seen more clearly in the $K^{*+} \bar{K}^{*0}$ channel (cf. Fig. 2c). Finally there are indications for the decays $D^+ \rightarrow K^- \pi^+ \pi^+ \pi^+ \pi^- \pi^0$, $D^+ \rightarrow \bar{K}^0 \pi^+ \pi^+ \pi^+ \pi^- \pi^-$ and the Cabibbo-suppressed $D^+ \rightarrow \pi^+ \pi^+ \pi^+ \pi^- \pi^- \pi^0$. The sum of the branching fractions for the nine new channels is $\sim 8\%$.

Our statistics for the D_s^+ meson is an order of magnitude smaller than that for the D^0 . Nevertheless, we establish a new strong channel $D_s^+ \rightarrow K^- K^0 \pi^+ \pi^+$ clearly seen in single- and double-tag samples. The former shows that this channel is dominated by the $K^{*+} \bar{K}^{*0}$ (i.e. a vector-vector) mode. This fits with the observation of a non-significant signal in the $D_s^+ \rightarrow K^+ \bar{K}^0 \pi^+ \pi^-$ channel which cannot proceed via the $K^* \bar{K}^*$ intermediate state. Thus the decay of D_s^+ into a non-resonant four-body system of charged kaon, neutral kaon and two charged pions is fairly weak, while the mode $D_s^+ \rightarrow K^{*+} \bar{K}^{*0}$ belongs to the strongest channels. Another new and fairly strong channel is the four-body $D_s^+ \rightarrow (K^+ K^-)_{nr} \pi^+ \pi^0$ mode clearly seen in the

double-tag sample. In the same sample one could claim new decays into pions only, i.e. $D_s^+ \rightarrow \pi^+\pi^+\pi^-\pi^0$ and $D_s^+ \rightarrow \pi^+\pi^+\pi^+\pi^-\pi^0$. There are also some indications for $D_s^+ \rightarrow (K^+K^-)_{nr}\pi^+\pi^+\pi^-$ and $D_s^+ \rightarrow \pi^+\pi^+\pi^+\pi^-\pi^-$ seen only in the single-tag sample. It is to be noted that the D_s^+ decays into pions only are clearly less abundant than those involving two kaons. This confirms the observation of the TPS collaboration [11] that the annihilation diagram is much weaker in D_s^+ decays than the spectator one. This was seen for the $D_s^+ \rightarrow \pi^+\pi^+\pi^-$ vs $D_s^+ \rightarrow \phi\pi^+$ channel; we extend this to higher decay multiplicities. The sum of branching fractions for four-body channels involving two kaons is several times larger than that for four-body decays into pions only and the same is true for five-body decays. Finally, we find the first indication for the Cabibbo-suppressed channels $D_s^+ \rightarrow K^+\pi^+\pi^-$ and $D_s^+ \rightarrow K^+\pi^+\pi^-\pi^0$ (note that these channels are doubly Cabibbo-suppressed for the D^+). The sum of the branching fractions for all new channels amounts to a value seven times larger than that of the $D_s^+ \rightarrow \phi\pi^+$ channel.

Finally, we turn to the Λ_c^+ decays for which the double-tag statistics is similar to that of the D_s^+ but the single-tag events are twice as numerous. Here we confirm the $\Lambda_c^+ \rightarrow pK^-\pi^+\pi^0$ mode seen by the NA1 collaboration [25] and determine its frequency. We also have the first evidence for the five-prong decay $\Lambda_c^+ \rightarrow pK^-\pi^+\pi^+\pi^-$ seen both in single- and double-tag samples. The decay $\Lambda_c^+ \rightarrow p\bar{K}^0\pi^+\pi^-\pi^0$ could be seen only in double-tag events but it looks like a good candidate for another five-body channel. There is a somewhat weaker evidence for the Cabibbo-suppressed mode $\Lambda_c^+ \rightarrow p\pi^+\pi^-\pi^0$. Nearly all remaining new channels listed in Table 9 are Cabibbo-suppressed and appear either in the single- or in the double-tag sample but never in both. All of them are less-than- 2σ effects and can be treated as indications only. The sum of the branching fractions for all new channels amounts to a value twice as big as that of the $\Lambda_c^+ \rightarrow pK^-\pi^+$ channel.

6.3 Branching Fractions of D_s^+ and Λ_c^+

As described in section 5.4, the fraction $\mathcal{L}_1$ of one-prong D_s^+ decays is not known. *A priori* it seems unlikely that $\mathcal{L}_1$ is much larger than $\mathcal{L}_3$ since there are but a few Cabibbo-favoured decays leading to such inequality. We mean here $D_s^+ \rightarrow K^+\bar{K}^0$ measured by the CLEO [26] and MARK III [27] collaborations to be just as strong as the $D_s^+ \rightarrow \phi\pi^+$. The $D_s^+ \rightarrow \pi^+\pi^0$, $D_s^+ \rightarrow \pi^+\pi^0\pi^0$ etc. should be fairly weak as they can only proceed via the annihilation diagram. Finally there is the $D_s^+ \rightarrow \eta\pi^+$ channel. The TPS group [11] has measured its branching ratio to be 0.6 ± 0.6 with respect to the $D_s^+ \rightarrow \phi\pi^+$, while the MARK II collaboration [23] has obtained 3.0 ± 1.3 for the ratio. Thus probably $D_s^+ \rightarrow \eta\pi^+$ and $D_s^+ \rightarrow \phi\pi^+$ are of similar strength. On the other hand, the strong channels $D_s^+ \rightarrow K^+\bar{K}^{*0}$, $D_s^+ \rightarrow K^{*+}\bar{K}^{*0}$ and $D_s^+ \rightarrow \eta'\pi^+$ contribute to $\mathcal{L}_1 < \mathcal{L}_3$ with weights of ~ 1.0 , 2.5 and ~ 4.0 , respectively (always in units of $D_s^+ \rightarrow \phi\pi^+$). The $D_s^+ \rightarrow \phi\pi^+$ itself and most of the remaining channels contribute to $\mathcal{L}_1 \equiv \mathcal{L}_3$. Thus an educated guess yields $\mathcal{L}_1$ smaller than $\mathcal{L}_3$ ($\mathcal{L}_1 \equiv 0.7 \mathcal{L}_3$). We tentatively assume $\mathcal{L}_1 \equiv$

$\mathcal{L}_3$ as for D^+ and recalculate the branching fractions (see last column in Table 8). This time we quote full branching fractions for channels involving ϕ or K^* (the values in the previous column referred to $\phi \rightarrow K^+K^-$ and $\bar{K}^{*0} \rightarrow K^-\pi^+$ only); the channel $D_s^+ \rightarrow K^-K^0\pi^+\pi^+$ is split into $K^*\bar{K}^{*0}$ and a non-resonant component. It is immediately seen that our branching fraction for the reference channel $D_s^+ \rightarrow \phi\pi^+$ is at the upper limit of 4.1% (90% C.L.) measured by the MARK II collaboration [12]. However, the reduction of the branching fraction to the level of $\sim 3\%$ would require $\mathcal{L}_1 \cong 2 \mathcal{L}_3$, which disagrees with our educated guess. Therefore either there are some strong D_s^+ decay channels, mainly in the one-prong domain, to be discovered or the branching fractions shown in the last column of Table 8 are not very far from reality.

The semileptonic branching fractions of D_s^+ amounting probably to $\sim 15\%$ are contaminating the channels with neutral decay products. As described above, we cannot discriminate between $D_s^+ \rightarrow K^+K^-\pi^+\pi^0$ and $D_s^+ \rightarrow K^+K^-e^+\nu_e$ or between $D_s^+ \rightarrow \pi^+\pi^+\pi^-\pi^0$ and $D_s^+ \rightarrow \pi^+\mu^+\pi^-\nu_\mu$. Considering also the one-prong decays there is ample space to accommodate the semileptonic decays. It should be remembered that they do not influence the branching fractions for fully reconstructed decays.

There is even less information for Λ_c^+ . The branching fraction for the best known channel $\Lambda_c^+ \rightarrow pK^-\pi^+$ is still controversial. Assuming the estimate of 4% from the extensive review of Klein [15] (cf. also ref.[28]), we calculate the branching fractions shown in the last column of Table 9. In such a case all three- and five-prong decay modes involving a proton amount to 20%. Adding about half of this value for one-prong decays like $\Lambda_c^+ \rightarrow p\bar{K}^0$, $\Lambda_c^+ \rightarrow p\bar{K}^0\pi^0$ etc., we obtain $\sim 30\%$ for all decays with a proton in the final state. This agrees with the statistical calculation of Bjorken as well as with the Lund model (both quoted in ref.[15]). In any case there are still many Λ_c^+ decay channels to be discovered and measured, mainly those involving a neutron or a Σ^+ . The latter need a special analysis of our data and will be the subject of a separate publication (see ref.[29] for preliminary results).

6.4 Comparison with Phenomenological Predictions

As has been already mentioned, this work finds new decay channels only in the field of many-body decays. On the other hand, most phenomenological studies deal exclusively with two-body decays. This restricts the field of comparison to only a few channels, namely the decays into a vector-vector final state where we obtain results which are either new or more accurate than those of other experiments. Unfortunately, we cannot identify both vector mesons in channels involving a π^0 . In such a case we calculate only an upper limit assuming that the π^0 always comes from the decay of a vector meson; e.g. we treat the branching ratio for the decay $D_s^+ \rightarrow \phi\pi^+\pi^0$ as an upper limit of the branching ratio for $D_s^+ \rightarrow \phi\rho^+$. In Table 12 we compare our results with the predictions of a factorization model of Bauer, Stech and Wirbel [24]. It should be remembered that neither the QCD sum rules used by Blok and

Shifman [30] nor the final-state-interaction approach of Kamal and Verma [31] yield predictions for charmed mesons decaying into a vector-vector final state. It is seen that most results are consistent with the predictions; there are however two channels for which there is a significant discrepancy. The discrepancy for the $D_s^+ \rightarrow \phi \rho^+$ channel has been already reported by the TPS collaboration [21]. We show that also for the mode $D^+ \rightarrow K^{*+} \bar{K}^{*0}$ the experimental result is significantly below the prediction of Ref.[24].

7. CONCLUSIONS

High-precision silicon detectors and a purely topological approach allowed an investigation of a wide range of charmed decays, in particular an extensive study of events with two secondary vertices. This has yielded branching ratios and branching fractions for numerous decay modes of D^0 , D^+ , D_s^+ and Λ_c^+ .

The results obtained in this work are generally consistent with those of other experiments.

We determine branching fractions for twenty decay channels of the D^0 . Among them we demonstrate new channels $D^0 \rightarrow \pi^+ \pi^+ \pi^- \pi^- \pi^0$, $D^0 \rightarrow \bar{K}^0 \pi^+ \pi^+ \pi^- \pi^-$, $D^0 \rightarrow K^- \pi^+ \pi^+ \pi^- \pi^0$, $D^0 \rightarrow (K^+ K^-)_{nr} \pi^+ \pi^-$ and $D^0 \rightarrow \phi \pi^+ \pi^-$.

Branching fractions are measured for eighteen D^+ decay channels including new decay modes $D^+ \rightarrow K^- \pi^+ \pi^+ \pi^+ \pi^-$, $D^+ \rightarrow \pi^+ \pi^+ \pi^- \pi^0$, $D^+ \rightarrow (K^+ K^-)_{nr} \pi^+ \pi^0$ and $D^+ \rightarrow \phi \pi^+ \pi^0$.

Sixteen decay modes of D_s^+ are studied; among them the new ones $D_s^+ \rightarrow K^- K^0 \pi^+ \pi^+$ (dominated by $D_s^+ \rightarrow K^{*+} \bar{K}^{*0}$) and $D_s^+ \rightarrow (K^+ K^-)_{nr} \pi^+ \pi^0$. There are also first indications of Cabibbo-suppressed decays into a single kaon and pions. We attempt an estimation of branching fractions by channel counting.

Among eleven decay channels of Λ_c^+ we observe for the first time the five-body decays $\Lambda_c^+ \rightarrow p K^- \pi^+ \pi^+ \pi^-$ and $\Lambda_c^+ \rightarrow p \bar{K}^0 \pi^+ \pi^- \pi^0$. There are numerous indications of Cabibbo-suppressed decay modes involving only pions or two kaons in addition to a proton.

ACKNOWLEDGEMENTS

The authors are very grateful to Mrs. Carole Ponting for her careful and patient typing of the compuscript. Two of us (K.R. and M.W.) would like to thank the Max-Planck-Institute for support.

REFERENCES

- [1] Particle Data Group, Review of Particle Properties, Phys. Lett. 204B (1988).
- [2] R.M. Baltrusaitis et al., Phys. Rev. Lett. 54 (1985) 1976;
R.M. Baltrusaitis et al., ibid. 55 (1985) 150;
R.M. Baltrusaitis et al., ibid. 56 (1986) 2136;
R.M. Baltrusaitis et al., ibid. 56 (1986) 2140;
J. Adler et al., Phys. Lett. 196B (1987) 107.
- [3] M. Aguilar et al., Z. Phys. C36 (1987) 559.
- [4] S. Barlag et al., contribution # 770 to the Int. Conf. on High Energy Physics, Munich, 1988; also CERN-EP/88-103.
- [5] S. Barlag et al., Phys. Lett. 232B (1989) 561.
- [6] C. Daum et al., Nucl. Instrum. Methods 217 (1983) 361.
- [7] S. Barlag et al., Phys. Lett. 218B (1989) 374.
- [8] S. Barlag et al., Phys. Lett. 233B (1989) 522.
- [9] S. Barlag et al., Phys. Lett. 236B (1990) 495.
- [10] S. Barlag et al., "*Production of the Charmed Λ_c^+ Baryon in π^- -Cu Interactions at 230 GeV/c*", submitted for publication to Phys. Lett. B;
S. Barlag et al., "*Production Properties of D^0 , D^+ , D^{*+} and D_s^+ in 230 GeV/c π^- and K^- -Cu Interactions*", submitted for publication to Z. Phys. C.
- [11] J.C. Anjos et al., Phys. Rev. Lett. 62 (1989) 125.
- [12] S.R. Wasserbaech, Ph.D. thesis, SLAC-345, 1989;
J. Adler et al., Phys. Rev. Lett. 64 (1990) 169.
- [13] W.Y. Chen et al., Phys. Lett. 226B (1989) 192.
- [14] J.C. Anjos et al., Phys. Rev. Lett. 60 (1988) 897.
- [15] S.R. Klein, Boston University preprint, BU-HEP-89-16, August 1989 (the author quotes preliminary results from the ARGUS and CLEO collaborations).
- [16] H. Albrecht et al., Phys. Lett. 207B (1988) 109.
- [17] J.C. Anjos et al., Fermilab-Pub-89/44E; submitted for publication to Phys. Rev. Lett.
- [18] D. Hitlin, Nucl. Phys. B (Proc. Suppl.) 3 (1988) 179.
- [19] J. Adler et al., Phys. Rev. Lett. 62 (1989) 1821.
- [20] J.C. Anjos et al., Phys. Rev. Lett. 62 (1989) 1587.
- [21] J.C. Anjos et al., Phys. Lett. 223B (1989) 267.
- [22] G. Wormser, Heavy Quark Symposium, Cornell University; also preprint LAL 89-27.

- [23] G. Wormser et al., Phys. Rev. Lett. 61 (1989) 1057.
- [24] M. Bauer, B. Stech and M. Wirbel, Zeit. Phys. C34 (1987) 103.
- [25] S.R. Amendolia et al., Zeit. Phys. C36 (1987) 513.
- [26] W.Y. Chen et al., Phys. Lett. 226B (1989) 192.
- [27] J. Adler et al., Phys. Rev. Lett. 63 (1989) 1211.
- [28] H. Albrecht et al., Phys. Lett. 210B (1988) 263.
- [29] P. Singer and B. Gad Eilan, eds., Proc. XII Int. Workshop on Weak Interactions and Neutrinos (Ginosar, Israel, April 1989), Nucl. Phys. B (Proc. Suppl.) 13A, to be published.
- [30] B. Yu Blok and M.A. Shifman, Yad. Fiz. 45 (1987) 211;
B. Yu Blok and M.A. Shifman, *ibid.*, 45 (1987) 478;
B. Yu Blok and M.A. Shifman, *ibid.*, 45 (1987) 841;
B. Yu Blok and M.A. Shifman, *ibid.*, 46 (1987) 1310;
{Sov. J. Nucl. Phys. 45 (1987) 135, 45 (1987) 301, 45 (1987) 522 and 46 (1987) 767}.
- [31] A.N. Kamal and R.C. Verma, Phys. Rev. D35 (1987) 3515;
A.N. Kamal and R.C. Verma, *ibid.* D36 (1987) 3527;
A.N. Kamal et al., Z. Phys. C41 (1988) 207.

Table 1

Branching ratios of the D^+ meson

Decay mode	# single tags	Acc (%)	Branching Ratio	Other results
$\pi^+\pi^+\pi^-$	$2.9^{+3.0}_{-2.6}$	7.01	$0.017^{+0.018}_{-0.015}$	0.042 ± 0.019 [1] $0.035\pm 0.007\pm 0.003$ [11]
$\bar{K}^0\pi^+\pi^+\pi^-$	15.4 ± 5.9	1.12	$0.549\pm 0.213\pm 0.047$	
$K^-K^0\pi^+\pi^+$	$4.6^{+5.1}_{-3.1}$	1.54	$0.119^{+0.132}_{-0.080}\pm 0.010^*$	
$K^-\pi^+\pi^+$	198.4 ± 11.0	7.97	1.000 (fixed)	
$\phi\pi^+$	11.8 ± 4.1	11.62	0.083 ± 0.028	0.073 ± 0.010 [1]
$K+\bar{K}^*0$	10.4 ± 4.3	8.98	0.070 ± 0.030	0.056 ± 0.010 [1]
$(K^+K^-\pi^+)_{nr}$	15.0 ± 4.8	9.00	0.067 ± 0.022	0.050 ± 0.009 [1]
$\pi^+\pi^+\pi^+\pi^-\pi^-$	2.4 ± 2.0	7.13	0.014 ± 0.012	< 0.019 [11]
$\bar{K}^0\pi^+\pi^+\pi^+\pi^-\pi^-$	1.0 ± 1.4	1.32	$0.009\pm 0.013\pm 0.001^*$	
$K^-\pi^+\pi^+\pi^+\pi^-$	15.5 ± 3.8	8.59	$0.072\pm 0.018^*$	

Table 2

Branching ratios of the D_s^+ meson

Decay mode	# single tags	Acc (%)	Branching Ratio	Other results
$\pi^+\pi^+\pi^-$	3.2 ± 2.9	4.76	0.13 ± 0.12	$0.44\pm 0.10\pm 0.04$ [11]
$K^+\pi^+\pi^-$	1.4 ± 1.8	4.82	$0.05\pm 0.07^*$	
$K^+\bar{K}^0\pi^+\pi^-$	2.2 ± 2.4	0.75	$0.56\pm 0.62^*$	
$(K^-K^0\pi^+\pi^+)_{nr}$	1.5 ± 1.6	0.89	$0.32\pm 0.35\pm 0.06^*$	
$K^*+\bar{K}^*0$	7.4 ± 3.0	1.05	$2.92\pm 1.37\pm 0.26^*$	
$\phi\pi^+$	23.1 ± 5.4	8.92	1.00 (fixed)	
$K^+\bar{K}^*0$	19.5 ± 5.6	5.96	0.94 ± 0.35	0.94 ± 0.19 [1] $0.84\pm 0.30\pm 0.22$ [12] $1.05\pm 0.17\pm 0.12$ [13]
$(K^+K^-\pi^+)_{nr}$	17.8 ± 5.5	6.65	0.51 ± 0.20	$0.25\pm 0.07\pm 0.05$ [14]
$\pi^+\pi^+\pi^+\pi^-\pi^-$	1.7 ± 1.6	4.56	$0.07\pm 0.07^*$	< 0.29 [11]
$\phi\pi^+\pi^+\pi^-$	7.7 ± 3.2	8.88	0.34 ± 0.16	0.48 ± 0.20 [1]
$(K^+K^-)_{nr}\pi^+\pi^+\pi^-$	2.9 ± 1.9	7.62	$0.07\pm 0.05^*$	< 0.32 [14]

Table 3

Branching ratios of the charmed baryon Λ_c^+

Decay mode	# single tags	Acc (%)	Branching Ratio	Other results
$p\pi^+\pi^-$	5.3 ± 2.7	4.44	$0.069\pm 0.036^*$	
$(p f_{975}^0$	4.3 ± 2.9	4.52	$0.055\pm 0.036^*)$	
$p\bar{K}^0\pi^+\pi^-$	12.4 ± 4.5	0.74	$0.98\pm 0.36\pm 0.08$	1.00 ± 0.42 [15] 0.51 ± 0.15 [15]
$pK^-\pi^+$	118.7 ± 8.9	6.45	1.00 (fixed)	
$(p\bar{K}^{*0}$	33.7 ± 10.0	8.47	$0.35\pm 0.11)$	0.22 ± 0.09 [1]
pK^+K^-	3.6 ± 2.0	5.90	$0.048\pm 0.027^*$	
$(p\phi$	2.8 ± 1.9	4.09	$0.040\pm 0.027^*)$	
$\Lambda^0\pi^+\pi^+\pi^-$	10.6 ± 4.5	0.65	$0.94\pm 0.41\pm 0.13$	$0.61\pm 0.16\pm 0.14$ [16] $0.82\pm 0.29\pm 0.27$ [17] 0.75 ± 0.14 [15]
$p\pi^+\pi^+\pi^-\pi^-$	3.0 ± 1.9	4.95	$0.036\pm 0.023^*$	
$pK^-\pi^+\pi^+\pi^-$	2.9 ± 1.9	7.80	$0.022\pm 0.015^*$	

Table 4

Some branching ratios of the D^0

Decay mode	Δm range (MeV)	Acc (%)	# events	Branching Ratio
$\pi^+\pi^-\pi^0$	141 - 151	0.68	5.2 ± 3.4	0.453 ± 0.300
$K^-\pi^+\pi^0$	141 - 153	1.49	25.6 ± 7.3	1.026 ± 0.313
$\pi^+\pi^+\pi^-\pi^-\pi^0$	141 - 153	3.06	5.7 ± 4.0	0.111 ± 0.079
$K^-\pi^+\pi^+\pi^-$	141 - 149	5.87	101.6 ± 11.1	1.000 (fixed)
$K^-\pi^+\pi^+\pi^-\pi^0$	141 - 153	6.15	27.9 ± 7.1	0.271 ± 0.075

Table 5

Separation of decay channels (see section 5.2 for definition of symbols)

Channel A	Channel B	m_{cut}^2 (GeV ²)	Confusion level s_i	t_i from single- tag events
$D^0 \rightarrow \bar{K}^0 \pi^+ \pi^-$	$D^0 \rightarrow \bar{K}^0 \pi^+ \pi^- \pi^0$	0.65	17%	1/17
$D^0 \rightarrow K^- \pi^+ \pi^0$	$D^0 \rightarrow K^- \pi^+ \pi^0 \pi^0$	0.25	21%	-
$D^0 \rightarrow K^+ K^- \bar{K}^0$	$D^0 \rightarrow K^+ K^- \bar{K}^0 \pi^0$	0.42	6%	0/17
$D^+ \rightarrow \bar{K}^0 \pi^+ \pi^+ \pi^-$	$D^+ \rightarrow \bar{K}^0 \pi^+ \pi^+ \pi^- \pi^0$	0.55	16%	7/25
$D_s^+ \rightarrow \pi^+ \pi^+ \pi^- \pi^0$	$D^+ \rightarrow \pi^+ \pi^+ \pi^- \pi^0$	0.08	35%	-
$D_s^+ \rightarrow K^+ K^- \pi^+ \pi^0$	$D^+ \rightarrow K^+ K^- \pi^+ \pi^0$	0.08	26%	-
$D_s^+ \rightarrow \pi^+ \pi^+ \pi^+ \pi^- \pi^- \pi^0$	$D^+ \rightarrow \pi^+ \pi^+ \pi^+ \pi^- \pi^- \pi^0$	0.07	25%	-
$\Lambda_c^+ \rightarrow p \bar{K}^0 \pi^+ \pi^-$	$\Lambda_c^+ \rightarrow p \bar{K}^0 \pi^+ \pi^- \pi^0$	0.44	14%	4/23
$\Lambda_c^+ \rightarrow p K^- \pi^+ \pi^0$	$\Lambda_c^+ \rightarrow p K^- \pi^+ \pi^0 \pi^0$	0.10	8%	-

Table 6
Branching fractions of D^0 meson

Decay mode	Branching ratio [5]	# double tag events	Correction factor f_i	Branching fraction (%) from this work	Branching fraction from RPP [1]
$\pi^+\pi^-$	0.009 ± 0.007	2	-	0.09 ± 0.06	0.13 ± 0.04
$\pi^+\pi^-\pi^0$	0.453 ± 0.300	15	1.07 ± 0.26	4.08 ± 1.11	1.1 ± 0.4
$\bar{K}^0\pi^+\pi^-$	0.444 ± 0.142	88	0.77 ± 0.10	$4.11\pm_{-1.05}^{+1.09}$	$5.6\pm_{-0.6}^{+0.7}$
$\bar{K}^0\pi^+\pi^-\pi^0$	-	120	0.72 ± 0.09	$14.0\pm3.4\pm2.1$	$10.6\pm_{-2.9}^{+7.3}\pm0.6$
$K^-\pi^+$	0.472 ± 0.039	26	0.92 ± 0.20	$3.75\pm_{-0.41}^{+0.44}$	$3.77\pm_{-0.32}^{+0.37}$
$K^-\pi^+\pi^0$	1.026 ± 0.313	109	0.86 ± 0.09	$8.67\pm_{-1.96}^{+2.00}$	$12.5\pm_{-1.3}^{+1.5}$
$K^-\pi^+\pi^0\pi^0$	-	173	0.77 ± 0.07	$16.1\pm2.5\pm1.7$	15 ± 5
K^+K^-	0.057 ± 0.016	4	-	0.47 ± 0.13	0.45 ± 0.08
$\phi\bar{K}^0$	0.170 ± 0.095	5	$0.80\pm_{-0.42}^{+0.48}$	1.75 ± 0.58	$0.83\pm_{-0.16}^{+0.18}$
$(K^+K^-)_{nr}\bar{K}^0$	0.194 ± 0.085	6	$0.33\pm_{-0.24}^{+0.39}$	$0.60\pm_{-0.36}^{+0.37}$	$0.74\pm_{-0.18}^{+0.23}$
$K^+K^-\bar{K}^0\pi^0$	-	3	-	$0.92\pm_{-0.44}^{+0.64*}$	-
$\pi^+\pi^+\pi^-\pi^-$	0.028 ± 0.012	3	-	0.25 ± 0.09	$1.01\pm_{-0.26}^{+0.71}$
$\pi^+\pi^+\pi^-\pi^-\pi^0$	0.111 ± 0.079	15	1.07 ± 0.26	$1.72\pm_{-0.46}^{+0.47*}$	-
$\bar{K}^0\pi^+\pi^+\pi^-\pi^-$	0.092 ± 0.055	25	0.80 ± 0.20	$1.02\pm_{-0.39}^{+0.40*}$	-
$K^-\pi^+\pi^+\pi^-$	1.000 (fixed)	42	1.10 ± 0.19	$7.72\pm_{-0.65}^{+0.68}$	$7.9\pm_{-0.9}^{+1.0}$
$K^-\pi^+\pi^+\pi^-\pi^0$	0.271 ± 0.075	40	0.80 ± 0.15	$2.56\pm_{-0.55}^{+0.57}$	seen
$\phi\pi^+\pi^-$	0.036 ± 0.010	0	-	$0.24\pm0.08^*$	-
$(K^+K^-)_{nr}\pi^+\pi^-$	0.019 ± 0.007	3	-	$0.16\pm0.06^*$	-
$K^+K^-\pi^+\pi^-\pi^0$	-	2	-	$0.28\pm_{-0.15}^{+0.25*}$	-
$\pi^+\pi^+\pi^+\pi^-\pi^-\pi^-$	0.005 ± 0.004	0	-	$0.04\pm0.03^*$	-

Table 7
Branching fractions of D^+ meson

Decay mode	# double tag events	Correction factor f_i	Branching fraction (%) from this work	Branching fraction from RPP [1]
$\pi^+\pi^+\pi^-$	2	-	$0.24^{+0.14}_{-0.13}$	$0.33^{+0.15}_{-0.14}$
$\pi^+\pi^+\pi^-\pi^0$	13	$0.31^{+0.24}_{-0.17}$	$2.3^{+2.0*}_{-1.3}$	-
$\bar{K}^0\pi^+\pi^+\pi^-$	79	0.84 ± 0.11	$6.1^{+2.3}_{-2.2}$	$7.0^{+2.3}_{-2.1}$
$\bar{K}^0\pi^+\pi^+\pi^-\pi^0$	115	0.68 ± 0.09	$11.3\pm 2.4\pm 2.8$	$4.4^{+5.2}_{-1.5}$
$K^-\pi^+\pi^+$	33	$1.03^{+0.17}_{-0.18}$	8.0 ± 1.4	$7.8^{+1.1}_{-0.8}$
$K^-\pi^+\pi^+\pi^0$ $K^-\pi^+\pi^+\pi^0\pi^0$ $K^-\bar{K}^0\pi^+\pi^+$	} 95	0.73 ± 0.09	$3.2\pm 2.4\pm 3.6$ < 1.5 $0.9^{+1.1*}_{-0.8}$	$3.7^{+1.5}_{-0.8}$ $2.2^{+5.0}_{-0.9}$ -
$\phi\pi^+$	0	-	$0.58^{+0.28}_{-0.26}$	$0.57^{+0.11}_{-0.09}$
$K+\bar{K}^*0$	3	-	$0.64^{+0.26}_{-0.23}$	$0.44^{+0.10}_{-0.08}$
$(K^+K^-\pi^+)_{nr}$	1	-	0.39 ± 0.16	$0.39^{+0.10}_{-0.08}$
$\phi\pi^+\pi^0$	6	$1.33^{+0.36}_{-0.43}$	$2.38^{+1.11*}_{-0.85}$	-
$(K^+K^-)_{nr}\pi^+\pi^0$	7	$1.43^{+0.31}_{-0.39}$	$1.52^{+0.65*}_{-0.51}$	-
$\pi^+\pi^+\pi^+\pi^-\pi^-$	1	-	$0.15\pm 0.11^*$	-
$\pi^+\pi^+\pi^+\pi^-\pi^-\pi^0$	1	-	$0.28^{+0.38}_{-0.20}$	seen
$\bar{K}^0\pi^+\pi^+\pi^+\pi^-\pi^-$	3	-	$0.10\pm 0.10^*$	-
$K^-\pi^+\pi^+\pi^+\pi^-$	1	-	$0.41^{+0.15*}_{-0.14}$	-
$K^-\pi^+\pi^+\pi^+\pi^-\pi^0$	1	-	$0.19^{+0.26*}_{-0.13}$	-

Table 8
Branching fractions of D_s^+ meson

Decay mode	# double tag events	Correction factor f_i	Branching fraction ¹ (%)	Normalized ² branching fraction
$\pi^+\pi^+\pi^-$	4	-	$2.6 \pm_{1.6}^{2.0}$	$1.3 \pm_{0.8}^{1.0}$
$\pi^+\pi^+\pi^-\pi^0$	14	$0.43 \pm_{0.19}^{0.25}$	$18.7 \pm_{8.9}^{13.0}$	$9.6 \pm_{4.5}^{6.6}$
$K^+\pi^+\pi^-$	1	-	$1.0 \pm_{0.8}^{0.9}$	$0.5 \pm_{0.4}^{0.5*}$
$K^+\pi^+\pi^-\pi^0$	3	-	$11.2 \pm_{5.3}^{7.8}$	$5.7 \pm_{2.7}^{4.0*}$
$K^+\bar{K}^0\pi^+\pi^-$	8	$1.25 \pm_{0.37}^{0.33}$	$8.5 \pm_{2.8}^{3.2}$	$4.4 \pm_{1.4}^{1.7*}$
$(K^-\bar{K}^0\pi^+\pi^+)_{nr}$ $K^{*+}\bar{K}^{*0}$	} 14		$15.6 \pm_{8.6}^{9.5}$ {	$1.6 \pm_{1.6}^{2.2*}$
				$14.3 \pm_{7.8}^{8.7*}$
$\phi\pi^+$	2	-	$5.8 \pm_{1.5}^{2.0}$	$6.0 \pm_{1.6}^{2.1}$
$K^+\bar{K}^{*0}$	2	-	$6.0 \pm_{2.3}^{2.6}$	$4.6 \pm_{1.7}^{2.0}$
$(K^+K^-\pi^+)_{nr}$	6	$1.33 \pm_{0.43}^{0.36}$	$7.3 \pm_{2.6}^{3.2}$	$3.7 \pm_{1.5}^{1.6}$
$\phi\pi^+\pi^0$	2	-	$4.2 \pm_{2.3}^{3.7}$	$4.3 \pm_{2.4}^{3.8*}$
$(K^+K^-)_{nr}\pi^+\pi^0$	9	$0.89 \pm_{0.33}^{0.35}$	$14.9 \pm_{4.4}^{5.5}$	$7.6 \pm_{2.3}^{2.8*}$
$\pi^+\pi^+\pi^+\pi^-\pi^-$	0	-	$0.7 \pm_{0.7}^{0.9}$	$0.4 \pm_{0.4}^{0.5*}$
$\pi^+\pi^+\pi^+\pi^-\pi^-\pi^0$	3	-	$6.4 \pm_{3.0}^{4.4}$	$3.3 \pm_{1.6}^{2.3*}$
$\phi\pi^+\pi^+\pi^-$	0	-	$1.7 \pm_{0.9}^{1.1}$	$1.8 \pm_{1.0}^{1.2}$
$(K^+K^-)_{nr}\pi^+\pi^+\pi^-$	0	-	0.8 ± 0.6	$0.4 \pm 0.3^*$

-
- 1) The branching fractions are normalized to a sum of unity within our sample of three- and five-prong events as explained in section 5.4.
- 2) The branching fractions in this column are normalized using the educated guess described in section 6.3.

Table 9
Branching fractions of Λ_c^+ baryon

Decay mode	# double tag events	Correction factor f_i	Branching fraction ¹ (%)	Normalized branching fraction(see section 6.3)
$p\pi^+\pi^-$	0	-	$1.4 \pm_{1.0}^{1.1}$	$0.26 \pm_{0.19}^{0.21*}$
$\{p f_{975}^0\}$	0	-	$1.1 \pm_{1.0}^{1.1}$	$0.21 \pm_{0.19}^{0.21*}$
$p\pi^+\pi^-\pi^0$	6	$0.67 \pm_{0.36}^{0.43}$	$14.3 \pm_{5.1}^{6.7}$	$2.7 \pm_{1.0}^{1.3*}$
$p\bar{K}^0\pi^+\pi^-$	25	0.88 ± 0.20	$29.6 \pm_{8.1}^{8.7}$	$5.6 \pm_{1.5}^{1.6}$
$p\bar{K}^0\pi^+\pi^-\pi^0$	6	1.00 ± 0.41	$12.1 \pm_{4.3}^{5.6}$	$2.3 \pm_{0.8}^{1.1*}$
$pK^-\pi^+$	9	$0.67 \pm_{0.29}^{0.35}$	$21.0 \pm_{5.5}^{5.7}$	4.0 (fixed)
$pK^-\pi^+\pi^0$	9	$1.11 \pm_{0.35}^{0.33}$	$15.0 \pm_{4.5}^{5.6}$	$2.9 \pm_{0.9}^{1.1}$
$pK^-\pi^+\pi^0\pi^0$	2	-	$4.9 \pm_{2.7}^{4.3}$	$0.93 \pm_{0.51}^{0.82*}$
pK^+K^-	0	-	$1.0 \pm_{0.8}^{0.9}$	$0.19 \pm_{0.15}^{0.16*}$
$\{p\phi\}$	0	-	$0.8 \pm_{0.6}^{0.7}$	$0.16 \pm_{0.12}^{0.13*}$
$\Lambda^0\pi^+\pi^+\pi^-$	0	-	$2.4 \pm_{2.4}^{7.3}$	$0.5 \pm_{0.5}^{1.4*}$
$p\pi^+\pi^+\pi^-\pi^-$	0	-	0.7 ± 0.5	$0.14 \pm 0.09^*$
$pK^-\pi^+\pi^+\pi^-$	1	-	0.6 ± 0.3	$0.11 \pm 0.06^*$

1) The branching fractions are normalized to a sum of unity within our sample of decays.

Table 10

Some inclusive branching fractions of D^0 and D^+ mesons

Decay mode	Branching fraction (%) this work	Branching fraction from RPP[1]
$D^0 \rightarrow K^- + \text{anything}$	42.2 ± 3.8	43 ± 5
$D^0 \rightarrow K^+ + \text{anything}$	$3.4^{+0.8}_{-0.7}$	$6.4^{+2.6}_{-1.7}$
$D^0 \rightarrow \bar{K}^0(K^0) + \text{anything}$	$21.6 \pm 4.2 (+10.0)$	33 ± 10
$D^+ \rightarrow K^- + \text{anything}$	$16.5^{+4.8}_{-3.7}$	16.2 ± 3.5
$D^+ \rightarrow K^+ + \text{anything}$	$3.8^{+0.9}_{-0.7}$	$5.8^{+2.9}_{-2.8}$

Table 11

Branching ratios of D^+ and D_s^+ for channels involving a π^0

Decay channel	D^+		D_s^+	
	This work	Other results	This work	Other results
$\pi^+\pi^+\pi^-\pi^0$	$0.29^{+0.26}_{-0.16}$	0.2 ± 0.1 [19]	$1.6^{+1.2}_{-0.9}$	$0.7 \pm 1.7 \pm 0.2$ [19]
$K^-\pi^+\pi^+$	1.00 (fixed)		-	-
$\phi\pi^+$	-	-	1.00 (fixed)	
$\phi\pi^+\pi^0$	$0.30^{+0.15}_{-0.12}$	< 0.28 [19]	$0.7^{+0.7}_{-0.5}$	$2.4 \pm 1.0 \pm 0.5$ [19] < 2.5 [24]
$(K^+K^-)_{nr}\pi^+\pi^0$	$0.19^{+0.09}_{-0.07}$	< 0.25 [19]	1.3 ± 0.6	< 2.4 [19]

Table 12

Branching ratios for vector-vector charmed meson decays;
comparison with phenomenological predictions

Vector-vector decay	Reference channel	This work	Prediction [24]
$D^0 \rightarrow K^{*0}\bar{K}^{*0}$	$D^0 \rightarrow K^-\pi^+$	0.03 ± 0.02	-
$D^0 \rightarrow \phi \rho^0$	$D^0 \rightarrow K^-\pi^+$	$0.009^{+0.006}_{-0.004}$	0.015
$D^+ \rightarrow K^{*+}\bar{K}^{*0}$	$D^+ \rightarrow \phi\pi^+$	1.7 ± 1.6	8.5
$D_s^+ \rightarrow \phi \rho^+$	$D_s^+ \rightarrow \phi \pi^+$	< 1.8	6.3
$D_s^+ \rightarrow K^{*+}\bar{K}^{*0}$	$D_s^+ \rightarrow \phi \pi^+$	2.4 ± 1.6	1.2
$D_s^+ \rightarrow K^{*0}\rho^+$	$D_s^+ \rightarrow \phi \pi^+$	< 1.3	0.4

Figure Captions

- Fig. 1 : Schematic drawing of the ACCMOR spectrometer with the details of the vertex detector in the inset. B1-B7 : silicon microstrip detectors forming a beam telescope; T : 2.5 mm Cu target; CCDs : charge-coupled devices; I : interaction counter; MSD1-MSD8 : silicon microstrip detectors used together with CCDs as a telescope for secondary particles and vertices (underneath are shown strip orientations with respect to the horizontal plane).
- Fig. 2 : Effective mass plots for the D^+ decay channels (in some of them D_s^+ is also seen) : a) $\pi^+\pi^+\pi^-$, b) $\bar{K}^0\pi^+\pi^+\pi^-$, c) $K^-K^0\pi^+\pi^+$ ($K^{*+}\bar{K}^{*0}$ events are cross-hatched), d) $K^-\pi^+\pi^+$, e) $K^+K^-\pi^+$ ($\phi\pi^+$ events are cross-hatched), f) $\pi^+\pi^+\pi^+\pi^-\pi^-$, g) $\bar{K}^0\pi^+\pi^+\pi^+\pi^-\pi^-$, h) $K^-\pi^+\pi^+\pi^+\pi^-$.
- Fig. 3 : Effective mass plots for some D_s^+ decay channels : a) $K^+\pi^+\pi^-$, b) $K^+\bar{K}^0\pi^+\pi^-$, c) $K^+K^-\pi^+\pi^+\pi^-$ ($\phi\pi^+\pi^+\pi^-$ events are cross-hatched).
- Fig. 4 : Effective mass plots for Λ_c^+ decay channels : a) $p\pi^+\pi^-$ ($p f_{975}^0$ events are cross-hatched), b) $p\bar{K}^0\pi^+\pi^-$, c) $pK^-\pi^+$, d) pK^+K^- ($p\phi$ events are cross-hatched), e) $\Lambda^0\pi^+\pi^+\pi^-$, f) $p\pi^+\pi^+\pi^-\pi^-$, g) $pK^-\pi^+\pi^+\pi^-$.
- Fig. 5 : Effective mass plots for the subsystems in the $\Lambda_c^+ \rightarrow pK^-\pi^+$ decay channel : a) $K^-\pi^+$, b) $p\pi^+$, c) pK^- .
- Fig. 6 : Δm plots for the $D^{*+} \rightarrow D^0\pi^+$ cascade with D^0 further decaying into : a) $\pi^+\pi^-\pi^0$, b) $\pi^+\pi^+\pi^-\pi^0$, c) $K^-\pi^+\pi^0$ (right combination), d) $K^-\pi^+\pi^0$ (wrong combination), e) $K^-\pi^+\pi^+\pi^-$ (right combination), f) $K^-\pi^+\pi^+\pi^-$ (wrong combination), g) $K^-\pi^+\pi^+\pi^-\pi^0$ (right combination), h) $K^-\pi^+\pi^+\pi^-\pi^0$ (wrong combination).
- Fig. 7 : Definition of angle θ between p_{proj}^{vis} and the direction of charmed particle P in the plane defined by the beam and the parent charmed particle P.

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6

Fig. 7