CERN-TH.5823/90

ERRATA

ET ADDENDUM

ERRATA ET ADDENDUM TO CERN-TH.5823/90:
RENORMALIZATION GROUP FLOWS IN
NON-PERTURBATIVE 2-D QUANTUM GRAVITY

F. Fucito^{* †}

CERN, 1211 Geneva 23, Switzerland

and

M. Martellini^{* ‡}

Theoretical Physics, ETH-Hönggerberg, 8093 Zurich, Switzerland

* On leave of absence from: Dipartimento di Fisica, Università di Roma II "Tor Vergata" and INFN, sezione di Roma "Tor Vergata", Via Carnevale, 00173 Roma, Italy, address from September 1, 1990.

† decnet address: vxcern::fucito

* On leave of absence from: Dipartimento di Fisica, Università di Milano, 20133 Milano, Italy and INFN, Sezione di Pavia, 27100 Pavia, Italy

‡ decnet address: vaxmi::martellini

We would like to add some comments on Equation (2.14) in the second chapter of our work, together with the analysis of the Painlevé equation for $x \mapsto -\infty$. For $\Delta > 0$ there are two real solutions one with poles on the lattice lines and one without poles on the half lattice lines. Both solutions are asymptotic to (2.14) whose poles are asymptotic to those of the two solutions. In the case we have described in this paragraph the poles are obviously not on the half lattice lines.

The existence of such a solution can also be inferred by another argument: if in (2.5) we set $E = -q$ we can rewrite it as $E = T + V$, interpreting the differential equation as a classical system with kinetic energy $T = (v')^2$ and potential $V = -2v^3 + 6v$. The potential energy of such a system shows a minimum allowing for the existence of bound states i.e. pole-free solutions of the equation.

We can now study with the same methods the behaviour of the solution for $x \mapsto -\infty$. To make sense of the change of variables (2.2) we send $x \mapsto -x$. In the asymptotic limit the potential becomes $V = -2v^3 - 6v$. In this case there are no bound states and no pole-free solutions. This can be deduced also from the analysis with the Weierstrass function. In the $x \mapsto -\infty$ case the defining equation of the Weierstrass function is (2.5) with $-6v$ replaced by $6v$. This leads to a discriminant which is always negative. We are thus in the case $\Delta < 0$ for which the only real solution has poles on the real negative axis.

There is also a more straightforward procedure to obtain the results of subsection 3.2. The equations starting from (3.13) can in fact be substituted by the following ones: let $T_3 \mapsto \infty$, i.e. $\varepsilon \mapsto 0$. The change of variables:

$$\begin{cases} z = y + 3t^2, \\ f(y, t) = t + W(z), \end{cases} \quad (1)$$

in (3.12) of our work leads to:

$$6W(z) \frac{\partial W(z)}{\partial z} - \frac{\partial^3 W}{\partial z^3} = 1. \quad (2)$$

Integrating (2) once we get the Painlevé type I equation whose asymptotic behaviour is $W(z) \sim \sqrt{z}$. Using the inverse of the change of variables defined in the first part of this subsection, and remembering that the parameter $T_3 \mapsto \infty$, we get $u(x) \sim \sqrt{x}$ for $x \mapsto \infty$. We thus conclude that the solution holds in a neighbourhood of the critical point $k = 2$.

In our chapter 4 there was a mistake in the last equation in (4.3). The correct equations look like:

$$\begin{aligned} L_{-1} &= \sum_{m=1}^{\infty} (m + \frac{1}{2}) t_m \frac{\partial}{\partial t_{m-1}} + \frac{1}{8} \lambda^{-2} x^2, \\ L_0 &= \sum_{m=0}^{\infty} (m + \frac{1}{2}) t_m \frac{\partial}{\partial t_m} + \frac{1}{16}, \\ L_n &= \sum_{m=0}^{\infty} (m + \frac{1}{2}) t_m \frac{\partial}{\partial t_{m+n}} + \frac{1}{2} \lambda^2 \sum_{m=1}^n \frac{\partial^2}{\partial t_{m-1} \partial t_{n-m}}, \quad n \geq 1, \end{aligned} \quad (3)$$

where λ is the renormalized string coupling constant and $t_0 = x$. $L_{-1} \tau = 0$ gives the string equation while $L_0 \tau = 0$ is the KdV-H.

The algebra to get to the final result is clearly changed but the conclusions still hold. Here we present the new algebra and the final analysis. We start with the set of differential equations to solve:

$$[\frac{7}{2} t_3 \frac{\partial}{\partial t_2} + \frac{1}{8} \lambda^2] \tau(\mu, t_1, \dots) = 0, \quad (4)$$

$$[\frac{1}{2} x \frac{\partial}{\partial x} + \frac{5}{2} t_2 \frac{\partial}{\partial t_2} + \frac{7}{2} t_3 \frac{\partial}{\partial t_3} + \frac{1}{16}] \tau(\mu, t_1, \dots) = 0, \quad (5)$$

$$[\frac{5}{2} t_2 \frac{\partial}{\partial t_3} + \frac{1}{2} \lambda^2 \frac{\partial^2}{\partial x^2}] \tau(\mu, t_1, \dots) = 0. \quad (6)$$

Here we limit ourselves, for the sake of simplicity, to the study of the case in which only the t_2, t_3 couplings are considered. This should be enough to give us information on the flows between the $k = 2$ and $k = 3$ models. We remark that if we neglect the couplings with index ≥ 4 the equations of the $SL(2, \mathbb{C})$ algebra are the only ones contributing to the flow in the t_2, t_3 space.

If we consider (4) and (5) only, the solution is:

$$\tau = \text{const}(t_3)^{-\frac{1}{56} e^{-\frac{x^2 t_2}{2 \lambda^2 t_3}}}, \quad (7)$$

which, as we shall see, will be changed by the addition of (6).

Let us now consider the entire system. The solution of (4) is:

$$\tau = C(x, t_3) e^{-\frac{x^2}{2\lambda^2 t_3}}. \quad (8)$$

Substituting (4) and (6) in (5) we get the second order differential equation:

$$\frac{\partial^2 \tau}{\partial x^2} - a x \frac{\partial \tau}{\partial x} + \frac{1}{4} a^2 (x^2 - \frac{1}{2a}) \tau = 0, \quad (9)$$

where $a = \frac{5t_2}{7\lambda^2 t_3}$. After substituting (8) in (9) and eliminating the term with the first derivative we get:

$$\begin{aligned} C(x, t_3) &= y(x, t_3) e^{\frac{3x^2}{16\lambda^2 t_3}}, \\ \frac{\partial^2 y}{\partial x^2} - \left(\frac{z^2}{2} + \frac{5}{16}\right) y &= 0, \\ z &= \sqrt{\frac{6t_2}{7t_3} x}. \end{aligned} \quad (10)$$

The solution of the differential equation in (10) is given by a parabolic cylinder function $D_p(z)$ where $p = -\frac{13}{16}, -\frac{3}{16}$. Our function τ now has the form $\tau = F(t_3) D_p(z) e^{\frac{3x^2}{16\lambda^2 t_3}}$. Finally, substituting into (6) we should be able to determine the final form of τ .

To integrate the last differential equation we have to compute an integral whose integrand is given by $t_3^{-\frac{3}{2}} \frac{D_p^2(z)}{z^p}$. We perform this integration in the asymptotic limit $x \rightarrow \infty$, using the asymptotic form of the parabolic cylinder functions. This leads to the following form of $F(t_3)$:

$$F(t_3) \sim \text{const.} t_3^{\frac{1}{70}} (12p + 1) e^{\frac{3x^2}{24\lambda^2 t_3}}. \quad (11)$$

The asymptotic form of τ is now:

$$\tau \sim \text{const.} (\pm \lambda)^{-1} \sqrt{\frac{6t_2}{7}} t_3^{-\frac{23}{70}} p + \frac{1}{70} x^p e^{\frac{3x^2}{24\lambda^2 t_3}}. \quad (12)$$

We can now proceed with the analysis of (12). The first observation to be made is that calling $\mu = \log x$ we get a dimensionless scale as the argument of a double exponential. It is almost compulsory to stress the similarity with what happens in 4-D QG when we consider topology transitions due to wormhole effects. The identification $\mu = \log x$, although arbitrary, emerges in a very natural way the framework of the R.G. approach invoked in our work at the beginning of this section.

Can we use the form of (12) to study the flows in the t_2, t_3 space? If (12) represents the perturbation to the critical model we can think of taking as initial condition the asymptotic form of the unperturbed model and see if the perturbation makes the system flow. It is straightforward to see that using this reasoning we get no flow in the t_2, t_3 space.

ACKNOWLEDGEMENTS

F.F. would like to thank R.Benzi and A.Sagnotti for many useful discussions.

RENORMALIZATION GROUP FLOWS IN
NON-PERTURBATIVE 2-D QUANTUM GRAVITYF. Fucito^{* †}*CERN, 1211 Geneva 23, Switzerland*

and

M. Martellini^{* †}*Theoretical Physics, ETH-Hönggerberg, 8093 Zurich, Switzerland*

ABSTRACT

We study the existence of solutions of the string equations which are real and pole-free, following the work of Boutroux. We exhibit such a solution but we also point out possible problems. We further study, using perturbation theory, the flows among different models and we find that there is no flow between the $k = 2$ and $k = 3$ theories. Moreover we develop a non-perturbative renormalization group scheme to extract the scaling behaviour of the partition function of the $k = 2$ model under perturbations. The key idea is to reinterpret the recent formulation, as Virasoro constraints, of the string loop equations as generalized Callan-Symanzik renormalization group equations.

^{*} On leave of absence from: Dipartimento di Fisica, Università di Roma II "Tor Vergata" and INFN, sezione di Roma "Tor Vergata", Via Carnevale, 00173 Roma, Italy, address from September 1, 1990.

[†] decnet address: vxcern::fucito

^{*} On leave of absence from: Dipartimento di Fisica, Università di Milano, 20133 Milano, Italy and INFN, Sezione di Pavia, 27100 Pavia, Italy

[†] decnet address: vaxmi::martellini

1. Introduction

The basic information of 2-D Quantum Gravity^[1] (2-D QG from now on) is summarized by the string equations.^[2] As is well known, they are differential equations for the specific heat of the theory. For the case of pure gravity, which is described by a matrix model with criticality index $k = 2$, the differential equation is known in the mathematical literature with the name of Painlevé type I. The physics of the problem constraints the solution of this equation to be real and singularity free on the real axis. Whether such a solution exists is still a matter of discussion. The evidence gathered until now seems to point out that for $k = 2$ such a solution does not exist while for $k = 3$ it does.

Let us carefully analyze the arguments which lead to such a conclusion. For $k = 2$ the matrix integral is ill defined, being the potential unbounded. There has recently been a great deal of work trying to find 2-D theories of pure quantum gravity with a sensible perturbation theory.^[3] For the $k = 3$ theory a numerical solution has been obtained.^[4] Moreover it has numerically been shown that there is no flow between the $k = 2$ and $k = 3$ theory.^[5]

In this paper we will discuss a different approach to these problems, which stems out of the close connection between the string equation and the KdV hierarchy (KdV-H) of partial differential equations^[6] and the observation that the loop equations of this theory can be rewritten as constraint conditions satisfied by certain Virasoro operators.^[7]

With a simple change of variables, the string equation (for any value of k) can be shown to be equal to the stationary KdV (or Novikov) equation plus terms which go to zero as $x \mapsto \infty$.^[8] This change of variables takes also care of the correct asymptotic behaviour if we can find a solution of the stationary KdV equation which is bounded in the asymptotic limit. As such solutions are known from the theory of soliton equations,^[9-11] this opens the way to the analytic study of the above mentioned problems.

The solutions are in fact given in terms of theta-functions, which are sections of certain line bundles whose pole and zero structure can be completely defined. Another advantage is that they admit a parametrization in terms of Riemann surfaces with genus $g \geq 1$, a fact which comfortably fits this formalism. Now the string equation for a model with generic k is a differential equation that can be classified according to its singularities as was shown by the Painlevé school.^[12] The computations involved in showing the absence of movable essential singularities, grow with the degree of the derivative. They have been performed up

to degree four and there is no problem to go to even higher degree. Now, given the existence of solutions which admit an infinite number of movable double poles, an exhaustive study of their properties (limited to the case corresponding to $k = 2$ for our models, i.e. the Painlevé type I equation) has been carried out by Boutroux.^[13]

The main finding of this work is the proof that the modulus of the difference between the solution of the Painlevé equation and its asymptotic limit is everywhere arbitrarily small in the x complex plane. This allows the author to infer all of the interesting properties of the solution at finite x from its asymptotic limit. This limit is performed after the perturbation theory has been resummed. Moreover there is a real and singularity-free solution (which is not the famous triply truncated solution) which is asymptotic to the degenerate Weierstrass function. This solution, though, has a pole which is arbitrarily close to the real axis. Similar conclusions were also drawn by Novikov.^[14] This could be the source of the problems with perturbation theory.^{*} As of now, we have no answer to this question. Before describing the plan of the paper we want to stress that the existence of a non-stable vacuum for the $k = 2$ theory, has no straightforward connections with the existence of a real and singularities-free solution of the Painlevé equation. Moreover, the perturbation theory we think of is not to be confused with the topological expansion. In particular the above mentioned asymptotic limit is not the first order of perturbation of the topological expansion.

In chapter one we briefly review Boutroux's theory and elucidate its connection with the theory of solitons. This allows us to argue that the difficulties of the $k = 2$ case, should not be there for higher k 's. In chapter two we study the flow of the $k = 3$ in the $k = 2$ theory solving the associated differential equations in the asymptotic limit, with an analysis inspired by Ref.5: the result is that there is no flow. In chapter four we give a non-perturbative formulation of this problem: we show that the subalgebra $SL(2, \mathbb{C})$ of the Virasoro operators found in Ref.7 leads to renormalization group type equations for the partition function of the theory. We solve this differential equation and show once again that there is no flow between these two theories. The partition function solution of the renormalization group equation has a non-perturbative functional dependence similar to that of 4-D QG where we observe topology transitions due to wormhole effects. Wormhole type effects have also been recently detected, in 2-D QG, in numerical simulations.^[15]

* The authors thank L. Alvarez-Gaumé for this remark.

2. Asymptotic Analysis of the Solutions of the String Equation.

For the sake of simplicity we will concentrate on the $k = 2$ case. The differential equation under study is:

$$x = \tilde{R}_2[u] = u^2 - \frac{1}{3}u'', \quad (2.1)$$

which, after the change of variables:

$$\begin{cases} u(x) = x^{\frac{1}{2}}v(\xi), \\ \xi(x) = \frac{5}{6}x^{\frac{5}{4}}, \end{cases} \quad (2.2)$$

becomes:

$$v'' - 3v^2 + 3 = \frac{4}{23}\xi^{\frac{2}{5}} - \frac{v'}{\xi}. \quad (2.3)$$

In the limit $\xi \mapsto \infty$ (2.3) becomes:

$$v'' - 3v^2 + 3 = 0. \quad (2.4)$$

If we multiply (2.4) by $2v'$ and integrate, we get:

$$(v')^2 = 2v^3 - 3v - q, \quad (2.5)$$

where q is the integration constant. The discriminant of the polynomial on the r.h.s. of (2.5) vanishes for $q = \pm 2$. (2.5) is solved by the Weierstrass function $\wp(\xi)$.^[16] Boutroux's analysis and the connection with the theory of solitons start from this point.

Now consider ξ to be the inverse function of the solution of (2.3) and ξ_0 the same for (2.4). We will show that along a path P the modulus of the difference between these two quantities is arbitrarily small.

Let $|\xi_0| > \varepsilon^{-1}$. Now let us place ourselves outside the circumference centered at the origin with radius ε^{-1} . It is then always possible to find a number a such that the coefficients of v, v' in (2.3) are bounded by $a\varepsilon$. Let us now take as initial conditions for the differential

equation:

$$\begin{cases} v(\xi_0) = \eta, \\ v'(\xi_0) = \eta', \\ q = -(\eta')^2 + 2\eta^3 - 3\eta. \end{cases} \quad (2.6)$$

Now while we go from ξ_0 to ξ , v will go along a path L of length l , from η to v . Multiplying (2.3) by $2v'$ and integrating:

$$(v')^2 = 2v^3 - 3v - q + 2 \int_{\xi_0}^{\xi} (s(v')^2 + tvv')d\xi = 2v^3 - 3v - q + 2 \int_{\eta}^v (sv' + tv)v dv. \quad (2.7)$$

s, t are the coefficients of the members in the r.h.s. of (2.3). In the r.h.s. of (2.7), v is considered as an independent variable. Along l is always possible to find numbers h, b, d such that $|v| < h$ and $b < |v'| = |\sqrt{2v^3 - 3v - q}| < d$ from which:

$$|(v')^2 - 2v^3 + 3v + q| < ale(d + h), \quad (2.8)$$

follows. Dividing (2.8) by $2v^3 - 3v - q$, multiplying it by v^{-1} and finally integrating leads to:

$$|\xi_0 - \xi| < \sqrt{\varepsilon}, \quad (2.9)$$

as it is always possible to find a path L such that $\frac{(d+h)ae^2}{d^2} < \varepsilon^{-\frac{1}{2}}$.

If v describes a loop starting from its initial conditions (2.6), the variable $\xi_0 \mapsto \xi_0 + \omega$, where ω is one of the two periods of $\wp(\xi_0)$. On the contrary $\xi \mapsto \xi_{10}$. In performing this operation we have to check that the loop does not get too close either to one of the three zeroes of the r.h.s. of (2.7) or to a pole. This condition can always be matched by deforming the path to go around the pole or the zeroes. Furthermore the loop has to encircle two of such zeroes. According to which zeroes are enclosed, ξ_0 is shifted by ω or ω' , the other period of $\wp(\xi_0)$ and we can define points ξ_{10} and ξ_{01} . Clearly $|\xi_{10} - \xi_0 - \omega| < \sqrt{\varepsilon}$. Turning more times around the loop in one direction or the other we can define additions of more periods and points $\dots, \xi_{20}, \xi_{10}, \xi_{-10}, \dots$. Joining these points we obtain the so-called periodic line.

Using (2.9) we can now identify the points $\xi_0 + \omega\omega'$ and $\xi_0 + \omega'\omega$ thus building a quadrilateral which tends to a parallelogram as ξ_0 becomes bigger. It is then not difficult

to prove that inside this parallelogram v has one double pole $\bar{\xi}_0$ which is isolated, that there are three zeroes of v' located in the neighbourhood of the three zeroes of the r.h.s. of (2.5) and that $|v - \varphi|$ is arbitrarily small with ε . As $q \mapsto \pm 2$ the quadrilateral gets stretched and finally becomes a single line as we will later see in more detail.

These results can easily be extended to all values of ξ : starting from any of the vertices of the quadrilateral we can in fact build new quadrilaterals to patch the space of complex ξ . To have (2.9) hold in all of the complex ξ plane it is enough to multiply the r.h.s. of (2.3) by a parameter $0 \leq \lambda \leq 1$ and substitute the bound $a\varepsilon$ for $|s|, |t|$ in (2.7) with $|s|, |t| < H$. Tuning λ properly, (2.9) can be extended to the entire complex ξ plane.

We can now study the problem of the reality and of the absence of singularities of v along the real ξ line. We first proceed to establish a connection with the theory of solitons which will later turn out to be very fruitful.^[17]

It was first noticed by Ince^[18] that the potential φ in the Schrödinger problem was leading to a single band solution. The function $\varphi(\xi_0)$ is defined on the period parallelogram that is on the complex torus, the Jacobian of our problem. Moreover from the theory of elliptic functions:

$$\varphi(\xi_0) = -2 \frac{\partial^2}{\partial \xi_0^2} \log \vartheta_{11}(\xi_0) + \text{constant}, \quad (2.10)$$

which is the basis for the argument that τ -functions can be expressed in terms of ϑ -functions. ϑ_{11} is the ϑ -function of characteristic $\{\frac{1}{2}, \frac{1}{2}\}$.

What are the reality properties of $\varphi(\xi_0)$? They can be listed according to the values of the discriminant Δ of the r.h.s. of (2.5):

1. $\Delta > 0$. In this case the periods are such that one is real and the other purely imaginary. The period lattice may be chosen to be rectangular. $\varphi(\xi_0)$ is real on the lattice lines $Re\xi_0 = m\omega, iIm\xi_0 = n\omega'$ and on the half-lines $Re\xi_0 = (m + \frac{1}{2})\omega, iIm\xi_0 = (n + \frac{1}{2})\omega'$. m, n are integer numbers. The solution on the lattice lines clearly exhibits poles while the other solution is the one we studied, as an example, in Ref.8 where we wrote (2.10) in terms of ϑ_{00} which has characteristic $[0, 0]$. The two ϑ functions can be expressed one in terms of the other, using simple identities for elliptic functions. We see here that the conditions on reality and absence of singularities, are reflected as conditions on the moduli and the Jacobian. In this simple example we see in fact that

our moduli have to be imaginary and that we have to take only a real section of the Jacobian (the line $Re\xi_0$).

2. $\Delta < 0$. The periods are complex conjugate and $\varphi(\xi_0)$ is real on the line $Re\xi_0 = \frac{\pi}{2}(\omega + \omega'), iIm\xi_0 = \frac{\pi}{2}(\omega - \omega')$. On these line we clearly find poles.
3. $\Delta = 0$. Two or all the roots coincide. $\varphi(\xi_0)$ degenerates in a simply periodic function as the periods go to ∞ . If the real period goes to infinity then $\varphi(\xi_0) = 1 + \frac{3}{\sin^2 \sqrt{3}(\xi_0 - \bar{\xi}_0)}$. If the imaginary period goes to infinity then $\varphi(\xi_0) = -1 + \frac{3}{\sin^2 \sqrt{3}(\xi_0 - \bar{\xi}_0)}$. This case will be important in the real solution exhibited by Boutroux. If both periods go to ∞ , then $\varphi(\xi_0) = (\xi_0 - \bar{\xi}_0)^{-2}$.

This is true in the asymptotic limit. Can we extend the real solution to all of $\xi \in \mathbb{R}$? Here we have to distinguish according to whether $k \geq 2$.

Let us turn on a perturbation of the KdV equation:

$$\frac{\partial u}{\partial t} - \frac{\partial}{\partial x} R_{k+1} = \varepsilon K[u], \quad (2.11)$$

where $K[u]$ is a differential polynomial in u .^[19] The effect of the perturbation is to couple the Jacobian with the moduli space as the matrix of periods and the moduli become ε -dependent. Notice that this conclusion still holds for the stationary case of (2.11). Now (2.2) fixes the asymptotic behaviour of the solution. The parameters which are left, have to take care of the reality and absence of poles. Now, given the coupling between the moduli space and the Jacobian, the solutions at $k = 2$, which is parametrized by a torus, might not have enough parameters to satisfy all the requirements. If we take surfaces with higher genus for $k = 2$, the situation does not change as the additional parameters give trivial relations. The space of parameters for solutions with $k > 2$, on the contrary, seems big enough to allow for real solutions.

After these heuristic arguments we proceed, for the case $k = 2$, to study what happens to the real solution of the $\varphi(\xi_0)$ function we discussed before following Boutroux's analysis.

Rewrite (2.3) and (2.4) as:

$$v'' = 3v^2 + 3 - \frac{5\mu}{\mu + 4} \frac{v'}{\xi} + \frac{4\mu(2 - \mu)}{(\mu + 4)^2} \frac{v}{\xi^2}, \quad (2.12)$$

where $0 \leq \mu \leq 1$. μ interpolates between (2.3) and (2.4). We have already seen that there is a real solution without poles for $\mu = 0$ with $-2 \leq q \leq 2$. When we turn μ on, this

solution gets deformed but it stays finite and oscillates with a series of maxima and minima which correspond to the addition of the periods. We want to know whether the number of oscillations is infinite: otherwise a pole is present on the real axis. The extrema are bounded by the two branches of the function $3v^2 - 3 + \frac{4(\mu(2-\mu)}{(\mu+1)^2} \frac{v}{\xi^2} = 0$. If q_0 was the original value for the asymptotic solution we find how q_0 is changed by the addition of a period. Using (2.6) we get:

$$q_1 = (v')^2 - 2\eta^3 + 3\eta = q_0 - 2 \int_{\xi_0}^{\xi} \frac{(v')^2}{\xi} d\xi + \frac{8}{25} \int_{\xi_0}^{\xi} \frac{vv'}{\xi^2} d\xi. \quad (2.13)$$

Inside the range of validity of our approximations it is the first integral, which is essentially positive, that dominates the second. Then $q_1 < q_0$ and in general $q_{n+1} < q_n$. We then deduce that the oscillations are an infinite number and as $-2 \leq q \leq 2$ and $q_0 < 2$, the infinite sequence $q_0, \dots, q_n, \dots$ tends to -2 . So for $v(\xi)$ the real axis is a periodic line and asymptotically:

$$\begin{cases} v(\xi) \mapsto -1 + \frac{3}{8\pi^2 \sqrt{\delta(\xi - \xi_0)}}, \\ q \mapsto -2, \\ \omega \mapsto \frac{2\pi}{\sqrt{3}}. \end{cases} \quad (2.14)$$

The pole ξ_0 gets arbitrarily close to the real axis.

3. A Perturbative Analysis of KdV flows

3.1 ASYMPTOTIC ANALYSIS IN THE SCALING VARIABLE AND COUPLING

In this subsection, following the strategy elaborated in Ref.5, we show how to use asymptotic methods to determine the existence or absence of a certain flow between two models with different index of criticality.

For massive one-matrix random models, the string equation becomes:

$$x = \sum_{k=0}^{\infty} (k + \frac{1}{2}) \epsilon_k R_k[u], \quad (3.1)$$

where $t_0 = x$ and the $R_k[u]$ are the Gelfand-Dikii differential polynomials.

In the case of the flux between a $k = 3$ and a $k = 2$ theory, (3.1) takes the form:

$$x = a\hat{R}_3[u] + b\hat{R}_2[u], \quad (3.2)$$

where $a = -\frac{35}{64}t_3$, $b = \frac{15}{32}t_2$ and $\hat{R}_2[u]$, $\hat{R}_3[u]$ are the differential polynomials defined as:

$$\begin{aligned} \hat{R}_2[u] &= u^2 - \frac{1}{3}u'' = \frac{16}{3}R_2, \\ \hat{R}_3[u] &= u^3 - uu'' - \frac{1}{2}(u')^2 + \frac{1}{10}u''' = -\frac{32}{5}R_3. \end{aligned} \quad (3.3)$$

Let us take as our initial condition the solution of (3.2) for $x \mapsto \infty$ with fixed t_3 and $t_2 = 0$ and study its evolution towards the region where $t_2 \mapsto \infty$. The change of variables:

$$\begin{cases} u(x) = x^{\frac{1}{3}}v(y), \\ v(x) = \frac{7}{6}x^{\frac{1}{6}}, \end{cases} \quad (3.4)$$

casts (3.2) in the form:

$$\begin{aligned} a[(v^3 - vv'' - \frac{1}{2}(v')^2 + \frac{1}{10}v''') - \frac{5}{6}x^{-\frac{7}{3}}v^2 - \frac{4}{27}x^{-\frac{14}{3}}v - \frac{1}{48}x^{-\frac{7}{3}}v' - \frac{1}{6}x^{-\frac{7}{6}}vv'] \\ - \frac{1}{15}x^{-\frac{7}{3}}v'' + \frac{7}{30}x^{-\frac{7}{6}}v'''] + bx^{-\frac{1}{3}}[v^2 - \frac{1}{3}v' + \frac{2}{27}x^{-\frac{7}{3}}v - \frac{2}{9}x^{-\frac{7}{6}}v'] = 1. \end{aligned} \quad (3.5)$$

Since $b = \frac{15}{32}t_2$ is a free parameter we can study the solution of (3.5) in the following limit:

$$\begin{cases} x \mapsto \infty, \\ t_2 \mapsto \infty, \\ \frac{15}{32}t_2x^{-\frac{1}{3}} = c, \end{cases} \quad (3.6)$$

where c is constant. (3.5) now becomes:

$$1 = a\hat{R}_3[v] + c\hat{R}_2[v], \quad (3.7)$$

the Novikov equation, whose solution is of the form:

$$v(y) \sim \frac{\partial^2}{\partial y^2} \log \vartheta(\vec{B}\vec{x} + \vec{c}|\tau). \quad (3.8)$$

As was argued in Ref.8, a suitable choice of the matrix period τ and of the shift vector $\vec{c}$ makes v real and pole-free on the real axis. Then we conclude that the asymptotic behaviour $x^{\frac{1}{3}}$ has not been spoiled by the "perturbation" $b\hat{R}_2[u]$: there is no flow between the $k = 2$ and $k = 3$ theories. The solution $v(y)$ though, is not exactly the solution for the $k = 3$ model: it is modulated by the stationary term $c\hat{R}_2[v]$.

The last thing we discuss in this subsection is the independence of our result from the t_2 behaviour of (3.6). Looking at (3.5) we realize that to keep a perturbation term we have only other two possibilities besides the one we used before:

1. $\frac{15}{32}t_2x^{-\frac{8}{3}} = \text{constant}$, $t_2 \sim x^{\frac{8}{3}}$, $x \mapsto \infty$,
2. $\frac{15}{32}t_2x^{-\frac{3}{2}} = \text{constant}$, $t_2 \sim x^{\frac{3}{2}}$, $x \mapsto \infty$.

In both cases it is still $bx^{-\frac{1}{3}}(v^2 - \frac{1}{3}v'')$ that dominates. Now $bx^{-\frac{1}{3}}$ is not a constant anymore, and the asymptotic limit has to be chosen such that $bx^{-\frac{1}{3}} \sim O(1)$. This is completely equivalent to the Whitham method (see Ref.9): its implementation, though, does not change our conclusion.

3.2 ASYMPTOTIC ANALYSIS IN THE SCALING VARIABLE

In this subsection we study the solution of the KdV flow interpolating between the $k = 2$ and $k = 3$ models. For this purpose we start from (3.2) and:

$$\frac{\partial u(x, t_k)}{\partial t_k} = \frac{\partial}{\partial x} \hat{R}_{k+1}[u(x, t_k)], \quad (3.9)$$

that, after a derivation is cast in the form:

$$-\frac{35}{64}t_3\frac{\partial^2}{\partial t_2} + \frac{15}{32}t_2\frac{\partial}{\partial x}\hat{R}_2[u] = 1. \quad (3.10)$$

Let us change variables:

$$\begin{cases} x = (\frac{2}{7})^{\frac{1}{3}}yt_3^{-\frac{1}{3}}, \\ t_2 = \sqrt{2t}, \\ u(x, t_2) = (\frac{2}{7})^{\frac{2}{3}}t_3^{\frac{2}{3}}f(y, t). \end{cases} \quad (3.11)$$

(3.10) becomes:

$$\frac{\partial f(y, t)}{\partial t} - 6f(y, t)\frac{\partial f(y, t)}{\partial y} + \frac{\partial^3 f(y, t)}{\partial y^3} = \varepsilon K(t), \quad (3.12)$$

where $\varepsilon = t_3^{-\frac{5}{3}}$, $K(t) = -\frac{\varepsilon}{\sqrt{t}}$ and $c = \frac{32}{5\sqrt{2}}(\frac{2}{7})^{\frac{5}{3}}$. (3.12) is the fundamental equation of this subsection. It describes a perturbation by $\varepsilon K(t)$ of the time-dependent KdV equation describing the dynamics of the solution interpolating between the $k = 2$ and $k = 3$ models.

Notice that this is not the usual perturbation of the KdV equation which is of the form $\varepsilon K(u)$ where the differential polynomial $K(u)$ is such that $K(u) \mapsto 0$ for $u \mapsto 0$.^[20] We look for a solution of (3.12) of the form:

$$f(y, t) = 2\sqrt{t}v(y). \quad (3.13)$$

Substituting in (3.12):

$$-24t^{\frac{3}{2}}v(y)\frac{\partial v(y)}{\partial y} + 2t\frac{\partial^3 v(y)}{\partial y^3} + (1 + \varepsilon c) = 0. \quad (3.14)$$

In (3.14) the time t is decoupled and can be reabsorbed by a further change of variables:

$$\begin{cases} y = t^{\frac{1}{10}}z, \\ v(y) = \frac{1}{2}t^{-\frac{7}{10}}W(z), \end{cases} \quad (3.15)$$

which leads to:

$$6W(z)\frac{\partial W(z)}{\partial z} - \frac{\partial^3 W}{\partial z^3} = 1 + \varepsilon c. \quad (3.16)$$

(3.16) is now a perturbation of the stationary KdV equation which is obtained in the limit $\varepsilon c \mapsto -1$. Integrating (3.16) once, we get a Painlevé type I equation of the form:

$$3\hat{R}_2[u] = (1 + \varepsilon c)z. \quad (3.17)$$

Using the techniques developed in Ref.8 we can study the asymptotic behaviour of (3.17). After performing a change of variables once again:

$$\begin{cases} \xi = \frac{4}{3}[\frac{1}{3}(1 + \varepsilon c)z]^{\frac{5}{4}}, \\ W(z) = \sqrt{\frac{1}{3}(1 + \varepsilon c)z}G(\xi), \end{cases} \quad (3.18)$$

where $G(\xi)$ satisfies the Novikov equations. To summarize we put all the pieces together to get the asymptotic behaviour of the $f(z, t_2)$:

$$f(z, t_2) \sim t_2^{-\frac{5}{2}}\sqrt{\frac{1}{3}(1 + \varepsilon c^{-\frac{5}{3}})}\sqrt{z}G(\frac{4}{13}(1 + \varepsilon c^{-\frac{5}{3}})^{\frac{5}{4}}z^{\frac{5}{4}}). \quad (3.19)$$

The effect of the perturbation is just to modulate the $k = 2$ solution. Even if we take the naive limit $\frac{t_2}{t_3} \gg 1$ the solution remains in a neighbourhood of the critical point $k = 2$.

4. A Renormalization Group Analysis of KdV flows

The analysis performed in the previous section is purely perturbative. Here we want to develop some non-perturbative analytic tools.

Our starting point is the intriguing relation among 2-D QG loop equations, the τ -function of the KdV-H and the twisted $A_1^{(1)}$ infinite dimensional Kac-Moody algebra.^[21] The main output of the works in Ref 7 is the equation:

$$Z(x, t_1, \dots) = \tau^2(x, t_1, \dots), \quad (4.1)$$

which holds true for even matrix potentials. The τ -function satisfies a system of homogeneous linear differential equations which take the form:

$$L_n \tau = 0, \quad n \geq -1, \quad (4.2)$$

where the differential operators L_n are given by:

$$\begin{aligned} L_{-1} &= \sum_{m=1}^{\infty} (m + \frac{1}{2}) t_m \frac{\partial}{\partial t_{m-1}} + \frac{1}{8} \lambda^{-2} x^2, \\ L_0 &= \sum_{m=0}^{\infty} (m + \frac{1}{2}) t_m \frac{\partial}{\partial t_m} + \frac{1}{16}, \\ L_n &= \sum_{m=0}^{\infty} (m + \frac{1}{2}) t_m \frac{\partial}{\partial t_{m+n-1}} + \frac{1}{2} \lambda^2 \sum_{m=1}^n \frac{\partial^2}{\partial t_{m-1} \partial t_{n-m}}, \quad n \geq 1, \end{aligned} \quad (4.3)$$

where λ is the renormalized string coupling constant and $t_0 = x$. $L_{-1} \tau = 0$ gives the string equation while $L_0 \tau = 0$ is the KdV-H.

Our approach begins with the observation that $L_0 \tau = 0$ is very similar to the Callan-Symanzik renormalization group (RG) equation^[22] where the β -function and the anomalous dimension are:

$$\beta_m = 2(m + \frac{1}{2}) t_m, \quad \gamma = \frac{1}{8}. \quad (4.4)$$

The t_m 's play the role of bare coupling constants and the RG scale is $\mu = \log x$. We also have to take in account, in a self-consistent way, the other constraints coming from (4.2).

The minimal requirement is to satisfy the $SL(2, \mathbb{C})$ subalgebra:

$$\begin{cases} L_{-1} \tau = 0, \\ L_0 \tau = 0, \\ L_1 \tau = 0, \end{cases} \quad (4.5)$$

which leads to:

$$[\frac{\partial}{\partial \mu} + \frac{x}{3!} \sum_{m \geq 1} \beta_{m+1} \frac{\partial}{\partial t_m} + \frac{1}{12} \frac{x^2}{t_1}] \tau(\mu, t_1, \dots) = 0, \quad (4.6)$$

$$[\frac{\partial}{\partial \mu} + \sum_{m \geq 1} \beta_m \frac{\partial}{\partial t_m} + \frac{1}{8}] \tau(\mu, t_1, \dots) = 0, \quad (4.7)$$

$$[\frac{\partial}{\partial \mu} + \sum_{m \geq 1} \beta_m \frac{\partial}{\partial t_m} + \lambda^2 \frac{\partial^2}{\partial x^2}] \tau(\mu, t_1, \dots) = 0. \quad (4.8)$$

Following the strategy of the RG we look for a solution $\tau(\mu, t_1, \dots)$ of the form:

$$\tau(\mu, t_1, \dots) = e^{-\frac{\mu}{8}} \bar{\tau}(\mu, \bar{t}_1(\mu), \dots), \quad (4.9)$$

where the functions $\bar{t}_m = \bar{t}_m(\mu, t_m)$ are chosen in such a way that (4.9) satisfies (4.7). This implies:

$$\begin{cases} \frac{\partial \bar{t}_m}{\partial \mu} = -\bar{\beta}_m = -(2m+1) \bar{t}_m, \\ \bar{t}_m(\mu=0) = t_m. \end{cases} \quad (4.10)$$

The solution of (4.10) is:

$$\bar{t}_m = t_m e^{-(2m+1)\mu}. \quad (4.11)$$

Up to now we have just exploited $L_0 \tau = 0$. We will now show that $L_1 \tau = 0$ will determine the functional dependence of $\bar{\tau}$ with respect to $\bar{t}_m$, while $L_{-1} \tau = 0$ will give a set of algebraic non-linear recursion relations among the couplings t_m 's.

We are now ready to solve the system of differential equations (4.5). Notice that with respect to the standard use of the RG analysis this solution will lead to an explicit form of the partition function (see (4.1)).

Plugging (4.7) into (4.8) and using (4.9) we get:

$$\frac{\partial \bar{\tau}}{\partial \mu} \simeq -\left(\frac{e^{2\mu}}{8\lambda^2} - \frac{1}{8}\right)\bar{\tau}, \quad (4.12)$$

where we have neglected the higher derivatives in μ . Changing variables as:

$$\frac{\partial \bar{\tau}}{\partial \mu} = \frac{\partial \bar{\tau}}{\partial \bar{\mu}} \frac{\partial \bar{\mu}}{\partial \mu} = -\bar{\beta}_m \frac{\partial \bar{\tau}}{\partial \bar{\mu}}, \quad (4.13)$$

we finally get:

$$\frac{\partial \bar{\tau}}{\partial \bar{\mu}} \simeq \frac{\left(\frac{e^{2\mu}}{8\lambda^2} - \frac{1}{8}\right)}{(2m+1)\bar{\mu}_m} \bar{\tau} = \frac{\bar{t}_m^\alpha + \delta_m}{\gamma_m \bar{\mu}_m} \bar{\tau}, \quad (4.14)$$

where we used (4.11) and $\alpha = \frac{-2}{(2m+1)}$, $\delta_m = -\lambda^2 \bar{t}_m^\alpha$, $\gamma_m = 8(2m+1)\delta_m$.

The announced recursive equation for the couplings $\bar{t}_m$ now follows from the substitution of (4.7) in (4.6):

$$\sum_{m \geq 1} \left[\frac{e^\mu}{\bar{t}_1} \bar{\beta}_{m+1} - \bar{\beta}_m \right] \bar{\tau} = \left(\frac{1}{8} - \frac{e^{3\mu}}{12\lambda^2 \bar{t}_1} \right) \bar{\tau}. \quad (4.15)$$

Expanding in μ and equating the coefficients with the same power of μ we have an infinite set of relations connecting different $\bar{t}_m$'s. The first equation of this set looks like:

$$\bar{t}_2 = \frac{2}{3}\bar{t}_1^2 + \frac{1}{120\lambda^2}. \quad (4.16)$$

Taking (4.11) into account, (4.16) leads to a non-trivial relation between t_2 and t_1 .

The solution of (4.14) is:

$$\bar{\tau}(\mu, \bar{t}_1, \dots) \simeq \bar{\tau}(\mu, 0, \dots) \bar{t}_m^{\frac{\delta_m}{\gamma_m}} e^{\frac{\bar{t}_m}{\gamma_m}}, \quad \forall \bar{t}_m. \quad (4.17)$$

Substituting (4.17) into (4.9) and using (4.11) leads to:

$$\tau(\mu, t_1, \dots) \simeq \tau(\mu, 0, \dots) \bar{t}_m^{\frac{\delta_m}{\gamma_m}} e^{\frac{1}{120\lambda^2} t_1^2}, \quad \forall t_m. \quad (4.18)$$

The effect of the factor $\rho = e^{\frac{1}{120\lambda^2} t_1^2}$ is typically of non-perturbative nature. It is almost compulsory to stress the similarity with what happens in 4-D QG when we consider topology transitions due to wormhole effects.^[23] For $\lambda^2 < 0$ ρ is maximized for $\mu \mapsto -\infty$ which is in that region of values of x where all topologies, according to the common wisdom, contribute to the topological expansion.

Let us see now how we can use this result to get non-perturbative information on the flows among models with different critical indices k . Recalling the discussion of the previous section we write:

$$Z_{k=2}^{R.G.}(\mu, t_2) \simeq Z_{k=2}(\mu, 0) t_2^{\frac{30}{2}} e^{-\frac{\mu}{4} \rho^2}, \quad (4.19)$$

where:

$$Z_{k=2}(\mu, x) \sim e^{-x^{\frac{1}{2}}} \sim e^{-\frac{x^{\frac{1}{2}}}{2}} (1 + O(\mu^2)), \quad x \mapsto \infty. \quad (4.20)$$

For the $k = 3$ model we get:

$$Z_{k=3}(\mu, x) \sim e^{-x^{\frac{1}{3}}} \sim e^{-\frac{x^{\frac{1}{3}}}{3}} (1 + O(\mu^2)), \quad x \mapsto \infty. \quad (4.21)$$

We now try to get (4.21) from (4.19) after having substituted (4.20) in the latter. This gives a condition at $O(\mu)$ on λ :

$$\lambda \approx \lambda_c^{R.G.} = \sqrt{\frac{3}{5}}. \quad (4.22)$$

This value does not coincide with the λ_c which is derived in the one-matrix random model for $k = 3$, which is $\lambda_c = 60$. This means we cannot match the critical value of the $k = 3$ model, starting from the $k = 2$ one, that is there is no R.G. trajectory between these two models.

ACKNOWLEDGEMENTS

One of the authors (M.M.) would like to thank J. Fröhlich, G. Felder and the Department of Theoretical Physics at ETH for their kind hospitality during part of this work.

REFERENCES

1. A. Polyakov, Phys. Lett. 103B (1981) 207; Mod. Phys. Lett. A2 (1987) 899.
2. D. Gross and A. A. Migdal, Phys. Rev. Lett. 64 (1990) 127; University of Princeton preprint PUPT-1159 (1989);
E. Brézin and V. A. Kazakov, Phys. Lett. 236B (1990) 144;
M. R. Douglas and S. H. Shenker, Nucl. Phys. B335 (1990) 635.
3. E. Marinari and G. Parisi, Phys. Lett. 240B (1990) 375; Università di Roma II "Tor Vergata" preprint ROM2F-90-29 (1990);
F. David, Mod. Phys. Lett. A5 (1990) 1019;
J. Ambjorn, J. Greensite and S. Varsted, Niels Bohr Institute preprint NBI-HE-90-39 (1990);
M. Karliner and A. Migdal, University of Princeton preprint PUPT-1191 (1990).
4. E. Brezin, E. Marinari and G. Parisi, Phys. Lett. 242B (1990) 35.
5. M. R. Douglas, N. Seiberg and S. H. Shenker, Rutgers preprint RU-90-19 (1990).
6. T. Banks, M. R. Douglas, N. Seiberg and S. H. Shenker, Rutgers University preprint RU-89-50 (1989);
M. R. Douglas, Rutgers University preprint RU-89-51 (1989);
P. Ginsparg and J. Zinn-Justin, Harvard University preprint HUTP-90/A004 (1990).
7. R. Dijkgraaf, H. Verlinde and E. Verlinde, University of Princeton preprint PUPT-1184 (1990);
M. Fukuma, H. Kawai and R. Nakayama, University of Tokyo preprint UT-562 (1990).
8. F. Fucito, A. Gamba and M. Martellini, CERN preprint CERN-TH.5767/90 (1990).
9. S. Novikov, S. V. Manakov, L. P. Pitaevskii and V. E. Zakharov, *Theory of Solitons*, Plenum Publishing Company (New York, 1984).
10. B. A. Dubrovin, V. B. Matveev and S. P. Novikov, Russ. Math. Surv. 31 (1976) 59.
11. A. Das, *Integrable Models*, World Scientific (Singapore, 1989);
M. J. Ablowitz and H. Segur, *Solitons and the Inverse Scattering Transform*, SIAM (Philadelphia, 1981).
12. E. L. Ince, *Ordinary Differential Equations*, ed. Dover (New York, 1956);
E. Hille, *Ordinary Differential Equations in the Complex Domain*, ed. J. Wiley & Sons (New York, 1976);
13. M. P. Boutroux, Ann. Ec. Norm. 30 (1913) 255; Ann. Ec. Norm. 31 (1914) 99.
14. unpublished work that we received from L. Alvarez-Gaumé.
15. M. E. Agishtein and A. Migdal, Princeton University preprint PUPT-1180 (1990).
16. H. Bateman and A. Erdélyi, *Higher Transcendental Functions*, vol. 3, ed. McGraw-Hill (New York, 1955).
17. for a review see: B. A. Dubrovin, Russ. Math. Surv. 36 (1981) 11.
18. E. L. Ince, Proc. Roy. Soc. Edinb. 60 (1940) 83.
19. I. M. Krichever, Russ. Math. Surv. 44 (1989) 145;
N. Ercolani and H. P. McKean, Invent. Math. 99 (1990) 483.
20. V. I. Karpman, Physica Scripta 20 (1979) 462.
21. see Ref. 7 and F. Fucito and M. Martellini, CERN preprint CERN-TH.5797/90 (1990).
22. see for example D. Gross in *Methods in Field Theory*, Les Houches session XXVIII, ed. by R. Balian and J. Zinn-Justin (North-Holland, 1976).
23. for a review see: I. Klebanov, *Wormholes and the Cosmological Constant Problem*, Lectures delivered at TASI 89, University of Colorado, Boulder, June 1989, Princeton University preprint PUPT-1153 (1989).