CERN 56-31

Theoretical Study Division
15th December 1956

ORGANISATION EUROPÉENNE POUR LA RECHERCHE NUCLÉAIRE
CERN EUROPEAN ORGANIZATION FOR NUCLEAR RESEARCH

Continuous groups in quantum mechanics

by

W. Pauli

G E N E V E

Contents

	<u>Page</u>
I. Continuous groups and their associated Lie algebra	1
II. Representations of Lie-rings and Lie-groups	5
- The theorem of complete reducibility of every finite reducible representation of semisimple Lie-rings (Lie-groups)	9
III. Lie algebra of inhomogeneous rotation groups	12
- Representations of the Lie algebra of the three- dimensional inhomogeneous rotation group	13
IV. The four-dimensional rotation group	19
V. The Lorentz group	23
- The inhomogeneous Lorentz group	25

Continuous groups in quantum mechanics *

by

W. Pauli **

I. Continuous groups and their associated Lie algebra

An abstract group G is a set of elements which possess a law of composition such that from any two elements a, b of G taken in a given order arises an element c . The following conditions must be satisfied :

1) Associative law :

$$c(ba) = (cb)a$$

2) Unit element : there exists in G an element I such that for any a of G

$$Ia = aI = a$$

3) Every element a of G possesses an inverse a^{-1} , the composition of which with a gives the unit element :

$$aa^{-1} = a^{-1}a = I.$$

The continuous groups which occur in physics appear, however, not with such abstract elements but as groups of transformations of variables. For these the associative law 1) and the existence of the unity element 2) are automatically fulfilled and 3) means the one-one valuedness of the transformation. The continuous transformation groups in question can be divided into two classes :

- 1) The finite continuous groups, the elements of which are determined by a finite number of parameters. Examples : rotation group, Lorentz group.
- 2) The infinite continuous groups in the elements of which occur arbitrary functions. Examples : gauge group, coordinate transformations in general relativity, canonical transformations in classical mechanics and in quantum mechanics.

* The present notes are based on a series of lectures given by Prof. W. Pauli, of the E.T.H., Zürich, to the members of the CERN Theoretical Study Division at Copenhagen in September 1955.

** The author is indebted to R. Jost and A.R. Edmonds for help in the preparation of these notes.

These lectures will deal with the first class only; I wish, however, to emphasize that according to my opinion the second class will presumably turn out to be of greater importance for physics in future.

In order to treat the continuous groups of the first class we use the infinitesimal methods of S. Lie.^(1,2) For the groups of physics the conditions of differentiability are indeed always fulfilled for suitably chosen parameters. Many interesting questions can be treated already in the frame of the infinitesimal elements which define the so-called Lie-ring.

To define it we use as a starting point the fact that the group elements which deviate not too largely from the unit element I , can be parametrized in the following way: to every r -tuple of numbers $s^1, \dots, s^r$ in an environment of $s^\alpha = 0$, ($\alpha = 1, \dots, r$) corresponds in a one-one valued way a group element which we shall simply denote by $(s^1, \dots, s^r)$ or (s) , with $I \sim (0)$. The number r is uniquely determined and is called the dimension of the group. To the product of two elements (s) and (t) there corresponds again a point u in the considered environment.

One has

$$u^\alpha = \varphi^\alpha(s^1, \dots, s^r, t^1, \dots, t^r) \quad (1)$$

where the functions φ^α can be assumed here to be differentiable an arbitrary finite number of times with respect to the arguments $s^1, \dots, s^r, t^1, \dots, t^r$. From $I \sim (0)$ it follows

$$\varphi^\alpha(0, t) = t^\alpha; \quad \varphi^\alpha(s, 0) = s^\alpha. \quad (2)$$

The development of $\varphi^\alpha(s, t)$ is, therefore, (compare chap. 9 of Pontrjagin⁽³⁾)

$$\varphi^\alpha(s, t) = s^\alpha + t^\alpha + s^\alpha \rho_\gamma^\alpha s^\beta t^\gamma + s^\alpha \rho_{\gamma\delta}^\alpha s^\beta s^\gamma t^\delta + h^\alpha \rho_{\gamma\delta}^\alpha s^\beta t^\gamma t^\delta + \dots \quad (3)$$

The commutator $q = (s)(t)(s^{-1})(t^{-1})$ of the two elements s and t is of fundamental importance. The development

$$q^\alpha(s, t) = c^\alpha \rho_\gamma^\alpha s^\beta t^\gamma + \dots \quad (4)$$

gives rise to the occurrence of the structure constants $c^\alpha \rho_\gamma^\alpha$ of the group,

connected with the coefficients $a^\alpha_{\beta\gamma}$ in (3) by

$$c^\alpha_{\beta\gamma} = a^\alpha_{\beta\gamma} - a^\alpha_{\gamma\beta}.$$

Hence

$$c^\alpha_{\beta\gamma} = -c^\alpha_{\gamma\beta}. \quad (5)$$

One uses here that in the reciprocal $(s)^{-1} = (\tilde{s})$ the following development holds

$$\tilde{s}^\alpha = -s^\alpha + a^\alpha_{\beta\gamma} s^\beta s^\gamma + \dots$$

A further condition for the $c^\alpha_{\beta\gamma}$ follows from the associative law which is for our parametrisation formulated by

$$\varphi(\varphi(s, t), u) = \varphi(s, \varphi(t, u)).$$

Inserting the development (3) for φ one sees that the equation is identically fulfilled in first and second order; in third order, however, one finds for the coefficient of $s^\beta t^\gamma u^\delta$ the condition

$$a^\alpha_{\sigma\delta} a^\sigma_{\beta\gamma} - a^\alpha_{\beta\sigma} a^\sigma_{\gamma\delta} = h^\alpha_{\beta\gamma\delta} + h^\alpha_{\beta\delta\gamma} - g^\alpha_{\beta\gamma\delta} - g^\alpha_{\gamma\beta\delta}. \quad (6)$$

Antisymmetrisation with respect to the indices β, γ, δ makes the right side vanish, while the left side can be expressed with the help of (4) by the $c^\alpha_{\beta\gamma}$ and gives rise to the condition

$$c^\alpha_{\beta\sigma} c^\sigma_{\gamma\delta} + c^\alpha_{\gamma\sigma} c^\sigma_{\delta\beta} + c^\alpha_{\delta\sigma} c^\sigma_{\beta\alpha} = 0. \quad (7)$$

Besides this no further condition follows.

We can now proceed to the definition of the Lie-ring of the infinitesimal group elements.

If $(s(\tau))$ is a one-dimensional submanifold (curve) of the group, containing the unit element $s^\alpha = 0$ for $\tau = 0$, one considers the tangent of the curve for $\tau = 0$, supposed as existent and defined by

$$\xi^\alpha = \left(\frac{ds^\alpha}{d\tau} \right)_{\tau=0}.$$

The r -dimensional vector space of these tangent vectors ξ^α is the Lie-ring.

From the two curves $(s(\tau))$ and $(t(\tau))$ a third curve with the parameter

$$\tau^1 = \tau^2$$

can be constructed by

$$(u(\tau^1)) = (s(\tau))(t(\tau))(s(\tau))^{-1}(t(\tau))^{-1}.$$

With $\xi^\alpha = \left(\frac{ds^\alpha}{d\tau}\right)_{\tau=0}$ and $\eta^\alpha = \left(\frac{dt^\alpha}{d\tau}\right)_{\tau=0}$ one has according to (4) the development

$$u^\alpha(\tau^1) = e^\alpha_{\beta\gamma} \xi^\beta \eta^\gamma \tau^1 + \dots$$

The tangent vector of this third curve is therefore given by

$$\zeta^\alpha = [\xi, \eta]^\alpha = e^\alpha_{\beta\gamma} \xi^\beta \eta^\gamma. \quad (8)$$

With the help of r linearly independent basis vectors e_α ($\alpha = 1, \dots, r$), one can also write a vector ξ with the components ξ^α as

$$\xi = \xi^\alpha e_\alpha.$$

Defining then

$$[e_\beta, e_\gamma] = e^\alpha_{\beta\gamma} e_\alpha \quad (8a)$$

and considering ξ^α, η^α as ordinary commuting c -numbers, one obtains agreement with (8) for

$$\xi = \xi^\alpha e_\alpha; \eta = \eta^\alpha e_\alpha; \zeta = [\xi, \eta] = \zeta^\alpha e_\alpha.$$

The position of the indices in (8) or (8a) shows the transformation law by a transition from the basis e_α to a new basis e'_α . The existence of the bracket-operation is characteristic for a Lie-ring. It has the properties

$$\left. \begin{aligned} [\xi, c_1 \eta_1 + c_2 \eta_2] &= c_1 [\xi, \eta_1] + c_2 [\xi, \eta_2] \\ [\xi, \eta] + [\eta, \xi] &= 0 \\ [[\xi, \eta] \zeta] + [[\eta, \zeta] \xi] + [[\zeta, \xi] \eta] &= 0 \end{aligned} \right\} \quad (9)$$

The last property is the Jacobi-identity which follows from the associative law and its consequence (7) for the $c^\alpha_{\beta\gamma}$. It distinguishes a Lie-ring from the vectors and its vector product in a three dimensional vector space.

The mathematicians have proved that Lie-groups with the same structure constants $c^\alpha_{\beta\gamma}$ are isomorphic in the small (i.e. in a sufficiently small environment of the unit element) and that to every Lie-ring (the product $[\xi, \eta]$ of which has the properties (9)) there exists a Lie-group. (See for instance Pontrjagin⁽³⁾).

II. Representations of Lie-rings and Lie-groups

Suppose that to each element ξ^α of the Lie-ring there corresponds a linear operator $A(\xi)$ (linear transformation of a vector space, the dimension of which is in general different from r) with the properties :

$$\left. \begin{aligned} A(\xi + \eta) &= A(\xi) + A(\eta) \\ A(c \cdot \xi) &= c \cdot A(\xi) \\ A([\xi, \eta]) &= A(\xi)A(\eta) - A(\eta)A(\xi) = [A(\xi), A(\eta)] \end{aligned} \right\} \quad (10)$$

the last of which fulfils the Jacobi-identity identically, as the multiplication of linear operators is associative. Such a correspondence of operators (linear transformations) with the elements of a Lie-ring is called a representation of the Lie-ring.

In all cases considered here a representation of the Lie-ring determines uniquely also a representation of the Lie-group, namely a one to one correspondence of linear operators $G(s)$ to the group-elements (s)

$$(s) \rightarrow G(s)$$

with the properties

$$(s)(t) \rightarrow G(s)G(t)$$

and

$$(s)^{-1} \rightarrow G^{-1}(s)$$

from which also follows $G(0) = I$.

We do not give here a general proof for the possibility of this extension, but the reverse passage from a representation of the Lie-group to a representation of the Lie-ring is trivial: if $(s(\tau))$ is a "smooth" curve, then $G(s(\tau))$ associates with it a curve $G(\tau)$ in the space of the operators considered. If now

$$\xi^\alpha = \left(\frac{ds^\alpha}{d\tau} \right)_{\tau=0}$$

then the operator

$$A(\xi) = \left(\frac{dG}{d\tau} \right)_{\tau=0}$$

can be made by definition to correspond to ξ^α . It can easily be seen that this correspondence fulfils all the conditions (10).

In the part of the lecture contained in these notes we are dealing with representations of Lie-rings rather than Lie-groups. The Lie-ring permits always a particular representation by r -dimensional linear transformations called the adjoint representation. One obtains it by interpreting $[\alpha, \xi]$ as a linear transformation of the vector ξ where $\alpha, \beta, \dots$ denote elements of the Lie-ring, too :

$$[\alpha, \xi] = A(\alpha)\xi . \tag{11}$$

This is a representation because

$$A(\beta)A(\alpha)\xi = [\beta[\alpha, \xi]]$$

$$A(\alpha)A(\beta)\xi = [\alpha[\beta, \xi]]$$

hence, in view of the Jacobi-identity,

$$[A(\alpha), A(\beta)]\xi = [\alpha[\beta, \xi]] - [\beta[\alpha, \xi]] = [[\alpha, \beta]\xi] = A([\alpha, \beta])\xi .$$

The adjoint representation is not necessarily faithful, which means that it is possible that $A(\alpha) = 0$ for $\alpha \neq 0$. This is so for those α 's, for which $[\alpha, \xi] = 0$. These particular α 's are again a Lie-ring to which the "centre" of the Lie-group corresponds.

Written in matrices the adjoint representation is given by the correspondence

$$\xi^k \rightarrow o_{k\beta}^\alpha$$

where α, β are row and column-indices of the matrix for each given k .

We mention here that to every representation $A(\xi)$ there exists a contravariant representation given by

$$A'(\xi) = -A^T(\xi)$$

where T is the transposed matrix obtained by permutation of rows and columns (for operators : the adjoint operator). For the Lie-group this corresponds to the substitution $(G^{-1})^T(s)$ for $G(s)$.

We use the following general definitions of group theory. A group is Abelian if all its elements commute. Clearly for an Abelian group all commutators and hence all structure constants vanish.

A subgroup of a group G is a subset of elements of G , which satisfy the group postulates.

An invariant subgroup N of a group G is a subgroup which contains all the conjugates of its elements: i.e. if n is any element of N and s any element of G , then sns^{-1} is in N .

A group is simple if it contains no invariant subgroup besides the unit element.

A group is semisimple if it has no Abelian invariant subgroup besides the unit element. In this case the adjoint representation is faithful.

We also say that the Lie-ring associated with a semisimple subgroup is semisimple. Cartan has given a sufficient and necessary condition for the group or Lie-ring to be semisimple, which uses only properties of the Lie-ring:

construct from the structure constants $c^{\alpha}_{\beta\gamma}$ the symmetric tensor

$$g_{ik} = g_{ki} = c^{\rho}_{i\sigma} c^{\sigma}_{k\rho} . \quad (12)$$

The condition in question is then, that the determinant of the g_{ik} is different from zero :

$$\det |g_{ik}| \neq 0 . \quad (13)$$

Summarizing we give the following key of translation of concepts related to Lie-groups to concepts related to Lie-rings :

subgroup $\leftrightarrow$ subring : $[\alpha, \beta]$ is in the subring together with α and β ;
 invariant subgroup $\leftrightarrow$ Ideal J : if α in J, ξ in Lie-ring R, then $[\xi, \alpha]$ in J;
 Abelian group $\leftrightarrow [\xi, \eta] = 0$;
 semisimple group $\leftrightarrow \det |g_{ik}| \neq 0$.

For semisimple groups there exists a theorem of Casimir, which we shall use in the following. To formulate it, one considers the normalized subdeterminants g^{ik} of g_{ik} , defined by

$$g^{i\rho} g_{\rho k} = \delta_i^k \quad (14)$$

which exist if and only if $\det |g_{ik}| \neq 0$. Then one considers the quadratic form

$$F = g^{\rho\sigma} e_{\rho} e_{\sigma} \quad (15)$$

where the e_{ρ} are again r linearly independent elements of the Lie-ring. The theorem of Casimir states then : F commutes with all elements of the Lie-ring or

$$[F, e_{\tau}] = g^{\rho\sigma} (e_{\rho} [e_{\sigma}, e_{\tau}] + [e_{\rho}, e_{\tau}] e_{\sigma}) = 0 . \quad (16)$$

Before we indicate the proof, we remark first that the ordinary product $e_{\rho} e_{\sigma}$ is not defined inside the Lie-ring. It is only defined either for representations of the Lie-ring by linear operators (or matrices) or in some amplified algebra. We adopt here the first point of view which also justifies the

definition of $[F, e_\tau]$, used in (16).

To prove Casimir's theorem, one shows first, using the condition (7) for the structure constants and the definition (12) of the g_{ik} , that

$$e_{ijk} = g_{i\rho} e^{\rho}_{jk} \quad (17)$$

is antisymmetric in i and j ,

$$e_{ijk} = -e_{jik} \quad (18)$$

Raising the indices i and j with help of the g^{ik} by

$$e^{ij}_{,k} = g^{\sigma i} g^{rj} e_{\sigma rk} = g^{rj} e^i_{rk} \quad (19)$$

it follows then that $e^{ij}_{,k}$, too is antisymmetric in i and j ,

$$e^{ij}_{,k} = -e^{ji}_{,k} \quad (20)$$

From the expression for $[F, e_\tau]$ given in (16) it follows, however, by (8a) and (19)

$$[F, e_\tau] = e^{\rho\sigma}_{,\tau} (e_\rho e_\sigma + e_\sigma e_\rho)$$

which vanishes in view of (20), q.e.d.

The theorem of complete reducibility of every finite reducible representation of semisimple Lie-rings (Lie-groups).

We wish to recall, firstly, the general concepts concerning representations which also apply to Lie-rings and Lie-groups. (cf. ^(4,5,6,7).)

Two representations $r \rightarrow A_1(r)$, $r \rightarrow A_2(r)$ are called equivalent if for every element r of the group or ring considered there is a constant matrix U such that

$$U A_1(r) U^{-1} = A_2(r).$$

A representation is reducible if it leaves a subspace of the space

R of the representation invariant. If so, there is a transformation U which puts all the matrices into the form

$$\begin{pmatrix} P & 0 \\ R & Q \end{pmatrix}$$

where the submatrix P performs a transformation on the subspace.

If the representation leaves invariant two subspaces which together span the total original space, then the matrices may be written in the form

$$\begin{pmatrix} P & 0 \\ 0 & Q \end{pmatrix}$$

and the representation is called decomposable or completely reducible. For semisimple Lie-groups Peter and Weyl^(*) proved the famous theorem, that all finite reducible representations are also decomposable. The proof of these authors contains essentially an integration over the whole (finite) volume of the group ("unitary trick"). As the theorem is a purely algebraic statement concerning Lie-rings alone, a proof was desirable which only uses algebraic properties of the Lie-ring.

For the three dimensional rotation group (for which the quadratic form F, defined by (15) can be simply written $L_x^2 + L_y^2 + L_z^2$) such a proof was first given by Casimir and van der Waerden^(*).

A very elegant, purely algebraic proof for all semisimple Lie-rings was given by Brauer^(*) later on.

Brauer's proof is based on his general theorem, which can be formulated as follows for Lie-rings :

Theorem of Brauer : If all representations of a Lie-ring of the particular form

$$\begin{pmatrix} A & 0 \\ C & 0 \end{pmatrix}$$

* Professor van der Waerden kindly informed me that the generalisation of this proof to all semisimple groups in this paper contains an error which was first pointed out by Freudenthal.

where A is an irreducible representation, are completely reducible, then all finite representations of the Lie-ring are completely reducible.

We cannot give here the proof of Brauer's theorem. Once it is given, one can use Casimir's quadratic form F to prove the premises of this theorem for all semisimple Lie-rings with the help of the following:
Auxiliary theorem : The eigenvalue λ of Casimir's quadratic form F defined for any semisimple Lie-ring, is different from zero for every non-trivial finite irreducible representation of the Lie-ring.

By the "trivial representation" of the Lie-ring we understand the one in which 0 corresponds to every element. We do not give here the proof of the auxiliary theorem either, which uses the construction of all irreducible representations for all semisimple Lie-algebras by Cartan and Weyl.

Using both Brauer's general theorem and the auxiliary theorem, one can proceed in the following way. Since the form F of Casimir commutes with every element of the Lie-ring, it is (by Schur's well known Lemma, see Weyl⁽¹¹⁾ p.83) a multiple of the unit matrix, $\lambda \cdot I$, where according to the auxiliary theorem $\lambda \neq 0$, except for the trivial representation. Hence F can be written

$$F = \begin{pmatrix} \lambda \cdot I & 0 \\ K & 0 \end{pmatrix}$$

For $\lambda \neq 0$, by application of the transformation

$$T = \begin{pmatrix} I & 0 \\ \frac{K}{\lambda} & 0 \end{pmatrix}$$

one obtains

$$T F T^{-1} = \begin{pmatrix} \lambda \cdot I & 0 \\ 0 & 0 \end{pmatrix}$$

Since the elements of the Lie-ring commute with F , this transformation decomposes them, i.e. the reducible representation is decomposable.

III. Lie algebra of inhomogeneous rotation groups

Such a group consists of the linear transformations on an n-dimensional space which leave invariant forms of the type

$$\sum_{i=1}^n (x_i - y_i)^2$$

where $(x_1, \dots, x_n), (y_1, \dots, y_n)$ are the coordinates of points in this space. The typical transformation is thus

$$x_i \rightarrow x'_i = \sum_{j=1}^n x_j a_{ji} + \sum_{i=1}^n b_i$$

where the coefficients a_{ji} form an orthogonal matrix; the coefficients a_{ji} and b_i are usually supposed to be real.

Thus this group includes translations as well as rotations; it is not semisimple, for the translations form an invariant Abelian subgroup. The infinitesimal operators are obtained by consideration of the appropriate Taylor series associated with functions on the n-dimensional space. We obtain :

$$e_{\lambda\mu} = x_\lambda \frac{\partial}{\partial x_\mu} - x_\mu \frac{\partial}{\partial x_\lambda} ; d_\lambda = \frac{\partial}{\partial x_\lambda}$$

It is sufficient to use this special representation of the Lie-algebra in order to derive its computation relations :

- i) $[e_{\lambda\mu}, e_{\rho\sigma}] = \delta_{\lambda\sigma} e_{\mu\rho} + \delta_{\mu\rho} e_{\lambda\sigma} - \delta_{\lambda\rho} e_{\mu\sigma} - \delta_{\mu\sigma} e_{\lambda\rho}$
- ii) $[d_\lambda, e_{\mu\nu}] = \delta_{\lambda\mu} d_\nu - \delta_{\lambda\nu} d_\mu$
- iii) $[d_\lambda, d_\mu] = 0.$

The Lie algebra for the homogeneous rotation group is given by omitting the d_λ .

We write conventionally $e_{\mu\nu} = -e_{\nu\mu} = iJ_{\mu\nu} ; d_\mu = ip_\mu$. $J_{\mu\nu}$ and p_μ are then Hermitian operators.

Representations of the Lie algebra of the three-dimensional inhomogeneous rotation group

In the case of $n = 3$ we also write $J_{12} = J_3$, $J_{23} = J_1$, $J_{31} = J_2$ and $J_1 \pm iJ_2 = J_{\pm}$, $p_1 \pm ip_2 = p_{\pm}$, (J_3 and p_3 remain).

The commutation relations are thus :

$$\begin{aligned} [J_3, J_+] &= J_+ ; \quad [J_3, J_-] = -J_- ; \quad [J_+, J_-] = 2J_3 ; \\ [J_+, p_-] &= [J_-, p_+] = 2p_3 ; \quad [J_3, p_{\pm}] = [p_3, J_{\pm}] = \pm p_{\pm} . \end{aligned}$$

Other commutator brackets are zero.

There are two invariants, (i.e. quantities which commute with all 6 operators), which may be constructed from the infinitesimal operators, namely

$$\vec{p}^2 = p_1^2 + p_2^2 + p_3^2 \quad \text{and} \quad \vec{p} \cdot \vec{J} = \vec{J} \cdot \vec{p} = p_1 J_1 + p_2 J_2 + p_3 J_3$$

$\vec{J}^2 = J_1^2 + J_2^2 + J_3^2$ is not an invariant of the inhomogeneous group. We now consider the value of $[\vec{J}^2, p_j]$. Take for example

$$\begin{aligned} [\vec{J}^2, p_1] &= J_2 [J_2, p_1] + [J_2, p_1] J_2 + J_3 [J_3, p_1] + [J_3, p_1] J_3 \\ &= +i[-J_2 p_3 - p_3 J_2 + J_3 p_2 + p_2 J_3] = 2i[p_2 J_3 - J_2 p_3] . \end{aligned}$$

Similar results of course will be obtained for $[\vec{J}^2, p_2]$ and $[\vec{J}^2, p_3]$. We obtain in a similar way $[\vec{J}^2, [\vec{J}^2, p_j]] = -4(\vec{J} \cdot \vec{p})J_j + 2(\vec{J}^2 p_j + p_j \vec{J}^2)$, a relation which will be employed in the investigation of the representations of the Lie algebra.

We shall assume J_3 is represented by a diagonal matrix :

$$(m|J_3|m) = m \delta_{mm} ,$$

The commutation rules give

$$(m|J_{\pm}|m') \neq 0 \quad \text{only for} \quad m' = m \mp 1 .$$

i.e. the m values in a given representation differ by integers. We see

also that J_+J_- is diagonal in m .

Now let us write $\varphi(m)$ as the (m,m) matrix element of J_+J_-

$$(m|J_+|m-1)(m-1|J_-|m) = \varphi(m)$$

and obtain from the commutation relations

$$\varphi(m) - \varphi(m+1) = 2m ; \text{ i.e. } \varphi(m) = \text{constant} -m(m-1).$$

Now

$$\mathbf{J}^2 = \frac{1}{2}(J_+J_- + J_-J_+) + J_z^2.$$

The eigenvalue of $\mathbf{J}^2$ in this representation is therefore

$$\frac{1}{2}(\varphi(m) + \varphi(m+1)) + m^2$$

since $(m|J_-J_+|m) = (m|J_-|m+1)(m+1|J_+|m) = \varphi(m+1)$. Now assume the representation is of finite degree; the trace of the commutator $[J_+, J_-]$ is then zero, i.e. $\sum m = 0$ which implies the m values are either integer or half-odd-integer. The finiteness of the representation may be proved from the assumption that it is Hermitian (i.e. that $\varphi(m) \neq 0$). We obtain in either case

$$-j \leq m \leq j ; \quad \varphi(-j) = \varphi(j+1) = 0$$

and the above constant $= j(j+1)$ = eigenvalue of $\mathbf{J}^2$.

The above discussion allows us to write down the matrices of J_z, J_+ and J_- in the j, m representation. They may be taken with the usual choice of an arbitrary phase, as

$$(jm|J_z|jm) = m$$

$$(jm|J_+|jm-1) = \sqrt{(j+m)(j-m+1)}$$

$$(jm|J_-|jm+1) = \sqrt{(j-m)(j+m+1)}$$

Let us now consider the representation of the p_i ; no reference

will be made at this stage to the fact that the p_i commute with each other; the theory developed will thus be applicable to the representation of the homogeneous rotation group in four dimensions. The commutation relations with the p_i then show that the non-zero matrix elements of the p_i may be factorized, the left-hand components being independent of m (cf. Weyl⁽¹²⁾ p.200). To determine the dependence on m of the matrix element of p_- , we refer to the relation $[p_-, J_-] = 0$. For example, if

$$j' = j$$

we have

$$\frac{(jm|p_-|jm+1)}{[(j-m)(j+m+1)]^{\frac{1}{2}}} = \frac{(jm-1|p_-|jm)}{[(j-m+1)(j+m)]^{\frac{1}{2}}}$$

i.e. each ratio is independent of m , and we may write

$$(jm|p_-|jm+1) = (j|p_-|j) \sqrt{(j-m)(j+m+1)}.$$

The dependence of the matrix of p_z on m is determined by use of the relation $[J_+, p_-] = 2p_z$, and that of p_+ by reference to the Hermitian nature of the representation.

We obtain thus :

$$(jm|p_z|jm) = (j|p|j)m$$

$$(jm|p_+|jm-1) = (j|p|j) \sqrt{(j+m)(j-m+1)}$$

$$(jm|p_-|jm+1) = (j|p|j) \sqrt{(j-m)(j+m+1)}$$

$$(jm|p_z|j+1, m) = (j|p|j+1) \sqrt{(j+m+1)(j-m+1)}$$

$$(jm|p_+|j+1, m-1) = (j|p|j+1) \sqrt{(j-m+2)(j-m+1)}$$

$$(jm|p_-|j+1, m+1) = - (j|p|j+1) \sqrt{(j+m+2)(j+m+1)}$$

$$\left. \begin{aligned} (jm|p_z|j-1, m) &= (j|p|j-1) \sqrt{(j+m)(j-m)} \\ (jm|p_+|j-1, m-1) &= - (j|p|j-1) \sqrt{(j+m)(j+m-1)} \\ (jm|p_-|j-1, m+1) &= (j|p|j-1) \sqrt{(j-m)(j-m-1)} \end{aligned} \right\} \text{N.B. } \begin{aligned} (j|p|j-1) &= 0 \\ \text{if } j &= 0, \frac{1}{2} \end{aligned}$$

In particular the diagonal matrix element of $p_+p_- - p_-p_+$ is :

$$(jm|p_+p_- - p_-p_+|jm) = 2m \{ -|(j|p|j+1)|^2 (2j+3) + \\ + |(j|p|j-1)|^2 (2j-1) + |(j|p|j)|^2 \} .$$

Let us define

$$(j|p|j+1)(j+1|p|j)(2j+3)(2j+1) = \varphi(j) .$$

Then

$$(j|p|j-1)(j-1|p|j)(2j+1)(2j-1) = \varphi(j-1)$$

$$\{\varphi(-1) = 0\}$$

and

$$(jm|p_+p_- - p_-p_+|jm) = 2m \left\{ \frac{-\varphi(j) + \varphi(j+1)}{2j+1} + (j|p|j)^2 \right\} .$$

The invariant

$$\vec{p}^2 = p_j^2 + \frac{1}{2}(p_+p_- + p_-p_+)$$

has the eigenvalue

$$p^2 = \left\{ \varphi(j) \cdot \frac{j+1}{2j+1} + \varphi(j-1) \frac{j}{2j+1} + |(j|p|j)|^2 j(j+1) \right\}$$

and the invariant

$$\vec{J} \cdot \vec{p} = \vec{p} \cdot \vec{J}$$

the eigenvalue

$$(j|p|j) = j(j+1).$$

These two quantities are constants for a given representation. Now $4(\vec{J} \cdot \vec{p})\vec{J}$ has clearly non-zero matrix elements only when they are diagonal in j . It follows from this and the relation

$$[\vec{J}^2, [\vec{J}^2, \vec{p}]] = -4(\vec{J} \cdot \vec{p})\vec{J} + 2(\vec{J}^2\vec{p} + \vec{p}\vec{J}^2)$$

that

$$(j'm' | (\vec{J}^2)^2 \vec{p} - 2\vec{J}^2 \vec{p} \vec{J}^2 + \vec{p}(\vec{J}^2)^2 - 2(\vec{J}^2 \vec{p} + \vec{p} \vec{J}^2) | j'' m')$$

is zero for $j' \neq j''$.

If $(j'|p|j'') \neq 0$ we have then

$$(j' + j'' + 2)(j' + j'')(j' - j'' + 1)(j' - j'' - 1) = 0,$$

i.e. if $j' \neq j''$ we have the selection rule $j' - j'' = \pm 1$.

We have not up to now made use of the fact that the p 's commute with each other. We now take this into account. We distinguish two cases according to whether the eigenvalue

$$(j|p|j)j(j+1) = c$$

of $\vec{J} \cdot \vec{p}$ is zero or not.

a) $c = 0$. The matrix element of $p_+p_- - p_-p_+ (=0)$ gives

$$\varphi(j) = \varphi(j-1) = \text{constant}.$$

The constant is clearly equal to the eigenvalue p^2 of $\vec{p}^2$. It follows from the definition of $\varphi(j)$ that

$$|(j|p|j+1)|^2 = \frac{p^2}{(2j+3)(2j+1)}$$

Now if the lowest possible value of j , namely j_0 , were non-zero, we would have a contradiction in the relations

$$0 = 2m \left(\frac{-\varphi(j) + \varphi(j-1)}{2j+1} \right)$$

if we suppose $p^2 \neq 0$. (i.e. due to the supposed $\varphi(j_0 - 1) = 0$.) If we take $j_0 = 0$ we have in this case $m = 0$ also, avoiding the contradiction. Hence we have an infinite representation with j taking all values from zero.

b) $C \neq 0$. The matrix element of $p_+p_- - p_-p_+$ now gives

$$\frac{-\varphi(j) + \varphi(j-1)}{2j+1} + \frac{C^2}{j^2(j+1)^2} = 0 ;$$

$$\text{i.e.} \quad \varphi(j) - \varphi(j-1) = C^2 \left(\frac{1}{j^2} - \frac{1}{(j+1)^2} \right)$$

$$\text{Hence} \quad \varphi(j) + \frac{C^2}{(j+1)^2} = \text{constant.}$$

Substitution into the expression for the eigenvalue p^2 of $\vec{p}^2$ gives

$$\varphi(j) \frac{j+1}{2j+1} + \left[\varphi(j) - C^2 \left(\frac{1}{j^2} - \frac{1}{(j+1)^2} \right) \right] \frac{j}{2j+1} + \frac{C^2}{j(j+1)} = \varphi(j) + \frac{C^2}{(j+1)^2} ,$$

i.e. the constant just mentioned equals p^2 .

We now determine the minimum value of j , namely j_0 , by setting $\varphi(j_0 - 1) = 0$. Then

$$C^2 = j_0^2 p^2 \quad \text{and} \quad C = \pm \sqrt{p^2} \cdot j_0.$$

We obtain therefore two non-equivalent representations corresponding to the eigenvalues of $\vec{J} \cdot \vec{p}$ having the two values $\pm \sqrt{p^2} \cdot j_0$. Now j_0 is the component of $\vec{J}$ parallel to $\vec{p}$:

$$\pm j_0 = \frac{\vec{J} \cdot \vec{p}}{\sqrt{p^2}}$$

and can be integer or half-odd-integer. If we consider the group which

includes reflections, the representation includes both signs of C , i.e. the reflection reverses the sign of $\vec{J} \cdot \vec{p}$.

IV. The four-dimensional rotation group

We consider first the Lie algebra for the proper orthogonal group, with the invariant form $x_1^2 + x_2^2 + x_3^2 + x_4^2$. We write down the commutation relations for the 6 infinitesimal operators

$$J_{\lambda\mu} = i \left\{ x_\mu \frac{\partial}{\partial x_\lambda} - x_\lambda \frac{\partial}{\partial x_\mu} \right\}$$

which we rewrite for convenience in the following way :

$$J_{23} \rightarrow M_1, \quad J_{31} \rightarrow M_2, \quad J_{12} \rightarrow M_3$$

$$J_{41} \rightarrow N_1, \quad J_{42} \rightarrow N_2, \quad J_{43} \rightarrow N_3.$$

Note that the M_i have the same commutation relations as the infinitesimal operators of the homogeneous 3-dimensional orthogonal group.

$$[M_1, M_2] = iM_3, \quad [M_2, M_3] = iM_1, \quad [M_3, M_1] = iM_2$$

$$[M_1, N_2] = [N_1, M_2] = iN_3$$

$$[M_2, N_3] = [N_2, M_3] = iN_1$$

$$[M_3, N_1] = [N_3, M_1] = iN_2$$

$$[M_1, N_1] = [M_2, N_2] = [M_3, N_3] = 0$$

$$[N_1, N_2] = iM_3, \quad [N_2, N_3] = iM_1, \quad [N_3, N_1] = iM_2.$$

We see further that the relations between the M 's and the N 's are the same as those between the J 's and p 's in the case of the inhomogeneous 3-dimensional rotation group, whose representation we have already considered. The only difference is that the N 's do not commute among themselves as do the components of $\vec{p}$.

We have again two invariants :

$$F = \frac{1}{2} \sum_{\mu < \nu} J_{\mu\nu}^2 = \frac{1}{2} (\vec{M}^2 + \vec{N}^2)$$

$$G = J_{41} J_{23} + J_{42} J_{31} + J_{43} J_{12} = \vec{M} \cdot \vec{N} = \vec{N} \cdot \vec{M}$$

(regarding the triples (M_1, M_2, M_3) and (N_1, N_2, N_3) as sets of vector components).

A certain linear combination of the infinitesimal operators is of interest :

$$\vec{K} = \frac{1}{2}(\vec{M} + \vec{N}) \quad ; \quad \vec{L} = \frac{1}{2}(\vec{M} - \vec{N}).$$

The components of $\vec{K}$ and $\vec{L}$ commute with each other; i.e.

$$[K_i, L_j] = 0 \quad \text{for all } i, j.$$

In addition

$$[K_1, K_2] = iK_3 \quad \text{etc.}$$

$$[L_1, L_2] = iL_3 \quad \text{etc.}$$

and

$$F = \vec{K}^2 + \vec{L}^2 \quad , \quad G = \vec{K}^2 - \vec{L}^2 .$$

This shows that the four-dimensional rotation group may be considered as the direct product of two 3-dimensional rotation groups. This relation is only valid for the proper rotation group and not for the Lorentz group. If we take $\vec{K}$ and $\vec{L}$ as Hermitian operators we get, using the known theory of the homogeneous 3-dimensional group, a set of finite representations of the 4-dimensional group, each representation with

$$\vec{K}^2 = k(k+1) \quad , \quad \vec{L}^2 = \ell(\ell+1)$$

and degree $(2k+1)(2\ell+1)$. Each representation may be labelled by the number pair (k, ℓ) and the invariants F and G have the values

$$F = k(k+1) + \ell(\ell+1)$$

$$G = k(k+1) - \ell(\ell+1) = (k-\ell)(k+\ell+1).$$

The mirror representations (k, ℓ) and (ℓ, k) correspond to a change of sign $\vec{N}$, i.e. to a reflection of the spacial coordinates.

We shall now adapt the representation theory of the inhomogeneous 3-dimensional rotation group to give that of the homogeneous 4-dimensional group. We start by making $\vec{M}^2$, $\vec{N}^2$ and M_3 diagonal; we know already that the eigenvalues of $\vec{M}^2$ and M_3 are $j(j+1)$ and m respectively, where $-j \leq m \leq j$. From the previous theory, in analogy with the treatment of $(p_+p_- - p_-p_+)$,

$$\langle jm | N_+ N_- - N_- N_+ | jm \rangle = 2m \{ -\langle j | N | j+1 \rangle \langle j+1 | N | j \rangle (2j+3) + \langle j | N | j-1 \rangle \langle j-1 | N | j \rangle (2j-1) + \langle j | N | j \rangle^2 \}$$

$$\langle jm | N^2 | jm \rangle = \{ \langle j | N | j+1 \rangle \langle j+1 | N | j \rangle (2j+3) (j+1) + \langle j | N | j-1 \rangle \langle j-1 | N | j \rangle (2j-1) j + \langle j | N | j \rangle^2 j(j+1) \}.$$

We write

$$\langle j | N | j-1 \rangle \langle j-1 | N | j \rangle (2j+1) (2j-1) = \varphi(j)$$

hence

$$\langle j | N | j+1 \rangle \langle j+1 | N | j \rangle (2j+1) (2j+3) = \varphi(j+1)$$

On evaluating $G = \vec{M} \cdot \vec{N}$ we get $G = \langle j | N | j \rangle j(j+1)$. Now $N_+ N_- - N_- N_+ = 2M_3$; hence we may rewrite the above equations :

$$\varphi(j+1) - \varphi(j) = -(2j+1) + \frac{G^2(2j+1)}{j^2(j+1)^2} \quad \text{when } j \neq 0$$

$$2F = \vec{M}^2 + \vec{N}^2 = \varphi(j+1) \left(\frac{j+1}{2j+1} \right) + \varphi(j) \left(\frac{j}{j(j+1)} \right) + \frac{G^2}{j(j+1)} + j(j+1).$$

The first of the new equations gives us

$$\varphi(j) = \text{constant} - \frac{G^2}{j^2} - j^2$$

and putting this result into the second equation gives

$$\text{constant} = 2F + 1 .$$

We discuss first the finite representations; these are for the proper orthogonal group also Hermitian.

There are two cases :

a) $G = 0$: a maximum $j = n$ must exist, i.e. $\varphi(n+1) = 0$. Hence

$$\varphi(j) = 2F + 1 - j^2 \quad ; \quad 2F + 1 = (n+1)^2 \quad ;$$

thus

$$0 \leq j \leq n$$

(We get no boundary condition for $j = 0$).

b) $G \neq 0$: now we get a minimum $j = j_0 \neq 0$; i.e. $j_0 \leq j \leq n$. Then

$$\varphi(j_0) = \varphi(n+1) = 0 ,$$

therefore

$$\frac{G^2}{(n+1)^2} + (n+1)^2 = \frac{G^2}{j_0^2} + j_0^2 = 2F + 1$$

and

$$G^2 \left(\frac{1}{j_0^2} - \frac{1}{(n+1)^2} \right) = (n+1)^2 - j_0^2 \quad ;$$

i.e.

$$G^2 = j_0^2 (n+1)^2 .$$

Hence

$$2F = (n+1)^2 + j_0^2 - 1 = n(n+2) + j_0^2 .$$

The degree of the representation is

$$\sum_{j=j_0}^n (2j+1) = (n+1)^2 - j_0^2 = (n+1+j_0)(n+1-j_0) .$$

The representation just found must be equivalent to one of those already derived; the degrees and the values of the invariants must coincide, hence we get $n = k + \ell$; $j_0 = |k - \ell|$. i.e.

- a) if $G > 0$ $k = \frac{1}{2}(n+j_0)$; $\ell = \frac{1}{2}(n-j_0)$
- b) if $G < 0$ $k = \frac{1}{2}(n-j_0)$; $\ell = \frac{1}{2}(n+j_0)$
- c) if $G = 0$ $j_0 = 0$ and $k = \ell = \frac{1}{2}n$.

Thus the mirror representations have different signs of G .

The representations just derived are also representations of the Lorentz group; they are, however, not in this case Hermitian representations; we shall see that the Hermitian representations are of infinite dimension.

V. The Lorentz group*

The Lorentz transformations differ from the orthogonal transformations by the reality conditions which can be expressed by putting $x_4 = i \cdot x_0$ so that now x_0 is real instead of x_4 . The Lorentz transformations therefore leave the indefinite quadratic form $-x_0^2 + x_1^2 + x_2^2 + x_3^2$ of the real variables x_0, x_1, x_2, x_3 invariant. Correspondingly the Hermitian infinitesimal operators of the Lorentz group are given by the unchanged J_{23}, J_{31}, J_{12} and

$$J_{0k} = \frac{1}{i} \left(x_0 \frac{\partial}{\partial x_k} + x_k \frac{\partial}{\partial x_0} \right) = \frac{1}{i} J_{4k} (k = 1, 2, 3).$$

In the following we substitute therefore always $i \cdot \vec{N}$ for $\vec{N}$ and correspondingly $i \cdot G$ for the earlier invariant G , putting

$$\vec{N} = (J_{23}, J_{31}, J_{12})$$

$$\vec{N} = (J_{01}, J_{02}, J_{03}) .$$

* See also ⁽¹³⁻¹⁷⁾.

We therefore obtain now the new commutation rules

$$[N_1, N_2] = -i M_3, \dots$$

the other commutation rules remaining unchanged, and the invariants

$$F = \frac{1}{2}(\vec{M}^2 - \vec{N}^2)$$

$$G = \vec{M} \cdot \vec{N}.$$

We define now

$$\psi(j) = (j|N|j-1)(j-1|N|j)(2j-1)(2j+1)$$

instead of the earlier $\varphi(j)$. By changing the appropriate signs in the expression obtained before for $\varphi(j)$, we have now

$$\psi(j) = - (2F+1) - \frac{G^2}{j^2} + j^2.$$

We now consider the possible Hermitian representations. We shall find that they are of infinite dimension, for there is now no upper limit on the value of j due to the requirement that $\psi(j)$ takes non-negative values.

a) $G \neq 0$. There must clearly be a minimum non-zero value of j since we recall that $G = j(j+1)(j|N|j)$. We call this minimum j -value j_0 ; and we see from the definition of $\psi(j)$ that

$$\psi(j_0) = 0.$$

Hence

$$2F = j_0^2 - 1 - \frac{G^2}{j_0^2},$$

and

$$\psi(j) = j^2 - j_0^2 + G^2 \left(\frac{1}{j_0^2} - \frac{1}{j^2} \right)$$

which is non-negative for all $j \geq j_0$.

b) $G = 0, j \geq j_0 > 0$. We have again $\psi(j_0) = 0$, and hence

$$2F + 1 = j_0^2 ; \quad \psi(j) = j^2 - j_0^2 .$$

c) $G = 0, j \geq 0$. This case is particularly interesting as here the vanishing of $\psi(0)$ is not required but only $\psi(j) > 0$ for $j = 1, 2, \dots$ which means $F < 0$.

A different partition of these cases is obtained by putting

1) Principal series :

$$2F = j_0^2 - 1 - \nu^2 ; \quad G = j_0 \cdot \nu$$

$j_0 = 0, 1, 2, \dots$ or $1/2, 3/2, \dots$; ν real.

2) Complementary series or critical strip :

$$2F = -1 + \alpha^2 ; \quad G = j_0 = 0$$

$$0 < \alpha < 1 .$$

A part of case c) is contained in 1), another part in 2), while the cases a) and b) are contained in 1). It turns out that due to the infinite volume of the group space of the Lorentz group only the part 1) of the irreducible representations is needed to form a complete orthogonal set. The other part, 2), is not used in physics.

The inhomogeneous Lorentz group

We must add the infinitesimal operators of translations; these involve the commutation relations

$$\left. \begin{aligned} [p_\lambda, J_{\mu\nu}] &= i(\delta_{\lambda\nu} p_\mu - \delta_{\lambda\mu} p_\nu) \\ [p_\mu, p_\nu] &= 0 \end{aligned} \right\} \quad \lambda, \mu, \nu = 1, 2, 3, 4 .$$

Since we chose previously to put $ix_0 = x_4$ we now have $+ip_0 = p_4$. Thus we have the original operators with their original commutation relations from the 3-dimensional inhomogeneous group, namely J_1, J_2, J_3 and $p_1, p_2,$

p_3 plus the operators p_0 and N_1, N_2 , and N_3 . The new commutation relations involving p_0 and N_1, N_2, N_3 are thus

$$\begin{aligned} [p_1, N_1] &= +ip_0 \dots\dots & [p_0, N_1] &= +ip_1 \dots\dots \\ [p_2, N_1] &= [p_1, N_2] = \dots\dots = 0 & [p_0, N_1] &= \dots = 0 \end{aligned}$$

F and G are no longer invariants; we now have as invariants the square of the four-momentum

$$P = -p_\nu p_\nu = p_0^2 - \vec{p}^2,$$

and another invariant which we construct by first introducing the skew-symmetric tensor

$$v_{k\lambda\mu} = p_k J_{\lambda\mu} + p_\lambda J_{\mu k} + p_\mu J_{k\lambda} = J_{\lambda\mu} p_k + J_{\mu k} p_\lambda + J_{k\lambda} p_\mu.$$

We define the four-vector

$$(w_1, w_2, w_3, w_4) = -i(v_{234}, v_{314}, v_{124}, v_{321})$$

We have now the relations

$$[w_\lambda, J_{\mu\nu}] = i(\delta_{\lambda\nu} w_\mu - \delta_{\lambda\mu} w_\nu);$$

$$[w_\lambda, p_\mu] = 0;$$

$$[w_\mu, w_\nu] \neq 0 \quad \text{for } \mu \neq \nu;$$

$$w_\nu p_\nu = 0.$$

The second invariant is thus

$$W = w_\nu w_\nu = -\frac{1}{6} v_{k\lambda\mu} v_{k\lambda\mu} = \frac{1}{2}(p_\lambda p_\lambda)(J_{\mu\nu} J_{\mu\nu}) - J_{k\mu} J_{k\nu} p_\mu p_\nu.$$

We consider the significance of these two invariants in the rest

system. For the first we have $\vec{p} = 0$, $p_4 = im$. Hence $\underline{p = m^2}$.

For the second, we have, since $\vec{p} = 0$,

$$(w_1, w_2, w_3) = im(J_{23}, J_{31}, J_{12}) \text{ and } w_4 = 0.$$

Hence $\vec{w} = im \times (\text{angular momentum in rest system})$

$$\text{i.e. } -W = m^2 s(s+1) \text{ and } -\frac{W}{P} = s(s+1),$$

where s is the spin, provided that $m \neq 0$. There are clearly $2s+1$ independent states for a given momentum vector. The cases $s = 0, \frac{1}{2}$ and 1 correspond to the Klein-Gordon, Dirac and Proca equations, respectively.

If $P = 0$, there are two cases :

a) $\underline{W = 0}$. We have $w_\nu = \lambda p_\nu$ from $w_\nu p_\nu = 0$. Here λ is essentially the spin s . To show this we make use of the expressions in the 3-dimensional vector operators :

$$\vec{w} = + p_0 \vec{M} - (\vec{p} \times \vec{N}) = \lambda \vec{p},$$

$$w_0 = \vec{p} \cdot \vec{M} = \lambda p_0.$$

Since $p_0 = \pm \sqrt{p_1^2 + p_2^2 + p_3^2} = \pm p$, we have $\lambda = \pm \frac{\vec{p} \cdot \vec{M}}{p}$.

The representation theory of the 3-dimensional inhomogeneous group shows us that the magnitude of this quantity (referred to there as $j_0 = \pm \frac{\vec{J} \cdot \vec{p}}{p}$) is the minimum j value possible in the given representation; for a given momentum vector there are thus 2 independent states if $\lambda \neq 0$, corresponding to two different states of polarization, and one state when $\lambda = 0$. The cases $s = 0, \frac{1}{2}$ and 1 correspond to the scalar wave equation, the neutrino equation, and Maxwell's equations respectively.

In the particular case $W = P = 0$ considered here the representation of the 3-dimensional inhomogeneous rotation group can easily be supplemented to a representation of the inhomogeneous Lorentz group. (Here we are dealing with their infinitesimal transformations only).

From the commutation rules of the p_i with the N_k follows

$$[\vec{p}, (\vec{p} \cdot \vec{N})] = i p_0 \vec{p}$$

and with $p_0 = \pm p$,

$$[p, \vec{N}] = \pm i \vec{p}.$$

Hence for an arbitrary function $f(p)$ of the absolute value p

$$[f(p), \vec{N}] = \pm i f'(p) \cdot \vec{p}$$

and

$$\left[\frac{1}{p}, \vec{N} \right] = \pm i - \frac{1}{p^2} \cdot \vec{p}.$$

The unit vector $\vec{n} = \frac{\vec{p}}{p}$ has therefore the commutation relations

$$[n_1, N_1] = p_1 \left[\frac{1}{p}, N_1 \right] + \left[p_1, N_1 \right] \frac{1}{p} = \pm i (1 - n_1^2), \dots$$

$$[n_1, N_2] = p_1 \left[\frac{1}{p}, N_2 \right] = \mp i n_1 n_2, \dots$$

Therefore, with

$$N_{\mu} = \frac{1}{2} [(\vec{n} \times \vec{N}) + (\vec{N} \times \vec{n})] = (\vec{n} \times \vec{N}) \mp i,$$

$$[n_1, N_{\mu}] = 0,$$

but

$$[p, N_{\mu}] = \pm i p.$$

As

$$0 = (\vec{p} \times \vec{w}) = p^2 \{ \pm (\vec{n} \times \vec{M}) + \vec{N} - \vec{n} (N_{\mu} \pm i) \}$$

the quantities N_{μ} and p are the only ones with which the quantities $\vec{M}$ and $\vec{n}$ of the 3-dimensional inhomogeneous rotation group have to be supplemented in order to obtain the case $\vec{w} = P = 0$ of the inhomogeneous Lorentz group. As $\vec{M}$ and $\vec{n}$ commute with p and N_{μ} they retain their matrix representation of the 3-dimensional group.

The continuous variable p with the volume element $p \cdot dp$ (obtained from the invariant volume element $\frac{d^3 p}{(p_0)} = p \cdot dp \cdot dM$ by integration over the directions, $d\Omega$) has then to be incorporated (besides the integers, j, m) in the space of the representation of the inhomogeneous Lorentz group.

In agreement with the commutation rule already mentioned,

$$[p, N_\mu] = \pm i p$$

the quantity N_μ has then to be represented by the operator

$$(N_\mu)_{op} F(p) = \mp i \frac{\partial}{\partial p} (pF)$$

which is Hermitian with respect to the volume element $p \cdot dp$.

b) $W \neq 0$. There is a continuous spectrum of W instead of the spin; these representations do not seem to have any physical significance.

For application of this representation theory to physical problems see refs. (18-21).

References

- (1) Lie, S. and Scheffers, G. Vorlesungen über kontinuierliche Gruppen. Leipzig, 1893.
- (2) Racah, G. Group theory and spectroscopy. Princeton, 1951. (mimeographed notes.)
- (3) Pontrjagin, L. Topological groups. Princeton, University press, 1939. (see also pp. 181 and 334.)
- (4) Weyl, H. The structure and representation of continuous groups. Princeton, 1934-5. (mimeographed notes. See Chapter 2, Pt. 1.)
- (5,6) Weyl, H. Theorie der Darstellung kontinuierlicher halb-einfacher Gruppen durch lineare Transformationen. Mathematische Zeitschrift, 23, p. 271-309, 1925; 24, p. 328-95, 1926. (Reprinted in : Selecta Hermann Weyl, Basel, Birkhauser, 1956. p. 262.)
- (7) Boerner, H. Darstellung von Gruppen. Berlin, Springer, 1955.
- (8) Peter, F. and Weyl, H. Die Vollständigkeit der primitiven Darstellungen einer geschlossenen kontinuierlichen Gruppe. Mathematische Annalen, 27, p. 737-55, 1927.
- (9) Casimir, H. and van der Waerden, B.L. Algebraischer Beweis der vollständigen Reduzibilität der Darstellungen halbeinfacher Liescher Gruppen. Mathematische Annalen, 111, 1-12, 1935.
- (10) Brauer, R. Eine Bedingung für vollständige Reduzibilität von Darstellungen gewöhnlicher und infinitesimaler Gruppen. Mathematische Zeitschrift, 41, p. 330-9, 1936.
- (11) Weyl, H. The classical groups, their invariants and representations. Princeton, University press, 1939.
- (12) Weyl, H. Gruppentheorie und Quantenmechanik. 2e Aufl. Leipzig, Hirzel, 1931.
- (13) Wigner, E. On unitary representations of the inhomogeneous Lorentz group. Annals of mathematics, 40, p. 149-204, 1939.
- (14) Harish-Chandra. Infinite irreducible representations of the Lorentz group. Royal Society. Proceedings A, 189, p. 272-401, 1947.
- (15) Bargmann, V. and Wigner, E.P. Group theoretical discussion of relativistic wave equations. National Academy of Sciences, Washington. Proceedings, 34, p. 211-23, 1948.
- (16) Bargmann, V. Irreducible unitary representations of the Lorentz group. Annals of Mathematics, 48, p. 568-640, 1947.

- (17) Gelf, I. and Neumark, M. Unitary representation of the Lorentz group. Journal of physics, USSR, 10, p. 93-4, 1946.
- (18) Pauli, W. Über das Wasserstoffspektrum vom Standpunkt der neuen Quantenmechanik. Zeitschrift für Physik, 36, p. 336-63, 1926.
- (19) Hulthén, L. Über die quantenmechanische Herleitung der Balmerterme. Zeitschrift für Physik, 86, p. 21-3, 1933.
- (20) Fock, V. Zur Theorie des Wasserstoffatoms. Zeitschrift für Physik, 98, p. 145-54, 1936.
- (21) Bargmann, V. Zur Theorie des Wasserstoffatoms. Zeitschrift für Physik, 99, p. 576-82, 1936.