Scanned with permission from the publisher:

Fourth-order magnetic moment of the electron / Petermann, Andreas

CERN-57-53

Published in: Helv. Phys. Acta: 30 (1957) 407-408

Date: Thu, 27 Jan 2005

From: Etienne Liv <etienne@birkhauser.ch>

To: vigen@mail.cern.ch

Subject: WG: Request for permission to reproduce published material

Dear Mr. Vigen

Thank you for your permission requests from "Helvetica Physica Acta" and "Experientia Supplementa" which Dr. Fund forwarded to me.

We herewith grant you the desired permission under condition that full reference to the original source of publication is made.

Please be informed that Helvetica Physica Acta is not published anymore and that Experientia was renamed to Cellular and Molecular Life Sciences.

Sincerely,

Liv Etienne
Rights & Licences

A. PETERMANN

HELVETICA PHYSICA ACTA

1957. Vol. 30. Fasc. 5, p. 407-408

Fourth order magnetic moment of the electronby **A. Petermann.**

CERN. Theoretical Study Division. Institute for theoretical Physics. Copenhagen.

(17. VIII. 1957.)

In connection with the upper and lower bounds analysis done by the author¹⁾, which indicated a clear discrepancy with the Karplus and Kroll's result for the 4th order magnetic moment²⁾, we have performed an analytic evaluation of the five independent diagrams contributing to this moment in fourth order*). The results are the following:

$$\mu_{\text{I}} = \frac{\alpha^2}{\pi^2} \left(\frac{1}{6} + \frac{13}{36} \pi^2 + \frac{5}{4} \zeta(3) - \frac{5}{6} \pi^2 \text{Log } 2 \right) = -0.467 \frac{\alpha^2}{\pi^2}. \quad (1)$$

$$\mu_{\text{II}_a} = \frac{\alpha^2}{\pi^2} \left(\frac{11}{48} + \frac{\pi^2}{18} \right) = 0.778 \frac{\alpha^2}{\pi^2}. \quad (2)$$

$$\begin{aligned} \mu_{\text{II}_c} = \frac{\alpha^2}{\pi^2} \left(-\frac{67}{24} + \frac{\pi^2}{18} - \frac{1}{2} \zeta(3) + \frac{1}{3} \pi^2 \text{Log } 2 - \frac{1}{2} \text{Log } \frac{\lambda^2}{m^2} \right) = \\ -0.564 \frac{\alpha^2}{\pi^2} - \frac{1}{2} \frac{\alpha^2}{\pi^2} \text{Log } \frac{\lambda^2}{m^2}. \end{aligned} \quad (3)$$

$$\mu_{\text{II}_d} = \frac{\alpha^2}{\pi^2} \left(\frac{11}{24} - \frac{\pi^2}{18} + \frac{1}{2} \text{Log } \frac{\lambda^2}{m^2} \right) = -0.090 \frac{\alpha^2}{\pi^2} + \frac{1}{2} \frac{\alpha^2}{\pi^2} \text{Log } \frac{\lambda^2}{m^2}. \quad (4)$$

$$\mu_{\text{II}_e} = \frac{\alpha^2}{\pi^2} \left(\frac{119}{36} - \frac{\pi^2}{3} \right) = 0.016 \frac{\alpha^2}{\pi^2}. \quad (5)$$

$$\mu_{\text{total}}^{(4)} = \frac{\alpha^2}{\pi^2} \left(\frac{197}{144} + \frac{\pi^2}{12} + \frac{3}{4} \zeta(3) - \frac{1}{2} \pi^2 \text{Log } 2 \right) = -0.328 \frac{\alpha^2}{\pi^2}. \quad (6)$$

Compared with the values given in their original paper by KARPLUS and KROLL, one can see that two terms were in error: μ_{I} differs by

$$\frac{\alpha^2}{\pi^2} \frac{1}{32} = 0.031 \frac{\alpha^2}{\pi^2};$$

$$\mu_{\text{II}_c} \text{ by } \frac{\alpha^2}{\pi^2} \left(\frac{32}{3} - \frac{61}{8} \pi^2 + \frac{17}{2} \pi^2 \text{Log } 2 - \frac{109}{4} \zeta(3) \right) = 2.614 \frac{\alpha^2}{\pi^2}.$$

The three other terms check. The error in μ_{I} remained of course undetected in the upper and lower bound analysis owing to its small-

*) The terminology of ref. 2 is used throughout this paper.

ness. But the large discrepancy in μ_{IIc} was that pin-pointed out in the previous paper.

A summary of the most important electromagnetic observables, the theoretical values of which are modified by the new value of the magnetic moment, is now given:

$$\text{Moment of the electron: } \frac{\mu_e}{\mu_0} = 1.0011596 = 1 + \frac{\alpha}{2\pi} - 0.328 \frac{\alpha^2}{\pi^2}.$$

FRANKEN and LIEBES' value for it: $\mu_e/\mu_0 = 1.001167 \pm 0.000005^*)$.
 g -factor of the μ -meson (electromagnetic):

$$2 (1.0011654) = 2 \left(1 + \frac{\alpha}{2\pi} + 0.75 \frac{\alpha^2}{\pi^2} \right).$$

Last Lederman's value: $2 (1.0021 \pm 0.0008)^*)$.

$2^2 S_{1/2} - 2^2 P_{1/2}$ (Hydrogen): (1057.94 ± 0.15) Mc/s; observed:
 (1057.77 ± 0.10) Mc/s.

$2^2 S_{1/2} - 2^2 P_{1/2}$ (Deuterium): (1059.22 ± 0.15) Mc/s; observed:
 (1059.00 ± 0.10) Mc/s.

Fine structure constant: $1/\alpha = 137.0384$; (previously: 137.0365).

The new fourth order correction given here is in agreement with:

a) The upper and lower bounds given by the author¹⁾.

b) A calculation using a different method, performed by C. SOMMERFIELD³⁾.

c) A recalculation done by N. M. KROLL and collaborators^{*)}.

The author thanks Prof. NIELS BOHR for the hospitality at the Institute.

References.

- ¹⁾ A. PETERMANN. Nuclear Physics **3**, 689 (1957) and Nuclear Physics in the press.
- ²⁾ R. KARPLUS and N. M. KROLL, Phys. Rev. **77**, 536 (1950).
- ³⁾ C. SOMMERFIELD, Phys. Rev. In the press.

^{*)} Private Communication.