The 8th workshop in a serie

Excited QCD 2016

Costa da Caparica, Lisbon, Portugal
6-12 March 2016

QCD at high temperatures & finite densities

- heavy-ion collisions, jets, diffraction, hadronisation
- quark-gluon plasma
- holography, colour-glass condensate
- compact stars, applications to astrophysics

QCD at low energies

- excited hadrons
- new resonances
- glueballs, multiquarks

Searches for Supersymmetry and Exotic phenomena with the ATLAS Detector

N. Chr. Benekos (NTUA)
Representing the ATLAS collaboration

More information
<https://indico.cern.ch/event/453434/>
excitedqcd@th.physik.uni-frankfurt.de

Organising Committee
 Pedro Bicudo, IST, Lisbon
 Francesco Giacosa, UJK, Kielce
 Jelena Jovicevic, TRIUMF
 Robert Kaminski, IJF PAN, Krakow
 Marina Marinkovic, CERN

- No other clear indications of new physics beyond the SM
 - Need to cast wide a net as possible even with some theoretical guidance
 - Need to cover broad phase spaces: many different detector signatures, large range of masses, large span in production rates
- Run 1 results have been extremely useful to kill some models or narrow down their allowed phase spaces

(for illustration only)

ATLAS Exotics Searches* - 95% CL Exclusion
Status: July 2015

ATLAS Preliminary
 $\sqrt{s} = 7, 8 \text{ TeV}$
 $\int \mathcal{L} dt = (4.7 - 20.3) \text{ fb}^{-1}$

*Only a selection of the available mass limits on new states or phenomena is shown.

2015 dataset at 13 TeV :

- 3.9 fb⁻¹ recorded
- 3.3 fb⁻¹ "all good" for BSM searches
- $L_{inst} = 5.1 \times 10^{33} \text{ cm}^2 \text{ s}^{-1}$

New 4th innermost layer of pixel in ATLAS : the IBL :

at 3.3 cm vs 5.05 cm for the 2nd layer

- improve *b*-tagging performance which is important for several BSM searches

less pileup than in Run-1 : $\langle \mu \rangle = 13$ vs 21

ATLAS performance close to or exceeding design specs in all compartments

Strategy:

- Define selection based on signal signatures and background (bkg) kinematics
- Compare data to Standard Model bkg (Monte Carlo (MC) + data driven) and MC signal predictions → data consistent with bkg+signal would be evidence for new physics

No evidence for new physics:

- Limits typically set on cross-section x branching fraction ($\sigma \times \text{BR}$)
- Comparisons provided for specific models, but usually possible for reader to constrain additional models

Global symmetry between fermions & bosons:
all SM particles have SUSY partners

$$Q|\text{fermion}\rangle = |\text{boson}\rangle$$

$$Q|\text{boson}\rangle = |\text{fermion}\rangle$$

$$s_{\text{SUSY}} = s_{\text{SM}}^{-1/2}$$

$$R = (-1)^{2s}(-1)^{3B}(-1)^L$$

Why SUSY?

- Solves the hierarchy problem
- Provides the dark matter candidate: if R-parity is conserved, lightest SUSY particle (LSP, neutralino) is stable
- Extends the Poincare group, provides unification with gravity
- Required for the string theory
- ... and more others ...

Electroweak

Strong

- Inclusive searches for $\tilde{q}$ and $\tilde{g}$
- Large cross-section
- Rich final states in case of $\tilde{g}-\tilde{g}$ production

- Dedicated searches for $\tilde{t}_1$ and $\tilde{b}_1$
- final state similar to the SM bkg

- Searches for $\tilde{\chi}_1^0$ and $\tilde{\chi}_1^\pm$
- low cross-section
- multi-leptons final states with low SM bkg

+ dedicated searches for long-lived particles and RPV SUSY

From <http://inspirehep.net/record/1326406>

• large increase of SUSY cross-section from 8 to 13 TeV :

• $\sigma(\tilde{g}\tilde{g}) \times 30$ for $m(\tilde{g}) = 1.4$ TeV

• $\sigma(t\tilde{t}) \times 8$ for $m(\tilde{t}) = 700$ GeV

• $\sigma(\tilde{\chi}\tilde{\chi}) \times 4$ for $m(\tilde{\chi}) = 500$ GeV

□ focus on gluino and third generation squarks searches with 2015 data, with a discovery potential beyond Run-1 limits even with 3 fb^{-1} of 13 TeV data

□ discovery potential of EW SUSY beyond Run-1 limits will be reached with 2016 data

Electrons :

- ID efficiencies measured in 2015 data to derive MC scale factors
- calibration based on run 1 with MC extrapolations

Muons :

- ID and reco efficiencies measured in 2015 data
- energy scale and resolution taken also from 2015 data

excellent agreement between data and MC

- Electrons and Muons calibrated with Z events
- Differences in efficiency and energy scale corrected

jets :

- Use Run-1 knowledge to extrapolate systematic uncertainties for run 2
- JES studies in 2015 data with photon-jet and multi-jets balance

b -tagging :

- improvements in algorithms and with new IBL
- b -tagging efficiency increased by 10% for the same light jet rejection
- MC calibration validated in data with $t\bar{t}$ events

- new E_T calculation with a track-based soft term (TST) combined with calorimeter-based measurements for hard objects
- less sensitivity to pile-up than with calorimeter-based soft term (CST)
- systematic uncertainties based on MC, and validated in data

Results and interpretations for 7 analyses

$1-l + \text{jets} + E_T$ analysis [ATLAS-CONF-2015-076](#)

$0-l + 2-6 \text{ jets} + E_T$ analysis [ATLAS-CONF-2015-062](#)

$0-l + 7-10 \text{ jets} + E_T$ analysis [ATLAS-CONF-2015-077](#)

$0-1-l + 3-4 b\text{-jets} + E_T$ analysis [ATLAS-CONF-2015-067](#)

$2-l$ same-sign / $3-l$ analysis [ATLAS-CONF-2015-078](#)

$Z \rightarrow ll + E_T$ analysis [ATLAS-CONF-2015-082](#)

$0-l + 2 b\text{-jets} + E_T$ analysis [ATLAS-CONF-2015-066](#)

Search for gluino pair production in final states with 1 lepton, jets and E_T^{miss}

ATLAS-CONF-2015-076

- 4 hard- ℓ SRs:
 - $p_T > 35$ GeV lepton
 - No additional leptons with $p_T > 10$ GeV
 - 4-6 jets
- 2 soft- ℓ SRs:
 - $p_T > 7(6)$ GeV for $e()$ and $p_T < 35$ GeV
 - No additional $e(\mu)$ with $p_T > 7(6)$ GeV
 - 2 or 5 jets

□ Main backgrounds : $t\bar{t}$ and W +jets events

□ Suppressed with cuts on transverse mass

✧ Estimated by normalising the MC in CRs

Ex : soft $e+2j$: split CR with low m_T and E_T^{miss} :

• $t\bar{t}$ CR : ≥ 1 b -jet

• W +jets CR : $= 0$ b -jet

□ Subdominant background from MC

- Overall good agreement between expected background and observation
- Largest deviation of 2σ observed in the SR with 1 hard lepton + 6 jets :

- 1 electron : $\text{exp} = 1.9 \pm 0.6$, $\text{obs} = 2$
- 1 muon : $\text{exp} = 2.5 \pm 0.8$, $\text{obs} = 8$

- Search for gluino and squark pair production in final states with 0 lepton, jets and E_T

[ATLAS-CONF-2015-062](#)

- Define 7 signal regions to probe all scenarios
- From 2 to ≥ 6 jets to target different models
- different m_{eff} cuts to probe different mass splittings

7 SRs with 2-6 jets & different cuts

- Targeting different models

Veto leptons with $p_T > 10$ GeV

4 CRs for each SR, to obtain background from

- Multi-jet
- $Z(\rightarrow \nu\nu)$ +jets
- $W(\rightarrow \ell\nu)$ +jets
- $t\bar{t}$, single- t

Background from MC

- Di-boson

Requirement	Signal Region						
	2jl	2jm	2jt	4jt	5j	6jm	6jt
$E_{\text{T}}^{\text{miss}} \text{ [GeV]} >$	200						
$p_{\text{T}}(j_1) \text{ [GeV]} >$	200	300	200				
$p_{\text{T}}(j_2) \text{ [GeV]} >$	200	50	200	100			
$p_{\text{T}}(j_3) \text{ [GeV]} >$	–			100			
$p_{\text{T}}(j_4) \text{ [GeV]} >$	–			100			
$p_{\text{T}}(j_5) \text{ [GeV]} >$	–				100		
$p_{\text{T}}(j_6) \text{ [GeV]} >$	–					100	
$\Delta\phi(\text{jet}_{1,2,(3)}, \mathbf{E}_{\text{T}}^{\text{miss}})_{\text{min}} >$	0.8	0.4	0.8	0.4			
$\Delta\phi(\text{jet}_{i>3}, \mathbf{E}_{\text{T}}^{\text{miss}})_{\text{min}} >$	–			0.2			
$E_{\text{T}}^{\text{miss}}/\sqrt{H_{\text{T}}} \text{ [GeV}^{1/2}] >$	15		20	–			
Aplanarity $>$	–			0.04			
$E_{\text{T}}^{\text{miss}}/m_{\text{eff}}(N_{\text{j}}) >$	–			0.2	0.25		0.2
$m_{\text{eff}}(\text{incl.}) \text{ [GeV]} >$	1200	1600	2000	2200	1600	1600	2000

□ Main backgrounds : $(Z \rightarrow \nu\nu) + \text{jets}$, $W + \text{jets}$, $t\bar{t}$ and single top events with hadronic taus

➤ normalise the MC in 3 dedicated CRs with same jets p_T and m_{eff} cuts as the SRs :

$(Z \rightarrow \nu\nu) + \text{jets}$ CR

- $\gamma + \text{jets}$
- $p_T(\gamma) > 130 \text{ GeV}$
- treat the photon as ν in E_T

$W + \text{jets}$ CR

- 1-lepton treated as a jet
- $30 < m_T < 100 \text{ GeV}$
- b -jet veto

top CR

- 1-lepton treated as a jet
- $30 < m_T < 100 \text{ GeV}$
- $\geq b$ -jet

□ sub-dominant diboson contribution estimated from MC

□ residual contribution from multi-jets events with fake E_T estimated in a CR with reverted cuts on $\Delta\phi_{\text{min}}(E_T, \text{jets})$ and E_T/m_{eff}

Expected and observed event count in each SR:

interpretation with gluinos
 exclude gluino masses up to $\sim 1.5\text{ TeV}$

Search for gluino pair production with complex decay chains

- 6 SRs with 7-8 $p_T > 80$ GeV jets, incl. 0-2 b -jets
(trigger : 5 jets with $p_T > 70$ GeV)
- 9 SRs with 8-10 $p_T > 50$ GeV jets , incl. 0-2 b -jets
(trigger : 6 jets with $p_T > 45$ GeV)
- No leptons with $p_T > 10$ GeV

- $t\bar{t}$, V+jets background obtained from CRs containing a lepton with $p_T > 20$ GeV
- Multijet background from CRs with 1 jet less.
- Utilize near invariance of $E_T^{miss}/\sqrt{H_T}$ wrt. N_{jets} (shape is almost invariant wrt N_{jets}) when MET originates from calorimeter mismeasurement

- Checked in VRs

[ATLAS-CONF-2015-077](#)

- SR: $\frac{E_T^{miss}}{\sqrt{H_T}} > 4 \sqrt{\text{GeV}}$

- Distribution normalization:

$$\frac{E_T^{miss}}{\sqrt{H_T}} < 1.5 \sqrt{\text{GeV}}$$

Overall good agreement
between expected background
and data in the 15 SR

Expected and observed event count in each SR:

Signal region	Fitted background			Obs events
	Multijet	Leptonic	Total	
8j50	109.3 $\pm$ 6.8	79 $\pm$ 25	189 $\pm$ 26	157
8j50-1b	76.7 $\pm$ 2.6	61 $\pm$ 21	138 $\pm$ 21	97
8j50-2b	33.8 $\pm$ 2.1	33 $\pm$ 13	67 $\pm$ 13	39
9j50	16.8 $\pm$ 1.2	12.8 $\pm$ 5.4	29.6 $\pm$ 5.6	29
9j50-1b	13.5 $\pm$ 1.9	10.2 $\pm$ 4.9	23.8 $\pm$ 5.3	21
9j50-2b	6.4 $\pm$ 1.6	5.8 $\pm$ 3.3	12.1 $\pm$ 3.6	9
10j50	2.61 $\pm$ 0.60	1.99 $\pm$ 0.62	4.60 $\pm$ 0.86	6
10j50-1b	2.42 $\pm$ 0.62	1.44 $\pm$ 0.49	3.86 $\pm$ 0.79	3
10j50-2b	1.40 $\pm$ 0.87	0.83 $\pm$ 0.37	2.23 $\pm$ 0.94	1
7j80	40.0 $\pm$ 5.1	30 $\pm$ 12	70 $\pm$ 13	70
7j80-1b	29.1 $\pm$ 3.2	20.8 $\pm$ 10	50 $\pm$ 10	42
7j80-2b	11.5 $\pm$ 1.6	11.0 $\pm$ 4.9	22.5 $\pm$ 5.2	19
8j80	4.5 $\pm$ 1.9	4.9 $\pm$ 2.1	9.3 $\pm$ 2.8	8
8j80-1b	3.9 $\pm$ 1.5	3.8 $\pm$ 2.1	7.6 $\pm$ 2.6	4
8j80-2b	1.72 $\pm$ 0.92	2.3 $\pm$ 1.1	4.1 $\pm$ 1.4	2

- for $m(\tilde{\chi}_1^\pm) < 500$ GeV and $m(\tilde{g}) > 1200$ GeV :
→ decay chain above is dominant
- for $m(\tilde{\chi}_1^\pm) > 500$ GeV and $m(\tilde{g}) < 1200$ GeV :
→ $\tilde{g} \rightarrow qq\tilde{\chi}_1^0$ becomes dominant

$$\bullet \quad m(\tilde{\chi}_1^\pm) = \frac{m(\tilde{g}) + m(\tilde{\chi}_1^0)}{2}$$

$$\bullet \quad m(\tilde{\chi}_2^0) = \frac{m(\tilde{\chi}_1^\pm) + m(\tilde{\chi}_1^0)}{2}$$

- ☐ Gluino-mediated stop pair production : [ATLAS-CONF-2015-067](#)
- ☐ very rich final state
- 2 channels with $= 0$ or ≥ 1 lepton
- require high jets and b -jets multiplicities
- use jet substructure techniques to probe boosted topologies with large- R jets

- ☐ Gluino-mediated sbottom pair production :
- ☐ 0-lepton channel only
- ☐ require high b -jets multiplicity
- ☐ Main discriminant variables to reject backgrounds are E_T and m_{eff}

- 8 SRs:
- ≥ 3 b -jets
 - 0 or 1 lepton
 - 0 or ≥ 1 top ($R = 1$ jet)

MET distributions in some SRs:

Results of background-only fit in VRs

systematics dominated by $t\bar{t}$ modeling and b -tagging unc.

No significant excess in data
 exclude gluino masses up to ~ 1.7 TeV

[ATLAS-CONF-2015-067](#)

No significant excess in data
 exclude gluino masses up to ~ 1.8 TeV

very low SM bkg, broad sensitivity to many SUSY scenarios with leptons in the decay chain :

Signal region	$N_{\text{lept}}^{\text{signal}}$	N_{bjets}^{20}	N_{jets}^{50}	E_T^{miss} [GeV]	m_{eff} [GeV]
SR0b3j	≥ 3	$= 0$	≥ 3	> 200	> 550
SR0b5j	≥ 2	$= 0$	≥ 5	> 125	> 650
SR1b	≥ 2	≥ 1	≥ 4	> 150	> 550
SR3b	≥ 2	≥ 3	-	> 125	> 650

- Events selected with a logical OR combination of E_T and di-lepton triggers
- 4 signal regions defined according to the leptons, b -jets, jets, E_T and m_{eff} :

3 main sources of backgrounds :

fake+non-prompt leptons electrons charge flip

- from jet misID, HF decay, or γ conversion
- data-driven estimate with the loose to tight matrix method

- from hard brem γ conversion
- charge flip rate from data in $Z/\gamma^* \rightarrow e^+ e^-$ events
apply this rate in 2L OS SR to get the 2L SS bkg in corresponding SR

irreducible bkg

- $t\bar{t} + V, VV, VVV$ events with 2 SS or 3 real leptons :
- estimated from MC and Validated with data in VRs
- largest deviation of 1.3σ in the $t\bar{t} + V$ VR

ATLAS-CONF-2015-078

ATLAS Collaboration

	SR0b3j	SR0b5j	SR1b	SR3b
Observed events	3	3	7	1
Total bkg events	2.4 ± 0.7	0.98 ± 0.32	4.3 ± 1.0	0.78 ± 0.24
$p(s=0)$	0.33	0.06	0.12	0.36
Fake/non-prompt leptons	< 0.2	$0.04^{+0.17}_{-0.04}$	0.8 ± 0.8	0.12 ± 0.16
Charge flip	–	0.02 ± 0.01	0.60 ± 0.12	0.19 ± 0.06
$t\bar{t}W, t\bar{t}Z$	0.13 ± 0.06	0.11 ± 0.06	2.0 ± 0.7	0.21 ± 0.09
WZ	1.5 ± 0.5	0.61 ± 0.25	0.17 ± 0.09	< 0.02
$W^\pm W^\pm jj$	–	0.11 ± 0.05	0.03 ± 0.01	< 0.01
ZZ	0.6 ± 0.4	< 0.14	< 0.03	< 0.03
Triboson	0.09 ± 0.05	0.02 ± 0.01	0.02 ± 0.01	< 0.01
Rare	0.05 ± 0.04	0.05 ± 0.04	0.7 ± 0.4	0.26 ± 0.14

ATLAS-CONF-2015-078

Search for gluino (or squark) pair production with Z in the decay chains

Dominant backgrounds:

- tt, t :
 - Using CR with different-flavor leptons
- $Z/\gamma^* + \text{jets}$:
 - Using CR with $\gamma + \text{jets}$

E_T^{miss} [GeV]	H_T [GeV]	n_{jets}	$m_{\ell\ell}$ [GeV]	SF/DF	$\Delta\phi(\text{jet}_{12}, p_T^{\text{miss}})$
> 225	> 600	≥ 2	$81 < m_{\ell\ell} < 101$	SF	> 0.4

- Check excess observed in the ATLAS
- Run-1 8 TeV analysis :
- 3 (1.7) σ excess in the $ee (\mu\mu)$ channel

➤ reproduce the Run-1 signal region :

- 2 $e (e^+ e^- \text{ or } \mu^+ \mu^-)$ with $p_T > 50, 25$ GeV and $81 < m(ee) < 101$ GeV
- ≥ 2 jets with $\Delta\phi_{\min}(E_T, \text{jets}) > 0.4$
- $E_T > 225$ GeV and $H_T > 600$ GeV

[ATLAS-CONF-2015-082](https://atlas.conf.cern.ch/2015/082)

10.3 ± 2.3 events expected
 21 observed ($10 ee + 11 \mu\mu$)
 excess of 2.2σ
 CMS result : $\text{Exp} = 12^{+4.0}_{-2.8}$, $\text{obs} = 12$

[ATLAS-CONF-2015-082](#)

	SRZ
Observed events	21
Total expected background events	10.3 ± 2.3
Flavour symmetric ($t\bar{t}$, Wt , WW and $Z \rightarrow \tau\tau$) events	5.1 ± 2.0
WZ/ZZ events	2.9 ± 0.8
$Z/\gamma^* + \text{jets}$ events	1.9 ± 0.8
Rare top events	0.4 ± 0.1
p -value	0.013
Significance	2.2
Observed (Expected) S^{95}	20.0 ($10.2^{+4.4}_{-3.0}$)

Search for direct sbottom squark production

[ATLAS-CONF-2015-066](#)

4 SRs:

- 3 “SRA” target pair production
- SRB target pair production with ISR jet

[ATLAS-CONF-2015-066](#)

Background:

- tt and + heavy flavor
 - Studied with CRs containing leptons

Variable	SRA	SRB
Event cleaning	Common to all SR	
Lepton veto	No e/μ with $p_T > 10$ GeV after overlap removal	
E_T^{miss}	> 250 GeV	> 400 GeV
Leading jet $p_T(j_1)$	> 130 GeV	> 300 GeV
2nd jet $p_T(j_2)$	> 50 GeV	> 50 GeV
Fourth jet $p_T(j_4)$	vetoed if > 50 GeV	
$\Delta\phi_{\text{min}}^j$	> 0.4	> 0.4
$\Delta\phi(j_1, E_T^{\text{miss}})$	-	> 2.5
b -tagging	j_1 and j_2	j_2 and (j_3 or j_4)
$E_T^{\text{miss}}/m_{\text{eff}}$	> 0.25	> 0.25
m_{CT}	$> 250, 350, 450$ GeV	-
m_{bb}	> 200 GeV	-

Main discriminator: contraverse mass,

$$m_{\text{CT}}^2(v_1, v_2) = [E_T(v_1) + E_T(v_2)]^2 - [\mathbf{p}_T(v_1) - \mathbf{p}_T(v_2)]^2$$

Can be used to measure m_b

Signal region channels	SRA250	SRA350	SRA450	SRB
Observed events	22	6	1	5
Fitted bkg events	40 ± 8	9.5 ± 2.6	2.2 ± 0.6	13.1 ± 3.2
Fitted $t\bar{t}$ events	0.9 ± 0.4	0.37 ± 0.16	0.06 ± 0.03	5.9 ± 2.4
Fitted single top events	2.1 ± 1.3	0.54 ± 0.37	0.15 ± 0.10	1.2 ± 0.8
Fitted W +jets events	6.3 ± 2.4	1.3 ± 0.6	0.41 ± 0.23	1.2 ± 0.6
Fitted Z +jets events	30 ± 7	7.1 ± 2.4	1.5 ± 0.5	3.3 ± 1.4
(Alt. method Z +jets events)	(33 ± 7)	(7.2 ± 1.9)	(2.7 ± 0.9)	
Fitted “Other” events	0.7 ± 0.6	0.1 ± 0.1	0.02 ± 0.02	1.4 ± 0.4

[ATLAS-CONF-2015-066](#)

Leptonic final states:

- ❑ e/μ and missing energy [ATLAS-CONF-2015-063](#)
- ❑ e^+e^- , $\mu^+\mu^-$ [ATLAS-CONF-2015-070](#)
- ❑ $e^+\mu^-$ [ATLAS-CONF-2015-072](#)

Di-jet/photon mass spectra:

- ❑ jet-jet [arXiv:1512.01530v2](#)
- ❑ γ -jet [arXiv:1512.05910](#)
- ❑ $\gamma\gamma$ [ATLAS-CONF-2015-081](#)

- ❑ “Exotics” means direct searches for particles/phenomena:
 - beyond the Standard Model
 - not supersymmetry or BSM Higgs

- ❑ Leaves a huge range of hypotheses that explain one or more of the mysteries in the SM

- ❑ These searches benefit greatly from the increased LHC energy
- ❑ with ~ 3 fb $^{-1}$ at 13 TeV, we have already exceeded Run-1 sensitivity in many cases

- Reconstruct exactly one isolated lepton with $p_T > 65\text{ GeV}$.
- The missing transverse energy in the event must exceed 55 GeV.
- Search in the transverse mass:

$$m_T = \sqrt{2p_T E_T^{\text{miss}}(1 - \cos \phi_{\ell\nu})},$$

- Test for heavy spin-1 W' bosons.
Event with one electron

ATLAS-CONF-2015-063

- Select a pair of leptons
 - well isolated
 - matched to the primary vertex
 - E_T or $p_T > 30$ GeV
- Main background from Drell-Yan Z/γ production.
- Test di-lepton masses up to 5 TeV
- Search for heavy Z' bosons or contact interactions.

New physics could alter the ee and $\mu\mu$ mass distributions in two ways

ATLAS-CONF-2015-070

a new resonance (Z') would create a bump

non-resonant effects could change the shape

p values:

Z' and
contact
interaction
limits:

Highly sensitive channel for new neutral resonance with lepton-flavor violating decays

- Require different flavor leptons
- Main background from $t\bar{t} \rightarrow W^+W^-$
- Test for LFV in Z' models and quantum black holes in RS and ADD models.
- No excess observed.

di-jet, $\gamma\gamma$, γ -jet mass spectra

Tests for q^* , quantum
black holes, Z'/W' ,
heavy Higgs models

- ❑ Sensitive to new particles in s- and t-channels, respectively
 - any new particle produced at the LHC must couple to quarks/gluons
 - dijet search is sensitive to all such particles

[doi:10.1016/j.physletb.2016.01.032](https://doi.org/10.1016/j.physletb.2016.01.032)

- Jet selections:
 - $p_T(j_1) > 440 \text{ GeV}$, $p_T(j_2) > 50 \text{ GeV}$
 - compatibility with primary vertex
- Multi-jet background reduced with requirement on rapidity difference $|y^*| = |(y_1 - y_2)/2| < 0.6$
- Di-jet mass resolution $\sim 2\%$ over full mass range
- Background modeled with a power law function.
- Look for a narrow resonance or an excess.

Limits on benchmark
Gaussian signals:

- The $m(jj)$ region above 5.4 TeV was previously unexplored.
- Large improvements in the exclusion limits over Run I.

Model	95% CL Exclusion limit		
	Run 1 Observed	Observed 13 TeV	Expected 13 TeV
Quantum black holes, ADD (BLACKMAX generator)	5.6 TeV	8.1 TeV	8.1 TeV
Quantum black holes, ADD (QBH generator)	5.7 TeV	8.3 TeV	8.3 TeV
Quantum black holes, RS (QBH generator)	–	5.3 TeV	5.1 TeV
Excited quark	4.1 TeV	5.2 TeV	4.9 TeV
W'	2.5 TeV	2.6 TeV	2.6 TeV
Contact interactions ($\eta_{LL} = +1$)	8.1 TeV	12.0 TeV	12.0 TeV
Contact interactions ($\eta_{LL} = -1$)	12.0 TeV	17.5 TeV	18.1 TeV

- Test quantum black holes (QBH) and excited quark models
- Events are selected with a well isolated photon. Both the photon and jet must have $E_T > 150$ GeV
- γ +jet mass resolution is $\sim 2.5\%$ up to 6 TeV
- Exclusion limits surpass Run I analysis.

- Select two isolated photons
- Photon energy calibrated using $Z \rightarrow ee$ events
- Energy selections relative to the signal mass
- Signal efficiencies range between 20-45% depending on the mass and production mechanism
- Main background is from non-resonant di-photon (90%) and jet (10%) production

- Run II shows some excess at $m(\gamma\gamma) \sim 750$ GeV
- Significance: local $\sim 3.6\sigma$, global ~ 2.0 sigma, estimated using a narrow width signal model.
- Using width $\Gamma \sim 45$ GeV gives larger significances: 3.9σ local (**2.3σ global**)
- Run I analysis did not see excess, but is still compatible within 2.3σ of the current result assuming gluon fusion production.

ATLAS-CONF-2015-081

Exclusion Limits

Local p-value in NWA

- successful commissioning of the ATLAS detector upgrade for Run-2
- early SUSY searches performed with 3.3 fb^{-1} at 13 TeV

□ exclude $m(\tilde{g})$ up to 1.8 TeV

□ exclude $m(\tilde{q}_{1,2})$ up to $\sim 1 \text{ TeV}$

□ exclude $m(b_1)$ up to $\sim 840 \text{ GeV}$

ATL-PHYS-PUB-2015-005

$\Rightarrow$ significantly improves exclusion limits from Run-1

□ Excess of 2.2σ in the $Z+E_T$ analysis, after 3σ in Run-1

- ❑ ATLAS is pursuing a broad search for exotic phenomena
- ❑ Some searches have already been updated using the 2015 data
 - thanks to the increased LHC energy, Run 1 sensitivity to many signals has already been exceeded with $\sim 3 \text{ fb}^{-1}$
- ❑ Unfortunately, few surprises so far but there is the diphoton excess
- ❑
- ❑ Still to come:
 - updated searches for many other new phenomena
 - including follow up on some of the most interesting results from Run 1

BACK UP SLIDES

- **Jets:**
 - Reconstructed from calorimeter energy clusters using the anti- k_T algorithm with radius parameter $R = 0.4$
 - Jets are reclustered with $R = 1$ to search for boosted top quarks
 - Corrected for avg. energy deposition from pile-up (= multiple collisions, averaging 14 in 2015)
 - Jet energy scale calibrated with detector response from MC and 8 TeV data
 - Event rejected if contains jet identified as due to noise or non-collision
- **b-jets:**
 - Tagged by multivariate algorithm using the impact parameters of tracks in the jet, and the presence and flight paths of displaced vertices from b/c hadrons
- **Electrons:**
 - Matching EM calorimeter clusters to inner-detector tracks & TRT threshold
- **Muons:**
 - Matching tracks in the muon spectrometer and inner detector

- Physics-object overlap removal:
 - If 2 objects (e , μ , jet, or b-jet) are nearby, indicating mis-identification, one of them is discarded according to an optimized algorithm

- Missing transverse energy:

$$- \quad \vec{p}_T^{miss} = - \left[\sum_{\text{physics objects}} \vec{p}_T + \sum_{\text{other PV tracks}} \vec{p}_T \right]$$

$$- \quad \text{MET} \equiv E_T^{miss} = |\vec{p}_T^{miss}|$$

- Scalar p_T sum

$$- \quad H_T = \sum_{\text{physics objects}} p_T$$

- Effective mass:

$$- \quad m_{\text{eff}} = \sum_{\text{physics objects}} p_T + E_T^{miss}$$

- Define signal regions (SRs)
 - Based on N_{leptons} , N_{jets} , $N_{b\text{-jets}}$ with T cuts, H_T , MET, m_{eff} , etc.
 - Targeting different regions in SUSY parameter space
- Estimate background for each SR in control regions (CRs)
 - Usually using Monte Carlo distributions to relate CR yields to SR yields
 - Background estimate from CRs validated using validation regions (VRs)
 - Smaller backgrounds often obtained from MC
- If no excess, set limits using the CLs prescription, accounting for systematic uncertainties:
 - Finite MC statistics
 - Theory, e.g., models used for background shapes
 - Jet energy scale and resolution
 - Lepton / b-jet ID efficiencies and purities