THE $C_{\mathcal{F}}$ -THEOREM**G.M. SHORE***CERN - Geneva***Abstract**

The C -theorem corresponding to the spin 2 spectral function in the energy-momentum tensor Green function in $n > 2$ dimensions is derived. In 4 dimensions, renormalisation group methods are used to relate this $C_{\mathcal{F}}$ -function to the coefficient $\beta_a(g)$ of the $\mathcal{F}$ term in the curved space conformal anomaly, where $\mathcal{F}$ is the square of the Weyl tensor. This complements the recent formulation of the 4-dim $C_{\mathcal{H}}$ -theorem, where $C_{\mathcal{H}}(g)$ is characterised by the spin 0 spectral function and related to the R^2 term in the conformal anomaly. The potential application of the $C_{\mathcal{F}}$ -theorem to dynamical symmetry breaking is briefly discussed.

1. Introduction

An important advance in the search for a natural generalisation of the Zamolodchikov C -theorem[1] beyond 2 dimensions has been the realisation that the potential C -functions are intimately related to the coefficients in the curved space conformal anomaly[2-7]. Recall that in 2 dimensions, the conformal anomaly is simply (see, e.g., refs.[8,3])^{*}

$$\Theta = \beta^i O_i + \beta_b R, \quad (1.1)$$

where O_i are a basis of dimension 2 operators occurring in the Lagrangian of the particular field theory, g^i and $\beta^i(g)$ are the corresponding couplings and beta functions, and $\beta_b(g)$ is a beta function for the extra counterterm proportional to R needed to renormalise the theory in curved spacetime. (Explicitly, $\beta_b = \frac{1}{6} \frac{1}{4\pi} (N_S + \frac{1}{2} N_F) + O(g^i)$, where N_S and N_F are the number of scalar and Majorana fermion fields.) The Zamolodchikov C -theorem states[1]^{**}

$$-\beta^i \frac{\partial C_R}{\partial g^i} = -12 G_{ij} \beta^i \beta^j, \quad (1.2)$$

where G_{ij} is a positive definite metric on the space of coupling constants, defined by the Green function $x^2 \langle 0 | O_i(x) O_j(0) | 0 \rangle_R$. The function $C_R(g)$ is a positive function of the couplings, and is constrained by eq.(1.2) to decrease monotonically along an IR directed renormalisation group flow. At a fixed point, $C_R(g) = -96(2\pi)^{-1} \beta_b$, and is proportional to the central charge of the Virasoro algebra of the corresponding conformal field theory.

In 4 dimensions, the conformal anomaly is (see, e.g., refs.[8,9,3,4])

$$\Theta = \beta^i O_i - \beta_a \mathcal{F} - \beta_b \mathcal{G} - \beta_c \mathcal{H} - \frac{1}{3} d \nabla^2 R, \quad (1.3)$$

where $\mathcal{F}$ is the square of the Weyl tensor, $\mathcal{G}$ is the Euler density and $\mathcal{H}$ is $\frac{1}{9} R^2$. The functions $\beta_a(g)$ and $\beta_b(g)$, like β_b in the 2-dim theory, are non-zero even for free field theories (explicitly, $\beta_a = -\frac{1}{120} \frac{1}{(4\pi)^2} (N_S + 6N_F + 12N_V) + O(g)$ and $\beta_b = \frac{1}{360} \frac{1}{(4\pi)^2} (N_S + 11N_F + 62N_V) + O(g)$ where N_S , N_F and N_V are the number of scalar, Dirac fermion and vector fields resp.) while $\beta_c(g)$ is of $O(\beta^i)$ and so vanishes at a fixed point. Notice that in 2 dimensions there is no analogue of the $\mathcal{F}$ term, since the Weyl tensor vanishes identically, while the analogue of both the $\mathcal{G}$ and $\mathcal{H}$ terms is the single R term, R itself being the Euler density in 2 dimensions.

In a recent paper[7], it was shown that in 4 dimensions there exists a new C -theorem, the “ $C_{\mathcal{H}}$ -theorem”, describing the renormalisation group flow of a positive semi-definite

^{*} Notice that here we have denoted the coefficient of the R term in the 2-dim conformal anomaly by β_b , whereas $-\beta_c$ was used in ref.[7].

^{**} We introduce the following notation. C_R is the Zamolodchikov C -function in 2 dimensions. In 4 dimensions, the three C -functions are labelled by the corresponding terms in the conformal anomaly, viz. $C_{\mathcal{F}}$, $C_{\mathcal{G}}$ and $C_{\mathcal{H}}$. In the general n -dim analysis, the C -functions corresponding to the spin 0 and spin 2 spectral functions are denoted by $C^{(0)}$ and $C^{(2)}$. Thus, in 4 dimensions, $C^{(0)} \equiv C_{\mathcal{H}}$ and $C^{(2)} \equiv C_{\mathcal{F}}$, while in 2 dimensions $C^{(0)} \equiv C_R$ and $C^{(2)}$ does not exist.

scalar function $C_{\mathcal{H}}(g)$, viz.*

$$-\beta^i \frac{\partial C_{\mathcal{H}}}{\partial g^i} = 2C_{\mathcal{H}} - 8G_{ij}\beta^i\beta^j. \quad (1.4)$$

The function $C_{\mathcal{H}}(g)$, which, like its 2-dim counterpart, is defined in terms of Green functions of the energy-momentum tensor and its trace, is closely related to the $\mathcal{H}$ term in the conformal anomaly, and as a consequence of the 4-dim result $\beta_c = O(\beta^i)$ satisfies $C_{\mathcal{H}}(g) = O(\beta^i)$.

In ref.[7], we gave a unified derivation of the 2-dim Zamolodchikov theorem and the 4-dim $C_{\mathcal{H}}$ -theorem. In fact, both are special cases of a single n -dim formula,

$$-\beta^i \frac{\partial C^{(0)}}{\partial g^i} = (n-2)C^{(0)} - \frac{1}{2}(n-1)^{-1}(n+2)(n+4)G_{ij}\beta^i\beta^j, \quad (1.5)$$

whose proof assumes only the existence of a conserved energy-momentum tensor, with a trace anomaly determined by the beta functions of the field theory, and unitarity to ensure positivity of G_{ij} and $C^{(0)}(g)$.

The physical interpretation of this 4-dim theorem is nevertheless quite different from that of Zamolodchikov. $C_{\mathcal{H}}(g)$ need not decrease along a RG flow, but its rate of increase is bounded according to eq.(1.4) by virtue of the positivity of the metric G_{ij} . At a fixed point, rather than providing a measure of the degrees of freedom of the theory, $C_{\mathcal{H}}(g)$ vanishes.

It was conjectured in ref.[7] that this is the first of three C -theorems in 4 dimensions, corresponding to the three different terms in the conformal anomaly. These three theorems would all be natural generalisations of the unique 2-dim C -theorem, but each would have a distinct physical interpretation and carry different information about the field theory.

The purpose of this note is to prove the second of these conjectured theorems, the $C_{\mathcal{F}}$ -theorem. $C_{\mathcal{F}}(g)$ is again defined in terms of appropriate 2-point functions of the energy-momentum tensor, chosen this time to isolate structures related to the $\mathcal{F}$ term in the conformal anomaly. This theorem therefore has no direct 2-dim analogue. In 4 dimensions, we find

$$-\beta^i \frac{\partial C_{\mathcal{F}}}{\partial g^i} = 2C_{\mathcal{F}} - 24G_{\mathcal{F}}, \quad (1.6)$$

where here $G_{\mathcal{F}}$ is a positive definite function defined in terms of the Green function $x^8(\langle 0|T_{\mu\nu}(x) T^{\mu\nu}(0)|0\rangle_R - \frac{1}{3}\langle 0|\Theta(x) \Theta(0)|0\rangle_R)$. Again, $C_{\mathcal{F}}(g)$ may increase, but with a rate of increase bounded according to eq.(1.6). At a fixed point, $C_{\mathcal{F}}(g)$ is proportional to β_a , and so, as in the original Zamolodchikov theorem, provides a (weighted) measure of the degrees of freedom of the theory.

The $C^{(0)}$ -theorem was proved in ref.[7] using the spectral decomposition of the 2-point Green functions of the energy-momentum tensor, an approach pioneered by Friedan

* Notice that we have changed the normalisation of the $C^{(0)}$ and $C_{\mathcal{H}}$ -functions relative to ref.[7] (see eq.(2.1))

et al[5,6]. In general, this decomposition contains two distinct Lorentz structures corresponding to spin 0 and spin 2 intermediate states. The function $C^{(0)}(g)$ was defined in terms of the spin 0 spectral function $\rho^{(0)}(\lambda^2)$. This of course exists in 2 dimensions (where $C^{(0)} \equiv C_R$) as well as in 4 dimensions (where $C^{(0)} \equiv C_H$). In this paper, we define the function $C^{(2)}(g)$ similarly in terms of the spin 2 spectral function $\rho^{(2)}(\lambda^2)$, which exists only in dimensions greater than 2, and use the methods of ref.[7] to derive the $C^{(2)}$ -theorem in $n > 2$ dimensions. Then, using the renormalisation group equations for energy-momentum tensor Green functions in 4 dimensions, we show how this $C^{(2)}$ -function is related to the $\mathcal{F}$ term in the anomaly, and finally show explicitly that $C_{\mathcal{F}}(g) = -2^{11} 3^2 (2\pi)^{-2} \beta_a(g) + O(\beta^i)$.

2. Derivation of the C -theorems

The spectral function derivation of the C -theorems follows precisely the method already described in ref[7]. The starting point is the spectral representation of the Green function of two energy-momentum tensors, which, rescaling by a factor of x^{2n} to remove subtractions and other contact terms, implies

$$x^{2n} \langle 0 | T_{\mu\nu}(x) T_{\lambda\rho}(0) | 0 \rangle_R = x^{2n} \left(\Pi_{\mu\nu,\lambda\rho}^{(0)}(\partial) \Omega^{(0)} + \Pi_{\mu\nu,\lambda\rho}^{(2)}(\partial) \Omega^{(2)} \right), \quad (2.1)$$

with

$$\Omega^{(0)} = (2\pi)^{-\frac{n}{2}} \int_0^\infty d\lambda^2 \lambda^{-4} \rho^{(0)}(\lambda^2) (\lambda^{-1} x)^{1-\frac{n}{2}} K_{\frac{n}{2}-1}(\lambda x), \quad (2.2)$$

and

$$\Omega^{(2)} = (2\pi)^{-\frac{n}{2}} \int_0^\infty d\lambda^2 \lambda^{-4} \rho^{(2)}(\lambda^2) (\lambda^{-1} x)^{1-\frac{n}{2}} K_{\frac{n}{2}-1}(\lambda x), \quad (2.3)$$

where

$$\Pi_{\mu\nu,\lambda\rho}^{(0)} = \frac{1}{(n-1)} \Delta_{\mu\nu} \Delta_{\lambda\rho}, \quad (2.4)$$

with $\Delta_{\mu\nu} = (-\partial^2 g_{\mu\nu} + \partial_\mu \partial_\nu)$, and

$$\Pi_{\mu\nu,\lambda\rho}^{(2)} = \frac{1}{2} \frac{1}{(n-1)} (-2\Delta_{\mu\nu} \Delta_{\lambda\rho} + (n-1)(\Delta_{\mu\lambda} \Delta_{\nu\rho} + \Delta_{\mu\rho} \Delta_{\nu\lambda})), \quad (2.5)$$

are projection operators onto the spin 0 and spin 2 tensors $f_{\mu\nu} = \Delta_{\mu\nu} f$ and $h_{\mu\nu}$ (symmetric, traceless, transverse) respectively. $\Pi_{\mu\nu,\lambda\rho}^{(2)}$ is traceless, and vanishes identically for $n = 2$.

Performing the differentiation in eq.(2.1) gives

$$\begin{aligned} & x^{2n} \langle 0 | T_{\mu\nu}(x) T_{\lambda\rho}(0) | 0 \rangle_R \\ &= \frac{1}{(n-1)} (2\pi)^{-\frac{n}{2}} \int_0^\infty d\lambda^2 \lambda^{n-2} x^{2n} \rho^{(0)}(\lambda^2) \left(A^{(0)}(\lambda x) g_{\mu\nu} g_{\lambda\rho} + \dots + E^{(0)}(\lambda x) x^{-4} x_\mu x_\nu x_\lambda x_\rho \right) \\ &+ \frac{(n-2)}{(n-1)} (2\pi)^{-\frac{n}{2}} \int_0^\infty d\lambda^2 \lambda^{n-2} x^{2n} \rho^{(2)}(\lambda^2) \left(A^{(2)}(\lambda x) g_{\mu\nu} g_{\lambda\rho} + \dots + E^{(2)}(\lambda x) x^{-4} x_\mu x_\nu x_\lambda x_\rho \right). \end{aligned} \quad (2.6)$$

Contracting indices, we find the other two Green functions considered by Zamolodchikov, viz.

$$x^{2n} \langle 0 | T_{\mu\nu}(x) \Theta(0) | 0 \rangle_R = -x^{2n} \Delta_{\mu\nu} \partial^2 \Omega^{(0)}, \quad (2.7)$$

$$= -(2\pi)^{-\frac{n}{2}} \int_0^\infty d\lambda^2 \lambda^{n-2} x^{2n} \rho^{(0)}(\lambda^2) \left(P^{(0)}(\lambda x) g_{\mu\nu} + Q^{(0)}(\lambda x) x^{-2} x_\mu x_\nu \right), \quad (2.8)$$

and

$$x^{2n} \langle 0 | \Theta(x) \Theta(0) | 0 \rangle_R = (n-1) x^{2n} \partial^4 \Omega^{(0)}, \quad (2.9)$$

$$= (n-1) (2\pi)^{-\frac{n}{2}} \int_0^\infty d\lambda^2 \lambda^{n-2} x^{2n} \rho^{(0)}(\lambda^2) R^{(0)}(\lambda x), \quad (2.10)$$

which isolate the spin 0 spectral function. A different contraction isolates the spin 2 spectral function[6], viz.

$$\begin{aligned} & x^{2n} \left(\langle 0 | T_{\mu\lambda}(x) T_\nu{}^\lambda(0) | 0 \rangle_R - \frac{1}{(n-1)} \langle 0 | T_{\mu\nu}(x) \Theta(0) | 0 \rangle_R \right) \\ &= -\frac{1}{2} \frac{(n+1)(n-2)}{(n-1)} x^{2n} \Delta_{\mu\nu} \partial^2 \Omega^{(2)}, \end{aligned} \quad (2.11)$$

$$= -\frac{1}{2} \frac{(n+1)(n-2)}{(n-1)} (2\pi)^{-\frac{n}{2}} \int_0^\infty d\lambda^2 \lambda^{n-2} x^{2n} \rho^{(2)}(\lambda^2) \left(P^{(2)}(\lambda x) g_{\mu\nu} + Q^{(2)}(\lambda x) x^{-2} x_\mu x_\nu \right), \quad (2.12)$$

and

$$\begin{aligned} & x^{2n} \left(\langle 0 | T_{\mu\nu}(x) T^{\mu\nu}(0) | 0 \rangle_R - \frac{1}{(n-1)} \langle 0 | \Theta(x) \Theta(0) | 0 \rangle_R \right) \\ &= \frac{1}{2} (n+1)(n-2) x^{2n} \partial^4 \Omega^{(2)}, \end{aligned} \quad (2.13)$$

$$= \frac{1}{2} (n+1)(n-2) (2\pi)^{-\frac{n}{2}} \int_0^\infty d\lambda^2 \lambda^{n-2} x^{2n} \rho^{(2)}(\lambda^2) R^{(2)}(\lambda x). \quad (2.14)$$

The explicit expressions for these coefficient functions may be calculated using the recursion relations for the modified Bessel functions $K_\nu(x)$, as explained in ref.[7], and with the normalisations chosen as above we find simply

$$E^{(0)}(\lambda x) = E^{(2)}(\lambda x) = (\lambda x)^{1-\frac{n}{2}} K_{\frac{n}{2}+3}(\lambda x), \quad (2.15)$$

$$Q^{(0)}(\lambda x) = Q^{(2)}(\lambda x) = (\lambda x)^{1-\frac{n}{2}} K_{\frac{n}{2}+1}(\lambda x), \quad (2.16)$$

$$R^{(0)}(\lambda x) = R^{(2)}(\lambda x) = (\lambda x)^{1-\frac{n}{2}} K_{\frac{n}{2}-1}(\lambda x). \quad (2.17)$$

The dimensionless Zamolodchikov functions $F^{(s)}$, $H^{(s)}$ and $G^{(s)}$ corresponding to spins $s = 0, 2$ are defined from these Green functions as follows,

$$F^{(s)} = (2\pi)^{-\frac{n}{2}} \int_0^\infty d\lambda^2 \rho^{(s)}(\lambda^2) \lambda^{n-2} x^{2n} E^{(s)}(\lambda x), \quad (2.18)$$

$$H^{(s)} = (2\pi)^{-\frac{n}{2}} \int_0^\infty d\lambda^2 \rho^{(s)}(\lambda^2) \lambda^{n-2} x^{2n} Q^{(s)}(\lambda x), \quad (2.19)$$

$$G^{(s)} = (2\pi)^{-\frac{n}{2}} \int_0^\infty d\lambda^2 \rho^{(s)}(\lambda^2) \lambda^{n-2} x^{2n} R^{(s)}(\lambda x). \quad (2.20)$$

The derivation of the C -theorem corresponding to the spin 2 spectral function now follows precisely that already given in ref.[7] for the spin 0 case. We define the $C^{(s)}$ -functions as

$$C^{(s)} = F^{(s)} + \left(\frac{n}{2} + 1\right)H^{(s)} - \left(\frac{n}{2} + 2\right)G^{(s)}. \quad (2.21)$$

These satisfy the equation,

$$x \frac{\partial C^{(s)}}{\partial x} = (n-2)C^{(s)} - \frac{1}{2}(n+2)(n+4)G^{(s)}, \quad (2.22)$$

and using dimensional analysis and the RGE for $C^{(s)}(g)$ (noting, as in ref.[7], that the inhomogeneous terms present in the RGEs for the 2-point functions of the energy-momentum tensor do not appear in the RGEs for the rescaled functions $F^{(s)}$, $H^{(s)}$, $G^{(s)}$) we find

$$-\beta^i \frac{\partial C^{(s)}}{\partial g^i} = (n-2)C^{(s)} - \frac{1}{2}(n+2)(n+4)G^{(s)}. \quad (2.23)$$

This is the basic result, describing the RG flow of the $C^{(s)}$ -functions defined in terms of the spectral functions by

$$C^{(s)} = (2\pi)^{-\frac{n}{2}} \int_0^\infty d\lambda^2 \lambda^{-n-2} \rho^{(s)}(\lambda^2) (\lambda x)^{\frac{3n}{2}+1} \times \left(K_{\frac{n}{2}+3}(\lambda x) + \left(\frac{n}{2} + 1\right) K_{\frac{n}{2}+1}(\lambda x) - \left(\frac{n}{2} + 2\right) K_{\frac{n}{2}-1}(\lambda x) \right). \quad (2.24)$$

Unitarity, which implies that the spectral functions are positive definite, together with the property $K_{\nu+1}(x) > K_\nu(x) > 0$ ($\nu \geq 0$), ensures that both the $G^{(s)}$ and $C^{(s)}$ -functions are positive (semi-)definite. However, we can only write $G^{(s)}$ in terms of the metric $G_{ij} = x^{2n} \langle 0 | O_i(x) O_j(0) | 0 \rangle_R$ in the spin 0 case, where $G^{(0)} = (n-1)^{-1} G_{ij} \beta^i \beta^j$.

An alternative representation of the $C^{(s)}$ -functions, independent of spectral functions, can be given by simply writing

$$W_{\Theta\Theta}^R(x) \equiv \langle 0 | \Theta(x) \Theta(0) | 0 \rangle_R = \int \frac{d^n k}{(2\pi)^n} e^{ik \cdot x} k^n \hat{F}_{\Theta\Theta}(t), \quad (2.25)$$

where $t = \frac{1}{2} \log \frac{k^2}{\mu^2}$, and

$$W_{TT}^R(x) \equiv \langle 0 | T_{\mu\nu}(x) T^{\mu\nu}(0) | 0 \rangle - \frac{1}{(n-1)} \langle 0 | \Theta(x) \Theta(0) | 0 \rangle_R = \int \frac{d^n k}{(2\pi)^n} e^{ik \cdot x} k^n \hat{F}_{TT}(t). \quad (2.26)$$

Then,

$$\Omega^{(0)} = \frac{1}{(n-1)} (2\pi)^{-\frac{n}{2}} \int_0^\infty dk k^{\frac{n}{2}-4} x^{1-\frac{n}{2}} J_{\frac{n}{2}-1}(kx) k^n \hat{F}_{\Theta\Theta}(t) + \text{c.t.}, \quad (2.27)$$

and

$$\Omega^{(2)} = \frac{2}{(n+1)(n-2)}(2\pi)^{-\frac{n}{2}} \int_0^\infty dk k^{\frac{n}{2}-4} x^{1-\frac{n}{2}} J_{\frac{n}{2}-1}(kx) k^n \hat{F}_{TT}(t) + \text{c.t.}, \quad (2.28)$$

where c.t. denotes contact terms which vanish when acted on by $x^{2n}\Pi^{(s)}$.

We can now repeat the steps above leading to the C -theorem, this time using the recursion relations for the ordinary Bessel functions $J_\nu(x)$ and expressing the Zamolodchikov functions in terms of $\hat{F}_{\Theta\Theta}$ and $\hat{F}_{TT}$ (see ref.[7]). We recover the C -theorem (2.22), where the $C^{(s)}$ -functions are now represented by,

$$C^{(0)} = \frac{1}{(n-1)}(2\pi)^{-\frac{n}{2}} \int_0^\infty dk k^{-1}(kx)^{\frac{3n}{2}+1} \hat{F}_{\Theta\Theta}(t) \\ \times \left(J_{\frac{n}{2}+3}(kx) - \left(\frac{n}{2}+1\right)J_{\frac{n}{2}+1}(kx) - \left(\frac{n}{2}+2\right)J_{\frac{n}{2}-1}(kx) \right), \quad (2.29)$$

and

$$C^{(2)} = \frac{2}{(n+1)(n-2)}(2\pi)^{-\frac{n}{2}} \int_0^\infty dk k^{-1}(kx)^{\frac{3n}{2}+1} \hat{F}_{TT}(t) \\ \times \left(J_{\frac{n}{2}+3}(kx) - \left(\frac{n}{2}+1\right)J_{\frac{n}{2}+1}(kx) - \left(\frac{n}{2}+2\right)J_{\frac{n}{2}-1}(kx) \right) \quad (2.30)$$

The functions $G^{(0)}$ and $G^{(2)}$ are defined similarly with the linear combinations of Bessel functions in brackets being replaced simply by $J_{\frac{n}{2}-1}(kx)$. The positivity properties of the $G^{(s)}$ and $C^{(s)}$ -functions, which were manifest in the representation in terms of spectral functions and the modified Bessel functions K_ν , are no longer clear from eqs.(2.29) and (2.30). However, this is the form in which it is easiest to relate $C^{(s)}(g)$ to the RG functions occurring in the conformal anomaly.

3. RGEs and the $C_{\mathcal{F}}$ -theorem

We are now in a position to consider the RGEs for the 2-point Green functions of the energy-momentum tensor and relate the $C^{(2)}$ -function in 4 dimensions to the coefficient β_a of the $\mathcal{F}$ term in the curved space conformal anomaly. The close relation between the spin 2 spectral function and the $\mathcal{F}$ term has been previously observed in ref.[6]. Following refs.[3,4], we write the dimensionally regularised 4-dim action in curved space, with position-dependent couplings to allow a systematic treatment of the renormalisation of composite operator Green functions, as

$$S_{\text{reg}} = S_B - \int d^n x \sqrt{g} \left(a_B \mathcal{F} + b_B \mathcal{G} + c_B \mathcal{H} + \frac{1}{2} A_{ij}^B \nabla^2 g^i \nabla^2 g^j + \dots \right), \quad (3.1)$$

where a_B , b_B , c_B and A_{ij}^B are divergent counterterms, and the precise forms of $\mathcal{F}$, $\mathcal{G}$ and $\mathcal{H}$ convenient for dimensional regularisation are defined in ref.[4].

Renormalised Green functions are defined by functional differentiation of the generating functional W , and the counterterms above contribute inhomogeneous terms which cancel the new divergences which arise in 2-point functions of renormalised composite operators. Thus,

$$\tilde{W}_{ij}^R(k) \equiv \langle 0|O_i(k) O_j(-k)|0\rangle_R = \langle 0|O_i(k) O_j(-k)|0\rangle + A_{ij}^B k^4. \quad (3.2)$$

Differentiating twice w.r.t. the metric $g_{\mu\nu}$ and taking the flat space limit, we find the renormalised energy-momentum tensor Green function,*

$$\langle 0|T_{\mu\nu}(k) T_{\lambda\rho}(-k)|0\rangle_R = \langle 0|T_{\mu\nu}(k) T_{\lambda\rho}(-k)|0\rangle + \frac{8}{(n-1)} c_B \Pi_{\mu\nu,\lambda\rho}^{(0)}(k) + \frac{8(n-3)}{(n-2)} a_B \Pi_{\mu\nu,\lambda\rho}^{(2)}(k). \quad (3.3)$$

This is one of the key results of this paper. It shows clearly how the spin 0 part of the energy-momentum tensor Green function is related to the $\mathcal{H}$ term in the regularised action, while the spin 2 part is related to the $\mathcal{F}$ term.

The different traces isolating the spin 0 and spin 2 contributions therefore give,

$$\tilde{W}_{\Theta\Theta}^R(k) \equiv \langle 0|\Theta(k) \Theta(-k)|0\rangle_R = \langle 0|\Theta(k) \Theta(-k)|0\rangle + 8c_B k^4, \quad (3.4)$$

and

$$\begin{aligned} \tilde{W}_{TT}^R(k) &\equiv \langle 0|T_{\mu\nu}(k) T^{\mu\nu}(-k)|0\rangle_R - \frac{1}{(n-1)} \langle 0|\Theta(k) \Theta(-k)|0\rangle_R \\ &= \langle 0|T_{\mu\nu}(k) T^{\mu\nu}(-k)|0\rangle - \frac{1}{(n-1)} \langle 0|\Theta(k) \Theta(-k)|0\rangle + 4(n+1)(n-3)a_B k^4. \end{aligned} \quad (3.5)$$

The RGEs follow immediately,

$$\mathcal{D}\tilde{W}_{ij}^R + \partial_i \beta^k \tilde{W}_{kj}^R + \partial_j \beta^k \tilde{W}_{ik}^R + \chi_{ij}^a k^4 = 0, \quad (3.6)$$

$$\mathcal{D}\tilde{W}_{\Theta\Theta}^R + 8\beta_c k^4 = 0, \quad (3.7)$$

$$\mathcal{D}\tilde{W}_{TT}^R + 20\beta_a k^4 = 0. \quad (3.8)$$

The solutions of the RGEs for $\tilde{W}_{ij}^R$ and $\tilde{W}_{\Theta\Theta}^R$ have been discussed in ref.[7], where a comparison is also made of the relation between the C -theorems and the RG consistency conditions of Osborn and Jack[3,4]. Here, we focus on the $\tilde{W}_{TT}^R$ equation. The solution is

$$\tilde{W}_{TT}^R(k) = k^4 \left(F_{TT}(\mu, g(t)) + 20 \int_0^t dt' \beta_a(g(t')) \right), \quad (3.9)$$

* In eqs.(3.3), (3.4) and (3.5) we have omitted contact terms proportional to the VEV $\langle \Theta \rangle$. These do not contribute to the RGEs, and are eliminated when we pass from the Green functions themselves to the rescaled Zamolodchikov functions $F^{(s)}$, $H^{(s)}$, $G^{(s)}$ and the $C^{(s)}$ -functions.

which, expanding for small t , gives

$$\tilde{W}_{TT}^R(k) = k^4 \left(F_{TT}(\mu, g) + t(20\beta_a + \beta^k \partial_k F_{TT}) + O(t^2) \right). \quad (3.10)$$

The higher order terms in this expansion are all of $O(\beta^i)$. Substituting into the expression (2.30) for the function $C^{(2)}(g)$, and with the notation $C^{(2)}|_{n=4} \equiv C_{\mathcal{F}}$, we find*

$$C_{\mathcal{F}}(g) = \frac{1}{5}(2\pi)^{-2} \int_0^\infty dk k^{-1} (kx)^7 \hat{F}_{TT}(t) \left(J_5(kx) - 3J_3(kx) - 4J_1(kx) \right), \quad (3.11)$$

$$= -\frac{1}{5} 2^9 3^2 (2\pi)^{-2} (20\beta_a + \beta^k \partial_k F_{TT}) + \dots \quad (3.12)$$

To summarise, therefore, we have found in 4 dimensions the “ $C_{\mathcal{F}}$ theorem”,

$$-\beta^i \frac{\partial C_{\mathcal{F}}}{\partial g^i} = 2C_{\mathcal{F}} - 24G_{\mathcal{F}}, \quad (3.13)$$

where,

$$C_{\mathcal{F}}(g) = -2^{11} 3^2 (2\pi)^{-2} \beta_a + O(\beta^i). \quad (3.14)$$

4. Discussion

As already noted, because its definition involves the Green function $\langle 0|T_{\mu\nu}(x) T^{\mu\nu}(0)|0\rangle_R$ rather than just $\langle 0|\Theta(x) \Theta(0)|0\rangle_R$, the function $G_{\mathcal{F}}$ occurring in eq.(3.13) cannot be written in terms of the metric G_{ij} . Nevertheless, it is positive, and so the rate of increase of $C_{\mathcal{F}}(g)$ along an IR directed RG flow is bounded.

Despite the presence of two terms of opposite sign on the RHS of eq.(3.13), it could still be possible that $C_{\mathcal{F}}(g)$ changes monotonically if we could show that $C_{\mathcal{F}}$ is always less (or greater) than $12G_{\mathcal{F}}$. We can write the difference as the single integral,

$$C_{\mathcal{F}} - 12G_{\mathcal{F}} = (2\pi)^{-2} \int_0^\infty d\lambda^2 \lambda^{-6} \rho^{(2)}(\lambda^2) (\lambda x)^7 \left(K_5(\lambda x) + 3K_3(\lambda x) - 16K_1(\lambda x) \right). \quad (4.1)$$

* Here, we use the standard integrals

$$\int_0^\infty dk k^p J_\nu(kx) = 2^p x^{-1-p} \frac{\Gamma(\frac{1}{2}(\nu + p + 1))}{\Gamma(\frac{1}{2}(\nu - p + 1))},$$

$$\int_0^\infty dk k^p J_\nu(kx) \log k = 2^{p-1} x^{-1-p} \frac{\Gamma(\frac{1}{2}(\nu + p + 1))}{\Gamma(\frac{1}{2}(\nu - p + 1))} \left(\psi(\frac{1}{2}(\nu + p + 1)) + \psi(\frac{1}{2}(\nu - p + 1)) - \log \frac{1}{4} x^2 \right).$$

The combination of Bessel functions in brackets has the shape of a van der Waal's potential – at fixed x , it tends to $+\infty$ for $\lambda \rightarrow 0$, dips to a minimum below zero for some finite λ , and approaches zero from below as $\lambda \rightarrow \infty$. Varying x simply changes the horizontal scale. $C_{\mathcal{F}} - 12G_{\mathcal{F}}$ does not therefore have a definite sign independent of the form of $\rho^{(2)}(\lambda^2)$. It is therefore difficult to come to hard conclusions about the evolution of $C_{\mathcal{F}}(g)$ without making some assumptions about $\rho^{(2)}(\lambda^2)$. However, for any reasonable spectral function, the integral in eq.(4.1) is positive for x sufficiently small. We can therefore conclude that as we move away from the UV limit $C_{\mathcal{F}}(g)$ initially rises. The same analysis would of course apply to $C_{\mathcal{H}}(g)$, which we recall is constrained to be zero at a fixed point.

The original suggestion by Cardy[2] was that in 4 dimensions $\beta_b(g)$ should play the role of the Zamolodchikov C -function. This conjecture was supported by an application to QCD. It is interesting to discuss the function $C_{\mathcal{F}}(g)$ in the same light. Assume QCD has an UV fixed point at zero coupling, and that the coupling then grows to infinity under a RG flow into the IR. Then, at the UV fixed point, $C_{\mathcal{F}}(g)$ is proportional to $\beta_a(g^*)$ where $g^* = 0$. $C_{\mathcal{F}}|_{UV}$ therefore measures the weighted degrees of freedom $(N_S + 6N_F + 12N_V)|_{QCD}$ of the original Lagrangian. In the infinite coupling IR limit, the spectrum of QCD is described by a free effective chiral Lagrangian (CL) describing only the states which remain massless in the confined, chiral symmetry broken phase, all other states becoming infinitely massive. We can therefore evaluate $C_{\mathcal{F}}(g)$ in the IR limit as proportional to $\beta_a(0)|_{CL}$, i.e. the combination $(N_S + 6N_F + 12N_V)|_{CL}$ in the effective chiral Lagrangian. For massless QCD with N colours and M flavours, $\beta_a(0)|_{QCD} \propto 12(N^2 - 1) + 6NM$, from the quarks and gluons, while $\beta_a(0)|_{CL} \propto M^2 - 1$, from the pions. In the asymptotically free theory, where $M < \frac{11}{2}N$ we therefore find $C_{\mathcal{F}}|_{UV} - C_{\mathcal{F}}|_{IR} > 0$. This condition is of course more stringent with β_a than with β_b . Another well-controlled example is supersymmetric QCD with $N = M$, where all condensates are finite as the quark mass tends to zero and the theory can be described similarly to ordinary QCD[10]. Here, the chiral symmetry $SU(M)_L \times SU(M)_R \times U(1)_V \times U(1)_X$ is spontaneously broken to $SU(M-P)_V \times SU(P)_V \times U(1)_X$. However, (see e.g. ref.[11]), independently of the value of P , the Goldstone bosons and their scalar and fermion supersymmetry partners fall into M^2 chiral multiplets. Since the quarks and squarks form NM chiral multiplets, it is clear that for $N = M$ SQCD we again find $C_{\mathcal{F}}|_{UV} - C_{\mathcal{F}}|_{IR} > 0$.

If this result were generally true, it would provide a very interesting constraint on the allowed patterns of dynamical symmetry breaking in 4-dim field theories, reminiscent of, and perhaps not unrelated to, the 't Hooft anomaly matching conditions[12]. However, as we have shown above, it is not clear how this could be proved from the $C_{\mathcal{F}}$ -theorem (3.13) without further assumptions on the form of the spectral function.

It is therefore uncertain at present how useful these new C -theorems will prove to be in analysing non-perturbative properties of field theories, either in constraining dynamical symmetry breaking (for which $C_{\mathcal{F}}$ seems most suited) or providing information on the fixed point structure (where $C_{\mathcal{H}}$ may play a rôle). It also remains to be seen what is the precise form of the conjectured $C_{\mathcal{G}}$ -theorem, and in particular whether this exhibits the monotonic behaviour which we expect of an entropy-like function characterising the information loss associated with an IR RG flow.

Acknowledgements

I would like to thank Ian Jack and Hugh Osborn for many useful discussions.

References

- [1] A.B. Zamolodchikov, JETP Lett 43 (1986) 930, Sov J Nucl Phys 46 (1988) 1090
- [2] J.L. Cardy, Phys Lett B215 (1988) 749
- [3] H. Osborn, Phys Lett B222 (1989) 97
- [4] I. Jack and H. Osborn, Nucl Phys B343 (1990) 647
- [5] D. Friedan, lectures at the Nordisk Forskarkurs, Recent Developments in Quantum Field Theory, July 1990, Langarvatn, Iceland, unpublished
- [6] A. Cappelli, D. Friedan and J.I. Latorre, “*C*-theorem and Spectral Representation”, Rutgers-Barcelona preprint, RU-90-43, UB-ECM-PF 11/90
- [7] G.M. Shore, “A New *C*-Theorem in Four Dimensions”, CERN preprint TH.5886/90
- [8] M.J. Duff, Nucl Phys B125 (1977) 334
- [9] L.S. Brown and J.C. Collins, Ann Phys 130 (1981) 215
S.J. Hathrell, Ann Phys 139 (1982) 136, *ibid* 142 (1982) 34
M.D Freeman, Ann Phys 153 (1984) 339
- [10] D. Amati, K. Konishi, Y. Meurice, G.C. Rossi and G. Veneziano, Phys Rep 162 (1988) 169
- [11] A.C.W. Kotcheff and G.M Shore, Int J Mod Phys A4 (1989) 4391
- [12] G. 't Hooft, in Recent Developments in Gauge Theories, Cargèse 1979, ed. G. 't Hooft et al (Plenum, New York, 1980)