CERN-TH/6078/91
CPT-91/P.2537

**REAL PART OF THE $\bar{p}p$ ELASTIC SCATTERING
AMPLITUDE AT $\sqrt{s} = 546$ GeV**

Claude Bourrely and Jacques Soffer
Centre de Physique Théorique, C.N.R.S. Luminy, Case 907,
13288 MARSEILLE Cedex 9

and

Tai Tsun Wu*
CERN - Geneva
and
Gordon McKay Laboratory, Harvard University
CAMBRIDGE, MA 02138

Abstract

In order to get an accurate determination of the real part of the $\bar{p}p$ elastic scattering amplitude at $\sqrt{s} = 546$ GeV, the Genoa-Palaiseau-Roma-Valencia collaboration plans to take experimental data on $d\sigma/dt$ in the Coulomb interference region. We present detailed theoretical predictions for this cross section. These predictions can be compared directly with the forthcoming data.

CERN-TH/6078/91
CPT-91/P.2537
June 1991

* Work supported in part by the US Department of Energy under Grant n° DE-FG02-84ER40158.

Seven years ago, it was predicted in the framework of the impact-picture phenomenology [1] that the ratio of the real to imaginary parts of the forward $\bar{p}p$ elastic amplitude

$$\rho = \frac{\text{Re } a^N(s, t=0)}{\text{Im } a^N(s, t=0)} \quad (1)$$

is about 0.13 [2] at energies of CERN $S\bar{p}pS$ collider. At that time there was only one accurate data point [3] for total cross section above ISR energies.

Realizing the difficulty of extrapolating the experimental data on the differential cross section $d\sigma/dt$ to the forward direction $t=0$, we utilized the impact-picture phenomenology of ref.[2] to predict the values of $d\sigma/dt$ in the Coulomb interference region [4]. This prediction is shown as a curve in fig.1. Shortly thereafter the UA4 collaboration published the data shown in fig.1 [5]. It is seen that the data gives a good confirmation of the theoretical prediction ^{a)}

Since this experimental result of UA4 was based on two days of running, the Genoa-Palaiseau-Roma-Valencia collaboration [6] proposed last year a more precise measurement in the Coulomb interference region. This proposal has been approved and the plan is to take data near the end of this year. It is therefore an opportune time to present here more detailed theoretical predictions of the impact-picture. It should be emphasized that, aside from using the Coulomb amplitude with the West-Yennie phase [7], no new parameter or modification has been introduced beyond ref.[2] of 1984.

Table 1 shows the values of $d\sigma/dt$ in the Coulomb interference region at $\sqrt{s} = 546$ GeV which is the energy of the forthcoming experiment.

^{a)} In spite of the good agreement shown in fig.1, the extrapolation used by the UA4 collaboration gives $\rho = 0.24 \pm 0.04$. Thus a comparison of theoretical predictions and experimental data should be carried out directly on $d\sigma/dt$ in the Coulomb interference region.

Acknowledgments

T.T. Wu is grateful to John Ellis, Maurice Jacob and many other members of the Theory Division for their kind hospitality at CERN.

Figure Caption

Fig.1 Comparison of theoretical prediction of ref.[4] and experimental measurement by UA4 at $\sqrt{s} = 546$ GeV in the small t -region from ref.[5].

Table Caption

Table 1 Theoretical predictions for the $\bar{p}p$ elastic differential cross section at $\sqrt{s} = 546$ GeV near the forward direction.

Table 1

$ t (GeV)^2 \quad d\sigma/dt \text{ mb}/(GeV)^2$		$ t (GeV)^2 \quad d\sigma/dt \text{ mb}/(GeV)^2$		$ t (GeV)^2 \quad d\sigma/dt \text{ mb}/(GeV)^2$	
0.0004	2149.10	0.0215	139.57	0.0520	84.71
0.0006	1134.70	0.0220	138.38	0.0530	83.38
0.0008	749.36	0.0225	137.20	0.0540	82.07
0.0010	557.99	0.0230	136.03	0.0550	80.78
0.0015	355.06	0.0235	134.88	0.0560	79.52
0.0020	279.99	0.0240	133.74	0.0570	78.28
0.0025	244.79	0.0245	132.61	0.0580	77.06
0.0030	225.62	0.0250	131.50	0.0590	75.86
0.0035	213.93	0.0255	130.39	0.0600	74.68
0.0040	206.10	0.0260	129.30	0.0610	73.52
0.0045	200.43	0.0265	128.22	0.0620	72.38
0.0050	196.04	0.0270	127.15	0.0630	71.26
0.0055	192.44	0.0275	126.08	0.0640	70.15
0.0060	189.36	0.0280	125.04	0.0650	69.07
0.0065	186.64	0.0285	124.00	0.0660	68.01
0.0070	184.18	0.0290	122.97	0.0670	66.96
0.0075	181.90	0.0295	121.95	0.0680	65.93
0.0080	179.77	0.0300	120.94	0.0690	64.92
0.0085	177.74	0.0305	119.94	0.0700	63.92
0.0090	175.81	0.0310	118.95	0.0720	61.98
0.0095	173.96	0.0315	117.97	0.0740	60.10
0.0100	172.16	0.0320	117.00	0.0760	58.29
0.0105	170.43	0.0325	116.04	0.0780	56.53
0.0110	168.74	0.0330	115.09	0.0800	54.83
0.0115	167.09	0.0335	114.15	0.0820	53.19
0.0120	165.49	0.0340	113.21	0.0840	51.60
0.0125	163.91	0.0345	112.29	0.0860	50.05
0.0130	162.38	0.0350	111.37	0.0880	48.56
0.0135	160.87	0.0360	109.56	0.0900	47.12
0.0140	159.38	0.0370	107.79	0.0920	45.73
0.0145	157.92	0.0380	106.05	0.0940	44.37
0.0150	156.49	0.0390	104.34	0.0960	43.06
0.0155	155.08	0.0400	102.66	0.0980	41.79
0.0160	153.69	0.0410	101.01	0.1000	40.57
0.0165	152.32	0.0420	99.40	0.1020	39.38
0.0170	150.97	0.0430	97.81	0.1040	38.23
0.0175	149.64	0.0440	96.24	0.1060	37.11
0.0180	148.32	0.0450	94.71	0.1080	36.03
0.0185	147.03	0.0460	93.21	0.1100	34.98
0.0190	145.74	0.0470	91.73	0.1120	33.97
0.0195	144.48	0.0480	90.27	0.1140	32.98
0.0200	143.23	0.0490	88.85	0.1160	32.03
0.0205	142.00	0.0500	87.44	0.1180	31.11
0.0210	140.78	0.0510	86.06	0.1200	30.21

References

- [1] H. Cheng and T.T. Wu, Phys. Rev. Lett. **24** (1970) 1456;
C. Bourrely, J. Soffer and T.T. Wu, Phys. Rev. **D19** (1979) 3249.
- [2] C. Bourrely, J. Soffer and T.T. Wu, Nucl. Phys. **B247** (1984) 15.
- [3] UA4 Collaboration, M. Bozzo et al., Phys. Lett. **B147** (1984) 392.
- [4] C. Bourrely, J. Soffer and T.T. Wu, Phys. Lett. **B196** (1987) 237.
- [5] UA4 Collaboration, D. Bernard et al., Phys. Lett. **B198** (1987) 583.
- [6] M. Bozzo, G. Sette, D. Bernard, M. Haguenauer, L. Cerrito, G. Matthiae, E. Sanchis
and J. Velasco, CERN/SPSC 90-9, SPSC/P250 (1990).
- [7] G.B. West and D.R. Yennie, Phys. Rev. **172** (1968) 1413.

Fig. 1