CERN-TH.6184/91

IEM-FT-33/91

Correlation between $M_{Z'}$ and m_t bounds: II. All data*

F. del Aguila[†]

*CERN, Geneva and
Departamento de Física Teórica y del Cosmos
Universidad de Granada, E-18071 Granada, Spain*

W. Hollik*

*Max-Planck-Institut für Physik
- Werner-Heisenberg-Institut -
P.O. Box 40 12 12, Munich, Germany*

J.M. Moreno

*Departement de Physique Théorique
Université Libre de Bruxelles, Boulevard du Triomphe
CP 225, B-1050 Bruxelles, Belgium*

M. Quirós[‡]

*Instituto de Estructura de la Materia
Serrano 119, E-28006 Madrid, Spain*

ABSTRACT

We present a global fit to neutral current data (including improved parity-violation data), M_W and the last year LEP results in the standard model and its minimal gauge extensions. In the standard model, for $m_H = 100 \text{ GeV}$ [$m_H = 1 \text{ TeV}$] and $\alpha_s(M_Z) = 0.12$, $m_t = 112^{+22}_{-23} \text{ GeV}$ [$m_t = 140^{+19}_{-21} \text{ GeV}$] (1σ). For gauge extensions with a minimal (arbitrary) Higgs doublet content, the $M_{Z'}$ lower bound improves (with respect to our previous analysis) by ~ 200 (50) GeV . In the case with an arbitrary Higgs doublet content, the bound on the $Z^0 Z'$ mixing is reduced by a factor of ~ 4 . No data set seems to favour a new Z' , meaning that present Z' limits are governed by the data error size only. The relevance of the different data in the fits is discussed, as well as the correlation between $M_{Z'}$ and $m_{t,H}$ bounds. Present limits on *magic* observables and mixing parameters are given.

* Work partly supported by CICYT under contracts AEN90-0139, AEN90-0683.

† Electronic mail: faguila@ugr.es and vxcern::aguila

* Electronic mail: woh@dm0mpi11

‡ Electronic mail: imtma27@cc.csic.es and quiros@cernvm

1 Introduction

Due to the absence of any positive signal of new Physics beyond the standard model at low energy, our empirical knowledge of the possible extensions of the standard model reduces to bounds. In this paper we complete the analysis of present limits on new neutral gauge interactions [1] by including in the global fits the improved parity-violation data [2] and the last year LEP results on Z widths and asymmetries [3]. We follow the notation and hypotheses of Ref. [1]. (There a more exhaustive list of references was also given.) The necessary equations to extend the global fit to LEP data are given in the Appendix, which completes the Appendix of Ref. [1]. The one-loop radiative corrections within the minimal standard model are taken into account, whereas only tree-level Z' contributions are included [4].

In Section 2 we review the influence of the different data pieces, including the variation with α_s , on a global fit to the standard model. The corresponding constraints on the minimal gauge extension of the standard model, as well as on illustrative (popular) models are presented in Section 3. The correlation between the top and Higgs mass bounds and the M_Z limits are discussed in Section 4, together with the experimental limits on *smart* observables already introduced in the literature [5]. Section 5 is devoted to conclusions. For recent related analyses, see Refs. [6, 7, 8].

Although different data sets constrain differently the standard model (or its extensions), we may say as a summary that νq and parity-violation constrain m_t similarly, once M_Z is taken into account in the fits. Analogously, all neutral current data and last year LEP results give a similar m_t bound. M_W is less restrictive at present. On the other hand, the data precision implies that m_H already plays an important rôle in the m_t determination. As a matter of fact, the error in m_t is of the same order as the uncertainty introduced by m_H . Finally the error in α_s introduces a sizeable uncertainty in the m_t value. We find in the minimal standard model (at 90% c.l. for two variables, if we consider M_Z as an adjustable variable) ¹

$$\begin{array}{ll} 89 \text{ GeV} < m_t < 155 \text{ GeV} & \text{for } m_H = 100 \text{ GeV} \text{ and } \alpha_s(M_Z) = 0.11, \\ 89 \text{ GeV} < m_t < 147 \text{ GeV} & 100 \text{ GeV} \quad 0.13, \\ 98 \text{ GeV} < m_t < 178 \text{ GeV} & 1 \text{ TeV} \quad 0.11, \\ 90 \text{ GeV} < m_t < 169 \text{ GeV} & 1 \text{ TeV} \quad 0.13. \end{array} \quad (1.1)$$

¹ m_t must be larger than its direct production limit [9].

In the minimal gauge extensions of the standard model the improvement in the parity-violation data and in the last year LEP results translate into a large improvement in the M_Z bound ($\sim 200 \text{ GeV}$). For models where the $Z^0 Z'$ mixing angle is a free parameter the improvement is of $\sim 50 \text{ GeV}$, but the limit on the mixing angle gets reduced by a factor ~ 4 .

2 Standard model constraints

In the minimal standard model only the top and Higgs masses, m_t and m_H , remain as unknown parameters. They are bounded from above, however, by their indirect effects [10] and also from below by direct production experiments [9, 11]. In this Section we discuss their present bounds from precise electroweak data. In our fits we take α and G_μ as input parameters [12]. The precise data we use are summarized in Table 1. νq , $\nu_\mu e$ and eH data were collected by P. Langacker [12]. We also use the improved parity-violation value of Q_W [2], the world average value of M_W [12, 13], and the LEP data presented in Ref. [3]. Although different data pieces help to constrain different types (or parameters) of standard model extensions, each piece alone gives similar top upper bounds in the minimal standard model (taking the central value of M_Z as an input parameter). The parity-violation data, summarized by the C coefficients in Table 1, are dominated by the Q_W combination

$$Q_W = -2[C_{1u}(2Z + N) + C_{1d}(Z + 2N)]. \quad (2.1)$$

For this reason, when combined with other data, it is enough to consider Q_W . We will use the improved Q_W value quoted in Table 1 [2]. (To be compared to the Q_W value from Eq. (2.1) and the C coefficients in Table 1 for ${}^{133}_{55}\text{Cs}$, $Q_W = -69.87 \pm 3.81$.) The 90% c.l. regions (for two variables) in the $\sin^2\theta_W - m_t$ plane, allowed by νq and Q_W data, respectively, are shown in Figs. 1a, 1b for $m_H = 100, 1000 \text{ GeV}$ and $\alpha_s(M_Z) = 0.12$. Due to the precision with which M_Z is known, when included in the fits it is like fixing M_Z to its central experimental value, and then the bounds implied by the other data pieces are given to a large extent by the intersection of the corresponding data band with the narrow M_Z band in the figure. Hence, Fig. 1 tells us that νq and Q_W imply similar m_t bounds. In this figure we also show the global fit (solid contour) to all ($\nu q, \nu_\mu e, Q_W, M_W$ and the last year LEP results, including M_Z) data. Comparing Figs. 1a and 1b we see that the central m_t value increases

with m_H , and similarly the m_t bounds. A change of m_H from 100 GeV to 1 TeV translates into an increase of ~ 25 GeV for m_t . In Figs. 2a, 2b we compare the neutral current ($\nu q, \nu_e e$ and Q_W) bounds with the M_W ones and with the bounds due to the recent LEP results on Z decay rates, Γ , and asymmetries, A . Thus, whereas M_W (and M_Z) can not constrain m_t , the m_t limits resulting from each of the other data sets are comparable.

In next Sections we fix M_Z to its central value and $\alpha_s(M_Z)$ to 0.12. However, one may wonder about the effect of the α_s variation on the m_t limits. This is shown in Figs. 3a, 3b where we plot for the Z widths and asymmetries and for all data, respectively, the 90% c.l. (for two variables) allowed regions for $\alpha_s(M_Z) = 0.11$ (solid) and 0.13 (dashed) (for $\alpha_s(M_Z) = 0.12$ the results were shown in Figs. 1, 2). The neutral current, M_W and M_Z regions are unchanged. The global fit gives the m_t values quoted in Eq. (1.1). Finally, in Figs. 4a, 4b we plot the χ^2 distributions as functions of m_t .

3 Extended gauge model constraints

The simplest class of extended gauge models depend on the extra parameters λ, θ_1 and $M_{Z'}$ [1]. In Figs. 5, 6 we plot for $m_H = 100, 1000$ GeV, respectively, and $m_t = 125$ GeV the $M_{Z'}$ lower bound (90% c.l. for one variable) for this class of models. Comparing Fig. 5 with Fig. 5 of Ref. [1] we can see the effect on the $M_{Z'}$ lower bound of the improved parity-violation data and the last year LEP results (in particular Z widths and asymmetries). The net result is to increase the $M_{Z'}$ bound by ~ 200 GeV. In Figs. 7a-e we plot the allowed regions (90% c.l. for two variables) by all data in the $s_3 - M_{Z'}$ plane, where s_3 measures the $Z^0 Z'$ mixing, for $m_{(t,H)} = (100 \text{ and } 130, 100 \text{ and } 1000) \text{ GeV}$ and for the illustrative (popular) models $\chi, LR, \psi, \eta, \chi'$, respectively. (See Ref. [1] for definitions. We must emphasize that taking s_3 free we are implicitly assuming an arbitrary Higgs doublet content. Then our equations which only include the minimal standard model radiative corrections are only approximate.) Comparing the $m_{t,H} = 100$ GeV case with Figs. 7a-e of Ref. [1] we see that the inclusion in particular of LEP data drastically improves the Z' limits. As stressed in Refs. [6, 7, 8], this is due to the sensitivity of LEP data to s_3 , for in the Z peak new Z' contributions are proportional to s_3 . For the minimal case of one Higgs doublet s_3 and $M_{Z'}$ are not independent and the present bounds

correspond to the intersection of the different curves with the dashed line. The $M_{Z'}$ values in this case are comparable to those of Figs. 5, 6.

We may wonder about the implications of these bounds for future Z' searches. Obviously, more precise data will further constrain $s_3, M_{Z'}$ and the Z' couplings to fermions. About the possibility of directly detecting some effect, the minimal class of models is less promising, except for a large hadron collider [14]. (The reduction of the Z' bounds for large λ in Figs. 5, 6 is due to the corresponding reduction of the Z' couplings.) For a model with an arbitrary number of Higgs doublets (Fig. 7), $M_{Z'}$ and s_3 are independent parameters, allowing for the possibility of lower $M_{Z'}$ upper bounds for negligible s_3 , and then for direct production, for instance, at e^+e^- (500 GeV) for $M_{Z'} < 500$ GeV [15].

4 Correlated $M_{Z'} - m_t$ bounds

In Figs. 8, 9 we give the $m_t - M_{Z'}$ allowed regions (90% c.l. for two variables) for $m_H = 100$ and 1000 GeV for models χ and LR and for several λ, θ_1 values (models), respectively. Comparing them with Figs. 2, 3 of Ref. [1] we see the effect of the improved parity-violation data and the last year LEP results on the $m_t - M_{Z'}$ bounds.

A global fit takes into account all the experimental data. It is also interesting, however, to look for observables which measure the unknown parameters. Two such observables, D and M , are particularly adequate to measure the presence (mass) of a new Z' and the mass of the top, respectively [5],

$$D = \frac{6}{23} \left(\frac{\Gamma_h - \Gamma_b}{4\Gamma_e} - \frac{26}{27} A_{FB}^b \right), \quad (4.1)$$

$$M = \frac{3}{13} \left(\frac{\Gamma_b}{\Gamma_e} - \frac{26}{27} A_{FB}^b \right), \quad (4.2)$$

where $\Gamma_{h,b,e}$ are the Z partial widths into hadrons, bottom quarks and electrons, respectively, and A_{FB}^b is the bottom forward-backward asymmetry. In Fig. 10 we plot the curves of constant D and M for models χ and LR and $m_H = 100, 1000$ GeV. Overimposing them to Figs. 8, 9 we can see the present limits on D and M , as well as the required precision on their determination to obtain new information. D and M in Fig. 10 incorporate the one-loop radiative corrections of the minimal standard model and the tree-level Z' contributions (see the Appendix).

5 Conclusions

This paper presents a global fit to the minimal standard model and its simplest gauge extension. The standard model contributions to the different observables include radiative corrections but the new Z' contributions are only considered at tree-level [4]. In a previous paper we studied the constraints resulting from neutral current data and gauge boson masses [1]. In this one we update them taking into account the improved parity-violation data and including the last year LEP results [2, 3].

In the minimal standard model the largest uncertainty to determine m_t results from our ignorance on m_H . We find that the experimental error and the error associated to m_H are comparable. The error due to the variation of α_s is of the order of 1/2 of the experimental error. For $\alpha_s(M_Z) = 0.12$ and $m_H = 100 \text{ GeV}$ [$m_H = 1 \text{ TeV}$] (90% *c.l.* for one variable) we find

$$m_t = 112^{+28}_{-23} \text{ GeV} \quad [m_t = 140^{+25}_{-27} \text{ GeV}] , \quad (5.1)$$

while $m_t = 112^{+22}_{-23} \text{ GeV}$ [$m_t = 140^{+19}_{-21} \text{ GeV}$] for 1 σ . In the minimal gauge extension the m_t limits are largely unchanged but correlated to the $M_{Z'}$ bounds. These are typically of the order of $\sim 700 \text{ GeV}$. The improvement in the bound (with respect to our previous analysis) is $\sim 200 \text{ GeV}$ for models with one Higgs doublet, varying with the model. For models with an arbitrary number of Higgs doublets the improvement is $\sim 50 \text{ GeV}$ with smaller $M_{Z'}$ bounds. In this case the limit on the $Z^0 Z'$ mixing is reduced by a factor of order ~ 4 .

The detailed numerical conclusions are model dependent and given in the figures. Other non-minimal E_6 models could be considered, and the results would be similar [6, 7, 8, 14, 15]. Due to the very good agreement between the data and the standard model, and the already stringent bounds on new Z' , any new improvement in the data (assuming no apparent departure of the standard model predictions) will translate into a moderate improvement in the Z' bounds (*proportional* to the improvement in the experimental error). A large hadron collider will allow for the final test for a very heavy Z' ($\sim$ several TeV).

Acknowledgements

We thank the Institute for Theoretical Physics at Santa Barbara for the invitation to participate at the *Workshop on Precise Electroweak Measurements* where part of this work was done.

Appendix

A LEP observables in extended electroweak models

To complete the Appendix of Ref. [1] we must add the expressions for the Z partial widths. These are obtained replacing the couplings v_f and a_f

$$\begin{aligned} v_f^{Z^0} &= I_{3f} - 2Q_f s_W^2 \\ a_f^{Z^0} &= I_{3f} \end{aligned} \quad (\text{A.1})$$

$$\begin{aligned} g_V^f &= v_f^{Z^0} c_3 - v_f^{Z'^0} s_3 \\ g_A^f &= a_f^{Z^0} c_3 - a_f^{Z'^0} s_3 \end{aligned} \quad (\text{A.2})$$

where [17]

$$\begin{aligned} v_u^{Z'^0} &= \frac{5}{6} s_1 c_1 s_W (\lambda - \frac{1}{\lambda}) \\ a_u^{Z'^0} &= -\frac{1}{2} s_1 c_1 s_W (\lambda - \frac{1}{\lambda}) - \frac{2}{3} c_2 (\frac{c_2^2}{\lambda} + \lambda s_1^2) s_W \\ v_d^{Z'^0} &= -\frac{1}{2} s_1 c_1 s_W (\lambda - \frac{1}{\lambda}) - \frac{1}{2} (\frac{c_2^2}{\lambda} + \lambda s_1^2) (c_2 + \sqrt{5/3} s_2) s_W \\ a_d^{Z'^0} &= \frac{1}{2} s_1 c_1 s_W (\lambda - \frac{1}{\lambda}) + \frac{1}{2} (\frac{c_2^2}{\lambda} + \lambda s_1^2) (-\frac{1}{3} c_2 + \sqrt{5/3} s_2) s_W \\ v_e^{Z'^0} &= -2v_u^{Z'^0} - v_d^{Z'^0} \\ a_e^{Z'^0} &= a_d^{Z'^0} \\ v_\nu^{Z'^0} &= -v_d^{Z'^0} - \frac{1}{2} v_u^{Z'^0} + \frac{1}{2} a_u^{Z'^0} \\ a_\nu^{Z'^0} &= v_\nu^{Z'^0} \end{aligned} \quad (\text{A.3})$$

On the other hand, the vector to axial ratio reads ³

$$\frac{g_V^f}{g_A^f} (M_Z) = \frac{(I_{3f} - 2Q_f s_f^2) c_3 - v_f^{Z'^0} s_3}{I_{3f} c_3 - a_f^{Z'^0} s_3} \quad (\text{A.4})$$

where s_f^2 is the corrected electroweak mixing angle [18].

²Note the different definition of the vector and axial couplings used here.

³Note that radiative corrections proportional to the $Z^0 Z'$ mixing have been neglected.

References

- [1] F. del Aguila, J.M. Moreno and M. Quirós, *Nucl. Phys. B* **361** (1991) 45.
- [2] M.C. Noecker, B.P. Masterson and C.E. Wieman, *Phys. Rev. Lett.* **61** (1988) 310; S.A. Blundell, W.R. Johnson and J. Sapirstein, *Phys. Rev. Lett.* **65** (1990) 1411; V. Dzuba, V. Flambaum and O. Sushkov, *Phys. Lett.* **141A** (1989) 147.
- [3] G.J. Bobbink, in *XXV Rencontres de Moriond*, Les Arcs, France, March 10-17, 1991; P. Ratoff, in the same Conference.
- [4] G. Degrassi and A. Sirlin, *Phys. Rev. D* **40** (1989) 3066.
- [5] F.M. Renard and C. Verzegnassi, *Phys. Lett.* **217B** (1989) 199; **260B** (1991) 225; A. Djouadi et al., *Nucl. Phys. B* **349** (1991) 48; W. Hollik, Max-Planck preprint, MPI-Ph/26-91 (1991).
- [6] K.T. Mahanthappa and P.K. Mohapatra, *Phys. Rev. D* **43** (1991) 3093; G. Altarelli et al., CERN preprints, CERN-TH. 6028/91; CERN-TH. 6051/91; see also, G. Altarelli, R. Barbieri and S. Jadach, CERN preprint, CERN-TH. 6124/91 (1991).
- [7] M.C. González-García and J.W.F. Valle, *Phys. Lett.* **259B** (1991) 365.
- [8] J. Layssac, F.M. Renard and C. Verzegnassi, LAPP preprint LAPP-TH-290/90 (1990) and updated version, January 1991; for a general discussion, see: P. Langacker, M. Luo and A.K. Mann, Pennsylvania preprint, UPR-0458T (1991).
- [9] CDF Collaboration, F. Abe et al., FERMILAB preprint, FERMILAB-PUB 90-1373 (1990).
- [10] J. Ellis and G.L. Fogli, *Phys. Lett.* **249B** (1990) 543.
- [11] ALEPH Collaboration, D. Decamp et al., CERN preprint PPE/91-019 (1991).
- [12] Particle Data Group, *Review of Particle Properties*, *Phys. Lett.* **239B** (1990) 1.
- [13] CDF Collaboration, F. Abe et al., *Phys. Rev. Lett.* **65** (1990) 2243; UA2 Collaboration, J. Alitti et al., *Phys. Lett.* **241B** (1990) 150.

Table Captions

Table 1 Experimental values of the fitted quantities.

- [14] P. Chiappetta et al., *New Vector Bosons*, in *Large Hadron Collider Workshop*, Aachen 4-9 October 1990, ECFA 90-133, Vol. II, p. 685.
- [15] C. Verzegnassi et al., *Extended Gauge Theories Working Group*, in *e^+e^- Collisions at 500 GeV*.
- [16] W. Hollik, *Fortschr. Phys.* **38** (1990) 3.
- [17] F. del Aguila, M. Quiros and F. Zwirner, *Nucl. Phys. B* **287** (1987) 419.
- [18] M. Böhm and W. Hollik, *Forward-Backward Asymmetries*, in *Z Physics at LEP 1*, CERN 89-08 (Ed. G. Altarelli, R. Kleiss and C. Verzegnassi), Vol. 1, p. 203; W. Hollik, in *XIII Warsaw Symposium on Elementary Particles*, Kazimierz 1990 (Ed. Z. Ajduk, S. Pokorski and A. Wróblewski).

Figure Captions

- Fig. 1a Allowed region (90% c.l.) in the $m_t - s_W^2$ plane for the standard model, by νq data, by Q_W , by M_Z and by ALL data, for $m_H = 100$ GeV.
- Fig. 1b The same as in Fig. 1a, but for $m_H = 1$ TeV.
- Fig. 2a The same as in Fig. 1a but by all neutral current (NC) data, by M_W , by the Z partial widths and asymmetries ($T + A$), by M_Z and by ALL data.
- Fig. 2b The same as in Fig. 2a, but for $m_H = 1$ TeV.
- Fig. 3a Variation with α_s of the allowed region (90% c.l.) in the $m_t - s_W^2$ plane, by the Z partial widths and asymmetries and by ALL data (the M_Z band does not change), for $m_H = 100$ GeV.
- Fig. 3b The same as in Fig. 3a, but for $m_H = 1$ TeV.
- Fig. 4a χ^2 distribution as a function of m_t for different data sets and α_s values, and $m_H = 100$ GeV.
- Fig. 4b The same as in Fig. 4a, but for $m_H = 1$ TeV.
- Fig. 5a Tridimensional plot of the M_Z lower bound (90% c.l. for one variable) from all data for $m_t = 125$ GeV and $m_H = 100$ GeV, as a function of λ and θ_1 .
- Fig. 5b Contours of constant M_Z from Fig. 5a.
- Fig. 6a The same as in Fig. 5a, but for $m_H = 1$ TeV.

Fig. 6b Contours of constant $M_{Z'}$ from Fig. 6a.

Fig. 7a Allowed region (90% c.l.) by all data in the $s_3 - M_{Z'}$ plane for $(m_t, M_H) = (100 \text{ GeV}, 100 \text{ GeV})$, $(130 \text{ GeV}, 100 \text{ GeV})$, $(100 \text{ GeV}, 1 \text{ TeV})$ and $(130 \text{ GeV}, 1 \text{ TeV})$, for the χ model [1]. The curve $s_3(M_{Z'})$ for models with minimal Higgs doublet content (dashed line) is also given.

Fig. 7b The same as in Fig. 7a, but for the LR model [1].

Fig. 7c The same as in Fig. 7a, but for the ψ model [1].

Fig. 7d The same as in Fig. 7a, but for the η model [1].

Fig. 7e The same as in Fig. 7a, but for the χ' model [1].

Fig. 8 Allowed region (90% c.l.) by all data in the $m_t - M_{Z'}$ plane for models χ and LR, and for $m_H = 100$ (1000) GeV, dashed (solid) line.

Fig. 9a The same as in Fig. 8, but for the illustrative set of (λ, θ_1) values: $(1.5, \pi/2)$, $(1.2, \pi/2)$, $(1.2, 0)$, and $(1.5, 0)$.

Fig. 10a Contours of constant D and M in the $M_{Z'} - m_t$ plane for the χ model and $m_H = 100 \text{ GeV}$.

Fig. 10b The same as in Fig. 10a, but for $m_H = 1 \text{ TeV}$.

Fig. 10c The same as in Fig. 10a, but for the LR model.

Fig. 10d The same as in Fig. 10c, but for $m_H = 1 \text{ TeV}$.

Quantity		Experimental Value
νq	g_L^2	0.2977 ± 0.0042
	g_R^2	0.0317 ± 0.0034
	θ_L	2.50 ± 0.03
	θ_R	$4.59^{+0.44}_{-0.27}$
$\nu_\mu e$	g_A^e	-0.513 ± 0.025
	g_V^e	-0.045 ± 0.022
eH	C_{1u}	-0.253 ± 0.071
	C_{1d}	0.391 ± 0.064
	$C_{2u} - \frac{1}{2}C_{2d}$	0.22 ± 0.36
	$Q_W(^{133}\text{Cs})$	-71.04 ± 1.81
W	M_W	$80.1 \pm 0.3 \text{ GeV}$
Z	M_Z	$91.174 \pm 0.020 \text{ GeV}$
	Γ_Z	$2485 \pm 9 \text{ MeV}$
	Γ_h	$1740 \pm 9 \text{ MeV}$
	Γ_t	$83.31 \pm 0.40 \text{ MeV}$
	$\frac{s_W^2}{s_A^2}(M_Z)$	0.0052 ± 0.0016

Table 1

Fig. 1a

Fig. 1b

Fig. 2a

Fig. 2b

Fig. 3a

Fig. 3b

Fig. 4a

Fig. 4b

Fig. 5a

Fig. 5b

Fig. 6a

Fig. 6b

Fig. 7a

Fig. 7b

Fig. 7c

Fig. 7d

Fig. 7e

Fig. 8

Fig. 9a

Fig. 9b

Fig. 10a

Fig. 10b

Fig. 10c

Fig. 10d