The Compact Muon Solenoid Experiment
Conference Report

Mailing address: CMS CERN, CH-1211 GENEVA 23, Switzerland

13 February 2018 (v3, 22 March 2018)

Highlights from CMS

Christian Autermann
on behalf of the CMS Collaboration

Abstract

This article summarizes the latest highlights from the CMS experiment as presented at the Lepton Photon conference 2017 in Guangzhou, China. A selection of the latest physics results, the latest detector upgrades, and the current detector status are discussed. CMS has analyzed the full dataset of proton-proton collision data delivered by the LHC in 2016 at a center-of-mass energy of 13 TeV corresponding to an integrated luminosity of 40 fb^{-1} . The leap in center-of-mass energy and in luminosity with respect to the 7 and 8 TeV runs enabled interesting and relevant new physics results. A new silicon pixel tracking detector was installed during the LHC shutdown 2016/17 and has successfully started operation.

Presented at *LP2017 The 28th International Symposium on Lepton Photon Interactions at High Energies*

Recent Highlights from the CMS Experiment

Christian Autermann

on behalf of the CMS Collaboration

I. Phys. Inst. B, RWTH Aachen University

Sommerfeldstr. 14, 52074 Aachen, Germany

E-mail: christian.autermann@cern.ch

This article summarizes the latest highlights from the CMS experiment as presented at the Lepton Photon conference 2017 in Guangzhou, China. A selection of the latest physics results, the latest detector upgrades, and the current detector status was shown. CMS has analyzed the full dataset of proton-proton collision data delivered by the LHC in 2016 at a center-of-mass energy of 13 TeV corresponding to an integrated luminosity of 40 fb^{-1} . The leap in center-of-mass energy and in luminosity with respect to the 7 and 8 TeV runs enabled interesting and relevant new physics results. A new silicon pixel tracking detector has been installed during the LHC shutdown 2016/17 and has successfully started operation.

Keywords: CMS; LHC; Highlights; Upgrades; Detector; 13 TeV

1. Introduction

The CMS detector¹ is a multipurpose detector located at the LHC accelerator at CERN. The LHC started operation in 2010 and has delivered until the end of 2012 proton-proton collisions at center-of-mass energies of 7 TeV and 8 TeV corresponding to a total integrated luminosity of 5 fb^{-1} and 20 fb^{-1} , respectively. In 2015 the operations at 13 TeV started and delivered about 4 fb^{-1} . The dataset delivered in 2016 at 13 TeV corresponding to a luminosity of 40 fb^{-1} is used for the majority of physics results discussed here.

CMS is characterized by the large superconducting solenoid magnet with 6 m inner diameter providing a magnetic field of 3.8 T. Inside this magnet the silicon pixel and strip tracking detectors and the calorimeter systems are installed. The magnet itself is located within the iron return yoke, which houses also the muon system of drift tube and resistive plate chambers. The tracking system is the largest silicon detector of any high energy detector and allows excellent particle momentum measurement at high efficiency and resolution. In particular the innermost pixel vertex detector enables precise b jet and τ tagging. The calorimeters are divided into the electromagnetic (ECAL) and hadronic calorimeters (HCAL). The ECAL is made out of dense but clear and scintillating lead tungstate crystals and silicon avalanche photodiodes for readout. The HCAL is a sampling calorimeter using brass and steel plates as absorbers and plastic scintillators as active material. A particle-flow (PF) event reconstruction algorithm is used to identify and reconstruct any particle of a proton–proton collision event, using the best possible combination of all detector subsystems. The PF event reconstruction leads to an improved

performance for the reconstruction of jets and missing transverse momentum p_T^{miss} , and for the identification of electrons, muons, and taus.

2. Status of the CMS detector

The CMS status² during the successful data-taking period 2016, the upgrades during the extended end-of-year technical stop 2016/17, and the startup in spring 2017 will be discussed briefly in the following.

2.1. Detector upgrades

Several upgrades were installed during the technical stop 2016/17. The main changes were the improvements of the hadronic forward calorimeter readout, the installation of a muon-endcap GEM prototype, and most importantly, of the new pixel vertex detector.

The readout electronics of the hadronic forward calorimeter were upgraded in order to make full use of the photomultiplier tubes (PMTs) replaced in the long shutdown of 2013. The PMTs read out the Cherenkov light produced in quartz fibers embedded in the iron absorbers of the hadronic calorimeter. However, occasionally charged particles produce Cherenkov light directly in the PMT window mimicking high-energetic particles. The new PMTs have 1/6 windows thickness, twice the quantum efficiency and gain, and dual readout that required the electronics to be upgraded.

Gas electron multiplier (GEM) detectors are being considered for an upgrade of the CMS muon system. A GEM demonstrator was installed in the forward high-eta region to test the system during LHC run conditions. GEMs are micro-pattern gaseous detectors that are expected to improve muon tracking and triggering utilizing a good spatial and time resolution, high efficiency, and radiation hardness.

The silicon pixel tracker is the innermost CMS sub-detector and crucial for precise track reconstruction and for vertex reconstruction with high resolution. The pixel tracker was replaced during the end-of-year shutdown 2016/17 due to radiation damage and significant data losses at instantaneous luminosity larger than $10^{34} \text{ cm}^{-2}\text{s}^{-1}$. The maximum LHC instantaneous luminosity is routinely exceeding this threshold by a factor of two, corresponding to a mean number of primary vertices of 50, causing too high occupancy in the old pixel readout chip. The upgraded pixel detector uses a digital readout chip with higher rate capability. The main new features are four instead of three barrel layers and three instead of two end-cap disks. The mechanical structure is designed to be very light and a new powering scheme and CO₂ cooling system is used in order to further reduce the amount of material.

Operation of the newly installed pixel detector has started. Layer 1 and 2 share a common programmable time delay. The time alignment was difficult to achieve due a faster Layer 1 readout chip. An optimal common plateau of efficiency with values

close to 99% for all pixel layers and disks at luminosities $\mathcal{L} = 1.6 \cdot 10^{34} \text{ cm}^{-2}\text{s}^{-1}$ as shown in Fig. 1 has been established. The timing is chosen to favor the Layer 1 performance. Although not yet at the ultimate detector performance, more complex functions like vertexing, b-tagging, and high-level trigger electron reconstruction are significantly better than with the old detector, which would not have been able to cope with the rates in the first place.

2.2. Detector performance

The LHC restarted collisions after the end-of-year technical stop in 2017 very successfully, delivering fast peak instantaneous luminosities near the design goal of $\mathcal{L} = 2 \cdot 10^{34} \text{ cm}^{-2}\text{s}^{-1}$ as shown in Fig. 2. Until the Lepton-Photon conference already 9 fb^{-1} of data at a center-of-mass energy of 13 TeV had been delivered³. While this 2017 data set is not yet used for physics analyses, the detector performance studies discussed in this section, in particular of the recently installed upgrades make use of this data.

The CMS Level 1 trigger system reduces the event rate from the proton bunch crossings rate at 40 MHz down to 100 kHz without detrimental impact on the physics acceptance, in particular for measurements and searches at the electroweak scale. The trigger system was upgraded during the long shutdown 1 (LS1) in 2013/14 to cope with the increased rates due to the larger instantaneous luminosity and the increase in center-of-mass energy. A significant increase in trigger threshold with negative impact on the physics program was avoided. The performance of the level 1 electron or photon and muon trigger in $\sqrt{s} = 13 \text{ TeV}$ data was measured⁴ and is shown exemplary in Fig. 3 (left) for isolated electron or photon candidates. The efficiency was measured using the tag&probe method on $Z \rightarrow ee$ events. The

Fig. 1. Hit efficiencies per layer of the upgraded CMS pixel detector in 2017

Fig. 2. Delivered and recorded instantaneous (top) and integrated luminosity (bottom) to CMS for the proton-proton runs since 2010³.

higher 2017 trigger efficiency is due to an improved relaxation scheme of the isolation selection as a function of E_T of the e/γ candidate. By recalibrating the calorimeter level-1 object also the transverse energy resolution of L1 e/γ candidates with respect to the offline reconstructed E_T could be improved in 2017, as shown in Fig. 3 (right).

To maintain good quality track reconstruction with high resolution for charged particles the position and the orientation of the silicon pixel and strip modules need to be known with a precision of several micrometers. The tracker alignment follows a global fit approach using Millepede-II and a local fit approach using the HipPy fit⁵. For the 2016 data taking, the modul-level alignment was obtained from cosmic-ray data collected before and a small amount of pp collision data. Time-dependent movements of the pixel detector large-scale structures were monitored. The tracker alignment in data taking, shown as red histogram in Fig. 4 (left), was corrected if necessary by an automated prompt calibration loop, as shown in Fig. 4 (right).

The missing transverse momentum p_T^{miss} is a measure of the momentum imbalance caused by undetectable particles such as neutrinos and is crucial for high precision measurements and in particular for searches for new physics with undetectable dark matter candidate particles. On the other hand, because the missing transverse momentum is calculated using the vectorial sum of all energy depositions reconstructed in the detector, it is a quantity that is highly sensitive to

Fig. 3. L1 electron or photon candidate trigger efficiency versus the transverse energy for an average of 24 simultaneous interactions (left) and the L1 E_T resolution with respect to the offline reconstruction⁴ (right).

Fig. 4. Alignment of the pixel detector. The median residuals of the y -positions of all pixel barrel detector modules are shown for simulation, during data taking, and for the end-of-year alignment data (left). The movements of the pixel barrel detector in y -direction for the second half of the 2016 data taking are shown (right). The grey bands indicate runs during which the CMS magnet was not at 3.8 T. Vertical dashed lines illustrate updates of the pixel reference geometry, after which mis-alignments are cured⁵.

the precision of the energy measurement in all sub-detectors, in particular of the calorimeter energy resolution. The missing transverse momentum performance and its dependence on the simultaneous luminosity, i.e., the amount of simultaneous proton-proton interactions called pileup, is studied using $Z \rightarrow ll$ events⁶. The well reconstructed leptonically decaying Z boson allows to probe the detector response of the hadronic system, and therefore to measure the scale and the resolution of the missing transverse momentum. The hadronic recoil is projected onto the axis of the well measured Z boson transverse momentum and its parallel and perpen-

Fig. 5. Comparison of resolution of parallel (left) and perpendicular (right) components of hadronic recoil measured using jet-energy-scale corrected PF p_T^{miss} (red markers) and corrected PUPPI (blue markers) as a function of number of vertices⁶.

dicular components are used to study the performance of p_T^{miss} as shown in Fig. 5 for particle-flow and „pileup per particle identification” (PUPPI) corrected p_T^{miss} .

3. Recent CMS physics highlights

The LHC started operations at a center-of-mass energy of 13 TeV in 2015 delivering an integrated luminosity of 3 fb⁻¹. In 2016 the 13 TeV data run continued, CMS recorded 36 fb⁻¹ of data, almost twice the amount of data collected before the long shutdown in 2012 at 8 TeV corresponding to an integrated luminosity of 20 fb⁻¹, as shown in Fig. 2. In combination with the leap in center-of-mass energy the 2016 data set therefore provides the basis for significant updates of physics analyses in terms of precision, sensitivity, and reach. The physics results discussed in the following are selected highlights of recent publications typically based on this data set, significantly extending and exceeding previous CMS results.

3.1. QCD multijet measurements

QCD multijet production processes dominate the spectrum of physics produced at the LHC and are therefore substantial standard model backgrounds to many precision measurements and searches for new physics. Direct measurements of QCD multijet production can provide valuable information for physics at higher energy scales, for example by extracting the energy scale depending strong coupling α_s ⁷ allowing for more precise fits to higher scales, as shown in Fig. 6.

The differential QCD cross-section measurement in dijet event topologies as a function of jet masses and transverse momenta is sensitive to QCD parton showering and in particular relevant for searches for new physics with boosted objects. The dijet topology was unfolded to correct for the effect of the detector reconstruction

Fig. 6. The running of the strong coupling parameter α_s and its uncertainty⁷. Previous measurements from the H1, ZEUS, and D0 experiments at the HERA and Tevatron colliders are shown together with recent CMS measurements.

using 2.3 fb^{-1} of 13 TeV data⁷. Jet grooming algorithms were used to remove low energy portions from the jets arising from soft radiations that are difficult to model. The Monte Carlo simulation of the jet mass spectrum was found to be improved for groomed jets, as shown in Fig. 7.

3.2. Electroweak production

The effective weak mixing angle was measured in 18.8 fb^{-1} of 8 TeV data using the forward-backward asymmetry of Drell-Yan events in the dielectron and dimuon final state⁸. The forward-backward asymmetry A_{FB} is defined as $A_{\text{FB}} = (\sigma_{\text{F}} - \sigma_{\text{B}})/(\sigma_{\text{F}} + \sigma_{\text{B}})$, where σ_{F} and σ_{B} are the cross sections in the forward and backward hemispheres in the Collins-Soper frame of the dilepton system. The measured asymmetry A_{FB} , as shown in Fig. 8 (left), allows to extract the most precise measurement of leptonic effective weak mixing angle $\sin^2 \theta_{\text{eff}}^{\text{lept}} = 0.23101 \pm 0.00052$ at the LHC. Uncertainties of the proton density functions (PDFs) translate into sizable variations of A_{FB} . Since the PDF uncertainties affect A_{FB} and $\sin^2 \theta_{\text{eff}}^{\text{lept}}$ differently, a Bayesian χ^2 reweighting method can be applied to constrain the PDF uncertainties and to reduce the uncertainty of the extracted value of $\sin^2 \theta_{\text{eff}}^{\text{lept}}$. The CMS measurement precision is comparable to Tevatron measurements and is compared in Fig. 8 (right) to other LHC, Tevatron, and LEP results.

The CMS collaboration has measured a wide range of standard model production cross sections, branching ratios, particle masses, and couplings. In Fig. 9 a selection of electroweak, top quark, and Higgs boson production cross section measurements is summarized. Even without QCD multijet processes cross sections over nine orders of magnitude have been measured and compared to theory predictions

Fig. 7. Differential dijet cross section unfolded to particle level as a function of the jet mass⁷.

Fig. 8. Measurement of the forward-backward asymmetry of Drell-Yan events (left) and the extracted electroweak mixing angle fitted to several previous and current analyses (right)⁸.

for the $\sqrt{s} = 7$ TeV, 8 TeV, and 13 TeV runs. A remarkable agreement is observed. The consistency of experimental observations and theoretical predictions over this wide range of cross sections, production mechanisms, and final states illustrates nicely the excellent control over the data reconstruction, the precision of theoretical calculations, and in result the yet unchallenged validity of the standard model at

Fig. 10. Top production cross sections depending on the center of mass energy for $t\bar{t}$ pair production (left) and single- t (right)².

- Recently the cross section of top quark pair production in association with a W or Z boson has been measured in the 13 TeV data set collected in 2016⁹. The measurement was performed in the same-sign dilepton, three- and four-lepton final states where the jet and b-tagged jet multiplicities were exploited to enhance the signal-to-background ratio. The resulting cross section is included in the summary plot in Fig. 9.
- One of the most promising channels for the direct measurement of the Higgs-top quark Yukawa coupling strength is the standard model production of the Higgs boson in association with a $t\bar{t}$ pair, where the top quarks decay leptonically (e or μ) and the Higgs boson as $H \rightarrow b\bar{b}$. The inclusive top quark pair plus dijet $t\bar{t} + b\bar{b}$ and $t\bar{t} + jj$ production cross sections as well as their ratio $\sigma_{t\bar{t}b\bar{b}}/\sigma_{t\bar{t}jj}$ have been measured in dileptonic events using the first 2.3 fb⁻¹ of 13 TeV data¹⁰. The ratio in the full phase space was measured to be $\sigma_{t\bar{t}b\bar{b}}/\sigma_{t\bar{t}jj} = 0.022 \pm 0.007$ in agreement with the prediction from the standard model. The contribution from $Ht\bar{t}$ was found to be negligibly small.
- The measurement of the semileptonic $t\bar{t}\gamma$ production cross section¹¹ is directly probing the coupling of the top quark and the photon. A deviation between the measurement and the standard model prediction could hint at exotic quarks with anomalous electric charge or electric dipole moment. The fiducial cross section for associated $t\bar{t}$ and photon production per semileptonic final state was found to be 127 ± 27 fb in agreement with standard model predictions.
- The single top quark production cross section in association with a Z boson in final states with three leptons (e or μ) was measured to 10^{+8}_{-7} fb at a significance of 2.4σ in the 8 TeV data set corresponding to 19.7 fb⁻¹ of integrated luminosity. This measurement allows to constrain flavor-changing neutral current interactions to $\mathcal{B}(t \rightarrow Zu) < 0.022\%$ and $\mathcal{B}(t \rightarrow Zc) < 0.049\%$ at 95% confidence level¹².

Fig. 11. Differential $t\bar{t}$ cross section measurements in missing transverse momentum (left) and in the jet multiplicity variable (right) in 35.9 fb^{-1} of 13 TeV data compared to different Monte Carlo simulations¹³.

Differential cross section measurements can provide highly sensitive tests to verify theoretical models of standard model processes. These precision tests allow to tune Monte Carlo generators and to improve for example the standard model background estimation relevant for search analysis. The top differential cross section in the semileptonic channel with one electron or muon has been measured with respect to variables that do not require to reconstruct the $t\bar{t}$ system¹³. The 13 TeV data collected in 2016 corresponds to an integrated luminosity of 35.9 fb^{-1} and to about $3 \cdot 10^7$ $t\bar{t}$ pairs that are used for this analysis. The resulting differential cross sections are unfolded to particle level in a phase space resembling the fiducial volume of CMS and are compared to state-of-the-art leading order and next-to-leading order $t\bar{t}$ simulations, as shown in Fig. 11.

One important example for top quark property measurements is the measurement of the top mass. Plots summarizing the various top quark mass measurement analyses at CMS in the traditional and in alternative channels using data recorded at $\sqrt{s} = 7, 8$ and 13 TeV are shown in Fig. 12. It is remarkable that the top mass measurement in the single semileptonic decay channel using 19.7 fb^{-1} of 8 TeV data exceeds the precision of the Tevatron combination. Worth mentioning is also the compatible precision of the various alternative channels on the top quark mass.

3.4. Higgs physics

The Higgs boson is the last particle discovered by the ATLAS and CMS experiments in 2012 and completes the standard model of particle physics. The discovery channels are the fully reconstructible $H \rightarrow \gamma\gamma$ diphoton and $H \rightarrow l^+l^-l^+l^-$ four lepton (electron or muon) decay channels, where the Higgs boson decays via a charged par-

Fig. 12. Summary plot of the top mass measurements in the classic channels (left) and in alternative channels with compatible sensitivity (right)¹⁴.

title loop or two Z bosons, respectively. Despite the small branching fractions of $\text{BF}_{\gamma\gamma} = 0.23\%$ and $\text{BF}_{4l} = 0.013\%$, these channels offer a clear fully reconstructible invariant mass resonance signal over a continuous background and allow for measurements of the Higgs boson properties, such as the mass, spin, CP -parity, and couplings.

The analysis of the diphoton final state using 35.9 fb^{-1} of 13 TeV data has measured the Higgs signal strength relative to the standard model prediction for different production modes and effective couplings to fermions and bosons as well as to photons and gluons¹⁵. The results are in agreement with the expectations from the standard model, in particular the $Ht\bar{t}$ production was measured relative to the expectation to $\mu_{Ht\bar{t}} = 2.2^{+0.9}_{-0.8}$, corresponding to a 3.3σ excess with respect to the absence of $Ht\bar{t}$ production. The analysis of the four-lepton final state in the same data set lead to similar measurements of cross section and coupling strengths that were also found to be consistent with standard model expectations¹⁶. The decay width of the Higgs boson was measured to be $\Gamma_H < 1.10 \text{ GeV}$ at 95% confidence level.

A combined ATLAS and CMS analysis of the LHC pp collision data at $\sqrt{s} = 7$ and 8 TeV¹⁷ has allowed to measure the Higgs boson production and decay rates and to constrain the Higgs couplings. In particular the couplings to the top and bottom quarks, to the Z and W vector bosons, and the tau lepton have been measured and the coupling to muons has been constrained. Recently, individual

Fig. 13. Invariant dilepton mass for the $H \rightarrow W^+W^-$ analysis¹⁸ using 15.2 fb⁻¹ of 13 TeV data (left) and the distribution of the decimal logarithm of the ratio between the expected signal and the sum of expected signal and expected background in each bin of the mass distributions to extract the results, in all signal regions for the $H \rightarrow \tau\tau$ analysis¹⁹ using 35.9 fb⁻¹ of data (right).

analyses of the 13 TeV data in the $H \rightarrow W^+W^-$, $H \rightarrow \tau\tau$, and $H \rightarrow b\bar{b}$ channels have been published, as will be briefly discussed in the following.

The measurement of the Higgs to WW decay channel uses 15.2 fb⁻¹ of 13 TeV data recorded in 2015 and 2016¹⁸. The W^+W^- candidate events are selected by requiring an opposite charged $e\mu$ pair and large missing transverse momentum, as shown in Fig. 13 (left). The analysis categorized the selection depending on the Higgs production mode, i.e. gluon or vector boson (meaning in the following W or Z) fusion and vector boson associated production, distinguished by the jet and lepton multiplicity. The observed cross section times branching fraction is measured to be 1.05 ± 0.26 times the standard model prediction, corresponding to an observed significance of 4.3σ .

The first observation of the Higgs boson decay to a pair of tau leptons by a single experiment has been reported¹⁹. The 13 TeV data set corresponding to 35.9 fb⁻¹ has been analyzed simultaneously for gluon and vector boson fusion production categories as shown in Fig. 13 (right), using maximum likelihood fits in two-dimensional planes to extract the results. The production cross section times branching fraction was measured $1.09^{+0.27}_{-0.26}$ times the standard model expectation. Combining this measurement with previously published CMS results using the 7 and 8 TeV data sets leads to an observed significance of 5.9σ for the observation of the $H \rightarrow \tau\tau$ decay channel by the CMS experiment.

A recent CMS highlight result is the observed evidence for the Higgs boson decay to a bottom quark-antiquark pair using 35.9 fb⁻¹ of data recorded at 13 TeV²⁰. The $H \rightarrow b\bar{b}$ decay directly tests the coupling to fermions, i.e. to down-type quarks.

Fig. 14. Combination of all channels of the VH search into a single BDT distribution (left) and weighted invariant dijet mass distribution (right) for 35.9 fb^{-1} of data recorded at 13 TeV²⁰.

In spite of the large Higgs branching fraction of 58% into b quarks, this channels has not yet been established with observation level significance. The combined analysis of the 7 and 8 TeV data by the ATLAS and the CMS experiments observed this channel at a significance of 2.6σ . The Tevatron experiments CDF and D0 reported an excess of events at $m(H) = 125 \text{ GeV}$ corresponding to a local significance of 3.0σ for the combined searches, that were dominated by VH production and the $H \rightarrow b\bar{b}$ decay channel. Here, the same VH production channel is studied, where the Higgs boson decays to bottom quarks and the vector bosons W or Z decays fall into one of the five categories $Z \rightarrow \nu\nu$, ee , $\mu\mu$, and $W \rightarrow e\nu$, $\mu\nu$. The category with two charged leptons is further divided into a low and a high boson p_T region. For each of the seven signal selections boosted decision tree (BDT) discriminators are trained using a large number of kinematic variables, such as transverse momenta, invariant masses, angular distances, and b-tagging discriminants. Figure 14 (left) shows the gathered BDT output values for these seven channels and Fig. 14 (right) shows the dijet invariant mass distribution, combined for all channels, with all background processes other than VH and VZ subtracted and weighted by $s/(s+b)$, where s and b are the numbers of expected signal and background events in the bin of the BDT output distribution. The observed signal strength for $m(H) = 125.09 \text{ GeV}$ is $\mu = 1.2 \pm 0.4$ corresponding to an observed significance of 3.3σ . Combining this measurement with the 7 and 8 TeV data recorded by CMS yields a signal strength of $\mu = 1.06^{+0.31}_{-0.29}$ and observed and expected significances of 3.8σ .

3.5. Search for exotic new physics

The standard model of particle physics is until today successfully describing all physics processes studied at particle accelerator experiments. Also most recent re-

sults as discussed before are in good agreement with the standard model predictions. And yet, astrophysical observations like dark mass and dark energy, the observation of non-negligible neutrino masses, and theoretical considerations imply that the standard model cannot be complete or valid up to arbitrary high energy scales. The standard model does not offer a candidate dark matter particle and cannot describe quantum gravity effects that become relevant at the Planck scale Λ_P . An important objective of the LHC experiments is to search for physics beyond the standard model (BSM physics). Analyses of the 8 TeV data have been found to be in agreement with standard model expectations²¹. The sensitivity of searches for new physics profits greatly from the increased center-of-mass energy to 13 TeV and the large integrated luminosity of data taken during 2016.

A popular BSM theory is the scenario of extra dimensions that would lower the effective Λ_P scale, as predicted for example by the Arkani-Hamed–Dimopoulos–Dvali model. A non-resonant enhancement of dijet production through virtual graviton exchange leads to signatures differing from the predictions of QCD. The analysis of dijet angular distributions using 35.9fb^{-1} of 13 TeV data is sensitive to the graviton exchange, as well as to other BSM scenarios such as the production of quantum black holes, dark matter, and quark contact interaction models²². The differential dijet production cross section dominated by QCD Rutherford scattering is flat in $\chi_{\text{dijet}} = \exp |y_1 - y_2| = (1 + |\cos \theta^*|)/(1 - |\cos \theta^*|)$, where θ^* is the jet angle to the beam axis in the dijet rest frame. The χ_{dijet} analysis is complementary to dijet narrow bump searches and allows to set for the first time limits on the universal quark coupling to large dark matter mediator masses $2.5 < M_{\text{Med}} < 5$ TeV, as shown in Fig. 15.

3.6. Searches for supersymmetry

Supersymmetry (SUSY) is probably the most popular model of physics beyond the standard model. SUSY is the last possible space-time symmetry that can provide a dark matter candidate particle, if the R -parity is conserved and the lightest SUSY particle (LSP) is neutral. It provides a solution to the Higgs hierarchy problem by virtue of the symmetry between fermions and bosons. For each known standard model particle a new supersymmetric partner is predicted differing in spin by half a unit, and in mass. Since no SUSY particle has been discovered so far²³, the mass differences between the supersymmetric partners must be sufficiently large to escape detection by previous searches. The doubled luminosity of the data taken in 2016 and in particular the increased center-of-mass energy have favored the sensitivity of searches for supersymmetry. The jump in sensitivity and reach within a short time period will be unmatched in future, until a collider with significantly more energy will be build, as the LHC experiments will now only gradually increase their sensitivity to BSM physics with the amount of integrated luminosity. In the following, the most recent result of three different flavors of supersymmetry will be discussed.

Fig. 15. The 95% CL upper limits on the universal quark coupling g'_q as a function of the mass of a DM mediator that only decays to quarks from the CMS dijet narrow resonance search, boosted dijet resonance search, and the χ_{dijet} analysis²².

A search for supersymmetry, with gauge mediated SUSY breaking (GMSB), has been carried out using events with at least one photon, missing transverse momentum, and large hadronic event activity (H_T) in the transverse plane²⁴. R -parity conservation is assumed, so that the LSP, the almost massless Gravitino $\tilde{G}$, is stable. The final state signature in signal events is determined by the nature of the next-to-lightest SUSY particle, the NLSP. In the studied signal scenarios the NLSP is a gaugino that decays promptly as $\tilde{\chi}_1^{0/\pm} \rightarrow V\tilde{G}$ depending on its mixing, where V is a SM boson γ , Z^0 , $W^\pm$, or H . Different regions in the p_T^{miss} spectrum are used to normalize and validate the SM background estimation and to extract the signal acceptance, as shown in Fig. 16 (left). The analyses excludes squark masses up to 1.65 TeV and gluino masses up to 2.0 TeV, as compared in Fig. 16 (right) to the exclusion contours of a search for GMSB supersymmetry in events with at least one photon and missing transverse momentum²⁵ that is specialized for electroweak production of GMSB SUSY with little hadronic event activity.

CMS searches for electroweak production of charginos and neutralinos in several previously published final states using the full 13 TeV data set recorded in 2016 and corresponding to 35.9 fb^{-1} have been statistically combined, including a new analysis of the final state with three or more charged leptons (e or μ)²⁶. Strongly coupling SUSY particles might be too massive to be directly produced at the LHC, motivating direct searches for the electroweakly produced gauginos. Assuming R -parity conservation and depending on the production channel, an abundance of SM bosons is produced through cascade decays of the heavy gauginos into the LSP. The analyses have statistically independent signal regions $1l2b$, $4b$, $2l$ on- Z , $2l$ soft, $\geq 3l$,

Fig. 16. Missing transverse momentum distribution as observed in the search for gauge mediated supersymmetry breaking in final states with at least one photon and jets²⁴ (left). The resulting exclusion contour for in a simplified signal scenario of gluino pair production and bino-like neutralino decays to photons and gravitinos is compared to a contour from the analysis with at least one photon and p_T^{miss} but without jet requirements²⁵ (right).

and $H(\gamma\gamma)$ targeting final states with two SM bosons WZ , WH , ZZ , ZH , or HH . The 4b search drives the exclusion at large branching fraction values of $\tilde{\chi}_1^0 \rightarrow H\tilde{G}$ while the on-Z dilepton search and the multilepton search are competing at lower branching fraction values. The search with three or more charged leptons targets $\tilde{\chi}_1^\pm \tilde{\chi}_2^0$ production, where the mass difference between $\tilde{\chi}_1^0$ and $\tilde{\chi}_2^0$ is approximately equal to the Z boson mass. The statistical combination of all final states is able to exclude for chargino-neutralino production $m(\tilde{\chi}_1^\pm) = m(\tilde{\chi}_2^0)$ up to 650 GeV. Similar limits are derived for electroweak production in the GMSB scenario²⁶.

Most searches for supersymmetry assume the R -parity to be conserved, causing the lightest SUSY particle to be stable. Typically the LSP is the neutral neutralino $\tilde{\chi}_1^0$, the sneutrino $\tilde{\nu}$, or the Gravitino $\tilde{G}$ that are dark matter candidate particles. A common feature of R -parity conserving SUSY signal events is missing transverse momentum, as the dark matter candidate particles do not interact with the detector. However, there is no fundamental theoretical reason for R -parity conservation. In particular in the light of the most recent increasingly stringent limits on SUSY particle masses that challenge „natural SUSY” scenarios, R -parity violation (RPV) can ease this tension, as the limits obtained from R -parity conserving analyses do not generally apply here, because depending on the RPV coupling, little or no p_T^{miss} is created in the signal events. Baryon and meson lifetime measurements place strong indirect limits on the combination of two RPV couplings. Therefore, it is typically assumed that all RPV couplings except one are negligibly small or zero.

Minimal flavor-violating SUSY motivates the search for R -parity violating supersymmetry involving the third generation via the $U_i D_j D_k$ coupling λ_{ijk}'' ²⁷, where

Fig. 17. Observed exclusion contours at the 95% CL in the plane of the higgsino mass and the branching fraction $\tilde{\chi}_1^0 \rightarrow H\tilde{G}$ (left) and exclusion contours at the 95% CL in the plane of $m(\tilde{\chi}_1^0)$ and $m(\tilde{\chi}_2^0)$ for different assumed decay modes for the statistical combination of electroweak searches²⁶ (right).

Fig. 18. The b -tagged jet multiplicity of the R -parity violating search for SUSY in gluino pair production and prompt decays $\tilde{g} \rightarrow tbs$ using 35.9 fb^{-1} of 13 TeV data (left) and the corresponding cross section and resulting limit (right)²⁷.

the U_i and D_j are the up-type and down-type quark $SU(2)$ singlet fields. Gluinos can decay through λ''_{332} into third generation SM quarks, i.e. $\tilde{g} \rightarrow t\bar{t} \rightarrow tbs$. The single-lepton (e or μ) final state depending on the jet and b -tagged jet multiplicity, as shown in Fig. 18 (left), and the sum of masses of large-radius jets is analyzed for gluino pair-production that both decay promptly as $\tilde{g} \rightarrow tbs$. No p_T^{miss} requirement is applied, enabling the re-interpretation of this result in the context of other BSM scenarios with large jet or b -tagged jet multiplicities and little or no missing transverse momentum. An upper limit of approximately 10 fb at 95% confidence level is determined for gluino pair production, allowing to exclude gluinos below $m(\tilde{g}) < 1.61 \text{ TeV}$, as shown in Fig. 18 (right).

4. Conclusion and Outlook

The outstanding performance of the Large Hadron Collider and the CMS detector have allowed to collect a 13 TeV data set corresponding to an integrated luminosity of 40 fb^{-1} in 2016. The most recent result highlights derived using this data have been presented here. The leap in center-of-mass energy and the increase in integrated luminosity by a factor of two compared to the data recorded during the Run I have resulted in standard model measurements of excellent precision. For the first time strong evidence for the $H \rightarrow b\bar{b}$ decay channel has been observed at a single experiment. Searches for physics beyond the standard model are one of the main objectives of the LHC experiments that likewise have profited from the increase in center-of-mass energy in the new data set. The large amount of available data as well as the excellent performance of the detector make more complicated or specialized search analyzes possible and worthwhile.

During the end-of-the-year shutdown 2016/17 different detector upgrades were installed, in particular a new silicon pixel detector that has since then successfully started operation. The new pixel tracking and vertexing detector is able to cope with the large number of simultaneous interactions expected at the maximum design instantaneous luminosities of $\mathcal{L} = 2 \cdot 10^{34} \text{ cm}^{-2}\text{s}^{-1}$.

So far, only of the order of one percent of the total expected luminosity during the lifetime of the LHC has been recorded. The data that is yet to come will allow for many more precision measurements and direct or indirect searches for new physics.

References

1. CMS Collaboration, The CMS Experiment at the CERN LHC, *JINST* **3**, p. S08004 (2008).
2. CMS Collaboration, The CMS Public Physics Results and Detector Performance Web-pages, <http://cms.web.cern.ch/news/cms-physics-results> (2017).
3. CMS Collaboration, The CMS Public Luminosity Results and Measurement Web-pages, <https://twiki.cern.ch/twiki/bin/view/CMSPublic/LumiPublicResults> (2017).
4. CMS Collaboration, Level-1 E/Gamma and Muon performance on 2017 data, Tech. Rep. CMS-DP-2017-024 (2017), cds.cern.ch/record/2273270.
5. CMS Collaboration, CMS Tracker Alignment Performance Results 2016, Tech. Rep. CMS-DP-2017-021 (2017), cds.cern.ch/record/2273267.
6. CMS Collaboration, Missing transverse energy performance in high pileup data collected by CMS in 2016, Tech. Rep. CMS-DP-2017-028 (2017), cds.cern.ch/record/2275227.
7. CMS Collaboration, Measurement of the differential jet production cross section with respect to jet mass and transverse momentum in dijet events from pp collisions at $\sqrt{s} = 13 \text{ TeV}$, Tech. Rep. CMS-PAS-SMP-16-010 (2017), cds.cern.ch/record/2273393.
8. CMS Collaboration, Measurement of the weak mixing angle with the forward-backward asymmetry of Drell-Yan events at 8 TeV, Tech. Rep. CMS-PAS-SMP-16-007, CERN (Geneva, 2017), cds.cern.ch/record/2273392.
9. A. M. Sirunyan *et al.*, Measurement of the cross section for top quark pair production in association with a W or Z boson in proton-proton collisions at $\sqrt{s} = 13 \text{ TeV}$,

- hep-ex:1711.02547*, subm. to JHEP (2017).
10. CMS Collaboration, Measurements of $t\bar{t}$ cross sections in association with b jets and inclusive jets and their ratio using dilepton final states in pp collisions at $\sqrt{s} = 13$ TeV (2017).
 11. CMS Collaboration, Measurement of the semileptonic $t\bar{t} + \gamma$ production cross section in pp collisions at $\sqrt{s} = 8$ TeV, *JHEP* **10**, p. 006 (2017).
 12. CMS Collaboration, Search for associated production of a Z boson with a single top quark and for tZ flavour-changing interactions in pp collisions at $\sqrt{s} = 8$ TeV, *JHEP* **07**, p. 003 (2017).
 13. CMS Collaboration, Measurement of the differential cross sections of top quark pair production as a function of kinematic event variables in pp collisions at $\sqrt{s} = 13$ TeV, Tech. Rep. CMS-PAS-TOP-16-014, CERN (Geneva, 2017), cds.cern.ch/record/2273459.
 14. CMS Collaboration, Measurement of the top quark mass using proton-proton data at $\sqrt{s} = 7$ and 8 TeV, *Phys. Rev.* **D93**, p. 072004 (2016).
 15. CMS Collaboration, Measurements of properties of the Higgs boson in the diphoton decay channel with the full 2016 data set, Tech. Rep. CMS-PAS-HIG-16-040, CERN (Geneva, 2017), cds.cern.ch/record/2264515.
 16. CMS Collaboration, Measurements of properties of the Higgs boson decaying into the four-lepton final state in pp collisions at $\sqrt{s} = 13$ TeV (2017).
 17. ATLAS and CMS collaborations, Measurements of the Higgs boson production and decay rates and constraints on its couplings from a combined ATLAS and CMS analysis of the LHC pp collision data at $\sqrt{s} = 7$ and 8 TeV, *JHEP* **08**, p. 045 (2016).
 18. CMS Collaboration, Higgs to WW measurements with 15.2 fb^{-1} of 13 TeV proton-proton collisions, Tech. Rep. CMS-PAS-HIG-16-021, CERN (Geneva, 2017), cds.cern.ch/record/2273908.
 19. CMS Collaboration, Observation of the Higgs boson decay to a pair of tau leptons, *hep-ex:1708.00373*, subm. to PLB (2017).
 20. CMS Collaboration, Evidence for the Higgs boson decay to a bottom quark-antiquark pair, *hep-ex:1709.07497*, subm. to PLB (2017).
 21. T. Golling, LHC searches for exotic new particles, *Prog. Part. Nucl. Phys.* **90**, 156 (2016).
 22. CMS Collaboration, Search for new physics with dijet angular distributions in proton-proton collisions at $\sqrt{s} = 13$ TeV and constraints on dark matter and other models, Tech. Rep. CMS-PAS-EXO-16-046, CERN (Geneva, 2017), cds.cern.ch/record/2273455.
 23. C. Autermann, Experimental status of supersymmetry after the LHC Run-I, *Prog. Part. Nucl. Phys.* **90**, 125 (2016).
 24. CMS Collaboration, Search for supersymmetry in events with at least one photon, missing transverse momentum, and large transverse event activity in proton-proton collisions at $\sqrt{s} = 13$ TeV (2017).
 25. A. M. Sirunyan *et al.*, Search for gauge-mediated supersymmetry in events with at least one photon and missing transverse momentum in pp collisions at $\sqrt{s} = 13$ TeV, *hep-ex:1711.08008*, subm. to PLB (2017).
 26. CMS Collaboration, Combined search for electroweak production of charginos and neutralinos in proton-proton collisions at $\sqrt{s} = 13$ TeV (2018).
 27. CMS Collaboration, Search for R -parity violating supersymmetry in pp collisions at $\sqrt{s} = 13$ TeV using b jets in a final state with a single lepton, many jets, and high sum of large-radius jet masses, *hep-ex:1712.08920*, subm. to PLB (2017).