RENORMALIZATION GROUP ANALYSIS OF A CLASS OF 2-DIMENSIONAL VORTEX SYSTEMS

Steven Thomas¹

Theory Division, C.E.R.N.
CH-1211 Geneva 23 Switzerland.

ABSTRACT

The statistical mechanical properties of a class of 2-dimensional vortex systems, described by grand canonical partition functions containing certain $\mathcal{Z}_n$ -valued adiabatic phases, are investigated. It has been shown previously that such systems describe 2-dimensional incompressible superfluids, in which an additional (non-thermal) vortex is present, and which is restricted to move along a closed curve Γ enclosing the bulk fluid. By generalizing the well known connection between the Coulomb gas partition function and the 2-dimensional Sine-Gordon model, to include adiabatic phases, we derive a series of renormalization group equations, the asymptotic solutions of which determine the macroscopic superfluid density. Numerical solutions to these equations are given, from which it is concluded that although the density is discontinuous along a line of critical temperatures T_c , the size of the discontinuity is reduced by a factor of $1/n$ compared to the case where adiabatic phases are absent.

CERN-TH.6558/92

June 1992

¹Bitnet: sthomas@cernvm

1 Introduction

In a recent paper [1], a class of 2-dimensional superfluids was considered in which quantum effects due to an additional, non-thermal vortex, moving in the fluid were investigated. When the motion of this non-thermal vortex is restricted to a curve Γ enclosing the bulk fluid, quantization of the vortex leads to the appearance of certain adiabatic phases [2,3] in the partition function of thermally created vortices. More specifically, if one considers states of the superfluid in which N vortices/antivortices are present as a result of thermal fluctuations, the phases take the form $\gamma = e^{2\pi i N(\pi \xi^2 \rho_0)}$ [3], where ρ_0 is the macroscopic superfluid (number) density and ξ is a length scale characterizing the size of the vortex cores. By integrating out the quantum degrees of freedom (i.e. the superfluid particles and the vortex confined to Γ) from the partition function (in the low temperature approximation), one is left with the grand canonical partition function (GCPF) describing vortex-antivortex pair creation in the superfluid, but now including the N -dependent phases γ . An important assumption leading to this result was that the superfluid be incompressible in the interior of the curve Γ .

In ref. [1], the classical statistical mechanical properties of the vortex system described by this modified GCPF were investigated in the simplest non-trivial case where $\gamma = e^{i\pi N/2}$ (where $N = N_+ + N_-$ is always even, since it is energetically favorable for vortices to be created in neutral pairs in the thermodynamic limit, $N_{\pm}$ being the total number of vortices (+) and antivortices (-)). In particular, possible modifications of the Kosterlitz-Thouless (K-T) phase transition [4], normally associated with such vortex systems in the absence of phases, were studied. This problem was simplified by exploiting the connection between the GCPF and the generating function of the Sine-Gordon model in 2-spatial dimensions. Indeed for the simple case of Z_2 -valued phase γ , it was found that the GCPF was proportional to the sum of partition functions corresponding to those of the S-G model with real and imaginary couplings respectively.

By studying the renormalization group (R.G.) behavior of the parameters in the latter two theories, the macroscopic superfluid density $\rho_{s.f.}$ could be determined as a function of the fugacity z and the quantity $(T - T_c(z))/T$, at least for small z and $T \approx T_c(z)$. It was found that a line of critical temperatures $T_c(z)$ exist at which the ratio $\rho_{s.f.}/T$ suffered a discontinuous change, the size of which was universal, a phenomenon that had previously been discovered in the case where the phase γ was

absent [5]. However, two important differences arose, namely that the size of the jump in $\rho_{s.f.}/T$ was half its usual value, and furthermore the value of $\rho_{s.f.}$ when $T > T_c$ (but T still sufficiently close to T_c to justify perturbative R.G. methods) was non-zero. The physical interpretation of this last phenomenon was that even though the density $\rho_{s.f.}$ is discontinuous at T_c , the Debye screening length remained infinite for $T > T_c$. This implies that the Coulomb force between a vortex-antivortex pair although decreased by a factor of a half in the region $T > T_c$ compared to $T \leq T_c$, is still long-range and decreases like $\ln|x|$ for large inter-vortex separation. This should be contrasted with the case of vortices in the normal superfluid, where for $T > T_c$, the screening length λ is finite so that the inter-vortex potential falls off like $e^{-|x|/\lambda}$ leading to unbinding of vortex- antivortex pairs [6]. Thus we can say that the presence of Z_2 - valued phases produces incomplete screening in the vortex gas, but still allows for a jump discontinuity in $\rho_{s.f.}$ at T_c . In the sense that the unbinding transition is suppressed one might say that the usual K-T phase transition is circumvented, although its physical manifestation as a discontinuity in $\rho_{s.f.}$ remains.

The purpose of the present paper is to consider the general case of Z_n -valued adiabatic phases $\gamma = e^{\frac{\pi i N}{n}}$ (we shall only consider rational phases in this paper), and specifically to compute the behavior of $\rho_{s.f.}$ as a means of investigating possible modifications to the K-T phase transition. We shall see that the case of general n is rather more involved than in the simple case of $n=2$, because even though one can again exploit the connection with Sine-Gordon (S-G) field theory, this now involves defining the latter theories with complex couplings. From this connection we derive a set of R.G. equations, that with appropriate boundary conditions, determine the macroscopic superfluid density. For general n we analyse the fixed points of the lowest order R.G.E.'s and find for each value of n a unique (infrared stable) fixed point, corresponding to a definite value of the macroscopic density $\rho_{s.f.}$. We present numerical solutions to the lowest order equations for various initial conditions and for the first few values of n . The solutions that emerge, show a similarity in behavior that can be extrapolated to any finite value of n . The main conclusion is that $\rho_{s.f.}$, for general n , undergoes a discontinuity at $T = T_c(z)$ (where we will specify $T_c(z)$ later) the size of which is universal and reduced by a factor of $1/n$ compared to the case where adiabatic phases are absent.

Moreover, as in the $n=2$ case, $\rho_{s.f.}$ is non-vanishing for $T > T_c$, and an explicit

formula is given for its value, valid for $T \approx T_c$ and small fugacity z . It is worth pointing out that the expression for $T_c(z)$ is independent of n , and is the same as the one derived in the case of normal 2-d superfluidity [6].

The structure of this paper is as follows. In section 2 starting from the GCPF of vortex-antivortex pair creation including $\mathcal{Z}_n$ -valued adiabatic phases (as derived in ref. [1]), we present the connection to S-G models with complex parameters, from which we derive an expression for $\rho_{s.f.}$ in terms of the renormalized two-point functions. Generalizing some well-known properties of the renormalized S-G model with real parameters to the present context, we obtain a set R.G.E.'s for the various parameters which determine the macroscopic density $\rho_{s.f.}$ with appropriate boundary conditions. In section 3 we analyse these equations and present numerical solutions for the first few values of n . Finally we end the paper with a discussion and interpretation of the results.

2 Vortex partition function and the superfluid density

In this section we will derive a formula for the density $\rho_{s.f.}(z, T)$ as a function of temperature T and fugacity z , where the latter is associated with vortex-antivortex creation in the superfluid. Our starting point will be the regularized GCPF of vortex-antivortex pair creation in the superfluid including the $\mathcal{Z}_n$ -valued adiabatic phase contributions coming from the quantization of the vortex restricted to the closed curve Γ , surrounding the system. This partition function, for general n , was derived in ref. [1] from the microscopic point of view, and where further details may be found. For the purposes of this paper one can regard this partition function as a particular kind of extension of that describing a Coulomb gas of charges [6], whose properties have been studied in the past literature. Having said this, it is worth remembering that this extension is not adhoc and can be derived from the physical model described in [1].

The GCPF which we start from is given by [1]

$$\begin{aligned}
 Z(z, T, \omega) &= \frac{D(\omega, T)}{2} \sum_{N=0}^{\infty} \frac{z^N}{N!} \prod_{a=1}^N \sum_{q_a} \int \frac{d^2 \mathbf{R}_a}{\pi \xi^2} e^{-E(\mathbf{R}_a, q_a, N)/T} \\
 &\times \sum_{p \in \mathcal{Z}} e^{-\omega p^2/T - \pi i N p q/n}
 \end{aligned} \tag{2.1}$$

In eq.(2.1) z is the fugacity associated with vortex-antivortex creation, N the number of vortices and antivortices which have charges q_a and 2-dimensional position vectors $\mathbf{R}_a, a = 1, \dots, N$. We only consider vorticities of unit magnitude $|q_a| = 1$ as these are the stable, lowest energy configurations. q is the charge of the vortex moving along Γ which we also take to be of unit magnitude, (it does not matter if $q = \pm 1$ since either sign gives the same contribution to Z). ξ is a length scale introduced earlier which characterizes the core region of each vortex. $E(\mathbf{R}_a, q_a, N)$ is the interaction energy of the N -vortex configuration described above and is of Coulomb type

$$E(\mathbf{R}_a, q_a, N) = 2\pi\rho_0 \left(- \sum_{\substack{a,b=1 \\ a \neq b}}^N \gamma_a \gamma_b \ln(|\mathbf{R}_a - \mathbf{R}_b|/\xi) \right) \quad (2.2)$$

where ρ_0 is the bulk superfluid density at zero temperature (i.e. in the absence of thermal vortex-antivortex pair creation), $\gamma_a \equiv q_a \hbar/m$ is the quantum of circulation or vorticity, and m the mass of the superfluid particles. Note that eq. (2.1) describes the system in the thermodynamic limit where the extent of the curve Γ is macroscopic. In such a limit it was shown in [1] that the terms one might expect in eq. (2.2) coming from interactions between the thermal and non-thermal vortices become negligible. The terms in eq. (2.1) that are a departure from the usual Coulomb gas GCPF are the prefactor $D(T, \omega)$ and the sum over winding modes labelled by integers p , coming from contributions of the non-thermal vortex restricted to Γ (which without loss of generality we take to be a circle) enclosing the system. $D(T, \omega)$ is a determinant arising from spin-wave fluctuations of the latter, i.e.

$$D(T, \omega) = \int D\tilde{\phi} \ e^{-\omega \int_0^\beta d\tau_0 (\dot{\tilde{\phi}})^2} \quad (2.3)$$

where the $\cdot$ in eq. (2.3) represents the derivative with respect to the imaginary time parameter τ_0 , $\tilde{\phi}$ are the spin wave fluctuations and β is the usual inverse temperature. The ω dependent factors in eqs. (2.1), (2.3) are inserted by hand to regulate the divergences coming from the infinite sum over winding modes. However the insertion of such a term into the Euclidean action of the vortex confined to Γ , with position vector $\mathbf{R}$, can be thought of as given the latter a mass proportional to ω , since it arises from a term of the form $\omega \int_0^\beta d\tau_0 \dot{\mathbf{R}}^2$. The physical limit corresponds to $\omega \rightarrow 0$ since it is well known that physical vortices carry no inertia and may be thought of as massless particles in this respect [3]. Associated with the winding modes of periodicity p in the GCPF (2.1) are the dynamical or adiabatic phases $e^{\pi i N p q/n}$ which depend on the number of (thermally created) vortices and antivortices enclosed by the path Γ .

In the limit where the regulator is removed, the divergence in Z is an overall constant multiplying the partition function. This constant factor will cancel in normalized statistical averages of quantities involving the thermal vortices, for example later we will see this in calculating the superfluid density. To verify this factorization property

of the divergence, first we decompose the sum over p in eq. (2.1) into separate sums over the n sets of integers $\mathcal{N}_i \equiv [i \bmod n]$, $i = 1 \dots n$. This can be done by inserting the projection operators $\mathcal{P}_j(n, p)$ into the sum over p in eq. (2.1), where the $\mathcal{P}_j(n, p)$, $j=1 \dots n$ are defined by

$$\mathcal{P}_j(n, p) = \frac{1}{n} \left[1 + e^{\frac{2\pi i}{n}(n+p-j)} + e^{\frac{2\pi i}{n}(n+2p-2j)} + \dots + e^{\frac{2\pi i}{n}(n+(n-1)p-(n-1)j)} \right] \quad (2.4)$$

The partition function Z can then be expressed as

$$\begin{aligned} Z(z, T, \omega) = & \frac{D(\omega, T)}{2n} \sum_{N=0}^{\infty} \frac{z^N}{N!} \sum_{\mathbf{q}_a} \int \frac{d^2 \mathbf{R}_a}{\pi \xi^2} e^{-E(\mathbf{R}_a, \mathbf{q}_a, n)/T} \left(\left[\theta \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0|i\omega) \right. \right. \\ & + e^{\frac{2\pi i}{n}(n-1)} \theta \begin{bmatrix} 0 \\ \frac{1}{n} \end{bmatrix} (0|i\omega) + \dots + e^{\frac{2\pi i}{n}} \theta \begin{bmatrix} 0 \\ \frac{n-1}{n} \end{bmatrix} (0|i\omega) \left. \right] e^{-\frac{\pi i N}{n}} \\ & + \left[\theta \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0|i\omega) + e^{\frac{2\pi i}{n}(n-2)} \theta \begin{bmatrix} 0 \\ \frac{1}{n} \end{bmatrix} (0|i\omega) + \dots \right. \\ & + \dots + e^{\frac{4\pi i}{n}} \theta \begin{bmatrix} 0 \\ \frac{n-1}{n} \end{bmatrix} (0|i\omega) \left. \right] e^{-\frac{2\pi i N}{n}} + \dots + \\ & \left. + \left[\theta \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0|i\omega) + \theta \begin{bmatrix} 0 \\ \frac{1}{n} \end{bmatrix} (0|i\omega) + \dots + \theta \begin{bmatrix} 0 \\ \frac{n-1}{n} \end{bmatrix} (0|i\omega) \right] \right) \end{aligned} \quad (2.5)$$

The generalized theta-functions $\theta \begin{bmatrix} \chi \\ \phi \end{bmatrix} (0|\tau)$ appearing in eq. (2.5) are defined by

$$\theta \begin{bmatrix} \chi \\ \phi \end{bmatrix} (0|\tau) = \sum_{p \in \mathbb{Z}} e^{i\pi(p+\chi)^2 \tau + 2\pi i(p+\chi)\phi} \quad (2.6)$$

In the $\lim_{\omega \rightarrow 0}$, the theta-functions $\theta \begin{bmatrix} 0 \\ \frac{1}{n} \end{bmatrix} (0|i\omega)$, $i=1\dots n-1$ vanish, whilst

$\theta \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0|i\omega)$ diverges like $\sum_{p \in Z}$. Thus the divergent terms inside the [] brackets in eq. (2.5) are of the latter type, and so Z in this limit approaches

$$Z(z, T, \omega) \approx \frac{\Omega(T, \omega)}{n} \sum_{j=1}^n Z_j(z, T) \quad (2.7)$$

where the divergent factor Ω is

$$\Omega(T, \omega) = \frac{D(T, \omega)}{2} \theta \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0|i\omega) \quad (2.8)$$

and where the ‘sub’-partition functions Z_j are defined as

$$Z_j(z, T) = \sum_{N=0}^{\infty} \frac{z^N}{N!} \prod_{a=1}^N \sum_{q_a} \int \frac{d^2 \mathbf{R}_a}{\pi \xi^2} e^{-E(\mathbf{R}_a, q_a, N)/T - (\frac{\pi i N}{n})j} \quad (2.9)$$

We see from eq. (2.9) that Z_j is just the usual Coulomb gas partition function with the fugacity rescaled by a phase $z \rightarrow ze^{\frac{i\pi j}{n}}$, $j = 1\dots n$. The divergent factor in eq. (2.8) are overall constants which cancel in computing physical quantities involving the ensemble of thermal vortices. Thus, the statistical mechanical properties of the thermal vortices, including the effects of adiabatic phases will be described equally well by the ω -independent partition function $\sum_{j=1}^n Z_j = nZ/\Omega$.

Next, we shall derive a formula for the superfluid density $\rho_{s.f.}$ in terms of the partition function Z . The derivation of this formula for the case of a normal 2-d superfluid can be found in ref. [7]. The generalization to the present case is quite straightforward. This is partly because the expression for the energy in eq. (2.2) is independent of the adiabatic phases. As such, it can be interpreted as the hydrodynamical hamiltonian associated with the normal component of the superfluid mass current, in the presence of an N -vortex configuration [6],[7] i.e.

$$E(\mathbf{R}_a, q_a, N) = \frac{1}{2\rho_0} \int d^2x \quad \mathbf{g}_{\perp} \cdot \mathbf{g}_{\perp} \quad (2.10)$$

where the perpendicular component of the mass current, $\mathbf{g}_\perp$, in the presence of an N-vortex configuration is

$$\mathbf{g}_\perp(\mathbf{x}) = \rho_0 \sum_{a=1}^N \frac{\gamma_a(\tilde{\mathbf{x}} - \tilde{\mathbf{R}}_a)}{|\mathbf{x} - \mathbf{R}_a|^2} \quad (2.11)$$

where the vectors $\tilde{\mathbf{x}}, \tilde{\mathbf{R}}_a$ are dual to $\mathbf{x}$ and $\mathbf{R}_a$. The total mass current $\mathbf{g}$ in 2-dimensions may be decomposed as $\mathbf{g} = \mathbf{g}_\perp + \mathbf{g}_\parallel$, with $\nabla \cdot \mathbf{g}_\perp = 0$, $\nabla \times \mathbf{g}_\parallel = 0$, ∇ being the two dimensional gradient operator. Note that the parallel component of $\mathbf{g}$ decouples from that of $\mathbf{g}_\perp$ in the expression for the energy in eq. (2.10) because of the overall neutrality of vortex configurations created by thermal fluctuations. Hence we can discard the $\mathbf{g}_\parallel$ contribution as it plays no role in determining the behavior of the superfluid mass flow which only depends on $\mathbf{g}_\perp$ [6],[7].

To define $\rho_{s.f.}$, (see ref. [7]), we imagine imposing an infinitesimal constant field $\mathbf{u} = \delta\mathbf{v} \cdot \mathbf{x}$ in the superfluid, which induces the change $\mathbf{g}_\perp \rightarrow \mathbf{g}_\perp + \rho_0\delta\mathbf{v}$. Strictly speaking the applied field induces a change in the full mass current $\mathbf{g}$ but again one can check that the parallel component decouples following the previous reasoning. Corresponding to the applied flow $\rho_0\delta\mathbf{v}$ there will be an induced infinitesimal mass current $\delta\mathbf{J}_u$ defined by

$$\delta\mathbf{J}_u = \frac{\partial}{\partial\delta\mathbf{v}} \ln(Z(z, T, \delta\mathbf{v})) \quad (2.12)$$

Here, $Z(z, T, \delta\mathbf{v})$ is the partition function of eq. (1.1) in the presence of the infinitesimal flow $\rho_0\delta\mathbf{v}$. The presence of $\delta\mathbf{v}$ modifies the configurational energy E which can easily be calculated from eq. (2.10) by substituting $\mathbf{g}_\perp \rightarrow \mathbf{g}_\perp + \rho_0\delta\mathbf{v}$. Following ref. [7], the definition of $\rho_{s.f.}$ we take is

$$\rho_{s.f.}^{-1} = \frac{1}{A} \left[\frac{\partial^2}{\partial|\delta\mathbf{v}|^2} \ln(Z(z, T, \delta\mathbf{v})) \right]_{\delta\mathbf{v}=0} \quad (2.13)$$

with A in eq. (2.13) being the area of the superfluid sample, which in this case is the interior area πR^2 bounded by the circle Γ , of radius R . Strictly speaking the expression for the partition function given in eq. (2.1) is true only in the thermodynamic limit R

$\rightarrow \infty$, [1] . However we shall assume that the ratio $R/\xi \gg 1$ so as to make eq. (2.1) a good approximation. Shortly we shall see that the factors of A will, infact, cancel in the expression for $\rho_{s.f.}$ allowing one to take the limit $R \rightarrow \infty$ in which eq. (2.1) is strictly valid. Now using the property that $\delta J_u = 0$ when $\delta \mathbf{v} = 0$, (which essentially follows again from overall neutrality of vortex configurations) and the fact that δJ_u is canonically conjugate to $\delta \mathbf{v}$ (see eq. (2.12)), one can show that

$$\rho_{s.f.} = \frac{1}{A} \left[\frac{\partial |\delta J_u|}{\partial |\delta \mathbf{v}|} \right]_{\delta \mathbf{v}=0} \quad (2.14)$$

To proceed we use the connection between $\mathbf{g}_\perp(\mathbf{x} - \mathbf{R}_a, \mathbf{q}_a)$ given in eq. (2.11) and the vortex charge density $n(\mathbf{x}, \mathbf{R}_a) = \sum_{a=1}^N \gamma_a \delta^{(2)}(\mathbf{x} - \mathbf{R}_a)$ via the relation

$$\mathbf{g}_\perp(\mathbf{x} - \mathbf{R}_a) = \rho_0 \nabla_{\mathbf{x}} \times \hat{\mathbf{z}} \int d^2 \mathbf{x}' \ln(|\mathbf{x} - \mathbf{x}'|) n(\mathbf{x} - \mathbf{x}', \mathbf{R}_a) \quad (2.15)$$

where $\hat{\mathbf{z}}$ is the unit vector perpendicular to the plane of the superfluid. From eqs. (2.12),(2.14),(2.15) it follows that

$$\frac{\rho_{s.f.}}{\rho_0}(z, T) = \lim_{k^2 \rightarrow 0} \left[\sum_{j=1}^n \frac{Z_j}{Z} \left(1 - \frac{2\pi\alpha}{T} \langle n(\mathbf{k})n(-\mathbf{k}) \rangle_{(j)} \right) \right] \quad (2.16)$$

where in eq. (2.16) it is convenient to express the result in terms of momentum space quantities, with $\mathbf{k}$ being the 2-dimensional momentum. It is also convenient to introduce the constant $\alpha = 2\pi\rho_0\hbar^2/m^2$ in eq. (2.16), while the brackets $\langle \rangle_{(j)}$ indicate normalized averaging with respect to the (sub)partition function Z_j defined earlier in eq. (2.9). The factors of the area A have cancelled in the formula for $\rho_{s.f.}$ as promised, and moreover simple inspection shows that the divergent factors in Z also cancel, so that we can safely remove the regulator ($\omega \rightarrow 0$).

Finally, the calculation of the density-density correlation functions in eq. (2.16) is simplified by introducing field theoretic representations of Z_j and hence of the correlation functions defined with respect to them. In the case of the standard Coulomb gas GCPF (which corresponds to $Z_{j=n}$), such a representation has been known for some time [8], and may be expressed as

$$Z_{j=n} = \frac{\int \mathcal{D}\chi e^{-S(\chi, T, z)}}{\int \mathcal{D}\chi e^{-S_0(\chi, T)}} \quad (2.17)$$

where

$$S = S_0 + S_{\text{int}} = \int d^2x \left(\frac{1}{2} (\nabla \chi)^2 - \frac{2z}{\pi \xi^2} \text{Cos} \left[\sqrt{\frac{2\pi\alpha}{T}} \chi \right] \right) \quad (2.18)$$

is the action of the Sine-Gordon model in 2 spatial dimensions. Since Z_j for $j = 1 \dots n-1$ is just the Coulomb gas partition function with rescaled fugacity $z \rightarrow e^{\frac{i\pi j}{n}} z$, their representation in terms of the Sine-Gordon generating functional is straightforward, namely

$$Z_j = \frac{\int \mathcal{D}\chi e^{-S(\chi, T, e^{\frac{i\pi j}{n}} z)}}{\int \mathcal{D}\chi e^{-S_0(\chi, T)}} \quad (2.19)$$

where the action $S(\chi, T, e^{\frac{i\pi j}{n}} z)$ is the same as in eq. (2.18) but with the fugacity z rescaled appropriately.

It should be mentioned that in passing from the (sub)partition functions Z_j to their field theory representations, as given in eq. (2.19), one strictly speaking obtains a non-local version of the Sine-Gordon model with the argument of the cosine $\chi(x)$ replaced by the smeared field $\int d^2y f_\xi(|y-x|/\xi) \chi(y)$ [6]. Here, f_ξ is a distribution function which describes the charge density of a vortex, and is maximized around the core region. Eqs. (2.17), (2.19) hold for δ -function charge distributions only. Whilst it is true that realistic vortices have a finite core region the extent of which is of order ξ , we expect that the long distance critical properties of the system in the thermodynamic limit will be described by local versions of the Sine-Gordon model. However, in this case there are logarithmic divergences associated with model which must be removed by renormalization. Hence we should understand the field theory representation of Z_j in eqs. (2.17), (2.19) as being expressed in terms of the renormalized Sine-Gordon models and their correlation functions. It is not difficult to show that in terms of the renormalized fields $\tilde{\chi}$, the formula for the superfluid density becomes

$$\frac{\rho_{\text{s.f.}}}{\rho_0}(z, T) = \lim_{k^2 \rightarrow 0} \left[k^2 \sum_{j=1}^n \frac{Z_j}{Z} \langle \tilde{\chi}(k) \tilde{\chi}(-k) \rangle_{\text{S.G.}}^{(j)} \right] \quad (2.20)$$

where the brackets $\langle \rangle_{\text{S.G.}}^{(j)}$ indicate correlations with respect to the renormalized Sine-Gordon model with bare self coupling $e^{\frac{i\pi j}{n}} z$.

In order to proceed with the evaluation of $\rho_{\text{s.f.}}$, it is necessary to understand some of the well known properties [9] of the renormalized Sine-Gordon model in 2-dimensions, and by implication those of the same models but with bare fugacity scaled by the phase factors $e^{\frac{i\pi j}{n}}$, $j=1\dots n-1$. Firstly, all divergences are removed by wavefunction and coupling constant renormalization which in the present context is equivalent to a renormalization of fugacity and temperature, (see eq. (2.18)). Also it is known that the Sine-Gordon partition function and correlations involving derivatives of the renormalized fields $\tilde{\chi}$ are even functions of the renormalized fugacity z_R . This a consequence of the $z_R \rightarrow -z_R$ symmetry of the model, which results from the periodicity $\tilde{\chi} \rightarrow \tilde{\chi} + 2n\pi\sqrt{T}$.

It is clear that the renormalization properties of Sine-Gordon models with bare coupling rescaled by phase factors will mirror those described above. Of course in these cases, the effect of renormalization will be to produce complex renormalized temperature and fugacity, which we shall denote by z_R^j and T_R^j , $j=1\dots n$ respectively, (although they are real for $j = n$, since in this case the phase is unity). The presence of complex renormalization constants does not destroy the periodicity described above so that there will still remain a symmetry $e^{\frac{i\pi j}{n}} z_R^j \rightarrow -e^{\frac{i\pi j}{n}} z_R^j$. From this symmetry one may deduce the following conjugation properties of Z_j and the two-point functions $\langle \tilde{\chi}(k)\tilde{\chi}(-k) \rangle_{\text{S.G.}}^j$.

$$Z_j^* = Z_{n-j}$$

$$\left(\langle \tilde{\chi}(k)\tilde{\chi}(-k) \rangle_{\text{S.G.}}^{(j)} \right)^* = \langle \tilde{\chi}(k)\tilde{\chi}(-k) \rangle_{\text{S.G.}}^{(n-j)} \quad (2.21)$$

Consequently, it is clear that $\rho_{\text{s.f.}}/\rho_0$ in eq. (2.20) is a real quantity (we shall return to the question of its positivity later). Consider then the j^{th} such Sine-Gordon theory with renormalized fugacity which we write as $e^{\frac{i\pi j}{n}} z_R^j(l)$, and renormalized temperature $T_R^j(l)$, where l is a parameter associated with a change of scale. Since of course, there is a single real physical temperature T of the underlying statistical system of vortices,

which does not scale, we write $T_R^j(l) = e^j(l)T$, for a set of parameters $e^j(l)$. The quantities $z_R^j(l)$ and $e^j(l)$ satisfy R.G.E's which can be obtained from those of the standard Sine-Gordon model [9], by the substitution $z_R \rightarrow e^{\frac{in_j}{n}} z_R^j$; $T_R \rightarrow T_R^j$. To lowest order in a double expansion of the respective parameters the resulting R.G.E.'s are

$$dz_R^j/dl = 2z_R^j(1 - \alpha/4e^jT) \quad (2.22a)$$

$$de^j/dl = \frac{2\pi^2\alpha}{T}(z_R^j)^2 e^{\frac{2in_j}{n}} \quad (2.22b)$$

The boundary conditions on the solutions of these equations are chosen to be $z_R^j(l=0) = z$, $j=1\dots n$, where z is the real fugacity appearing in the vortex partition function of eq. (2.1), and $e^j = 1$ (i.e. $T_R^j = T$), $j=1\dots n$. The choice of these boundary conditions reflects the physical situation present in the vortex system where scaling of the temperature and fugacity z is a consequence of (length dependent) screening of the interaction between vortices. At very short scales $l \approx 0$, screening effects become negligible and so the values of z_R^j and T_R^j should be equal to the fugacity and temperature z , T appearing in the vortex partition function. Also we note that the conjugation properties described in eq. (2.21) imply that the solutions to the R.G.E's should satisfy the constraints $z_R^{j*} = z_R^{n-j}$, $e^{j*} = e^{n-j}$.

In terms of the renormalized parameters $e^j(l)$, one may obtain a formula for the superfluid density from eq. (2.20), to leading order in a small z expansion

$$\frac{\rho_{s.f.}}{\rho_0}(z, T) \approx \lim_{l \rightarrow \infty} \left[\frac{4T}{n\alpha} \sum_{j=1}^n [1 - t^j(l)] \right] \quad (2.23)$$

where we have introduced the conventions

$$y^j(l) \equiv \frac{\pi\alpha z_R^j(l)}{2T} \quad ; \quad t^j(l) \equiv \left(1 - \frac{\alpha}{4Te^j(l)}\right) \quad (2.24)$$

Again it is clear that the r.h.s. of eq.(2.23) is real since $t^{j*} = t^{n-j}$. The factors of $1/n$ appearing in eq. (2.23) come from the small z expansion of $Z_j \approx 1 + O(z)$. It is convenient to decompose the (in general) complex parameters t^j, y^j into real and

imaginary parts $y^j = y_0^j + iy_1^j$, $t^j = t_0^j + it_1^j$, in terms of which, the first order R.G.E's of eq. (2.22) become

$$dt_0^j/dl = 2((y_0^j)^2 - (y_1^j)^2) \cos(2\pi j/n) - 4y_0^j y_1^j \sin(2\pi j/n) \quad (2.25a)$$

$$dt_1^j/dl = 4y_0^j y_1^j \cos(2\pi j/n) + 2((y_0^j)^2 - (y_1^j)^2) \sin(2\pi j/n) \quad (2.25b)$$

$$dy_0^j/dl = 2(y_0^j t_0^j - y_1^j t_1^j) \quad (2.25c)$$

$$dy_1^j/dl = 2(t_0^j y_1^j + t_1^j y_0^j) \quad (2.25d)$$

The boundary conditions on the variables $y_0^j, \dots, t_1^j$ are $y_0^j = \pi\alpha z/2T$, $y_1^j = 0$, $t_0^j = (1 - \alpha/4T)$, $t_1^j = 0$, $j=1, \dots, n$. It can be checked that the constraints $t_0^j = t_0^{n-j}$, $t_1^j = -t_1^{n-j}$, $y_0^j = y_0^{n-j}$, $y_1^j = y_1^{n-j}$ which follow from the previous conjugation properties, are consistent with the $4n$ coupled equations (2.25a), ..., (2.25d) for $j = 1, \dots, n$. Note that even though the formula for $\rho_{s.f.}$ in eq. (2.23) depends only on the variables t_0^j , one has to solve, in principle, the coupled set of equations (2.25) in all 4 variables $y_0^j, y_1^j, t_0^j, t_1^j$. In the next section, we shall study the fixed points of the R.G.E's (2.25) for general n, j . We have been unable to obtain analytic solutions to these equations for arbitrary n and j , so instead we shall present numerical results for the first few values of n and j , which as we shall see, allow extrapolation to the general case. Having said this, $\rho_{s.f.}$ depends only on the infrared (IR) asymptotic values of these solutions, which as we shall see in the next section, are just the IR (stable) fixed points for which an analytic formula is given.

3 Fixed points and Numerical Solutions.

Before discussing the fixed points and numerical solutions to the lowest order R.G.E's eqs. (2.25), it is convenient to introduce the variables

$$Y_j \equiv 2((y_0^j)^2 - (y_1^j)^2) \quad ; \quad \dot{Y}_j \equiv 4y_0^j y_1^j \quad (3.1)$$

Eqs. (2.25) then become

$$d(t_0^j \cos(2\pi j/n) + t_1^j \sin(2\pi j/n))/dl = Y_j \quad (3.2a)$$

$$d(-t_0^j \sin(2\pi j/n) + t_1^j \cos(2\pi j/n))/dl = \dot{Y}_j \quad (3.2b)$$

$$dY_j/dl = 4(t_0^j Y_j - t_1^j \dot{Y}_j) \quad (3.2c)$$

$$d\dot{Y}_j/dl = 4(t_0^j \dot{Y}_j + t_1^j Y_j) \quad (3.2d)$$

Using eqs. (3.2a,b) to substitute for $Y_j, \dot{Y}_j$ in eqs. (3.2c, d) yields the following two second order equations in the variables t_0^j, t_1^j

$$\begin{aligned} d^2(t_0^j \cos(2\pi j/n) + t_1^j \sin(2\pi j/n))/dl^2 = \\ d\left(2((t_0^j)^2 - (t_1^j)^2) \cos(2\pi j/n) + 4t_0^j t_1^j \sin(2\pi j/n)\right)/dl \end{aligned} \quad (3.3a)$$

$$\begin{aligned} d^2(-t_0^j \sin(2\pi j/n) + t_1^j \cos(2\pi j/n))/dl^2 = \\ d\left(-2((t_0^j)^2 - (t_1^j)^2) \sin(2\pi j/n) + 4t_0^j t_1^j \cos(2\pi j/n)\right)/dl \end{aligned} \quad (3.3b)$$

Now since the r.h.s. of eqs. (3.3) are total derivatives, one can integrate these equations once to yield solutions for $Y_j, \dot{Y}_j$ in terms of the variables t_0^j and t_1^j

$$Y_j = 2((t_0^j)^2 - (t_1^j)^2) \cos(2\pi j/n) + 4t_0^j t_1^j \sin(2\pi j/n) + c_j \quad (3.4a)$$

$$\dot{Y}_j = 2((t_1^j)^2 - (t_0^j)^2) \sin(2\pi j/n) + 4t_0^j t_1^j \cos(2\pi j/n) + \dot{c}_j \quad (3.4b)$$

where the integration constants $c_j, \dot{c}_j$ in eqs. (3.4) are obtained from imposing the boundary conditions on the solutions i.e.

$$c_j = 2(y_0^j(0))^2 - 2(t_0^j(0))^2 \cos(2\pi j/n) \quad (3.5a)$$

$$\dot{c}_j = 2(t_0^j(0))^2 \sin(2\pi j/n) \quad (3.5b)$$

Finally, substituting the solutions $Y_j(t_0^j, t_1^j)$ and $\dot{Y}_j(t_0^j, t_1^j)$ into eqs. (3.2a,b) we see that the original set of R.G.E's (2.25) of section 2 are equivalent to the following

$$dt_0^j/dl = 2((t_0^j)^2 - (t_1^j)^2 + (y_0^j(0))^2 \cos(2\pi j/n) - (t_0^j(0))^2) \quad (3.6a)$$

$$dt_1^j/dl = 4t_0^j t_1^j + 2(y_0^j(0))^2 \sin(2\pi j/n) \quad (3.6b)$$

$$dy_0^j/dl = 2(y_0^j t_0^j - y_1^j t_1^j) \quad (3.6c)$$

$$dy_1^j/dl = 2(t_0^j y_1^j + t_1^j y_0^j) \quad (3.6d)$$

The problem of determining the R.G. behavior of t_0^j , (which appears in the formula for $\rho_{s.f.}$) is reduced to solving a pair of coupled equations, (3.6a,b) in the variables t_0^j, t_1^j . However we are also interested in solving for y_0^j, y_1^j to see if they exhibit strong coupling behaviour where they increase without bound along a R.G. trajectory that starts close to the origin.

Let us consider the fixed points of eqs. (3.6), which are given by the solutions of the algebraic equations $dt_0^j/dl = dt_1^j/dl = dy_0^j/dl = dy_1^j/dl = 0$. The solutions fall into two classes depending on whether the quantity $(y_0^j(0))^2 \sin^2(2\pi j/n)$ is zero or not. Defining the constants τ, ν by

$$\tau \equiv y_0^j(0) = \frac{\pi \alpha Z}{2T} \quad ; \quad \nu \equiv t_0^j(0) = (1 - \frac{\alpha}{4T}) \quad (3.7)$$

the fixed points are found to be

$$\tau^2 \text{Sin}^2(2\pi j/n) \neq 0$$

$$t_{0\pm}^j = \pm \frac{(-4\nu^2 + 4\tau^2 \text{Cos}(2\pi j/n) - \sqrt{\Delta}) \sqrt{-4\nu^2 + 4\tau^2 \text{Cos}(2\pi j/n) + \sqrt{\Delta}}}{8\sqrt{2}\tau^2 \text{Sin}(2\pi j/n)}$$

$$t_{1\pm}^j = \pm \frac{\sqrt{-4\nu^2 + 4\tau^2 \text{Cos}(2\pi j/n) + \sqrt{\Delta}}}{2\sqrt{2}}$$

$$y_0^j = y_1^j = 0$$

$$\Delta \equiv (4\nu^2 - 4\tau^2 \text{Cos}(2\pi j/n))^2 + 16\tau^4 \text{Sin}^2(2\pi j/n) \quad (3.8)$$

$$\tau^2 \text{Sin}^2(2\pi j/n) = 0$$

$$\tau \neq 0$$

$$t_{0\pm}^j = \pm \sqrt{\nu^2 - \tau^2 \text{Cos}(2\pi j/n)}, \quad t_1^j = y_0^j = y_1^j = 0,$$

$$j = n/2, n \quad n \text{ even} \quad \text{or} \quad j = n, \quad n \text{ odd}$$

$$\tau = 0$$

$$t_{0\pm}^j = \pm \nu, \quad t_1^j = y_0^j = y_1^j = 0, \quad \forall j, n \quad (3.9)$$

Note that the position of the fixed points depends on the initial conditions through the parameters ν and τ . To be consistent with the truncation of the R.G.E.'s to lowest

order, these fixed points should be close to the origin, for all values of j and n . In fact one can always choose initial values ν and τ sufficiently close to the origin, so that all fixed points are consistent with perturbation theory.

In the case when $\tau^2 \sin^2(2\pi j/n) \neq 0$, fixed points exist for all initial values of τ, ν . In the special case where $\tau^2 \sin^2(2\pi j/n) = 0$, $\tau \neq 0$ (eq. (3.9)), and $j = n$, real fixed points only exist if $|\nu| \geq |\tau|$. We shall return to this case later.

In a previous paper [1], the R.G.E's corresponding to the $n=2$ case have already been studied. The fixed points in this case are given in eqs. (3.9). Here we see that since the initial values of y_1^j , and t_1^j , namely $y_1^j = t_1^j = 0$ coincide with the value of these variables at the fixed point, the 4 equations (3.6a,...d) reduce to just two in the variables t_0^j and y_0^j , $j=1,2$. The latter two equations are exactly those derived in ref. [1] for the $n=2$ case. In fact for this simple case (as well as for $j = n$, n odd, and $j = n/2$, n even), one can solve the two coupled equations exactly. Before discussing these solutions, and those of the more general case it is instructive to study the stability of the fixed points listed in eqs. (3.8), (3.9). This can be done by linearizing the R.G.E's in regions close to these points. The linearized equations corresponding to the fixed points in eq. (3.8) are

$$d(t_0^j - t_{0\pm}^j)/dl = 4((t_0^j - t_{0\pm}^j)t_{0\pm}^j - (t_1^j - t_{1\pm}^j)t_{1\pm}^j) \quad (3.10a)$$

$$d(t_1^j - t_{1\pm}^j)/dl = 4((t_0^j - t_{0\pm}^j)t_{1\pm}^j + (t_1^j - t_{1\pm}^j)t_{0\pm}^j) \quad (3.10b)$$

$$dy_0^j/dl = 2(y_0^j t_{0\pm}^j - y_1^j t_{1\pm}^j) \quad (3.10c)$$

$$dy_1^j/dl = 2(y_1^j t_{0\pm}^j + y_0^j t_{1\pm}^j) \quad (3.10d)$$

where in obtaining the last two equations (3.10c,d) we have used the fact that $y_0^j = y_1^j = 0$ at the fixed points. Similar equations corresponding to the fixed points in eq. (3.9) can be obtained from eqs. (3.10) by setting $t_{1\pm}^j = 0$. The general solution of the linearized equations (3.10) is given by

$$(t_0^j - t_{0\pm}^j) = e^{4t_{0\pm}^j l} \left((\nu - t_{0\pm}^j) \text{Cos}(4t_{1\pm}^j l) + t_{1\pm}^j \text{Sin}(4t_{1\pm}^j l) \right) \quad (3.11a)$$

$$(t_1^j - t_{1\pm}^j) = e^{4t_{0\pm}^j l} \left(-t_{1\pm}^j \text{Cos}(4t_{1\pm}^j l) + (\nu - t_{0\pm}^j) \text{Sin}(4t_{1\pm}^j l) \right) \quad (3.11b)$$

$$y_0^j = e^{2t_{0\pm}^j l} \left(\tau \text{Cos}(2t_{1\pm}^j l) \right) \quad (3.11c)$$

$$y_1^j = e^{2t_{0\pm}^j l} \left(\tau \text{Sin}(2t_{1\pm}^j l) \right) \quad (3.11d)$$

It is clear from these solutions that in the case where $t_{1\pm}^j \neq 0$, the projection of the R.G. trajectories in the $t_0^j - t_1^j$ plane are spirals centred about the point $(t_{0\pm}^j, t_{1\pm}^j)$, and in the $y_0^j - y_1^j$ plane, spirals centred about the origin, for each value of j . In the $t_0^j - t_1^j$ plane, the radius of the spiral is $e^{4t_{0\pm}^j l} \sqrt{(\nu - t_{0\pm}^j)^2 + (t_{1\pm}^j)^2}$, whilst in the $y_0^j - y_1^j$ plane it is $\tau e^{2t_{0\pm}^j l}$. Solutions corresponding to the fixed points eq. (3.9) are just the lines $t_1^j = y_1^j = 0$, in the $t_0^j - t_1^j$, and $y_0^j - y_1^j$ planes respectively. Whether a point spirals towards or away from the fixed point (or whether it moves towards or away from the fixed points along lines, in the case when $t_{1\pm}^j = 0$) depends only on the sign of $t_{0\pm}^j$. It is easily checked from eqs. (3.8), (3.9) that the sign of $t_{0\pm}^j$ has the following properties

(i) $\tau^2 \text{Sin}^2(2\pi j/n) \neq 0$

$$\begin{array}{llll} t_{0+}^j < 0, & t_{0-}^j > 0, & j = 1 \dots n/2 - 1, & n \text{ even} \\ & & j = 1 \dots (n-1)/2, & n \text{ odd} \\ t_{0+}^j > 0, & t_{0-}^j < 0, & j = n/2 + 1 \dots n-1, & n \text{ even} \\ & & j = (n+1)/2 \dots n-1, & n \text{ odd} \end{array} \quad (3.12)$$

$$(ii) \quad \tau^2 \sin^2(2\pi j/n) = 0,$$

$$\tau \neq 0, \quad t_{0+}^j > 0, \quad t_{0-}^j < 0, \quad \begin{array}{ll} j = n/2, n, & n \text{ even} \\ j = n, & n \text{ odd} \end{array}$$

$$\tau = 0, \quad t_{0+}^j > 0, \quad t_{0-}^j < 0 \quad \nu \neq 0, \quad \forall j, n$$

(3. 13)

Thus we see that in case (i), the fixed points $t_{0-}^j(t_{0+}^j)$, $j=1, \dots, n/2$, n even or $j=1, \dots, (n-1)/2$, n odd, and $t_{0+}^j(t_{0-}^j)$, $j=(n/2+1), \dots, n-1$, n even or $j=(n+1)/2, \dots, n-1$, n odd, are unstable(stable) respectively as $l \rightarrow \infty$. In the special cases where $j = n/2$ or n (n even) or $j = n$, n odd, the fixed points $t_{0+}^j(t_{0-}^j)$ are unstable(stable).

Having studied the stability of the fixed points as $l \rightarrow \infty$, we now wish to discuss the behaviour of the solutions to the R.G.E.'s (3.6). In particular, we want to know if there are strong coupling regions in the space of R.G. trajectories. By strong coupling we mean a region where a point starting near the origin is driven along a trajectory, out to indefinitely large distances, as $l \rightarrow \infty$. Of course before reaching arbitrarily large values, the couplings violate the region of validity of perturbation theory. Conversely, a weak coupling regime is where an initial point arbitrarily close to the origin, remains in its vicinity as $l \rightarrow \infty$.

At first sight one may think that a region in the space of coupling constants near an unstable fixed point would constitute strong coupling, since all trajectories move away from such a region. However, this depends on whether or not there is what one might call an 'accessible' stable fixed point nearby. Trajectories moving away from unstable fixed points then either go off to infinity (strong coupling) or terminate at an accessible stable fixed point. The latter case will constitute weak coupling because the fixed points in eqs. (3.8), (3.9) are only valid near the origin where perturbation theory holds. To understand what is meant by accessible, consider the simplest case where $j = n$, n arbitrary, which as mentioned before corresponds to the usual superfluid without adiabatic phases present. The R.G.E.'s in this case have been studied in the literature [5], [9], [10]. We have seen that fixed points only exist if $|\nu| > |\tau|$. So in the $y_0 - t_0$

plane (here we drop the superscript $j = n$ on the parameters y_0^j, t_0^j), there are 3 regions (I-III), defined by $y_0 < |t_0|$, ($t_0 < 0, t_0 > 0$) which correspond to regions (I,II), and $y_0 > |t_0|$ corresponding to III. All 3 regions are separated by the lines $y_0 = \pm t_0$, and fixed points only exist in regions I and II, along the t_0 -axis (see fig. 1).

The R.G. trajectories also fall into 3 classes depending on which of the 3 regions they are in. For initial points lying in I, there exist stable fixed points along the t_0 -axis, towards which they are driven as l increases. In the case of initial points in region II, the fixed points here are unstable and so they are driven off to infinity. They are not driven to the stable fixed points of region I because this would require the existence of trajectories that cross from region II to that of I, via III. However, such trajectories cannot exist. This is because (see for example ref [9]) the quantity $(y_0)^2 - (t_0)^2$ is a constant, C say, along trajectories, with $C > 0$ in region III while $C < 0$ in regions I and II. The separatrix between these regions is given by the curve with $C=0$. So we can say that the existence of a strong coupling region II ($t_0 > 0$ in fig. 1) is a consequence of the fact that R.G. trajectories which move away from the unstable fixed points with $t_{0+} > 0$ cannot approach the stable points $t_{0-} < 0$ which are therefore ‘inaccessible’. This itself may be thought of as a kind of topological obstruction which is connected to the property that fixed points only exist in certain regions of the $y_0 - t_0$ plane. Region III also exhibits strong coupling behaviour, which is a consequence of the absence of any fixed points at finite values of the couplings.

Let us contrast this situation with that of the next simplest case when $j = n/2$, n even. Again in this case the fixed points have the property that $t_1 = y_0 = y_1 = 0$, (for convenience we have once more suppressed the superscripts $j = n/2$ in the y and t parameters.) However, there is a crucial sign change in the equation for t_0 at the fixed points (see eq. (3.9)), namely $t_{0\pm} = \pm\sqrt{\nu^2 + \tau^2}$. This means that $t_{0\pm}$ exists for all initial values of y_0, t_0 , (subject to the restriction that such initial values are close to the origin to justify a perturbative expansion of the R.G.E.’s). Therefore we see that there is no obstruction in this case for trajectories which move away from the unstable fixed points having $t_{0+} > 0$ to approach asymptotically the stable points $t_{0-} < 0$, and not go off to infinity as in the previous example.

In fig. 2 the R.G. trajectories are plotted for various initial conditions, and we see that they are sections of semi circles of radius $\sqrt{\nu^2 + \tau^2}$. These are in fact exact solutions to the lowest order R.G.E.’s (3.6) as was shown in ref [1] for the case where

$j = 1, n = 2$ but which also applies to the case $j = n/2, n$ even. They confirm the behaviour previously described so that no strong coupling is found, in contrast to that of fig. 1.

We now move on to consider the remaining cases, namely $j = 1 \dots n-1, n$ odd and $j = 1 \dots n-1, (j \neq n/2), n$ even. Unfortunately we have not been able to find analytic solutions in these cases, (other than those of the linearized equations described earlier, which are valid near the fixed points). However we shall present numerical solutions for the first few values of j, n , and it will become apparent that an extrapolation based on these solutions to general values of j, n is possible. From the formula for the fixed points Eqs.(3.8), (3.9) (for $\tau \neq 0$), we see that since $\Delta \geq 0$ and $(\sqrt{\Delta} - 4\nu^2 + 4\tau^2 \cos(2\pi j/n)) \geq 0$, fixed points again exist for all initial conditions. We should expect therefore, that the R.G. trajectories, which we may picture as simultaneous curves in the $t_0^j - y_0^j$ and $t_1^j - y_1^j$ planes have similar properties to those in the case $j = n/2, n$ even, described above.

Figs. 3a,b show the trajectories for $n = 3$ and $j = 1$. We have not plotted the trajectories when $j=2$ because they are trivially obtained from those in the $j = 1$ case by the reflection $t_1^j \rightarrow -t_1^j, y_1^j \rightarrow -y_1^j$. This is a consequence of the constraints satisfied by the solutions as discussed earlier. Initial values all lie on the t_0^j and y_0^j axes since the boundary conditions dictate that $t_0^j(l=0) = y_0^j(l=0) = 0$. In fig. 3a we again see the trajectories moving away from the unstable fixed points having $t_{0-}^{j=1} > 0, t_{1+}^{j=1} > 0$ and terminate at the stable fixed points $t_{0+}^{j=1} < 0, t_{1+}^{j=1} > 0$. Note that although there are in fact two possible values of $t_{1\pm}^j$ lying in the upper and lower half planes for each value of j , only $t_{0+}^j > 0$ is relevant for $j = 1$ since the trajectories all lie in the upper half $t_0^j - t_1^j$ plane. Equally, for the $j = 2$ case (which we have not plotted), the relevant fixed point has $t_{1-}^{j=2} < 0$ since all trajectories lie in the lower half plane. In fig. 3b the trajectories begin and end on the y_0^j axis. The spiral nature of the trajectories near the fixed points is more pronounced in fig. 3b than in 3a. This is because the speed at which the trajectories approach the fixed points is exponentially faster in the former case compared to the latter (see eqs. (3.11)).

Further plots are given in figs.4 and 5. Figs. 4a,b illustrate the solutions for $n = 4$ and $j=1$. The case $j = 2$ is equivalent to $j=n/2, n$ even, as plotted in fig 2. Again the case where $j = 3$ can be derived from that of $j = 1$ by the usual inversion and is not plotted. $j = n = 4$ is of course equivalent to that illustrated in fig. 1 since the adiabatic

phase is just unity in this case. Figs. 5a,b correspond to $n=5, j=1$, while those of 5c,d the case $n=5, j=2$. In all of the plots corresponding to non-trivial adiabatic phases, we see a similarity in behaviour namely the lack of strong coupling regions. All trajectories flow to stable fixed points, which for consistency have to be near the origin. These observations together with our previous findings about the existence of fixed points for general n, j lead one to conclude that the only case where strong coupling is observed is when $j=n$, i.e. when adiabatic phases are absent. It should be mentioned that we have also found numerical solutions for the next to leading order R.G.E.'s which are obtained from the corresponding equations for the S-G model [9]. We have found no significant departures from the solutions of the lowest order equations given in figs.2-5 (this was already known for the usual S-G R.G.E.'s [9]) which therefor remain qualitatively correct.

Note that the apparent crossing of separate trajectories in some of the plots is an artifact of their projection onto the $t_0^j - t_1^j$ and $y_0^j - y_1^j$ hyperplanes. The system of equations (3.6) constitute what is termed an autonomous system in the literature (see e.g. [11], chapter 9), since the r.h.s have no explicit l dependence. Such systems are known to have the property that (in the present context) every point in the 4-dimensional space of couplings $t_0^j, t_1^j, y_0^j, y_1^j$ for fixed j , lies on a unique trajectory.

With these results in mind, let us now return to the original goal of this section which was to calculate the superfluid density from the formula given in eq. (2.23) of section 2. In terms of the real parameters t_0^j this formula can be rewritten as

$$\begin{aligned}
\frac{\rho_{s.f.}}{\rho_0} &= \lim_{l \rightarrow \infty} \frac{4T}{\alpha n} \left[\sum_{j=1}^{(n-1)/2} 2(1 - t_0^j(l)) + (1 - t_0^{j=n}(l)) \right] \quad n \quad \text{odd} \\
&= \lim_{l \rightarrow \infty} \frac{4T}{\alpha n} \left[\sum_{j=1}^{n/2-1} 2(1 - t_0^j(l)) + (1 - t_0^{j=n/2}(l)) \right. \\
&\quad \left. + (1 - t_0^{j=n}(l)) \right] \quad n \quad \text{even}
\end{aligned} \tag{3.14}$$

Fortunately, the formula in eq. (3.14) for the superfluid density does not involve the explicit solutions $t_0^j(l)$ but rather their asymptotic limit as $l \rightarrow \infty$. Our previous conclusion concerning the R.G. trajectories was that for $j=1 \dots n-1$ and arbitrary initial

conditions (always subject to being within the region of validity of the lowest order R.G.E.'s), points are driven along trajectories which finally end on the stable fixed points, (listed in eqs. (3.12), (3.13)) as $l \rightarrow \infty$. Moreover, we have not observed any non-analytic behaviour in the solutions as the initial conditions are varied.

On the other hand, the solutions for $j = n$ correspond to the situation studied previously in the literature, since this is the case when the adiabatic phase is absent. The solutions are well known to show non-analytic behaviour in the $l \rightarrow \infty$ limit as the initial point in the $y_0 - t_0$ plane (see fig. 1) crosses from say region I into region III which are separated by the line $y_0 = -t_0$. As one approaches this line from region I, $t_0(\infty) \rightarrow 0$. Approaching the same line from region III, it has been shown previously [5,7] that $t_0(\infty) \rightarrow 1$, since in this region the corresponding Sine-Gordon field theory has a non-vanishing mass gap.

Hence, $t_0(\infty)$ shows a discontinuity at $T = T_c(z)$ where the latter line of critical temperatures corresponds to the line of initial values $\tau(z, T) = -\nu(T)$ or equivalently to $T_c(z) = \frac{\alpha}{2}(1/2 - \pi z)$. This discontinuity manifests itself as a discontinuity in $\rho_{s.f.}$ at T_c . In the $n=1$ case, the only contribution to $\rho_{s.f.}$ comes from $t_0^{j=n=1}$ in eq. (3.14), and the corresponding jump was first described by Nelson and Kosterlitz [5] (but see also ref [7]). For $n > 1$, $t_0^{j=n}$ is only one of n different contributions to $\rho_{s.f.}$ in eq. (3.12), but it is the only one showing non-analytic behaviour in the initial conditions ν, τ .

Thus, the discontinuity in $\rho_{s.f.}/T$ at $T = T_c$ which can be derived from eq. (3.14) is

$$\lim_{T \rightarrow T_c^-} \left(\frac{\rho_{s.f.}}{T} \right) - \lim_{T \rightarrow T_c^+} \left(\frac{\rho_{s.f.}}{T} \right) = \frac{2m^2}{\pi \hbar^2 n} \quad (3.15)$$

For normal superfluids i.e. $n=1$, the size of the jump in $\rho_{s.f.}/T_c$ is $2m^2/\pi \hbar^2$ and is universal since it only depends on fundamental constants and the mass m of the superfluid particles, (e.g. He^4 atoms in the case of superfluid helium layers).

Thus our first conclusion concerning the superfluid density is that the presence of $\mathcal{Z}_n$ adiabatic phases in the vortex partition function as described in the last section, reduces the size of the jump at $T_c(z)$ by a factor of $1/n$ compared to the case when such phases are absent. We see from eq. (3.15) that the size of this jump is still universal.

The second, and somewhat surprising conclusion concerns the nature of $T_c(z)$ itself. For the case of normal superfluids, $T_c(z)$ is the critical temperature beyond which the

Kosterlitz-Thouless phase transition occurs. The system of vortices changes from a medium of bound vortex-antivortex pairs ($T < T_c(z)$) to one of free vortices and antivortices ($T > T_c(z)$). This transition reflects itself in the behaviour of $\rho_{s.f.}$ which is finite for $T < T_c(z)$ but which vanishes discontinuously for $T > T_c(z)$, i.e. the free vortices and antivortices completely destroy the superfluid condensate. However for $n > 1$, we find that when $T > T_c(z)$ (but still close to $T_c(z)$), $\rho_{s.f.}$ is non-vanishing because of the contributions from $t_0^j(\infty)$, $j = 1 \dots n-1$ in eq. (3.14). Since $t_0^j(\infty)$ are just given by their values at the stable fixed points, the superfluid density for $T > T_c(z)$ becomes

$$\begin{aligned}
\frac{\rho_{s.f.}}{T} &\approx \frac{4m^2}{\pi\hbar^2 n} \sum_{j=1}^{(n-1)/2} \left[1 + \frac{(\sqrt{-4\nu^2 + 4\tau^2 \cos(2\pi j/n)} + \sqrt{\Delta})}{8\sqrt{2}\tau^2 \sin(2\pi j/n)} \right. \\
&\quad \times \left. (4\nu^2 - 4\tau^2 \cos(2\pi j/n) + \sqrt{\Delta}) \right] \quad n \text{ odd} \\
&\approx \frac{4m^2}{\pi\hbar^2 n} \left(\sum_{j=1}^{n/2-1} \left[1 + \frac{(\sqrt{-4\nu^2 + 4\tau^2 \cos(2\pi j/n)} + \sqrt{\Delta})}{8\sqrt{2}\tau^2 \sin(2\pi j/n)} \right. \right. \\
&\quad \times \left. \left. (4\nu^2 - 4\tau^2 \cos(2\pi j/n) + \sqrt{\Delta}) \right] + (1 + \sqrt{\nu^2 + \tau^2}) \right) \quad n \text{ even}
\end{aligned} \tag{3.16}$$

where Δ in eq. (3.16) was defined in eq. (3.8). It is easily verified that the r.h.s. of eqs.(3.16) are non-negative for arbitrary values of ν and τ as should be the case since $\rho_{s.f.}$ is a density.

So, while $T_c(z)$ (as in the usual case) marks the point at which the superfluid density undergoes a discontinuity, the latter does not vanish for $T > T_c(z)$. In the discussion that follows in the next section, we shall see how one can understand these results in terms of the screening length of the system and how they modify the usual picture of charge unbinding.

Discussion

To understand the results of the previous section, it is helpful to consider the notion of the screening length of the system. For a normal Coulomb gas of free charges, the screening length and screened potential have been discussed widely in the literature (see e.g. ref [12]). In appendix A of [1], these quantities were discussed in the case where $\mathcal{Z}_2$ -valued adiabatic phases were also present. The present case is a straightforward generalization of this to include $\mathcal{Z}_n$ -valued phases, and we shall simply state the results.

The physical linearly screened potential associated with the vortex system, or rather its momentum space representations $V_L^{\text{phys}}(k, \lambda_{\text{phys}})$, turns out to be proportional to the sum over Sine-Gordon type two point correlations given on the r.h.s. of eq. (2.20) of section 2. Here, λ_{phys} is the physical screening length of the system. We can write the small momentum dependence of the screened potential as $V_L^{\text{phys}}(k, \lambda_{\text{phys}}) \approx 2\pi/(k^2 + \lambda_{\text{phys}}^{-2})$. For the simplest case $n = 1$, where there is only one contribution to the sum in eq. (2.20), the quantity $\lambda_{\text{phys}}^{-2}$ can be associated with the mass gap of the corresponding Sine-Gordon model. A zero mass gap is equivalent to an infinite screening length in the vortex system, in which case there is a non-vanishing contribution from the r.h.s. of eq. (2.20) in the limit $k^2 \rightarrow 0$. This case describes the region $T < T_c(z)$, and a non-vanishing $\rho_{\text{s.f.}}$. On the other hand for $T > T_c(z)$, a mass gap develops so λ_{phys} is finite and the superfluid density vanishes. In the former case where the screening length is infinite, the potential between vortices which is proportional to $V_L^{\text{phys}}(|\mathbf{x}|)$, falls off logarithmically and so maintains long range order (i.e. the superfluid condensate). In the latter case such order is destroyed since the potential falls off like $e^{-|\mathbf{x}|/\lambda_{\text{phys}}}$ for $|\mathbf{x}| \gg \lambda_{\text{phys}}$.

The situation where $n > 1$ is more complicated in that there are now n contributions to the sum in eq. (2.20) and λ_{phys} is no longer directly related to the mass gap of any one particular Sine-Gordon model. This is to be expected since the correlation functions for $n > 1$ are in general complex, satisfying the conjugation relations given in eq. (2.21), whilst λ_{phys} is of course real. Thus even though the $j = n$ contribution involves two point correlations of the standard Sine-Gordon theory with a (non)zero mass gap for ($T > T_c(z)$) $T < T_c(z)$, the physical screening length in fact remains infinite in both these cases. This is because, as our R.G. analysis has shown, there

are non vanishing contributions from the $j = 1 \dots n-1$ other terms in eqs. (2.20), (2.23) in both of these temperature ranges, with no non-analytic behaviour observed in going from $T < T_c(z)$ to $T > T_c(z)$. The fact that the physical screening length remains infinite, at least for temperatures close to the critical line, means that the potential between vortices remains logarithmic and so maintains a non-vanishing condensate, i.e. long range order.

Having said this, there is nevertheless a jump in the density $\rho_{s.f.}$ at T_c even for $n > 1$. From the above arguments we conclude that the inter-vortex potential remains logarithmic before and after the discontinuity occurs, but that there is an effective dielectric function, which is discontinuous at T_c . That is, for $k^2 \approx 0$ we can write $V_L^{\text{phys}}(k, T) = 2\pi/(\epsilon(T)k^2)$ with $\epsilon(T)$ a temperature dependent (dimensionless) dielectric function. From the size of the jump in $\rho_{s.f.}$ as calculated previously, (see eq. (3.14)), it follows that $1/\epsilon(T = T_c^+) - 1/\epsilon(T = T_c^-) = 1/n$.

This latter property of the inter-vortex potential suggests that what may be described as ‘incomplete’ screening takes place in the presence of adiabatic phases. The effective dielectric function is constant at long wavelengths, and has the discontinuity at $T_c(z)$ as described. By contrast, the effective dielectric in the absence of such phases has a very different long wavelength behaviour, namely $1/\epsilon(k, T) \approx k^2/(k^2 + \lambda_{\text{phys}}^{-2})$ for $T \rightarrow T_c^+$, and $1/\epsilon(k, T) \approx 1$ for $T \rightarrow T_c^-$.

A final comment in the comparison of the physics in the presence or absence of adiabatic phases, concerns the heuristic arguments based on an analysis of the free energy, which have been used previously to anticipate the K-T phase transition [4]. In the argument, one considers a free vortex-antivortex pair, whose total energy is then just the sum of their self energies (assuming the system is finite in extent-i.e. before taking the thermodynamic limit). Whilst self energies become logarithmically large as the size of the system is increased, there is an additional (negative) logarithmic entropy term in the free energy which can compensate. Thus it is argued that when the entropy is sub dominant to the self energy, it is statistically unfavorable to form free vortex-antivortex pairs in the thermodynamic limit (the free energy is positive and logarithmically divergent in this limit), whilst the opposite is true when the entropy is dominant. The transition between the two regimes marks the K-T phase transition, and indeed this simple argument does reproduce the correct value of T_c , but only for $z = 0$ [4].

Since, as we have seen earlier (eq. (2.2)) the energy of a neutral pair of vortices is unaffected by the presence of adiabatic phases (strictly speaking in the thermodynamic limit), it might appear at first sight that our conclusion that no charge unbinding takes place at T_c is inconsistent with free energy considerations. However, it should be remembered that in the partition function $Z = \frac{\Omega}{n} \sum_{j=1}^n Z_j$, from which the density $\rho_{s.f.}$ is derived, there is a phase factor $e^{i\pi j N/n}$ present in Z_j . Thus in computing the contribution of a free vortex-antivortex pair to the partition function, apart from the usual integration over the centre of mass position of each vortex, there is a sum over n different ‘sectors’, weighted by the above phase. Now precisely because the energy of the vortex-antivortex pair is independent of n , we see that the resulting contribution is proportional to $\sum_{j=1}^n e^{2i\pi j/n}$ which is therefore zero. The conclusion is that the free energy of an unbound neutral pair of vortices is infinitely large and positive when the adiabatic phase is present, and hence charge unbinding is suppressed. The reason why the free energy is divergent is because the arguments we have just presented only really hold in the thermodynamic limit (in particular as we have mentioned before, the vanishing of interactions between thermally created vortices and the non-thermal vortex at the boundary only strictly holds in this limit). So we see that the results of the previous section concerning the absence of charge unbinding, which were obtained through R.G. methods, are also consistent with heuristic free energy arguments.

The conclusion that the critical line $T = T_c(z)$ does not correspond to the superfluid/normal fluid phase transition begs the question at what temperature does this transition take place when adiabatic phases are present? Unfortunately, it seems that an answer to this question requires going beyond the perturbative R.G. approach adopted in this paper. Further complications could also arise if this new critical temperature, corresponding to charge-unbinding in the vortex system, is high enough to excite quasi-particles in the superfluid. In this case, the implicit assumption made in the derivation of the vortex partition function of eq. (2.1) [1], that one can ignore quasi-particle contributions, is no longer valid.

Figure Captions

Figure 1- Figure 1 is a plot of the R.G. trajectories in the parameters t_0^j, y_0^j for $j = n$, and n arbitrary. A representative trajectory is illustrated for each of the three regions I-III described in the text. Arrows on these (and later) trajectories indicate the direction an initial point moves as l increases.

Figure 2- Figure 2 plots the R.G. trajectories in the variables t_0^j, y_0^j for $j = n/2$, n even, for various initial conditions.

Figures 3a,b- Figures 3a and 3b are plots of the R.G. trajectories in the 4 variables $(t_0^j, t_1^j, y_0^j, y_1^j)$ for $j = 1, n=3$, projected onto the $t_0^j - t_1^j$ and $y_0^j - y_1^j$ planes, again for various initial conditions.

Figures 4a,b -Figures 4a and 4b are as in 3a, 3b but with $j = 1, n=4$.

Figures 5a-5d -Figures 5a and 5b are as in 3a, 3b but with $j=1, n=5$; Figures 5c and 5d are the same plots with $j=2, n=5$.

References

- [1] S. Thomas, *A Modified Kosterlitz-Thouless Phase Transition in 2-Dimensional Vortex Systems* CERN preprint CERN-TH. 6450/92.
- [2] M.V. Berry, *Proc. R. Soc. London Ser. A* **392** (1984) 45; B. Simon, *Phys. Rev. Lett.* **51** (1983) 2167.
- [3] F.M. Haldane and Y-S. Wu, *Phys. Rev. Lett.* **55** (1985) 2887. (1984) 722.
- [4] J.M. Kosterlitz and D.J. Thouless, *J. Phys.* **C6** (1973) 1181.
- [5] D.R. Nelson and J.M. Kosterlitz, *Phys. Rev. Lett.* **39** (1977) 1201.
- [6] P. Minnhagen, *Rev. Mod. Phys.* **59** (1987) 1001.
- [7] P. Minnhagen and G.G. Warren, *Phys. Rev.* **B24** (1981) 2526.
- [8] J. Frolich in *Renormalization Theory*, Proc. of the NATO Advanced Study Institute, Erice 1975, eds. G. Velo and A.S. Wightman (Reidel, Dordrecht/Boston 1976, p.371); S. Samuel, *Phys. Rev.* **D18** (1978) 1916.
- [9] D.J. Amit, Y.Y. Goldschmidt and G. Grinstein, *J. Phys.* **A13** (1980) 585.
- [10] J.M. Kosterlitz, *J. Phys.* **C7** (1974) 1046.
- [11] *Elementary Differential Equations and Boundary Value Problems* W.E. Boyce and R.C. DiPrima, (John Wiley, New York 1969).
- [12] P. Minnhagen, *Phys. Rev.* **B23**(1981) 5745.

Fig. 1

Fig. 2

Fig. 3a

Fig. 3b

Fig. 4a

Fig. 4b

Fig. 5a

Fig. 5b

Fig. 5c

Fig. 5d