The π NN Coupling Constant*

Torleif E. O. Ericson

CERN, CH-1211 Geneva 21, Switzerland

Abstract

A report on a panel discussion exploring the possible sources of the small value for the coupling constant recently reported by several groups, the principal quantities constraining the coupling constant as well as the actions which should be taken to elucidate the situation.

CERN-TH. 6405/92
February 1992

---

*To appear in Nuclear Physics A, Proceedings of the XIIIth Conference on Few-Body Problems in Physics, Adelaide, Australia, January 5-11, 1992.

1. INTRODUCTION

The purpose of the panel session was to critically examine the present evidence for the value of f^2 , to identify the critical constraints both from theory and from the data as well as to identify the most profitable next steps which will clarify the situation. It was originally planned that the principal experts would all be able to contribute to the discussion. Regrettably four of them were unable to attend, namely Profs. Arndt, Bugg, Höhler and de Swart. Their views may not be correctly reflected in the following.

For some years the Nijmegen group¹ has consistently reported that their semi-phenomenological phase shift analysis of pp and np scattering scattering data as well as recently a similar analysis of the less accurately known antiproton charge exchange reaction on the proton all indicate that the πNN coupling constant is substantially smaller by approximately 5% than the classical value $f^2 = 0.079(1)$ deduced from πN scattering and from the pp forward dispersion relation. Their analyses have a quite low χ^2 and should thus describe the data well.

Recently, the Virginia Polytechnical Institute (VPI) group of Arndt and coworkers² have produced new sets of phase shift of πN scattering incorporating recent experimental data. These phase shifts have been used to produce the input for the so-called 'Hüperplot' for the spin flip πN amplitude for given t both for π^+ and π^- . They obtain by this interpolation method $f^2=0.0735(15)$ at $t=-0.2 \text{ GeV}^2$. This value for the coupling constant reported is even smaller than the one reported by the Nijmegen group and at first glance seems to confirm their conclusion. These results have caused considerable concern. The reason is that f^2 is a crucial constant for a several other relations measured to very high precision. In addition it is a scale factor in some of the most accurately measured and investigated quantities in nuclear physics. The issue of its value must necessarily be seen in a broader context. The following are the key areas to be watched:

- the Goldberger-Treiman relation between the weak axial coupling constant g_A and the πNN coupling constant f , both taken at zero momentum transfer (and not at the pion-nucleon pole at $t=\mu^2$, where μ is the pion mass). The standard values for the coupling constant and the pion decay constant describe the relation to $(3.5 \pm 1)\%$ without any form factor corrections; the older discrepancies have gradually decreased over the years with new values for the input. Recently a new correction to the usual value for the pion decay constant as deduced from experiments further reduces the discrepancy by another percent³. It is therefore easy with any reasonable form factor effect to come within 2 standard deviations or less of fulfilling the relation. The newly proposed values would do even better and yield near perfect agreement with no form factor included. On the other hand, in certain non-linear σ -models this would even overshoot the goal due to small additional quark mass terms. Thus the Goldberger-Treiman relation is not a crucial constraint in spite of its precision.

- the threshold photo-production cross sections of charged pions are known to a few % and are well described by the Kroll-Ruderman theorem. This means they are proportional to f^2 . A change in this quantity by 5% therefore provokes a discrepancy in both the π^+ and π^- cross sections by about 2.5 standard deviations in the same direction, i.e. by about 3.5 standard

deviations. This is an uncomfortable number and would therefore require an explanation if the smaller coupling is accepted.

- similarly, constraints are imposed by the accurately known quadrupole moment of the deuteron and its D/S ratio, which are linearly related.

In order to define the constants to be discussed, I first remind that it is convenient to use the πNN coupling with the following charge independent definition (μ_c is the charged pion mass and M the nucleon mass) as schematically indicated in Figure 1.:

Figure 1. The pseudoscalar and pseudovector couplings.

As far as the determination of the coupling constant is concerned the two are completely equivalent and the relation $g/2M=f/\mu_c$ holds. A word of warning: in the literature one will often see the pseudovector coupling constant for π^0 proportional to the π^0 mass and not to μ_c ; this is so because with the definition above a factor μ_c/μ_c is added separately. The two definitions are both charge independent, but it is important not to mix them in taking information from different sources! In the discussion on the πNN coupling constant there is presently no indication of any charge dependence of the coupling. To avoid any possible confusion: this issue does not enter the present discussion and the various groups agree on this point. In an earlier stage of the discussion this possibility was evoked by the Nijmegen group, but the ground has now shifted.

2. INFORMATION FROM THE πN INTERACTION

Dr. Locher first reviewed the methods for deducing f^2 from πN data, concentrating in particular his discussion on the recent VPI analysis. The analysis is based on the spin flip amplitude $B(v,t)$, which nearly everywhere has a dominant or, at least, an important contribution, from the direct and crossed nucleon Born terms, i.e. from the terms which give f^2 . The amplitude is taken at fixed t and corrected by a dispersion relation. This produces a linear relation with v (for π^- the negative of v is used). The crucial data with high precision come from the resonance region and are dominated by the accurately known phase shift combination $P_{33}-P_{31}$. The intercept with $v=0$ should give f^2 independent of t in this Hüperplot representation.

Figure 2. Typical Hüperplot for the determination of f^2 .

Dr. Locher reported that he had access to the VPI phase shifts and that he therefore had been able to repeat their analysis not only at $t = -0.2 \text{ GeV}^2$, but also for other values of t verifying the consistency of the results. The conclusion is that f^2 varies substantially with t in a range from about 0.072 to 0.082 and that the value given for $t = -0.2 \text{ GeV}^2$ is one of the lowest. The error of ± 0.0015 seems therefore strongly underestimated and should rather be ± 0.0050 . Several persons criticized the VPI determination arguing that it was in principle unhappy to deduce the Born residue from a phase shift analysis in which the Born terms do not appear explicitly. In my opinion this criticism is as such unjustified, since f^2 is extracted to $\pm 7\%$ precision which is not so bad after all, but it is likely that precision can be improved by including the Born terms in the phase shift analysis. Similar conclusions to the ones of Dr. Locher have also been reached by Prof. Höhler.⁴

Dr. Locher had in addition evaluated the forward charge exchange dispersion relation using the VPI phase shift solution as input. Here, the isovector scattering length and a total cross section integral dominated by the Δ region contribute about equally and add coherently; the cross section integral should be quite accurately determined and the crucial point is in the scattering lengths. According to Dr. Locher the VPI solution reproduces in this case the standard value of 0.079, essentially demonstrating that it is consistent with empirical values for scattering lengths

and total cross sections. The conclusion is therefore that the VPI analysis cannot be used to support the claim of a low value for f^2 .

Next, the input data in the analysis of πN scattering analysis was examined. Prof. Ristinen gave a comprehensive survey of the experimental data. Every analysis depends critically on the very high precision data on elastic, charge exchange and total cross sections in the region 100-250 MeV determined by Bugg et al. in the early '70's at CERN. While the high quality of these data is certain, the error bars are extremely small, reaching even 0.3% precision in some cases. Not only do these data give an overwhelming contribution in any χ^2 analysis, but no experimental group since has been able to reach such a precision in spite of the knowledge of these experiments and in spite of the developments in techniques from that time. He compared the measurements with the predictions of the Karlsruhe analysis and demonstrated that the cross sections and analyzing powers indicate numerous deviations from these below 100 MeV. In the resonance region he drew attention to the fact that the experiments of Pedroni et al. were systematically 1-2% below those of Bugg et al., although the precision indicated was less. In the discussion Dr. Sainio in particular pointed out that the cross sections of Bugg et al. agree with the unitarity limit at resonance to 0.5%, so that at least at this point there is an independent justification for his cross sections. He also made it quite clear that data below 100 MeV could not be effectively used at present for the analysis leading to the extraction of the σ -term. While this had little to do with the present discussion of the coupling constant, the discrepancies and inconsistencies between the different groups was not theoretically acceptable.

Dr. Seviør presented the πN situation at 67 MeV and showed that the differential cross sections were in good to approximate agreement with the single energy analysis of Brack et al., while there were very large deviations in the asymmetry parameter A_y . This would be investigated at Los Alamos, TRIUMF and PSI in the next two years.

From the discussion of the experimental data we can conclude that the forward dispersion relation for isovector πN scattering can also be directly evaluated using the measured cross sections and the experimental isovector scattering length. The cross sections can marginally decrease f^2 by 1% if they are wrongly normalized by 2% in the Δ region; if one uses the value of the isovector scattering length from the Panofsky ratio of 0.088(2) instead of 0.091(2), the coupling constant would decrease but not more than by 2% at most from its classical value. This would then be in minor conflict with the corresponding scattering length from the πp atom. It seems difficult to believe from the data that the coupling constant can be lowered by more than a few % at the very most.

3. INFORMATION FROM THE NN INTERACTION

The discussion then turned to the information from NN scattering. Dr. Haidenbauer first gave a survey of the assumptions in the Nijmegen partial wave analysis. It uses a relativistic Schrödinger equation with an energy dependent potential

$$V = V_{EM} + V_{OPE} + V_{HBE} + V_{SR}.$$

At short range it is parametrized for each wave separately by an energy dependent square well potential of range b (typically and mostly $b=1.4$ fm); the electromagnetic part includes Coulomb, magnetic moment interaction and vacuum polarization, which are important at low energies. The key part for the present discussion is the One-Pion-Exchange (OPE) potential

$$V_{OPE} = 1/3 \left(f^2/4\pi \right) (\mu/\mu_c) M/E \, e^{-\mu r} / r \left\{ \sigma_1 \sigma_2 + S_{12} [1 + 3/(\mu r) + 3/(\mu r)^2] \right\}.$$

This potential is applied outside the distance b of the square well region. For this OPE potential the nucleon finite size is accounted for by the Gaussian form factor

$$F(k^2) = \exp[-(k^2 + \mu^2)/\Lambda^2]$$

chosen so that $F(-\mu^2)=1$ at the pion pole; the results are nearly insensitive to this form factor. In addition, outside the distance b the OPE potential is supplemented by V_{HBE} given by the intermediate range Nijmegen Heavy Boson Exchange soft core potential⁵ or, nearly as well, by the Paris non-OPE potential⁶. This approach leads to the best χ^2 in any of the analyses of pp and np scattering data, in both cases with a small value for f^2 . Dr. Haidenbauer first asked the question: what possibilities are there to fit the data for pp scattering and the deuteron observables when f^2 is varied? For this he reported on the results of an investigation with K. Holinde and A. Thomas⁷ based on a modified Bonn potential which in particular used a relatively soft πNN vertex with $\Lambda=800$ MeV/c and a rather heavy π meson. He showed that by varying the π contribution it was possible to accommodate changes of the coupling constant of the magnitude suggested by the Nijmegen group particularly in the case of the deuteron observables.

Several participants took up the point of the π meson which is introduced *ad hoc*. Its role in practice is clearly to compensate for the soft πNN vertex, so that it partly simulates a hard form factor, an effective result for the pion dominated tensor interaction on which there has always been an insistence on from specialists on the Bonn potential so as not to spoil the $^3D_1$ phase shifts and as well as from the deuteron specialists so as not to spoil the D/S ratio and the quadrupole moment. Although this point was conceded by the authors, Prof. Thomas made the point that the effective hardening of the form factor did not mean that the interaction was effectively that of a very hard form factor. Instead there was still an effect of about 5-7 % in the critical region 1.5-2 fm. All this means that the NN force in this area is not as well pinned down via the 2π dispersion approach as is generally believed. Further, he drew attention to the fact the 2π dispersive approach à la Paris always subtracts the iterated OPEP from the weight function and that this subtraction procedure is an important one in some channels; in the presence of a form factor it is then reasonable to make this subtraction including the form factor in the vertices and this is then physically a separate issue from the phenomenological π . In the absence of

some of the main protagonists on this point the further discussion of the significance of the π meson ended without a clear conclusion, but Prof. Thomas point is doubtlessly of importance..

Dr. Kukulin pointed out the deuteron observables provided a very strong constraint on the coupling constant and illustrated this with the results from the Moscow group. In the discussion I pointed out that while this is undoubtedly the case as such, it is essential to locate exactly the source of the dependence with f^2 and to quantify it. This can be done using the reduced representation of the quadrupole moment and the D/S parameter, carefully keeping the effective range at its experimental value⁸. For a vast class of models, in particular for all reasonable energy independent potentials, there is a linear relation between these quantities to very high precision which permits one to state that a change in f^2 by 2% leads to a 1% change of the D/S ratio keeping the quadrupole moment at its experimental value. For example the model of Haidenbauer et al. follows approximately this relation. The problem is that while the D/S ratio is claimed to be measured to about 2% precision, different experimental approaches do not agree to this precision. The constraint on the coupling constant from this source is thus not very strong, although it is certainly true that the deuteron is a very nice example of pion physics.

Prof. Thomas and Prof. Bugg emphasized the importance of the πNN form factor for the conclusion of the Nijmegen group. In particular, Dr. Bugg in a comment transmitted before the session to me in view of the upcoming discussion stated that it is important to make the difference between determinations at the pole and in the physical region of t . In particular, when he analyzes data using no form factor, his own experience is that his fits of the neutral coupling constant using pp data in the region 30 MeV and of 210 to 515 MeV give values consistent with the Nijmegen value. He is confident, however, that this is a form factor effect. His statement seems fair enough, since data on total cross sections in these energy regions are major ingredients in the forward dispersion relation determination of the neutral coupling constant by extrapolation to the near-lying π^0 pole (see Figure 3). His statements are completely in line with the emphasis that has previously been given to the form factor effect by Drs. Holinde and Thomas⁹, in the context of the earlier suggestion by the Nijmegen group that the $pp\pi^0$ had a large charge dependence, now replaced by the suggestion of a generally smaller coupling.

Figure 3. Extrapolation of the singlet discrepancy function to the π^0 pole.

To this the Nijmegen group would undoubtedly answer that they have explored the sensitivity to the form factor by varying the cut-off parameter Λ from 500 MeV to ∞ in a Gaussian representation $\exp[-(k^2+\mu^2)/\Lambda^2]$ with practically no effect. The crux of the difference at this point is that practically every other group uses a parametrization $[1+(k^2+\mu^2)/\Lambda^2]^{-1}$, which for low k^2 is identical to the Nijmegen one but which falls off far less rapidly with k^2 . The effect of this in r -space is major, particularly in the tensor force, which is affected strongly⁸. For example, for $\Lambda=500$ MeV the tensor force is reduced by over 30% at 2 fm and even at 3 fm the reduction is still 10%; for $\Lambda=800$ MeV the reduction is still about 15% at $r=2$ fm (see Figure 4)!

It is thus clear that if the real form factor has a monopole shape, the Nijmegen group would unwittingly determine a too weak OPE interaction and identify it with the asymptotic pion field in the belief that they would be unaffected by the form factor. In spite of some statements to the contrary, it is important to realize that it is not technically necessary to extrapolate analytically to the pion pole position and the Nijmegen group do not do this; it is sufficient to identify correctly the coefficient of the asymptotic pion field as $r \rightarrow \infty$, which is completely equivalent. It would therefore seem that the whole discussion of the value of coupling constant depends critically on the shape of the form factor. From the discussion it was obvious that very strong opinions are held by various groups on this issue. It will not be easy to reconcile the disagreeing groups and obtain the degree of consensus necessary for the determination an objective physical quantity like the coupling constant. I therefore suggested an objective physical test, that should avoid subjective judgements. The Nijmegen phase shifts could be taken to represent set of pseudo-data from which the physical observables can be explicitly constructed. One can therefore explicitly

calculate the Coulomb corrected pp total cross sections for different spin orientations. These are exactly the crucial data that in a particular combination, enter the pole extrapolation

Figure 4. Dependence of the OPE tensor potential in r -space on the monopole formfactor.

procedure which usually determines the neutral coupling constant. The procedure uses the so called discrepancy function which also marginally uses information on total cross sections at high energy, but this is hardly an important source of uncertainty. Such a procedure should thus give a coupling constant, which would be an appropriate starting point for the discussion. It goes without saying that the same procedure should be used to compare the Nijmegen phase shift results also with the measured spin dependent total cross sections as well as with the direct empirical singlet discrepancy function. In view of the excellent χ^2 , one expects however *a priori* that these are well described.

Finally, the issue of the experimental data and their relation to the Nijmegen phase solution was discussed. A general complaint was that it was not easy to identify the critical data for the conclusion about the coupling constant. This complaint is only partially correct. The Nijmegen group show explicitly in the case of their analysis of pp scattering¹⁰ that the triplet P waves are

very important in the determination. Presumably the P waves in the region of 10 to 30 MeV have been very important in their earlier analysis of np data, although in the most recent work more emphasis is put on higher energy data. This seems to be borne out by a private discussion of the group with Dr. Bugg in which they have stated that the differential cross sections of Bonner et al. are an important constraint. Dr. Bugg draws attention to the fact that the np differential cross section data are subject to large systematic uncertainties of order of 7% at best. There are large (6%) systematic discrepancies between different groups and even in total cross sections. There are large systematic discrepancies of order of 6 % between measurements of total np cross sections with different techniques: a) with monochromatic beams and liquid hydrogen targets which have best systematics and worst statistics, b) with broad band beams and CH₂ targets with best statistics and worst systematics. As a consequence, the differential cross section data of Bonner et al., which are important to the Nijmegen analysis, have a normalization which is largely unknown by up to maybe 15%. These data are very useful for the separation of the relative strength of the P-waves, but not directly for their normalization, which must be obtained from some other source. There are two conclusions to be made from these statements. First, since charge independence is no longer questioned in the present context and at the present precision, the pp data can be used alone for the discussion of the coupling constant; these data are not contested and they have good precision. Second, there is clearly an issue of absolute normalization in the neutron-proton data. Probably, the Nijmegen constrains the normalization in the analysis, but the exact source of the constraint seems not to be identified. It would be extremely useful to have a few highly accurately normalized neutron-proton cross sections at some selected energies for calibration purposes which would then allow the use of high statistics methods methods calibrated to these points.

Finally, the antiproton-proton charge exchange reaction is not a very precise constraint on the coupling constant with present precision, as the Nijmegen group will doubtless readily agree. It serves mainly to demonstrate that such data can now be analyzed consistently with other information. The extracted coupling constant has a quite large error bar, in particular since the normalization of the data is probably incorrect by about 20% according to Dr. Bugg. On the other hand, Dr. Bradamante states that it should not be difficult to improve such experiments by a factor of 5 or so in the near future, which would make this reaction a valuable source of information. Drs. Haidenbauer and Holinde remarked that an overall factor M/E should be included in the Nijmegen potential in this case also and that its effect is not negligible. However, as of now the antiproton data have not yet any real bearing on the coupling constant discussion and this whole issue should be left aside for the time being.

4. CONCLUSION

- the π NN coupling constant from the VPI analysis has large systematics and inconsistencies at the level of precision which is claimed. It cannot presently be used as an argument for a weak π NN coupling.
- there are various inconsistencies in the experimental π N scattering data. The old very high precision data of Bugg et al., although accurate, are probably overclaimed in precision and there is a need for high precision modern data. The precision in the standard dispersion analysis may thus possibly be overstated by a factor of up to two.
- additional constraints are imposed by the charged photo-production amplitude at threshold and the Panofsky ratio, which do not readily admit large changes in the coupling constant.
- the controversy in the determination from NN data hinges on a different assessment of the effect and structure of the π NN form factor. It is urgent to perform the standard dispersion evaluation using the Nijmegen phase shifts as pseudodata, which yields a form factor independent result. The same procedure should also be used for an explicit comparison between the spin dependent total cross sections in this evaluation with the experimental ones most crucial to the dispersion relation.
- the precision of the np cross sections is contested and it is claimed that they are subject to large systematic uncertainties. Although not crucial to the present discussion, this point must be cleared up. It would appear that a few absolute precision calibration points for neutron-proton cross sections would greatly clarify the issue.
- the deuteron observables impose secondary constraints on the coupling constant, but these are not the most critical data at present.

- 1 R. A. M. Klomp et al., Phys. Rev. C 44 (1991) 1258 and references therein.
- 2 R. A. Arndt, Z. Li, L. D. Roper, and R. I. Workman, Phys. Rev. Lett. 65 (1990) 157.
- 3 B. R. Holstein, Phys. Lett. B244 (1990) 83.
- 4 G. Höhler, π N Newsletter No. 3 (1991) 66.
- 5 M. M. Nagels, T. A. Rijken and J. J. de Swart, Phys. Rev. D 17 (1978) 768.
- 6 M. Lacombe et al., Phys. Rev. C 21 (1980) 861.
- 7 J. Haidenbauer, K. Holinde and A. W. Thomas, Preprint ADP-91-163/T100 to appear in Phys. Rev. D.
- 8 T. E. O. Ericson and M. Rosa-Clot, Ann. Rev. Nucl. Part. Sci. 35 (1985) 271.
- 9 A. W. Thomas and K. Holinde, Phys. Rev. Lett. 63 (1989) 2025.
- 10 J. R. Bergervoet et al., Phys. Rev. C 41 (1990) 1435.