CERN/LAA/MSL/92-011

CERN-PPE/92-145

15 July 1992

A STUDY OF THE VARIOUS APPROACHES TO M_{GUT} AND α_{GUT}

F. Anselmo¹⁾, L. Cifarelli^{1,2,3)} and A. Zichichi¹⁾

¹⁾ CERN, Geneva, Switzerland

²⁾ Physics Department, University of Pisa, Italy

³⁾ INFN-Bologna, Italy

Abstract

An exhaustive study is presented of the different M_{GUT} and α_{GUT} values obtained using the renormalization group equations for the evolution of the couplings, with and without the evolution of the masses, with and without light and heavy threshold effects, using different levels of theoretical accuracy: one and two loops, simple and detailed threshold effects. A structure of the heavy threshold with mass ratios as large as 10^6 and values of $\alpha_3(M_Z)$ as high as the world average plus 2σ are needed in order to reach high values of M_{GUT} , which are still an order of magnitude below M_{Planck} . A discussion concludes the paper.

Submitted to "Il Nuovo Cimento"

1. INTRODUCTION

One of the most interesting problems in high-energy physics is the energy level where the couplings of the known $SU(3) \times SU(2) \times U(1)$ gauge forces merge, M_{GUT} , and the value of the unified coupling, α_{GUT} . The results from “low-energy” data indicate, for the convergence of the couplings $(\alpha_1, \alpha_2, \alpha_3)$, an energy level $M_{GUT} \approx 10^{16}$ GeV, i.e. sufficiently away from M_{Planck} , where the gravitational coupling becomes strong. The interest in this gap sparks from the fact that it could have consequences on the energy range accessible to present, forthcoming and future experimental facilities, such as LEP I, LEP II, HERA, LHC, SSC and ELN.

The structure of the paper is as follows. In section 2 the definitions and input values needed for our different approaches are given. In section 3 we give the values of M_{GUT} derived from the different approaches. In section 4 we present our conclusions.

2. DEFINITIONS AND INPUT VALUES

In order to work out the value of the energy at which the known couplings $\alpha_1, \alpha_2, \alpha_3$ should become equal,

$$\alpha(M_{GUT}) = \alpha_3(M_{GUT}) = \alpha_2(M_{GUT}) = \alpha_1(M_{GUT}), \quad (1)$$

the basic ingredient is the renormalization group equations (RGEs). There are, however, different approaches which allow all couplings to be equal (1) at a given energy, M_{GUT} . Our group has studied these approaches in order to investigate the SUSY-threshold problem [1-5]. Here we focus our attention on M_{GUT} and α_{GUT} . Let us review the different approaches aimed at deriving the values of these basic quantities.

Definitions that we will use are the following:

ΔT_H : indicates that the heavy threshold, when SUSY [$SU(5)$] is broken into SUSY [$SU(3) \times SU(2) \times U(1)$], has been taken into account (using either a simple degenerate mass or a structure with three masses).

ΔT_L : means that we have taken into account a complex structure for the light threshold effect when SUSY [$SU(3) \times SU(2) \times U(1)$] breaks either into $SU(3) \times SU(2) \times U(1)$ or into $SU(3) \times U(1)_{em}$.

EGM: indicates that the simultaneous evolution of the couplings and of the gaugino masses has been studied.

The various approaches have as experimental inputs the world average values and their limits at $\pm 2\sigma$. We have taken for the world averages and $\pm 1\sigma$ the following values [1]:

$$\begin{aligned}\alpha_3(M_Z)^{WA} &= 0.118 \pm 0.008 \\ \sin^2\theta(M_Z)^{WA} &= 0.2334 \pm 0.0008 \\ 1/\alpha_{em}(M_Z)^{WA} &= 127.9 \pm 0.2 .\end{aligned}$$

Other input values are needed for the heavy and light thresholds calculations. For the case of the heavy threshold [2, 6-7], the input values are the ratios of M_V/M_Σ , which are allowed to vary in a wide range: 10^0 – 10^6 . For the light thresholds the inputs are the two ratios of the primordial masses: $(m_0/m_{1/2})^2$ and $(m_4/m_{1/2})^2$ [4], which are allowed to vary in a range 10^{-2} – 10^2 .

All results for M_{GUT} and α_{GUT} will always be quoted with uncertainties corresponding to $\pm 2\sigma$ of the experimental inputs.

For the Standard Model with three families and one Higgs doublet [8], the following matrix coefficients will be used in the text :

$$\{b_j'\} = \begin{pmatrix} 41/10 \\ -19/6 \\ -7 \end{pmatrix} \quad (2)$$

$$\{b_{ij}'\} = \begin{pmatrix} 199/50 & 27/10 & 44/5 \\ 9/10 & 35/6 & 12 \\ 11/10 & 9/2 & -26 \end{pmatrix} . \quad (3)$$

For the Minimal Supersymmetric Standard Model (MSSM) with three families and two Higgs doublets [9], the matrix coefficients will be:

$$\{b_j^s\} = \begin{pmatrix} 33/5 \\ 1 \\ -3 \end{pmatrix} \quad (4)$$

$$\{b_{ij}^s\} = \begin{pmatrix} 199/25 & 27/5 & 88/5 \\ 9/5 & 25 & 24 \\ 11/5 & 9 & 14 \end{pmatrix} . \quad (5)$$

3. THE M_{GUT} VALUES FROM DIFFERENT APPROACHES

3a The value of M_{GUT} using the simplest approach: evolution of $\alpha_1, \alpha_2, \alpha_3$ at one loop; only one M_{SUSY} (step function)

The evolution equations are (if $\mu > M_{\text{SUSY}} > M_Z \geq m_t$):

$$\frac{1}{\alpha_1(\mu)} = \frac{1}{\alpha_1(M_Z)} - \frac{b_1^s}{2\pi} \log(\mu/M_{\text{SUSY}}) - \frac{b_1'}{2\pi} \log(M_{\text{SUSY}}/M_Z) \quad (6)$$

$$\frac{1}{\alpha_2(\mu)} = \frac{1}{\alpha_2(M_Z)} - \frac{b_2^s}{2\pi} \log(\mu/M_{\text{SUSY}}) - \frac{b_2'}{2\pi} \log(M_{\text{SUSY}}/M_Z) \quad (7)$$

$$\frac{1}{\alpha_3(\mu)} = \frac{1}{\alpha_3(M_Z)} - \frac{b_3^s}{2\pi} \log(\mu/M_{\text{SUSY}}) - \frac{b_3'}{2\pi} \log(M_{\text{SUSY}}/M_Z) \quad (8)$$

The result for M_{GUT} is as follows:

$$M_{\text{GUT}} = \left(3.1 \begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 2.2 \\ 2.6 \end{smallmatrix}\right) \times 10^{16} \text{ GeV}; \quad (9a)$$

for α_{GUT} :

$$1/\alpha_{\text{GUT}} = 22.4 \begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 4.4 \\ 3.3 \end{smallmatrix}. \quad (9b)$$

Figure 1 shows the correlation between the possible input values for $\alpha_3(M_Z)$ and the results for M_{GUT} .

3b The value of M_{GUT} using the next to the simplest approach: evolution of $\alpha_1, \alpha_2, \alpha_3$ at two loops; only one M_{SUSY} (step function)

The next-to-simplest approach is based on a new [1] numerical solution applied directly to the two-loops coupled RGEs for the $\alpha_1, \alpha_2, \alpha_3$ evolution (previous approaches were based on numerical solutions of analytic approximations to the $\alpha_1, \alpha_2, \alpha_3$ evolutions):

$$\mu \frac{d\alpha_i}{d\mu} = \frac{b_i}{2\pi} \alpha_i^2 + \sum_j \frac{b_{ij}}{8\pi^2} \alpha_j \alpha_i^2 \quad (10)$$

with SUSY coefficients b_i, b_{ij} if $\mu > M_{\text{SUSY}}$ (and assuming $M_{\text{SUSY}} > M_Z \geq m_t$).

The results are:

$$M_{\text{GUT}} = \left(1.6 \begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 4.7 \\ 1.4 \end{smallmatrix}\right) \times 10^{16} \text{ GeV} \quad (11a)$$

$$1/\alpha_{\text{GUT}} = 23.6^{+4.6}_{-0.5} . \quad (11b)$$

Note that increasing accuracy lowers the values of M_{GUT} .

Figure 2 shows the correlation between the possible input values for $\alpha_3(M_Z)$ and the results for M_{GUT} .

3c The value of M_{GUT} with $\alpha_1, \alpha_2, \alpha_3$ evolution at one loop and introducing the evolution of the gaugino masses (at one loop)

Keeping the one loop for the evolution of the couplings, we introduce a one-loop treatment of the threshold when SUSY [$SU(3) \times SU(2) \times U(1)$] breaks either into $SU(3) \times SU(2) \times U(1)$ or into $SU(3) \times U(1)_{\text{em}}$. First we consider the light threshold in the simplest way, using only one primordial mass, $m_{1/2}$, which parametrizes the gauginos, and allowing these masses to evolve [3].

The equations are:

$$\frac{1}{\alpha_3(m_{\tilde{g}})} = \frac{1}{\alpha_{\text{GUT}}(m_{1/2})} - \frac{3}{2\pi} \log\left(\frac{M_{\text{GUT}}}{m_{\tilde{g}}}\right) \quad (12)$$

$$\frac{1}{\alpha_2(m_{\tilde{W}})} = \frac{1}{\alpha_{\text{GUT}}(m_{1/2})} - \frac{19}{12\pi} \log\left(\frac{m_{1/2}}{m_{\tilde{W}}}\right) + \frac{1}{2\pi} \log\left(\frac{M_{\text{GUT}}}{m_{1/2}}\right) \quad (13)$$

$$\frac{1}{\alpha_{\text{GUT}}(m_{1/2})} = \frac{1}{(\alpha_{\text{GUT}})_0} + \frac{73}{40\pi} \log\left(\frac{m_{1/2}}{M_Z}\right) \quad (14)$$

$$[(\alpha_{\text{GUT}})_0]^{-1} = \frac{1}{20} \left[\frac{3}{\alpha_{\text{em}}^{\text{exp}}(M_Z)} + \frac{12}{\alpha_3^{\text{exp}}(M_Z)} \right] \quad (15)$$

$$(M_{\text{GUT}})_0 = M_Z \exp\left[\frac{\pi}{10} \left(\frac{1}{\alpha_{\text{em}}^{\text{exp}}(M_Z)} - \frac{8}{3\alpha_3^{\text{exp}}(M_Z)} \right) \right] \quad (16)$$

$$m_{\tilde{g}} = \frac{\alpha_3(m_{\tilde{g}})}{\alpha_{\text{GUT}}(m_{1/2})} m_{1/2} \quad (17)$$

$$m_{\tilde{W}} = \frac{\alpha_2(m_{\tilde{W}})}{\alpha_{\text{GUT}}(m_{1/2})} m_{1/2} \quad (18)$$

$$M_{\text{GUT}} = (M_{\text{GUT}})_0 \left(\frac{m_{1/2}}{m_Z} \right)^{-7/60}. \quad (19)$$

The results, with one loop for the evolution of $\alpha_1, \alpha_2, \alpha_3$, one loop for the evolution of the gaugino masses, are:

$$M_{\text{GUT}} = \left(1.10 \begin{smallmatrix} + 4.00 \\ - 1.08 \end{smallmatrix} \right) \times 10^{16} \text{ GeV} \quad (20a)$$

$$1/\alpha_{\text{GUT}} = 27.7 \begin{smallmatrix} + 14.7 \\ - 4.6 \end{smallmatrix} \quad (20b)$$

Note: the introduction of the evolution of the gaugino masses lowers M_{GUT} [see (9a)]. Figure 3 shows the correlation between the possible input values for $\alpha_3(M_Z)$ and the results for M_{GUT} .

3d The value of M_{GUT} with $\alpha_1, \alpha_2, \alpha_3$ evolution at one loop introducing the evolution of all masses and a detailed spectrum for the light threshold (at one loop)

This approach should be compared with the approach shown in 3c, where we have one loop for the evolution of the couplings ($\alpha_1, \alpha_2, \alpha_3$) and one loop for the evolution of the gaugino masses. Here we extend the evolution [4] to all masses (squarks, sleptons, Higgsino, ...) and consider a detailed light-threshold spectrum.

The equations for the couplings are:

$$\begin{aligned} \frac{1}{\alpha_{\text{GUT}}} = & \frac{3}{20} \left(\frac{1}{\alpha_{\text{em}}^{\text{exp}}(M_Z)} + \frac{4}{\alpha_3^{\text{exp}}(M_Z)} \right) + \\ & + \frac{1}{2\pi} \left[\frac{6}{5} \log\left(\frac{m_{\tilde{g}}}{M_Z}\right) + \frac{65}{48} \log\left(\frac{m_{\tilde{q}}}{M_Z}\right) + \frac{43}{240} \log\left(\frac{m_{\tilde{t}_L}}{M_Z}\right) + \right. \\ & + \frac{1}{6} \log\left(\frac{m_{\tilde{t}_R}}{M_Z}\right) + \frac{3}{20} \log\left(\frac{m_{\tilde{t}_L}}{M_Z}\right) + \frac{3}{20} \log\left(\frac{m_{\tilde{t}_R}}{M_Z}\right) + \\ & + \frac{1}{5} \log\left(\frac{m_{\tilde{h}}}{M_Z}\right) + \frac{1}{20} \log\left(\frac{m_H}{M_Z}\right) + \frac{1}{5} \log\left(\frac{m_{\tilde{W}}}{M_Z}\right) + \\ & \left. + \frac{83}{120} \log\left(\frac{m_t}{M_Z}\right) \right] \end{aligned} \quad (21)$$

$$\begin{aligned} \frac{1}{\alpha_1(\mu)} = & \frac{1}{\alpha_{\text{GUT}}} + \frac{1}{2\pi} \left[\frac{33}{5} \log\left(\frac{M_{\text{GUT}}}{\mu}\right)^* - \frac{49}{60} \log\left(\frac{m_{\tilde{q}}}{\mu}\right)^* + \right. \\ & - \frac{17}{30} \log\left(\frac{m_t}{\mu}\right)^* - \frac{1}{60} \log\left(\frac{m_{\tilde{t}_L}}{\mu}\right)^* - \frac{4}{15} \log\left(\frac{m_{\tilde{t}_R}}{\mu}\right)^* + \\ & - \frac{3}{10} \log\left(\frac{m_{\tilde{t}_L}}{\mu}\right)^* - \frac{3}{5} \log\left(\frac{m_{\tilde{t}_R}}{\mu}\right)^* - \frac{2}{5} \log\left(\frac{m_{\tilde{h}}}{\mu}\right)^* + \\ & \left. - \frac{1}{10} \log\left(\frac{m_H}{\mu}\right)^* \right] \end{aligned} \quad (22)$$

$$\begin{aligned} \frac{1}{\alpha_2(\mu)} = & \frac{1}{\alpha_{\text{GUT}}} + \frac{1}{2\pi} \left[\log\left(\frac{M_{\text{GUT}}}{\mu}\right)^* - \log\left(\frac{m_{\tilde{q}}}{\mu}\right)^* + \right. \\ & - \log\left(\frac{m_t}{\mu}\right)^* - \frac{1}{2} \log\left(\frac{m_{\tilde{t}_L}}{\mu}\right)^* - \frac{1}{2} \log\left(\frac{m_{\tilde{t}_R}}{\mu}\right)^* + \\ & \left. - \frac{2}{3} \log\left(\frac{m_{\tilde{h}}}{\mu}\right)^* - \frac{1}{6} \log\left(\frac{m_H}{\mu}\right)^* - \frac{4}{3} \log\left(\frac{m_{\tilde{w}}}{\mu}\right)^* \right] \end{aligned} \quad (23)$$

$$\begin{aligned} \frac{1}{\alpha_3(\mu)} = & \frac{1}{\alpha_{\text{GUT}}} + \frac{1}{2\pi} \left[-3 \log\left(\frac{M_{\text{GUT}}}{\mu}\right)^* - 2 \log\left(\frac{m_{\tilde{g}}}{\mu}\right)^* + \right. \\ & - \frac{2}{3} \log\left(\frac{m_t}{\mu}\right)^* - \frac{5}{3} \log\left(\frac{m_{\tilde{q}}}{\mu}\right)^* - \frac{1}{6} \log\left(\frac{m_{\tilde{t}_L}}{\mu}\right)^* + \\ & \left. - \frac{1}{6} \log\left(\frac{m_{\tilde{t}_R}}{\mu}\right)^* \right] , \end{aligned} \quad (24)$$

where:

$$\log(x)^* \equiv \Theta(x-1) \cdot \log(x) \quad \text{and} \quad \Theta(y) \equiv \begin{cases} 0 & \text{if } y < 0 \\ 1 & \text{if } y > 0 \end{cases} \quad (25)$$

For the masses the equations are:

$$\begin{aligned} \frac{M_{\text{GUT}}}{(M_{\text{GUT}})_0} = & \left(\frac{m_{\tilde{w}}}{M_Z}\right)^{1/15} \cdot \left(\frac{m_{\tilde{g}}}{M_Z}\right)^{-4/15} \cdot \left(\frac{m_{\tilde{h}}}{M_Z}\right)^{1/15} \cdot \left(\frac{m_H}{M_Z}\right)^{1/60} \cdot \\ & \cdot \left(\frac{m_{\tilde{q}}}{M_Z}\right)^{-5/48} \cdot \left(\frac{m_{\tilde{t}_L}}{M_Z}\right)^{1/240} \cdot \left(\frac{m_{\tilde{t}_R}}{M_Z}\right)^{1/20} \cdot \\ & \cdot \left(\frac{m_{\tilde{t}_R}}{M_Z}\right)^{1/20} \cdot \left(\frac{m_t}{M_Z}\right)^{1/120} \end{aligned} \quad (26)$$

$$\begin{aligned} \log\left(\frac{m_{1/2}}{M_Z}\right) = & F(X) + \frac{28}{19} \log\left(\frac{\alpha_3(m_{\tilde{g}})}{\alpha_{\text{GUT}}}\right) - \frac{32}{19} \log\left(\frac{\alpha_2(m_{\tilde{w}})}{\alpha_{\text{GUT}}}\right) + \\ & + \frac{3}{19} \left\{ \frac{5}{2} \log\left(\frac{m_{\tilde{q}}}{M_Z}\right) - \frac{19}{6} \log\left(\frac{m_{\tilde{t}_L}}{M_Z}\right) + \frac{5}{3} \log\left(\frac{m_{\tilde{t}_R}}{M_Z}\right) + \right. \\ & \left. - 3 \log\left(\frac{m_{\tilde{t}_L}}{M_Z}\right) + 2 \log\left(\frac{m_{\tilde{t}_R}}{M_Z}\right) - 3 \log\left(\frac{m_t}{M_Z}\right) + \right. \end{aligned}$$

$$\begin{aligned}
& + 6 \log\left(\frac{\alpha_2(m_H)}{\alpha_{\text{GUT}}}\right) + \frac{2}{11} \log\left(\frac{\alpha_1(m_H)}{\alpha_{\text{GUT}}}\right) - 4 \log\left(\frac{m_4}{m_{1/2}}\right) + \\
& + 6r^2 - \frac{1}{2} \log\left[\frac{3}{2} \left[1 - \left(\frac{\alpha_2(m_H)}{\alpha_{\text{GUT}}}\right)^2\right] + \right. \\
& \left. \frac{1}{22} \left[1 - \left(\frac{\alpha_1(m_H)}{\alpha_{\text{GUT}}}\right)^2\right] + \left(\frac{m_0}{m_{1/2}}\right)^2 + \right. \\
& \left. + \left(\frac{\alpha_{\text{GUT}}}{\alpha_2(m_H)}\right)^3 \cdot \left(\frac{\alpha_{\text{GUT}}}{\alpha_1(m_H)}\right)^{1/11} \cdot \left(\frac{m_4}{m_{1/2}}\right)^2\right] \} , \quad (27)
\end{aligned}$$

where the function $F(X)$ is defined as:

$$\begin{aligned}
F(X) = \frac{3}{19} \left\{ \frac{20\pi}{\alpha_{\text{em}}^{\text{exp}}(M_Z)} (0.2029 + \right. \\
\left. + \frac{7\alpha_{\text{em}}^{\text{exp}}(M_Z)}{15\alpha_3^{\text{exp}}(M_Z)} - \sin^2\theta^{\text{exp}}(M_Z)) \right\} \quad (28)
\end{aligned}$$

$$\log\left(\frac{(M_{\text{GUT}})_0}{M_Z}\right) = \frac{\pi}{10} \left(\frac{1}{\alpha_{\text{em}}^{\text{exp}}(M_Z)} - \frac{8}{3\alpha_3^{\text{exp}}(M_Z)} \right) \quad (29)$$

[eq. (29) is the same as eq. (16)];

$$m_{\tilde{g}} = \frac{\alpha_3(m_{\tilde{g}})}{\alpha_{\text{GUT}}} m_{1/2} \quad (30)$$

$$m_{\tilde{W}} = \frac{\alpha_2(m_{\tilde{W}})}{\alpha_{\text{GUT}}} m_{1/2} \quad (31)$$

$$\left(\frac{m_{\tilde{q}}}{m_{1/2}}\right)^2 = -\frac{8}{9} \left[1 - \left(\frac{\alpha_3(m_{\tilde{q}})}{\alpha_{\text{GUT}}}\right)^2\right] + \left(\frac{m_0}{m_{1/2}}\right)^2 \quad (32)$$

$$\begin{aligned}
\left(\frac{m_{\tilde{t}_L}}{m_{1/2}}\right)^2 = & -\frac{8}{9} \left[1 - \left(\frac{\alpha_3(m_{\tilde{q}})}{\alpha_{\text{GUT}}}\right)^2\right] + \\
& + \frac{w1}{6} \left[1 - \left(\frac{\alpha_3(m_{\tilde{q}})}{\alpha_{\text{GUT}}}\right)^2\right] + \left(\frac{m_0}{m_{1/2}}\right)^2 \quad (33)
\end{aligned}$$

$$\begin{aligned}
\left(\frac{m_{\tilde{t}_R}}{m_{1/2}}\right)^2 = & -\frac{8}{9} \left[1 - \left(\frac{\alpha_3(m_{\tilde{q}})}{\alpha_{\text{GUT}}}\right)^2\right] + \\
& + \frac{w1}{3} \left[1 - \left(\frac{\alpha_3(m_{\tilde{q}})}{\alpha_{\text{GUT}}}\right)^2\right] + \left(\frac{m_0}{m_{1/2}}\right)^2 \quad (34)
\end{aligned}$$

$$\begin{aligned} \left(\frac{m_{\tilde{I}_L}}{m_{1/2}}\right)^2 &= \frac{3}{2} \left[1 - \left(\frac{\alpha_2(m_{\tilde{I}_L})}{\alpha_{\text{GUT}}}\right)^2\right] + \\ &+ \frac{1}{22} \left[1 - \left(\frac{\alpha_1(m_{\tilde{I}_L})}{\alpha_{\text{GUT}}}\right)^2\right] + \left(\frac{m_0}{m_{1/2}}\right)^2 \end{aligned} \quad (35)$$

$$\left(\frac{m_{\tilde{I}_R}}{m_{1/2}}\right)^2 = \frac{2}{11} \left[1 - \left(\frac{\alpha_1(m_{\tilde{I}_R})}{\alpha_{\text{GUT}}}\right)^2\right] + \left(\frac{m_0}{m_{1/2}}\right)^2 \quad (36)$$

$$m_{\tilde{H}} = m_4 \left[\frac{\alpha_{\text{GUT}}}{\alpha_2(m_{\tilde{H}})}\right]^{3/2} \cdot \left[\frac{\alpha_{\text{GUT}}}{\alpha_1(m_{\tilde{H}})}\right]^{1/22} \cdot \exp\left(-\frac{3r^2}{2}\right) \quad (37)$$

where

$$r^2 = \frac{1}{4\pi} \int_0^T \alpha_t(\mu') d\log\left(\frac{\mu'^2}{\mu^2}\right) \quad (38)$$

with

$$\alpha_t = \frac{h_t^2}{4\pi} = \text{top-Yukawa coupling} \quad (39)$$

and

$$T = \log\left(\frac{M_{\text{GUT}}^2}{\mu^2}\right), \quad (40)$$

$$m_H = \sqrt{m_{\tilde{H}}^2 + m_{\tilde{I}_L}^2} \quad (\text{at scale } m_H) \quad (41)$$

The results, with one loop for the evolution of α_1 , α_2 , α_3 and one loop for the evolution of all masses with a detailed light threshold spectrum at one loop, are:

$$M_{\text{GUT}} = \left(1.3 \begin{smallmatrix} + 4.7 \\ - 1.1 \end{smallmatrix}\right) \times 10^{16} \text{ GeV} \quad (42a)$$

$$1/\alpha_{\text{GUT}} = 25.7 \begin{smallmatrix} + 3.9 \\ - 2.7 \end{smallmatrix} \quad (42b)$$

The result (42a) should be compared with (20a): a detailed light threshold spectrum slightly pushes M_{GUT} to higher values.

Figure 4 shows the correlation between the possible input values for $\alpha_3(M_Z)$ and the results for M_{GUT} .

3e The value of M_{GUT} with $\alpha_1, \alpha_2, \alpha_3$ evolution at two loops introducing the evolution of the gaugino masses (at two loops) with a detailed spectrum of the light threshold at one loop

In this approach we have two loops for the evolution of the couplings ($\alpha_1, \alpha_2, \alpha_3$) and of the gaugino masses [5].

The equations are:

$$\begin{aligned} \frac{1}{\alpha_1(M_Z)} = \frac{3}{5} \alpha_{\text{em}}^{-1}(1-\sin^2\theta) = \frac{1}{\alpha_{\text{GUT}}} + \frac{1}{2\pi} [& \frac{33}{5} \log\left(\frac{M_{\text{GUT}}}{M_Z}\right)^* + \\ & - \frac{49}{60} \log\left(\frac{m_{\tilde{q}}}{M_Z}\right)^* - \frac{17}{30} \log\left(\frac{m_t}{M_Z}\right)^* - \frac{1}{60} \log\left(\frac{m_{\tilde{t}_L}}{M_Z}\right)^* + \\ & - \frac{4}{15} \log\left(\frac{m_{\tilde{t}_R}}{M_Z}\right)^* - \frac{3}{10} \log\left(\frac{m_{\tilde{t}_L}}{M_Z}\right)^* - \frac{3}{5} \log\left(\frac{m_{\tilde{t}_R}}{M_Z}\right)^* + \\ & - \frac{2}{5} \log\left(\frac{m_{\tilde{H}}}{M_Z}\right)^* - \frac{1}{10} \log\left(\frac{m_H}{M_Z}\right)^*] + \frac{C_1}{4\pi} \end{aligned} \quad (43)$$

$$\begin{aligned} \frac{1}{\alpha_2(M_Z)} = \frac{\sin^2\theta}{\alpha_{\text{em}}} = \frac{1}{\alpha_{\text{GUT}}} + \frac{1}{2\pi} [& \log\left(\frac{M_{\text{GUT}}}{M_Z}\right)^* - \log\left(\frac{m_{\tilde{q}}}{M_Z}\right)^* + \\ & - \log\left(\frac{m_t}{M_Z}\right)^* - \frac{1}{2} \log\left(\frac{m_{\tilde{t}_L}}{M_Z}\right)^* - \frac{1}{2} \log\left(\frac{m_{\tilde{t}_R}}{M_Z}\right)^* + \\ & - \frac{2}{3} \log\left(\frac{m_{\tilde{H}}}{M_Z}\right)^* - \frac{1}{6} \log\left(\frac{m_H}{M_Z}\right)^* - \frac{4}{3} \log\left(\frac{m_{\tilde{W}}}{M_Z}\right)^*] + \frac{C_2}{4\pi} \end{aligned} \quad (44)$$

$$\begin{aligned} \frac{1}{\alpha_3(M_Z)} = \frac{1}{\alpha_{\text{GUT}}} + \frac{1}{2\pi} [& -3 \log\left(\frac{M_{\text{GUT}}}{M_Z}\right)^* - 2 \log\left(\frac{m_{\tilde{g}}}{M_Z}\right)^* + \\ & - \frac{2}{3} \log\left(\frac{m_t}{M_Z}\right)^* - \frac{5}{3} \log\left(\frac{m_{\tilde{q}}}{M_Z}\right)^* - \frac{1}{6} \log\left(\frac{m_{\tilde{t}_L}}{M_Z}\right)^* + \\ & - \frac{1}{6} \log\left(\frac{m_{\tilde{t}_R}}{M_Z}\right)^*] + \frac{C_3}{4\pi} \end{aligned} \quad (45)$$

The C_i stand for the following combinations of two-loop matrix coefficients:

$$C_i = \sum_j \frac{b_{ij}^s}{b_j^s} \log\left(\frac{\alpha_j^{-1}(M_Z)}{\alpha_{\text{GUT}}^{-1}}\right) + \left(\sum_j \frac{b_{ij}^{s'}}{b_j^{s'}} - \sum_j \frac{b_{ij}^s}{b_j^s} \right) \log\left(\frac{\alpha_j^{-1}(M_Z)}{\alpha_j^{-1}(m_{1/2})}\right) \quad (46)$$

for $i = 1, 2, 3$.

The results, with two loops for the evolution of $\alpha_1, \alpha_2, \alpha_3$ and for the evolution of the gaugino masses and light threshold at one loop, are:

$$M_{\text{GUT}} = \left(1.0 \begin{smallmatrix} + 2.8 \\ - 0.8 \end{smallmatrix}\right) \times 10^{16} \text{ GeV} \quad (47a)$$

$$1/\alpha_{\text{GUT}} = 25.3 \begin{smallmatrix} + 3.9 \\ - 2.8 \end{smallmatrix} \quad (47b)$$

The result (47a) should be compared with (42a). Again, increasing accuracy favours lower M_{GUT} values.

Figure 5 shows the correlation between the possible input values for $\alpha_3(M_Z)$ and the results for M_{GUT} .

3f The value of M_{GUT} introducing the simplest structure in ΔT_H (at one loop)

We now study the approach of 3d, where we allow ΔT_H to have a structure. The simplest one is in terms of three masses: M_V, M_H, M_Σ . Two of them need to be very high and quasi-degenerate: M_V and M_H , owing to the limits on τ_p . The one which could be low is M_Σ . Let us consider for ΔT_H the ratio M_V/M_Σ . We let this ratio be in the range 10^0 – 10^6 .

The equations are the same as in sect. 3d, but with the following changes: Eqs. (21), (23), (24) should be modified respectively into (48), (49), (50):

$$\begin{aligned} \frac{1}{\alpha_{\text{GUT}}} = & \frac{3}{20} \left(\frac{1}{\alpha_{\text{em}}^{\text{exp}}(M_Z)} + \frac{4}{\alpha_3^{\text{exp}}(M_Z)} \right) + \\ & + \frac{1}{2\pi} \left[\frac{6}{5} \log\left(\frac{m_{\tilde{g}}}{M_Z}\right) + \frac{65}{48} \log\left(\frac{m_{\tilde{q}}}{M_Z}\right) + \frac{43}{240} \log\left(\frac{m_{\tilde{t}_L}}{M_Z}\right) + \right. \\ & + \frac{1}{6} \log\left(\frac{m_{\tilde{t}_R}}{M_Z}\right) + \frac{3}{20} \log\left(\frac{m_{\tilde{t}_L}}{M_Z}\right) + \frac{3}{20} \log\left(\frac{m_{\tilde{t}_R}}{M_Z}\right) + \\ & + \frac{1}{5} \log\left(\frac{m_{\tilde{H}}}{M_Z}\right) + \frac{1}{20} \log\left(\frac{m_H}{M_Z}\right) + \frac{1}{5} \log\left(\frac{m_{\tilde{W}}}{M_Z}\right) + \\ & \left. + \frac{83}{120} \log\left(\frac{m_t}{M_Z}\right) - \frac{21}{10} \log\left(\frac{M_V}{M_\Sigma}\right) \right] \quad (48) \end{aligned}$$

$$\begin{aligned} \frac{1}{\alpha_2(\mu)} = \frac{1}{\alpha_{GUT}} + \frac{1}{2\pi} \Big[\log\left(\frac{M_{GUT}}{\mu}\right)^* - \log\left(\frac{m_{\tilde{q}}}{\mu}\right)^* + \\ - \log\left(\frac{m_t}{\mu}\right)^* - \frac{1}{2} \log\left(\frac{m_{\tilde{t}_L}}{\mu}\right)^* + \\ - \frac{1}{2} \log\left(\frac{m_{\tilde{t}_R}}{\mu}\right)^* - \frac{2}{3} \log\left(\frac{m_{\tilde{b}}}{\mu}\right)^* + \\ - \frac{1}{6} \log\left(\frac{m_H}{\mu}\right)^* - \frac{4}{3} \log\left(\frac{m_{\tilde{W}}}{\mu}\right)^* + 2 \log\left(\frac{M_V}{M_\Sigma}\right) \Big] \quad (49) \end{aligned}$$

$$\begin{aligned} \frac{1}{\alpha_3(\mu)} = \frac{1}{\alpha_{GUT}} + \frac{1}{2\pi} \Big[-3 \log\left(\frac{M_{GUT}}{\mu}\right)^* - 2 \log\left(\frac{m_{\tilde{g}}}{\mu}\right)^* + \\ - \frac{2}{3} \log\left(\frac{m_t}{\mu}\right)^* - \frac{5}{3} \log\left(\frac{m_{\tilde{q}}}{\mu}\right)^* + \\ - \frac{1}{6} \log\left(\frac{m_{\tilde{t}_L}}{\mu}\right)^* - \frac{1}{6} \log\left(\frac{m_{\tilde{t}_R}}{\mu}\right)^* + 3 \log\left(\frac{M_V}{M_\Sigma}\right) \Big] \quad (50) \end{aligned}$$

The right-hand side of eq. (26) should be multiplied by $(M_V/M_\Sigma)^{3/10}$.
To the right-hand side of eq. (27) should be added a term $\frac{6}{19} \log\left(\frac{M_V}{M_\Sigma}\right)$.

If $(M_V/M_\Sigma) = 10^2$ the results, using the $\alpha_1, \alpha_2, \alpha_3$ evolutions at two loops, ΔT_L at one loop; ΔT_H at two loops, are:

$$M_{GUT} = \left(4.8 \begin{smallmatrix} + 10.9 \\ - 3.9 \end{smallmatrix}\right) \times 10^{16} \text{ GeV} \quad (51a)$$

$$1/\alpha_{GUT} = 24.8 \begin{smallmatrix} + 4.0 \\ - 2.8 \end{smallmatrix} \quad (51b)$$

This is the highest possible value for M_{GUT} so far.

Figure 6 shows the correlation between the possible input values for $\alpha_3(M_Z)$ and the results for M_{GUT} .

Note that:

1) The ratio $M_{GUT}^{MAX}/M_{GUT}^{MIN}$ changes by a factor of almost 2 when $(m_4/m_{1/2})^2$ goes from 10^{-2} to 10^2 . This ratio remains practically unchanged when M_V/M_Σ goes from 1 to 10^6 . The M_{GUT} value obtained when

$$\frac{M_V}{M_\Sigma} = 10^6 \text{ and } \left(\frac{m_4}{m_{1/2}}\right)^2 = 10^2$$

is:

$$M_{\text{GUT}} = 8.87 \times 10^{17} \text{ GeV} .$$

2) The ratio $M_{\text{GUT}}^{\text{MAX}}/M_{\text{GUT}}^{\text{MIN}}$ changes by 20% when $(m_0/m_{1/2})^2$ goes from 10^{-2} to 10^2 . This ratio remains practically unchanged when M_V/M_Σ goes from 1 to 10^6 .

In order to see how high M_{GUT} can be, we have used as inputs the values:

$$\left(\frac{m_4}{m_{1/2}}\right)^2 = \left(\frac{m_0}{m_{1/2}}\right)^2 = 10^2 , \quad \frac{M_V}{M_\Sigma} = 10^6 ,$$

with $\alpha_3(M_Z) = 0.134$ (WA+2 σ). The result is:

$$M_{\text{GUT}} = 3.14 \times 10^{18} \text{ GeV} ,$$

which represents the highest possible value for the energy level where the “low-energy” measurements allow $\alpha_1, \alpha_2, \alpha_3$ to merge.

Table 1 summarizes the results of our approach 3f.

3g The value of M_{GUT} introducing the smooth convergence towards M_{GUT} and the unified evolution above M_{GUT} (one loop)

We now follow a different approach [2]. The evolution of the couplings $(\alpha_1, \alpha_2, \alpha_3)$ is worked out from above, starting from the high side and descending down to 10^2 GeV. Going from the high-energy side down, a supergroup, for example SUSY-SU(5), will break into SUSY $[SU(3) \times SU(2) \times U(1)]$ crossing a heavy threshold at M_X . In this approach we ignore the light threshold problems and concentrate on the heavy threshold, which is treated in the simplest way: assuming only one heavy mass and taking as unique input the value of α_3 at the Z^0 mass: $\alpha_3(M_Z)$. We then allow the program to choose the value for M_{GUT} that best fits the other two experimentally measured quantities: $\sin^2\theta(M_Z)$ and $\alpha_{\text{em}}(M_Z)$.

The equations are:

$$\frac{1}{\alpha_1(\mu_1^2)} = \frac{1}{\alpha_1(\mu_0^2)} - \frac{1}{24\pi} [39.6 \log(\mu_1^2/\mu_0^2) + 57.6 F^{(1)}(\mu_1^2, \mu_0^2, M_X^2)] \quad (52)$$

$$\frac{1}{\alpha_2(\mu_1^2)} = \frac{1}{\alpha_2(\mu_0^2)} - \frac{1}{24\pi} [6 \log(\mu_1^2/\mu_0^2) + 24 F^{(1)}(\mu_1^2, \mu_0^2, M_X^2)] \quad (53)$$

$$\frac{1}{\alpha_3(\mu_1^2)} - \frac{1}{\alpha_3(\mu_0^2)} = \frac{18}{24\pi} \log(\mu_1^2/\mu_0^2), \quad (54)$$

where μ_1 is a scale above M_X where unification is assumed to have taken place, μ_0 is a scale below the M_X threshold, and:

$$F^{(1)}(x, y, z) = \log \left(\frac{x + 4z}{y + 4z} \right) \quad (55)$$

The result for M_{GUT} , introducing the simplest possible heavy threshold to allow the smooth convergence of $\alpha_1, \alpha_2, \alpha_3$ towards M_{GUT} and the unified evolution above M_{GUT} , at one loop, is:

$$M_{GUT} = \left(3.6 \begin{smallmatrix} +5.0 \\ -2.2 \end{smallmatrix} \right) \times 10^{16} \text{ GeV} \quad (56)$$

Figure 7 shows the correlation between the possible input values for $\alpha_3(M_Z)$ and the results for M_{GUT} .

3h The value of M_{GUT} introducing the smooth convergence towards M_{GUT} and the unified evolution above M_{GUT} (two loops)

Here we repeat the same approach as in 3g, improving the theoretical approximation via the two loop calculation.

The equations are:

$$\begin{aligned} \frac{1}{\alpha_1(\mu_1^2)} = \frac{1}{\alpha_1(\mu_0^2)} - \frac{1}{24\pi} [39.6 \log(\mu_1^2/\mu_0^2) - \\ - 57.6 F^{(1)}(\mu_1^2, \mu_0^2, M_X^2)] + T_1 + F_1^{(2)}(\mu_1^2, \mu_0^2, M_X^2) \end{aligned} \quad (57)$$

$$\begin{aligned} \frac{1}{\alpha_2(\mu_1^2)} = \frac{1}{\alpha_2(\mu_0^2)} - \frac{1}{24\pi} [6 \log(\mu_1^2/\mu_0^2) - \\ - 24 F^{(1)}(\mu_1^2, \mu_0^2, M_X^2)] + T_2 + F_2^{(2)}(\mu_1^2, \mu_0^2, M_X^2) \end{aligned} \quad (58)$$

$$\frac{1}{\alpha_3(\mu_1^2)} = \frac{1}{\alpha_3(\mu_0^2)} - \frac{18}{24\pi} \log(\mu_1^2/\mu_0^2) + T_3, \quad (59)$$

where

$$T_i = \sum_{j=1}^3 \frac{b_{ij}}{4\pi b_j} \log \frac{\alpha_j(\mu_1^2)}{\alpha_j(\mu_0^2)} \quad (60)$$

and

$$F_j^{(2)}(\mu_1^2, \mu_0^2, M_X^2) = (T_3 - T_j) \frac{F^{(1)}(\mu_1^2, M_Z^2 \cdot \mu_1^2/\mu_0^2, M_X^2)}{F^{(1)}(\mu_1^2, M_Z^2, M_X^2)}, \quad (61)$$

The results, with ΔT_H at two loops, are:

$$M_{GUT} = \left(3.0 \begin{smallmatrix} +4.6 \\ -1.8 \end{smallmatrix} \right) \times 10^{16} \text{ GeV} \quad (62a)$$

$$1/\alpha_{GUT} = 24 \pm 1 \quad (62b)$$

As for the approach 3b, increasing accuracy lowers the value for M_{GUT} .

Figure 8 shows the correlation between the possible input values for $\alpha_3(M_Z)$ and the results for M_{GUT} .

4. DISCUSSION AND CONCLUSIONS

All M_{GUT} values calculated using different approaches and theoretical levels of accuracy are reported in fig. 9. We report in fig. 10 the correlation between these values of M_{GUT} and the associated α_{GUT} . Finally in fig. 11 we show how M_{GUT} depends on $\alpha_3(M_Z)$ and on the ratio M_V/M_Σ . The lowest value of M_{GUT} corresponds to a value of $\alpha_3(M_Z)$ that is equal to the world average minus 2σ and to a ratio $M_V/M_\Sigma = 10^0$. This value is $M_{GUT} = 2.5 \times 10^{15} \text{ GeV}$; it is lower than the M_X allowed by the experimental lower limit of τ_p . If we assume a quasi-degenerate structure for the heavy threshold, the ratios of the primordial masses equal to 1 and the central values for the experimentally measured quantities [$\alpha_3(M_Z)$, $\sin^2\theta(M_Z)$, $\alpha_{em}(M_Z)$], the value of M_{GUT} is $1.3 \times 10^{16} \text{ GeV}$. However, this estimate depends on the ratio of the primordial masses and is dominated by the ratio M_V/M_Σ . If this ratio is 10^6 and $\alpha_3(M_Z)$ has the world average value $+2\sigma$, the result is $M_{GUT} = 2.1 \times 10^{18} \text{ GeV}$, a value below M_{Planck} by almost one order of magnitude. However, with the same inputs, if we take 10^2 for the ratios of the primordial masses $(m_4/m_{1/2})^2$ and $(m_0/m_{1/2})^2$, the result for M_{GUT} is $3.1 \times 10^{18} \text{ GeV}$. This is the extreme limit where “low-energy” data can push M_{GUT} . The existence of a gap between M_{GUT} and M_{Planck} depends on the knowledge of these key

quantities.

Thanks to our studies on M_{GUT} as a function of experimental and theoretical inputs and uncertainties, the trends shown in table 2 can be worked out. The gap between M_{GUT} and M_{Planck} appears to be on safe grounds. In fact even with extreme input conditions M_{GUT} remains safely below M_{Planck} . The consequences of the existence of this gap in the range of accessible energies should be extensively studied.

Table 1

The values of M_{GUT} for the approach described in section 3f when $\alpha_3(M_Z)$ ranges from the world average (WA) value to $\pm 1\sigma$, $\pm 2\sigma$, and the ratio M_V/M_Σ goes from 10^0 to 10^6 . The quoted values are for $(m_4/m_{1/2})^2 = (m_0/m_{1/2})^2 = 1$.

$\alpha_3(M_Z)$	$\log_{10}(M_V/M_\Sigma)$	$\log_{10}(M_{\text{GUT}})$
0.118 (WA)	0	16.12
	2	16.68
	4	17.24
	6	17.80
0.126 (WA+1 σ)	0	16.39
	2	16.96
	4	17.52
	6	18.09
0.134 (WA+2 σ)	0	16.61
	2	17.19
	4	17.76
	6	18.33
0.110 (WA-1 σ)	0	15.79
	2	16.35
	4	16.90
	6	17.46
0.102 (WA-2 σ)	0	15.41
	2	15.96
	4	16.51
	6	17.06

Table 2

The trends are as follows: increasing accuracy (from one loop to two loops) lowers M_{GUT} . The introduction of ΔT_L has no effect on M_{GUT} . The introduction of a structure in the heavy threshold pushes M_{GUT} upwards and the introduction of EGM lowers M_{GUT} . Higher values of $\alpha_3(M_Z)$, $\sin^2\theta(M_Z)$ and $\alpha_{\text{em}}(M_Z)$, all push M_{GUT} up.

$M_{\text{GUT}} \uparrow$	TRENDS	$M_{\text{GUT}} \downarrow$	Reference
	From 1 loop to 2 loops ΔT_L (one mass)		§§ 3a/3b
	From 1 loop to 2 loops ΔT_H (one mass)		§§ 3g/3h
	Introducing EGM (one loop) with $\alpha_1 \alpha_2 \alpha_3$ (one loop)		§§ 3a/3c
	Introducing EGM (two loops) with $\alpha_1 \alpha_2 \alpha_3$ (two loops)		§§ 3b/3c
	Introducing ΔT_L		§§ 3c/3d
	Introducing ΔT_H (one mass) (2 loops)		§§ 3b/3h
	Introducing $\Delta T_H (M_V/M_Z)$ in (EGM + ΔT_L) (one loop)		[4]
	$\alpha_3(M_Z) \uparrow$		[4]
	$\sin^2\theta(M_Z) \uparrow$		[4]
	$\alpha_{\text{em}}(M_Z) \uparrow$		§§ 3d/3f

References

- [1] F. Anselmo, L. Cifarelli, A. Peterman and A. Zichichi, *Il Nuovo Cimento* 104A (1991) 1817. We refer the reader to this work for notations, formulae and other details not discussed in the present paper.
- [2] F. Anselmo, L. Cifarelli, A. Peterman and A. Zichichi, "The Convergence of the Gauge Couplings at EGUT and Above: Consequences for $\alpha_3(M_Z)$ and SUSY Breaking", preprint CERN/LAA/MSL/91-026, December 1991, submitted to *Il Nuovo Cimento*.
- [3] F. Anselmo, L. Cifarelli, A. Peterman and A. Zichichi, *Il Nuovo Cimento* 105A (1992) 581.
- [4] F. Anselmo, L. Cifarelli, A. Peterman and A. Zichichi, "The Simultaneous Evolution of Masses and Couplings: Consequences on SUSY Spectra and Thresholds", preprint CERN/LAA/MSL/92-008, 2 April 1992 and CERN-PPE/92-103, 26 June 1992.
- [5] F. Anselmo, L. Cifarelli, A. Peterman and A. Zichichi, "Analytic Study of the Supersymmetry-breaking Scale at Two Loops", preprint CERN-TH.6543/92, June 1992.
- [6] J. Ellis, S. Kelley and D.V. Nanopoulos, *Nucl. Phys.* B373 (1992) 55.
- [7] R. Barbieri and L.J. Hall, *Phys. Rev. Lett.* 68 (1992) 752.
- [8] D.G. Unger and Y.P. Yao, University of Michigan, Report No. UMHE 81-30 (unpublished). See also D.R.T. Jones, *Phys. Rev.* D25 (1982) 581.
- [9] M.B. Einhorn and D.R.T. Jones, *Nucl. Phys.* B196 (1982) 475.

Figure captions

- Figure 1 Correlation between M_{GUT} and $\alpha_3(M_Z)$ for the approach 3a: $\alpha_1, \alpha_2, \alpha_3$ evolution at one loop; only one M_{SUSY} (step function).
- Figure 2 Correlation between M_{GUT} and $\alpha_3(M_Z)$ for the approach 3b: $\alpha_1, \alpha_2, \alpha_3$ evolution at two loops; only one M_{SUSY} (step function).
- Figure 3 Correlation between M_{GUT} and $\alpha_3(M_Z)$ for the approach 3c: $\alpha_1, \alpha_2, \alpha_3$ evolution at one loop; $(m_{1/2})$ EGM (one loop) (step function). No ΔT_L .
- Figure 4 Correlation between M_{GUT} and $\alpha_3(M_Z)$ for the approach 3d: $\alpha_1, \alpha_2, \alpha_3$ evolution at one loop; $(m_{1/2})$ EGM (one loop) (step function) plus ΔT_L (one loop). The ratios $M_V/M_\Sigma = M_V/M_H = 1$. The leftmost curve corresponds to $(m_0/m_{1/2})^2 = (m_4/m_{1/2})^2 = 10^{-2}$, the central one to $(m_0/m_{1/2})^2 = (m_4/m_{1/2})^2 = 1$, and the rightmost one to $(m_0/m_{1/2})^2 = (m_4/m_{1/2})^2 = 10^2$.
- Figure 5 Correlation between M_{GUT} and $\alpha_3(M_Z)$ for the approach 3e: $\alpha_1, \alpha_2, \alpha_3$ evolution at one loop; $(m_{1/2})$ EGM (two loops) (step function) plus ΔT_L (one loop).
- Figure 6 Correlation between M_{GUT} and $\alpha_3(M_Z)$ for the approach 3f: $\alpha_1, \alpha_2, \alpha_3$ evolution at one loop; $(m_{1/2})$ EGM (one loop) (step function) plus ΔT_L (one loop). The ratios $M_V/M_\Sigma = 10^2$, $M_V/M_H = 1$. The leftmost curve corresponds to $(m_0/m_{1/2})^2 = (m_4/m_{1/2})^2 = 10^{-2}$. The central to $(m_0/m_{1/2})^2 = (m_4/m_{1/2})^2 = 1$. The rightmost curve corresponds to $(m_0/m_{1/2})^2 = (m_4/m_{1/2})^2 = 10^2$.
- Figure 7 Correlation between M_{GUT} and $\alpha_3(M_Z)$ for the approach 3g: $\alpha_1, \alpha_2, \alpha_3$ evolution at one loop; ΔT_H (one M_X , threshold smooth) (one loop).
- Figure 8 Correlation between M_{GUT} and $\alpha_3(M_Z)$ for the approach 3h: $\alpha_1, \alpha_2, \alpha_3$ evolution at two loops; ΔT_H (one M_X , threshold smooth) (two loops).
- Figure 9 The values of M_{GUT} from different approaches, as specified.

The uncertainties are computed taking $\pm 2\sigma$ around the world average for $\alpha_3(M_Z)$. For the two approaches: ($\alpha_1, \alpha_2, \alpha_3$ one loop; EGM& ΔT_L one loop) and (ΔT_H two masses & two loops; EGM& ΔT_L one loop) the central values are computed with $m_{\text{top}} = 125$ GeV; error bands also include uncertainties on m_{top} ($95 \text{ GeV} < m_{\text{top}} < 180 \text{ GeV}$). The primordial mass ratios are $m_0/m_{1/2} = m_4/m_{1/2} = 1$.

Figure 10 The α_{GUT} values corresponding to the different values of M_{GUT} resulting from the different approaches described in the text. The uncertainties are computed taking $\pm 2\sigma$ around the world average for $\alpha_3(M_Z)$. For the two approaches: ($\alpha_1, \alpha_2, \alpha_3$ one loop; EGM& ΔT_L one loop) and (ΔT_H two masses & two loops; EGM& ΔT_L one loop), the central values are computed with $m_{\text{top}} = 125$ GeV; error bands also include uncertainties on m_{top} ($95 \text{ GeV} < m_{\text{top}} < 180 \text{ GeV}$). The primordial mass ratios are $m_0/m_{1/2} = m_4/m_{1/2} = 1$.

Figure 11 These data show the dependence of M_{GUT} from $\alpha_3(M_Z)$ and the ratio of the two crucial heavy threshold masses M_V/M_Σ , as described in the text. Note that the extreme value for M_{GUT} is above 10^{18} GeV.

Approach 3a

Figure 1

Approach 3b

Figure 2

Approach 3c

Figure 3

Approach 3d

Figure 4

Approach 3e

Figure 5

Approach 3f

Figure 6

Approach 3g

Figure 7

Approach 3h

Figure 8

Figure 9

Figure 10

- World Average: WA
- WA $\pm 1\sigma$
- WA $\pm 2\sigma$
- WA $\pm 3\sigma$

$\square$ Extreme values
 $\left\{ \begin{array}{l} \text{WA} - 2\sigma \quad \& \quad \frac{M_V}{M_\Sigma} = 10^0 \quad \& \quad \left(\frac{m_0}{m_{1/2}}\right)^2 = \left(\frac{m_4}{m_{1/2}}\right)^2 = 1 \\ \text{WA} + 2\sigma \quad \& \quad \frac{M_V}{M_\Sigma} = 10^6 \quad \& \quad \left(\frac{m_0}{m_{1/2}}\right)^2 = \left(\frac{m_4}{m_{1/2}}\right)^2 = 10^2 \end{array} \right.$

Figure 11