Archives

EUROPEAN ORGANIZATION FOR NUCLEAR RESEARCH

CERN LIBRARIES, GENEVA

CM-P00057475

CERN TH.6655/92

UITP 02/92

September 1992

PARTICLE PRODUCTION NEAR THRESHOLD

T.E.O. Ericson

CERN, CH 1211 Geneva 23, Switzerland

and

Institute for Theoretical Physics, S-75238 Uppsala, Sweden

To be published in Physica Scripta

*Conference Summary of Second International Symposium on Particle Production Near
Threshold, Uppsala, Sweden, August 26-29, 1992.*

6653

1 Introduction

This has been a very interesting conference with many new results and suggestions for the future. I have enjoyed listening to the talks and the discussions. Now, when I must summarize all this, I realize that it is not possible to do justice to it all, since there have been over 40 talks. I will therefore make no detailed comments, but only take up some general points. In particular I will not comment on the talks in the last session, since they anyway are fresh in your mind. Let me follow approximately the order of topics in the programme.

2 Facilities, detectors and targets

The conference started out discussing facilities, special detectors and targets. These are technical matters on which I can make only general comments, being a theoretical physicist. However, I found it highly appropriate to spend some time on these. With the cooler rings that have just started to operate here in Uppsala and at Indiana, and with those which will come into operation in the next few years, we are dealing with qualitatively different facilities as compared to normal accelerators. True, we have already had such storage rings available at LEAR for antiprotons, and that has been most instructive. Still, the exceptional beam quality, the exceptional energy resolution, and the luminosities which are possible both for polarized and unpolarized beams are so different in character that we must get used to exploiting them maximally. This is of course always the goal, but here we are dealing with instruments of such a power that we do not always perceive the possibilities. It is therefore an important task to develop dedicated, powerful, multipurpose detectors such as EDDA and the time-of-flight system at COSY, SPHERE, the WASA 4 π detector, etc. The facilities themselves were discussed in a general perspective by Reistad and the set-ups at COSY by Schult. The targets were not explicitly discussed. They are also important for we must learn how to make collisions in the rings without destroying the beams. This is why the molecular beam targets, gas-jet targets, dust targets, etc. were touched on in various presentations. They represent a new generation of targets quite distinct from traditional accelerator targets. In this context of technical developments let me also remind you of the challenges of the kinematics in this novel environment. A splendid example of this is provided by thresholds at which all the reaction products are projected into a very small cone at 0° , so that small counters become 4π detectors. I will not dwell on these aspects, important as they are, for they are only the tools for the physics we wish to achieve. Let me single out just one novel item which may be of great importance. One of the basic problems in storage rings is that of cooling the beam as fast as possible. Normally this is done either by adjusting the individual particles (stochastic cooling) or by immersing the 'hot' beam in a 'cold' electron beam. In the latter case, if you look at the situation in the rest frame of the electrons, the protons move about like peas on a mat among the stationary electrons and come to rest as a result of 'friction' after a rather long time. This time is crucial for the operation and for the flexible use of the cooler ring, and a lot of advantages follow if it can be reduced. Ikegami proposed a new method for quick cooling of the beam. It is based on the idea of coupling the proton cyclotron frequency to the discrete frequencies of an applied radio frequency field under conditions such that the heat can be rapidly transferred between them. Depending on the exact conditions of the applied field the protons will either heat or cool. Ikegami compares this to a coherent maser effect. He has tested it on a small device and, indeed, some of

Figure 1: The development of a cooling focus according to Ikegami

the beam cools as seen in the outer ring in Fig. 1. The expected cooling time is 100 to 1000 times shorter than normal. It must now be established by research and development whether this can be scaled up and applied to a real ring.

3 Particle production at threshold in few-body systems

The next topic under discussion was threshold production phenomena. These issues are most specific to the cooler rings, since the kinematical conditions at threshold are quite unique and cannot be reproduced otherwise. In addition, the kinematical singularity, together with the very high energy resolution and the possibility to vary the energy, permits one to explore the detailed physics very close to the threshold with reasonable count rates in spite of the vanishing phase space at exact threshold. Here we heard three experimental talks of which the ones by Meyer and by Rohdjes concerned π^0 production, while Tegnér discussed an upcoming study of 2π production. There were theoretical talks by Ueda, Krippa, and Kondratyuk, inspired by the Saclay η production experiments, discussing what could be learned from them. The outstanding news concerned the first detailed study of the process $pp \rightarrow pp\pi^0$ with good statistics down to only about 2 MeV above threshold. Interestingly, there is still a non-negligible p wave even at these very low π^0 energies. More importantly, Meyer showed that the attempts to describe this process theoretically both by himself, by Niskanen, and by others, based on multiple scattering on the one hand and on three-body equations on the other, failed to reproduce the data by a factor of five in the absolute cross-section. This is a very serious matter indeed and there must be an important message here. The reason is that pion threshold phenomena are very closely linked to Chiral Symmetry, i.e., the symmetry which results in the limit of a vanishing pion mass or, alternatively, the limit in which the up- and down-quark masses can be considered to be vanishingly small. There are numerous consequences of Chiral Symmetry and even theorems. The most well-known examples are the Tomozawa-Weinberg theorems for the pion-anything scattering lengths, the Kroll-Ruderman theorem for threshold pion

Machner and Nieves discussed the prospects of how to form the very deep 'inaccessible' bound states in pionic atoms and how such states can be used as a tool for learning about the behaviour of the pion in the medium. The highlight of this was the experimental information in Gaarde's talk on the 0° scattering. Here, the Δ peak observed in the charge exchange scattering appears throughout the periodic system but this experimental ' Δ ' peak is shifted downward by a nearly universal amount. The interpretation of this shift is that it is evidence for the predicted pion mode, which is a novel type of collective nuclear mode quite distinct from the collective modes of nuclear spectroscopy.

You may wonder what is so special about 0° charge exchange reactions. The reason is that in this case the energy-momentum transfer ω, q acts like an incident pion with this energy and momentum, but this pion is a virtual one with $\omega^2 - q^2 \neq 0$. This is why charge exchange reactions with different projectiles like (p,n) , $(^3\text{He},t)$, ..., $(^{40}\text{Ar}, ^{40}\text{K})$ and at quite different incident energies have such similar features. There are, of course, distortions of various kinds which will change that, and the interaction gets confined to the edges of the nucleus, but in essence they are all more or less the same. The cleanest such examples are expected to come in neutrino physics from the forward (ν, e) and (ν, μ) reactions. In this case there are no distortions and there is even a theorem by Adler that states that the amplitude for any reaction is identical to the corresponding (unphysical) pion reaction with the same energy-momentum transfer but for a scale factor. So, in principle, neutrinos would be the ideal probes for this physics but for the killing small cross-sections and the poor energy definition of about 100 MeV, which hold back their use at present. To see qualitatively how the pion works, take the example of the (p,n) reaction and go to the nucleon rest system. The projectile is then surrounded by a virtual cloud of positive pions, since it converts virtually into a neutron by the emission of a π^+ with energy $\omega = 0$. The pion has a momentum distribution with momenta k . If now the momentum transfer in the reaction is due to the pion exchange, that momentum must be all in the beam direction for a 0° reaction. Now let us accelerate the projectile to its beam velocity β and let it undergo an energy loss ω . Now the pion is transformed relativistically so that its energy-momentum relation is

$$\omega = \beta \gamma k, \quad q_z = \gamma k.$$

So one has made a beam of virtual pions! The interest in this is that virtual pions explore properties of the nucleus which are inaccessible by real pions. In particular, one can now match the kinematical conditions for the pionic mode. All the 0° reactions we see on complex nuclei have the peak in the Δ region shifted down by about 70 MeV. This does not mean that the Δ mass has been shifted down in nuclear matter by such an amount. The Δ mass can be explored very well by photons and then the peak is nearly unshifted in the total photon cross-sections as seen in Fig. 3. This is so, since the photon is transverse. A coherent pion wave cannot propagate coherently in the forward direction to leading order. Then the Δ stands out very cleanly. Pions on the other hand like to go forward in the medium since they are longitudinal, but the coherent wave is strongly damped in the Δ region for physical pions. This is not so for virtual pions. If the energy-momentum relation for the pionic mode is matched, we will see the mode explicitly and that is just what the forward charge exchange reactions are good for. This collective state is neither a pure pion nor just a modification of the Δ . You should look at it as a coherent superposition

Figure 2: Single- and double-scattering mechanism in threshold π^0 production

photoproduction, and the Goldberger-Treiman relation relating the weak axial coupling constant to the πNN one. In other contexts it gives a deep insight into mesonic corrections to nuclear structure such as nuclear exchange currents. It also furnishes the link between nuclear axial currents and pion physics or, in other words, between nuclear spin physics, Gamow-Teller modes, and pionic modes. Not surprisingly, it is therefore a very important matter if something is not understood about a chiral phenomenon. In the present case of pion production the momentum transfer is about 600 MeV/c. This is rather large for chiral physics and we are on less familiar ground for the chiral manifestations. This is what makes it interesting. The basic processes can be viewed as direct production and as a cooperative two-nucleon effect in which the pion rescatters on the other nucleon as illustrated in Fig. 2. In the theoretical approaches so far, no special attention has been given to the chiral constraints on the rescattering process. In my opinion the crux of the matter lies in the proper understanding of the off-mass-shell structure of the process $\pi^0 p \rightarrow \pi^0 p$ for s wave pions. This issue must clearly be given careful theoretical attention. I believe the intrinsic uncertainty in the theory due to model-dependent physics to be no more than 20%. We should certainly learn much from that factor of five. At the same time as the theoretical effort, it would be valuable to extend such experiments so close to the threshold that the p wave can be eliminated. Good statistics is required.

4 Δ 's, π 's and pionic modes in nuclei

In this area there were a number of interconnected talks and in fact some of the material in the session on heavy-ion collisions is directly related to the issues raised here. The talks fall into two major categories: first the talks in which the modifications of the properties of pions and deltas embedded in nuclear matter were discussed. Here Gaarde gave a survey of the large amount of information that is now coming from charge exchange reactions at 0° and the interpretations they suggest. The theoretical talks of Osterfeld, Oset, Arve and Schepkin had all to do with the theoretical aspects raised by such experiments, in particular the issue of a pionic mode. In the second group of talks Yamazaki,

Figure 3: Total cross-sections for π and photons on a nucleus. Note that the Δ resonance is not shifted for photons

of $\pi + NN^{-1} + \Delta N^{-1}$ states or, alternatively and qualitatively, as the effect of the pion optical potential. In fact, the pion branch and the Δ branch cross at higher momenta transfers so there is a lot of mixing expected between the two branches as always when there is a level crossing. The issues that were discussed at the meeting were in particular:

Which are the decay channels and what are the characteristics?
How much of the process goes into an incoherent cascade process?
What is the density dependence of the pionic mode?

etc.

One of the interesting new features reported by Gaarde is that the $1\pi^+$ channel can now be seen and that it is very forward peaked, characteristic of a coherent process. The present situation is that the universality of the physics is qualitatively clear. It is equally clear that these phenomena are only at the beginning of exploration and that a large effort on the detailed features will be required so as to understand the properties of virtual pions in the nuclear environment.

The second topic of these discussions concerned the deeply bound atomic states of pions. Negative pions like electrons can be captured in a Coulomb field and will have typical Bohr states, but with a scale very different from that of electronic atoms. When the pion wave function overlaps with the nucleus they are exposed to the pion-nuclear strong interaction and absorption. The normal way to study these systems is via the spectroscopy of the X-rays emitted in the cascade from outer orbits to the innermost ones. Much has been learned from this, but there has always been a serious difficulty. The information in heavy nuclei always comes from high orbits and the outer skin of the nucleus and does not probe the pion-nuclear interaction in regions of higher densities.

The reason is that the pion in the nuclear Coulomb field is like a satellite in orbit around the earth. It spirals in towards the nucleus but is 'burned up' in the upper atmosphere by absorption before reaching the innermost orbits. There are sound reasons to believe that there will be deeply bound pion states of moderate width which live half inside the nucleus, half outside in the attraction of the Coulomb field. The probability of finding the pion in the most absorbing central region of the nucleus is quite small, so the absorptive width is expected to remain moderate. The problem is now the following: How can one implant a pion into such an orbit and directly determine its energy? The basic strategy is to make use of some reaction which produces a pion on a nucleon in the nucleus and get the pion captured directly into a bound state instead of being produced as a free particle. Its energy is readily measured for simple reactions. The snag is the cross-section, for there is usually a bad momentum mismatch for the capture into the bound state. Searches have so far been unsuccessful. There were two suggestions to overcome this. Yamazaki intends to use the (n,d) reaction making use of the fact that the neutron energy can be chosen such that there is vanishing momentum transfer in the (n,d) vertex. This does not mean that the momentum transfer can be neglected in the formation of the atom, since a nucleon is picked up in addition, but it can be kept within reason and calculations indicate that there is hope. Nieves suggested the (π, γ) reaction with the photon in the 30 MeV range, which guarantees low momentum transfer, but it may have experimental problems. Clearly the whole trick is to find the right practical conditions and in a sense it cannot really fail. After all, bound states exist, at least the pure Coulomb states.

Another point which touches on particle physics was raised by Schepkin. He pointed to the anomaly in pion double-charge exchange reactions on complex nuclei near the forward direction for pions of about 50 MeV energy. One possible explanation is that there exists a d' deuteron-like state, which then should have quantum numbers $J=0^-$, $T=0$ to have escaped detection. Some supporting arguments for such a low-lying two-nucleon state can be found from QCD. The real issue is an experimental one in my opinion. The evidence for such a state should come from simple reactions and it is there that the issue will be decided. For example, $pp \rightarrow d'\pi^+$ should directly display such a d' as a quasi two-body reaction.

5 Particle physics

In this area it is clear that cooler rings can contribute only to a limited number of problems, but some of them are essential tests of the Standard Model. This theory works extraordinarily well. The problem is that its success is just too great. It is not a unique theory, but really the simplest version of a class of theories. This is why it is necessary to test it up to the hilt to get a real feel for it in all circumstances. The contribution of cooler rings is that they can provide very high rates of tagged π^0 , η , η' , K^0 , etc. At the big particle accelerators, the tests are based on the idea that enough energy permits tests of forbidden processes by providing more phase space for them or by getting above thresholds. At the cooler rings, the tests are based on very high precision which also permits the exploration of short-distance phenomena, but of course the number of reactions is heavily restricted. Bergström reviewed the situation for rare decay modes and pointed in particular to the two-body decays of pseudoscalar mesons: $\pi^0 \rightarrow e^+e^-$ and $\mu^+\mu^-$, $\eta^0 \rightarrow e^+e^-$ and $\mu^+\mu^-$, etc. These are sensitive to non-standard components such as pseudoscalar Higgs-like particles or an observable CP-violating effect as exists in some

theories. The experimental challenge is the branching ratios of 10^{-7} to 10^{-8} which requires sophisticated 4π detectors like WASA. Imazato reviewed the K decay programme at KEK, while Cotanch described how the electromagnetic K production at CEBAF could be used to learn about the form factors of strange particles and hyperons.

Mizuno pointed out that there is a qualitative analogy between the spin-1 vector mesons and the spin-1 deuteron: the deuteron is a np system with a tensor interaction that gives it a d state, and a vector meson is predominantly a two-body $q\bar{q}$ system. It should therefore by analogy have a d-state component. He suggested to look for it in the radiative decay such as $\omega \rightarrow \pi\gamma$, which is analogous to the famous M1 threshold np capture reaction $np \rightarrow d\gamma$. This will not work, for there is no signature for the d state in a two-body decay and one must rely fully on dynamical evidence. In the M1 transition for the deuteron the rate is very insensitive to the d state. This is probably also so in the present case. One must find another signature. In principle three-body decays contain such evidence, but it is a quantitative question whether one can experimentally display it. Vector mesons should contain a d state like the deuteron simply because the QCD hyperfine interaction has a tensor component like the one-pion-exchange interaction in the deuteron. This also causes d-state components in the baryons of spin $3/2$ such as the Δ isobar, and there is a substantial effort going on to display it. Once the question is posed in this manner in terms of the d state and the tensor interaction, we can pose the question in a different way. The deuteron binding and its d state originates in the pion-exchange tensor interaction. Is it possible or not to produce mesonic 'molecules' of, say, two vector mesons held together by the strong tensor force of pion exchange just like for the deuteron? This should be further explored.

6 $\bar{N}N$ reactions

This session dealt mainly with two-body reactions of the type $\bar{N}N \rightarrow \bar{Y}Y$ or $B_i B_s$, where Y are hyperons and $B_{i,s}$ are mesons. These topics were covered by Johansson, Holinde and Oelert. The reactions $\bar{N}N \rightarrow \bar{Y}Y$ are particularly suited for cooler rings since they involve thresholds and the beautiful experiments from LEAR on $\Lambda\Lambda$ production are well known. The experimental situation was reviewed by Johansson. The theoretical issues here are concerned with the question whether the most appropriate and economical description is in terms of hadron physics or in terms of quark-gluon physics. It should always be kept in mind that these two descriptions are in no way in basic conflict. An exact description in terms of quark physics encompasses the hadron picture which is the practical description when we discuss strong interaction at long distance and with only moderate resolution. It is really at short range that the quark description comes fully into its own. Now, it was not pointed out that there is one reaction which has qualitatively similar features to the hyperon reaction and that is simply $p\bar{n} \rightarrow n\bar{p}$. This reaction has a strong peak in the 'forward' direction and it is very well described in terms of one-pion exchange to the extent that it would be foolish to attempt a description in explicit quark degrees of freedom. The hyperon reactions for Λ and Σ are similarly forward-peaked and it is then tempting to look at them in terms of meson exchange. Here is a list of the different possible reactions of this type with the corresponding exchange of lowest mass.

$\bar{\Lambda}\Lambda$	K, K^* exchange
$\bar{\Sigma}\Lambda$	K, K^* exchange
$\bar{\Sigma}\Sigma$	$\pi K, \pi K^*$ exchange
$\Xi\Xi$	2K, $2K^*$ exchange
$\bar{\Omega}\Omega$	3K, $3K^*$ exchange
$\bar{\Lambda}_c\Lambda_c$	D exchange

All of the last three involve the exchange of quite heavy objects, so that the interaction range will be quite short. These are the interactions which have not yet been explored experimentally. It seems to me very likely that there will be a real change of physics when we move over to the study of the heavy hyperons, and that in this case the meson exchange picture is likely to fail badly. As pointed out by Osterfeld both pictures describe the present data quite well. He also pointed to the difference in their consequences once you study polarization variables. In the present quark approaches there is no QCD hyperfine tensor force, while this is a key consequence of meson exchange. There are then huge differences in the predicted depolarization parameter D_{nn} . Polarization is a crucial test of the two pictures. While I agree, it is clear that the QCD tensor force exists also on the quark level, so the absence of this aspect is really a statement about the sophistication of present quark descriptions in this context.

Another hot topic was that the $\bar{\Lambda}\Lambda$ reaction has recently been measured accurately less than 1 MeV above threshold as reported by Johansson. The surprise is that there is a very rapidly changing, resonance-like structure near 1 MeV. At present not much is known about it but its narrow width indicates that it is not likely to be in the s wave. Perhaps it is the $^{-}^{-}^3D_1$ state suggested as a loosely bound $\bar{B}B$ state. Furthermore Johansson reported that the spin correlation in the $\bar{\Lambda}\Lambda$ system has just been measured, with the finding that they are always produced in a spin triplet state within experimental errors.

Oelert described the experimental situation in the reactions $\bar{N}N \rightarrow$ mesons. If we ask the question how such reactions should be described, it is then clear, because of the necessity of exchanging much heavier objects, that such processes are much more naturally described in terms of quarks than in terms of hadrons. However, from a practical viewpoint the situation is unclear.

Plendl described experiments on $\bar{p}$ absorption in nuclei. The interest in such experiments is that it is possible to deposit a large amount of energy in a localized region in the nucleus and that exotic phenomena may result. So far no such effects have been reported.

7 Heavy-ion reactions

A major part of this session was devoted to subthreshold production, in particular to pion production. By subthreshold we mean that the production occurs for an incident energy below the production for a free nucleon. This is, in a way, the earliest topic of meson physics but it has attracted renewed attention in recent years. In fact, at the end of the 1940s all the artificially produced pions were produced below threshold at accelerators like the one at Rochester with 225 MeV protons. It is a particularly appropriate topic here in Uppsala. The early accelerator here with 180 MeV protons was long known as the accelerator in the world with the highest energy not producing pions. Historically very important contributions to subthreshold production were made on this machine by a

particle interaction becomes negligible. The hope is now that such correlations are seen to eliminate the trivial effects such as quantum statistics and to control the less obvious contributions. For example, one neutron emitted may be overtaken by a slightly faster one with a rescattering outside the interaction region (proximity scattering). This effect is particularly pronounced for very small Δp , since the threshold NN interaction has very large scaling amplitudes.

8 Thanks

It is a great pleasure for me, on behalf of all the participants, to convey our thanks to all those here in Uppsala who have contributed to making this meeting such a success and who, at the same time, have provided us with a very enjoyable and friendly atmosphere. I know from experience that all these things that run so effortlessly and smoothly in reality correspond to a lot of planning, hard work, and even sleepless nights. A particular thank-you is due to Curt Ekström, who has been masterminding it all very efficiently. With this there only remains for me to take leave of you all. Here is a cartoon to remind you that on the road to the next workshop in Jülich, there is some homework to do and some problems to straighten out.

Reference

- [1] B. Höistad, Adv. Nucl. Phys. **11**, 135 (1979).

Figure 4: Coherent pion production in the ($p + {}^{12}\text{C}, \pi^0 + {}^{13}\text{C}$) reaction by 185 MeV protons [1]

couple of students preparing their thesis. Dahlgren and Höistad, with minimal equipment, succeeded from 1970 onwards in producing pions on many nuclei by the ($p, \pi^+ \pi^-$) reactions to specific states. In so doing they achieved at the time the best energy resolution of pions anywhere as Fig. 4 shows. The remarkable thing about these reactions is that the momentum transfer of 600 MeV/c is very large on the nuclear scale, but, even so, the nucleus acts coherently and there are various systematic features in the cross-sections.

It has become clear that there are also many interesting features in subthreshold collisions between complex nuclei. We heard about this in experimental and theoretical surveys by Jakobsson and Cassing, and in talks by Homolka and Laville. A theme that came back repeatedly in the discussions is the degree to which the nucleons conspire coherently to take up all the momentum necessary. Here one does not work all the way down to the lowest nuclear excitations. A comparison to establish the significance of these coherent effects is provided by the statistical approaches, which were discussed in particular by Zubkov. The statistical picture summarizes many features and indicates deviations that may point to new physics. All these subthreshold reactions are characterized by very rapid variation with energy in the threshold region, which reflects the fact that the phase space is rapidly constrained as the threshold is approached.

One of the most striking deviations is the quantum correlation when two like particles are emitted. The enormous degree of correlation that can result is apparent in the recent investigations reported by the CHIC collaboration. Here the neutrons correlate with a typical relative momentum of less than 10 MeV! Such correlations are closely related to the well-known Hanbury-Brown-Twiss effect between photons from far-away galaxies which are used to determine the size of the source. Similarly, here the aim is to use these correlations to learn about the size of the freeze-out region beyond which the

Figure 5