Protonium annihilation into $K_L^0 K_S^0 \pi^0$ and $K_L^0 K_S^0 \eta$

C. Amsler¹³, D. S. Armstrong¹, I. Augustin^{7,a}, C.A. Baker⁴, B.M. Barnett¹⁰,
C.J. Batty⁴, K. Beuchert², P. Birien¹, P. Blüm⁷, R. Bossingham¹, K. Braune¹¹,
J. Brose¹⁰, D.V. Bugg⁸, M. Burchell^{5,b}, T. Case¹, A. Cooper⁸, K.M. Crowe¹,
T. Degener², H.P. Dietz^{11,j}, S. von Dombrowski¹³, M. Doser⁵, W. Dünnweber¹¹,
D. Engelhardt⁷, M. Englert¹¹, M.A. Faessler¹¹, C. Felix¹¹, G. Folger^{11,c}, R. Glantz⁶,
R. Hackmann¹⁰, R.P. Haddock⁹, F.H. Heinsius⁶, N.P. Hessey⁵, P. Hidas³, P. Illinger¹¹,
D. Jamnik^{11,d}, Z. Jávorf³, H. Kalinowsky¹⁰, B. Kämmler⁶, T. Kiel⁶, J. Kisiel^{11,e},
E. Klempt¹⁰, M. Kobel⁵, H. Koch², C. Kolo¹¹, K. Königsmann^{11,f}, M. Kunze²,
R. Landua⁵, J. Lüdemann², H. Matthäy², M. Merkel^{10,c}, J.P. Merlo¹⁰, C.A. Meyer^{13,g},
L. Montanet⁵, A. Noble¹³, F. Ould-Saada¹³, K. Peters², C.N. Pinder⁴, G. Pinter³,
S. Ravndal², J. Salk², A.H. Sanjari^{8,h}, E. Schäfer¹⁰, B. Schmid^{13,i}, P. Schmidt⁶,
S. Spanier¹⁰, C. Straßburger¹⁰, U. Strohbush⁶, M. Suffert¹², D. Urner¹³, C. Voelcker¹¹,
F. Walter¹⁰, D. Walther², U. Wiedner⁶, N. Winter⁷, B. Zou⁸, J. Zoll⁵, Č. Zupančič¹¹

The Crystal Barrel Collaboration

Abstract

The annihilation of $\bar{p}p$ at rest into $K_L^0 K_S^0 \pi^0$ and $K_L^0 K_S^0 \eta$ was investigated with the Crystal Barrel detector at LEAR. These final states with negative C-parity are dominated by ϕ and K^{0*} resonances. The ratios of the branching ratios are $B(\bar{p}p \rightarrow \phi \pi^0)/B(\bar{p}p \rightarrow \phi \eta) = 8.3 \pm 2.1$ and $B(\bar{p}p \rightarrow K^0 \bar{K}^{0*} + \bar{K}^0 K^{0*} \rightarrow K_L^0 K_S^0 \pi^0)/B(\bar{p}p \rightarrow \phi \pi^0 \rightarrow K_L^0 K_S^0 \pi^0) = 2.04 \pm 0.21$.

To be submitted to Phys. Lett. B

¹ University of California, LBL, Berkeley, CA 94720, USA

² Universität Bochum, D-44780 Bochum, Germany

³ Academy of Science, H-1525 Budapest, Hungary

⁴ Rutherford Appleton Laboratory, Chilton, Didcot OX11, 0QX, UK

⁵ CERN, CH-1211 Genève, Switzerland

⁶ Universität Hamburg, D-22761 Hamburg, Germany

⁷ Universität Karlsruhe, D-76344 Karlsruhe, Germany

⁸ Queen Mary and Westfield College, London E14NS, UK

⁹ University of California, Los Angeles, CA 90024, USA

¹⁰ Universität Mainz, D-55099 Mainz, Germany

¹¹ Universität München, D-85748 München, Germany

¹² Centre de Recherches Nucléaires, F-67037 Strasbourg, France

¹³ Universität Zürich, CH-8001 Zürich, Switzerland

a) Now at University of Siegen, Germany

b) Now at University of Kent, Canterbury, UK

c) Now at CERN, Genève, Switzerland

d) On leave of absence from the University of Ljubljana, Ljubljana, Slovenia

e) On leave of absence from the University of Silesia, Katowice, Poland

f) Now at Max Planck Institut, Heidelberg, Germany

g) Now at Carnegie Mellon University, Pittsburgh, PA, USA

h) Now at State University of New York, Stony Brook, NY, USA

i) Now at University of California, Irvine, CA, USA

j) This work is part of the Ph.D. thesis of H.P. Dietz

The study of $\bar{p}p$ annihilation into final states with open or hidden strangeness is of special interest. Since, according to the Naive Quark Model [1], the $\bar{p}$ and p contain only $\bar{u}\bar{u}\bar{d}$ and uud constituent quarks, the s and $\bar{s}$ quarks in the final-state mesons must be created during the reaction. Alternatively, in a less naive model, they can be included as $\bar{s}s$ sea quark pairs in the wave function of the $\bar{p}$ and p . Thus, one question behind the special interest is how much $\bar{s}s$ pairs contribute to the (anti)baryon wave function at hadronic scales relevant to annihilation at rest (low Q^2) [2].

A related question is how well the OZI rule [3] applies to specific channels such as $\bar{p}p \rightarrow \phi\pi^0$ or $\bar{p}p \rightarrow \phi\pi^0\pi^0$. Indeed, the large size of the branching ratio $\bar{p}p \rightarrow \phi\pi^0$ [4, 5], which seems to violate the OZI rule quite strongly, has been considered to indicate either a large amount of $\bar{s}s$ in the proton wave function [2], or the existence of a $\bar{q}q\bar{s}s$ crypto-exotic intermediate state [6, 7] or, more mundanely, final-state interaction between K^0 and $\bar{K}^0$ produced via an OZI-allowed intermediate state [8].

Two channels which have never been studied before are

$$\bar{p}p \rightarrow K_L^0 K_S^0 \pi^0 \quad (1)$$

and

$$\bar{p}p \rightarrow K_L^0 K_S^0 \eta. \quad (2)$$

In the present experiment they are detected via the ‘zero-prong’ final states where only photons are detected, i.e. via the $K_S^0 \rightarrow 2\pi^0 \rightarrow 4\gamma$ and $\eta \rightarrow 2\gamma$ decay modes. The K_L^0 escaping detection with a probability of about 50% is identified by missing mass. We will show that $K_L^0 K_S^0 \pi^0$ is dominated by the two-body intermediate states $K^{0*}\bar{K}^0$, $\bar{K}^{0*}K^0$ and $\phi\pi^0$, and $K_L^0 K_S^0 \eta$ is dominated by $\phi\eta$. The latter has not been measured before in liquid hydrogen, but the $K^0\bar{K}^{0*} + \bar{K}^0 K^{0*}$ and $\phi\pi^0$ branching ratios are already fairly well known from charged final states [4, 5, 9, 10]. The measurement of absolute branching ratios requires a precise knowledge of the interaction probability of K_L^0 in the calorimeter. An analysis is in progress to determine this quantity and will be published in a forthcoming paper, together with absolute branching ratios into various channels containing K_L^0 ’s. The present investigation is restricted to obtaining relative branching ratios of $K^0\bar{K}^{0*} + \bar{K}^0 K^{0*}$, $\phi\pi^0$, and $\phi\eta$ with small statistical and systematic errors.

The Crystal Barrel detector [11] has been set up at the Low-Energy Antiproton Ring (LEAR) at CERN to investigate $\bar{p}p$ annihilations at rest and up to about 2 GeV/c incoming $\bar{p}$ momentum. It has been designed for high-statistics measurements of almost all possible final states and has access to many channels which have not been investigated before. For channels which are already known it can reduce the present statistical and systematic uncertainties.

The detector covers 98% of the full solid angle and is sensitive both to charged particles and photons. Charged particles are observed and measured in two proportional wire chambers (PWCs) and a cylindrical jet drift chamber (JDC). Photons, for example those

produced in the decays of $\pi^0 \rightarrow \gamma\gamma$, $\eta \rightarrow \gamma\gamma$, etc., are measured in the electromagnetic (e.m.) calorimeter which surrounds the drift chamber. As the results presented here primarily depend on the photon calorimeter, we briefly describe its relevant characteristics. A detailed description of the detector can be found elsewhere [11].

The calorimeter consists of 1380 CsI(Tl) crystals, 16 radiation lengths thick, read out by photodiodes. It covers polar angles θ between 12° and 168° and has complete coverage in azimuth. Photons which hit the crystals closest to the acceptance holes ($\theta < 12^\circ$ and $\theta > 168^\circ$) of the detector are rejected in the analysis (see below). This reduces the solid angle for photon detection, leading to an acceptance of 95% of 4π . The energy calibration of the crystals is achieved using the 2γ decay mode of the π^0 , abundantly produced in $\bar{p}p$ annihilations, by imposing the standard mass of $134.974 \text{ MeV}/c^2$ on the reconstructed π^0 . The photon energy resolution is

$$\frac{\sigma_E}{E} = \frac{2.45\%}{(E/\text{GeV})^{1/4}}$$

for the data used in this work. The angular resolution is $\sigma = 20 \text{ mrad}$ in the polar and azimuthal angles for isolated high-energy showers.

The trigger for an incident antiproton is provided by a silicon beam counter placed just before the liquid-hydrogen target. This trigger is combined with a veto against annihilations producing charged particles, using the on-line information from the PWCs. The resulting ‘zero-prong’ trigger permits a large increase in the data-acquisition rate for annihilations into final states with only neutral particles, which represent about 4% of all annihilations. In this way several million zero-prong events have been accumulated during several running periods.

The starting sample for the analysis of the reactions (1) and (2) contains $4.5 \cdot 10^6$ zero-prong events corresponding to 107 million $\bar{p}p$ annihilations at rest. First, $4 \cdot 10^5$ events are rejected which show charged tracks in the JDC after off-line reconstruction, having escaped on-line PWC detection. (A very small PWC inefficiency, γ conversions, and charged K_S^0 decays are the cause of these events). Next, all events not having exactly six well-isolated e.m. showers with a minimum energy of 20 MeV are rejected. This cut removes about 80% of the data sample. It also reduces the detection efficiency since not every photon will manifest itself as one separated e.m. shower: some of the photons escape through the holes of the detector; others merge with another photon or, as result of shower fluctuations, appear as two separated energy deposits.

Since among the products of reactions (1) and (2), only π^0 , η , and K_S^0 are detected while K_L^0 is unobserved, the events of interest are characterized by missing energy. Therefore, a cut on the total energy deposit in the calorimeter is imposed which corresponds to the possible range of visible energy in these reactions. Of the $8.8 \cdot 10^5$ events, kept up to this point, all events are rejected which do not fulfill the condition:

$$900 \text{ MeV} < \sum_{i=1}^6 E_i < 1400 \text{ MeV}.$$

In this way the bulk of background events such as $\bar{p}p \rightarrow 3\pi^0 \rightarrow 6\gamma$ or $\bar{p}p \rightarrow 2\pi^0\eta \rightarrow 6\gamma$ is removed, leaving 58000 events. In order to remove further background from events with higher photon multiplicities where one or more photons are lost, events with missing momentum pointing towards one of the detector holes are rejected. Furthermore we demand that an e.m. shower does not spread its energy among adjacent crystals in such a way that the maximum energy fraction is deposited in a crystal on the border of one of the holes. Applying these cuts reduces the number of events to 37270.

The quality of the data obtained at this level is illustrated by fig. 1a, which shows the distribution of the missing mass squared

$$m_{miss}^2 = (2m_p - \sum_{i=1}^6 E_i)^2 - (\sum_{i=1}^6 \vec{p}_i)^2,$$

where a K_L^0 signal shows up clearly. (The probability to loose a K_S^0 is negligible.) The hypothesis

$$\bar{p}p \rightarrow K_L^0 6\gamma$$

permits a one-constraint (1C) kinematic fit which is survived by 10305 events with a confidence level (CL) larger than 5%. At this stage there is clear evidence for a K_S^0 signal. This is illustrated by fig. 1b which is obtained for a subsample of the 10305 events where the six photons combine to give three neutral pions; the invariant mass spectrum of two of these pions is shown (3 entries per event).

In the next step a 4C kinematic fit is applied to the 10305 events according to the hypothesis

$$\bar{p}p \rightarrow K_L^0 K_S^0 \gamma\gamma \quad (K_S^0 \rightarrow \pi^0 \pi^0).$$

This fit yields 4395 events with $CL > 1\%$. Figure 1c shows the invariant mass distribution of the remaining two photons. A prominent π^0 signal and a small η signal can be seen, which correspond to the reactions (1) and (2). Finally, two 5C kinematic fits are applied to these events in order to separate reactions (1) and (2), making the approximation that K_S^0 decays at the primary vertex. In order to compensate for this approximation the errors on the photon direction were appropriately increased in the fit. The data are split into two subsamples:

- (I): 2834 events with $CL > 10\%$ for reaction (1) and $CL < 1\%$ for reaction (2).
- (II): 72 events with $CL > 10\%$ for reaction (2) and $CL < 1\%$ for reaction (1).

The probability distributions for these fits are flat above 10%, showing that the quality of the final event samples is acceptable and the measurement errors on the kinematical variables are correct. It was checked by Monte-Carlo simulation that the feed-through from reaction (1) to (2) is 4 ± 2 events while the feed-through from (2) to (1) is less than one event. The number of background events in these two subsamples is estimated from the numbers of entries in the invariant mass distribution of the two remaining photons, excluding the π^0 and the η signals. This gives 120 ± 7 background events in the $K_L^0 K_S^0 \pi^0$ subsample

and 26 ± 8 background events in the $K_L^0 K_S^0 \eta$ subsample. The K_S^0 and K_L^0 energy distributions of the events outside the π^0 and η signal suggest that the background events are homogeneously distributed over the allowed phase space. The overall efficiencies (ϵ) to reconstruct and identify reactions (1) and (2) in the neutral decay mode, assuming no K_L^0 interaction and including all cuts and kinematic fits, were calculated with a complete Monte-Carlo simulation based on GEANT [12]. They are $\epsilon = (21.3 \pm 0.5)\%$ for reaction (1) and $\epsilon = (22.0 \pm 2.0)\%$ for reaction (2). Their relative variation $\Delta\epsilon/\epsilon$ over the full phase space is less than 5%.

Figures 2a to 2d show the $K_L^0 K_S^0 \pi^0$ Dalitz plot and its three projections for the event sample (I) using the momentum values from the kinematic fit. This Dalitz plot exhibits the following prominent features: an accumulation of events in the two $K^*(892)$ regions and a sharp peaking of events for which the $(K_L^0 K_S^0)$ effective mass is in the $\phi(1020)$ region. We observe a slight asymmetry between the $m^2(K_L^0 \pi^0)$ and $m^2(K_S^0 \pi^0)$ projections. This asymmetry is due to a small systematic error in our analysis method. There is a threefold combinatorial ambiguity as to which two of the three π^0 s form the K_S^0 . The ambiguity affects the $K_L^0 \pi^0$ and the $K_L^0 K_S^0$ (but not the $K_S^0 \pi^0$) invariant mass. This combinatorial effect was studied with Monte Carlo simulations. From this study we determined that $4.8 \pm 0.6\%$ of the $\phi\pi^0$ events and $7.8 \pm 0.7\%$ of the $K_S^0 K^{0*}$ events give wrong $\pi^0\pi^0$ combinations. This explains the different heights of the two K^{0*} peaks. In figs. 2e and 2f the decay cosine distribution of $\phi \rightarrow K_L^0 K_S^0$ and $K^{0*} \rightarrow K^0 \pi^0$ are shown by selecting events with $m^2(K_L^0 K_S^0) < 1.1 \text{ GeV}^2$ and $0.7 \text{ GeV}^2 < m^2(K^0 \pi^0) < 0.9 \text{ GeV}^2$. They both exhibit the angular distribution expected for vector resonances produced from the $\bar{p}p$ initial state 3S_1 (see below).

Figures 3a and 3b show the $K_L^0 K_S^0 \eta$ Dalitz plot of the event sample (II) and its $m^2(K_L K_S)$ projection (fitted quantities). The ϕ signal is prominent, while the other regions of the Dalitz plot are sparsely populated – mainly by background events.

As the final state has C -parity = -1 , it is reasonable to assume that the production is dominated by the $\bar{p}p$ initial state 3S_1 . We fit the $K_L^0 K_S^0 \pi^0$ Dalitz plot with transition amplitudes of the channels $\bar{p}p(^3S_1) \rightarrow \phi\pi^0$ and $\bar{p}p(^3S_1) \rightarrow K^{0*}K$ using the maximum likelihood method. An intensity distribution of the form

$$I = |\alpha_1 \vec{A}(^3S_1, \phi(1020)) + \alpha_2 \vec{A}(^3S_1, K^{0*}(892))|^2$$

is assumed with complex parameters α_i . The transition amplitudes $\vec{A}(^3S_1, \phi(1020))$ and $\vec{A}(^3S_1, K^{0*}(892))$ consist of a function $\vec{\chi}$ which describes the decay distribution of the resonances, satisfying total angular momentum and parity conservation, and a relativistic Breit-Wigner factor $F(m)$. The decay distribution functions $\vec{\chi}$ are constructed according to the method developed by Zemach [13], where the three scalar amplitudes corresponding to the three magnetic substates of the initial $J=1$ protonium state are represented by the three components of a vector amplitude $\vec{A}$. We have

$$\vec{A}(^3S_1, \phi(1020)) = \vec{\chi}(\phi) \cdot F(m(K_L^0 K_S^0)). \quad (3)$$

The amplitude $\vec{A}({}^3S_1, K^{0*}(892))$ contains two terms, each corresponding to an eigenstate of charge conjugation with $C = -1$:

$$\vec{A}({}^3S_1, K^{0*}(892)) = \vec{\chi}(K_L^*) \cdot F(m(K_L^0\pi)) - \vec{\chi}(K_S^*) \cdot F(m(K_S^0\pi)). \quad (4)$$

Here, the symbols K_L^* and K_S^* mean that the invariant mass of π^0 and K_L^0 or K_S^0 is in the region of the $K^{0*}(892)$ resonance. We have $K_L^* = (K^{0*} + \bar{K}^{0*})/\sqrt{2}$ and $K_S^* = (K^{0*} - \bar{K}^{0*})/\sqrt{2}$. The minus sign between the two terms of eq. (4) projects the part with zero strangeness out of the final-state wave function containing $K_L^0 K_S^*$ and $K_S^0 K_L^*$. The angular distribution of the ϕ -decay is given by

$$\vec{\chi}(\phi) = \vec{q}_K \times \vec{p}_{KK},$$

where $\vec{q}_K$ is the 3-momentum of the K_L^0 in the $K_L^0 K_S^0$ rest frame and $\vec{p}_{KK}$ is the 3-momentum of the $K_L^0 K_S^0$ system in the laboratory. The angular distributions of the K^{0*} -decays are described by

$$\vec{\chi}(K^{0*}) = \vec{q}_K \times \vec{p}_{K\pi},$$

where $\vec{q}_K$ is the 3-momentum of the $K_L^0(K_S^0)$ in the $K_L^0\pi^0(K_S^0\pi^0)$ rest frame and $\vec{p}_{K\pi}$ is the 3-momentum of the $K_L^0\pi^0(K_S^0\pi^0)$ system in the laboratory.

The relativistic Breit-Wigner factors for the ϕ and the K^{0*} resonances are

$$F(m) = \frac{W(p) \cdot W(q)}{q \cdot p \cdot R^2} \frac{m_o \Gamma_o}{m^2 - m_o^2 + i \cdot m_o \Gamma(m)},$$

with

$$\Gamma(m) = \Gamma_o \frac{m_o}{m} \frac{q}{q_o} \frac{W^2(q)}{W^2(q_o)},$$

where m_o , Γ_o are the nominal mass and width of the resonances, $q = |\vec{q}|$ is the decay momentum in the resonance rest frame ($q = q_o$, if $m = m_o$), and $p = |\vec{p}|$ is the laboratory momentum of the resonance. The function $W(q)$ is the Blatt-Weisskopf damping factor for angular momentum $\ell = 1$ [14]. (The angular momentum is one unit between the resonances and the recoiling particle, as well as between the decay particles of the resonance). Therefore, W is

$$W(q) = (q \cdot R) \left(\frac{1}{1 + (q \cdot R)^2} \right)^{1/2}$$

with $R = (0.197327 \text{ GeV})^{-1}$ ($=1$ fermi).

The complex parameters $\alpha_i = |\alpha_i| \cdot \exp(i\phi_i)$ are left free in the fit. The phase of α_1 is arbitrary and therefore fixed at 0. Since the natural width of the ϕ -resonance, $\Gamma_\phi = 4.4$ MeV, is nearly of the same magnitude as the experimental resolution $\sigma = 2.5$ MeV, the intensity distribution I is convoluted with a Gaussian curve which represents the resolution along the $m^2(K_L^0 K_S^0)$ axis. A constant intensity is added in order to account for the 120 background events mentioned above. The results of the fit are given in table 1. The

Table 1: Amplitudes and phases of the fit

Initial state	Intermediate state	Amplitude and phase $\alpha_j = \alpha_j \cdot \exp(i\phi_j)$	Ratio (%) $= \alpha_j ^2 \cdot 100$
3S_1	$\phi(1020)$	$ \alpha_1 = 0.57 \pm 0.01$ $\phi_1 = 0$	32.9 ± 1.1
3S_1	$K^{0*}(892)$	$ \alpha_2 = 0.82 \pm 0.01$ $\phi_2 = -1.64 \pm 0.11$	67.1 ± 1.5

normalization is such that $|\alpha_1|^2 + |\alpha_2|^2 = 1$.

In order to test the consistency of our analysis method we generated a Monte-Carlo data set of $K_L^0 K_S^0 \pi^0$ events with a Dalitz plot distribution according to the fit result given in table 1. These Monte-Carlo data were then analysed exactly like the real data, as described above. The spectra that result from this analysis are superimposed onto the spectra of the real data in the figs. 2a to 2f. The agreement between the real data and the simulation is satisfactory. A χ^2 -test was applied to test the compatibility of this simulation and the data. The χ^2 per degree of freedom is 1.2. Since the data are well reproduced by the set of amplitudes that we have assumed, we conclude that the final state $K_L^0 K_S^0 \pi^0$ is produced almost entirely from the 3S_1 state. We do not need any substantial contribution from the 1P_1 initial state. A priori, the latter cannot be excluded if we assume that it feeds the resonance both via an $l = 0$ and an $l = 2$ transition. However, it would take an improbable accident of nature for the 1P_1 state to produce an angular distribution as close to $\sin^2\theta$ as we observe it.

Estimates, based on our own data, of the probability $\bar{P}_{int}$ that the K_L^0 does not interact or the interaction is not detected in the calorimeter, yield $\bar{P}_{int} \approx 50 \pm 10\%$. This number has a relatively large systematic error but it cancels in the ratios of branching ratios unless $\bar{P}_{int}$ depends on the K_L^0 momentum. Such a dependence would show up in a systematic deviation between the data and the fit which assumes $\bar{P}_{int}$ to be constant as a function of the $K_S^0 \pi^0$ invariant mass (or the K_L^0 momentum, alternatively). In fact, the data are consistent with no momentum dependence within the statistical errors and the relevant momentum range. They allow the extraction of upper limits for the momentum dependence of $\bar{P}_{int}$; these are taken into account as additional systematic uncertainty of 7.5% for the ratio $B(\bar{p}p \rightarrow K^{0*} \bar{K}^0 + \bar{K}^{0*} K^0)/B(\bar{p}p \rightarrow \phi \pi^0)$ and of 10% for the ratio $B(\bar{p}p \rightarrow \phi \pi^0)/B(\bar{p}p \rightarrow \phi \eta)$.

We define the number of $\phi \pi^0$ events as the product of $|\alpha_1|^2$ and the number of detected $K_L^0 K_S^0 \pi^0$ events. This yields 893 $\phi \pi^0$ events compared to 737 events if we simply count the number of events in the ϕ peak. The latter number is significantly lower, firstly because events in the high mass tail of the ϕ -resonance are not counted, secondly because events are lost from the peak due to wrong $\pi^0 \pi^0$ combinations as explained above. The ratio $B(\bar{p}p \rightarrow K^0 \bar{K}^{0*} + \bar{K}^0 K^{0*})/B(\bar{p}p \rightarrow \phi \pi^0)$ is obtained as $|\alpha_2|^2/|\alpha_1|^2$.

For the determination of the ratio of $B(\bar{p}p \rightarrow \phi \pi^0)$ and $B(\bar{p}p \rightarrow \phi \eta)$, we count the $\phi \pi^0$ and $\phi \eta$ events by setting a window on the invariant $K_L^0 K_S^0$ mass ($m^2 \leq 1.1 \text{ GeV}^2$), and sub-

tract the estimated background (obtaining $737 \pm 27 \phi\pi$ and $35 \pm 7 \phi\eta$ events). We use the branching ratios $B(\pi^0 \rightarrow \gamma\gamma) = 0.988$ and $B(\eta \rightarrow \gamma\gamma) = 0.389$ [15].

Systematic errors of the results (other than those due to $\bar{P}_{int}$) were investigated by varying the cuts on confidence levels, minimum shower energies, total energy deposits, and varying the number of allowed $\pi^0\pi^0 \rightarrow K_S^0$ combinations, by variation of the ϕ -window, the active solid angle, and the annihilation vertex in the Monte-Carlo simulation. These studies lead to a systematic error of $\sim 5\%$ of the $\phi\pi^0$ signal, and we assume the other signals to have a comparable systematic error. This is included in the final systematic errors of the ratios of branching ratios, although most effects considered above are expected to cancel to a substantial degree.

We find

$$\frac{B(\bar{p}p \rightarrow \phi\pi^0)}{B(\bar{p}p \rightarrow \phi\eta)} = 8.3 \pm 1.7_{stat} \pm 1.3_{syst}$$

$$\frac{B(\bar{p}p \rightarrow K^{0*}\bar{K}^0 + \bar{K}^{0*}K^0 \rightarrow K_L^0 K_S^0 \pi^0)}{B(\bar{p}p \rightarrow \phi\pi^0 \rightarrow K_L^0 K_S^0 \pi^0)} = 2.04 \pm 0.10_{stat} \pm 0.19_{syst}.$$

Here, the lower indices *stat* and *syst* indicate the statistical and the systematic error. The final systematic error squared of the first (second) ratio is the sum of the squares of the 10% (2%) statistical error of the efficiency, the 10% (7.5%) uncertainty concerning $\bar{P}_{int}$ variations, and the 5% (5%) systematic error mentioned last.

These ratios provide a useful additional input to $\bar{p}p$ annihilation models, in particular since we are dealing with a final state of specific C-parity -1 . This reduces the possible initial $\bar{p}p$ states to $J^{PC} = 1^{--}$ (s-wave) and $J^{PC} = 1^{+-}$ (p-wave). It has been shown that only the 1^{--} state is needed to describe the Dalitz plot in a satisfactory way. This result is plausible: s-wave annihilation is dominant, in general (for a recent publication see ref. [16]) and, in addition, dynamical selection rules appear to favour the $J^{PC} = 1^{--}$ initial state [17]. The isospin is 1 for the $\phi\pi^0$ channel and 0 for $\phi\eta$. A complete isospin analysis of the $K^{0*}K$ channel has been carried out in the classic paper on $\bar{p}p \rightarrow K\bar{K}\pi$ by Barash et al. [9]. This analysis shows that in our case the population density of the Dalitz plot is independent of isospin, which therefore, cannot be determined solely from studying the $K_L^0 K_S^0 \pi^0$ channel.

In conclusion, we observe that the branching ratios for the channels $\bar{p}p \rightarrow K_L^0 K_S^*$, $K_S^0 K_L^*$ and $\phi\pi^0$ are nearly equal, thus confirming a previous observation that the branching ratio into $\phi\pi$ is not OZI suppressed. However, in contrast, using the branching ratio $B(\bar{p}p \rightarrow \phi\pi) = (4.0 \pm 0.8) \cdot 10^{-4}$ from ref. [5] and our ratio $B(\phi\pi^0/\phi\eta)$ we obtain $B(\bar{p}p \rightarrow \phi\eta) = 0.48 \pm 0.17 \cdot 10^{-4}$. This value is as small, compared to the branching ratio $B(\bar{p}p \rightarrow \omega\eta) = (10.4 \pm 1.0) \cdot 10^{-3}$ [19] or $= (15.1 \pm 1.2) \cdot 10^{-3}$ [20], as expected from the OZI rule. Tentative explanations of this phenomenon were suggested by several authors. Furui et al. [8] argued that the kaon from K^{0*} decays rescatters off the directly produced kaon thus forming

ϕ mesons via an OZI-allowed intermediate state. The small ratio for $\phi\eta$ compared to $\omega\eta$ can be explained by the absence of a strange meson decaying into $K\eta$. However, other final states which can be reached by rescattering from OZI-allowed intermediate states should be enhanced as well –this does not seem to be the case. Similarly, a large amount of $\bar{s}s$ in $\bar{p}p$ cannot explain the peculiarity of the $\phi\pi/\omega\pi$ ratio; it would presumably simultaneously affect all branching ratios with a ϕ meson. Perhaps, the absence of an OZI suppression in the particular case of $\phi\pi^0$ can be explained by the absence of cancellation between various intermediate states as discussed by Lipkin et al. [18]. Alternatively, Close and Lipkin [6] and Dover and Fishbane [7] associate enhanced $\phi\pi$ production with the existence of a crypto-exotic 4-quark state. This $\bar{s}s\bar{q}q$ state is identified by the latter authors with the $\phi\pi$ resonance C(1480) [21], which is supposed to couple to the $\bar{p}p$ system. However, the small branching ratio into $\phi\eta$ excludes the hypothetical [7] crypto-exotic $I=0$ partner of the C(1480) meson. The present situation certainly asks for more data on ϕ production. An analysis, similar to the one described in this paper, is in progress on the channel $\bar{p}p \rightarrow K_L^0 K_S^0 \gamma$ which should provide interesting new information, when compared with the recently published branching ratio $\bar{p}p \rightarrow \omega\gamma$ [22].

Acknowledgements

We would like to thank the technical staff of the LEAR machine group and of all the participating institutions for their invaluable contributions to the success of the experiment. We acknowledge financial support from the German Bundesministerium für Forschung und Technologie, the Schweizerischer Nationalfonds, the British Science and Engineering Research Council, and the U.S. Department of Energy (contract No. DE-FG03-87ER40323 and DE-AC03-76SF00098). J.K. and K.M.C. acknowledge support from the A. von Humboldt Foundation. We wish to thank K. Steinberger and A. Loesch for the system management on the computer cluster in Garching.

References

- [1] J.J.J. Kokkedee, *The Quark Model* (Benjamin, New York 1969).
- [2] J. Ellis, E. Gabathuler, and M. Karliner, *Phys. Lett. B* 217 (1989) 173.
- [3] S. Okubo, *Phys. Lett. B* 5 (1963) 165; G. Zweig, CERN Report No.8419/TH412 (1964); I. Iizuka, *Progr. Theor. Phys. Suppl.* 37 (1966) 21.
- [4] M. Chiba et al., *Phys. Rev. D* 38 (1988) 2021; M. Chiba et al., *Phys. Lett. B* 202 (1988) 447.
- [5] J. Reifenröther et al., *Phys. Lett. B* 267 (1991) 299.
- [6] F.E. Close and H.J. Lipkin, *Phys. Rev. Lett.* 41 (1978) 1263.
- [7] C.B. Dover and P.M. Fishbane, *Phys. Rev. Lett.* 62 (1989) 2917.
- [8] S. Furui, *Z. Phys. C* 46 (1990) 621; S. Furui, G. Stubel, A. Faessler, and R. Vinh Mau, *Nucl. Phys. A* 516 (1990) 643.
- [9] N. Barash et al. *Phys. Rev.* 139B (1965) 1659; B. Conforto et al., *Nucl. Phys. B* 3 (1967) 469.
- [10] A. Bettini et al., *Nuovo Cimento* 63A (1969) 1199.
- [11] E. Aker et al., *Nucl. Instrum. Methods A* 321 (1992) 69.
- [12] R. Brun et al., GEANT3, Internal Report CERN DD/EE/84-1, CERN, 1987.
- [13] Ch. Zemach, *Phys. Rev. B* 140 (1965) 97 and 109.
- [14] J.M. Blatt and V. Weisskopf, *Theoretical Nuclear Physics* (Wiley, New York, 1952).
- [15] K. Hikasa et al., *Phys.Rev. D* 45 (1992) Part II: Review of Particle Properties.
- [16] C. Amsler et al., *Phys.Lett. B* 297 (1992) 214.
- [17] C.B. Dover, T. Gutsche, M. Maruyama and A. Faessler, *Prog. Part. Nucl. Phys.* 29 (1992) 87.
- [18] H.J. Lipkin, *Nucl. Phys. B* 244 (1984) 147 and *Nucl. Phys. B* 291 (1987) 720.
- [19] L. Adiels et al., *Z. Phys. C* 42 (1989) 49.
- [20] C. Amsler et al., *Z. Phys. C* 58 (1993) 175.
- [21] S.I. Bitukov et al., *Phys. Lett. B* 188 (1987) 383. L.G. Landsberg, *Sov. J. Nucl. Phys.* 55 (1992) 1051.
- [22] C.Amsler et al. *Phys. Lett. B* 311 (1993) 371.

Figure captions

Figure 1: Invariant mass distribution: a) missing mass of events with 6 e.m. showers after a cut on the total visible energy; b) $\pi^0\pi^0$ mass after the 1C kinematic fit for events with three π^0 ; c) and d) $\gamma\gamma$ mass after the 4C kinematic fit.

Figure 2: a) Dalitz plot and b) to d) projections for the reaction $\bar{p}p \rightarrow K_L^0 K_S^0 \pi^0$. Decay cosine distributions e) in the rest frame of ϕ ($m_{K_L^0 K_S^0}^2 \leq 1.1 \text{ GeV}^2$) and f) in the rest frame of K^{0*} ($0.7 \leq m_{K_L^0 K_S^0}^2 \leq 0.9 \text{ GeV}^2$); the peak at -0.4 is due to the crossing with the other K^{0*} band.

Figure 3: a) Dalitz plot and b) projection onto the $K_L^0 K_S^0$ invariant mass for the reaction $\bar{p}p \rightarrow K_L^0 K_S^0 \eta$.

Figure 1

Figure 2

Figure 3