CERN-TH.7416/94

HIGHER-ORDER CORRECTIONS TO THE LARGE- N_c BOUND ON $M_\eta/M_{\eta'}$

S. Peris^{*,†}

Theoretical Physics Division, CERN
CH - 1211 Geneva 23

ABSTRACT

Next-to-leading corrections to the upper bound on $M_\eta/M_{\eta'}$ recently obtained by Georgi are considered. These corrections are just what is needed to reconcile the bound with the observed η and η' masses.

^{*} Invited talk given at the workshop "QCD 94", Montpellier, France, July 1994. To appear in the Proceedings.

[†] Work partially supported by research project CICYT-AEN93-0474. On leave from Grup de Física Teòrica and IFAE. Universitat Autònoma de Barcelona, Spain. E-mail: PERIS@CERNVM.CERN.CH.

CERN-TH.7416/94
August 1994

Higher-order corrections to the large- N_c bound on $M_\eta/M_{\eta'}$

S. Peris^{*,†}

Theory Division, CERN, CH-1211 Geneva 23, Switzerland.

Next-to-leading corrections to the upper bound on $M_\eta/M_{\eta'}$ recently obtained by Georgi are considered. These corrections are just what is needed to reconcile the bound with the observed η and η' masses.

In a recent paper, Georgi[1] has noticed an amusing fact concerning the large- N_c limit of QCD[2] and the η - η' mass matrix[3]. To lowest nontrivial order in $1/N_c$ and in the quark masses he has found that

$$\begin{aligned} \frac{M_\eta^2}{M_{\eta'}^2} &\leq \frac{3-\sqrt{3}}{3+\sqrt{3}} \left[1 + \frac{\sqrt{3}}{2} \frac{m_u+m_d}{m_s} \right. \\ &\quad \left. + \mathcal{O}\left(\left(\frac{m_u+m_d}{m_s}\right)^2\right) \right] \\ &\approx 0.268 \left[1 + \frac{1}{10} + \dots \right]. \end{aligned} \quad (1)$$

The experimental number for this ratio, i.e. 0.328(7)[4], is higher than this upper bound. It is therefore impossible to fit the experimental masses within this approximation. As I shall show[5], higher orders eliminate this bound. Higher orders come with new operators and unknown coefficients that make the parameter space larger and, consequently, they may avoid the constraints that lead to the bound (1). However, consistency of the perturbative picture requires that these higher orders be "small" corrections to the lowest-order result. It is only because the bound (1) and the experimental masses are very close that this is in principle possible.

The large- N_c limit gives us an interesting way (if not the only one) to treat the physics of the η' from first principles, i.e. from QCD. In this limit, and in a world of massless quarks, the η' would

^{*}Invited talk given at the workshop "QCD 94", Montpellier, France, July 1994. To appear in the proceedings.
[†]Work partially supported by research project CICYT-AEN92-0474. On leave from Grup de Física Teòrica and IFAE. Universitat Autònoma de Barcelona, Spain. E-mail: PERIS@CERNVM.CERN.CH

become the Goldstone boson of the $U(1)_A$ symmetry present in the QCD Lagrangian[3]. It is then expected that the real world can be reached from this limit by a combined perturbative expansion in the quark mass, m_q , and in $1/N_c$.

The next-to-leading order involves operators of $\mathcal{O}(m_q)$, $\mathcal{O}(1/N_c^2)$ and $\mathcal{O}(m_q/N_c)$. We shall next see that only the latter are in fact relevant for overcoming the bound (1).

First I shall show that higher-orders in the quark mass alone cannot reconcile the bound (1) with the empirical masses because this bound persists, in fact, to all orders in m_q in the limit $N_c \rightarrow \infty$ and not only to first order, as originally derived in ref. [1].

Let us start with the QCD Lagrangian with massive quarks in the limit $N_c \rightarrow \infty$. In this limit OZI violating transitions are strictly forbidden. The mass matrix, $\mathcal{M}$, for the π^0 ($= \frac{u\bar{u}-d\bar{d}}{\sqrt{2}}$), η ($= \frac{u\bar{u}+d\bar{d}-2s\bar{s}}{\sqrt{6}}$) and η' ($= \frac{u\bar{u}+d\bar{d}+s\bar{s}}{\sqrt{3}}$) mesons must read, in the basis $(u\bar{u}, d\bar{d}, s\bar{s})$,

$$\mathcal{M}^2 = \begin{pmatrix} A & 0 & 0 \\ 0 & B & 0 \\ 0 & 0 & C \end{pmatrix}. \quad (2)$$

Non-diagonal entries vanish because they originate from OZI violating transitions.

I can now take the reasonable limit $m_u = m_d = 0$, $m_s \neq 0$. This makes things simpler. After all, we know that keeping m_u, m_d different from zero, as in eq. (1), does not solve the problem. In this case there exists a further $SU(2)_A$ symmetry rotating the up and down quarks under which the π^0 gets shifted by a constant amount (as a good Goldstone boson). This further forbids any mass term and hence A and B in eq. (2) have to vanish

in this limit.

Including now the anomaly as the lowest-order contribution in $1/N_c$, one obtains

$$M^2 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & C \end{pmatrix} + \frac{a}{N_c} \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}, \quad (3)$$

where a parametrises the strength of the anomaly. This mass matrix is exactly of the same form as that obtained by Georgi in ref. [1] and therefore leads to the same mass ratio:

$$\frac{M_\pi^2}{M_\eta^2} = \frac{3 + R - \sqrt{9 - 2R + R^2}}{3 + R + \sqrt{9 - 2R + R^2}}, \quad (4)$$

with $R \equiv CN_c/a$, which is maximized for $R = 3$ so that

$$\frac{M_\pi^2}{M_\eta^2} \leq \frac{3 - \sqrt{3}}{3 + \sqrt{3}}, \quad (5)$$

as in ref. [1]. However, from this derivation we see that this result is valid to all orders in the strange quark mass and leading order in $1/N_c$.

This discussion tells us that corrections of order m_s^2 will not affect Georgi's bound and may be consequently disregarded. One can also convince oneself that corrections of order $1/N_c^2$ can only come from the same diagrams that originate from a nonvanishing anomaly (i.e. second term in eq. (3)) but with higher topological complexity. Therefore they only redefine the value of a and will not affect Georgi's bound either. Only the mixed $O(m_s/N_c)$ can do this, and to these operators I now turn.

The quadratic part of the Lagrangian describing the $\eta - \eta'$ system to order m_s/N_c may be obtained from [6]

$$\begin{aligned} \mathcal{L} &= \mathcal{L}_0 + \delta\mathcal{L} \\ \mathcal{L}_0 &= \frac{f_\pi^2}{4} \left[\text{Tr} \partial_\mu U \partial^\mu U^\dagger + \text{Tr} (\chi U^\dagger + U \chi^\dagger) \right] \\ &\quad + \frac{a}{4N_c} (\text{Tr} \log U - \text{Tr} \log U^\dagger)^2, \\ \delta\mathcal{L} &= \frac{f_\pi^2}{4} \left[\frac{2a}{3N_c} \text{Tr} U^\dagger \partial_\mu U \text{Tr} U^\dagger \partial^\mu U \right. \\ &\quad + \frac{\epsilon}{2\sqrt{2}N_c} (\text{Tr} \log U - \text{Tr} \log U^\dagger) \\ &\quad \times \left. \text{Tr} (\chi U^\dagger - \chi^\dagger U) \right] + \dots, \end{aligned} \quad (6)$$

where, following the conventions of ref. [7], one has $U = \exp \left(\frac{-i\sqrt{3}\pi(x)}{f_\pi} \right)$ with $\Phi(x) = \frac{x^2}{\sqrt{3}} + \frac{\lambda}{\sqrt{3}}x$ and λ are the eight Gell-Mann matrices, $f_\pi \simeq 93$ MeV is the pion decay constant; $\chi = 2B_0 M$ where B_0 is a parameter related to the quark condensate in QCD and M is the quark mass matrix. In this Lagrangian a , α and ϵ are parameters of $O(N_c^0)$. Then the first two terms in $\mathcal{L}_0$ are of $O(p^2 N_c^0)$ and the term proportional to a is of $O(a N_c^{-1})$. The Lagrangian $\mathcal{L}_0$ is the leading order and $\delta\mathcal{L}$ can be considered as the $1/N_c$ correction.

Using this Lagrangian, it is very easy to compute the mass matrix in the (η, η') basis. Neglecting again terms proportional to the up and down quark masses but not to the strange quark mass, one finds

$$M^2 = \frac{4}{3} M_K^2 \begin{pmatrix} 1 & -\frac{\sqrt{3}}{2} \\ -\frac{\sqrt{3}}{2} & 2 + z \end{pmatrix}, \quad (7)$$

where

$$\begin{aligned} y &\equiv 1 + \delta y = \frac{1 - \frac{3\epsilon}{N_c \sqrt{2}}}{\sqrt{1 - \frac{2\epsilon}{N_c}}} \\ &\approx 1 - \frac{1}{N_c} \left(\frac{3\epsilon}{\sqrt{2}} - \alpha \right) + O\left(\frac{1}{N_c^2}\right), \end{aligned} \quad (8)$$

$$\begin{aligned} z &\equiv \frac{9a}{4M_K^2 N_c (1 - \frac{2\epsilon}{N_c})} \\ &\approx \frac{9a}{4M_K^2 N_c} \left(1 + \frac{2\epsilon}{N_c} + O\left(\frac{1}{N_c^2}\right) \right). \end{aligned} \quad (9)$$

In the spirit of the $1/N_c$ expansion one should take δy as a small parameter and expand in it. The η and η' masses are then determined by the conditions

$$\frac{3(M_\pi^2 + M_\eta^2)}{4M_K^2} = \frac{3}{2} + z + \delta y, \quad (10)$$

$$\frac{9M_\pi^2 M_\eta^2}{16M_K^2} = z. \quad (11)$$

These equations yield $\delta y \simeq -0.35$ and $z \simeq 2.57$ when the masses $M_K \simeq 495$ MeV, $M_\eta \simeq 547$

I would like to thank Stephan Narison for his kind invitation to this workshop.

REFERENCES

1. H. Georgi, Phys. Rev. D49 (1994) 1686. A similar observation has been made independently by J. Schechter, A. Subbaraman and H. Weigel, Phys. Rev. D48 (1993) 339.
2. G. 't Hooft, Nucl. Phys. B72 (1974) 461.
3. E. Witten, Nucl. Phys. B156 (1979) 269; G. Veneziano, Nucl. Phys. B159 (1979) 213; P. Di Vecchia and G. Veneziano, Nucl. Phys. B171 (1980) 253; E. Witten, Ann. of Phys. 128 (1980) 363; C. Rosenzweig, J. Schechter and G.G. Trahern, Phys. Rev. D21 (1980) 3368.
4. Particle Data Group, Phys. Rev. D45 (1992).
5. S. Peris, Phys. Lett. B324 (1994) 442.
6. P. Di Vecchia, F. Nicodemi, R. Pettorino and G. Veneziano, Nucl. Phys. B181 (1981) 318.
7. A. Pich and E. de Rafael, Nucl. Phys. B367 (1991) 313.

MeV, $M_{\eta'} \simeq 958$ MeV are used. It is clear that the system (10) + (11) has always one solution for z and δy , once the masses for the pseudoscalars are given; the mass matrix (7) can thus fit the η and η' masses, and the bound of ref. [1] is overcome. As a matter of fact, taking the mass matrix (7) one easily obtains

$$\frac{M_\pi^2}{M_{\eta'}^2} = \frac{1 + z + y^2/2 - \sqrt{(1 + z + y^2/2)^2 - 4z}}{1 + z + y^2/2 + \sqrt{(1 + z + y^2/2)^2 - 4z}}. \quad (12)$$

This expression has a maximum when varied with respect to z ($\sim$ the anomaly to quark mass ratio), keeping y fixed. One can understand this on physical grounds: for $z \rightarrow \infty$ the mass ratio (12) goes to zero because the η' mass becomes infinite. Furthermore, when $z \rightarrow 0$ the mass ratio also goes to zero because the η mass vanishes since the situation of the $U(1)_A$ problem is reproduced. So there must be an intermediate value of z at which eq. (12) has a maximum. Moreover, in this eq. (12) y equals unity only when next-to-leading terms in $1/N_c$ are neglected. In this case one can see that eq. 12 reaches its maximum at $z_0 = 3/2$ and one then obtains the bound (5). However, if next-to-leading terms are included one has instead that $y \approx 1 + \delta y$ with $|\delta y| \sim 1/N_c \ll 1$. Expanding eq. (12) in δy about z_0 one obtains

$$\frac{M_\pi^2}{M_{\eta'}^2} \leq \frac{3 - \sqrt{3}}{3 + \sqrt{3}} (1 - \frac{2}{\sqrt{3}} \delta y). \quad (13)$$

A negative value for δy of order $1/N_c \sim 0.3$, which is the natural size, is more than enough for eq. (13) to be satisfied experimentally.

In summary, I have shown that contributions of $O(m_s/N_c)$ are the ones capable of reconciling Georgi's bound with the experimental masses. At least, there is no problem already at this level. However there are lots of things that can still go wrong before we can treat, in practice, the η' as a good Goldstone boson as, let's say, the kaon. This is because the η' mass, although formally small as $1/N_c$, is numerically very large. Further studies are still clearly needed.