JS

CERN LIBRARIES, GENEVA

P00021279

ICRR-Report-305-93-17
KEK Preprint 93-176
TKU-PAP-93-02
KOB-AP-93-02
NGTHEP-93-02
OULNS-93-06
TIT-HPE-93-09
GIFU-PH-93-01
INS-Rep.-1018
UPR-0223E

SEARCH FOR LOW ENERGY NEUTRINOS FROM GALACTIC GAMMA RAY SOURCES

Y. Fukuda, T. Hayakawa, K. Inoue, T. Ishida, T. Kajita, Y. Koshio, M. Nakahata,
K. Nakamura, A. Sakai, M. Shiozawa, J. Suzuki, Y. Suzuki, Y. Totsuka, M. Mori,
K. S. Hirata, K. Kihara, Y. Oyama, A. Suzuki, M. Yamada, M. Koshihara, K. Nishijima,
T. Kajimura, T. Suda, A. T. Suzuki, H. Takei, M. Koga, K. Miyano, H. Miyata, T. Hara,
N. Kishi, Y. Nagashima, M. Takita, A. Yoshimoto, K. Kaneyuki, Y. Takeuchi,
T. Tanimori, S. Tasaka, K. Nishikawa, E. W. Beier, E. D. Frank, W. Frati, S. B. Kim,
A. K. Mann, F. M. Newcomer, R. Van Berg, and W. Zhang

(December 1993)

Submitted to the Astrophysical Journal

3-2-1 Midori-cho Tanashi, Tokyo 188 Japan

Telephone (0424)-61-4131, Telefax (0424)-68-1438

SEARCH FOR LOW ENERGY NEUTRINOS FROM GALACTIC GAMMA RAY SOURCES

Y. FUKUDA, T. HAYAKAWA, K. INOUE, T. ISHIDA, T. KAJITA, Y. KOSHIO, M.
NAKAHATA, K. NAKAMURA, A. SAKAI, M. SHIOZAWA, J. SUZUKI, Y. SUZUKI, AND
Y. TOTSUKA

Institute for Cosmic Ray Research, University of Tokyo, Tanashi, Tokyo 188, Japan

M. MORI¹, K. S. HIRATA, K. KIHARA, Y. OYAMA, A. SUZUKI², AND M. YAMADA³
National Laboratory for High Energy Physics (KEK), Tsukuba, Ibaraki 305, Japan

M. KOSHIBA AND K. NISHIJIMA
Tokai University, Shibuya, Tokyo 151, Japan

T. KAJIMURA⁴, T. SUDA⁵, AND A. T. SUZUKI
Department of Physics, Kobe University, Kobe, Hyogo 657, Japan

H. TAKEI⁶, M. KOGA, K. MIYANO, AND H. MIYATA
Niigata University, Niigata 950-21, Japan

T. HARA, N. KISHI, Y. NAGASHIMA, M. TAKITA, AND A. YOSHIMOTO⁷
Department of Physics, Osaka University, Toyonaka, Osaka 560, Japan

K. KANEYUKI, Y. TAKEUCHI, AND T. TANIMORI
Department of Physics, Tokyo Institute of Technology, Meguro-ku, Tokyo 152, Japan

S. TASAKA
Department of Physics, Gifu University, Gifu, Gifu 501-11, Japan

K. NISHIKAWA
Institute for Nuclear Study, University of Tokyo, Tanashi, Tokyo 188, Japan

AND

E. W. BEIER, E. D. FRANK, W. FRATI, S. B. KIM⁸, A. K. MANN, F. M.
NEWCOMER, R. VAN BERG, AND W. ZHANG⁹
Department of Physics, University of Pennsylvania, Philadelphia PA 19104, U. S. A.

¹Now at Department of Physics, Miyagi University of Education, Sendai 980, Japan.

²Now at Bubble Chamber Physics Laboratory, Tohoku University, Sendai 980, Japan.

³Now at Niigata Polytechnic College, Niigata, Japan.

⁴Now at Hydrographic Department, Maritime Safety Agency, Chuô-ku, Tokyo 104, Japan.

⁵Deceased.

⁶Now at Oarai Engineering Center, Power Reactor & Nuclear Fuel Development Corporation, Oarai, Ibaraki 311-13, Japan.

⁷Now at Public Systems Development Division, NEC Corporation, Minato-ku, Tokyo 105, Japan.

⁸Now at the University of Michigan, Ann Arbor, MI 48109, U.S.A.

⁹Now at NASA, Goddard Space Flight Center, Greenbelt, MD 20771, U.S.A.

ABSTRACT

Data from the Kamiokande detector are used to set upper limits on fluxes of low energy neutrinos from possible nearby sources within or near our galaxy. The search was conducted for 1557 live detector days from January 1987 to February 1993.

1. Introduction

Since the publication of the directional observation in real-time of neutrinos from the Sun by the Kamiokande-II collaboration (Hirata *et al.* 1991), questions relating to searches for other stellar sources of low energy ($\lesssim 30$ MeV) neutrinos have been asked of us. The Sun is roughly 10^5 times closer to the Earth than the next nearest stellar objects, and consequently a more distant source is required to exhibit extraordinary intensity if a neutrino signal from it is to be observed. Nevertheless, it may be useful to report the results of one such search, as we do here, to illustrate the generality of the method on which the Kamiokande-II and -III solar neutrino measurement is based, and because discoveries of very bright nearby objects (at least at certain electromagnetic wavelengths) continue to be made, *e.g.*, PSR J0437-4715 (Johnston *et al.* 1993).

2. Kamiokande Detector

The Kamiokande detector is located 1000 m underground (2700 mwe) in Gifu Prefecture in western Japan. The detector is a cylindrical (14.4 m ϕ , 13.1 m height) imaging water Cherenkov detector containing 2.14 metric kilotons of purified water, viewed by 948 (50 cm diameter) photomultiplier tubes (PMT) covering 20 % of the detector inner surface area. An optically separated 4π solid-angle water Cherenkov counter surrounds the detector and serves as both a passive and active shield against background events.

The Kamiokande-II solar neutrino observation was started in January 1987 and extended to April 1990, with 1040 detector live days. Further details of the Kamiokande-II detector are found in Hirata *et al.* (1991, 1988). The new phase of the experiment, Kamiokande-III, was started in December 1990, after replacement of the entire electronics, replacement of the dead PMTs, and installation of a reflecting cone on each PMT. The description of the Kamiokande-III detector is found in Nakamura (1993) and Inoue (1993).

Low energy neutrinos are detected by their elastic scattering off atomic electrons. The Cherenkov radiation from the recoiling electrons is observed by the PMT. For 10 MeV electrons typically, knowing the position of the individual PMT and using their 7 nanosecond (rms) timing resolution, the vertex of a neutrino-electron interaction can be located to 1.0 $\sim$ 1.2 m (rms), and the direction of the electron relative to the nominal incident neutrino direction can be determined to 27 degrees (rms), which is set principally by multiple Coulomb scattering. (In the absence of Coulomb interactions, $\theta_{e,\nu}^2 \leq 2m_e/E_e$.) To illustrate the angular resolution a comparison is shown in Figure 1 of data obtained with a collimated 9 MeV calibration γ -ray source with a Monte Carlo simulation of the electron angular distribution produced by that source (Ni-Cf).

Note that in searching for a signal from the Sun, the real-time observation of each electron is necessary to specify the radius vector from the sun to the detector for comparison with the observed electron direction. The detector electron energy resolution is $\sigma(E_e) = 0.63\sqrt{E_e(\text{MeV})}$ and, for example, a 10 MeV electron illuminates ≈ 30 PMT in Kamiokande-II and ≈ 40 PMT in Kamiokande-III.

3. Data Reduction and Analysis

The vast majority of events observed in the detector arise from radioactivity in the detector water and the rock surrounding the detector and from the inelastic interactions of cosmic ray muons in and nearby the detector. The number of these background events is reduced by the energy, spatial, and timing criteria used to select the final sample of events. The event selection criteria are the same as those in the solar neutrino analysis described in Hirata *et al.* (1991), except for a few unimportant differences in the reduction of the Kamiokande-III data.

In the present analysis, the threshold for the electron energy was taken to be 8 MeV and the final event samples from Kamiokande-II and -III were combined and binned in intervals of $\Delta(\cos \theta_{e, \text{Source}}) = 0.05$, where $\theta_{e, \text{Source}}$ is the electron angle relative to the direction of the Sun and each of several bright sources in the Galaxy, the Galactic Center, and LMC-X4.

To investigate the angular distribution with respect to bright sources other than the Sun, the solar neutrino signal becomes part of the “background” which exhibits an angular shape peculiar to the celestial position relative to the ecliptic, and symmetric with respect to $\theta_{e, \text{Source}} = 90^\circ$, if the data are accumulated throughout a year. Figure 2 shows the angular distribution with respect to each studied bright source of the Monte Carlo-generated solar neutrino events, where the energy threshold for electrons and live detector days are properly taken into account.

The angular distribution with respect to a point source at a fixed celestial position is, therefore, represented by an isotropic background component B and a solar-neutrino background component B_S , on which is superimposed a signal of strength S from a point source. The component B_S is shown in Figure 2.

Since we do not know the energy spectrum of possible low-energy neutrinos from these bright sources, our basic approach is to determine S assuming monochromatic electron-neutrinos of energy E_ν , and then to study the E_ν -dependence of S. For this purpose, the angular distribution of the expected signal is determined for $E_\nu = 10, 12, 14$, and 15 MeV using Monte Carlo-generated events. Electron-antineutrinos, if any, are assumed to give an isotropic angular distribution which is indistinguishable from the background. Muon- and tau-neutrinos are not considered, but the result for electron neutrinos only can be easily converted to the result for a given flux ratio of $\nu_e : \nu_\mu : \nu_\tau$ at each E_ν using the known cross section ratio of $\sigma(\nu_e e \rightarrow \nu_e e) : \sigma(\nu_\mu e \rightarrow \nu_\mu e) : \sigma(\nu_\tau e \rightarrow \nu_\tau e) \approx 6 : 1 : 1$, because the signal angular distribution is mainly determined by the angular resolution.

4. Results and Discussion

For the studied bright point sources other than the Sun the quantities B and S are determined using a maximum likelihood method from the best fits of the assumed angular distribution to the data. The numerical results for $E_\nu = 10$ MeV are given in Table 1. Figures 3(a) - 3(i) show the resulting angular distributions with the solar-neutrino component B_S subtracted. The overlaid dashed lines show the numerical results for $E_\nu = 10$ MeV. The 90 % confidence-level ν_e flux upper limits given in column 6 of Table 1 were calculated from the respective likelihood function by limiting the domain of the variable S to be the physical region, $S \geq 0$. The E_ν -dependence of the results is graphically summarized in Figure 4.

For comparison, Figure 3(j) shows the angular distribution with respect to the Sun for $E_e \geq 8$ MeV. The solid line shows the result of a maximum likelihood analysis. For the case of the Sun, as was described in Hirata *et al.* (1991), it was assumed that the distribution is represented by an isotropic component due to the background B on which is superimposed a signal of strength S from the Sun. The expected electron angular distribution is obtained by integrating above the analysis threshold the product of the ^8B solar neutrino energy spectrum and the energy dependent elastic scattering cross section. In Table 1 the numerical result for the Sun with $E_e \geq 8$ MeV is also given. As expected, the Sun is the only source exhibiting a statistically significant signal.

5. Conclusions

The data presented in this paper lead to the two following conclusions.

First, the upper limits on neutrino fluxes from different directions in the sky given in column 6 of Table 1, and indicated by right ascensions (RA) and declinations in columns 2 and 3, when compared with the positive result from the Sun, represent further validation of the method used by the Kamiokande-II collaboration to extract the published solar neutrino signal (Hirata *et al.* 1991).

Second, the upper limits themselves are of some intrinsic interest. For example, the upper limit on the recently discovered pulsar PSR J0437-4715, a very bright nearby binary pulsar with a low mass companion (Johnston *et al.* 1993), indicates that significant steady or rapidly pulsed emission of low energy neutrinos does not occur from that relatively bright source of electromagnetic radiation.

TABLE 1

SUMMARY OF UPPER LIMITS ON ELECTRON-NEUTRINO FLUXES AT $E_\nu = 10$ MEV
INCIDENT ON THE EARTH FROM THE INDICATED SOURCES OTHER THAN THE SUN

Source	R. A. (deg)	Decl. (deg)	Dist. (kpc)	Events	90% C.L. ν_e flux upper limit ($\text{cm}^{-2} \text{sec}^{-1}$)	90% C.L. ν_e luminosity upper limit (erg sec^{-1})
Geminga	97.7	17.8	0.1 ^(a)	-1_{-21}^{+22}	1.0×10^5	1.9×10^{42}
PSR J0437 -4715	69.32	-47.25	0.15	-12_{-19}^{+20}	8.0×10^4	3.4×10^{42}
Crab	82.9	22.0	2	-17_{-20}^{+22}	7.9×10^4	6.1×10^{44}
Vela Pulsar	128.4	-45	0.5	-4_{-18}^{+20}	9.7×10^4	4.6×10^{43}
SS433	285.1	4.8	3	-25_{-17}^{+19}	6.0×10^4	1.0×10^{45}
Her X-1	254.0	35.4	5	2_{-19}^{+21}	1.0×10^5	4.8×10^{45}
Cyg X-3	307.6	40.8	12	20_{-19}^{+21}	1.4×10^5	3.9×10^{46}
Gal Ctr	265.5	-28.6	10	-40_{-17}^{+19}	4.9×10^4	9.4×10^{45}
LMC X-4	81.0	-69.5	50	-23_{-18}^{+19}	6.4×10^4	3.1×10^{47}
Sun	—	—	5×10^{-9}	$358_{-28}^{+29(b)}$	—	—

^(a)See Bignami, Caraveo, and Mereghetti 1993.

^(b)The ^8B solar neutrino flux obtained from this result is statistically consistent with the published result by Kamiokande-II (Hirata *et al.* 1991), 0.46 ± 0.05 (stat.) ± 0.06 (syst.) times the standard solar model prediction (Bahcall and Ulrich 1988).

REFERENCES

- Bahcall, J. N. and Ulrich, R. K. 1988, *Rev. Mod. Phys.*, **60**, 297.
Bignami, G. F., Caraveo, P. A., and Mereghetti, S. 1993, *Nature*, **361**, 704.
Hirata, K. S., *et al.* 1988, *Phys. Rev. D*, **38**, 448.
———. 1991, *Phys. Rev. D*, **44**, 2241.
Inoue, K. 1993, in *Proc. 28th Rencontres de Moriond “Electroweak Interactions and Unified Theories”*, in press.
Johnston, S., *et al.* 1993, *Nature* **361**, 613.
Nakamura, K. 1993, *Nucl. Phys. B (Proc. Suppl.)* **31**, 105.

FIGURE CAPTIONS

Fig. 1. — Measured and calculated electron angular distribution produced by a collimated 9 MeV γ -ray source (Ni-Cf) in Kamiokande-II from Hirata *et al.* (1991).

Fig. 2. — The angular distributions of the Monte Carlo-generated solar neutrino events with respect to the directions from various bright sources to the Earth. These Monte Carlo events correspond to the data shown in Figure 3(j), *i.e.*, the energy threshold for electrons (8 MeV) and live detector days are properly taken into account.

Fig. 3. — (a) - (i) The data points indicated by the circles show the angular distributions from which the fitted solar-neutrino component is subtracted. The numerical results indicated by the dashed lines correspond to a superposition of a flat background and a possible source signal for monochromatic electron-neutrinos of $E_\nu = 10$ MeV. (j) The electron angular distribution for $E_e \geq 8$ MeV (combined data from Kamiokande-II and -III) with respect to the direction from the Sun to the Earth. The solid line is the best fit to a flat background plus a ^8B solar-neutrino signal.

Fig. 4. — Summary of the 90 % confidence level upper limits of the ν_e flux incident on the Earth from the directions of the various indicated sources, as a function of assumed monochromatic neutrino energy E_ν .

FIG.1

$\cos \theta_{e, Source}$

FIG.2

FIG.3(a)

FIG.3(b)

FIG.3(c)

FIG. 3(d)

FIG.3(e)

FIG. 3(f)

FIG.3(g)

FIG.3(h)

FIG. 3(i)

FIG.3(j)

FIG.4