AC

M/C.TH.93/16 (revised)

DO-TH.93/13

MC-TH 93-16 rev
Sc 94 10

Large p_T Prompt Photon Production at HERA

L.E.Gordon

February, 1994

Institut für Physik,

Universität Dortmund,

44221 Dortmund, Germany

J.K.Storow

Department of Theoretical Physics,

The University, Manchester M13 9PL, England.

Abstract

We examine the possibility of using direct photon production at HERA as a means of obtaining information about the photon structure and fragmentation functions. Estimates of the cross section for inclusive prompt photon production at the ep collider HERA are made completely in next-to-leading order QCD. Using parton distributions for the photon evolved in next-to-leading order QCD for the first time, we show that the $O(\alpha_s)$ corrections to the processes involving these functions are

significant in the regions where they dominate. This can have important consequences for the possibility of measuring the gluon content of the photon using these processes at HERA. Moreover, we show that it may not be straightforward to separate the various contributions to the cross section in the ep lab frame.

1 Introduction

In addition to providing other useful tests of perturbative QCD, large pt prompt photon production at the HERA ep collider offers the possibility for measurement of the quark and gluon structure of the real photon, and could yield further information about the gluon structure of the proton [1]. Perturbative QCD predicts that the photon possesses a quark and gluon structure [2], but unlike the case of the quark structure, which has been identified and measured in two photon processes at e^-e^+ colliders, the gluon structure has only very recently been confirmed to exist experimentally [3]. It is widely accepted that further more precise measurements are necessary and that HERA offers an excellent possibility of such measurements via various photoproduction processes. Many different processes, such as single and two jet production [4, 5], heavy quark, J/ψ and lepton pair production [6-9] have been proposed as a means for measuring g^γ . However except for heavy quark and lepton pair production which suffer from very low event rates, and single jet production [10-13], all have the disadvantage of being calculated to leading log accuracy only. The associated large uncertainties in the predictions due to factorisation/renormalisation scale sensitivity and the uncertainty about the size of the higher order corrections to some of the contributing processes mean that exact quantitative conclusions cannot be drawn from such calculations.

In the case of inclusive prompt photon production it is possible to perform a complete calculation in next-to-leading order (NLO) QCD because all the relevant matrix elements are available [14-17]. In addition, and for the first time, it is now possible to use fully regularised parton distributions for the photon evolved in NLO [18,19], and there are many different sets of parton distributions for the proton so that sensitivity to g^p can also be checked. This is important since it is known that deep inelastic scattering does not provide a tight constraint on g^p since it only en-

ters indirectly in determining the evolved structure functions through the (weak) coupling of the Altarelli-Parisi equations for the singlet quark and gluon sectors. Prompt photon production at HERA, on the other hand, offers the possibility of directly measuring both g^p and g^γ at x values as low as 10^{-3} . It has also been suggested that it may for the first time provide measurement of the photon fragmentation functions [20] which are very important for the extraction of information on g^p from prompt photon production hadron-hadron scattering, although in practical situations this importance is somewhat reduced due to the need to perform isolation cuts in order to identify the photon signal [21].

In analogy with jet photoproduction, we consider two classes of contributions to the photoproduction of direct photons at HERA: the direct contribution where the photon interacts directly with the partons in the proton through its electromagnetic interaction, and the resolved contributions where the photon interacts through its quark and gluon structure [4,22]. It has been demonstrated in leading order (LO) calculations for both jets and prompt photons [4,5,20,23], and in NLO for jets [10-13] that there are kinematic regions where either class of process may dominate, and thus be separated from the other. It was also shown that particular subprocesses which may be of special interest may be isolated and studied. This is, in fact, what provides the possibility of obtaining information on both the photon and proton distribution functions, and particularly the gluon distributions.

We note that in a complete NLO calculation this type of separation is no longer justified since the factorisation procedure introduces a dependence on the scale, Q^2 , at which the photon structure function is evaluated, into the 'direct' contributions [13,24]. We nevertheless, in the interest of continuity, still keep the two types of processes separate and set the factorisation scale, Q^2 , in the direct contribution equal to the scale at which we evaluate the photon structure function in the resolved

contribution. This avoids the possibility of scale mismatching which could lead to an unreliable estimate of the cross section when the two contributions are added together. It also leads to reasonably small higher order corrections to the direct contributions, as long as Q^2 is chosen in the region $(\tilde{p}_T^2)^2/4 \leq Q^2 \leq 4(\tilde{p}_T^2)^2$, where $\tilde{p}_T^2$ is the transverse momentum of the observed final state photon. Strictly speaking, only the sum of the two contributions has physical significance: evidence for this is provided by the fact that the direct contribution can have very large corrections for certain choices of the scale, as is found in the case of jet production [13,24].

The original studies which examined the possibility of measuring the parton content of the photon using inclusive prompt photon production at HERA relied on three main assumptions [1,20]:

- (a) the higher order $O(\alpha_s)$ corrections to the resolved processes are small in the regions where these contributions dominate,
- (b) the important processes can be separated in rapidity in the ep lab frame,
- (c) and if (b) does not hold, then the various final states can be distinguished from each other by their different topologies, making it possible to identify the individual contributing subprocesses.

Assumption (a) has been checked recently at HERA energies by Aurenche et al [25] though they work at a fixed γp energy, which we will see is not realistic for HERA. We will be taking their work one step further by working with a realistic spectrum of initial state photons. Another problem has been the lack of reliable parton distributions for the photon in NLO and we will also be examining the sensitivity of the results to different parametrisations of both photon and proton distribution functions. Assumption (b) has not been checked in any detail which takes into account the practical limitations on tagging of the scattered electrons at HERA, and it seems to us that assumption (c) depends to some extent on (a) being

true. It is difficult to see how one may extract information on parton distributions using processes which may suffer from significant higher order corrections without evaluating them.

To summarise, in this paper we check assumptions (a) and (b) in some detail, taking particular care over the effects of the spectrum of initial state photons. We also examine the effects of different choices of NLO distribution functions of both photon and proton.

We present our results in two parts; in section 2 we present the results of a consistent LO analysis in which we explore the issue of separation in rapidity of the contributions in the lab frame. Then in section 3 we present the full NLO analysis. In section 4 we draw our conclusions.

2 The ep Laboratory Frame: Electroproduction vs Photoproduction

At HERA, photoproduction processes will be initiated by the interaction of almost on shell photons emitted by a 30 GeV electron beam and 820 GeV protons. It has been suggested that performing the calculation in the asymmetrical ep lab frame rather than a fixed γp cm frame with an estimated cm energy, usually taken as $S_{\gamma p} \approx 30000 \text{ GeV}^2$ [15, 20], will aid in the separation in rapidity of the various subprocess contributing to the cross section. This should aid in the direct measurement of those that involve g^+ and g^0 . Calculation of electroproduction in the ep lab frame is usually performed by making use of the Weizsacker-Williams or Equivalent Photon Approximation (EPA) to estimate the flux of quasi-real photon emitted by the electron beam [26]. This leads to a convolution of the parton distribution functions of the photon $\hat{q}_i^{\gamma}(x, Q^2)$, with the 'electron distribution function' $f_{e/\gamma}(x)$ to produce an effective electron distribution function. Thus the distribution of parton i in the

electron is given by

$$f_i(x_e, Q^2) = \int_{x_e}^1 \frac{dz}{z} f_{e\gamma}(z) q_i^2(z/x_e, Q^2), \quad (1)$$

The specific form of the function $f_{e\gamma}(x)$ depends on whether the scattered electron is tagged or not. It has been shown that the use of this approximation does not introduce more than a $\pm 10\%$ error in the estimated cross section [4,27] for some specific processes, and thus should be preferred over a fixed γp cm energy since it more realistically reproduces the spread in the initial state photon energies.

It can also be argued that use of this approximation in the forms given in refs. [4] and [27] leads to the inclusion of photons with higher virtuality than can justify the use of the real photon structure function. The only practical alternative to using the approximations given in these references is to perform very precise tagging of the scattered electron. But as has been pointed out in ref. [28], tagging will only be possible in a certain (restricted) angular range, still leading to some spread in the initial photon energies, but with the added disadvantage that the observable cross section can fall by a factor of 10 – 1000, depending on the tagging parameters. If the scattered electrons are tagged very precisely it should be expected that the cross section will fall by even more than the estimates given in ref. [28], leading to poor statistics and reducing the usefulness of the measurements. This conclusion can be justified by the already relatively small sizes of the cross sections when compared to that of, for example, inclusive jet production [10-13,20]. We will examine how the separation of the processes in the ep lab frame is affected by the spectrum of initial state photon energies.

For this part of the calculation we make use of the first (LO) set of photon distribution functions of ref. [18], the GS(1) set. For the proton distributions we use those of ref. [29], called the MT(SL) set. Positive rapidity is taken in the direction of the electron beam, and unless otherwise stated we use $Q^2 = (p_T^*)^2$ as the scale for

the hard scattering, and take the electron structure function as

$$f_{\gamma/e}(x) = \frac{\alpha_{em}}{2\pi} \left[\frac{1 + (1-x)^2}{x} \right] \left(\ln \frac{\bar{E}^2}{m_e^2} \right) \quad (2)$$

where m_e is the electron mass and $\bar{E}$ is taken as the electron beam energy. We note that other choices are possible for $\bar{E}$ and the cross section can be sensitive to the choice made.

In LO the processes that contribute to the direct component are

$$\gamma + q \rightarrow \gamma + q \quad (3)$$

and

$$\begin{aligned} \gamma + q &\rightarrow q + g \\ \gamma + g &\rightarrow q + \bar{q} \end{aligned} \quad (4)$$

where the latter involve the photon fragmentation functions. We use the asymptotic parametrisations of ref. [30] throughout the paper. The resolved processes are

$$\begin{aligned} \bar{q} + q &\rightarrow \gamma + g \\ q + g &\rightarrow \gamma + q \end{aligned} \quad (5)$$

and

$$\begin{aligned} q + q' &\rightarrow q + q' \\ q + q &\rightarrow q + q \\ q + \bar{q} &\rightarrow q' + \bar{q}' \\ q + \bar{q} &\rightarrow q + \bar{q} \\ q + \bar{q} &\rightarrow g + g \\ g + g &\rightarrow q + \bar{q} \\ q + g &\rightarrow q + g \\ q + g &\rightarrow g + g. \end{aligned} \quad (6)$$

where the latter involve fragmentation processes. The matrix elements for the processes in eqs. (4-6) were taken from refs. [30] and [31] respectively.

In this section we present results for the processes given in eqs. (3) and (5) only after checking that the processes (4) and (6) give smaller contributions, particularly for the photoproduction processes, probably because of the convolution with the fragmentation functions. Also, our aim is to study how the EPA affects the rapidity distributions and similar conclusions can be drawn for the other processes. We first performed the calculation in the lab frame for a no tag situation and then for the 'golden tag' situation suggested in ref. [28]. The shape of the rapidity distributions did not change significantly, but we confirmed the result that the cross section was significantly reduced (by a factor of 10 in this case) when tagging is performed. We then performed the calculation in the ep lab frame but with a fixed energy of 9 GeV for the photon, and then in the fixed γp cm frame with fixed cm energy $S_{\gamma p} = 30000$ GeV². This is equivalent to a 9 GeV fixed photon beam energy. As may be expected the only difference between the two rapidity distributions was a shifting toward negative rapidity (i.e. the proton direction) in the case of the lab frame calculation.

In figs. 1 and 2 we will present results for the two extreme cases of a fixed γp cm energy, and a no tag situation in electroproduction in the ep lab frame, where the full EPA approximation is used. As stated earlier, we choose $\bar{E}$ as the electron beam energy in the EPA formula. Fig. 1(a) shows clearly that the minimum values of x_p , the Bjorken variable for the proton, probed as a function of the rapidity of the observed final state photon are different in the fixed γp and the ep lab frames. As expected, the whole distribution is shifted in rapidity, but more significantly, the value of $x_p^{\min}$ is about ten times smaller in the ep frame. This will have important consequences for the shape of the rapidity distributions since these depend on the

shape of the parton distributions with respect to x_p , and different x_p values are probed in each case. Similar conclusions can be drawn about x_e , the corresponding variable for the electron, and hence for x_γ , that for the photon.

In fig. 1(b) we show the u-quark distribution at $Q^2 = 100$ GeV² plotted as function of x , before and after convolution with the Weizsäcker-Williams function. As expected, the distribution is much smaller after the convolution and the characteristic large distribution at large x is lost.

Figs. 2(a) and 2(b) show the rapidity distributions for the various subprocesses in the γp and ep lab frames respectively. It is clear that there are very significant differences between the two. The most significant change is in the basic process of eq. (3), which had a sharp peak at positive rapidity in the γp cm frame, but does not have a well defined maximum in the lab frame. In the γp frame the two quark-gluon scattering processes $g^*q' \rightarrow \gamma q$ and $g^*q' \rightarrow \gamma q$, where the first involves a gluon from the photon and a quark from the proton and the second the opposite combination, are more or less separated from each other, the first dominating the negative and the second the positive rapidity regions. This separation, although not altogether lost, is significantly less pronounced when the calculation is performed in the ep lab frame. We are led to the conclusion that when the calculation is performed in the ep lab frame, it is more difficult to disentangle the various subprocess contributions to inclusive prompt photon production at HERA.

One could ask whether some sort of restricted tagging procedure, not as severe as the 'golden tag' procedure[28], might preserve the separation characteristic of a pure photoproduction process. The 'poor-tag' situation, also outlined in ref [28], is just such a procedure. In fig. 2(c), we compare the rapidity distributions for the basic direct process $\gamma q \rightarrow \gamma q$ (eq. 3) for all three scenarios. Compared to fig. 2(b), there is very little difference between the three curves apart from the obvious

significant differences in the sizes of the cross sections. The situation is very similar for the resolved processes. Thus it appears that mere tagging restrictions will not aid significantly in separating resolved and direct contributions to the production of prompt photons at HERA. Other methods would have to be used to achieve this.

3 Cross Sections

In this section we present the result of our full NLO calculation of the inclusive prompt photon cross section at HERA. We use the NLO parametrisation of the photon distribution functions of refs. [18] and [19], the GS(1)' and GRV parametrisations respectively, and the proton distributions of refs. [29], [32] and [33] which we refer to as MT(S), GRV' and KMRS(B_0), respectively. Unless otherwise stated results are quoted for the GS(1)' photon and the MT(S) proton. Our standard choice of renormalisation/factorisation scales are again taken as $Q^2 = (p_T^2)^2$. The matrix elements calculated by Aureuche *et al.* in refs. [14] and [15] as well as those calculated recently by Gordon and Vogelsang [16] are used for the resolved and direct non-fragmentation processes respectively. These matrix elements have been compared and found to agree except for small differences due to different conventions used for initial gluon spin averaging. We use both for convenience of programming only. We refer to refs. [14], [15] and [20] for a list of the additional subprocesses that have to be included in the full NLO calculation. The method of dealing with the soft and collinear singularities that arise from the higher order processes is well documented in refs. [14-16], so we do not repeat the discussion here.

3.1 p_T^2 Distribution

We first look at the p_T^2 distribution of the cross section integrated over all rapidities, $d\sigma/dp_T^2$. Throughout this subsection the matrix elements of refs. [14,15] are used.

Fig. 3(a-b) show the ratio, $R(p_T^2)$, of the LO to the full NLO processes for the resolved and direct contributions to the cross section. As explained in the introduction we keep resolved and direct contributions separate. The ratio, in the case of the direct processes, is defined by

$$R(p_T^2) = \frac{\frac{d\sigma}{dp_T}(\gamma q \rightarrow \gamma q)}{\frac{d\sigma}{dp_T}} \quad (7)$$

where the numerator is the full NLO direct cross section (which includes both the LO contributions and the NLO corrections to them) and the denominator is the basic lowest order contribution of eq. (3). We follow ref. [20] and include the $O(\alpha_s^2\alpha_{em}^2)$ 'box' diagram contribution, ($\gamma q \rightarrow \gamma q$), along with the $O(\alpha_s)$ corrections to the basic γq scattering process of eq. (3). The fragmentation contributions (eq. (4)) contribute in LO and should be included in the denominator of eq. (7) for the calculation of a proper K-factor. The situation is similar in the case of the resolved contributions and we discuss this in more detail below.

In the case of the resolved processes we may define the ratio relative to the two LO processes of eq. (5), plus the fragmentation processes of eq. (6) which also contribute to the same order in the strong and electromagnetic couplings, i.e. to $O(\alpha_s\alpha_{em})$. There is some ambiguity in our precise definition of the higher order corrections for the resolved processes. In refs. [14] and [34] the full NLO cross section was written as

$$\frac{d\sigma}{dp_T dy} = \frac{d\sigma^{LO}}{dp_T dy} + \frac{d\sigma^{HO}}{dp_T dy} + \frac{d\sigma^{A,V}}{dp_T dy} \quad (8)$$

where LO represents the leading order processes of eq. (5), HO is defined as the higher order corrections proper and include non-collinear photon bremsstrahlung of the final state quarks, but do not include those processes where the final state photon is collinear with a final state quark. These are correctly included in the fragmentation processes of eqs. (6), which in this case, strictly speaking, should be

convoluted with NLO fragmentation functions. The ambiguity arises because the LO fragmentation functions are $O(\alpha_{em}/\alpha_s)$ in the electromagnetic and strong couplings. When this is convoluted with the $O(\alpha_s^2)$ processes of eqs. (6), then they contribute in the leading order, $O(\alpha_{em}\alpha_s)$, also and thus can be included in the equivalent of the denominator of eq. (7) for the resolved processes. Thus when taking the ratio of the full NLO to the LO cross sections these contributions can be included in both, or only in one or the other. We will thus also take the ratio of the full NLO cross section to the LO processes of eq. (5) only to reflect the attitude of the authors of ref. [21], for example, that the fragmentation contributions contribute in NLO only, and also to be consistent with our definition of the ratio in the direct contributions. We stress that there is no real ambiguity in the definition of the differential cross section, only in our definition of the ratio of the NLO to LO cross sections of the direct and resolved components when we consider them separately.

We also point out that we are in a position to properly compare the leading and higher order predictions for the resolved contribution to the cross section in a consistent way, i.e., by using LO structure functions in the LO calculation which were obtained by fits to the same data sets as the corresponding NLO distributions. These are the GS(I) and GS(IV) distributions for the photon in LO and NLO respectively, and the MT(S) and MT(SL) distributions for the proton. The NLO distributions are both in the $\overline{MS}$ scheme. To check the contribution of the higher order matrix elements to the higher order corrections we will also follow the conventional procedure of using exactly the same structure functions in both calculations although in the LO case this is inconsistent.

It is clear from fig. 3(a) that the corrections to the direct processes are only significant, though not large, at low p_T^2 . At $p_T^2 = 5 \text{ GeV}/c$ they are 18% and decrease to 2.5% at $p_T^2 = 40 \text{ GeV}/c$. The corresponding figures for the $O(\alpha_s\alpha_{em})$

fragmentation contributions of eq. (4) are 30% and 5% respectively. It is thus safe to say that the higher order corrections are under control in this case. Fig. 3(b) shows that, for the resolved contribution, if we assume that the fragmentation processes contribute in NLO only, then the higher order corrections are very large. They also do not decrease rapidly with increasing p_T^2 . At $p_T^2 = 5 \text{ GeV}/c$ they are 90%, at $p_T^2 = 40 \text{ GeV}/c$ they are still 40% and even out to p_T^2 of 60 GeV/c they still contribute 20%. If, on the other hand, we assume, more properly, that the fragmentation processes contribute in LO also then the corresponding figures are 20%, 15% and 3% respectively. The second set of figures can be considered small in comparison to errors from other sources such as the uncertainty in the structure functions.

We note that if we use the LO distribution functions in the leading order part of the calculation and the NLO distributions in the full calculation the figures do not change by more than 1 – 4% from those given above. This suggests that the higher order corrections to the hard scattering cross sections are responsible for most of the differences between the LO and NLO results.

The full NLO resolved contributions completely dominate the cross section in the region $5 \text{ GeV}/c \leq p_T^2 \leq 15 \text{ GeV}/c$ but fall very steeply with increasing p_T^2 and are negligible compared to the direct ones beyond $p_T^2 = 20 \text{ GeV}/c$ (fig. 4). At $p_T^2 = 5 \text{ GeV}/c$ and $10 \text{ GeV}/c$ respectively, the ratio of the resolved to direct contributions is respectively 5.5 and 1.5. This indicates that it will be possible to obtain information on the parton distributions of the photon in the lower p_T^2 region only. Our results also indicate that the full cross section, i.e. resolved plus direct contributions, will be small beyond $p_T^2 = 25 - 35 \text{ GeV}/c$, and will thus be difficult to measure. Assuming an integrated luminosity of $100 \text{ pb}^{-1}\text{year}^{-1}$, there will be $O(35000)$ events per year at $p_T^2 = 5 \text{ GeV}/c$ for both resolved and direct contributions.

This decreases to 1200 events at $\hat{p}_T = 10 \text{ GeV}/c$, and at $\hat{p}_T = 30 \text{ GeV}/c$ there will only be $O(50)$ events.

All these results suggest that the measurement of the photon distribution functions using the inclusive direct photon cross section at HERA will not be as easy as previously anticipated from LO analyses [1,20]. It is far more likely that only indirect information on these functions will be obtainable from a study of this cross section. This could be obtained by comparing measurements on the cross section with QCD predictions and checking for sensitivity to these functions. There is also the possibility that the study of more cross sections, such as the cross section for isolated prompt photon plus jet production will give more detail. This is suggested from studies of jet production at HERA where the single jet inclusive cross section yields similarly limited information, but where the two-jet cross sections yield clear separation of the subprocesses of eqs.(6) [5,23].

If the KMRs(B_0) or GRV' parametrisations of the proton distribution functions are used instead, the results are not substantially different (fig. 5(a)). The KMRs(B_0) distributions give a 2% smaller and the GRV' an 8% larger cross-section at $\hat{p}_T = 5 \text{ GeV}/c$. These already small differences become even smaller at larger $\hat{p}_T$. These figures are not substantially different when the resolved and direct contributions are taken separately.

Fig. 5(b) shows the ratio of the cross sections predicted by the photon structure function parametrisations of refs. [18] and [19] for the full cross section and LO contribution to it, where the parametrisation of the right order was used in each case. In both cases the GRV' parametrisations give larger cross sections than the GS ones. The differences are substantially more significant in higher order where they are 60% and 45% respectively at $\hat{p}_T = 5$ and $60 \text{ GeV}/c$ respectively. The corresponding figures are 20% and 5% in the LO case. The reason for the larger

predictions of the GRV' parametrisations is probably because they started their evolutions at very low Q^2 ($\sim 0.25 \text{ GeV}^2$) thereby generating many more gluons than in the GS case where the evolution was started at $Q^2 = 5.3 \text{ GeV}^2$ with a similar gluon input. This is supported by the fact that the differences mentioned above are significantly larger at lower $\hat{p}_T$ where the quark and gluon distributions are probed at low x . This is where the gluon distribution most affects the quark distributions during the evolution [18]. The reason for the bumps in the curves at medium $\hat{p}_T$ is due to the differences in the way the charm contribution is included in each case [18]. It is also useful to note that the reason for the different ratios in leading and higher order can be explained by the fact that in ref. [19] no attempt was made to obtain agreement between the leading and higher order 'fits' to the data, whereas in ref. [18] such an attempt was made. Also in ref. [19] the NLO parametrisations are presented in the $D/\bar{S}_\gamma$ scheme and have to be transformed to the $\overline{MS}$ scheme. This may also be responsible for some of the differences seen between the leading order and higher order comparisons.

The cross section was found to be sensitive to the photon fragmentation function used, increasing at all $\hat{p}_T$ when the parton model fragmentation functions are used. This can not be taken as proof that information on these functions will be obtainable however, since the parton model fragmentation functions are completely unrealistic, partly because the gluon fragmentation function is zero in this approximation.

3.2 Rapidity Distribution

In this subsection we examine the cross section at different rapidity values exclusively in the EPA and include NLO corrections to all processes. We use exclusively the GS photon distributions and the GRV' proton distributions, both in LO and NLO. In the resolved and direct processes of eq. (3) and (5) the matrix elements of refs. [16] are

used. Although these matrix elements are calculated for hadron-hadron scattering and do not include the direct photon processes directly, these are easily obtained from the reactions $qg \rightarrow \gamma qg$ and $gq \rightarrow q\bar{q}g$ by removing non-Abelian couplings and adjusting colour factors. The matrix elements of Aversa *et al* [35] are used for the resolved fragmentation contributions (eq.(6)) and for the direct fragmentation processes of eq. (4) we use those recently calculated by Gordon [17]. As before, we refer to the references cited for details of the calculations of these matrix elements. The only inconsistency in our calculation is the use of the asymptotic fragmentation function.

Figs. 6(a) and (b) show the non-fragmentation and fragmentation contributions respectively in both LO and NLO at $p_T^\gamma = 5 \text{ GeV}/c$. In the case of the contribution without fragmentation (eqs. (3) and (5)) there is no clear separation between resolved and direct in both LO and NLO although the resolved is larger at negative rapidities in LO. The separation is somewhat better in the case of the fragmentation contribution although a clear separation only occurs at large negative rapidities which may not be accessible to experiment.

In fig. 6(c) we show a breakdown of the resolved contributions of eq. (5) and include the direct process of eq. (3) and both resolved and direct fragmentation contributions for comparison. We omit the contribution from the gq initial state because it is so small and by the initial state $q\bar{q}$ we mean all possible combinations of quarks and anti-quarks in the initial state. Unfortunately, it is clear that the processes are not separated in rapidity to any significant extent. We note that the fragmentation contributions will be somewhat reduced when the isolation criteria are taken into account, but this cannot affect our conclusions about the non-fragmentation contributions.

Lastly, in fig. 6(d) the $\hat{p}_T^\gamma$ distribution at rapidity $y = -2$ is shown. We chose this

value as one which will be accessible in the HERA experiments. The curves show that the expected relationship with the direct non-fragmentation contributions dominating as p_T^γ is increased and both resolved and direct fragmentation contributions falling away rapidly. The curves also show that the resolved contributions would dominate the cross section at very small p_T^γ ($\leq 3 \text{ GeV}/c$), but it is questionable whether this information is of any use since the use of perturbation theory must be doubtful at such small p_T^γ and it is also probably not possible for experimentalists to identify the prompt photon signal here.

4 Conclusions

We have performed an up-to-date estimate of the cross section for single prompt photon inclusive production at HERA fully in NLO QCD using the EPA. Our main conclusion is that it will be difficult to use this cross section to isolate those contributions involving a gluon from the photon, thereby directly measuring this distribution, unless the scattered electron can be tagged very precisely. This is due to a combination of factors. The lack of a fixed cm energy in the γp system due to the fact that the basic process is electroproduction as opposed to photoproduction leads to a spread in the energies of the initial state photons, and hence to difficulties in the angular separation of the final states of the various subprocess contributions. This may be avoidable by tagging of the scattered electron and hence fixing the energy of the initial state photon but this will significantly reduce the size of the cross section. Another factor which makes this separation more difficult is that the distinction between direct and resolved processes is an artefact of LO calculations, as we have discussed. We have demonstrated that the higher order corrections to the processes relevant for the measurement of the photon distribution functions, i.e. the resolved processes, are significant at the values of $\hat{p}_T^\gamma$ where these processes dominate, and

these corrections are not the same for all subprocesses. An interesting finding is that the parametrisation of ref. [19] for the photon distribution functions gives a larger predicted cross section than that of ref. [18], markedly so in NLO.

Our results appear to contradict the results of Aurenche et al [25] who compute higher order QCD corrections to the photoproduction of direct photons at HERA energies and conclude that the gluon content of the photon is measureable. However they work with a fixed photon energy (of 10 GeV) and present cross sections in the γp CM frame: we have demonstrated how misleading that can be.

We also point out that the fragmentation functions used in our work could and indeed should be improved. Some progress in this area has been reported in two recent papers [36,37].

Acknowledgements

We would like to thank J. Ph. Guillet for providing us with the FORTRAN code of ref. [35] and for helpful explanations.

References

- [1] A. C. Bawa and W. J. Stirling, J. Phys. **G14** (1988) 1353.
- [2] E. Witten, Nucl. Phys. **B120**, (1977) 189.
- [3] R. Tanaka *et al.*, AMY Collaboration, Phys. Lett. **B277** (1992) 215; H. Hayashii *et al.*, TOPAZ Collaboration, Phys. Lett **B314** (1993) 149.
- [4] M. Drees and R. Godbole Phys. Rev. Lett. **61** (1988) 682; Phys. Rev. **D39** (1989) 169.
- [5] H. Baer, J. Ohnemus and J. F. Owens, Z. Phys. **C42** (1988) 657.
- [6] R. S. Fletcher, F. Halzen and R.W. Robinett, Phys. Lett. **B225** (1989) 176.
- [7] Z. Kunszt and W. J. Stirling Phys. Lett. **B217** (1989) 563
- [8] R. S. Fletcher, F. Halzen S. Keller, and W.H. Smith, Phys. Lett. **B266** (1991) 183.
- [9] M. Glück, E. Reya and A. Vogt, Phys. Lett. **B285** (1992) 285; M. Glück, E. Reya and A. Weber, Phys. Lett. **B298** (1993) 176.
- [10] L. E. Gordon and J. K. Storrow, Phys. Lett. **B291** (1992) 320.
- [11] G. Kramer and S. G. Salesch, 'Single Jet Photoproduction at HERA in Next-to-Leading Order QCD' DESY preprint, DESY 93-010 (unpublished); F. Borzumati, B. Kniel and G. Kramer, Z. Phys. **C59** (1993) 341.
- [12] D. Bödeker, Phys. Lett. **B292** (1992); Z. Phys. **C59** (1993) 501.

- [13] M. Greco and A. Vicini, 'Photoproduction of High p_T jets in Next-to-Leading Order QCD', Frascati preprint LNF-93/017.
- [14] P. Aurenche, R. Baier, A. Douiri, M. Fontannaz, and D. Schiff, Nucl. Phys. **B286** (1987) 553.
- [15] P. Aurenche, A. Douiri, R. Baier, M. Fontannaz, and D. Schiff, Z. Phys. **C24** (1984) 309.
- [16] L. E. Gordon and W. Vogelsang, Phys. Rev. **D48** (1993) 3136.
- [17] L. E. Gordon, 'Next-to-leading order corrections to inclusive hadron photoproduction', Dortmund preprint DO-TH 93/25.
- [18] L. E. Gordon and J. K. Storrow, Z. Phys. **C56** (1992) 307.
- [19] M. Glück, E. Reya and A. Vogt, Phys. Rev. **D46** (1993) 1973; Phys. Rev. **D45** (1992) 3986.
- [20] A. C. Bawa, M. Krawczyk and W. J. Stirling, Z. Phys. **C50** (1991) 293.
- [21] E. L. Berger and J. Qiu, Phys. Rev. **D44** (1991) 2002.
- [22] J. K. Storrow, J. Phys. G **19** (1993) 1641.
- [23] L. E. Gordon and J. K. Storrow, 'Structure Function Sensitivity of Leading Order Jet Cross Sections at HERA and the Measurement of the Gluon Distribution g^* ', Manchester preprint M/C.TH.92/06, J. Phys. G (to be published).
- [24] J. Chyla, Phys. Lett. **B320** (1993) 186.
- [25] P. Aurenche, P. Chiappetta, M. Fontannaz, J. Ph. Guillet and E. Pilon, Z. Phys. **C56** (1992) 589.
- [26] C. F. von Weizsacker, Z. Phys., **88** (1934) 612 ; E. J. Williams, Phys. Rev. **45** (1934) 729.
- [27] A. C. Bawa and W. J. Stirling, J. Phys. G **15** (1989) 1339; A. C. Bawa, Ph.D. Thesis, University of Durham (1989).
- [28] B. Diekmann *et al.*, *Proceedings of the HERA Workshop*, Vol. 2 Hamburg (1988) 515. Ed. Pececi, R.D.
- [29] J. G. Morfin and W. K. Tung, Z. Phys. **C52** (1991) 13.
- [30] D. W. Duke and J. F. Owens, Phys. Rev. **D26** (1982) 1600.
- [31] B. L. Combridge, J. Kripfganz, and J. Raftl, Phys. Lett. **B70** (1977) 2
- [32] M. Glück, E. Reya and A. Vogt, Z. Phys. **C53** (1992) 127.
- [33] J. Kwiecinski, A. D. Martin, R. G. Roberts and W. J. Stirling, Phys. Rev. **D42** (1990) 3645.
- [34] P. Aurenche, R. Baier and M. Fontannaz, Phys. Rev. **D42** (1990) 1440
- [35] F. Aversa *et al.* Nucl. Phys. **B327** (1989) 105.
- [36] P. Aurenche, P. Chiappetta, M. Fontannaz, J. Ph. Guillet and E. Pilon, Nucl. Phys. **B399** (1993) 34.
- [37] M. Glück, E. Reya and A. Vogt, Phys. Rev. **D48** (1993) 116.

Figure Captions

- [1] (a) Minimum value of x_p , the Bjorken variable for the proton, probed when the calculation for inclusive prompt photon production at HERA is performed in the fixed γp cm frame with an estimated cm energy of $S_{\gamma p} = 30000 \text{ GeV}^2$ (broken line) and when the calculation is performed in the ep lab frame in the EPA (solid line), both as functions of rapidity and both corresponding to $p_T^\gamma = 5 \text{ GeV}/c$.
- (b) u -quark distribution function (at $Q^2 = 100 \text{ GeV}^2$) before (u^b ; solid line) and after (u^c ; broken line) convolution with the EPA formula.
- [2] Rapidity distribution of the leading contributions to inclusive prompt photon production at HERA at $p_T^\gamma = 5 \text{ GeV}/c$ using (a) a fixed cm energy in the γp system of $S_{\gamma p} = 30000 \text{ GeV}^2$ and (b) the same contributions in the ep lab system using the EPA. (c) Rapidity distribution for the direct process $\gamma q \rightarrow \gamma q$ for the full EPA (broken line), and various tagging situations [28]; 'poor tag' (solid line) and 'golden tag' (dashed line).
- [3] $R(p_T^\gamma)$, as defined in the text (eq. 7), vs p_T^γ for (a) direct and (b) resolved contributions to the single photon inclusive cross section at HERA. Legend: (a) (— — —) ratio of Born (eq. 3)) plus fragmentation processes (eq.(4)) to Born terms alone; (——) ratio of Born plus NLO corrections (including the box diagram process $\gamma g \rightarrow \gamma g$ [14]) to Born ($\gamma q \rightarrow \gamma q$). (b) (——) ratio of full NLO cross section relative to the basic processes of eq. 5, (— — —) ratio of full NLO cross section relative to the sum of basic processes of eq. 5 and the fragmentation processes of eq. 6.
- [4] p_T^γ distribution of the single photon inclusive cross section integrated over all rapidities showing resolved and direct contributions and their sum.

- Legend: (— — —) direct contributions, (— — —) resolved contributions, and (——) sum.
- [5] (a) Ratio of the cross section $d\sigma/dp_T^\gamma$ for inclusive prompt photon production at HERA using the GRV' and KMRS(B_0) proton structure function parametrisation to that using the MT(5) proton. (b) solid line (——) ratio of the full next-to-leading order resolved contribution to the cross section as predicted using the parametrisation of ref. [19] to that using those of ref. [18] for the photon distribution functions and broken line; (— — —) same but for the leading order contribution of eq. 4 only and using the LO parametrisations of the same references.
- [6] Plots of various contributions to the single photon inclusive cross section at HERA:
- (a) Rapidity distributions of non-fragmentation contributions, both direct (eq. 3) and resolved (eq. 5) in LO (broken curves) and NLO (solid curves) at $p_T^\gamma = 5 \text{ GeV}/c$.
- (b) Same as (a) except for fragmentation contributions (eqs. 4 and 6).
- (c) Rapidity distributions for the breakdown of the resolved non-fragmentation contributions (eq.5); we also show the direct non-fragmentation contribution of eq.3 and both resolved and direct fragmentation contributions for comparison. All at $p_T^\gamma = 5 \text{ GeV}/c$ and labelled.
- (d) p_T^γ distribution at $y^\gamma = -2$ for direct (broken line) and resolved (solid line) fragmentation (F) and non-fragmentation (N-F) contributions (labelled).

Fig.1

Fig.2

Fig.3

Fig.4

Fig.5

Fig.6