MESON PRODUCTION IN e^+e^- ANNIHILATION IN THE LEP ENERGY RANGE
INCLUDING GLUINO FRAGMENTATION

E.C. Katsoufis* and S.D.P. Vlassopoulos

Department of Physics, National Technical University,

GR-15773 Zografou, Athens, Greece

We calculate π and K meson production rates in e^+e^- collisions at LEP energies, taking into account contributions introduced by supersymmetric QCD below the squark production threshold, assumed to be further away. Although suppressed in the inclusive rate, we find a non-negligible probability for gluinos to fragment into slow ordinary mesons.

* Present address : EP Division, CERN, Geneva

CONTENTS

	<u>Page</u>
I. INTRODUCTION	1
II. MESON PRODUCTION CROSS SECTIONS	2
III. SQCD EVOLUTION EQUATIONS	5
IV. MODEL	7
V RESULTS AND CONCLUSIONS	10
ACKNOWLEDGEMENTS	11
REFERENCES	12
TABLE I	15
TABLE II	16
FIGURES	19

I. INTRODUCTION

Scaling violations in hadron production in e^+e^- collisions have been known experimentally to be present at currently available energies¹⁾ and to be small - in agreement with the leading logarithmic approximation of quantum chromodynamics (QCD-LLA), which is found to be working at surprisingly small values of $z=2E_h/Q$, where Q is the centre-of-mass energy of the collision and E_h is the energy of the observed hadron²⁾. It is of interest to know whether the introduction of supersymmetry (SUSY)³⁾, can possibly modify this picture, especially in the extended energy range to be explored by LEP. Indeed, SUSY gauge theories require the existence of a bosonic (fermionic) partner to every known fermion (boson) and in particular of new partons, namely squarks and gluinos, thereby enriching the final-state content of the process $e^+e^- \rightarrow$ hadrons⁴⁾.

Detection of squark jets (and slepton pairs and gauginos) would be direct experimental evidence for SUSY. In view of the diversity of various model estimates of the mass of SUSY partners⁵⁾, we assume in this paper, pessimistically, that the squark production threshold lies beyond LEP energies. We calculate the inclusive π and K production rates in e^+e^- interactions up to $Q \sim 200$ GeV assuming that the gluino production threshold lies near the presently highest PETRA-PEP energies ($Q \sim 40$ GeV). This is in qualitative accord with estimations based on recent CERN-Collider findings⁶⁾. Making use of the

good quality PETRA-PEP data⁷⁾ we extract the fragmentation functions of the ordinary partons into π and K mesons at $Q \sim 30$ GeV, where the gluino fragmentation functions are taken to be zero. We then extrapolate these functions to higher Q values, making use of the QCD-LLA evolution equations²⁾, which are extended to allow for gluino contributions, besides the standard quark and gluon fragmentation terms.

Supersymmetric QCD (SQCD) effects are, of course, suppressed in the inclusive cross section below the squark production threshold, being of the order a_s^2 (in contrast to $O(a_s)$ ordinary QCD effects). Nevertheless, the fragmentation functions of the gluino to slow mesons turn out to be non-negligible: For example, we find that the probability for a gluino to fragment into a pion is larger than the probability for a gluon to fragment into a kaon for $z \lesssim 0.05$ and $Q \gtrsim 150$ GeV. This may well affect the ratio of K/ π relative production rates in hadronic collisions in the TeV energy range.

II. MESON PRODUCTION CROSS SECTIONS

The inclusive cross section for hadron production in e^+e^- annihilation is given by²⁾

$$\frac{1}{\sigma_t(Q^2)} \frac{d\sigma^h}{dz}(z, Q^2) = \sum_q e_q^2 \left[D_q^h(z, Q^2) + D_{\bar{q}}^h(z, Q^2) \right] / \sum_q e_q^2, \quad (1)$$

where σ_t is the total cross-section for $e^+e^- \rightarrow \text{hadrons}$ and e_q is the quark charge. This relation stands in fact for the definition of the Q^2 -dependent fragmentation function D_q^h of the quark q into the hadron h and brings QCD-LLA in correspondence to the naive parton model. Since gluons and gluinos do not couple to electroweak currents in LLA, their fragmentation functions appear only implicitly in (1) through the Q^2 -evolution of D_q^h (see below). The ratio $\sigma_t^{-1} d\sigma/dz$, being insensitive to either electroweak effects or z -independent $O(a_s)$ corrections, is suitable for testing SQCD-LLA.

Taking into account isospin conservation, and assuming, as usual, that all quarks fragment with equal probability to a given hadron if they belong to its sea content⁸⁾, but neglecting the fragmentation of b into π and K mesons, we get from Eq. (1)

$$\frac{1}{\sigma_t} \frac{d\sigma^{\pi^\pm}}{dz} = \frac{1}{\sigma_t} \frac{d\sigma^{\pi^0}}{dz} = \frac{1}{4} \Sigma^\pi, \quad (2)$$

$$\frac{1}{\sigma_t} \frac{d\sigma^{K^\pm}}{dz} = \frac{1}{4} \Sigma^K + \frac{1}{4} \frac{e_u^2 - e_d^2}{e_u^2 + e_d^2} (\Delta^K - \Delta_S^K), \quad (3a)$$

$$\frac{1}{\sigma_t} \frac{d\sigma^{K^0, \bar{K}^0}}{dz} = \frac{1}{4} \Sigma^K - \frac{1}{4} \frac{e_u^2 - e_d^2}{e_u^2 + e_d^2} (\Delta^K + \Delta_S^K), \quad (3b)$$

where

$$\Sigma^h = \sum_q (D_q^h + D_{\bar{q}}^h) , \quad \{h = \pi, K\} \quad (4)$$

transform as siglet and

$$\Delta^K = D_u^{K^+} - D_{\bar{u}}^{K^+} , \quad \Delta_S^K = D_{\bar{s}}^{K^+} - D_s^{K^+} \quad (5)$$

transform as non-singlet combinations.

The data show that within errors, we have $d\sigma^{K^\pm}/dz \approx d\sigma^{K^0, \bar{K}^0}/dz$ (7), implying $\Delta^K, \Delta_S^K, < \Sigma^K$. In the following we set

$$\frac{1}{\sigma_t} \frac{d\sigma^{K^\pm}}{dz} = \frac{1}{\sigma_t} \frac{d\sigma^{K^0, \bar{K}^0}}{dz} = \frac{1}{4} \Sigma^K , \quad (3c)$$

which is consistent (within 10%) with the parametrizations of refs 8).

Energy conservation requires

$$\sum_h \int_0^1 z D_q^h(z, Q^2) dz = 1 \quad (6a)$$

at all Q^2 . The evolution equations preserve this constraint, once it is imposed at one value of $Q=Q_0$. Summed over all quarks and antiquarks, Eq (6a) gives

$$3 \int_0^1 z \Sigma^\pi dz + 4 \int_0^1 z \Sigma^K dz \lesssim 2f , \quad (6b)$$

where f is the number of quark flavours.

III. SQCD EVOLUTION EQUATIONS

In the presence of gluinos the QCD evolution equations for the fragmentation functions are modified to

$$Q^2 \frac{\partial}{\partial Q^2} D_q^h(z, Q^2) = \frac{a_s(Q^2)}{2\pi} \int_z^1 \frac{dy}{y} \left[P_{qq}\left(\frac{z}{y}\right) D_q^h(y, Q^2) + P_{gq}\left(\frac{z}{y}\right) D_g^h(y, Q^2) \right], \quad (7a)$$

$$Q^2 \frac{\partial}{\partial Q^2} D_g^h(z, Q^2) = \frac{a_s(Q^2)}{2\pi} \int_z^1 \frac{dy}{y} \left[P_{qg}\left(\frac{z}{y}\right) \sum_{q=1}^{2f} D_q^h(y, Q^2) + P_{gg}\left(\frac{z}{y}\right) D_g^h(y, Q^2) + P_{\lambda g}\left(\frac{z}{y}\right) D_\lambda^h(y, Q^2) \right], \quad (7b)$$

$$Q^2 \frac{\partial}{\partial Q^2} D_\lambda^h(z, Q^2) = \frac{a_s(Q^2)}{2\pi} \int_z^1 \frac{dy}{y} \left[P_{g\lambda}\left(\frac{z}{y}\right) D_g^h(y, Q^2) + P_{\lambda\lambda}\left(\frac{z}{y}\right) D_\lambda^h(y, Q^2) \right]. \quad (7c)$$

They are depicted in Fig. 1.

The coupling strength of QCD is modified by the new degrees of freedom of fermionic gluinos to⁹⁾

$$a_s(Q^2) = \frac{12\pi}{33-2f-2N_\lambda} \frac{1}{\ln Q^2/\Lambda_s^2} \quad (8)$$

where $N_\lambda=3$ for Majorana gluinos. The effective scale parameter Λ_s is chosen so that (8) matches with the coupling strength of the ordinary QCD ($N_\lambda=0$) at the gluino production threshold $Q_0=2m_\lambda$ ⁹⁾. Note that the scalar quarks decouple from the evolution kernel if we stay below their production threshold⁹⁾.

For $Q^2 \gg m_\lambda^2$ the new elements of the anomalous dimension matrix are simply related⁹⁾ to the ordinary ones, which can be found in the standard literature²⁾, by an appropriate change of the group factors, namely

$$P_{g\lambda} = \frac{9}{4} P_{gq}, \quad P_{\lambda g} = 3P_{qg}, \quad P_{\lambda\lambda} = \frac{9}{4} P_{qq}. \quad (9)$$

Taking the Mellin transform $\{\tilde{f}(n) \equiv \int_0^1 x^{n-1} f(x) dx\}$ of Eqs (7) we obtain, as usual, the coupled first-order differential equations

$$\frac{\partial}{\partial \xi} \begin{bmatrix} \tilde{\Sigma}^h(n, \xi) \\ \tilde{D}_g^h(n, \xi) \\ \tilde{D}_\lambda^h(n, \xi) \end{bmatrix} = \begin{bmatrix} P_{qq}(n) & 2fP_{gq}(n) & 0 \\ P_{qg}(n) & P_{gg}(n) & P_{\lambda g}(n) \\ 0 & P_{g\lambda}(n) & P_{\lambda\lambda}(n) \end{bmatrix} \cdot \begin{bmatrix} \tilde{\Sigma}^h(n, \xi) \\ \tilde{D}_g^h(n, \xi) \\ \tilde{D}_\lambda^h(n, \xi) \end{bmatrix} \quad (10)$$

where

$$\xi = \frac{6}{33-2f-2N_\lambda} \ln \frac{\ln Q^2 / \Lambda_s^2}{\ln Q_0^2 / \Lambda_s^2}. \quad (11)$$

Note that the elements $P_{\lambda q}$ and $P_{q\lambda}$ of the anomalous-dimension matrix given in terms of graphs involving squarks vanish in this model.

IV. MODEL

Fragmentation function parametrization and moment fitting

Following previous attempts^{8,10)} we parametrize the singlet fragmentation functions according to

$$z\Sigma^h(z, Q^2) = a_q^h(Q^2) z^{-\rho_q^h(Q^2)} (1-z)^{\mu_q^h(Q^2)}, \quad (12a)$$

$$zD_i^h(z, Q^2) = a_i^h(Q^2) z^{-\rho_i^h(Q^2)} (1-z)^{\mu_i^h(Q^2)}, \quad (12b)$$

where

$$a_i^h(Q^2) = a_{i0}^h + a_{i1}^h \xi + a_{i2}^h \xi^2 \quad \{i = q, g, \lambda\},$$

$$\rho_i^h(Q^2) = \rho_{i0}^h + \dots \text{ etc} \quad (13)$$

The reference point Q_0 will be taken at $Q_0 = 30$ GeV, i.e. near the top PETRA-PEP energies. Lacking experimental information on the gluino fragmentation function, we plausibly take

$$D_\lambda^h(z, Q_0^2) = 0.$$

We take $\Lambda = 0.4$ GeV but of course our parametric forms (12) are independent of this choice, which only affects the appearance of our final results at $Q > Q_0$ (Figs 2-4).

The parameters a_{ik}^h , ρ_{ik}^h , μ_{ik}^h ($k=1,2$) are found as usual²⁾ by fitting the $n=2,3,\dots,6$ moments of (12) (given in terms of Euler's B-functions) at $\xi = .02, .04, \dots$ to their SQCD values constructed by solving the coupled Eqs (10). The eigenvalues of the 3×3 evolution matrix in the right hand side of Eq.(10) are not in general given by simple

algebraic expressions as in standard QCD²⁾. Thus we must solve the system (10) numerically given the initial conditions for $\bar{\Sigma}^h(n,0)$, $\bar{D}_g^h(n,0)$ and that $\bar{D}_\lambda^h(n,0) = 0$.

Initial conditions

a. Quark singlet-combinations: To obtain an adequate expression of the initial forms of quark singlet fragmentation combinations we fit

$$\frac{1}{\sigma} \frac{d\sigma^h}{dz} = \frac{1}{4} a_{q0}^h z^{-1-\rho_{q0}^h} (1-z)^{\mu_{q0}^h} \quad (14)$$

to selected π and K inclusive production data at $Q_0=29-33\text{GeV}^7$). We have a good fit ($\chi^2/\text{NDF}=10.3/9$ for π 's and $20.3/19$ for K's) with the values of the parameters a_{q0}^h , ρ_{q0}^h , and μ_{q0}^h given in Tables I and II. The results of this fit are shown in Fig. 2.

The energy conservation constraint (6a) is satisfied by the results of our fit, the left hand side of (6b) being equal to 8.2 ± 1.7 , which shows the magnitude of the total baryon production contribution. Incidentally, we stress the point that the experimental data used for the fits include the decay products of the resonances ($\rho, \omega, K^*, \text{etc.}$): The fragmentation functions used in our analysis are defined in this effective sense.

b. Gluon and gluino fragmentation functions: Due to the lack of sufficient experimental information, the initial forms of the gluon and gluino fragmentation functions

$$zD_i^h(z, Q_0^2) = a_{io}^h z^{-\rho_{io}^h} (1-z)^{\mu_{io}^h} \quad (i=g, \lambda) \quad (15)$$

are chosen on a much less stringent basis. We take

$$\rho_{\lambda o}^h : \rho_{go}^h : \rho_{qo}^h = 1 \quad , \quad (16)$$

while the exponents

$$\mu_{go}^\pi \approx \mu_{go}^K \approx 2.5 \quad , \quad (17)$$

controlling the behaviour of the gluon fragmentation functions for $z \rightarrow 1$, are taken from the analysis of ref (10). For the gluino decay function we try both

$$\mu_{\lambda o}^\pi = \mu_{\lambda o}^K = \mu_{go}^h = 2.5 \quad (18a)$$

and

$$\mu_{\lambda o}^\pi = \mu_{\lambda o}^K = 5 \quad . \quad (18b)$$

The normalization constants a_{go}^h are determined from the energy conservation relation

$$3 \int_0^1 z D_g^\pi(z, Q_0^2) + 4 \int_0^1 z D_g^K(z, Q_0^2) \approx 1 \quad , \quad (19)$$

together with the estimation⁸⁾

$$a_{go}^K : a_{go}^\pi = 0.5 . \quad (20)$$

We have tested the sensitivity of our results against variation of the assumptions (16) and (20): Varying any of the numerical right hand sides of these equations by 10% produces no more than a 10% variation in $\Sigma^h(z, Q^2)$ and $D_i^h(z, Q^2)$ for $0.02 \lesssim z \lesssim 0.9$ and $30 \lesssim Q \lesssim 240$ GeV.

V. RESULTS AND CONCLUSIONS

The values of the parameters, which resulted from the moment fitting, are shown in Tables I, II. Fig. 3 shows our results for the singlet fragmentation functions Σ^h , D_g^h at LEP energies. As in standard QCD, more significant scaling violation is found in the gluon fragmentation function. The corresponding gluino fragmentation functions, generated by the full 3×3 kernel, are shown in the same figure: At LEP energies they turn out to be about 10% of the corresponding gluon fragmentation functions D_g^h at $z \approx 0.2$, and, being much steeper than D_g^h , they grow to significant values as $z \rightarrow 0$.

Figure 4 shows our output gluino fragmentation functions at $Q=60$ and 210 GeV for the two choices (18) of the initial value of the exponent of $1-z$ at $Q_0=30$ GeV: These choices give results differing by less than 25% for $z \lesssim 0.5$

In Fig. 2 we also present our predictions for π as well as K production cross section $\sigma^{-1} d\sigma/dz$ at LEP energies. These are practically identical with the predictions of standard QCD (i.e. without gluinos) with the same initial conditions (14) and (15). Very good quality data are required in order to test quantitatively these small scaling violations.

We conclude that, although suppressed in the inclusive cross section below the squark threshold, gluino fragmentation into ordinary mesons can be important at small z and large Q^2 . This can have direct consequences for meson production in pp , $\bar{p}p$ and γp interactions at very high energies.

ACKNOWLEDGEMENTS

We are grateful to the TASSO Collaboration for providing us with data, prior to their publication, at the early stages of this work. One of us (S.D.P.V.) wishes to thank the CERN TH-Division, where part of this work was done, for its hospitality. We are indebted to T.A. Filippas and G. Tiktopoulos for reading the manuscript.

REFERENCES

- 1) D. Lüke (TASSO results), Proc. 21st Int. Conf. on High-Energy Physics (Paris, 1982);
J.F. Patrick et al. (SPEAR and PEP results), Phys. Rev. Lett. 49, 1232(1982).
- 2) For example, A.J. Buras, Rev. Mod. Phys. 52, 199(1980);
Yu.L. Dokshitzer, D.I. Dyakonov and S.J. Troyan, Phys. Reports 58, 269(1980);
E. Reya, Phys. Reports 69, 195(1981);
G. Altarelli, Phys. Reports 81,1(1982).
- 3) Yu.A. Gel'fand and E.P. Likhtham, JETP Lett. 13, 323(1971);
D. Volkov and V.P. Akulov, Phys. Lett. 46B, 109(1973);
J. Wess and B. Zumino, Nucl. Phys. B70, 39(1974);
P. Fayet and S. Ferrara, Phys. Reports 32, 249(1977).
- 4) P. Fayet, Phys. Lett. 69B, 489(1977); 70B, 461(1977);
Proc. Conf. on Unification of the Fundamental Interactions, Erice 1980 (Plenum Press).
- 5) P. Fayet, Proc. 21st Int. Conf. on High-Energy Physics (Paris, 1982); also, LPTENS 83/19 (to be published in Phys. Reports).
- 6) G. Arnison et al. (UA1 Coll.), CERN-EP/84-42.
- 7) M. Althoff et al. (TASSO Results), Zeit. Phys. 17, 5(1983);
W. Bartel et al. (JADE Results), Zeit. Phys. 20, 187(1983);
H. Aihara et al. (PEP Results), LBL Preprint, TPC-LBL-83-31 (December 1983).

- 8) R.D. Field and R.P. Feynman, Phys. Rev. D15, 2590(1977);
Nucl. Phys. B136, 1(1978).
A.P. Contogouris, R. Gaskell and S. Papadopoulos, Phys.
Rev. D17, 2314(1978);
J. Owens, E. Reya and M. Glück, Phys. Rev. D18, 1501(1978).
- 9) B.A. Campbell, J. Ellis and S. Rudaz, Nucl. Phys. B198,
1(1982);
I. Antoniadis, C. Kounnas and R. Lacaze, Nucl. Phys. B211,
216(1983);
P.R. Harrison and C.H. Llewellyn-Smith, Nucl. Phys. B213,
223(1983);
C. Kounnas and D.A. Ross, Nucl. Phys. B214, 317(1983);
S.K. Jones and C.H. Llewellyn-Smith, Nucl. Phys. B217,
145(1983).
- 10) R. Baier, J. Engels and B. Petersson, Zeit. Phys. C2,
265(1979).

TABLE I

π	π_{aik}			ρ_{ik}^{π}			μ_{ik}^{π}		
	q	g	λ $\mu_{\lambda 0}^{\pi}=2.5, 5$	q	g	λ $\mu_{\lambda 0}^{\pi}=2.5, 5$	q	g	λ
0	4.33	.466	0.	.300	.300	0.300	3.45	2.5	2.5, 5.0
1	-4.96	-1.75	.716, 1.97	.616	1.98	10.8, 4.36	4.25	8.43	72.3, 23.2
2	-4.94	-1.30	2.06, -14.4	.661	1.83	-86.9, -6.05	2.68	-12.6	-394., -219.

TABLE II

K	μ_{aik}^K				ρ_{ik}^K				μ_{ik}^K			
$\begin{matrix} i \\ k \end{matrix}$	q	g	λ $\mu_{\lambda_0}^K=2.5, 5.0$		q	g	λ $\mu_{\lambda_0}^K=2.5, 5.0$		q	g	λ	
0	1.44	.233	0.		.14	.14	.14		2.61	2.5	2.5 , 5.0	
1	-2.83	-1.58	.365, 1.07		1.45	3.78	13.1, 4.94		3.52	5.09	72.3, 26.1	
2	-10.5	5.52	.570, -8.08		2.53	-11.2	-102., -4.19		-11.5	11.9	-410., -247.	

TABLE AND FIGURE CAPTIONS

Table I

Numerical values of the parameters entering our quark singlet, gluon and gluino fragmentation functions into pions according to Eqs (12) and (13).

Table II

Same as table I, for kaons

Figure 1

Illustration of the evolution equations (7): The intuitive interpretation of the ordinary Altarelli-Parisi equations is not affected by the presence of gluinos far away from their production threshold.

Figure 2

Fit to selected data for the inclusive cross sections $e^+e^- \rightarrow \pi^{\text{ch}} X$ and $e^+e^- \rightarrow K^{\text{ch}} X$ at centre-of-mass energy $Q \gtrsim 30$ GeV (where the gluino production threshold is assumed to be), together with our predictions at higher energies.

Figure 3

Comparison of the resulted fragmentation functions $\Sigma^h(z, Q^2)$ {labelled as $q \rightarrow h$ } and $D_i^h(z, Q^2)$ {labelled as $i \rightarrow h$ } for $i=g, \lambda$ and $h=\pi, K$ at $Q=90$ and 180 GeV. Note that $D_\lambda^\pi \sim D_g^K$ at very small z .

Figure 4

Dependence of the output gluino fragmentation functions on the initial value of the exponent μ_λ^h . Numbers give Q in GeV. {Left scale for D_λ^K , right scale for D_λ^π }.

Fig. 1

Fig. 2

Fig. 3

Fig. 4