CC

ISBN 0020-3866

CBPF-NF 93-064

sep 94/9

CBPF - CENTRO BRASILEIRO DE PESQUISAS FÍSICAS

Notas de Física

CERN LIBRARIES, GENEVA

P00022455

CBPF-NF-064/93

*Probing the Froissart Bound for
Models with Charged Vector Fields in
 $D=3$*

by

O.M. Del Cima

*Rio de Janeiro
1993*

NOTAS DE FÍSICA é uma pré-publicação de trabalho original em Física

NOTAS DE FÍSICA is a preprint of original works unpublished in Physics

Pedidos de cópias desta publicação devem ser enviados aos autores ou ã:

Requests for copies of these reports should be addressed to:

Centro Brasileiro de Pesquisas Físicas
Área de Publicações
Rua Dr. Xavier Sigaud, 150 - 4º andar
22.290 - Rio de Janeiro, RJ
BRASIL

CBPF-NF-064/93

*Probing the Froissart Bound for
Models with Charged Vector Fields in
 $D=3$*

by

O.M. Del Cima^{†}*

Centro Brasileiro de Pesquisas Físicas — CBPF/CNPq
Rua Dr. Xavier Sigaud, 150
22290-180 – Rio de Janeiro, RJ – Brasil

^{*}Bitnet e-mail:oswaldo@brlncc.bitnet

[†]Internet e-mail:oswaldo:oswaldo@cbfsul.cat.cbpf.br

Abstract

One discusses the tree-level unitarity and presents asymptotic behaviour of scattering amplitudes for 3-dimensional gauge-invariant models where complex Chern-Simons-Maxwell fields (with and without a Proca-like mass) are coupled to an Abelian gauge field.

The study of gauge field theories in 3 space-time dimensions has raised a great deal of interest since the early work of Deser, Jackiw and Templeton [1]. More recently, this line of investigation has been well-motivated in view of the possibilities of providing a gauge-theoretical foundation in the description of Condensed Matter phenomena, such as high- T_c superconductivity [2]. Also, the recent result on the Landau gauge finiteness of Chern-Simons theories is a remarkable property that makes 3-dimensional gauge theories so attractive [3].

In a recent work [4], some preliminary aspects of the dynamics concerning the interaction between Maxwell and complex Chern-Simons fields have been discussed. It was ascertained that the minimal coupling of the charged Chern-Simons-Maxwell (CSM*) and Chern-Simons-Maxwell-Proca (CSMP*) fields to a $U(1)$ -gauge field breaks a local symmetry that is crucial for the decoupling of negative-norm 1-particle states associated to the charged massive vector field. As a follow up, our purpose in the present letter is to probe more deeply the tree-level unitarity of the CSM* and CSMP* models in the presence of a minimal $U(1)$ coupling [4]. This shall be achieved through the computation of tree-level amplitudes for the scattering between a CSM* (and CSMP*) particle and the photon in $D = 3$.

The minimal coupling of a CSM*-field, B_μ , to a $U(1)$ -gauge field, A_μ , is described by the Lagrangian :

$$\begin{aligned} \mathcal{L}_{CSM} = & -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} - \frac{1}{2}G_{\mu\nu}^*G^{\mu\nu} + \frac{m}{2}\epsilon^{\alpha\mu\nu}B_\alpha^*G_{\mu\nu} + im\omega\epsilon^{\alpha\mu\nu}B_\alpha^*A_\mu B_\nu + \\ & -i\omega(G_{\mu\nu}^*A^\mu B^\nu - G_{\mu\nu}A^\mu B^{*\nu}) - \omega^2(A_\mu B_\nu - A_\nu B_\mu)A^\mu B^{*\nu}, \end{aligned} \quad (1)$$

where $G_{\mu\nu} \equiv \partial_\mu B_\nu - \partial_\nu B_\mu$ and $F_{\mu\nu} \equiv \partial_\mu A_\nu - \partial_\nu A_\mu$ are the field-strengths, m is a real parameter with dimension of mass and ω a coupling constant with dimension of $(\text{mass})^{\frac{1}{2}}$.

The Lagrangian that describes the $U(1)$ -invariant CSMP*-model reads :

$$\begin{aligned} \mathcal{L}_{CSMP} = & -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} - \frac{1}{2}G_{\mu\nu}^*G^{\mu\nu} + \frac{m}{2}\epsilon^{\alpha\mu\nu}B_\alpha^*G_{\mu\nu} + \hat{\mu}^2 B_\mu^* B^\mu + \\ & + im\omega\epsilon^{\alpha\mu\nu}B_\alpha^*A_\mu B_\nu - i\omega(G_{\mu\nu}^*A^\mu B^\nu - G_{\mu\nu}A^\mu B^{*\nu}) + \\ & -\omega^2(A_\mu B_\nu - A_\nu B_\mu)A^\mu B^{*\nu}, \end{aligned} \quad (2)$$

where $\hat{\mu}$ is a real parameter with dimension of mass.

Considering the Lagrangians above, we get the following free propagators for the B_μ -field :

$$\Delta_{CSM}^{\mu\nu}(k) = -i \frac{1}{k^2(k^2 - m^2)} \left[im \epsilon^{\mu\kappa\nu} k_\kappa + k^2 \left(\eta^{\mu\nu} - \frac{k^\mu k^\nu}{k^2} \right) \right], \quad (3.a)$$

$$\begin{aligned} \Delta_{CSMP}^{\mu\nu}(k) = & -i \frac{1}{(k^2 - m_+^2)(k^2 - m_-^2)} \left[im \epsilon^{\mu\kappa\nu} k_\kappa + (k^2 - \hat{\mu}^2) \left(\eta^{\mu\nu} - \frac{k^\mu k^\nu}{k^2} \right) \right] + \\ & + i \frac{1}{\hat{\mu}^2} \left(\frac{k^\mu k^\nu}{k^2} \right), \end{aligned} \quad (3.b)$$

where

$$m_\pm^2 = \frac{1}{2} [m^2 + 2\hat{\mu}^2 \pm \sqrt{m^2(m^2 + 4\hat{\mu}^2)}] \quad (4.a)$$

and

$$m_-^2 = \frac{1}{2}[m^2 + 2\hat{\mu}^2 - \sqrt{m^2(m^2 + 4\hat{\mu}^2)}] . \quad (4.b)$$

From the interaction pieces of Lagrangians (1) and (2), we can read the 3- and 4-vertex Feynman rules required for our tree-level calculations (see Fig.1) :

$$(V_3)_{\alpha\mu\nu} = m\omega\epsilon_{\alpha\mu\nu} , \quad (5.a)$$

$$(\bar{V}_3)_{\alpha\mu\nu} = i\omega(\eta_{\nu\alpha}k_\mu - \eta_{\mu\alpha}k_\nu + \eta_{\nu\alpha}m_\mu - \eta_{\mu\nu}m_\alpha) \quad (5.b)$$

and

$$(V_4)_{\alpha\nu\beta\mu} = i2\omega^2(\eta_{\alpha\beta}\eta_{\mu\nu} - \eta_{\alpha\nu}\eta_{\beta\mu}) . \quad (5.c)$$

At tree-level, even if negative-norm states are shown to decouple from physical amplitudes, unitarity may be jeopardised by processes governed by the exchange of a charged massive quantum (i.e., B_μ -field exchange)[5, 6]. We then select the Compton-like scattering amplitude to be analysed (Fig.2). If unitarity bound is to be violated, the process of Fig.2 already signals such a violation. The total scattering amplitude, $\mathcal{M}$, may be written according to :

$$\mathcal{M} = \mathcal{M}_{V_4} + \sum_{\substack{v, v' = \text{all} \\ \text{3-vertices}}} (\mathcal{M}_{vv'}^{(1)} + \mathcal{M}_{vv'}^{(2)}) , \quad (6)$$

where v and v' denote all possible 3-vertices in the model and $\mathcal{M}_{V_4}$, $\mathcal{M}_{vv'}^{(1)}$ and $\mathcal{M}_{vv'}^{(2)}$ are the partial amplitudes for the ‘‘Compton’’ scattering graphs presented in Fig.2.

For the calculation of the individual graphs and for the evaluation of the product $\mathcal{M}^*\mathcal{M}$, use has been made of FORM software, since the number of terms generated is extremely high and algebraic manipulations become very cumbersome.

For the CSM*-model, we have found that in the limit of very high center-of-mass energies ($\sqrt{s} \rightarrow \infty$) the squared amplitude behaves according to :

$$\lim_{\sqrt{s} \rightarrow \infty} |\mathcal{M}_{CSM}|^2 = 90 \omega^4 ; \quad (7)$$

this behaviour is not conflicting with unitarity. Nevertheless, for the CSMP*-model, $|\mathcal{M}|^2$ exhibits the following asymptotic behaviour :

$$\lim_{\sqrt{s} \rightarrow \infty} |\mathcal{M}_{CSMP}|^2 = \frac{\omega^4}{4\hat{\mu}^4} s^2 , \quad (8)$$

for both quanta, m_+^2 and m_-^2 . Then, one concludes by (8) that unitarity is clearly violated, since the probability amplitude increases with the quartic power of the center-of-mass energy.

In order to justify such a conclusion, we refer to the works of Chaichian *et al.* [7, 8], where upper-limits are derived for the two-particle scattering cross-section in a $D(\geq 3)$ -dimensional space-time [8, 9]

$$\lim_{\sqrt{s} \rightarrow \infty} \sigma_{tot} \leq c s^{\frac{(D-2)}{2}} (\ln s)^{D-2} . \quad (9)$$

Bearing in mind the optical theorem [10], we can make use of the limit (9) to read the asymptotic behaviour of $|\mathcal{M}|^2$ as dictated by unitarity. In $D = 3$, it turns out to be

$$\lim_{\sqrt{s} \rightarrow \infty} |\mathcal{M}|^2 \leq c s^{\frac{3}{2}} \ln s. \quad (10)$$

Therefore, the asymptotic behaviour given in eq.(8) is in conflict with the unitarity bound typical of 3 space-time dimensions.

It would be however interesting to check if it is possible, as it happens in $D = 4$ [6], to introduce a gauge-invariant non-minimal coupling that might restore Froissart bound at tree-level.

The most general gauge-invariant non-minimal coupling that respects renormalisability in $D = 3$ reads :

$$\mathcal{L}_g = ig F^{\mu\nu} B_\mu^* B_\nu, \quad (11)$$

where g is a real coupling constant with dimension of $(\text{mass})^{\frac{1}{2}}$. This interacting Lagrangian (11) yields the following 3-vertex Feynman rule (see Fig.1):

$$(\tilde{V}_3)_{\alpha\mu\nu} = ig(l_\alpha \eta_{\nu\mu} - l_\nu \eta_{\alpha\mu}). \quad (12)$$

Reconsidering the Lagrangian (2) and adding to it the Lagrangian $\mathcal{L}_g$ of eq.(11), one obtains, by using the equation (3.b), (6) and all vertex Feynman rules, the following high-energy behaviour for the squared amplitude, $|\mathcal{M}_{CSMP_g}|^2$, associated to ‘‘Compton’’ scattering :

$$\lim_{\sqrt{s} \rightarrow \infty} |\mathcal{M}_{CSMP_g}|^2 = \frac{1}{4\hat{\mu}^4} (\omega^4 - \frac{3}{2}\omega^2 g^2 + g^4) s^2. \quad (13)$$

As already mentionned, the purpose is to eliminate the s^2 -dependence, since it destroys the unitarity. This would be achieved whenever the condition

$$\omega^4 - \frac{3}{2}\omega^2 g^2 + g^4 = 0, \quad (14)$$

is fulfilled. Nevertheless, it can be quickly seen that

$$\omega^4 - \frac{3}{2}\omega^2 g^2 + g^4 > 0, \quad (15)$$

for all non-vanishing real values of ω and g . Then, equation (15) reveals the impossibility of restoring the Froissart bound of the CSMP*-model (2) by the addition of a non-minimal coupling (11) in order to cancel the s^2 -dependence of $|\mathcal{M}_{CSMP}|^2$ (8).

Our final conclusion is that the gauge-invariant CSM*-model discussed in ref.[4] exhibits a well-behaved asymptotic limit for the ‘‘Compton’’ scattering amplitudes at tree-level, even if ghosts do not decouple. On the other hand, the case of the CSMP*-field coupled to an Abelian gauge-field violates Froissart bound in $D = 3$ and, contrary to what happens in the case of 4-dimensional massive charged vector fields coupled to the Maxwell-field, Froissart bound cannot be restored at the expenses of a gauge-invariant non-minimal coupling.

The author expresses his gratitude to Dr. J.A. Helayel-Neto, Prof. M. Chaichian, Dr. O. Piguet, Dr. S.P. Sorella and Dr. F.A.B. Rabelo de Carvalho for patient and

helpful discussions. Thanks are also due to Prof. H.F. de Carvalho and my colleagues of CBPF-DCP for encouragement. CNPq-Brazil is acknowledge for invaluable financial help.

Figure 1: Interaction 3- and 4-vertices.

Figure 2: Compton scattering diagrams $\mathcal{M}_{V_4}$, $\mathcal{M}_{vv'}^{(1)}$ and $\mathcal{M}_{vv'}^{(2)}$.

References

- [1] S. Deser, R. Jackiw and S. Templeton, *Ann.Phys. (N.Y.)* **140** (1982) 372.
- [2] N. Dorey and N. E. Mavromatos, *Phys.Lett.* **B226** (1991) 163 ;
N. Dorey and N. E. Mavromatos, *Nucl.Phys.* **B386** (1992) 614.
- [3] F. Delduc, C. Lucchesi, O. Piguet and S.P. Sorella, *Nucl.Phys.* **B346** (1990) 313 ;
A. Blasi, O. Piguet and S.P. Sorella, *Nucl.Phys.* **B356** (1991) 154 ;
C. Lucchesi and O. Piguet, *Nucl.Phys.* **B381** (1992) 281.
- [4] O.M. Del Cima and F.A.B. Rabelo de Carvalho, *Some Quantum Aspects of Complex Vector Fields with Chern-Simons Term*, CBPF preprint, CBPF-NF-029/93 (1993), *submitted for publication*.
- [5] J.A. Helayël-Neto, O. Piguet and S.P. Sorella, *private communications*.
- [6] R. Finkelstein, *Rev.Mod.Phys.* **36** (1964) 632.
- [7] M. Chaichian and J. Fischer, *Nucl.Phys.* **B303** (1988) 557.
- [8] M. Chaichian, J. Fischer and Yu. S. Vernov, *Nucl.Phys.* **B383** (1992) 151.
- [9] M. Chaichian and J.A. Helayël-Neto, *private communications*.
- [10] R.J. Eden, P.V. Landshoff, D.I. Olive and J.C. Polkinghorne, *The Analytic S-Matrix*, Cambridge Univ. Press (Cambridge, 1966) ;
A.O. Barut, *Scattering Theory*, ed. A.O. Barut, Gordon and Breach Science Publishers (New York, 1969).