CERN-TH. 7313/94

SPACE-TIME DESCRIPTION OF ULTRA-RELATIVISTIC NUCLEAR COLLISIONS IN THE QCD PARTON PICTURE

K. Geiger

CERN, TH-Division, CH-1211 Geneva 23, Switzerland

(submitted to Physics Reports)

ABSTRACT

Recent developments in the dynamical description of the early stage of heavy ion collisions at collider energies on the basis of the improved parton model and kinetic theory are presented. The Parton Cascade Model is discussed which links firmly established perturbative QCD with classical transport theory for multiparton systems to give a comprehensive description of nuclear collisions in complete phase-space. The space-time evolution of these reactions is investigated in terms of the dissipative dynamics of short range (semi)hard quark and gluon interactions, by evolving the partons' phase-space distributions in real time according to a transport equation through multiple collisions, emission and absorption processes within perturbative QCD. The resulting parton cascade picture is applied to study multiparticle production in $p\bar{p}$, AA and pA collisions at collider energies ($\sqrt{s}/\text{nucleon} \gtrsim 100$ GeV), as well as issues of thermalization, chemical equilibration, and properties of plasma signatures in heavy ion collisions at the future RHIC and LHC experiments.

PACS Indices: 25.75.+r, 12.38.Bx, 12.38.Mh, 24.85.+p

e-mail: klaus@surya11.cern.ch

CERN-TH. 7313/94
June 1994

TABLE OF CONTENTS

1. INTRODUCTION: PERSPECTIVES OF NUCLEAR REACTIONS AT COLLIDER ENERGIES
 - 1.1 Space-time picture of ultra-relativistic nucleus-nucleus collisions
 - 1.2 Theoretical approaches to a dynamical description
 - 1.3 Organisation of the review
2. THE PARTON PICTURE OF HADRONIC INTERACTIONS
 - 2.1 Basic concepts
 - 2.2 Criterium for the applicability of the parton description
 - 2.3 The DGLAP evolution equations for the parton distributions
 - 2.4 Parton cascades and formfactors in nucleon-nucleon collisions
 - 2.4.1 Factorization of short- and long-range dynamics.
 - 2.4.2 The formfactor for space-like branchings
 - 2.4.3 The formfactor for time-like branchings
 - 2.5 Further developments of the parton cascade description
3. SEMI-CLASSICAL TRANSPORT THEORY OF MANY-PARTON SYSTEMS IN NUCLEAR COLLISIONS
 - 3.1 Transport equation for the kinetics of many-parton systems
 - 3.2 Physical quantities relating microscopic parton dynamics to macroscopic observables
 - 3.3 Detailed balance and equilibration
 - 3.4 Practical determination of the parton phase-space distributions
 - 3.4.1 Cellular phase-space and finite particle number effects
 - 3.4.2 Sensitivity of time-development to phase-space discretization
4. THE PARTON CASCADE MODEL FOR ULTRA-RELATIVISTIC NUCLEAR COLLISIONS
 - 4.1 The initial state

- 4.1.1 Initial momentum distribution
 - 4.1.2 Initial spatial distribution
 - 4.2 The perturbative evolution of parton cascades
 - 4.2.1 Elementary parton-parton collisions
 - 4.2.2 Virtual and real corrections to the elementary processes
 - 4.2.3 Space-like parton cascades
 - 4.2.4 Time-like parton cascades
 - 4.3 Cluster formation and hadronization
 - 4.3.1 Parton recombination and cluster formation
 - 4.3.2 Hadronization of clusters
 - 4.4 Model parameters from phenomenology and experiment
 - 4.4.1 Fixing $p_{\perp cut}$ from nucleon-nucleon cross-sections
 - 4.4.2 Fixing μ_0 from fragmentation in e^+e^- -annihilation
 - 4.5 Central assumptions and approximations of the model
- 5. MULTIPARTICLE PRODUCTION IN NUCLEAR REACTIONS AT COLLIDER ENERGIES
 - 5.1 $p\bar{p}$ collisions
 - 5.2 AA collisions
 - 5.3 pA collisions
- 6. HEAVY ION COLLISIONS AT RHIC AND LHC: FROM PRE-EQUILIBRIUM TO PARTON PLASMA FORMATION
 - 6.1 Characteristics of the partons' space-time evolution in AA collisions
 - 6.2 Thermalization
 - 6.3 Chemical equilibration
 - 6.4 The "hot glue scenario"
 - 6.5 Dilepton production
 - 6.6 Strangeness, charm, and bottom production
- 7. CONCLUSIONS AND OUTLOOK
- 8. APPENDIX: THE MONTE CARLO PROCEDURE

1. INTRODUCTION: PERSPECTIVES OF NUCLEAR REACTIONS AT COLLIDER ENERGIES

After a head-on collision between two heavy nuclei at an energy of at many GeV/nucleon several hundred or even thousand charged particles are emitted. No one would care to study such events, unless there existed a challenging reason. The current interest in nuclear collisions at very high energies (far above 1 GeV/nucleon in the center-of-mass system) [1, 2] is fueled by the expectation that a *quark gluon plasma* may be created temporarily in these events [3, 4, 5]. Moreover, such ultra-relativistic heavy ion collisions are expected to exhibit new *phenomena of QCD in dense matter*, associated with the microscopic dynamics of quarks and gluons in the hot, ultra-dense environment that may be created in the central collision region of these reactions.

There is general consensus among theorists that QCD at thermodynamic equilibrium exhibits a phase transition from the normal color-confined phase of hadronic matter with broken chiral symmetry to a deconfined, chirally symmetric phase at sufficiently high energy density. However, many aspects of this transition and the properties of QCD in hot, dense matter are still a matter of debate. Such details are actually the crux of the biscuit and concern, e.g., the order of the phase transition; the precise value of the critical energy density, the nature of experimentally observable signatures of the transition, the impact of dense medium effects, and how fast thermal equilibrium conditions are established in nuclear collisions over a sufficiently large space-time volume. These issues require much theoretical and experimental study.

The experimental programs at the future heavy ion (AA) colliders, the BNL Relativistic Heavy Ion Collider (RHIC) with maximum available beam energy $\sqrt{s} = 200$ A GeV (gold on gold) and the CERN Large Hadron Collider (LHC) with $\sqrt{s} = 6000$ A GeV (lead on lead), will provide for the first time the opportunity to study nuclear matter under extreme density compression and very high temperatures. With the advent of these accelerators it is an exciting challenge to recreate and explore aspects of the early universe when the hadron matter to quark gluon plasma transition occurred.

In this report I will give a (somewhat subjective) review on recent developments in the description of ultra-relativistic nuclear collisions on the level of the microscopic dynamics of quarks and gluons (the partons) and their interactions. The emerging picture of the early stage of heavy ion collisions at collider energies on the basis of the QCD improved parton model and kinetic theory will be discussed. The viewpoint advocated here is that perturbative QCD provides a firm basis for the description of the nuclear dynamics in terms of the evolution of the parton distributions in phase-space. I hope this review helps overcoming the still widespread dilution that perturbative QCD processes may be only rare fluctuations that have hardly anything to do with the global reaction dynamics. In fact,

most of the entropy and transverse energy is expected to be produced already within the first fm ¹ after nuclear contact by very frequent parton interactions with only a few GeV momentum transfer [6]. Therefore I believe that the space-time evolution of nuclear collisions at collider energies *must* be investigated in terms of the dissipative parton dynamics within perturbative QCD. To do so, a kinetic description is required that links together concepts of *relativistic transport theory* on the one hand and *renormalization group improved perturbative QCD* on the other hand. Instead of following ambitious attempts to establish a quantum kinetic formulation on QCD's first principles [7, 8, 9] (which up to date lacks applicability to realistic problems), I advocate a pragmatic, but transparent and widely applicable approach: it allows to follow the evolution of the parton phase-space densities in real time through multiple collisions, emission and absorption processes, according to a Lorentz invariant semiclassical transport equation. The resulting parton cascade picture can be applied to study the issues of thermalization, chemical equilibration, and particle production in *AA* collisions at RHIC and LHC. These topics are of great phenomenological interest and will be discussed in detail.

1.1 Space-time picture of ultra-relativistic nucleus-nucleus collisions

Let me briefly recall the general features of the dynamics of ultra-relativistic nucleus-nucleus collisions ($\sqrt{s}/\text{nucleon} \gtrsim 100$ GeV), particularly those involving very heavy nuclei. In such energetic reactions the physics of hadronic matter can be studied at extremely high energy densities [2]. For example, at RHIC a central $^{197}\text{Au} + ^{197}\text{Au}$ collision with $\sqrt{s} = 200 A$ GeV (A denotes the nuclear mass number) harnesses a total energy $E_{tot} = 40$ TeV; at LHC a head-on $^{208}\text{Pb} + ^{208}\text{Pb}$ collision with $\sqrt{s} = 6000 A$ GeV provides $E_{tot} = 1200$ TeV. These values imply initial energy densities at the moment of maximum nuclear overlap of $\varepsilon_i \approx 250$ (7500) GeV/ fm^3 at RHIC (LHC). Comparing this with the critical energy density for deconfinement [2], $\varepsilon_{crit} = 1 - 10$ GeV/ fm^3 , one sees that the collision system will initially be by orders of magnitude in the pure quark-gluon phase. The question is: what happens next?

It is expected that in the center-of-mass frame two highly Lorentz contracted nuclei pass through each other and deposit their kinetic energy gradually behind them [98, 99, 100, 101]. The commonly accepted qualitative space-time evolution of these reactions is schematically illustrated in Fig. 1 and can be summarized as follows. Immediately after the *first instant of nuclear contact*, in the region between the two fast receding discs, the partons of the colliding nuclei begin to interact frequently with each other, which results in a vehement materialization of excited quanta. The excited partons can rescatter and emit new particles and thereby evolve through a *pre-equilibrium* stage towards a thermalized *quark gluon*

¹Natural units $\hbar = c = k_B = 1$ will be employed throughout, except where noted otherwise.

plasma state, from which the system evolves further according to the laws of relativistic hydrodynamics [10, 11]. The plasma expands and cools, and eventually a phase transition - perhaps via a mixed parton-hadron phase - into a purely *hadronic gas* occurs. The *freeze-out* of the hadron matter subsequently produces the final hadron yield escaping from the collision region to the detectors. The time scales of these successive stages are very different. Although numbers are currently still uncertain, expected typical values are (depending on the collision energy) as follows: the pre-equilibrium stage takes about 0.1 - 0.3 *fm* [6, 12], the plasma phase may live for 1 - 5 *fm* [13], and the hadronization process might take 20 - 40 *fm* [14].

Of particular interest is the very early stage in these reactions, namely from initial parton materialization via pre-equilibrium to plasma formation, since it is expected that most of the entropy and transverse energy are produced already here. Moreover, the parton dynamics during the first 1 - 2 *fm* determines the initial and boundary conditions for a hydrodynamic expansion of the plasma phase, so that any characteristic signal for this state is seeded during the pre-equilibrium and thermalization stages. In the past it was commonly assumed that the quanta which are produced in the central rapidity region quickly reach a state of local thermodynamic equilibrium. Such an assumption is the convenient basis of most theoretical studies addressing the properties of a quark gluon plasma in terms of hydrodynamic expansion [11]. However, the validity of this assumption remains to be checked. In particular, one would like to know how the quanta evolve in phase-space, if and under what circumstances they eventually thermalize, the time scale for this thermalization, the typical energy density and entropy soon after equilibration, etc.. Without this knowledge there is the uncertainty in choosing the formation time and initial temperature of the plasma, two parameters which sensitively control the further evolution. To study these questions systematically in a realistic model, a comprehensive description of the dissipative processes occurring during very early stage of high energy nuclear collisions is necessary. A kinetic space-time description based on non-equilibrium kinetic theory is needed: starting from the measured parton structure of the colliding nuclei, a dynamical calculation must evolve the initial partons smoothly from their first hard scattering, through a pre-equilibrium stage, into a thermalized plasma.

1.2 Theoretical approaches to a dynamical description

Within QCD phenomenology there exist basically two kinds of approaches to describe the evolution of high energy nuclear collisions both of which have been extensively investigated in the last years, namely, a) the *string picture* [15, 16] and b) the *parton approach* [17, 18]. The conceptual visualization of these two approaches is illustrated in Fig. 2.

a) *String picture*: Based on phenomenological models of soft hadron-hadron interactions,

the collision of two nuclei is viewed as their interpenetration with only small rapidity loss (of the order of one unit). The wounded nucleons draw color flux tubes (strings) between each other, and these strings subsequently fragment by $q\bar{q}$ production within a typical proper time of $\simeq 1 \text{ fm}$. The production of entropy and transverse energy is thus associated with non-perturbative, phenomenological string dynamics.

- b) *Parton approach*: The description is founded on the fieldtheoretic parton picture of hadronic interactions within the firmly established framework of perturbative QCD. The colliding nuclei are visualized as clouds of quasireal quarks and gluons which penetrate through each other, leading to rapid materialization. Multiple short range scatterings between partons (minijet production) together with associated QCD radiation (gluon bremsstrahlung) produce the major part of the total entropy and transverse energy.

Both approaches have their conceptual problems and limitations. The *string picture* becomes questionable at very high energies, when the string density becomes large and the strings begin to overlap, so that they cannot be treated as individual entities that fragment independently. Concerning the issue of quark gluon plasma formation, it appears inconsistent to assume that the establishment of a truly perturbative QCD phase should proceed via non-perturbative dynamics. The *parton approach* on the other hand becomes invalid at lower beam energies, where most parton scatterings involve momentum transfers too small to be described by perturbation theory. Long range color forces are assumed to be neglectable, so that non-perturbative collective effects and color correlations are (at this point) beyond the scope of this description. One might therefore view the string picture and the parton approach as complementary to each other.

In view of the ultra-relativistic collider energies available at RHIC and LHC the parton approach appears to be more realistic than the string picture. It has been realized that short range parton interactions play the major role for the nuclear dynamics at collider energies [19] - at least during the early and most dissipative stage of the first few fm . Here copiously produced minijets [20, 21] cannot be considered as isolated rare events, but must be embedded in complicated multiple cascade-type processes. At the same time it is found [22] that color correlations among the initial partons randomize so rapidly as the nuclei interpenetrate, such that the long range color field effectively vanishes on a space-time scale of a small fraction of a fm . Thus, the short range character of the interactions implies that perturbative QCD can and must be used, and that for example the string picture does not apply anymore. From a more theoretical point of view, one may say that the parton approach is based on the explicit form of the QCD interaction between quarks and gluons, whereas on the contrary the string picture utilizes the QCD terminology and the QCD

degrees of freedom rather than the explicit form of the QCD Lagrangian.

In the framework of hadron fragmentation within the string picture numerous Monte Carlo event generators have been developed over the years, FRITIOF [23], ATILA [24], VENUS [25], RQMD [26], QGSM [27], SPM, [28], SPACER [29], and others (see e.g. Ref. [25] for a comparative overview). On the basis of the perturbative QCD parton model there exist many event generators to describe *hadronic collisions* at high energies, to mention a few, PYTHIA [30], HERWIG [31], FIELDJET [34], COJET [32], EUROJET [33], ISAJET [35], and others (see e.g. Ref. [36] for a comparison of these "high $p_\perp$ " Monte Carlo models).

Much fewer attempts have been made so far to simulate *nuclear collisions* at collider energies on the level of microscopic parton dynamics. A pioneer work was due to Boal [37] who for the first time computed with a numerical simulation the time evolution of the parton distributions in nucleus-nucleus collisions. In the same spirit, later B. Müller and myself developed the parton cascade model (PCM) [38, 39], which, implemented in a Monte Carlo program, provides a kinetic space-time description of parton evolution by utilizing refined techniques for perturbative QCD calculations, developed in high energy particle physics. A very related approach is the HIJING model [40, 41], that also incorporates the perturbative QCD approach to multiple minijet production, however it does not incorporate a direct space-time description. Also to mention in this context is DTUNUC [42] and the aforementioned VENUS model, which, although primarily based on the dual parton model [43], include some aspects of the perturbative QCD parton picture.

1.3 Organisation of the review

The remainder of this review is structured as follows. In Chapter 2 the conceptual ideas the parton picture of high energy hadronic interactions is briefly reviewed and the relevant formalism is introduced. Emphasis is put on the probabilistic parton language that provides an intuitive, almost classical approach to the description of hadronic and nuclear collisions. The very successful perturbative QCD description of many high energy physics aspects on the basis of multiplicative parton cascade evolution is illustrated and exploited as motivation for extending this approach to nuclear collisions.

Chapter 3 is devoted to introduce the key elements of relativistic kinetic theory necessary to generalize the momentum space description of the parton picture to a time evolution in full phase-space on the basis of a relativistic invariant transport equation for the parton distributions. The relation between microscopic parton dynamics and macroscopic thermodynamic observables is established. Also, issues concerning detailed balance of parton interactions and equilibration of a many-parton system are discussed.

Chapter 4 forms the core of the review. Here the PCM is explained in great detail. The

concepts of Chapters 2 and 3 are joined to establish a kinetic description of quark-gluon transport phenomena in hadron-hadron, hadron-nucleus, and nucleus-nucleus collisions at highly relativistic energies. The probabilistic parton cascade picture is translated to the space-time evolution of the parton distributions according to the relativistic version of the Uehling-Uhlenbeck equation, the collision kernel of which incorporates the specific perturbative QCD structure of parton interactions. The collision kernel is represented by effective matrix-elements that correspond to a set of elementary $2 \rightarrow 2$, $1 \rightarrow 2$ and $2 \rightarrow 1$ processes plus inclusively summed higher order corrections that are associated with real and virtual emission and absorption of partons. Embedded in the transport equation, this collision kernel generates the space-time evolution of nuclear collisions in terms of many interconnected parton cascades that consist of multiple parton scatterings plus associated emission and absorption processes. The parton cascade evolution is supplemented by additional phenomenological elements accounting for nuclear and medium effects, and the (separate) hadronization scheme that converts the final parton configuration into the asymptotic hadronic states.

In Chapter 5 the PCM approach is applied to investigate multiparticle production in nuclear collisions at collider energies. The model is first tested in the case of proton-antiproton ($p\bar{p}$) collisions for which exists a wide variety of experimental data to compare to. It follows an analysis of heavy ion (AA) and proton-nucleus (pA) collisions at RHIC and LHC energies, in which calculated predictions for particle spectra and multiplicities are presented. Experimental measures that may reflect characteristic nuclear and dense medium effects of the parton dynamics associated with "QCD in medium" are also discussed.

Chapter 6 deals with the expected thermodynamic behavior of the parton distributions in high energy heavy ion collisions. With regard to the fundamental interest in the the dynamics of quark gluon plasma formation and QCD phase transition, the time evolution of the many-parton system in central AA collisions is followed in phase-space from the first instant of collision, via pre-equilibrium, towards plasma formation. Specific properties of the equilibration process, such as relaxation time scales, time evolution of temperature, energy density, entropy production, and chemical composition are extracted and analysed. It follows a discussion of the emerging understanding of this early stage in heavy ion reactions, that is characterized by gluon dominated matter, being considerably hotter and denser as anticipated in the past. The optimistic implications for observables that should allow for an unambiguous detection of parton equilibration and plasma evolution, are exemplified by investigating lepton pair emission, as well as strangeness and charm production, from cascading partons during the pre-equilibrium and thermalization stages.

Finally, Chapter 7 is reserved for concluding remarks and a perspective outlook. A summarizing list is assembled of the main problems and open questions connected to the

PCM approach, which should be solved or at least better understood in the nearest future. I hope this will be useful to the reader and can balance to some extent the rather optimistic viewpoint of the author.

In the Appendix a more technical summary is given of the Monte Carlo generation procedure that was employed to "solve" the transport equation for the parton distributions by simulating nuclear collisions on the computer. The computer program of the PCM is obtainable from the author.

2. THE PARTON PICTURE OF HADRONIC INTERACTIONS

In this Chapter I will review the essential ideas and concepts that underly the field-theoretic description of strong interactions in the parton picture at high energies. The dynamical evolution of the parton distributions in terms of a probabilistic parton cascade description is exemplified for the case of nucleon-nucleon collisions as a prototype reaction for the extension to nucleus-nucleus collisions later on.

2.1 Basic concepts

It is the very successful description of perturbative QCD of the dynamics of relativistic hadronic collisions with numerous correct quantitative predictions (see e.g. Refs. [44, 45, 46]) that provides an encouraging motivation to extend and apply the parton model ideas [47, 48, 49] to hadron-nucleus and nucleus-nucleus collisions. Particularly the observation [17, 18] that the major part of the total nucleon-nucleon cross-section arises from the large probability for the production of quark and gluon "minijets" [20, 21] carrying only a small fraction of the initial energy has lead to the point of view that perturbative QCD provides a good basis for a global description of hadronic interactions and not only of rare fluctuations in the hadron wave function.

Pioneered to large extent by the Leningrad group many years ago (see reviews [17, 18]), an appropriate description of hadronic interactions at high energy was developed that is based on the microscopic dynamics of parton cascades within renormalization group improved perturbative QCD. The basic idea is that a fast hadron may be considered as a *coherent* cloud of partons with life times much larger than the typical interaction times (the time needed for the interaction with another particle). Each parton can be characterized by its type, the fraction x of the initial hadron energy which it carries, and by its transverse momentum $q_{\perp}$. Furthermore, the fact that these partons are confined inside the hadron and cannot exist as "free particles" implies that they are off-shell by some (small) *space-like virtuality* $q^2 = q_0^2 - (\vec{q}^2 + m_0^2) < 0$, where m_0 represents the rest mass. Thus the invariant mass q^2 is an additional independent variable. At any instant an individual parton can initiate a *space-like cascade* by branching into a number of partons with smaller x as

illustrated in Fig. 3a. The global probability for a branching of a parton with momentum q_{j-1} leading to the emission of parton with q_j ,

$$P_j \propto \int_{q_{j-1}^2}^{q_{j+1}^2} \frac{dq_j^2}{q_j^2} \frac{\alpha_s(q_j^2)}{\pi} \int_{x_{j+1}}^{x_{j-1}} \frac{dx_j}{x_j}, \quad (1)$$

can be of order unity, because the available phase-space $d\Omega_j \propto dx_j/x_j dq_j^2/(4\pi q_j^2)$ is very large so that the smallness of the coupling constant $\alpha_s(q_j^2) \ll 1$ at the branching vertex is compensated by the logarithmically large integration over the momentum of the daughter partons. A cascade such as depicted in Fig. 3a can be described in terms of a tree-like structure, composed of many branchings. First the partons in a cascade multiply, then they assemble back due to their coherence (since they do not have enough energy to exist as free particles). Thus, such a space-like cascade lives for a finite time, but new cascades are born to replace the former ones. One may interpret these branching mechanisms as fluctuations in the coherent hadron wave function.

However, if a parton in a cascade collides on its way with some "external" particle (e.g., a photon in deep inelastic scattering, or another parton in a hadron-hadron collision) then this interaction destroys the coherence of the cascade, as illustrated in Fig. 3b. The external particle acts like a probe that resolves the parton structure of the cascade down to a space-time extent $\Delta r \simeq 1/\sqrt{Q^2}$ where Q^2 is the momentum transfer of the interaction. The parton with space-like four-momentum q is struck out of the cascade and becomes a "real" excitation with time-like four-momentum p , because the interaction provides it with the necessary energy to get on mass shell or even acquire a *time-like virtuality* $m^2 \equiv p_0^2 - (\vec{p}^2 + m_0^2) > 0$, where m_0 denotes the rest mass of the parton. As a result the other partons in this particular cascade branch cannot reassemble, but continue to live. Due to their excitation, each of these secondary partons can now initiate a *time-like cascade* by emitting further partons through gluon bremsstrahlung or $q\bar{q}$ production. As a consequence the parton density grows locally, leading (in e.g. hadron-hadron collisions) to an increase of the cross-section and thus an enhancement of the probability for further parton scatterings accompanied by particle emission. It is evident that this breaking of initial coherence and the subsequent multiplicative cascading is the elementary source of entropy production in the picture, since the produced partons will at some point transform into secondary hadrons so that the parton multiplicity is the direct source of the experimentally measured particle yield.

2.2 Criterium for the applicability of the parton description

Since the secondary partons that are produced during the course of the cascade evolution must at some point convert into hadrons, an important question is, how do these off-spring partons (mostly gluons) influence the hadronic yield. To answer this, one has to realize

first what is the condition for a virtual parton to behave as an independent colored object and thus as an additional source of new particles. In a time-dependent description, the *formation time*, i.e. the time it takes for a virtual parton to emit a gluon, controls when this gluon can be considered as irreversibly emitted. Following [50], it can be simply estimated from the proper life-time $\tau_a \simeq 1/m_a^{virt}$ of a virtual parton a with invariant mass $m_a^{virt} = \sqrt{|q_a^2|}$ to decay by gluon emission into a parton b and a gluon c (a more rigorous argument is given in [51]). If parton a has momentum q_a , parton b and gluon c carry momenta $q_b = q_a - q$ and $q_c = q$, respectively then the uncertainty relation with account for time dilation effects gives for the gluon formation time in the laboratory frame:

$$t_g^{form} \simeq \tau_a \gamma = \frac{1}{m_a^{virt}} \frac{E_a}{m_a^{virt}} = \frac{E_a}{q_a^2} \approx \frac{E_a}{q E_a \theta^2} \approx \frac{q}{q_\perp^2} , \quad (2)$$

where θ is the angle between b and c and q now denotes the absolute value $\sqrt{|q^2|}$. Comparing this expression with the typical hadronization time of a gluon [50]

$$t_g^{had} \approx q R^2 , \quad (3)$$

(here R , the nucleon radius, sets the typical hadronic scale), one can conclude that it is the transverse momentum restriction

$$q_\perp \stackrel{!}{>} R^{-1} = \text{a few hundred MeV} \quad (4)$$

which guarantees the gluon's being, since it must be produced before it hadronizes:

$$t_g^{form} \sim \frac{q}{q_\perp^2} < t_g^{had} \sim q R^2 \quad (5)$$

The same reasonings can be applied to any parton branching process, so that the basic restriction (4) represents the *condition of the applicability of the parton language*, that is of the perturbative QCD description. Note that there exists another way of argumentation which approves this restriction as a natural limit for the perturbative approach: at finite values of q or $q_\perp$ the running coupling α_s in the expression (1) for the emission probability approaches the infrared pole, thus limiting the perturbative description at small momentum q or $q_\perp$. The boundary condition (4) being physically very important, leaves at the same time rather broad room for a perturbative analysis. In inelastic processes (e.g. deep inelastic lepton-nucleon scattering or high energy hadronic collisions) with a large momentum transfer Q the bulk of phase-space where the parton emission amplitudes are logarithmically large [c.f. eq. (1)],

$$R^{-1} \ll q_\perp \lesssim q \lesssim \sqrt{Q^2} , \quad (6)$$

remains in the regime of perturbative QCD. The parameter $q_\perp R$ can be said to regulate the "life-time" of a secondary parton. Therefore the main characters of the parton cascade play are those quanta with momenta satisfying the strong inequality

$$q_\perp R \gg 1, \quad (7)$$

since they will live for a long time, radiating in their turn new off-spring, and thus developing the cascade multiplication of partons.

To gain a large total emission probability (1) in spite of α_s restricted to be small, one has to pick up at least one logarithmic integration out of the phase-space. For completeness, let me list the possible orderings in eq. (6), each of which reveals itself in definite physical phenomena [50]:

- (i) $R^{-1} \ll q_\perp \ll q \sim \sqrt{Q^2}$: quasi-collinear hard parton branchings, leading to the well known scaling violations of the parton distributions;
- (ii) $R^{-1} \ll q_\perp \sim q \ll \sqrt{Q^2}$: large angle soft gluon emission, responsible for coherence properties of interfering gluons;
- (iii) $R^{-1} \ll q_\perp \ll q \ll \sqrt{Q^2}$: double-logarithmic (soft + collinear) gluon emission, causing strong multiplicity growth with energy and determining the QCD formfactors of partons.

2.3 The DGLAP evolution equations for the parton distributions

To quantify the description of this specific QCD parton multiplication an approach is required that accounts for the many particle aspect with the detailed structure of parton cascades. Starting from the above conceptual idea of representing a high energy nucleon as a semiclassical assembly of partons with varying number, confined inside the nucleon, one employs the statistical analogue of the wave function in momentum space, namely the Wigner function or phase-space distribution function of the partons with respect to their momenta. Let me distinguish in *space-like* and *time-like* partons and parametrize the four-momenta $k \equiv k^\mu = (k_0, k_z, \vec{k}_\perp)$ with $k \equiv q, p$ as

$$\begin{aligned} q &= \left(xP + \frac{q^2 + q_\perp^2}{2xP}; xP, \vec{q}_\perp \right), \quad Q^2 \equiv -q^2 \quad (q^2 = q_\mu q^\mu < 0, \text{ space-like}) \\ p &= \left(xP + \frac{m^2 + p_\perp^2}{2xP}; xP, \vec{p}_\perp \right), \quad Q^2 \equiv m^2 \quad (m^2 = p_\mu p^\mu > 0, \text{ time-like}), \end{aligned} \quad (8)$$

where P denotes the nucleon momentum assuming $P/M_N \gg 1$ (M_N the nucleon mass). Here the partons' rest masses are neglected and the space-like (time-like) virtualities

$Q^2 = -q^2$ ($Q^2 = m^2$) must satisfy, according to Sec. 2.1, $Q^2 \ll P^2$ in order that the parton approach is applicable. It is important to realize that energy and momentum are independent variables, because the partons are represented as off-shell particles of virtuality Q^2 with a continous mass spectrum.

The distribution of partons in longitudinal momentum (or energy) fraction x and virtuality Q^2 is the given by the transverse momentum integrated *number densities* of quarks q_i and antiquarks $\bar{q}_i$ of flavor i , or gluons g , between Q_0^2 and $Q^2 > Q_0^2$,

$$f_a(x, Q^2, Q_0^2) = \int_{Q_0^2}^{Q^2} dQ'^2 \int dk_\perp^2 \frac{dN_a}{dx dk_\perp^2 dQ'^2} , \quad (a \equiv q_i, \bar{q}_i, g) , \quad (9)$$

where N_a labels the number of partons of type a and the mass scale $Q_0 \gtrsim 1$ GeV is introduced to characterize the minimum value of virtuality Q^2 and to set the lower boundary for the perturbative description, requiring $\alpha_s(Q_0^2) \ll 1$. The experimentally measured *nucleon structure functions* can then formally be represented as the convolution of a hadronic structure function $h_N^a(x, Q_0^2)$ with a partonic structure function $f_a(x, Q^2, Q_0^2)$ [52]:

$$F_a^N(x, Q^2) \equiv \int_0^1 \frac{dy}{y} h_a^N(y, Q_0^2) f_a\left(\frac{x}{y}, Q^2, Q_0^2\right) , \quad (10)$$

expressing the probability for finding a parton of type a inside a nucleon N , carrying a fraction x of the total nucleon momentum when the nucleon is probed at a scale $Q^2 > Q_0^2$. The function $h_a^N(x, Q_0^2)$ embodies the non-perturbative dynamics prior to Q_0^2 , which cannot be predicted in perturbative QCD, but has to be extracted from experimental data of e.g. deep inelastic lepton-nucleon scattering. Thus, the absolute value the structure functions depends on the soft dynamics below Q_0^2 which must be absorbed in the function $h_a^N(x, Q_0^2)$ by parametrization of structure function measurements corresponding to a certain value of Q_0^2 . However, the change of the structure functions with varying Q^2 is solely described by $f_a(x, Q^2, Q_0^2)$ and is rigorously calculable by perturbative methods. Hence, with given initial values of h_N^a at Q_0^2 , the structure functions F_a^N can be obtained for any $Q^2 > Q_0^2$ by evolving the parton densities f_a in Q^2 . This is illustrated in Fig. 4.

The Q^2 -dependence of the structure functions (10) in connection with multiplicative parton cascades was first addressed many years ago in the *Leading Logarithmic Approximation* (LLA) of QCD [53, 54, 55, 56, 57, 58, 59]. The observation that the branching probability (1) involves a logarithmic distribution of the parton virtualities $\propto dq^2/q^2 = dQ^2/Q^2$ has lead to the logic to select and resum Feynman diagrams that contain the maximum power of $\ln Q^2$ in every order of perturbation theory. More accurately one requires

$$\frac{\alpha_s(Q_0^2)}{\pi} \ll 1 , \quad \frac{\alpha_s(Q_0^2)}{\pi} \ln \left(\frac{Q^2}{Q_0^2} \right) \simeq 1 . \quad (11)$$

where the running QCD coupling strength in one-loop order is given by

$$\alpha_s(Q^2) = \frac{12\pi}{(11n_c - 2n_f)\ln(Q^2/\Lambda^2)} \quad , \quad (12)$$

with $n_c = 3$ color degrees of freedom, and n_f the number of quark flavors that can be probed at scale Q^2 (Λ is the QCD renormalization scale).

On the basis of the requirement (11), the LLA framework keeps trace of contributions to the structure functions of the order

$$F_a^N(x, Q^2) \propto f_a(x, Q^2, Q_0^2) = \sum_{n=0}^{\infty} g_n(x) \left[\frac{\alpha_s}{\pi} \ln \left(\frac{Q^2}{Q_0^2} \right) \right]^n \quad , \quad (13)$$

while systematically neglecting corrections of relative order α_s that are *not* accompanied by a large log. In QCD, being a field theory with asymptotic freedom, such an approximation proves to be *asymptotically exact* as $Q^2 \rightarrow \infty$ [54].

The interpretation of the series representation (13) in terms of a probabilistic evolution of parton cascades with sequential branchings was first given by Lipatov [55, 60]. To obtain large logs in each subsequent branching, the momenta q_j^2 of the partons in each genealogical branch of the cascade must be strongly ² ordered:

$$\dots \ll q_{j-1}^2 \ll q_j^2 \ll \dots \ll Q^2 \quad ; \quad \dots \gg x_{j-1} \gg x_j \gg \dots \gg Q^2/P^2 \quad . \quad (14)$$

The ordering (14) leads to a picture, illustrated in Fig. 5, of successive fluctuations, one enclosed into the other: the parton q_j , being the "bare" particle from the point of view of its parent q_{j-1} , is a "dressed" particle - the parent of partons of the level q_{j+1} . The probability to find the parton q_j is the probability of finding a parton with momentum q_1 in the nucleon, times the probability that it branches into a pair of partons with q_2 and q'_2 , times the probability of the branching of q_2 into q_3 and q'_3 etc., up to the branching that yields q_j . Such a probabilistic (semiclassical) description proves valid, since in the LLA the ladder type diagrams (Fig. 6a) alone generate the branching cascade, whereas the interference of different diagrams (Fig. 6b), which violates the probabilistic picture, always leads to losing one or more logarithmically large regions over the integration over the intermediate parton momenta. The contribution of such interference processes is negligible at high energies, because they are of the order $\alpha_s \ll 1$, uncompensated by a large log. As a consequence, the density matrix of the many parton system is approximately diagonal so that a parton cascade can be described by the incoherent sum of *squared* amplitudes (probabilities) for elementary branchings in between which the intermediate partons are

²The *strong* ordering will be relaxed later and replaced by the much less restrictive *angular* ordering that extends the description to "soft" branching processes.

dressed with a specific formfactor (Fig. 6c). This fact provides the *property of locality* (i.e. no remembrance of prehistory) inherent to a probabilistic picture.

The essential implication is that when probing a nucleon at a larger Q^2 one finds an increase in resolution of the parton cascade that is being struck, i.e. a more detailed parton composition in the nucleon is revealed, as is shown in Fig. 7. At fixed value of x the total number of partons in a hadron increases with $Q^2 = |q^2|$ and their average transverse size decreases as $1/\sqrt{Q^2}$. However, only a rare fraction of these partons has large x ; due to the branching mechanism most of the partons are likely to be found at rather small values of x . This property is expressed by the Dokshitzer-Gribov-Lipatov-Altarelli-Parisi (DGLAP) equations [54, 55, 61, 62] for the Q^2 -evolution of the parton distributions (10):

$$Q^2 \frac{\partial}{\partial Q^2} F_b(x, Q^2) = \frac{\alpha_s(Q^2)}{2\pi} \sum_a \int_0^1 \frac{dz}{z} \left[F_a\left(\frac{x}{z}, Q^2\right) - d_{ac} F_b\left(x, Q^2\right) \right] \gamma_{a \rightarrow bc}(z). \quad (15)$$

Here $x_b \equiv x$, $x_a \equiv x/z$, and the variable z is the fraction of x -values of daughter to mother partons in the branching $a \rightarrow bc$, and the $\gamma_{a \rightarrow bc}(z)$ denote the usual branching functions [61, 62] for the three possible processes $q \rightarrow qg$, $g \rightarrow gg$, $g \rightarrow q\bar{q}$, which describe the energy distribution at the vertex,

$$\begin{aligned} \gamma_{q \rightarrow qg}(z) &= C_Q \left(\frac{1+z^2}{1-z} \right) \\ \gamma_{q \rightarrow gq}(z) &= C_Q \left(\frac{1+(1-z)^2}{z} \right) \\ \gamma_{g \rightarrow gg}(z) &= 2C_G \left(z(1-z) + \frac{z}{1-z} + \frac{1-z}{z} \right) \\ \gamma_{g \rightarrow q\bar{q}}(z) &= \frac{1}{2} (z^2 + (1-z)^2), \end{aligned} \quad (16)$$

where $C_Q = \frac{n_c^2-1}{2n_c} = 4/3$ and $C_G = n_c = 3$. Note that for a given particle type b the sum over a in (15) automatically fixes c . The factor $d_{ac} \equiv (1 + \delta_{ac})^{-1}$ in the second term on the right side of (15) accounts for the identical particle effect if the outgoing partons are indistinguishable (affecting only the $g \rightarrow gg$ process).

The above representation of the DGLAP evolution equations (15) exhibits clearly the balance of gain and loss terms on the right hand side, both of which individually diverge as $z \rightarrow 0$, but the sum of gain and loss terms cancels the soft singularity at $z = 0$ and renders the equations – and consequently the measurable parton distributions – finite. Fig. 8 illustrates this complete probabilistic picture of parton evolution [63]. For simplicity the diagram shows only three quark lines a, b, c . Suppose these partons represent part of the substructure of a hadron. When the hadron is probed by e.g. a virtual photon at scale Q^2 , the number of quarks seen by the probe with momentum fraction x to $x + dx$ is

$F_q(x, Q^2)dx$. When the scale changes from Q^2 to $Q^2 + \Delta Q^2$, the number of quarks seen by the probe changes to $F_q(x, Q^2 + \Delta Q^2)dx$. The difference between $F_q(x, Q^2 + \Delta Q^2)dx$ and $F_q(x, Q^2)dx$ comes from two contributions. A positive contribution arises when a new quark a with x is gained after the quark a with $x_1 > x$ at Q^2 has emitted a gluon. A negative contribution arises when quark b with x is lost due to its gluon radiation and has afterwards $x_2 < x$. Eq. (15) is nothing but the formal copy of this intuitive picture.

2.4 Parton cascades and formfactors in nucleon-nucleon collisions

It is useful to demonstrate the application of the probabilistic parton cascade description for the dynamical evolution of the parton distributions in Q^2 and x according to the evolution equation (15) with a specific example, namely a nucleon-nucleon collision which serves as a prototype reaction for the extension to nucleus-nucleus collisions later on.

2.4.1 Factorization of short- and long-range dynamics

To be specific, consider a proton-antiproton ($p\bar{p}$) collision at ultra-relativistic energy, e.g. at the CERN $p\bar{p}$ collider. In order to apply the parton picture to this reaction, one has to go into a frame where both the colliding nucleons are moving very fast, so that they can be resolved into individual partons. The visualization of a nucleon as an instantaneous distribution of partons at any time requires probing the nucleon over a time durations and spatial distance small on the scale of internal motions of the partons. This condition is fulfilled in any frame of reference in which the nucleon moves almost with the speed of light, because the time dilation effect slows the internal motions such that the wavefunction of the nucleon can be described as a simple quantum mechanical ensemble of quasireal partons that do not mix with vacuum fluctuations (except for the slowest gluons and sea quarks) [17]. Therefore the appropriate choice of Lorentz frame is the center-of-mass (CM) frame of the two nucleons such that

$$P_z^{(p)} = +P, \quad P_z^{(\bar{p})} = -P, \quad \vec{P}_\perp^{(p)} = \vec{P}_\perp^{(\bar{p})} = \vec{0}, \quad (17)$$

so that $\sqrt{s_{p\bar{p}}} = 2P \gg M_N$. Then the parton description the reaction can be factorized in three steps [52], as depicted in Fig. 9:

- 1) The initial parton distributions in the incoming p and $\bar{p}$ given by the measured nucleon structure functions (10) at the initial resolution scale Q_0^2 .
- 2) The perturbative evolution of the ensemble of partons during the course of the collision in terms of parton cascades.
- 3) The non-perturbative hadronization process, e.g. by recombination or fragmentation of the partons into final hadron states.

This factorization [64] rests upon two physical concepts: the Lorentz contraction and time dilation of internal states of the nucleon, and the long-time nature of hadronization. Both the "initial state" interactions between partons associated with the range color correlations, and the "final state" interactions between hadronizing fragments occur at space-time distances too large to affect the basic parton-parton scattering, and hence the inclusive cross-section. The elementary scattering is therefore directly proportional to the density of partons, which is frozen over the short scattering time scale $\approx 1/\sqrt{Q^2}$, set by the momentum transfer Q^2 . The parton density at a particular Q^2 can be obtained from the DGLAP equations (15), given the initial structure functions at Q_0^2 , by space-like cascade evolution. The additional increase of the parton density after the scattering due to gluon bremsstrahlung or $q\bar{q}$ production, until hadronization sets in around μ_0^2 , can be similarly described by time-like cascade evolution.

As I will explain now, the perturbative cascade evolution [step 2)] within the LLA framework means that the partons are not point-like particles, but rather of "extended size" with an internal substructure. This fieldtheoretic property can be represented by specific space-like and time-like formfactors that describe the partons as states with a fluctuating number of cascading daughter partons.

2.4.2 The formfactor for space-like branchings

The LLA approach allows build up the parton distributions of the incoming p and $\bar{p}$ by iterating elementary $1 \rightarrow 2$ parton branching processes, given the structure functions (10) at some small Q_0^2 . After the two colliding nucleons overlap, the partons may scatter off each other with a probability determined by their cross-section. Suppose a parton belonging to the p collides with another parton originating from the $\bar{p}$ with a momentum transfer $Q^2 > Q_0^2$, as in Fig. 9, where 'HS' denotes the hard scattering. Physically the scale Q^2 specifies how detailed the parton of the p resolves the substructure of the $\bar{p}$ and vice versa for parton of the $\bar{p}$. In other words, the scattering selects a certain Q^2 up to which the two scattering partons must have evolved by successively increasing their space-like virtualities up to $|q^2| \approx Q^2$, starting from the initial scale Q_0^2 . Recall from Sec. 2.2 that the amplitude squared for such a sequence of consecutive branchings when evaluated in a physical (any axial) gauge, can be represented diagrammatically by a simple ladder diagram as in Fig. 10a, where the virtualities of the intermediate partons are strongly ordered according to

$$Q_0^2 \leq |q_n^2| \ll \dots \ll |q_j^2| \ll \dots \ll |q_1|^2 \equiv |q^2| \lesssim Q^2. \quad (18)$$

Correspondingly the amplitude itself has the tree structure shown in Fig. 10b, consisting of a series of individual branchings $a \rightarrow bc$, each of which can be interpreted as a decay probability for a parton a of virtual mass q_a^2 to branch into two others, b and c , carrying fractions z and $(1 - z)$, respectively, of the momentum of a , and having masses $|q_{b,c}^2| > |q_a^2|$

[52, 65]. In the frame where $\vec{q}_a$ is along the a -axis, the kinematics of a single branching (Fig. 10c) yield the constraint ($q_{a,b}^2 < 0, q_c^2 > 0$):

$$-\frac{q_a^2}{z} = -q_b^2 + \frac{q_c^2}{1-z} + \frac{q_\perp^2}{z(1-z)}, \quad (19)$$

where $\pm \vec{q}_\perp$ are the momentum components of b and c transverse to the z -axis.

The probability for a single branching process is given by the branching functions $\gamma_{a \rightarrow bc}(z)$, given in (16). In the LLA the probability of a sequence of n branchings becomes the product of probabilities for each single branching. The evolution of a space-like cascade is governed by the same DGLAP equations (15), which describe the Q^2 -evolution of the structure functions $F_a(x, Q^2)$. The equations express that during a small increase dQ^2 in Q^2 , there is a probability for a parton a with momentum fraction x_a of its parent nucleon to become resolved into a parton b at $x_b = zx_a$ and a parton c at $x_c = x_a - x_b = (1-z)x_a$. Correspondingly, during a decrease dQ^2 a parton b may be "unresolved" into a parton a , i.e. when going to smaller values of Q^2 . The relative probability dP_b for this to occur is given by dF_b/F_b [66] which, using eq. (15), becomes

$$dP^{(s)} = \frac{dF_b(x_b, Q^2)}{F_b(x_b, Q^2)} = \frac{dQ^2}{Q^2} \frac{\alpha_s(Q^2)}{2\pi} \sum_a \int_{x_b}^1 \frac{dx_a}{x_a} \frac{F_a(x_a, Q^2)}{F_b(x_b, Q^2)} \gamma_{a \rightarrow bc} \left(\frac{x_b}{x_a} \right), \quad (20)$$

Summing up the cumulative effect of small changes $d \ln Q^2$, the probability for no branching to happen exponentates. Therefore one may define a formfactor

$$S_b(x_b, Q^2, Q_0^2) = \exp \left\{ - \int_{Q_0^2}^{Q^2} \frac{dQ'^2}{Q'^2} \frac{\alpha_s(Q'^2)}{2\pi} \sum_a \int_{z_{\min}(Q'^2)}^{z_{\max}(Q'^2)} \frac{dz}{z} \frac{F_a(x_a, Q'^2)}{F_b(x_b, Q'^2)} \gamma_{a \rightarrow bc}(z) \right\} \quad (21)$$

where $z = x_b/x_a$ and $Q^2 \simeq Q_b^2$ is identified with the scale of the scattering vertex of parton b and the values of $z_{\min}, z_{\max}$ are determined by the branching kinematics via (19). S is called the *Sudakov formfactor for space-like branchings* ('SB' in Fig. 9), giving the probability that parton b has *not* originated from a preceding branching $a \rightarrow bc$ between Q_b^2 and $Q_0^2 < Q_b^2 \simeq Q^2$. Conversely, $1 - S$ is the probability that parton b stems from the previous branching of parton a , which itself may have come from a still earlier branching, and so forth. This "backward evolution" [66, 67, 68] of the space-like cascade reconstructs the parton virtualities from the q_b^2 value implied by the scattering of the primary parton b back to the initial resolution scale Q_0^2 and takes into account the Q^2 evolution of the initial p and $\bar{p}$ structure functions.

2.4.3 The formfactor for time-like branchings

If the momentum transfer in the scattering in Fig. 9 provides one or both of the scattered partons with more energy than is necessary to be on mass shell, they acquire a time-like

virtual mass $m^2 > 0$ and may initiate a sequence of time-like branchings. That is, they "de-excite" by decreasing their time-like virtuality from the value $m^2 \approx Q^2$ obtained at the scattering vertex downwards via emission of a number of partons. This process of successive branchings can be described in a way similar to what was said about space-like branchings before. Again, in a physical gauge, the amplitude for a time-like cascade can be represented by a ladder diagram (Fig. 10a). However, now the evolution of the parton virtualities starts at the Q^2 -scale of the hard scattering, decreasing in a strongly ordered manner,

$$Q^2 \gtrsim m^2 \equiv m_1^2 \gg m_2^2 \gg \dots \gg m_j^2 \gg \dots \gg m_n^2 \geq \mu_0^2 \quad (22)$$

down to some final value μ_0^2 . The LLA probability for such a time-like n -branching process (Fig. 10b) is given by the product of probabilities for individual branchings $a \rightarrow bc$, where a parton of time-like mass m_a^2 decays into partons b, c with mass $m_{b,c}^2 < m_a^2$, and having fractions z and $(1 - z)$ respectively of the momentum p_a of the parent parton a [52, 65] (Fig. 10c). The time-like kinematics in the frame where $\vec{p}_a$ is along the z -axis require

$$m_a^2 = \frac{m_b^2}{z} + \frac{m_c^2}{1-z} + \frac{p_\perp^2}{z(1-z)} , \quad (23)$$

with $\pm \vec{p}_\perp$ being the transverse momentum of b, c with respect to the z -axis. Thus a sequence of time-like final state branchings can be represented, similar to the case of space-like initial state branchings, as a tree of successive single branchings connected to the scattering vertex. However, the two processes differ in that the virtualities and momentum fractions evolve in the opposite direction (Fig. 10). The time-like branchings are again associated with the branching functions $\gamma_{a \rightarrow bc}(z)$, eq. (16), characteristic for the type of branching. For a branching $a \rightarrow bc$, their behavior with varying m_a^2 is controlled by the probability distribution

$$dP^{(T)} = \frac{\alpha_s(\tilde{m}^2)}{2\pi} \frac{dm_a^2}{m_a^2} \gamma_{a \rightarrow bc}(z) dz . \quad (24)$$

here $\tilde{m}^2$, the argument in α_s , is generally a function of m^2 and z ; the proper choice of this scale is $\tilde{m}^2 = z(1-z)m^2 \simeq p_\perp^2$ [69, 70]. The probability that a parton a does not branch between the initial maximal mass $Q^2 \simeq m^2$ and a mass $\mu_0 < m^2$ is given by the exponentiation of (24),

$$T_a(Q^2, \mu_0^2) = \exp \left\{ - \int_{\mu_0^2}^{Q^2} \frac{dm'^2}{m'^2} \sum_b \int_{z_{\min}(m'^2)}^{z_{\max}(m'^2)} dz \frac{\alpha_s(\tilde{m}'^2)}{2\pi} \gamma_{a \rightarrow bc}(z) \right\} , \quad (25)$$

where $z = x_b/x_a$ and $Q^2 \simeq m_a^2$ is again identified with the scattering vertex. T is called the *Sudakov formfactor for time-like branchings* ('TB' in Fig. 9). It gives the probability that a parton a does *not* branch between invariant mass Q^2 , associated with the vertex at

which it was produced, and a minimal invariant mass $\mu_0^2 < m^2$. The limits z_{min}, z_{max} on the z -integration are constrained by the branching kinematics (23).

It is useful to compare the formfactor T_a (25) with the corresponding expression S_b for space-like branchings S_b , eq. (21). The most obvious difference between S_b and T_a is the explicit appearance of structure functions in the former. This expresses the fact that the probability for a parton b to originate from a branching of a parton a is proportional to the number of partons of type a present in the nucleus of the incoming nuclei. In contrast to that, the probability for a given parton a to branch, once it is present, is independent of this number.

2.5 Further developments of the parton cascade description

Following the logic of the LLA approach, in the 1980's the concepts and techniques outlined in the previous sections were extended and improved to design an accurate description of the properties of particles with very small momenta ($x \ll 1$) produced in parton interactions. The practical importance of the subject stems simply from the fact that such soft particles, which take away a neglegable portion of the total energy, form at the same time the bulk of multiplicity. Applied to relativistic hadron collisions it was found [17, 18] that the cross-section and the reaction dynamics arises mainly from the dynamics of frequent so-called *semihard* parton interactions with momentum transfers Q of only a few GeV. Such processes involve the production of quark and gluon minijets with transverse momenta $q_\perp$ large compared to the QCD scale Λ , but with a small fraction x of the initial hadron energy, that is

$$\sqrt{Q^2} \simeq q_\perp \gg \frac{1}{R_N}, \quad x \ll 1, \quad (26)$$

where the nucleon radius R_N sets the typical hadronic scale. Due to the dense parton phase-space population at small x (c.f. Sec. 2.1), the global cross-section for these semihard processes (being proportional to the number of partons) proves to be rather large,

$$\sigma^{(semihard)} \propto \frac{n^{(parton)}}{q_\perp^2} \simeq R_N^2, \quad (27)$$

reaching values comparable with the geometrical dimensions of nucleons. Also in many other respects these semihard processes prove to be similar to truly "soft" (non-perturbative) nucleon-nucleon interactions. But unlike "soft" interactions the semihard processes occur at short distances,

$$r^{(semihard)} \simeq \frac{1}{q_\perp} \ll R_N, \quad (28)$$

and can be calculated in perturbative QCD. Therefore it is possible to study rather typical hadronic interactions ($\sigma \sim R_N^2$) by perturbative methods.

In order to take into describe quantitatively the specific properties of parton cascades at small x , techniques were developed to account for important interference graphs that involve soft gluon lines connecting harder partons of quite different generations in a cascade. To preserve the classical cascade picture, which implies that the structure of elementary parton branchings should only depend on the nearest ancestors of a considered parton, the task was to absorb all essential interference terms into a local probabilistic scheme. This was achieved by a simple generalization of the LLA scheme, so that this extended approach is often termed the *Modified Leading Logarithmic Approximation* (MLLA). For detailed reviews I refer to Refs. [52, 73, 74, 45]. The essential idea of the MLLA is an appropriate choice of evolution variable, namely the opening angle of successive branchings rather than the virtuality. This choice makes it possible to incorporate all substantial interference contributions in Markov chains of sequential angular ordered parton branchings that preserve the property of locality and thereby the probabilistic picture of parton cascade evolution [65]. This angular ordered parton evolution will be of relevance for the PCM approach explained in Chap. 4.

The approach of semihard physics within the MLLA has been very successful in describing the dynamics of high energy hadron-hadron collisions. It predicts the correct behaviour of the total and inelastic cross-sections, their energy ($\sqrt{s}$) dependence and the inclusive spectra of hadrons with $\langle q_{\perp} \rangle \gtrsim 400$ MeV with good accuracy. In particular, the most important predictions [17, 18] are: (i) a significant growth of the characteristic transverse momenta of partons $\bar{q}_{\perp} \sim \exp(c\sqrt{\ln s})$; (ii) a rapid increase in the total multiplicity $N \sim \bar{q}_{\perp}^2 \sim \exp(2c\sqrt{\ln s})$; and, (iii) a strong increase of the total hadronic cross-section $\sigma_{tot} \propto (a \ln s)^2$.

In Fig. 11 some calculated predictions from detailed numerical studies of minimum bias pp ($p\bar{p}$) collisions within the perturbative QCD description are shown in comparison to experimental data where existent, as examples for the predictive power of the parton approach at collider energies. Fig. 11a shows the growth with $\sqrt{s}$ of the average transverse momentum of produced charged particles, Fig. 11b the corresponding increase of the average charged multiplicity, and Fig. 11c the strong rise of the total and inelastic pp ($p\bar{p}$) cross-sections. It is important to realize that the general growth pattern of these quantities is only partly due to the increasing amount of available energy $\sqrt{s}$; the main contribution arises from the increase of the number of minijets produced. These above predictions result essentially from the fact that the average total multiplicity of partons in a cascade increases with beam energy, $\bar{n} \sim \exp[const. \alpha_s \ln s]$, most of them carrying small energy fractions x (c.f. Sec. 2.3). Therefore, at high energies the interaction probability approaches unity and a nucleon appears to be increasingly "black". This property gives rise to the experimentally observed *BEL* effect [75] which is illustrated in Fig. 12. The probability distribution $P_{inel}(b)$

for an inelastic pp ($p\bar{p}$) collision is plotted versus the impact parameter b of the colliding particles, for various energies $\sqrt{s}$. It is obvious that not only approaches P_{inel} unity for central collisions, but most strikingly the distribution widens significantly with increasing collision energy $\sqrt{s}$. Therefore the term *BEL* effect: at high energies a nucleon appears to become blacker, edgier, and larger. Asymptotically, a nucleon looks like a black disc with an energy dependent radius $R \propto \ln s$.

3. SEMI-CLASSICAL TRANSPORT THEORY OF MANY-PARTON SYSTEMS IN NUCLEAR COLLISIONS

So far I have only dealt with the general features of the perturbative QCD description of high energy hadron interactions and the implications following from the underlying parton cascade picture. The evolution of the parton densities (9) with respect to the variables x and Q^2 , integrated over transverse momentum, is a pure momentum space formulation. No reference to the space-time development was made. For hadronic collisions this is (at least at current collider energies) a sufficient concept, since the parton dynamics in these reactions occurs on such short space-time distances that it is very rare that the same parton scatters more than once, or that particles from different cascades interact with each other. For nuclear collisions however, in particular heavy ion reactions, the duration of a collision is comparably rather long, so that the perturbative parton evolution can extend over space-time distances of several fm and the assumption of a single parton scattering per nucleon-nucleon collisions does not apply anymore. Moreover the parton density of the highly Lorentz contracted nuclei is rather large already in the initial state and is further increased by materialization and multiplication of partons. Therefore multiple scatterings (including sequential scatterings) of partons can easily lead to a large number of simultaneously evolving cascades that also can interact with each other. In order to resolve such an internetted structure of parton interactions, the space-time dynamics of the system *must* be taken into account. Therefore, to investigate the space-time evolution of the parton distributions, one needs to extend the momentum space description to full six dimensional phase-space and time. In this direction ambitious efforts [7, 8, 9] have been attempted, to derive from first principles a rigorous quantum kinetic theory for QCD, but due to the non-abelian character of the theory and the great theoretical complexity, a number of unresolved conceptual and technical problems prevent up to now a practical application to realistic problems.

An alternative, more pragmatic approach is realized in the parton cascade model (PCM) [38, 39] within semi-classical kinetic theory and the field theoretic QCD parton picture. In this approach, as will be explained in more detail in Chap. 4, the partons are considered as "extended" quanta with a QCD formfactor, that propagate on classical trajectories *in*

between their vertices, but the interaction amplitudes *at* the vertices are calculated quantum theoretically according to perturbative QCD and quantum statistics. The basic picture that underlies the PCM is that immediately after the contact of two colliding nuclei the partons, originally confined inside the nucleons of the incoming nuclei, are frequently scattering on each other in independent parton-parton collisions, exchanging energy and momentum and emitting other partons due to their excitation. These multiple interactions are visualized as the space-time evolution of many simultaneous parton cascades, interacting with each other. The dynamical evolution of the parton distributions is described in a probabilistic manner as a succession of binary parton-parton collisions with associated radiative emission and absorption processes. Before explaining and discussing the model in Chap. 4, it is useful to recall beforehand some general elements of semi-classical transport theory. In this chapter I will elucidate how the space-time evolution of a many-parton system can be dynamically described on the basis of single particle phase-space distributions that obey a Lorentz invariant transport equation of Uehling-Uhlenbeck type. Embedding the momentum-space parton cascade description outlined in the previous chapter into this framework, one obtains then a formulation for the time development in full phase-space.

3.1 Transport equation for the kinetics of many-parton systems

In order to quantitatively describe the dynamics of a statistical many-parton configuration in complete phase-space and time along the parton picture, I will now set up a kinetic formulation that is suitable to describe the space-time development of high energy nuclear collisions in terms of parton cascades. A relativistic transport equation for the phase space density of partons is introduced, which can be solved numerically by means of Monte Carlo techniques, i.e. by simulating many collision events and taking their average (c.f. Appendix). The basic idea is to embody the perturbative parton cascade picture into a relativistic version of the semiclassical kinetic theory, originally introduced by Uehling and Uhlenbeck [76]. For a review of various transport equation approaches to nuclear collisions and their relationship, see Ref. [77].

Consider the partons as *off-shell* particles of various species. Accounting for the off-shellness (or virtuality) [78] is a novum, as it is a purely quantum theoretical concept which is commonly avoided in statistical transport theory of quasiclassical systems. However, as explained before, in QCD field theory the partons must be viewed as "extended" objects with a quark-gluon substructure, which is embodied in the specific QCD formfactors of the quanta. The various types of partons are gluons g , quarks q_i and antiquarks $\bar{q}_i$ appearing generally in six flavors $i = u, d, s, c, b, t$ with corresponding (current) masses m_i (and of course $m_g = 0$). The spin and color degrees of freedom are not explicitly taken into account; they enter only in the spin and color averaged interaction amplitudes and in a

heuristic quantum statistical factor (explained below).

The state of a parton will thus be characterized by its species $a = q_i, \bar{q}_i, g$, virtuality Q^2 , momentum $\vec{p}$ and position $\vec{r}$. The energy is determined by $E^2 = p^2 + m_a^2 + Q^2$ where m_a is the rest mass of parton a and Q^2 - as in (8) - a possible space-like ($Q^2 = -q^2$; $q^2 < 0$) or time-like ($Q^2 = m^2$; $m^2 > 0$) virtual mass ($Q^2 = 0$ for partons on mass shell). Thus, energy and momentum are independent variables, because of the continuous dynamical mass spectrum of the partons. Let me introduce the following short notation for the four vectors,

$$p \equiv p^\mu = (E, \vec{p}) , \quad r \equiv r^\mu = (t, \vec{r}) . \quad (29)$$

As before natural units are employed. Then one can define the *Lorentz invariant single particle distribution function* for the time-dependent phase space density of partons of species a [79, 80]:

$$F_a(p, r) d^4p d^3r = \begin{cases} \text{expected number - at time } t - \text{ of partons of type } a \\ \text{in } d^3r \text{ around } \vec{r} \text{ with momentum in } d^3p \text{ around } \vec{p} \\ \text{and virtuality } Q^2 = E^2 - \vec{p}^2 - m_a^2 \end{cases} . \quad (30)$$

This function ³ contains all the information that is required for the statistical description of the time-evolution of a many-parton system in six-dimensional phase-space. The distribution function for the complete parton system is given by the sum over all species

$$\mathcal{F}(p, r) \equiv \sum_{a=g, q_i, \bar{q}_i} F_a(p, r) . \quad (31)$$

Two issues are worth mentioning here:

(i) *Covariance of the phase-space distributions*: The distribution functions F_a or $\mathcal{F}$ themselves are Lorentz scalars, although the variables $E, \vec{p}$ and $t, \vec{r}$ refer to a certain Lorentz frame. The scalar character becomes evident when one represents F_a according to the definition (30) in terms of the off-shell condition $Q^2 = E^2 - \vec{p}^2 - m_a^2$:

$$F_a(p, r) = F_a(Q^2, \vec{p}; \vec{r}, t) \delta \left[Q^2 - (E^2 - \vec{p}^2 - m_a^2) \right] \theta(E) . \quad (32)$$

Now consider the particle distribution in the rest frame L_{rest} of the system,

$$F_a(p, r) d^4p d^4r \Big|_{rest} = F_a(Q^2, \vec{p}; \vec{r}, t) \frac{d^3p}{2E} dQ^2 dt dz d^2r_\perp \Big|_{rest} . \quad (33)$$

When transforming the system to a different Lorentz frame, say L_{lab} , by a boost γ along the z -axis, one observes that both $d^3p/(2E)$ and dQ^2 are Lorentz invariant elements and $t_{lab} = \gamma t_{rest}$, $z_{lab} = z_{rest}/\gamma$, $\vec{r}_{\perp lab} = \vec{r}_{\perp rest}$, and consequently

$$F_a(p, r) d^4p d^4r \Big|_{lab} = F_a(p, r) d^4p d^4r \Big|_{rest} , \quad (34)$$

³I emphasize that the parton phase-space densities $F_a(p, r)$ are not to be confused with the nucleon structure functions $F_a^N(x, Q^2)$, eq. (10), although they are closely related (see item (ii) below).

which proves the scalar character of the distribution function $F_a(p, r)$.

(ii) *Relation between phase-space distributions and structure functions:* Recall from Chap. 2 that the short distance behavior of QCD interactions manifests itself in a miraculous "decoherence" [81] of partons in a nucleon when probed at scales $1/\sqrt{Q^2} \ll R_N$. Long range color forces corresponding to soft gluon modes with wavelengths $\sim R_N$ are practically unobservable by means of experimental triggering. Thus, the structure functions mirror a nucleon as an ensemble of uncorrelated, independent partons that interact binarily almost like classical particles. In other words, the structure functions contain information about the single-particle aspect of the parton distributions. However, the nucleon structure functions $F_a^N(x, Q^2)$, eq. (10), correspond to the single-particle distributions $F_a(p, r)$ of eq. (30) only after integration of the complete space-time history as well as transverse momentum, and consequently contain solely information about the parton densities in terms of the subset of variables (x, Q^2) out of the full set $(\vec{p}, Q^2, \vec{r}; t)$. This relation between $F_a^N(x, Q^2)$ and $F_a(\vec{p}, Q^2, \vec{r}; t)$ is evident by employing eqs. (9) and (10),

$$F_a^N(x, Q^2) = \int_0^{Q^2} dQ'^2 \frac{dN_a(x, Q^2)}{dx dQ'^2} \quad (35)$$

where according to the definition (30) and eq. (32)

$$N_a = \int_{-\infty}^{\infty} dt \int d^3r \int dQ^2 \int \frac{d^3p}{2E} E F_a(Q^2, \vec{p}; \vec{r}, t) \quad (36)$$

Using $d^3p/(2E) = (\pi/2)(dx/x)dp_{\perp}^2$ and $d^3r|_{lab} = xM_N d^3r|_{rest}$, one obtains,

$$\frac{dN_a}{dx dQ^2} = \frac{2\pi^2}{3} R_N^3 M_N \int dp_{\perp}^2 F_a(Q^2, \vec{p}; \vec{r}, t), \quad (37)$$

where $\vec{p} = (xP, \vec{p}_{\perp})$ and $Q^2 = E^2 - \vec{p}^2 - m_a^2$. The nucleon radius and mass are denoted as R_N , respectively M_N .

Within a statistical description in terms of the single-particle distributions $F_a(p, r)$, the time evolution of the many-parton system can now be described on the basis of a relativistic *transport equation for the parton distribution functions* (30),

$$p^{\mu} \partial_{\mu} F_a(p, r) = \sum_{\text{processes } k} I_a^{(k)}(p, r), \quad (38)$$

where $\partial_{\mu} \equiv \partial/\partial r^{\mu}$. The left-hand side of this equation describes the free propagation of partons, whereas the collision term on the right-hand side $\sum_k I_a^{(k)} \equiv \sum_k (I_a^{(k)gain} - I_a^{(k)loss})$ represents a sum of Lorentz invariant phase-space integrals containing the matrix-elements for the different kinds of interaction processes in which at least one parton of species a is involved in either the initial or final state of a process. A term corresponding to a particular

n -body process at $\vec{r}$ and time t is given by an integral over $(n - 1)$ -particle phase space (see Fig. 13):

$$I_{a_i}^{(k)}(p_i, \vec{r}, t) = (\pm) \frac{1}{S_i} \int \prod_{\substack{j=1 \\ j \neq i}}^n d^4 p_j \left| \langle a_n, \dots, a_{m+1} | \hat{S}^{(k)} | a_m, \dots, a_1 \rangle \right|^2 D \{F(j)\} , \quad (39)$$

where the sign in front is $-(+)$ for the case where parton i is coming in (emerging from) the interaction process, and the corresponding S -matrix element is related to the invariant amplitude $\mathcal{M}^{(k)}$ by

$$\langle a_n, \dots, a_{m+1} | \hat{S}^{(k)} | a_m, \dots, a_1 \rangle = \mathcal{M}^{(k)}(p_1, \dots, p_n)_{a_1, \dots, a_m \rightarrow a_{m+1}, \dots, a_n} (2\pi)^4 \delta^4(P_{in} - P_{out}) . \quad (40)$$

Here P_{in} and P_{out} are the sums of the four momenta of the incoming, respectively outgoing partons. The quantum statistical distribution factor D in (39) is

$$D \{F(j)\} \equiv \prod_{in} F(j) \prod_{out} [1 \pm F(j)] - \prod_{in} [1 \pm F(j)] \prod_{out} F(j) , \quad (41)$$

and the statistical factor S_i in front of the integral is given by

$$S_i = \prod_{j \neq i} K_{a_j}^{in} K_{a_j}^{out} , \quad (42)$$

where there are $K_a^{in,out}$ identical partons of species a in the initial or in the final state of the process, excluding the i -th parton.

At this point it is important to make the following precautions remarks. The transport equation is, as it stands, a semiclassical formulation: it describes the evolution of a many-particle system in terms of single-particle distribution functions and classical particle trajectories. The quantum nature of the system is inherent exclusively in the collision term (its specific form will be introduced in Sec. 4.2). The collision term models the space-time structure of parton interactions embodied in the relevant matrix-elements and takes into account quantum statistical effects heuristically by using Fermi-Dirac and Bose-Einstein statistics for quarks and gluons, respectively. Note that long range color forces are neglected and that no color mean field is included. It is important to realize that this transport equation is not derived from first principles of QCD. Rather it is a special case of Landau's kinetic equation which can be deduced from the general many-body Schrödinger equation within the statistical approach of particle transport theory, or alternatively from classical many-body theory (for a review see Ref. [77]). Therefore the kinetic framework introduced here is based on the usual assumptions [79, 80] that underly the derivation of this type of transport equation.

In order to solve the transport equation (38) for the time evolution of the many-parton system under consideration, one needs to specify (i) the initial parton distributions $F_a^{(0)}(p, \vec{r}, t_0)$ at time t_0 and (ii) the specific form of the collision term, that is the class of parton interaction processes to be considered. Both ingredients depend on the physical problem under consideration. The application to collisions of highly relativistic hadrons or nuclei will be explained in detail in the following Chap. 4.

3.2 Physical quantities relating microscopic parton dynamics to macroscopic observables

Starting from the initial parton distributions at $t = t_0$, the time evolution of the phase-space densities $F_a(p, r)$ according to the transport equation (38) reflects the state of the system around $\vec{r}$ and $\vec{p}$ at any time $t > t_0$. Therefore, it is desirable to extract the functions $F_a(r, p)$ from eq. (38) and to calculate, in a Lorentz invariant manner, relevant macroscopic quantities that are directly or indirectly observable by an experimental apparatus. Relativistic transport theory [80] relates physical quantities to phase-space integrals over products of combinations of four-momenta or tensors and the single-particle distributions $F_a(p, r)$. For example, the local space-time dependent parton currents n_a are given by

$$n_a^\mu(r) = \gamma_a \int dQ^2 \int \frac{d^3 p}{(2\pi)^3 2p^0} p^\mu F_a(p, r) \quad . \quad (43)$$

where [c.f. (30) and (32)]

$$p^0 \equiv E = +\sqrt{\vec{p}^2 + m_a^2 + Q^2} \quad . \quad (44)$$

The factors γ_a are the products of spin and color degeneracy, $\gamma_g = 2 \times 8$ for gluons and $\gamma_q = \gamma_{\bar{q}} = 2 \times 3$ for quarks and antiquarks. The normalization factor $(2\pi)^3$ in the denominator of the invariant momentum space element has been introduced for convenience. In correspondence to (31), the total parton current is obtained by summing over all species,

$$n^\mu(r) = \sum_{a=g,q_i,\bar{q}_i} n_a^\mu(r) \quad . \quad (45)$$

Correspondingly, the energy-momentum tensor is given by

$$T^{\mu\nu}(r) = \sum_{a=g,q_i,\bar{q}_i} \gamma_a \int dQ^2 \int \frac{d^3 p}{(2\pi)^3 2p^0} p^\mu p^\nu F_a(r, p) \quad , \quad (46)$$

and the total entropy current of the parton system is

$$s^\mu(r) = - \sum_{a=g,q_i,\bar{q}_i} \gamma_a \int dQ^2 \frac{d^3 p}{(2\pi)^3 2p^0} p^\mu \left\{ F_a(r, p) \ln[F_a(r, p)] - \theta_a (1 + \theta_a F_a(r, p)) \ln[1 + \theta_a F_a(r, p)] \right\} \quad , \quad (47)$$

where $\theta_g = +1$ and $\theta_q = \theta_{\bar{q}} = -1$ take into account Bose-Einstein statistics for gluons and Fermi-Dirac statistics for quarks and antiquarks, respectively (note that $\theta_a = 0$ recovers the classical Maxwell-Boltzmann limit).

In order to formulate a Lorentz invariant description of these macroscopic quantities, one must introduce the *hydrodynamical four-velocity* $u^\mu(r)$. It is a vector field, defined as a time-like vector in each space-time point: $u_\mu(r)u^\mu(r) = 1$. To specify the hydrodynamic velocity various definitions may be employed [80]. For example,

$$u^\mu(r) = \frac{n^\mu(r)}{\sqrt{n_\nu(r)n^\nu(r)}} \quad (\text{Eckardt's definition}), \quad (48)$$

which relates u^μ directly to the local particle four-flow. A different choice is

$$u^\mu(r) = \frac{T^{\mu\nu}(r)u_\nu(r)}{\sqrt{u_\rho(r)T^{\rho\sigma}(r)T_{\sigma\tau}(r)u^\tau(r)}} \quad (\text{Landau-Lifshitz's definition}), \quad (49)$$

which relates the hydrodynamical velocity to the flow of energy, or equivalently the momentum density.

By contracting the quantities (43)-(47) with the local flow velocity u^μ , one can now obtain corresponding invariant scalars of particle density, pressure, energy density and entropy density:

$$n(r) = n_\mu(r) u^\mu(r) \quad (50)$$

$$p(r) = -\frac{1}{3} T_{\mu\nu}(r) \left(g^{\mu\nu} - u^\mu(r)u^\nu(r) \right) \quad (51)$$

$$\varepsilon(r) = T_{\mu\nu}(r) u^\mu(r)u^\nu(r) \quad (52)$$

$$s(r) = s_\mu(r) u^\mu(r), \quad (53)$$

where $g^{\mu\nu} = \text{diag}(1, -1, -1, -1)$. Due to the scalar character of these quantities, they provide local, Lorentz invariant measures of the many-particle system. It is left to convenience in which Lorentz frame the calculation is performed, but in general it is considerably simplified in the local restframe of the matter where $u^\mu = (1, \vec{0})$. The total number, the free energy and the entropy of the particles at a given time can be obtained by integrating over coordinate space. Furthermore, one can always define a space-time dependent *effective temperature*, e.g. as

$$T(r) := \frac{\varepsilon(r) + p(r)}{s(r)}, \quad (54)$$

which coincides with the expression for the temperature of an equilibrated plasma of partons, but of course has not the strict meaning of temperature for a system away from thermal equilibrium. Nevertheless, this quantity generally serves as a measure of dissipation and transverse energy production due to particle interactions, and its space-time dependence can indicate a possible approach to local thermal equilibrium.

It is important to realize that the quantities (50)-(54) are well defined in terms of phase-space integrals for *any* distribution $F_a(p, r)$ within non-equilibrium kinetic theory. In particular, there is *no* assumption about a thermal or chemical equilibrated state of the system implicit. In fact, the strength of this formulation lies in the ability to follow the system in time from a highly non-equilibrated state as it evolves eventually into an equilibrium. This issue will be studied in Chap. 6 for the realistic case of nuclear collisions.

3.3 Detailed balance and equilibration

It is useful for later considerations to point out here some properties of *detailed balance* of the various interaction processes k taken into account in the collision integrals $I_a^{(k)gain}$ and $I_a^{(k)loss}$, and the implications for thermal and chemical equilibration of the many-parton system. When the collision term on the right hand side of eq. (38) becomes zero, the parton distribution functions provide a stationary solution to the transport equation. In the case of high energy nucleus-nucleus collisions there are basically two possibilities for the collision term to vanish locally [82]: (i) in the central region, when the system of partons reaches a state of detailed balance with respect to the contributing $2 \rightarrow 2$, $2 \rightarrow 1$, $1 \rightarrow 2$ processes, implying equilibration, and (ii) when the partons in the beam fronts are fast receding from the collision region and are approaching free streaming.

In general the interaction processes contributing to the collision term (39) fall into two categories; namely, processes that do not change the number of quarks and antiquarks, and those that alter the quark flavor composition. For the $2 \rightarrow 2$ processes these are

$$a + b \rightarrow a + b \quad , \quad (55)$$

$$a + \bar{a} \rightarrow c + \bar{c} \quad , \quad (56)$$

where $a, b = q, \bar{q}, g$ and $c = q$. The process (55) corresponds to elastic scattering and the annihilation process (56) is a 'chemical reaction' that changes the number of quarks. The inelastic processes $2 \rightarrow 1$ and $1 \rightarrow 2$, which produce or annihilate one gluon, are

$$a \leftrightarrow a + g \quad , \quad (57)$$

$$g \leftrightarrow c + \bar{c} \quad , \quad (58)$$

with $a = q, \bar{q}, g$ and $c = q$. Here (57) is a gluon emission ($\rightarrow$) or an absorption ($\leftarrow$) process with parton a appearing in both, initial and final state. The process (58) is again a 'chemical reaction' that increases (reduces) the number of quark-antiquark pairs.

Semiclassical kinetic theory [80] states that when the system is in local equilibrium with respect to the elastic binary collisions (55) and the gluon emission and absorption processes

(57), then the distribution functions in phase-space for the various species a , $F_a(p, r)$, take the form

$$F_a^\pm(p, r) = \frac{1}{\exp[\beta_\nu(p^\nu - \lambda_a^\nu)] \pm 1} . \quad (59)$$

Here p_ν is the four-momentum of the particle and the upper sign refers to fermions (quarks) and the lower sign to bosons (gluons). $\lambda_\nu(r)$ and $\beta_\nu(r)$ are four-vectors constructed from the local flow velocity $u_\nu(r)$, the local chemical potentials $\mu_a(r)$ (one for each quark species) and the inverse local temperature $\beta(r) = 1/T(r)$:

$$\lambda_a^\nu(r) = \mu_a(r) u^\nu(r) , \quad \beta^\nu(r) = \beta(r) u^\nu(r) . \quad (60)$$

The non-vanishing chemical potentials μ_a for flavor a are the parameters which measure the number of quarks minus antiquarks. In a baryon-free central rapidity region they give the deviation from complete chemical equilibrium. In addition, equilibrium with respect to the ‘chemical reactions’ (56) and (58) requires that the sum of the chemical potentials of the particles on the left hand side equals the sum on the right hand side for each process [83]. This can easily be verified by inserting the distribution functions (59) into the collision integrals and using the identity $1 \mp F^\pm = \exp[\beta_\nu(p^\nu - \lambda_a^\nu)] F^\pm$. For the processes (56), the equilibrium conditions are

$$\mu_a + \mu_{\bar{a}} = \mu_c + \mu_{\bar{c}} , \quad (61)$$

$$\mu_a + \mu_{\bar{a}} = 2\mu_g , \quad (62)$$

and for the processes (58)

$$\mu_a + \mu_{\bar{a}} = \mu_g . \quad (63)$$

Consequently, from (62) and (63), one sees that the gluon chemical potential must vanish and therefore $\mu_a = -\mu_{\bar{a}}$. On the other hand, in a baryon-free plasma particle-antiparticle symmetry imposes $\mu_a = \mu_{\bar{a}}$, so that the conditions (61)-(63) imply

$$\mu_g = 0 , \quad \mu_a = \mu_{\bar{a}} = 0 , \quad (64)$$

and the latter equality must hold for each quark flavor individually.

Thus, in nuclear collisions where one may distinguish between an approximately baryon-free central region dominated by gluons and $q\bar{q}$ pairs, and a baryon-rich nuclear fragmentation region containing most of the fast valence quarks, the implications of detailed balance are as follows: In the approximately baryon-free central region, a complete thermal and chemical equilibrium can (and, provided the system has time enough, eventually will) be established, if the system is characterized by detailed balance with respect to the processes (55)-(58), that is by vanishing chemical potentials for all quark flavors. As far as

the baryon-rich nuclear fragmentation region is concerned, the collision term becomes very small at times when the fast partons in the beam fronts approach free streaming and move collinearly away from the central region. In this case the parton densities become dilute and the probability of two partons a and b to collide, $dP_{ab} \propto \sqrt{(p_a \cdot p_b)^2 - m_a^2 m_b^2} d\hat{\sigma}_{ab}$, tends to zero. Once the system of partons has reached an equilibrium state, its further evolution can be described within the framework of relativistic hydrodynamics according to the hydrodynamic equations:

$$\partial_\mu T^{\mu\nu}(r) = 0, \quad \partial_\mu n^\mu(r) = 0. \quad (65)$$

Whereas the first equation is a consequence of energy-momentum conservation, the second equation follows from (38) by integrating over d^3p and the fact that the collision term vanishes, if the system is in a thermal and chemical equilibrium state with the distribution functions taking the form (59). In the absence of dissipation, the energy-momentum tensor can be written in terms of the proper energy density $\varepsilon(r)$, the pressure $p(r)$ and the local flow velocity $u^\mu(r)$, using eqs. (51) and (52):

$$T^{\mu\nu}(r) = -g^{\mu\nu} p(r) + \left(\varepsilon(r) + p(r) \right) u^\mu(r) u^\nu(r). \quad (66)$$

The details of the space-time development of the system will depend on its equation of state $\varepsilon(p)$, or equivalently, $p(T)$, interrelating the local energy density ε , pressure p and temperature T . For a quark-gluon plasma consisting of partons with chemical potentials μ_a for quarks and antiquarks, the equation of state may be derived from the thermodynamic potential

$$\Omega(T, V, \mu_a) = V a(T, \mu_a, g) T^4, \quad (67)$$

with T the temperature, V the volume and g the strong coupling strength taken at a typical momentum $\max(T, \mu_a)$. The coefficient function $a(T, \mu_a, g)$ has been calculated [84, 85] with the inclusion of parton interactions up to order $g^3 T^4$ within perturbative QCD. The pressure $p = -(d\Omega/dV)_{T, \mu}$, the entropy $S = -(d\Omega/dT)_{V, \mu}$, etc., follow from the standard thermodynamic relations.

The initial conditions for the hydrodynamic description $\varepsilon_0 = \varepsilon(t_i, \vec{r})$ or $p(t_i, \vec{r})$, at a specified time $t = t_i$ when the fluid dynamical expansion of the plasma starts, depend sensitively on the preceding space-time evolution during the pre-equilibrium stage. Thus, by following the system from some initial non-equilibrium state up to the formation of a thermalized plasma as governed by the transport equation (38), one can extract the initial and boundary conditions for the subsequent evolution according to the laws of relativistic fluid dynamics [86].

It is illustrative to exemplify the hydrodynamic evolution with the famous Bjorken model [100] for relativistic nucleus-nucleus collisions. The underlying picture of this model

is the one sketched in Sec. 1.1. It is presumed that from the point of time when equilibration is attained, the expansion of the system proceeds exclusively in longitudinal direction along the beam (z) axis and is invariant under Lorentz boosts along this axis. This implies that all local thermodynamic quantities are functions of the proper time $\tau = \sqrt{t^2 - z^2}$ only, i.e. $\varepsilon(t, \vec{r}_\perp, z) = \varepsilon(\tau)$, $T(t, \vec{r}_\perp, z) = T(\tau)$ etc.. The fluid four-velocity is taken to be $u^\mu(r) \equiv (\gamma, \gamma \vec{v}) = (t/\tau, 0, 0, z/\tau)$, so that $v_z = z/t$. Another assumption of the Bjorken scenario is that the baryon number of the initial nuclei is assumed to reside completely in the receding beam fronts and that the central region is ideally baryon free. Moreover, residual interactions between the partons are neglected and the particles are treated as massless, so that the system can be treated within the ideal gas approximation. Hence, if one concentrates on the central region and sets in eq. (67) $g = 0$ and $\mu_a = 0$, one obtains the equation of state for an ideal gas of non-interacting, massless quarks and gluons [$a(T, \mu_a = 0, g = 0) \equiv a(T)$]:

$$p = a(T) T^4 \quad (68)$$

and for the energy density, entropy density and particle number density, respectively,

$$\begin{aligned} \varepsilon &= 3p \left(1 + \frac{T}{a(T)} \frac{\partial a(T)}{\partial T} \right) \simeq 3 a(T) T^4 \\ s &= \frac{\varepsilon + p}{T} \simeq 4 a(T) T^3 \end{aligned} \quad (69)$$

$$n = b(T) T^3$$

where

$$a(T) = \frac{\pi^2}{90} \left(16 + \frac{21}{2} n_f \right) = \begin{cases} 4.06 & (n_f = 2) \\ 5.20 & (n_f = 3) \\ 6.36 & (n_f = 4) \end{cases} \quad (70)$$

$$b(T) = \frac{\zeta(3)}{\pi^2} (16 + 9n_f) = \begin{cases} 4.14 & (n_f = 2) \\ 5.24 & (n_f = 3) \\ 6.33 & (n_f = 4) \end{cases} \quad (71)$$

Here $n_f = n_f(T)$ is the effective number of quark flavor degrees of freedom in the deconfined phase, depending on the temperature T above the critical temperature $T_c \approx 180$ MeV.

The total entropy produced by the partons contained in an rapidity interval Δy around $y = 0$ at a given time is obtained by integrating the local entropy density $s(r)$ from eq. (53) over position space, Using $y = 1/2 \ln [(E + p_z)/(E - p_z)]$ and $\eta = 1/2 \ln [(t + z)/(t - z)]$ one obtains:

$$S(t) = \int dz \int d^2 r_\perp u^\mu s_\mu \simeq \pi R_A^2 \int_{-\Delta y/2}^{+\Delta y/2} d\eta \tau s(\tau) = \Delta y \pi R_A^2 4 a T^3 \tau, \quad (72)$$

where $a \equiv a(T)$ from eq. (70), $T = T(\tau)$, $R_A = 1.12 A^{1/3}$ is the nuclear radius and the range of η is given by $-\Delta y/2 \leq \eta \leq +\Delta y/2$. Note that the approximate equality on the right hand side of (72) holds only under the assumptions of ideal longitudinal flow and of a homogeneous density in transverse direction up to R_A , that is, $s(r) = s(\tau, \eta, \vec{r}_\perp) \rightarrow s(\tau)$. The conservation of the total entropy follows then as a consequence of the thermodynamic equations (50)-(53):

$$\frac{d}{d\tau} \left(\frac{dS}{dy} \right) = 0 \quad . \quad (73)$$

I emphasize that the Bjorken model is an idealized scenario that attracts by its simplicity. However, it neglects effects due to flavor dependent quark masses, interactions among the quarks and gluons, chemical potentials for the light quarks, non-vanishing baryon density in the central region, and the time-dependence of the relative admixtures of the gluons and the various quark flavors. These aspects are included in the PCM, and in Chap. 6 the numerical results for the realistic space-time evolution will be compared to the idealized Bjorken model.

3.4 Practical determination of the parton phase-space distributions

The statistical description of the complex space-time evolution of a very large parton system, e.g. involving several thousand particles as is typical in relativistic collisions of two heavy nuclei, requires a computer simulation and therefore numerical extraction of the parton distributions. The basic concept is here to introduce a discretization of phase-space [86], similar in spirit to earlier analyses within microscopic models of nuclear dynamics [87, 88].

Consider for the sake of definiteness a central collision of two heavy nuclei at very high energy, in which a large number of partons evolves in a volume of more than 100 fm^3 . To calculate the distribution functions $F_a(p, r)$ of the partons a , the six-dimensional phase-space volume $\propto d^3p d^3r$ may be divided into cells labeled by an index k . The number of partons of type a , $n_a(k)$, in each cell is found at some fixed time in the total center-of-mass frame of the colliding nuclei. Statistics is improved by summing $n_a(k)$ over several collision events of the parton cascade simulation. The distribution functions $F_a(r, p)$ are then estimated as

$$\bar{F}_a(k) = \frac{1}{\nu_{ev}} \left(\frac{n_a(k)}{\Delta\Gamma_a(k)} \right) \quad , \quad (74)$$

where ν_{ev} is the number of collision events accumulated and

$$\Delta\Gamma_a(k) = \gamma_a \int_k \frac{d^3r d^3p}{(2\pi)^3} \quad (75)$$

is the phase-space volume of cell k in natural units ($\hbar = c = 1$) of the elementary volume $h^3 = (2\pi\hbar)^3 \equiv (2\pi)^3$ for partons of species a with degeneracy factors $\gamma_g = 16$ for gluons

and $\gamma_q = \gamma_{\bar{q}} = 6$ for each quark flavor. The total parton distribution summed over all species is thus given by

$$\mathcal{F}(k) = 16 F_g(k) + 6 \sum_{i=1}^{n_f} \left\{ F_{q_i}(k) + F_{\bar{q}_i}(k) \right\}. \quad (76)$$

The division of phase-space into cells must be done with some care. This is especially important for the entropy density (53), measured in experiment by the accumulated multiplicity of produced particles, since it is - unlike particle density or energy density - not additive in the particles. In Ref. [86] a detailed analysis concerning this issue has been carried out, which is briefly summarized in the following.

3.4.1 Cellular phase-space and finite particle effects

There are two essential requirements for an appropriate phase-space discretization [88]:

- (i) The cell size must be small enough so that $\bar{F}(k)$ is approximately uniform within the six-dimensional volume of the cells. A too large cell size underestimates the particle density and overestimates the entropy.
- (ii) In the average the cells must include enough particles so that the statistical errors in the particle numbers $n(k)$ are not significant. If, for a given cell size, the number of particles is too small such that most cells are empty and the occupied cells contain only very few particles, then the density in cells containing particles overestimates the true density. Since only the occupied cells contribute in the calculation of the entropy, the computed value of the latter would be too low.

To choose a phase-space division that satisfies these two conflicting criteria such that the computed entropy is estimated with sufficient accuracy, it is important to quantitatively understand the errors associated with a certain cell configuration and the number of particles involved in the analysis. In Fig. 14 the effect of the non-uniformity of F within the cells is shown as examined within a analytically solvable model problem [86]. Concerning the error due to counting statistics with a finite number of particles, it was found that in order to achieve an accuracy for the entropy per particle of e.g. 10 %, one needs only 5 particles in each occupied cell.

It is important to check whether these considerations of errors associated with the cellular phase-space and the number of particles distributed over it are also true for the practical computation of the entropy of partons in heavy ion collisions. This can be done by considering the situation where a large number of particles are confined inside a cylindrical volume (mimicking the nuclear collision geometry) of length $2 fm$ and radius $4 fm$, i.e. $V_{box} \simeq 100 fm^3$, and distributed homogeneously throughout this volume. Fig. 15 shows the absolute error in the entropy per particle for this system as a function of the cell size (in units of Plank's volume h^3) for various numbers of particles $N = 200 - 50000$. Evidently,

the effect of the number density on the error is far less important than the choice of the cell size. For instance, an accuracy of 0.1 per particle can be achieved with a cell size of $\approx 4 h^3$ and a total number of particles $N \geq 2500/\nu_{ev}$, accumulated in ν_{ev} events, or, an average particle density of $N/V_{box} \geq 25/\nu_{ev} fm^{-3}$.

3.4.2 Sensitivity of time-development to phase-space discretization

Finally there is the question of how the time evolution of the produced entropy is affected by the cell division and the particle configuration. This is of interest because the time span for the relaxation of the entropy production determines the equilibration time scale. In the case of a nuclear collisions, the initially sparsely populated phase-space will be rapidly filled with partons that are ‘freed’ by mutual scatterings. Therefore one may expect that the statistical error in computing the entropy at certain points of time is relatively large in the beginning and decreases as more and more particles are produced, improving the statistics. Thus, the entropy would be underestimated most drastically in the early stage and consequently the rise of produced entropy with time would be too steep. Fortunately, this effect is not very significant, as can be examined quantitatively from the simple system of particles in the cylindrical box considered above. In Fig. 16 the time evolution of the estimated specific entropy $(S/N)_{est}$ for this toy model is shown for a total of $N = 10000$ particles and various cell sizes $\Delta\Gamma = 1 h^3 - 8 h^3$. The entropy per particle is initially zero (because all the particles are assigned to reside in the central cell), rises steeply as the particles spread out to occupy the phase-space and finally reaches its equilibrium value. The curves corresponding to the various phase-space divisions differ insignificantly, except around $t = 1-2 fm$, where the curvature is the largest. The relaxation times range from $\simeq 0.5 fm$ ($\Delta\Gamma = 1 h^3$) to $\simeq 0.6 fm$ ($\Delta\Gamma = 8 h^3$), implying a rather small variation of $\simeq 0.1 fm$ compared to the total equilibration time of about $6 fm$, that is a relative error of $\simeq 1 - 2 \%$.

4. THE PARTON CASCADE MODEL FOR ULTRA-RELATIVISTIC NUCLEAR COLLISIONS

The PCM [38, 39] is a relativistic kinetic model for high energy nuclear collisions which is inspired by the parton picture of hadronic interactions outlined in Chap. 2. It provides a comprehensive description of hadron-hadron, hadron-nucleus and nucleus-nucleus collisions in terms of the space-time evolution of parton cascades, i.e. the nuclear dynamics is traced back to the microscopic level of quark and gluon interactions in the framework of perturbative QCD. Embedded in the kinetic formulation of Chap. 3, this approach allows to study the time evolution of the nuclear collisions in complete phase-space from the instant of nuclear contact, through the perturbative QCD evolution of parton distributions, up

to the formation of final hadronic states. The model as a whole consists of three major building blocks, each of which is illustrated schematically in Fig. 17:

1. *Initial state*: Phenomenological construction of the initial state in terms of the partons' phase-space distributions in the incoming nuclei (or hadrons) on the basis of the experimentally measured nucleon structure functions and elastic nucleon form-factors, as well as nuclear shadowing corrections.
2. *Parton cascade development*: Time development of parton cascades in complete six-dimensional phase-space according to a semiclassical, relativistic transport equation with a collision kernel that contains the asymptotically dominant perturbative QCD interaction processes among the partons. The time variation of the parton distributions, the number and type of interactions is determined by the dynamics itself.
3. *Hadronization*: Formation of color neutral clusters from the ensemble of final partons at the end of the perturbative QCD phase. Independent hadronization of each cluster through isotropic two-body decay yields primary hadron resonances which subsequently decay into final hadrons according to the particle data tables.

In this approach the first and last components, describing initial state and final hadronization, contain essential non-perturbative QCD effects associated with the confinement properties, whereas the second component in between, the parton cascade development, is almost exclusively based on the firmly established perturbative QCD picture of Chap. 2, except where current lack of better knowledge requires supplementary phenomenological input, as e.g. the influence of dense matter environment. Attempts to model non-perturbative color exchange forces among the partons (e.g. color strings, flux tubes or color background fields) are avoided; rather the perturbative evolution is cut off when it reaches its limits of validity. Note that the third element, the non-perturbative hadronization mechanism, is considered within a completely separate model framework, and may be replaced by any other phenomenological scheme that is suitable to describe the dynamics beyond the perturbative regime.

In the present Chapter I will review the most important aspects of the PCM exemplarily for the case of nucleus-nucleus collisions $A + B \rightarrow \text{hadrons}$. For hadron-nucleus or hadron-hadron collisions the replacement of one or both of the beam nuclei by a single hadron is straightforward. Original presentations of the model can be found in [38, 39, 82].

4.1 The initial state

It is most convenient (for describing collider experiments) to choose the overall center-of-mass (CM) frame of the colliding nuclei as the reference frame, with the collision axis in the

z -direction. In the CM frame of the incoming nuclei A and B have longitudinal momenta $P_{A,B} = \pm P_{CM}$, where P_{CM} denotes the absolute value of total CM momentum. The nucleons then carry equal fractions of P_{CM} . That is, neglecting their Fermi motion, they only have nonvanishing longitudinal momentum along the z -axis, that is, $\vec{P}_{N_i} = (0, 0, P_{N_i})$ with

$$P_{N_i} = \pm \frac{1}{N_{nuc}} \frac{[s - (M_A + M_B)^2]^{\frac{1}{2}} [s - (M_A - M_B)^2]^{\frac{1}{2}}}{2\sqrt{s}} = \pm \frac{P_{CM}}{N_{nuc}}, \quad (77)$$

where $s = E_{CM}^2 = (E_A + E_B)^2$, $M_{A,B}$ are the nuclear masses, $N_{nuc} = A + B$ is the total number of nucleons in the collision system, and $+$ $(-)$ is for the nucleons N_i ($i = 1, \dots, N_{nuc}$) in nucleus $A(B)$.

In the CM frame, where the two incoming nuclei A and B and their constituent nucleons are moving close to the speed of light, the parton picture is applicable and the parton substructure of the nucleons can be resolved with reference to some *initial resolution scale* Q_0 (c.f. Sec. 2.4). This resolution scale is generally beam energy dependent and is identified with the statistically estimated expectation value for the interaction scale Q^2 of all *primary* parton-parton collisions, those in which at least one initial state parton is evolved:

$$Q_0^2(s) = \langle Q_{prim.coll.}^2 \rangle. \quad (78)$$

Here the choice for the scale associated with each primary parton scattering is $Q^2 \equiv p_\perp^2$, with $p_\perp$ the relative transverse momentum of the parton pair. Recall from Secs. 2.3 and 2.4 that the scale Q_0 defines the initial point in momentum space beyond which the system of partons is evolved perturbatively. The dynamics prior to this point must be absorbed in the initial phase-space distributions of the partons. Clearly, the convention (78) yields only an average value dominated the most probable semihard parton collisions with relatively small momentum transfers of a few GeV. However, primary parton collisions with a momentum transfer $p_\perp^2 \gg Q_0^2$, which correspond to comparably rare fluctuations, are taken into account individually by the Q^2 -evolution of the nucleon structure functions through space-like radiation processes, as will be explained in Sec. 4.2.2 below.

The actual form of the initial phase-space distribution of partons needs to be phenomenologically specified at the reference point Q_0 . It is constructed as the following superposition of the parton distributions in the individual nucleons at Q_0 and at time $t = t_0$ when the two nuclei begin to overlap:

$$F_a^{(0)}(Q_0^2; p, r) = \sum_{i=1}^{N_{nuc}} P_a^{N_i}(Q_0^2; \vec{p}, \vec{P}) \cdot R_a^{N_i}(\vec{p}, \vec{r}, \vec{R}) \Big|_{t=t_0}. \quad (79)$$

The right hand side is a sum over all N_{nuc} nucleons of the incoming nuclei A and B , each with a parton distribution given by a convolution of an initial momentum distribution

$P_a^{N_i}$ and a spatial distribution $R_a^{N_i}$. The subscript $a = g, q_j, \bar{q}_j$ ($j = 1, \dots, n_f$) labels as before the species of partons, and N_i refers to the type of the i -th nucleon, proton or neutron. The four-vectors $p = (E, \vec{p})$ and $r \equiv r = (t, \vec{r})$ refer to the partons. The nucleons' momenta and initial positions are denoted $\vec{P}$, respectively $\vec{R}$. All vectors are understood with respect to the CM , at time $t = t_0$. The partons' energies $E \equiv E_a(q^2) = \sqrt{\vec{p}^2 + m_a^2 + q^2}$ take into account their initial space-like virtualities $q^2 < 0$ which are distributed around $\langle |q^2| \rangle = -Q_0^2$, under the constraint that for each nucleon separately their total invariant mass equals the nucleon mass. This mimics the fact that the initial partons are confined inside their parent nucleons and cannot be treated as free particles, i.e. they have not enough energy to be on mass shell, but are off mass shell by a space-like virtuality $q^2 \approx -Q_0^2$.

4.1.1 Initial momentum distribution

For each nucleon the number of partons, the distribution of the flavors, their momenta and associated initial space-like virtualities, are obtained from the function $P_a^{N_i}$ in (79). Denoting $P \equiv P_{CM}/N_{nuc}$, it is represented in the form

$$P_a^{N_i}(Q_0^2, \vec{p}, \vec{P}) = \left(\frac{x}{\tilde{x}} \right) F_a^{N_i}(x, Q_0^2) \rho_a^A(x) g(\vec{p}_\perp) \delta(P_z - P) \delta^2(\vec{P}_\perp) \quad (80)$$

with the momentum and energy fractions

$$x = \frac{p_z}{P} ; \quad \tilde{x} = \frac{E}{P} = \sqrt{x^2 + \frac{p_\perp^2 + m_a^2 + q^2}{P^2}} \quad (81)$$

and the normalizations

$$\sum_a \int_0^1 dx x F_a^{N_i}(x, Q_0^2) = 1 \quad (82)$$

$$\int_0^\infty d^2 p_\perp g(\vec{p}_\perp) = 1 \quad (83)$$

$$\sum_a \int d|q^2| \frac{d^3 p}{(2\pi)^3 2E} E P_a^{N_i}(Q_0^2, \vec{p}, \vec{P}) = n^{N_i}(Q_0^2, P) . \quad (84)$$

The physics behind the ansatz (80) is the following:

(i) The functions $F_a^{N_i}(x, Q_0^2)$ are the scale-dependent measured nucleon structure functions (10) with x being the fraction of the parent nucleons longitudinal momentum P carried by the parton. The transverse momentum distribution $g(p_\perp)$ is specified below. The factor $x/\tilde{x}$ in (82) is included to form the invariant momentum integral $\int d^3 p / [(2\pi)^3 2E]$ out of the distribution $P_a^{N_i}$ [89]. In (84) the quantity n^{N_i} is the total number of partons in a given nucleon with momentum $P = |\vec{P}|$ at Q_0^2 .

(ii) The function $\rho_a^A(x)$ in (80) takes into account nuclear shadowing effects affecting mainly soft (small x) partons in a nucleus. These shadowing effects are evident in

the observations of the European Muon Collaboration [90] as a depletion of the nuclear structure functions at small x relative to those of a free nucleon. Several mechanisms have been proposed to explain this nuclear shadowing effect on the basis of the parton model [91, 92, 93, 94]. However, here instead the phenomenological approach of Wang and Gyulassy [41] is adopted which is based on the following parametrization [93] for the A dependence of the shadowing for both quarks and gluons:

$$\rho_a^A(x) \equiv \frac{F_a^{A,N_i}(x, Q_0^2)}{A F_a^N(x, Q_0^2)} = 1 + 1.19 \left(\ln A \right)^{1/6} \left[x^3 - 1.5(x_0 + x_L)x^2 + 3x_0x_Lx \right] (85) \\ - \left[\alpha_A(r) - \frac{1.08(A^{1/3} - 1)}{\ln(A + 1)} \sqrt{x} \right] \exp(-x^2/x_0^2) .$$

where $F_a^{A,N}$ and F_a^N is the structure functions of a nucleus A , respectively those of a free nucleon, and $x_0 = 0.1$, $x_L = 0.7$. The function

$$\alpha_A(r) = 0.1 \left(A^{1/3} - 1 \right) \frac{4}{3} \sqrt{1 - \frac{r^2}{R_A^2}} \quad (86)$$

takes into account the impact parameter dependence, with r labeling the transverse distance of a nucleon from its nucleus center and R_A the radius of the nucleus. Fig. 18 shows the parametrization (86) for $\rho_a^A(x)$ for various nuclei in comparison to the corresponding experimental data.

(iii) The distribution $g(p_\perp)$ specifies the primordial transverse momenta of partons according to a Gaussian $\propto \exp(-|\vec{p}_\perp|^2/p_0^2)$, independent of the type of parton or nucleon. It takes into account the uncertainty of momentum ("Fermi motion") due to the partons confinement within the nucleon. The width $p_0 = 0.42$ GeV corresponds to a mean $\langle p_\perp \rangle \approx 0.38$ GeV, in agreement with values inferred from experimental data [96].

4.1.2 Initial spatial distribution

The initial spatial distribution of the partons, $R_a^{N_i}$, appearing in eq. (79), depends on the magnitude of their momenta, the positions of their parent nucleons and on the spatial substructure of the latter. It is represented as

$$R_a^{N_i}(\vec{p}, \vec{r}, \vec{R}) = \delta^3(\vec{R} - \vec{R}_{AB}) \left[h_a^{N_i}(\vec{r}) H_{N_i}(\vec{R}) \right]_{\text{boosted}} , \quad (87)$$

where the momentum dependence is here purely due to boosting the distributions to the nuclear CM frame (see below).

The components of the ansatz (87) have the following meaning:

(i) The incoming nuclei are assumed to have initial CM positions $\vec{R}_{AB} = (\pm \Delta Z_{AB}/2, \vec{B}_{AB})$, corresponding to a chosen impact parameter $\vec{B}_{AB} = (B_x, B_y)$ and a minimum longitudinal separation $\Delta Z_{AB} (\approx 1 \text{ fm})$. Then, in the rest frame of each nucleus, the individual nucleons are assigned positions around the centers of their parent nuclei according to a Fermi

distribution for nuclei with mass number $A \geq 12$ and a Gaussian distribution for nuclei $A < 12$,

$$H_{N_i}(\vec{R}) = \begin{cases} \frac{1}{4\pi} (1 + \exp[(R - c)/a])^{-1} & (A \geq 12) \\ \frac{1}{4\pi} \frac{2}{\sqrt{\pi b}} \exp[-R^2/b^2] & (A < 12) \end{cases} \quad (88)$$

The parameters are $c = r_0 A^{1/3}$, $r_0 = 1.14 \text{ fm}$, $a = 0.545 \text{ fm}$ and $b = \sqrt{2/3} R_A^{ms}$, where R_A^{ms} is the mean square radius of the nucleus.

(ii) Next, the partons are distributed around the centers of the nucleons, still in the rest frame of the actual nucleus, with an exponential distribution

$$h_a^{N_i}(\vec{r}) = \frac{1}{4\pi} \frac{\nu^3}{8\pi} \exp[-\nu r] \quad , \quad (89)$$

where $\nu = 0.84 \text{ GeV}$ corresponds to the measured elastic formfactor of the nucleon with a mean square radius of $R_N^{ms} \equiv \sqrt{12/\nu} = 0.81 \text{ fm}$.

(iii) Finally, as indicated by the subscript ‘boosted’ in eq. (87), the positions of the nucleons and their valence quarks are boosted into the CM frame of the colliding nuclei. This is illustrated in Fig. 19: The valence quarks then occupy the Lorentz contracted region $(\Delta z)_v \approx 2R_A M_N/P$, whereas The sea quarks and gluons are smeared out in the longitudinal direction by an amount $(\Delta z)_{g,s} \approx \hbar/p_z < 2R_A$ ($R_A = r_0 A^{1/3}$) around the valence quarks. This procedure yields the *distributed Lorentz contraction* which is an important feature of the partons when boosting a nucleus to high rapidities [97, 98, 102]. As a consequence, the parton positions are correlated in longitudinal direction with their momenta, as required by the uncertainty principle. This leads to an enhancement of the densities of gluons and sea quarks with $x < 1/(2R_N M_N)$ proportional to $A^{1/3}$, because such partons from different nucleomns overlap spatially when the nucleons are at the same impact parameter.

4.2 The perturbative evolution of parton cascades

Recall from Chap. 3 that the time development of the phase-space distribution $F_a(p, r)$ for the partons of species a , starting from its given initial form $F_a^{(0)}(Q_0^2, p, r)$ at time $t = t_0$, eq. (79), is governed by the relativistic transport equation (38),

$$p^\mu \partial_\mu F_a(p, r) = \sum_{\text{processes } k} I_a^{(k)}(p, r) \quad , \quad (90)$$

with a collision kernel on the right hand side that interplays with the free streaming term on the left hand side and generates the space-time evolution of the many-parton system according to the specific perturbative QCD interactions of the partons.

The task is therefore to specify the form of collision term $\sum_k I_a^{(k)}$, i.e. the parton interaction processes to be considered. The general form of the collision term, eq. (39),

includes all possible $n \rightarrow n'$ interaction processes. Each term of the sum over contributing processes is a phase-space integral involving the corresponding squared matrix elements $|\mathcal{M}_{n \rightarrow n'}|^2$. In the PCM a subset of $n \rightarrow n'$ processes is considered which consists of $2 \rightarrow 2$ collisions, $2 \rightarrow 1$ fusions and $1 \rightarrow 2$ decays in lowest order perturbation theory, plus higher order corrections which are associated with multiple emissions and absorptions of partons. Correspondingly, the collision term is represented as a sum over all possible elementary $2 \rightarrow 2$, $2 \rightarrow 1$ and $1 \rightarrow 2$ parton interactions, in which each of the partons coming in and going out of the vertices is "dressed" by a specific formfactor that inclusively sum up the higher order corrections associated with the leading logarithmic collinear and infrared singularities of QCD. This concept - schematically illustrated in Fig. 20 - is the logic extension of the picture underlying the (M)LLA framework sketched in Chap. 2, namely the very idea that the structure of parton interactions is build up from elementary "blocks" that depend on just the nearest ancestors. The rate of time change exhibits the property of locality without any remembrance of the pre-history.

As explained in Sec. 3.1, the collision term $\sum_k I_a^{(k)} = \sum_k (I_a^{(k)gain} - I_a^{(k)loss})$ locally changes the phase-space distribution $F_a(p, r)$ of parton species a at the space-time point $r = (t, \vec{r})$ due to interactions in which a parton a may be gained or lost in a phase-space element $dQ^2 d^3p / [(2\pi)^3 2E]$. Embedding the above concept into the collision term, one can represent eq. (39) in the form

$$\sum_{\text{processes } k} I_a^{(k)}(p_1, r) = \sum_{b,c,d} J_{abcd}(p_1, r) + \sum_{b,c} K_{abc}(p_1, r) . \quad (91)$$

The two distinct types of interaction integrals J and K are given by the difference of gain and loss terms (with respect to partons of species a with four-momentum p_1) for $2 \rightarrow 2$ and for $2 \rightarrow 1$, $1 \rightarrow 2$ processes, respectively. Specifically,

$$\begin{aligned} J_{abcd}(p_1, r) &= j_{cd \rightarrow ab}^{gain}(p_1, r) - j_{ab \rightarrow cd}^{loss}(p_1, r) \\ &= -\frac{1}{2} \frac{1}{S_{ab}} \frac{1}{S_{cd}} \int \frac{d^3 p_2}{(2\pi)^3 2E_2} \int \frac{d^3 p_3}{(2\pi)^3 2E_3} \int \frac{d^3 p_4}{(2\pi)^3 2E_4} (2\pi)^4 \delta^4(p_1 + p_2 - p_3 - p_4) \\ &\quad \times \left\{ F_a(1) F_b(2) [1 \pm F_c(3)] [1 \pm F_d(4)] |\overline{\mathcal{M}}_{ab \rightarrow cd}|_{eff}^2 \right. \\ &\quad \left. - F_c(3) F_d(4) [1 \pm F_a(1)] [1 \pm F_b(2)] |\overline{\mathcal{M}}_{cd \rightarrow ab}|_{eff}^2 \right\} \end{aligned} \quad (92)$$

and

$$\begin{aligned} K_{abc}(p_1, r) &= k_{c \rightarrow ab}^{gain}(p_1, r) - k_{ab \rightarrow c}^{loss}(p_1, r) \\ &= -\frac{1}{2} \frac{1}{S_{ab}} \int \frac{d^3 p_2}{(2\pi)^3 2E_2} \int \frac{d^3 p_3}{(2\pi)^3 2E_3} (2\pi)^4 \delta^4(p_1 + p_2 - p_3) \\ &\quad \times \left\{ F_a(1) F_b(2) [1 \pm F_c(3)] |\overline{\mathcal{M}}_{ab \rightarrow c}|_{eff}^2 \right. \end{aligned} \quad (93)$$

$$- F_c(3) [1 \pm F_a(1)] [1 \pm F_b(2)] |\overline{\mathcal{M}}_{c \rightarrow ab}|_{eff}^2 \} .$$

Here the notation is $F_\alpha(i) \equiv F_\alpha(p_i, r)$ for the distribution functions of parton species $\alpha = a, b, c, d$ and corresponding four-momenta $p_i = p_1, p_2, p_3, p_4$ at space-time point $r = (t, \vec{r})$. The squared matrix elements, characteristic for the types of interactions, are weighted by a distribution function F_α for each of the particles coming into the interaction vertex and a factor $[1 \pm F_\alpha]$ for each of the outgoing ones. The $-$ sign refers to quarks and antiquarks with $[1 - F_q]$ indicating Pauli blocking, and the $+$ sign is for gluons so that $[1 + F_g]$ results in a Bose enhancement. The statistical factors $1/S_{ab} = 1/(1 + \delta_{ab})$ and $1/S_{cd} = 1/(1 + \delta_{cd})$ account for the identicle particle effect when the two incoming and/or outgoing partons are indistinguishable. Finally, the factors $1/2$ in front of the right hand side of eqs. (92) and (94) result from the normalization $1/(2E_1)$.

The *effective squared matrix elements* $|\overline{\mathcal{M}}|_{eff}^2$ are expressed in terms of the lowest order matrix elements, $|\overline{M}^{(0)}|^2$ summed over spin and color, and are multiplied by S (space-like) and T (time-like) formfactors for the partons coming in, respectively going out of, the vertex (see Fig. 20). That is,

$$|\overline{\mathcal{M}}_{ab \rightarrow cd}|_{eff}^2 = S_a(p_a; Q^2, Q_0^2) S_b(p_b; Q^2, Q_0^2) \left[|\overline{M}_{ab \rightarrow cd}^{(0)}|^2(Q^2) \right] T_c(Q^2, \mu_0^2) T_d(Q^2, \mu_0^2) \quad (94)$$

$$|\overline{\mathcal{M}}_{ab \rightarrow c}|_{eff}^2 = S_a(p_a; p_c^2, Q_0^2) S_b(p_b; p_c^2, Q_0^2) \left[|\overline{M}_{ab \rightarrow c}^{(0)}|^2(p_c^2) \right] T_c(p_c^2, \mu_0^2) \quad (95)$$

$$|\overline{\mathcal{M}}_{c \rightarrow ab}|_{eff}^2 = \left[|\overline{M}_{c \rightarrow ab}^{(0)}|^2(p_c^2) \right] T_a(p_c^2, \mu_0^2) T_b(p_c^2, \mu_0^2) \quad (96)$$

The lowest order invariant squared matrix elements $|\overline{M}^{(0)}|^2$ are well known for the elementary $2 \rightarrow 2$, $2 \rightarrow 1$ and $1 \rightarrow 2$ processes and the formfactors will be specified in Secs. 4.2.3 and 4.2.4. The distribution functions $F_\alpha(p, r)$ in (92) and (94) on the other hand are not available in analytic form since they are determined by the dynamics itself. However, they can be dynamically calculated during the cascade evolution at each point of time with the cellular phase-space method described in Sec. 3.4.

The physical significance of the effective matrix elements (94)-(96) is the following:

- (i) In (94), Q^2 specifies the invariant scale at which the interaction occurs. It generally depends on the four-momenta of the scattering partons and is not unambiguous in lowest order perturbation theory. It is chosen to be the transverse momentum of the scattered partons in their center-of-mass frame, $Q^2 = p_\perp^2$, except for $q\bar{q}$ annihilation where $Q^2 = (E_{q\bar{q}}^2)_{cm}$. The value of Q^2 determines the strength of the QCD coupling $\alpha_s(Q^2)$ associated with the vertex.

- (ii) Q_0^2 is the resolution scale (78) at which the colliding nuclei initially were resolved into their parton substructure and μ_0^2 is an invariant mass cut-off below which parton emission is considered to be unresolvable. The values of Q_0^2 and μ_0^2 define the initial and final point, respectively, in between which the evolution of parton cascades is followed perturbatively. The non-perturbative dynamics outside this regime must be absorbed in the initial parton distributions (Sec. 4.1) depending on Q_0^2 and in the final hadronization of partons (Sec. 4.3) which takes effect below μ_0^2 .
- (iii) The representations (94)-(96) may be pictured as elastic processes with the incoming and outgoing partons being off mass shell or "dressed", corresponding to higher order inelastic contributions. Setting the formfactors S and T equal to unity therefore corresponds to treating the external particles as bare quanta on mass shell (see Fig. 20).
- (iv) The phenomenological difference between the S and T formfactors is important to realize. The S formfactors are of relevance only for primary space-like partons that evolve directly from the initial nuclear parton clouds and encounter their first interaction. For all other (secondary) partons, those which have either emerged from a scattering or been produced in a decay process, S is equal to unity. Furthermore, one might get the impression that the T formfactors only "dress" those partons that emerge from an interaction vertex. However, these "dressed" partons can subsequently interact again, so that the T formfactors are implicitly present also in those interactions where time-like partons are incoming to the vertex. To avoid double-counting, therefore only the partons receding from a vertex are "dressed" by a T formfactor in (94)-(96) as illustrated in Fig. 20.

To give an idea how the space-time evolution of the parton distributions in a nuclear collision according to the transport equation (90) is described in terms of parton cascades, a typical parton cascade development is illustrated in Fig. 21. It is important to realize that in a nuclear collision there are many such cascades, which are internetted and simultaneously evolving. Each cascade can be subdivided into elementary $2 \rightarrow 2$ scatterings, $1 \rightarrow 2$ branchings (emissions), and $2 \rightarrow 1$ fusions (absorptions). In Fig. 21, a primary parton a that originates from one of the incident nuclei, collides with another parton b with some momentum transfer $Q_{ab}^2 = p_\perp^2$. The parton a has evolved from the initial scale Q_0^2 , at which it was resolved in its parent nucleus, up to Q_{ab}^2 by successive space-like branchings. From the scattering of a and b the partons c and d emerge, both of which can initiate sequences of time-like branchings. These newly produced partons can themselves radiate, rescatter, and/or absorb (be absorbed by) other partons. This cascade development corresponds to

the formal representation of the effective matrix-elements (94)-(96), depicted in Fig. 20. These effective matrix-elements enter the collision integrals in the transport equation (90) and generate the evolution of the parton system in terms of multiple parton scatterings and associated emission/absorption processes.

4.2.1 Elementary parton-parton collisions

For the elementary parton scatterings $a + b \rightarrow c + d$ as in Fig. 21, two distinct classes of processes are considered: (i) truly perturbative QCD hard and semihard parton scatterings; (ii) phenomenological treatment of soft parton interactions with as low momentum transfer. The motivation for this differentiation of parton-parton collisions is to smoothen the singular behavior of the collision integrals (91) which results from the divergence of the squared matrix elements (94) for small momentum transfers (except for $q\bar{q}$ annihilation). To render the parton-parton cross-sections finite, an invariant cut-off scale $Q_{cut}^2 \equiv p_{\perp cut}^2$ is introduced such that collisions with $p_{\perp} \geq p_{\perp cut}$ are treated as (semi)hard scatterings within QCD perturbation theory, whereas for $p_{\perp} < p_{\perp cut}$ a soft interaction is assumed to occur. That is, the total parton-parton cross-section for a collision between two partons a and b is represented as

$$\hat{\sigma}_{ab}(\hat{s}) = \sum_{c,d} \left\{ \int_0^{p_{\perp cut}^2} dp_{\perp}^2 \left(\frac{d\hat{\sigma}_{ab \rightarrow cd}^{soft}}{dp_{\perp}^2} \right) + \int_{p_{\perp cut}^2}^{\infty} dp_{\perp}^2 \left(\frac{d\hat{\sigma}_{ab \rightarrow cd}^{hard}}{dp_{\perp}^2} \right) \right\} . \quad (97)$$

Here $p_{\perp}^2 \simeq \hat{t}\hat{u}/\hat{s}$ is the relative transverse momentum produced by the scattering, defining the interaction scale $Q^2 \equiv p_{\perp}^2$, the variables $\hat{s}$, $\hat{t}$ and $\hat{u}$ are the usual Mandelstam variables and the sum over c, d corresponds to summing over all possible reaction channels (processes). The specific values of $p_{\perp cut}$ generally can depend on beam energy as well as nuclear mass number, and must be fixed from existing experimental data (as will be explained in Sec. 4.4 below).

The complementation of both hard and soft contributions renders the parton cross-section finite and well defined for *all* $p_{\perp}$. This regularization scheme for the cross-section (97) is illustrated in Fig. 22. It must be emphasized that the phenomenological inclusion of soft scatterings serves here only for the purpose of having a smooth behavior of $d\hat{\sigma}/dp_{\perp}^2$ around the cut-off $p_{\perp cut}^2$. The effect of soft scatterings on the global dynamics is not essential, since these processes occur considerably less frequent than the (semi)hard scatterings. Moreover, the soft scatterings naturally involve comparably small momentum transfer, so that their contribution to transverse energy production is neglectable.

The following $2 \rightarrow 2$ interaction processes are included:

$$\begin{aligned} gg \rightarrow gg , \quad & gg \rightarrow q\bar{q} , \quad & qg \rightarrow qg , \quad & \bar{q}g \rightarrow \bar{q}g , \\ qq \rightarrow qq , \quad & \bar{q}\bar{q} \rightarrow \bar{q}\bar{q} , \quad & q\bar{q} \rightarrow q'\bar{q}' , \quad & q\bar{q} \rightarrow gg . \end{aligned} \quad (98)$$

In accord with the two-component structure of the cross-section (97), these scattering processes are distinguished depending on the momentum transfer (Fig. 22):

(i) **Hard and semihard collisions above $p_{\perp cut}$:** For these perturbatively calculable processes the momentum transfer $\hat{t}$ is determined by the For the *perturbative (semi)hard collisions* above $p_{\perp cut}$, the momentum transfer $\hat{t}$ is determined by the corresponding differential cross-sections

$$\frac{d\hat{\sigma}_{ab \rightarrow cd}^{hard}}{d\hat{t}} = \frac{1}{16\pi\hat{s}^2} |\overline{M^{(0)}}_{ab \rightarrow cd}|^2 (Q^2, \hat{s}, \hat{t}, \hat{u}) , \quad (99)$$

where $\hat{s} = (p_a + p_b)^2$, $\hat{t} = (p_a - p_c)^2$, $\hat{u} = (p_a - p_d)^2$, $Q^2 = p_{\perp}^2 \simeq \hat{t}\hat{u}/\hat{s} \simeq -\hat{t}$ as before, and $|\overline{M^{(0)}}|^2$ are the spin and color averaged squared matrix elements in lowest order perturbation theory that appear in eqs. (94)-(96). For the processes (98) the cross-sections (99) are published in the literature (see e.g. Refs. [46, 103] and for massive quarks Refs. [104, 105, 106]).

(ii) **Soft collisions below $p_{\perp cut}$:** In the case of a non-perturbative soft collision between two partons it is assumed that a very low energy double gluon exchange occurs. This provides a natural continuation to the harder collisions above $p_{\perp cut}$ where the dominant one gluon exchange processes $gg \rightarrow gg$, $gq \rightarrow gq$ and $qq \rightarrow qq$ have the same overall structure [107]. The momentum transfer $\hat{t}$ between two soft interacting partons is chosen according to the simple phenomenological distribution

$$\frac{d\hat{\sigma}_{ab \rightarrow cd}^{soft}}{d\hat{t}} = f(P) \exp \left[\frac{\hat{t} - \hat{t}_{cut}}{\hat{t}_0} \right] , \quad (100)$$

where $\hat{t}_0 = 1 \text{ GeV}^2$ and $\hat{t}_{cut} = -\hat{s}/2 \left(1 - \sqrt{1 - 4p_{\perp cut}^2/\hat{s}} \right)$. The normalization factor $f(P)$ depends on $P = P_{CM}/N_{nuc}$ and is determined by the requirement that the sum $\hat{\sigma}^{inel} = \hat{\sigma}^{soft} + \hat{\sigma}^{hard}$ per nucleon corresponds to the measured inelastic pp ($p\bar{p}$) cross-sections for given beam energy P . Thus, the soft contribution adds to the total inelastic cross-section what is left out by (semi)hard parton collisions.

4.2.2 Virtual and real corrections to the elementary processes

The important perturbative QCD radiative corrections associated with the collision $a + b$ occur in two distinct contexts, both of which are evident in Fig. 21: *space-like and time-like parton cascades*. First, if the scattering partons a or/and b are primary quanta, they may have undergone a sequence of space-like branchings before their interaction and thereby have evolved from the initial resolution scale Q_0^2 up to the interaction scale associated with the scattering vertex. Second, each of the secondary quanta c and d emerging from the vertex $a + b \rightarrow c + d$ may subsequent to its production initiate its own time-like cascade by multiple successive emissions of gluons or branchings into $q\bar{q}$ pairs. The subtle difference

between space-like and time-like parton cascades is that in the former a primary parton evolves from the initial scale Q_0^2 (78), at which it was originally resolved in the nuclear structure functions, up to the scale Q^2 of the scattering vertex, whereas in the time-like case a secondary parton produced at the vertex of the collision evolves, starting from Q^2 , down to the cut-off scale μ_0^2 . In any case, the produced secondary partons can either themselves emit additional partons, or absorb an already present close-by parton, or, as a third possibility collide with another parton as described before.

Associated with this intuitive picture are three types of corrections to the elementary free-space parton cross-section which corresponds to the lowest order matrix-elements $|\overline{M}^{(0)}|^2$ in the effective matrix-elements (94)-(96). These corrections are embodied in the formfactors of the partons coming in and going out of an elementary vertex as shown in Fig. 20, and originate from:

- (i) the coherence of soft gluon emission,
- (ii) the competition between emission and absorption processes,
- (iii) the Landau-Pomeranchuk-Migdal effect.

(i) **Coherence of soft gluon emission:** As repeatedly emphasized before, one of the main QCD properties of the partons is that they are, loosely speaking, extended off-shell quanta with an inner quark and gluon substructure. Formally this fact is inherent in the Sudakov formfactors that correspond to the inclusive summation of (leading) virtual graphs to all orders and that control the parton branching probabilities for real emission processes. The dominant contributions to the formfactors arise from both *collinear* and *infrared* singularities of the branching amplitudes. The LLA framework introduced in Chap. 2 only accounts for collinear, hard (large x) parton branchings. However, the bulk of gluon radiation comes from the infrared region of soft (small x) branchings. With the extension of the MLLA framework, mentioned in Sec. 2.5, it is straightforward to include this "coherent soft gluon emission" in the probabilistic scheme of successive branchings, including essential interference effects. These issues have been studied extensively in particle physics. For details I refer to e.g. the reviews [52, 73, 74, 50]. The main point is that coherent soft gluon radiation can be consistently accounted for by so-called *angular ordering* of successive emissions. That is, instead of ordering a sequence of branchings according to the parton virtualities Q^2 as in (18) or (22), one employs the angular variable ξ that orders the emission angles as one moves along the branching cascade. This automatically incorporates the coherence of different emissions which leads to a significant suppression of soft gluon production due to destructive interference [70, 68]. This effect, for which exists strong experimental evidence in e^+e^- annihilation and in $p\bar{p}$ collisions [108], is of particular

importance when the number of soft gluons becomes large, as is typically the case during the course of a heavy ion collision.

(ii) **Competitive emission and absorption:** In a collision of two heavy nuclei the density of partons in the central region can become rather large during the reaction: First, partons from different nucleons overlap spatially, and second, there is an enhanced production of additional, new partons through multiple emission processes. Therefore the reverse processes of parton absorption become important. The net emission rate will decrease until emissions and absorptions locally reach a detailed balance. In Ref. [82] it was explained how one can take into account absorptive processes depending on the parton densities in phase-space by modifying the emission probability of partons in sequential branchings to an effective probability for competitive emission and absorption. The resulting expression recovers the ordinary emission probability given by the branching functions (16), when the density of partons in phase-space goes to zero. However, if the parton density grows during the nuclear collision, the effect is that emission processes will be increasingly suppressed until a balance between emissions and absorptions is reached.

(iii) **Landau-Pomeranchuk-Migdal effect:** The radiation emitted by a virtual parton has a specific formation time which results in the Landau-Pomeranchuk-Migdal effect [109]. If the density of partons becomes large, so that there is a good chance that a parton undergoes multiple scatterings in a short time interval, then the radiation this parton wants to emit is suppressed. The Landau-Pomeranchuk-Migdal effect is a consequence of the specific space-time evolution of radiation emitted by an accelerated charge and can be observed in the bremsstrahlung of electrons [109, 110]. When a charge undergoes multiple scatterings in a medium and the time between the successive scatterings is shorter than the ‘formation time’ of the radiation, then the contributions from intermediate scatterings cancel out, resulting in an effective suppression of radiation. This effect is expected to occur also for the radiative gluon emissions of partons in relativistic heavy ion collisions [110], where partons may encounter multiple scatterings shortly after each other. A heuristic way to incorporate this effect into the framework of multiple parton emissions was introduced in Ref. [82] on the basis of the formation time of radiation from a virtual parton such that this finite time scale delays an instantaneous, irreversible emission and mimics the possibility that the parton rescatters before it radiates. For details see Ref. [82].

In the following I will introduce generalized parton formfactors that extend the LLA expressions of Chap. 2 to include also these elements (i)-(iii).

4.2.3 Space-like parton cascades

Recall that space-like parton cascades are associated with the QCD radiative corrections due to bremsstrahlung emitted by primary space-like partons that encounter their very first collision and are thereby struck out of the coherent nuclear wavefunction to become free.

A typical space-like cascade is illustrated in Fig. 21, where the parton a_0 , originating from the initial nuclear wavefunction, undergoes successive space-like branchings $a_0 \rightarrow a_1 k_1$, $a_1 \rightarrow a_2 k_2$, ..., $a_{n-1} \rightarrow a_n k_n$ to become the parton $a \equiv a_n$ which then actually collides with another parton b . The branching chain proceeds by increasing the virtualities of the partons a_i in the cascade, starting from a_0 with $q_0^2 \simeq -Q_0^2$ up to $q^2 \simeq -Q^2$ the space-like virtuality of the scattering parton a . The partons k_i on the side branches on the other hand have, due to energy-momentum conservation at the branching vertices, time-like virtualities and each of them can initiate a time-like cascade as will be described below.

The proper inclusion of both collinear, hard and coherent, soft branchings is achieved by simply replacing parton virtualities $Q_i^2 = -q_i^2$ as evolution variable, with the new variable [68, 111]

$$\tilde{Q}_i^2 \equiv E_i^2 \tilde{\xi}_i, \quad \tilde{\xi}_i = \frac{q_0 \cdot p_{i+1}}{E_0 \omega_{i+1}}, \quad (101)$$

where $q_i = (E_i, \vec{q}_i)$ and $p_i = (\omega_i, \vec{p}_i)$ refer to the partons a_i , respectively k_i in Fig. 21. This corresponds to replacing the ordering in virtualities $Q_{i+1}^2 > Q_i$ by $\tilde{Q}_{i+1}^2 > \tilde{Q}_i^2$ which is equivalent to the angular ordering $E_i \theta_{q_0 p_{i+1}} > E_{i-1} \theta_{q_0 p_i}$.

The method used here to evolve a space-like cascade is the "backward evolution scheme" adopted from Ref. [66], where a detailed presentation can be found. To sketch the procedure, consider the space-like branching $a_{n-1} \rightarrow a_n k_n$ in Fig. 21, where the virtualities of the partons a_{n-1} , a_n and k_n are q_{n-1}^2 , q_n^2 and p_n^2 , respectively, with $|q_n^2| < |q_{n-1}^2|$, and $q_n^2, q_{n-1}^2 < 0$ (space-like) but $p_n^2 > 0$ (time-like). The relative probability for this branching to occur between $\tilde{Q}^2$ and $\tilde{Q}^2 + d\tilde{Q}^2$ is given by

$$d\mathcal{P}_{a_{n-1}a_n}^{(S)}(x_{a_{n-1}}, x_{a_n}, \tilde{Q}^2) = \frac{\alpha_s ((1-\tilde{z})\tilde{Q}^2)}{2\pi\tilde{Q}^2} \left(\frac{F_{a_{n-1}}^N(x_{a_{n-1}}, \tilde{Q}^2)}{F_{a_n}^N(x_{a_n}, \tilde{Q}^2)} \right) \frac{1}{\tilde{z}} P_{a_{n-1} \rightarrow a_n k_n}(\tilde{z}) d\tilde{Q}^2 d\tilde{z}. \quad (102)$$

Here $x_\alpha = (p_\alpha)_z/P$ for $\alpha \equiv a_{n-1}, a_n$ are the fractional longitudinal momenta of the parent nucleons carried by the partons a_{n-1} and a_n , $F_\alpha^N(x_\alpha, \tilde{Q}^2)$ are the corresponding nucleon structure functions that enter the initial parton distributions (79). The variables

$$\tilde{z} = \frac{E_n}{E_{n-1}} \simeq \frac{x_{a_n}}{x_{a_{n-1}}}, \quad 1 - \tilde{z} = \frac{\omega_n}{E_{n-1}} \simeq \frac{x_{a_{n-1}} - x_{a_n}}{x_{a_{n-1}}} \quad (103)$$

specify the fractional energy or longitudinal momentum of parton a_n and k_n , respectively, taken away from a_{n-1} .

The "backwards evolution" of the space-like branching $a_{n-1} \rightarrow a_n + k_n$ is expressed in terms of the probability that parton a_{n-1} did *not* branch between the lower bound given by the initial resolution scale Q_0^2 and $\tilde{Q}^2$. In that case parton a_n can *not* originate from this branching, but must have been produced otherwise or already been present in the initial

parton distributions. This non-branching probability is given by the *Sudakov form-factor for space-like branchings* [c.f. (21)]:

$$S_{a_n}(x_{a_n}, \tilde{Q}^2, Q_0^2) = \exp \left\{ - \sum_{a_{n-1}} \int_{Q_0^2}^{\tilde{Q}^2} \int_{\tilde{z}_-(\tilde{Q}')}^{\tilde{z}_+(\tilde{Q}')} d\mathcal{P}_{a_n a_{n-1}}^{(S)}(x_{a_n}, \tilde{z}, \tilde{Q}'^2) \right\} . \quad (104)$$

The upper limit of the $\tilde{Q}^2$ -integration is set by $\tilde{Q}^2 \approx Q_{ab}^2$ associated with the scale of the scattering vertex of parton $a_n \equiv a$ with another parton b in Fig. 21. The limits $\tilde{z}_\pm$ are determined by kinematics [68]: $\tilde{z}_-(\tilde{Q}) = Q_0/\tilde{Q}$ and $\tilde{z}_+(\tilde{Q}) = 1 - Q_0/\tilde{Q}$.

I would like to mention at this point that in nuclear collisions the contribution of space-like cascades to the dynamics of the parton system as a whole turns out to be insignificant relative to the time-like cascades (as can be read off Tables 1 and 3 in Chap. 5), because they are only of relevance for primary partons involved in their first collisions with a interaction scale $Q^2 \gg Q_0^2$. Since the value of Q_0^2 is estimated according to eq. (78), only rather hard primary scatterings, which are rare fluctuations, give rise to a space-like cascade.

4.2.4 Time-like parton cascades

Time-like parton cascades are initiated by secondary partons that either emerge from a scattering or fusion process, or are produced by a radiative emission. Consider the time-like cascade initiated by the parton c in Fig. 21. This parton has been produced in the collision $a + b \rightarrow c + d$ with a time-like off-shellness $m_c^2 \simeq Q^2 (\equiv Q_{ab}^2)$. An ‘effective’ probability $d\mathcal{P}_{cf}^{(T)}$ for the subsequent branching $c \rightarrow fg$ of parton c with virtuality m_c^2 into partons f and g with $m_f^2, m_g^2 < m_c^2$, has been proposed in Ref. [82] as an extension of the usual description of the branching process (c.f. Chap. 2), namely,

$$d\mathcal{P}_{cf}^{(T)}(\xi, z) = \Pi_c(m_c^2, \Delta t) \left(1 - \frac{F_f(z, \xi)}{1 \pm F_f(z, \xi)} \right) \frac{\alpha_s(Q^2)}{2\pi\xi} \gamma_{c \rightarrow fg}(z) dz d\xi . \quad (105)$$

It is formulated in terms of the energy fractions carried by the partons f and g ,

$$z = \frac{E_f}{E_c} , \quad 1 - z = \frac{E_g}{E_c} , \quad (106)$$

the angular evolution variable [52, 69, 65]

$$\xi = \frac{p_f \cdot p_g}{E_f E_g} \simeq 1 - \cos \theta_{fg} , \quad (107)$$

and the virtuality m_c^2 of parton c , which is related to z and ξ through

$$m_c^2 = m_f^2 + m_g^2 + 2E_c^2 z(1-z)\xi . \quad (108)$$

It has been shown [69, 65] that for the branching process $c \rightarrow fg$ described by eq. (105), the scale Q^2 in the running coupling strength $\alpha_s(Q^2)$ must be identified with

$$Q^2 = 2z^2(1-z)^2 E_c^2 \xi \simeq k_\perp^2 , \quad (109)$$

which is approximately the relative transverse momentum squared between the products f and g at the vertex.

The branching probability (105) consists of three characteristic terms:

- a) $\Pi_c(m_c^2, \Delta t)$ is the probability that parton c with virtuality m_c^2 and a corresponding ‘life-time’ $\tau \sim 1/m_c$ decays (branches) within a time interval Δt , taken to be the discrete time steps $\Delta t = 10^{-2} fm$ along which the cascade development is followed in time.
- b) The term in brackets accounts for the possibility of parton c to absorb a parton f to yield a parton g , that is $c + f \rightarrow g$, rather than to branch into partons f and g via $c \rightarrow f + g$. The relative probability for this to happen is determined by the phase-space density $F_f(z, \xi)$ of partons of species f within $d\xi dz$ around ξ and z .
- c) The function $\alpha_s \gamma_{c \rightarrow fg} / (2\pi\xi)$ is the branching probability (16) associated with the DGLAP evolution equations (15), where $\gamma_{c \rightarrow fg}(z)$ is the appropriate branching kernel that describes the energy distribution in the process.

It is important to realize that the specific form (105) of $d\mathcal{P}_{cf}^{(T)}$ incorporates the physics elements explained in Sec. 4.2.2. It is based on the coherent description of the branching in terms of the variable ξ , taking into account the destructive interference of soft emitted gluons which has been shown to significantly reduce the available phase-space for the branching to occur. Furthermore the expression (105) extends the usual description of time-like parton cascades, developed originally for QCD jet studies in e^+e^- -annihilation and high energy pp ($p\bar{p}$) collisions, to include two new features that become very important in highly relativistic nucleus-nucleus collisions. These are, first, the Landau-Pomeranchuk-Migdal effect resulting from a finite formation time of radiative gluon emissions (effectively described by the function Π_c), and second, the fact that in heavy ion collisions a dense phase-space occupancy of partons is expected to lead to an enhancement of absorption processes relative to emissions which eventually establishes a detailed balance (expressed by the term in brackets containing the phase-space density F_f of partons f). In addition to the soft gluon interference, both of these latter features also lead to an effective suppression of radiative emissions and consequently to a reduced particle multiplicity. Whereas the time delay of a parton emission connected with the Landau-Pomeranchuk-Migdal effect allows a parton to rescatter before it is able to radiate, the increasingly probable absorption of partons in dense phase-space regions leads to a decreasing branching probability.

The probability that parton c does *not* branch between ξ and a minimum value ξ_{min} is given by the exponentiation of (105), yielding the specific *Sudakov form-factor for time-like*

branchings. [c.f. (25)]:

$$T_c(\xi, \xi_{min}) = \exp \left\{ - \int_{\xi_{min}}^{\xi} \sum_f \int_{z_-(\xi', E_c)}^{z_+(\xi', E_c)} d\mathcal{P}_{cf}^{(T)}(\xi', z) \right\} . \quad (110)$$

The cut-offs ξ_{min} and $z_{\pm}$ have to be introduced to regularize collinear divergences (in z) and infrared divergences (in ξ). They are determined by the requirement that the branching must terminate when the virtual parton mass m_c^2 drops below some critical invariant scale μ_0^2 . This condition leads to a dependence on the cut-off μ_0 , namely, $\xi_{min} = 4\mu_0^2/E_c^2$ and $z_+(\xi, E_c) = 1 - z_-(\xi, E_c) = \mu_0/\sqrt{\xi}E_c$, so that for $z_+(\xi_{min}, E_c) = z_-(\xi_{min}, E_c) = 1/2$ the phase-space for the branching vanishes. To specify a process independent cut-off mass μ_0 , effective parton masses may be defined as [70]

$$m_{eff}^{(g)} = \frac{1}{2} \mu_0 ; \quad m_{eff}^{(q)} = \sqrt{\frac{1}{4} \mu_0^2 + m_q^2} , \quad (111)$$

where m_q are the ordinary flavor-dependent current quark masses. The minimum masses for gluons and quarks are then given by

$$\mu_0^{(g)} = 2 m_{eff}^{(g)} = \mu_0 ; \quad \mu_0^{(q)} = m_{eff}^{(q)} + m_{eff}^{(g)} . \quad (112)$$

The Sudakov formfactor (110) governs the development of a time-like cascade as sequential branchings, until the partons rescatter or are being absorbed, in which cases the respective branch of the cascade terminates. On the other hand, an individual branch within a cascade also terminates when the virtualities of the partons at a branching vertex fall below the infrared cut-off μ_0 .

4.3 Cluster formation and hadronization

The perturbative evolution of the ensemble of parton cascades proceeds until the majority of produced partons have harnessed most of their initial kinetic energy in collisions or branchings and their virtualities have dropped to $\simeq \mu_0$. At this point the residual parton interactions reduce to purely elastic collisions and the system eventually approaches free streaming, with the partons freely moving apart. However, the presence of long range color forces implies that a set of partons can evolve independently only if they form a color singlet system. Therefore the task is to identify appropriate sets of partons which evolve beyond the perturbative cut-off scale μ_0 , defined by eqs. (111) and (112), and separately form hadrons. The important assumption is here that the process of hadron formation depends only on the local structure of the parton system over small space-time volumes.

The scheme to model the non-perturbative mechanism of hadron formation is based on the approaches of Refs. [112, 69]. The central idea is that at the end of the perturbative

phase of parton cascade development, color singlet clusters of partons are formed which subsequently decay independently into hadrons. This concept follows naturally from the *preconfinement* property [113]. Preconfinement is the tendency of the partons produced in the individual cascades to be arranged in color singlet clusters with limited extension in both position and momentum space. It is natural to suppose that these clusters are the basic units out of which hadrons arise non-perturbatively. The preconfinement picture is based on the observation that partons which are correlated in color space due to mutual interactions, have in perturbation theory a damped invariant mass distribution [10] with an average mass of the order of μ_0 . The usual approach to identify sets of partons that are "close in color", is to employ a description in terms of color diagrams [69, 68] which connect interacting partons among each other and keep track of the color flow in and out of a vertex. Such color connected systems of partons are the minimal color neutral clusters [114] that may be taken to form hadrons independently. Exploiting the property that partons which are close in color (in particular minimal color singlets) are also close in phase-space [52], the approach adopted here is, instead of using a color flow description, to forget about the color structure during the cascade development. Rather, at the end of the perturbative evolution color neutral clusters are formed from the final partons which have a minimal separation in coordinate and momentum space. Since the cluster-hadronization model is presented in very detail in Ref. [39], I will here only briefly summarize the phenomenology.

Both, the cluster formation scheme from the collection of quarks and gluons at the end of the perturbative phase, as well as the subsequent cluster decay into final hadrons, consists of two components:

- (i) The rearrangement of the secondary partons, being time-like or on mass shell, their conversion into color singlet clusters (*central clusters*) and the subsequent decay into secondary hadrons.
- (ii) The recombination of the primary, space-like partons that remained spectators throughout the nuclear reaction (*beam clusters*) and the fragmentation of these clusters.

4.3.1 Cluster formation

(i) **Central clusters:** Central clusters are formed from secondary partons, i.e. those that have interacted at least once or have been created during the perturbative evolution. The first step in the cluster formation scheme is to forcibly split each secondary gluon into a collinear quark-antiquark pair. It is assumed that this occurs through a non-perturbative enhancement below μ_0 of the splitting $g \rightarrow q\bar{q}$, which is relatively uncommon in the perturbative branching. This leaves an ensemble of final state quarks and antiquarks which will have to be rearranged to form color singlet clusters. The next step is to employ the

preconfinement picture mentioned before: each quark is assigned to an antiquark sibling (distinct from its antiquark partner in the gluon splitting), which is closest to it in phase-space and therefore in the average also the closest one in color space. The phase-space distance measure Δ_{ij} between two partons, labelled with indices i and j , is chosen to be [115] $\Delta_{ij} = \sqrt{r_{ij}^\mu r_{ij,\mu}} \sqrt{p_{ij}^\nu p_{ij,\nu}}$, where $r_{ij} = r_i - r_j$ and $p_{ij} = p_i - p_j$. Accordingly quark-antiquark clusters $q_i \bar{q}_j$ are formed by measuring for each quark q_i its phase-space separation Δ_{ij} with respect to every antiquark $\bar{q}_j$ and selecting the one that has the minimum distance $\min(\Delta_{i1}, \dots, \Delta_{ij}, \dots, \Delta_{in})$ to be the sibling antiquark of q_i . This scheme, illustrated in Fig. 23, leads to a particular phase-space distribution of color singlet $q_i \bar{q}_j$ clusters. In Fig. 24 a typical invariant cluster mass distribution calculated for $p\bar{p}$ collisions with various beam energies $\sqrt{s} = 200, 546, 900, 1800$ GeV is shown. It exhibits the characteristic features of preconfinement: the distribution is peaked at low mass, strongly damped at large mass and is roughly independent of the beam energy.

(ii) **Beam clusters:** The remaining fraction of the longitudinal momentum and energy that has not been redirected and harnessed in parton interactions is carried by the primary partons of the initial nuclei. These initial state partons that remained spectators throughout the nuclear collision have maintained their originally assigned momenta and their space-like virtualities (except for the valence quarks which are on mass shell) and make up the two fast receding beam fronts. Whereas the primary valence quarks represent the baryon number density that remains in the the beam fronts, the space-like sea quarks and gluons are to be interpreted as virtual excitations of the color field around the valence quarks and therefore must contribute to the internal energy of the beam fragments. They may be pictured as the coherent relics that remained from the original wave-functions of the initial nuclei. Therefore the primary virtual partons must be treated differently than the secondary partons which have been scattered out of the coherent initial state and have become real excitations. With respect to the nuclear center-of-mass frame the primary partons are divided into two sets according to their direction of momentum along the beam axis and their positions. Each set of partons is then assumed to form a beam cluster C_+ (C_-) with positive (negative) longitudinal momentum moving apart from each other on the right (left) side of the center-of-mass, and four momentum given by the sum of parton momenta. The cluster center-of-mass positions are simply taken to be the vector mean of the respective partons' positions.

4.3.2 Hadronization of clusters

(i) **Central clusters:** The invariant mass distribution of central clusters (Fig. 24) may be interpreted as a 'smeared out' version of the spectrum of primordial resonances formed in the early stages of the confinement mechanism [52]. It seems therefore reasonable to treat the fragmentation of these central clusters as a kind of averaged resonance decay which, as

implied by the locality assumption, must be determined entirely by their invariant masses, flavors and total angular momentum. Each cluster in the resonance spectrum may either represent a single hadron resonance that converts directly into a physical hadron with a definite mass, or else fragments through a two-body decay into a pair of final state hadrons. From the particle spectra obtained in e^+e^- -annihilation experiments it appears that quasi-two-body final states are universally dominant, so that the latter possibility seems more favorable if kinematically allowed.

The hadronization scheme for the central clusters employed here follows closely the scheme of Marchesini and Webber [69, 68]. Three regions of cluster mass have to be distinguished, each of which is illustrated in Fig. 25:

- a) Any cluster C that is too light to decay into two hadrons is taken to represent the lightest single hadron (meson) H of its flavor,

$$C(q_i \bar{q}_j) \longrightarrow H(q_i, \bar{q}_j) , \quad (113)$$

with its mass shifted to the appropriate value by adjusting its energy through exchange with a neighbouring cluster.

- b) Each cluster massive enough to decay, but below a fission threshold decays isotropically in its rest-frame into a pair of mesons or baryons as

$$\begin{aligned} C(q_i \bar{q}_j) &\longrightarrow H_1(q_i, \bar{q}_a) + H_2(q_a, \bar{q}_j) \\ C(q_i \bar{q}_j) &\longrightarrow H_1(q_i, q_a q_b) + H_2(\bar{q}_a \bar{q}_b, \bar{q}_j) , \end{aligned} \quad (114)$$

by pulling a quark-antiquark pair $(q_a \bar{q}_a)$ or a diquark-antidiquark pair $(q_a q_b)(\bar{q}_a \bar{q}_b)$ from the vacuum. These created pairs need not be viewed as physical entities with dynamical attributes. Rather, they function as a label for counting and conserving flavor and baryon number.

- c) Occasionally the final state parton configuration yields clusters with masses that are too large for an isotropic decay to be a reasonable ansatz. Even though the mass spectrum falls rapidly at large masses (Fig. 24), a small fraction (about 5-10 percent) of the clusters have masses $M_{cluster} > 4$ GeV. These clusters are fragmented by using the iterative fission model proposed in Ref. [69, 68]. For a cluster C of mass $M_C > M_{fiss}$ the fission $C \rightarrow C_A + C_B$ is assumed to yield two clusters C_A and C_B with $M_A^2 \simeq M_B^2 \simeq \mu_0 M_C$. The fission products $C_{A,B}$ then decay as in a) or b) if $M_{A,B} < M_{fiss}$,

$$\begin{aligned} C(q_i \bar{q}_j) &\longrightarrow C_A(q_i \bar{q}_k) + C_B(q_k \bar{q}_j) \\ &\longrightarrow H_1(q_i, \bar{q}_a) + H_2(q_a, \bar{q}_k) + H_3(q_k, \bar{q}_b) + H_4(q_b, \bar{q}_j) , \end{aligned} \quad (115)$$

or else, if M_A and/or M_B are still $> M_{fiss}$, the fission procedure is repeated until all cluster masses fall below M_{fiss} .

(ii) **Beam clusters:** The two beam clusters that are formed from the spectator partons representing the receding beam fronts carry a significant part of the original longitudinal beam momentum and are characterized by large rapidities. This leads to a specific momentum distribution of final hadrons in the high rapidity region which is assumed to be generated by a soft underlying beam-beam interaction. The fragmentation of the beam clusters proceeds in two steps as follows. First, each of the two clusters is split into an ensemble of collinear nucleons carrying equal fractions of the four-momentum of the cluster. The number of nucleons per beam cluster is determined by the requirement that the sum of nucleon masses equals the cluster mass. The nucleon species are chosen as protons or neutrons with the same probability. If the sum of the nucleon charges does not match the total cluster charge, then one or more neutrons are replaced by protons or antiprotons until the total charge is adjusted. Next, soft interactions between the nucleons of the incoming nuclei and the nucleons resulting from the beam cluster splitting are assumed to have occurred during the collision. This is modelled as low momentum transfers between pairs of nucleons, each pair consisting of a nucleon from one beam cluster and one from the other cluster to which it is closest in phase-space. Each of these pairwise interactions involves a squared momentum transfer t with a distribution corresponding to a diffractive nucleon-nucleon scattering [30].

The motivation to take into account such a soft beam-beam interaction underlying the reaction dynamics of partons is, that this ansatz naturally mimics the structure and number of diffractive events in pp ($p\bar{p}$) collisions (which are the subject of Sec. 5.1). In the simulation of these reactions it occasionally happens (especially in peripheral collisions) that the partons of the incoming p and p ($\bar{p}$) pass through each other without any interaction. Such events are then simply pictured as diffractive events as opposed to non-diffractive events.

4.4 Model parameters from phenomenology and experiment

Before applying the model framework to study the dynamics of nuclear collisions, reasonable values for the two central parameters have to be determined from phenomenological considerations and current experimental knowledge: (i) the scale $p_{\perp cut}$ separating hard and soft parton-parton scatterings, and (ii) the minimum virtuality μ_0 in branching processes acting as an infrared cut-off that limits the soft gluon bremsstrahlung. Furthermore the QCD scale Λ has to be specified. These parameters of the model are fixed by experimental data for the pp ($p\bar{p}$) cross-sections [116] and e^+e^- -annihilation [39], respectively.

4.4.1 Fixing $p_{\perp cut}$ from nucleon-nucleon cross-sections

Little is known about the correct choice of the momentum cut-off $p_{\perp cut}$, since it cannot be studied in QCD-processes triggered by electromagnetic interactions. However, its value can be determined from consideration of the total non-diffractive pp and $p\bar{p}$ cross section as was done in Ref. [116]. Similar analyses have been made in Refs. [70, 117, 118, 119]. The basic idea is to employ the eikonal formalism [120] for the impact parameter representation of the nucleon-nucleon scattering amplitude. At high energies, corresponding to large invariant $s = (p_{N_1} + p_{N_2})^2$ of the colliding nucleons, the total, elastic, and inelastic cross sections can be expressed as integrals over impact parameter b , involving the (complex valued) eikonal function $\chi(b, s)$:

$$\sigma_{tot}(s) = 2 \int d^2b \left[1 - \text{Re} \left(e^{-\chi(b,s)} \right) \right] \quad (116)$$

$$\sigma_{el} = \int d^2b \left| 1 - e^{-\chi(b,s)} \right|^2, \quad (117)$$

$$\sigma_{inel} = \int d^2b \left(1 - \left| e^{-\chi(b,s)} \right|^2 \right). \quad (118)$$

Whereas σ_{tot} and σ_{el} depend on the real as well as the imaginary part of the eikonal, the inelastic cross section is determined by the real part of $\chi(b, s)$ alone.

Now, there is ample experimental evidence that the inelastic nucleon-nucleon interactions can be divided into two classes, diffractive and nondiffractive events:

$$\sigma_{inel} = \sigma_D + \sigma_{ND}. \quad (119)$$

Assuming that σ_{ND} can be calculated from perturbative QCD and the measured parton distributions, while σ_D is determined by soft interactions outside the domain of perturbative QCD, the eikonal function may be separated in two corresponding parts,

$$\chi(b, s) = \chi_0(b, s) + \chi_{QCD}(b, s), \quad (120)$$

and the real part of χ_{QCD} can be identified with the contribution of perturbative parton-parton scattering. In the impulse approximation $\text{Re } \chi(b, s)$ factorizes:

$$\text{Re } \chi(b, s) = \frac{1}{2} \sigma(s) A(b), \quad (121)$$

where $A(b) = \int d^2b' \rho(|\vec{b} - \vec{b}'|) \rho(b')$ is the so-called profile function, related to the impact parameter distribution $\rho(b)$ of partons in the nucleon.

The perturbative QCD contribution to $\text{Re } \chi$ is in lowest order given by

$$\begin{aligned} \sigma_{QCD}(s, p_{\perp cut}^2) = & \sum_{a,b} \sum_{\text{processes } k} \int_0^1 dx_a \int_0^1 dx_b \int_{p_{\perp cut}^2}^{\infty} dp_{\perp}^2 f_a(x_a, Q^2) f_b(x_b, Q^2) \\ & \times \left(\frac{d\sigma_{ab}^{(k)}}{dp_{\perp}^2} \right) \theta(Q^2 - p_{\perp}^2), \end{aligned} \quad (122)$$

where a , b , and k label the type of scattering partons, respectively the scattering process, f_a , f_b are the parton structure functions of the colliding nucleons, and $d\sigma_{ab}^{(k)}/d\hat{t} = (\pi\alpha_s^2/\hat{s}^2)|\overline{M}_{ab}^{(k)}(\hat{s}, \hat{t}, \hat{u})|^2$ are the corresponding parton-parton cross sections [46]. Furthermore, $Q^2 = p_T^2 = \hat{u}\hat{t}/\hat{s}$, where $\hat{s} = x_1x_2s$. Note that σ_{QCD} explicitly depends on the scale $p_{\perp cut}^2$, defining the perturbative domain.

The non-diffractive and the diffractive cross section then take the forms

$$\sigma_{ND}(s) = \int d^2b \{1 - \exp[-\sigma_{QCD}(s)A(b)]\} \quad (123)$$

$$\sigma_D(s) = \int d^2b \{1 - \exp[-\sigma_0(s)A(b)]\} \exp[-\sigma_{QCD}(s)A(b)] . \quad (124)$$

The relation between σ_{QCD} and σ_{ND} , determined from eq. (123), can be used to deduce $\sigma_{QCD}(s)$ from the experimental value of $\sigma_{ND}(s)$. From the so determined value of σ_{QCD} one can then obtain a value for $p_{\perp cut}$ by solving eq. (122) for it. This analysis requires knowledge of the nondiffractive part σ_{ND} of the inelastic nucleon-nucleon cross section as function of the total center-of-mass energy $\sqrt{s}$. Since this quantity is not measured directly, one relies on phenomenological fits [121] to the total, elastic, and diffractive cross sections, σ_{tot} , σ_{el} , σ_D , respectively, to determine

$$\sigma_{ND} = \sigma_{tot} - \sigma_{el} - \sigma_D . \quad (125)$$

For the highest presently accessible energies the elastic and diffractive cross sections saturate the so-called Pomplin bound [122] $\sigma_{el} + \sigma_D \leq \sigma_{tot}/2$. Here the equality sign is approached in the limit of very many open diffractive channels [123] and implies that the eikonal phases χ_0 and χ_{QCD} in (120) become of equal size. In view of eq. (121) one can write this asymptotic limit as $\sigma_0(s) \xrightarrow{s \rightarrow \infty} \sigma_{QCD}(s)$. The analysis shows that this asymptotic limit is already well satisfied at Tevatron energies ($\sqrt{s} = 1.8$ TeV).

The eikonal cross sections $\sigma_{QCD}(s)$ and $\sigma_0(s)$ can now be deduced from σ_{ND} (123) and σ_D (124). The resulting curves are exhibited in Fig. 26. The value $\sigma_{QCD} + \sigma_0 \approx 50$ mb for center-of-mass energies below 100 GeV is in good agreement with other, independent determinations [117, 118]. However, the essential difference is to those is that here most of this value is attributed to parton-parton scatterings. The resulting values for $p_{\perp cut}(s)$ are shown in Fig. 27 for two different sets of nucleon structure functions. Least-square fits through the data points show that the functional relation

$$p_{\perp cut}(s) = a \left(\frac{s}{s_0} \right)^b \quad (126)$$

provide a good description, where $s_0 = 1$ GeV² and $a = 0.35$, $b = 0.14$ for the GRV structure functions [127], and $a = 0.48$, $b = 0.11$ for the (outdated) EHLQ structure functions [126].

For instance, on account of (126) one has for the GRV structure functions:

$$p_{\perp cut}(s) = \begin{cases} 1.55 \text{ GeV} , & \sqrt{s} = 200 \text{ GeV} \\ 2.85 \text{ GeV} , & \sqrt{s} = 1800 \text{ GeV} \\ 4.00 \text{ GeV} , & \sqrt{s} = 6000 \text{ GeV} \\ 6.80 \text{ GeV} , & \sqrt{s} = 40000 \text{ GeV} \end{cases} . \quad (127)$$

For the case of nuclear collisions $A + B$ the values of $p_{\perp cut}(s)$ associated with the nucleon-nucleon cross-section simply require to replace in eq. (126) the total CM energy by the energy per nucleon pair,

$$\sqrt{s} \longrightarrow \sqrt{s_{NN}} \equiv \frac{2\sqrt{s}}{A+B} , \quad (128)$$

where on the right hand side $\sqrt{s}$ is now the total invariant mass of the nuclear system and $\sqrt{s_{NN}}$ refers to a pair of nucleons.

It is important to realize that this approach to determine $p_{\perp cut}$ is based on the lowest-order parton cross-section in Born approximation, eq. (122), which implies a "single collision" approximation, neglecting multiple scatterings, radiative processes, and higher twist effects. Therefore the parametrization (126) represents strictly speaking only an estimate for the initial primary parton scatterings. This is presumably sufficient for nucleon-nucleon collisions, but in nuclear collisions, particularly heavy ion reactions, where multiple interactions, parton cascading, screening and dense medium effects play an important role, this simple parametrization is very likely to be modified. Note however, that $p_{\perp cut} \geq 1 \text{ GeV}$ in the entire range of energies $\sqrt{s} \geq 25 \text{ GeV}$ where the model might be considered to be applicable and the probability that a parton interaction with momentum transfer of 1 GeV (corresponding to $\alpha_s \approx 0.4$) leads to inelastic nucleon scattering is greater than 99.9 percent [118]. Nonetheless, $p_{\perp cut}$ must be regarded as a phenomenological parameter which describes in a way the screening of the QCD interaction at large distances.

4.4.2 Fixing μ_0 from fragmentation in e^+e^- -annihilation

Recall that in the PCM the parameter μ_0 defines the boundary for the perturbative evolution of the parton distributions up to the point when the majority of cascading partons have de-excited to invariant masses $m \simeq \mu_0$. On the other hand, as illustrated in Fig. 17, the evolution of the final parton configuration beyond the perturbative cut-off scale μ_0 is intimately connected with the pre-confinement mechanism [113] that ultimately yields to the hadronization associated with the QCD scale Λ . The commonly accepted parton-hadron duality hypothesis [128] views the hadronization process as a local phenomenon which implies that the conversion of partons into hadrons is universal (independent of the specific reaction). Therefore it is at hand to determine the interplay of μ_0 and Λ from precision data of e^+e^- -annihilation experiments, in which the fragmentation of parton cascades can

be studied most cleanly and has been analysed in great detail [129]. The dimensionless ratio μ_0/Λ defines the maximum value of the effective coupling α_s/π , which can occur in the perturbative cascade evolution of parton branchings. A third parameter, entering the hadronization scheme of Sec. 4.3, is the fission threshold M_{fiss} for the fission of heavy clusters before they decay isotropically. As it turns out, this parameter is, in contrast to Λ and μ_0 , of little influence on global observables such as multiplicities and momentum distributions.

The most characteristic observable measures to study the significance of a variation of the parameters Λ and μ_0 are the multiplicities and the particle composition of the final state, and the momentum distributions of the produced hadrons. In Ref. [39] a detailed examination of the dependence on the parameters μ_0 , Λ and M_{fiss} has been carried out by comparing the predictions of the model with the very well measured e^+e^- data of PETRA/PEP and LEP [129]. In this analysis, e^+e^- annihilation events were simulated as the production of $q\bar{q}$ jets through a virtual γ with invariant mass Q , followed by the perturbative evolution of the jets as parton cascades to Sec. 4.2, and the hadronization of the final partons using the cluster hadronization of Sec. 4.3. The parameter dependence was analyzed in detail in Ref. [39].

The following Figs. 28-31 show a few examples that summarize the effects of the choice of Λ and μ_0 for the four combinations $(\Lambda, \mu_0) = (0.2, 1); (0.4, 1); (0.4, 0.5); (0.6, 1.5)$ GeV and compare the resulting predictions for various observables in e^+e^- annihilation with experimental data. In Fig. 28 the $p_\perp$ -distribution of the final charged hadrons is plotted for the two center-of-mass energies $Q = 15$ GeV and $Q = 29$ GeV, whereas Fig. 29 shows $\langle p_\perp^2 \rangle$ as a function of Q . The significance of the tail in the $p_\perp$ -distribution and the magnitude of the resulting $\langle p_\perp^2 \rangle$ are governed by two effects [112]: First, the higher the invariant mass of the individual cluster, the larger will be the transverse momentum generated in their decays into hadrons. Second, the larger Λ is, the higher is the probability for the emission of large transverse momentum gluons during the perturbative cascade evolution, which leads to a larger power-law tail in the partons' $p_\perp$ -distribution and hence in the $p_\perp$ -spectrum of the resulting hadrons. Thus, the largest $\langle p_\perp^2 \rangle$ is obtained with a large Λ (hence large α_s) and large μ_0 (hence large cluster masses).

The multiplicities of produced hadrons and the sensitivity of the model predictions to a variation of Λ and μ_0 are exhibited in Fig. 30 where the overall multiplicity of final charged particles is shown as a function of Q , and in Fig. 31 which depicts the behaviour of the average $\pi^+ + \pi^-$, $K^+ + K^-$ and $p + \bar{p}$ multiplicities with increasing center-of-mass energy. As expected from the previous discussion, a parameter combination with Λ small and μ_0 small results in a larger overall multiplicity and a stronger increase with Q than one with both Λ and μ_0 being large. Also, pion production dominates in this case, because

kaon and baryon production is suppressed by the larger K and p masses and a larger number of π than K is produced in the decays of low-lying meson resonances. Note that the measured approximate constancy of the total charged hadron multiplicity over the range $1.5 \leq Q \leq 4$ GeV in Fig. 30 supports the assumption that clusters with masses in this range directly convert into pairs of light mesons or baryons without cascading through clusters of intermediate mass, i.e. without cluster fission. Accordingly, the fission threshold of $M_{fiss} = 4$ GeV, as employed in the calculations, seems to be a reasonable choice.

The conclusion drawn in Ref. [39] was that the following choice of the parameters μ_0 , Λ , and M_{fiss} , yields the best overall agreement with the experimental data:

$$\mu_0 = 1 \text{ GeV} , \quad \Lambda = 0.4 \text{ GeV} , \quad M_{fiss} = 4 \text{ GeV} . \quad (129)$$

The measured momentum spectra as well as the particle multiplicities are most accurately reproduced with this particular parameter combination. In the PCM these values are considered to be universally fixed and independent of the type of nuclear reaction, i.e. the same for $p\bar{p}$, pA , or AA collisions, which are the subject of the following Chapter.

4.5 Central assumptions and approximations of the model

At this point it is important to point out clearly the limitations and shortcomings of the PCM approach. There are a number of assumptions and open questions [38, 82] that remain to be improved or resolved, if possible, in the future:

1) The PCM is a semiclassical approach to model some of the numerous complex mechanisms that predominantly govern the dynamics of high energy nuclear collisions. It is based on a classical transport equation for the single-particle distributions of partons, in which the collision term is constructed on the basis of improved QCD perturbation theory. The central assumption is here, that (at least during the early stage of the nuclear reactions) the dynamics is essentially driven by binary parton-parton collisions in which the partons are treated as quasiclassical particles that interact through independent scatterings with associated radiative emissions and absorptive processes. Accordingly, long range color field and color correlations among the partons are assumed to be negligible for the initial dissipative dynamics.

2) The model combines the rather well understood dynamics of parton cascades in the framework of perturbative QCD with a number of relatively bold phenomenological elements regarding the extrapolation from hadronic collisions to nuclear reactions: (i) the ansatz for the initial parton distribution of the colliding nuclei; (ii) the extrapolation of (semi)hard parton collisions to a phenomenological soft scattering region; (iii) the use of free-space matrix elements together with medium independent infrared cut-off parameters $p_{\perp cut}$ and μ_0 , which stand in for a dynamical regularization due to Debye screening in dense

matter environment; (vi) the almost classical prescription to account for the formation time of radiative emissions and the Landau-Pomeranchuk-Migdal effect by treating the virtual partons as unstable excitations with a finite life-time.

3) It is well known that a Lorentz invariant many-body theory must necessarily be of many-time nature, even at the classical level: each particle carries its own proper time. On the other hand, the structure of any conventional cascade calculation is, that the (three-dimensional) variation of spatial distance for every possible pair of particles is followed to arrange the ordering of binary collisions taking place in cascades (i.e. which pair collides first, which next, etc.). However, it turns out [82], that the prescription to determine whether a collision occurs (c.f. Appendix) by comparing the distance of closest approach $(r_{ab})_{min}$ between two particles a and b with the interaction radius $\sqrt{\hat{\sigma}_{ab}}/\pi$, is almost a Lorentz invariant quantity in any Lorentz frame moving along the beam direction. Similar arguments hold for the radiative emission and absorption processes of partons.

4) Concerning the question of gauge invariance, the PCM approach meets a fundamental problem: the parton-parton scattering amplitudes with off-mass shell partons in the initial or final state of the vertex are not gauge invariant. This is of relevance when time-like virtual partons propagate and rescatter before they have returned on mass shell. However, the majority of time-like partons have relatively small virtualities, close to the mass shell, because most of the parton-parton scatterings, which provide them with this virtual excitation, involve a moderate momentum transfer. Only in the rare truly hard scatterings partons get scattered far off-mass shell. Therefore the violation of gauge invariance should be small in the mean. Nevertheless, a satisfactory implementation of gauge invariance in the parton cascade approach remains a conceptual problem.

5. MULTIPARTICLE PRODUCTION IN NUCLEAR REACTIONS AT COLLIDER ENERGIES

This Chapter is devoted to illustrate the application of the PCM approach to proton-antiproton ($p\bar{p}$), nucleus-nucleus (AA) and proton-nucleus (pA) collisions at beam energies from RHIC ($\sqrt{s} = 100$ GeV/nucleon) to LHC ($\sqrt{s} = 3000$ GeV/nucleon). The time evolution in phase-space of the collisions is analysed from the instant of nuclear contact, via the parton cascade development, up to the formation of final hadronic states. The model is tested first for $p\bar{p}$ collisions at $\sqrt{s} = 200$ -1800 GeV where a decent agreement with the experimentally measured momentum and multiplicity distributions is found. Encouraged by that, the approach is subsequently applied to AA and pA collisions with focus on the impact of a number of nuclear and medium effects on the inclusive hadron production. The hypothesis of the hadronization process being a local phenomenon implies that the conversion of partons into hadrons must be universal and independent of the specific reaction;

thus the rationale is that the PCM approach should work equally well for hadronic and nuclear reactions. However, when proceeding from hadron-hadron to heavy ion collisions new phenomena arise on the parton level due to nuclear effects and dense medium environment, e.g. nuclear shadowing, multiple interactions, collective behavior. In fact, the model calculations of charged particle spectra and multiplicities in heavy ion collisions exhibit a non-trivial interplay between such effects with a general tendency to a suppression of particle production; still the results for central rapidity densities are of a factor of ~ 2.5 per nucleon larger than in pp or $p\bar{p}$ collisions at corresponding energies.

5.1 $p\bar{p}$ collisions

With the model parameters $p_{\perp cut}$, μ_0 and Λ assumed to be universally fixed, it is useful to test the applicability of the PCM in $p\bar{p}$ collisions which have been studied experimentally as well as theoretically in great detail. Many high statistics data have been collected by the collaborations UA1, UA2 and UA5 at CERN ($\sqrt{s} = 200, 546, 900$ GeV) and the CDF collaboration at the TEVATRON ($\sqrt{s} = 1800$ GeV) to which the model results for particle multiplicities, momentum spectra, etc., can be compared. A detailed analysis has been carried out in Ref. [39]. Similar studies have been done by others [136, 70, 68, 71].

The simulation of $p\bar{p}$ collisions in the PCM consists of generating the initial parton distributions (Sec. 4.1) in the incoming p and $\bar{p}$ with a randomly chosen impact parameter, evolving the parton distributions in phase-space by following the cascade development (Sec. 4.2), and hadronizing the final partons with the hadronization scheme (Sec. 4.3). Tables 1 and 2 list some average properties of $p\bar{p}$ collision events generated in this fashion for different collider energies. Fig. 32 shows the inclusive transverse momentum spectra at various beam energies, a) for the final state parton configuration just before hadronization and b) for the resulting hadrons. Correspondingly, Fig. 33 displays the (pseudo)rapidity distributions of a) final partons and b) produced hadrons, respectively. The most important aspects of the Figs. 32 and 33 can be summarized as:

- (i) The shapes of the final hadron spectra are essentially predetermined by the corresponding distributions at the parton level. The evident "softening" of both $p_{\perp}$ - and $y(\eta)$ -spectra in the parton-hadron conversion is generated by the hadronization and is determined by the average energy release in cluster decays (and in the subsequent decay chain of produced hadron resonances).
- (ii) In the $p_{\perp}$ -spectra, one observes for large $p_{\perp} \gtrsim 2$ GeV a characteristic power law tail, reflecting contribution of comparably hard parton collisions and branchings, whereas at low $p_{\perp} \lesssim 2$ GeV, soft physics is dominant, mostly due to soft and collinear gluons emitted in bremsstrahlung processes.

- (iii) The rapidity spectra exhibit a depletion of partons around zero rapidity which is a consequence of the destructive interference of soft gluons in the cascades (c.f. Sec.4.2). This effect persists also in the hadron spectra (even though the kinematic difference between true rapidity (y) and pseudorapidity (η) also contributes here).
- (iv) The increase of the central rapidity density around $\eta = 0$ is totally due to the growing number of parton-parton collisions and the multiplicity increase of partons produced in the associated cascades, whereas the increase of the overall width of the rapidity distributions proportional to $\sqrt{s}$ is essentially the consequence of a larger energy and longitudinal momentum brought into the reactions.

The naive parton model assumes that single parton-parton scatterings clearly dominate in hadron-hadron collisions. In the PCM calculations however multiple parton scatterings (even rescatterings) occur with significant probability (see Table 1). This effect is due to the increase in the number of parton cascades as the beam energy grows. It is also clearly reflected when analyzing the multiplicity distributions of final charged hadrons produced in $p\bar{p}$ collisions. Fig. 34 shows the total charged particle multiplicity distributions from the calculations and the corresponding data [137, 138] at $\sqrt{s} = 200, 546$ and 900 GeV. The model results (full histograms) decently reproduce the systematic widening of the measured distributions with increasing collider energy. It is illuminating to see how the contributions from events with $k = 0, 1, 2$, and ≥ 3 parton-parton collisions (indicated by the patterned curves) add up to give the total distributions: the low multiplicity region is dominated by events with no parton scatterings above $p_{\perp cut}$, whereas the high multiplicity tail is due to events with several interactions. Furthermore, the average charged multiplicity $\langle n_{ch} \rangle$ also increases with the mean number of interactions, as is obvious from Table 2 and from the shift of the peak in the plots of Fig. 34. However, each additional interaction gives a smaller contribution than the preceeding one, so that there seems to be a saturation. These features have independently been predicted within perturbative QCD before [70, 71]. The model independence of the predictions and the agreement with the measured distributions therefore provide strong support for the evidence of multiple parton interactions and manifests the important role of semihard perturbative QCD for the event structure.

Further support for the expectation that a perturbative QCD description in terms of semihard parton interactions and minijet production is not only satisfactory, but also necessary to explain many features of multiparticle production of hadron collider physics, comes from detailed Monte Carlo studies within different models, e.g. HIJING [71], PYTHIA [70], and HERWIG [68] (for a general overview and compilation, see Ref. [72]). For example, in Fig. 35 the increase of the central rapidity density with beam energy $dN^{ch}/dy|_{y=0}$ in

$p\bar{p}$ collisions is shown as predicted by HIJING calculations. The full curve corresponds to truly inelastic collisions including minijet production. The dashed line on the other hand is the calculated central density without minijets. Comparing those two curves, one sees that the jet production becomes substantial at collider energies $\sqrt{s} \gtrsim 50$ GeV and clearly dominates at higher energies, implying that a large fraction of produced particles in the central region comes from minijets. The agreement with experimental data in Fig. 35 is impressive. Another suggestive manifestation of copious minijet production arises when studying the $p_{\perp}$ -dependence of the two-particle correlation function in minimum biased events of hadron collisions at collider energies. Because particles produced by jet fragmentation tend to cluster together in phase-space, the two particle correlation function must be enhanced due to minijet production. Especially for the correlation in azimuthal angle ϕ , the contribution from back-to-back minijets should be strongly peaked in both forward ($\Delta\phi = 0$) and backward ($\Delta\phi = \pi$) directions. Contrary to that, particles from soft (non-perturbative) production are known to be approximately isotropically distributed in the transverse plane and would only show some nominal correlation due to momentum conservation. Fig. 36 nicely demonstrates this, by showing calculations of HIJING [147] for the two-particle correlation function in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV for selected particles with a different $p_{\perp}$ cut. A strong correlation is obvious, peaked around $\Delta\phi = 0$ and $\Delta\phi = \pi$, with a valley at $\Delta\phi \approx \pi/3$. A larger $p_{\perp}$ cut extracts this signal naturally clearer from the flat soft particle background.

The essential conclusion from the study of $p\bar{p}$ collisions is that multiple jet production and parton cascading is increasingly important in many aspects of particle production at collider energies. The results of various independent analyses and their good agreement with experimental data clearly show that the parton picture provides a consistent framework to explain the increase (with collider energy) of the central rapidity density and the total cross-section, to describe correctly the shapes of rapidity, transverse momentum, and multiplicity spectra, as well as to predict the characteristics of two-particle correlations.

5.2 AA collisions

Nucleus-nucleus collisions at collider energies, in particular those involving heavy ions, are expected to exhibit novel effects in the dynamical evolution [82, 148] and multiparticle production [39, 149]. Of particular interest are nuclear and dense medium effects associated with the microscopic dynamics of the many-parton systems during these reactions, since they offer to study QCD in hot, ultra-dense matter and may signal some evidence for the predicted QCD phase transition to a quark gluon plasma. In this context the PCM approach provides a special opportunity to study the relevance of these new physical features for heavy ion collisions, features that are not visible or significant in dilute and small-size

systems as hadronic collisions. Aside from the mentioned modifications of the perturbative parton evolution, it is clear that the conversion of partons into hadrons is a much more complicated process than the hadronization of final state partons in hadron-hadron collisions. Little is known, but the transition is expected to involve a rather long time scale ($\simeq 10 fm$) and might proceed first as an expansion and cooling of the plasma and then through a mixed parton-hadron phase to a purely hadronic phase. Several models have been proposed to describe the hadronization in ultra-relativistic heavy ion reactions (see e.g. Refs. [150, 151] and references therein).

Different Monte Carlo models based on the perturbative QCD parton picture [41, 42, 25, 39] have been applied to investigate multiparticle production in high energy AA collisions. However, up to now only in HIJING [41] and the PCM the dynamics is manifestly driven by perturbative QCD minijet production and copious materialization of partons. Although there is qualitative agreement in the predictions for the particle spectra and the multiplicity increase with collider energy, a substantial uncertainty exists in global observables, e.g. the absolute number of produced particles. This is made obvious by Fig. 37, which compares results of a) the HIJING model and b) the PCM for the $p_{\perp}$ - and η -spectra of charged hadrons produced in heavy ion collisions at RHIC and LHC. The similarity of the spectra from the two independent calculations is obvious and is a direct consequence of the perturbative QCD description common to both models, but the predictions for the absolute magnitude of the particle production or the central rapidity density differ by factors of 2 or more. This discrepancy is mainly due to two reasons: first, the different regularization procedures, i.e. the interpretation of the cut-off parameters $p_{\perp cut}$ and μ_0 , and second, the different (and in HIJING partly absent) treatment of multiple parton interactions and collective effects.

The sobering conclusion is: a useful quantitative predictions for observables in heavy ion collisions will not be obtainable, before the role of infrared regularization by screening mechanisms and the details of dense medium effects in QCD are better understood. The impact of nuclear and dense matter modifications was investigated within the PCM framework, both on the level of the microscopic parton kinetics [82], and in the context of hadron production in central $Au + Au$ collisions at $\sqrt{s} = 200$ A GeV (RHIC) and $\sqrt{s} = 6300$ A GeV (LHC) [39, 148]. The time evolution of the partons' phase-space distributions was followed from the moment of nuclear contact up to the production of hadronic final states, as described in Chap. 4. However, instead of attempting to model a detailed space-time description of the complex parton-hadron transition, the partons were simply hadronized locally at the end of the perturbative phase provided they had separated sufficiently to be considered as distinct, non-interfering quanta. The Tables 3 and 4 summarize some average properties of simulated $Au + Au$ events on the basis of the full model description

(i.e. including nuclear and medium effects discussed in Sec. 4.1 and 4.2). Table 3 lists the mean numbers of the various parton interactions (c.f. corresponding Table 1 for $p\bar{p}$) and Table 4 gives the average numbers of initial and final partons as well as the resulting charged hadron multiplicities (c.f. corresponding Table 2 for $p\bar{p}$).

The question, how significant various nuclear and dense medium effects are reflected in the final hadron spectra and whether it is possible to disentangle the significance of the different phenomena, was addressed in Refs. [39, 82] by considering five different evolution scenarios for the parton distributions. The *evolution schemes* are as follows:

- a) *'Naive evolution'*: Parton shadowing in the initial parton distributions is neglected. Parton scatterings occur instantaneously without time delay. Associated cascades of successive parton branchings happen immediately at the same space-time point as the scattering, i.e. with zero formation time for radiative emissions. Absorptive processes and soft gluon interference are switched off.
- b) *Nuclear shadowing*: As a), but now the initial parton distributions are modified to account for nuclear shadowing effects as explained in Sec. 4.1.
- c) *Parton fusion and absorption*: As b), but in addition the time scale for each individual parton-parton collision is taken into account. Parton fusion and absorptive processes compete with the emission of partons in the cascades.
- d) *Landau-Pomeranchuk-Migdal effect*: As c), but furthermore, successive parton branchings in the cascade evolution are delayed according to the characteristic formation time for gluon emissions.
- e) *Soft gluon interference*: As d), but moreover, the destructive interference of soft gluons decreases the available phase-space for low energetic gluon bremsstrahlung.

The influence of the above effects on the evolution of AA collisions is rather vivid on the parton level [82]. Fig. 38 displays the average cumulative number of secondary partons per nucleon a function of time measured in the total CM frame for $S + S$ ($A=32$) and $Au + Au$ ($A=197$) collisions. The dotted and full lines correspond to the evolution scenarios (a) and (e), respectively (except for nuclear shadowing). Comparing the curves (a) and (e) one sees that in the latter case the multiplicity of secondary partons is significantly lower and the evolution of the system is dilated. Also, these effects are clearly more prominent for $Au + Au$ than for $S + S$. Furthermore, an interesting feature of scenario (e) is the sudden temporary suppression of parton production following the explosive increase at $t \simeq 0.6 fm$. The reason for this behaviour is the abrupt production of a large number of partons within a very short time which gives rise to a dense phase-space population. The particle emission

from excited partons is effectively suppressed because many partons encounter multiple scatterings and cannot emit radiation immediately. In addition the absorption of partons by others becomes important as the system responds to the high density.

The new physics associated with the effects in a) - e) is also drastically evident in the charged particle distributions in pseudorapidity (η), calculated for central $Au + Au$ collisions with $\sqrt{s} = 200$ A GeV and $\sqrt{s} = 6300$ A GeV. The spectra are shown in Fig. 39, where the various curves a) - e) exhibit the characteristics of the different evolution scenarios. The particle density in the central rapidity region is reduced successively from the 'naive' calculation (case a) to the default result (case e) by more than a factor of 3. For both collider energies the behaviour is roughly the same, although the impact of nuclear shadowing (curve b) and the Landau-Pomeranchuk-Migdal effect (curve d) seem to become more prominent at $\sqrt{s} = 6300$ A GeV. Obviously the drastic reduction of the charged particle yield is most significant at $\eta \approx 0$, because the inclusion of delayed gluon emission associated with the Landau-Pomeranchuk effect (case d) and soft gluon interference (case e) leads to a specific depletion of softer partons which is reflected by the dip appearing in the range $|\eta| \leq 1$. Although clearly mirrored in the hadron distributions, both these effects originate from the parton level [39]. From RHIC to LHC energies, i.e. increasing in $\sqrt{s}$ by a factor of 30, the central rapidity density roughly doubles (from $\simeq 1200$ to $\simeq 2500$ in case e). At the same time the total charged multiplicity grows from $\simeq 6600$ to $\simeq 21500$ from RHIC to LHC, i.e. by more than a factor of 3, because not only the height but also the width of the spectra increases.

The corresponding transverse momentum ($p_{\perp}$) distributions of charged particles $Au+Au$ collisions at RHIC and LHC energies have also been computed and are presented in Ref. [39]. It was found that the $p_{\perp}$ -spectra corresponding to the evolution schemes a) - e) are hardly distinguishable in their shape, in contrast to the rapidity spectra in Fig. 39. The slope is roughly the same for all five cases, at least for moderate $p_{\perp} \leq 3 - 4$ GeV. However, in the region $p_{\perp} \geq 6$ GeV the spectra appear to become slightly harder when comparing the cases from a) to e), in that this high $p_{\perp}$ -domain becomes more populated. At both RHIC and LHC energies, the low and moderate $p_{\perp}$ -region ≤ 4 GeV is characterized by an approximate exponential distribution $\sim \exp(-p_{\perp}/p_0)$ with $p_0 \simeq 0.35$ GeV ($\sqrt{s} = 200$ A GeV) and $p_0 \simeq 0.39$ GeV ($\sqrt{s} = 6300$ A GeV). From RHIC to LHC the average $p_{\perp}$ however increases strongly, namely from $\langle p_{\perp} \rangle \simeq 0.39$ GeV ($\sqrt{s} = 200$ A GeV) to $\langle p_{\perp} \rangle \simeq 0.54$ GeV ($\sqrt{s} = 6300$ A GeV).

The comparison of the η - and $p_{\perp}$ -spectra in $Au + Au$ collisions with the corresponding momentum distributions in $p\bar{p}$ collisions provides another instructive insight. In Fig. 40

the ratios

$$R_{Au/p}(\eta) = \frac{dn_{ch}/d\eta (Au + Au)}{dn_{ch}/d\eta (p + p)}, \quad R_{Au/p}(p_{\perp}) = \frac{dn_{ch}/dp_{\perp}^2 (Au + Au)}{dn_{ch}/dp_{\perp}^2 (p + p)} \quad (130)$$

of the η -distribution and the inclusive $p_{\perp}$ -spectrum for the evolution scheme e). Even for the case e) with the lowest multiplicities, one has significantly more more than twice the particle density per nucleon around $\eta = 0$. On the other hand, there is obviously a strong suppression of particles at large $p_{\perp}$ which is mainly due to the increased energy loss especially of high $p_{\perp}$ partons in the range $p_{\perp} = 4 - 8$ GeV. This phenomenon is due to so-called of jet quenching [152] resulting from the energy loss of high- $p_{\perp}$ partons as they traverse dense nuclear matter, as has independently been studied before [149] with qualitatively the same results. The enhanced probability of multiple scatterings accompanied by stimulated gluon radiation leads to dissipation of the energy-momentum of jets in medium as compared to vacuum. As a consequence, produced high- $p_{\perp}$ jets in heavy ion collisions will increasingly be "washed out". This effect is vividly suggestive in Fig. 41, where the transverse energy distribution $dE_{\perp}/dyd\phi$ in rapidity y and azimuthal angle ϕ is plotted for $p\bar{p}$ and $Au + Au$, respectively. Whereas in $p\bar{p}$ the jets, once produced, persist to stick out clearly, in $Au + Au$ collisions the original jets "evaporate" with time and disappear in a vast background of noisy fluctuations.

5.3 pA collisions

A systematic study of pA collisions in comparison with pp or $p\bar{p}$ collisions at the same energies is essential to gain insight into the structure of the medium effects discussed in Sec. 5.2, since these are absent in pp ($p\bar{p}$) collisions, but become increasingly prominent in pA collisions, and are of major importance in AA reactions. Hence, by increasing the target mass A and the collider beam energy $\sqrt{s}$ in pA collisions, one may probe nuclear matter as a function of its density and study the gradually growing impact of medium and nuclear effects by comparing to pp collisions. Once the physics of 'QCD in medium' is better understood, then collective phenomena and medium effects in heavy ion collisions can be disentangled from the basic hadronic effects. In this context, one of the most controversial issues is the huge entropy production due to very intense gluon multiplication in nuclear matter [6, 153] that is triggered by the frequent semihard parton collisions and parton cascading. The crucial questions are: first, is there an upper limit on the amount of gluon multiplication imposed by unitarity, and second, what is the effect of Debye screening in dense nuclear matter for non-equilibrium systems. In Ref. [154] both these issues were discussed in the framework of the PCM.

Recall from Chap. 2 that the applicability of the parton picture requires a description in a Lorentz frame where both the projectile and target are moving close to the speed of

light, so that both their substructure can be resolved into individual partons. Due to the asymmetry in beam and target masses in pA collisions, the overall CM frame (17) and (77) employed before for $p\bar{p}$ and AA collisions, is not appropriate, because here the nucleus is essentially at rest. Instead one may choose the *nucleon-nucleon center-of-mass frame* (CM_{NN}) in which each nucleon has the same value of longitudinal momentum P [154],

$$P_z^{(p)} = +P, \quad P_z^{(A)} = -AP, \quad \vec{P}_\perp^{(p)} = \vec{P}_\perp^{(A)} = \vec{0}, \quad (131)$$

so that $\sqrt{s} = \sqrt{4AP^2 + M_N^2(1 + A^2)}$ and $\sqrt{s_{NN}} = 2P$, where A is the nuclear mass number, M_N the nucleon mass. The condition $P/M_N \gg 1$ is certainly satisfied at the colliding beam accelerators RHIC and LHC. For example, at RHIC, the maximum P is 250 GeV for $p + p$, 125 GeV for $p + {}^{16}\text{O}$ and 100 GeV for $p + {}^{197}\text{Au}$ [155]. At LHC one has generally a factor of 30 larger energy available [156]. The choice of the CM_{NN} frame may appear uncommon for proton-nucleus collisions. However, at the collider facilities RHIC and LHC both proton and nucleus beams are accelerated in opposite direction with the same Lorentz factor. Therefore the center-of-mass energy per nucleon pair $\sqrt{s_{NN}}$, or equivalently P , is the actual measure of the beam energy. Note that the beam momenta $P = 100$ (3000) GeV at RHIC (LHC) transform to about $P_{Lab}^{(p)} = 21$ (19000) TeV incident proton momentum in the Lab frame. For comparison, at the currently highest energy pA experiments performed at the FNAL accelerator, the incident proton has at most $P_{Lab}^{(p)} = 800$ GeV momentum in the laboratory, corresponding to $P \simeq 20$ GeV for $p + {}^{208}\text{Pb}$.

(i) **Parton saturation:** First let me discuss the issue of parton multiplication constrained by unitarity. In the parton picture a pA collision may be viewed as a diffusion of the partons of the projectile proton, traversing the nucleus and interacting frequently with partons belonging to different target nucleons [18, 157]. As has become clear from the investigation of AA collisions in Sec. 5.2, multiple scattering and bremsstrahlung, accompanying the partons' paths through the nuclear medium, lead to a rapid entropy production that increases the local density and consequently enhance the probability for further interactions. The fundamental question is, whether the intense parton multiplication that results in a dense phase-space population at small energy fractions $x \ll 1$, leads at some point to a probability larger than unity for a parton to interact with this dense parton matter, in which case the unitarity principle would be violated. This occurs when the partons are so densely packed that they spatially overlap in transverse direction [158, 159] and the nucleus appears to be completely black. However, detailed balance between scattering, branching and fusion processes must yield a local saturation of the parton density and stop a further increase, thereby ensuring unitarity. The requirement of unitarity conservation can be satisfied by demanding that the sum of probabilities of all elastic plus inelastic reaction channels must not exceed unity. When phrased in terms of the *Gribov-Levin-Ryskin*

unitarity condition [17, 18]: the total interaction probability W_a of a given parton of species a to collide with another parton must always satisfy (locally!) $W_a \leq 1$.

In Ref. [154] this condition was implemented in the kinetic approach of the PCM by computing the probability W_a as a function of time in individual phase-space cells and from this determining the relative probability of parton production to parton fusion processes in a self-consistent, dynamic manner. The interaction probability W_a can be expressed as a sum of contributions from scatterings of a parton a with all kinds of partons b , that become scattering candidates within Δt . It is integrated over the volume which the parton a traverses during the time span Δt , as well as integrated over the momentum and weighted by the phase-space distribution of those partons b present in this volume (see Fig. 42). Thus, the *unitarity condition* is:

$$W_a(p_a, \vec{r}, \Delta t) := \Delta t \sum_b \int_z^{z+\Delta z} dz \int_0^{R_A} d^2 r_\perp w_{ab}(p_a, \vec{r}, \Delta t) \stackrel{!}{\leq} 1, \quad (132)$$

with

$$w_{ab}(p_a, \vec{r}, \Delta t) = \frac{1}{2E_a} \frac{1}{\gamma_a \gamma_b} \frac{1}{S_{ab}} \int \frac{d^3 p_b}{(2\pi)^3 2E_b} \gamma_b F_b(p_b, \vec{r}, t) \quad (133)$$

$$\times \int \frac{d^3 p_c}{(2\pi)^3 E_c} \int \frac{d^3 p_d}{(2\pi)^3 E_d} \left[\sum |\mathcal{M}_{ab \rightarrow cd}|_{eff}^2(p_a, p_b, Q^2) \right] \delta^4(p_a + p_b - p_c - p_d).$$

The notation is here the same as in Sec. 4.2, eqs. (92) and (94). The physical meaning of this constraint is very simple. Since W_a is proportional to the density of scatterers $n_b \propto \int d^3 p_b F_b(p_b, \vec{r}, t)$, the relation $W_a \ll 1$ will be satisfied as long as n_b is sufficiently dilute. On the other hand, when W_a approaches 1, absorption or fusion processes become essential. Thus, W_a gives the time-dependent probability for the fusion of a given parton a with some other parton b , depending on the local density n_b and the scale of interaction, $Q^2 = p_\perp^2$.

In Fig. 43 the time evolution of the probability W_g for gluons at small rapidities $|y| \leq 1.5$, obtained from the PCM simulations of pA collisions with $P = 100$ GeV and $P = 3000$ GeV, where P is defined in eq. (131). The quarks and antiquarks play a minor role here, so that W_g can be identified to good approximation with the total $W = \sum_a W_a$. Common to both beam energies is that for light nuclei $A \lesssim 30$ the fusion probability is only $\simeq 0.2 - 0.3$, but it already reaches more than 70% for nuclei around $A = 100$, and 90 %, respectively 100 % for ^{238}U at $P = 100$ (3000) GeV. It is also interesting that the time it takes to increase W_g from 0.1 to 0.9 of its final value is typically a factor of 2 smaller at $P = 3000$ GeV than at $P = 100$ GeV. Furthermore, at fixed P this time scale decreases significantly when proceeding from ^{32}S to ^{238}U nuclei. Note that the saturating behaviour that is evident even for e. g. ^{32}S although W_g is far from unity, results from the fact that

the duration of the reaction is the smaller the lighter the nucleus is, so that the partons naturally cease to interact (except for some small amount of residual interactions) when the system flows apart before it finally hadronizes.

(ii) **Screening effects:** Now let me turn to the related question concerning Debye screening [160] of parton interactions in dense nuclear matter. In the semiclassical approach of the PCM screening can be described microscopically as controlled by the dynamical evolution of the many-parton system, i.e. the space-time dependent phase-space occupation. In Ref. [154] it was investigated how this classical screening mechanism can accomplish a selfconsistent regularization of the infrared singularity of the effective parton cross-section, because long range interactions corresponding to large impact parameters of partons are naturally cut off provided the surrounding matter is sufficiently dense.

The idea is to generalize the usual cascade prescription to evaluate the probability of a particle a to scatter off another particle b which traditionally is expressed in terms of an effective area of interaction with radius

$$\hat{b}_{ab}^{max}(p_{\perp cut}^2) = \sqrt{\frac{\hat{\sigma}_{ab}(\hat{s}, p_{\perp cut}^2)}{\pi}} \quad (134)$$

which sets the *maximum* allowed impact parameter for the pair a, b under consideration. In the PCM, $\hat{\sigma}_{ab}$ is the perturbative parton-parton cross-section (97). Now the unpleasant feature of this approach is that the maximum allowed impact parameter (134) depends sensitively on $p_{\perp cut}$, because the cross-section $\hat{\sigma}_{ab}$ rises at small momentum transfers as $1/p_{\perp cut}^2$ (c.f. discussion in Sec. 4.2.1). Therefore already a small variation of $p_{\perp cut}$ results in a sizeable change of the collision probability, since the effective interaction area varies as $\pi(\hat{b}_{ab}^{max})^2 = \hat{\sigma}_{ab}(\hat{s}, p_{\perp cut}^2)$. However, this sensitivity can be overcome, provided the density of scatterers is sufficiently large, by implementing a dynamical screening of the divergent long wavelength contributions to the interaction probability. In nuclear collisions, pA or AB , involving at least one heavy nucleus ($A \gg 1$), the population of partons in the central region becomes very dense [82, 86], so that the probability of finding $n > 1$ collision candidates is almost unity. One can therefore define a *minimum* impact parameter

$$\hat{b}_{ab}^{min} = \min_{b=1, \dots, n} [\hat{b}_{ab}] , \quad (135)$$

as illustrated in Fig. 44. Then, if $n \gg 1$, it is almost always possible to find a collision partner much closer than the distance $\hat{b}_{ab}^{max}$ set by the cross-section $\hat{\sigma}_{ab}$, so that

$$\frac{\hat{b}_{ab}^{min}}{\hat{b}_{ab}^{max}(p_{\perp cut}^2)} \ll 1 . \quad (136)$$

As a consequence the parameter $p_{\perp cut}$ loses its sensitive influence and the cross-section $\hat{\sigma}_{ab}$ is naturally cut off at

$$(p_{\perp})_{ab} \equiv \frac{1}{b_{ab}^{min}} > p_{\perp cut} . \quad (137)$$

In other words, the long range part of the Coulomb type color field is screened below $(p_{\perp})_{ab}$, where each collision pair a, b has its individual value of $(p_{\perp})_{ab}$, depending on the local density of surrounding partons.

Fig. 45 shows the A -dependence of the average impact parameter $\langle \hat{b} \rangle$, obtained from accumulated statistics of all parton-parton scatterings during the first 3 fm in pA collisions at the two considered energies $P = 100$ GeV and $P = 3000$ GeV. Evidently the values of $\langle \hat{b} \rangle$ change insignificantly for light nuclei ($A \leq 32$), then significantly decrease roughly as $A^{-1/6}$ as one proceeds to heavier nuclei, and finally seem to saturate for very heavy nuclei. This behavior is consistent with what one would expect from a simple geometrical consideration [154]. The essence is, that this classical screening mechanism considered here indeed cuts off scattering at impact parameters $\hat{b} < \hat{b}^{max}$ which is determined by the parton dynamics. Thus $\langle \hat{b} \rangle$ can be associated with a screening length. In plasma physics the natural regularization of Coulomb-type infrared divergence is provided by the Debye screening length λ^D [160]. Therefore an identification of $\langle \hat{b} \rangle$ with λ^D appears to be plausible. Although this concept is a rather simple minded approach to the screening problem, it is satisfactory that the extracted values of $\langle \hat{b} \rangle$ coincide approximately with results for the Debye screening length λ^D in a quark-gluon plasma [160] formed in heavy ion collisions at RHIC and at LHC with typical values $\lambda^D \simeq 0.39$ (0.13) fm at $P = 100$ (3000) GeV [161].

(iii) Hadron production: Finally let me address the question of how the properties of parton saturation and dynamical screening might affect hadronic observables in pA collisions. In Fig. 46 the dependence of the number dN_{pA}^{ch}/dy of produced charged particles on the nuclear mass number A around $y = 0$ is exhibited for $P = 100$ GeV and $P = 3000$ GeV, respectively. The solid lines with full circles display the results of the calculations including the effects of parton saturation and dynamical screening discussed before, whereas the dashed lines with open circles display the neglect of these effects. The multiplicity of secondary hadrons with $|y| < 2$ (more than 80 % percent of which are pions) shows for light nuclei a strong increase $\sim A^{0.62}$ ($A^{0.65}$), but for $A \gtrsim 40$ the multiplicity growth becomes progressively slower, tending to a $\sim A^{0.33}$ ($A^{0.15}$) behaviour for $P = 100$ (3000) GeV. The comparison between the full and dashed curves shows that parton saturation and screening effects significantly reduce the particle production the more the heavier the nuclei. For ^{238}U the relative suppression is about 15 (20) %, and for very heavy nuclei ($A \gtrsim 200$) the multiplicity growth around $y = 0$ clearly saturates.

In Fig. 47 the calculated average $\langle p_{\perp} \rangle^{ch}$ per secondary charged hadron is plotted versus

A. One can extract that the average $\langle p_{\perp} \rangle^{ch}$ per particle increases $\propto A^{\alpha}$ with $\alpha = 0.11$ (0.18) for $P = 100$ (3000) GeV which is close to a $A^{1/6}$ -scaling. This growth pattern is however non-trivial [154]. It is strongly correlated with the amount of particle production by cascading partons. The latter increases the slower with A the heavier the nucleus, so that for a fixed amount of total transverse momentum produced, the average $\langle p_{\perp} \rangle^{ch}$ per particle is enhanced.

Although the above considerations concerning nuclear and medium effects are rather qualitative at this point, they show that it is very important to study and understand in detail, both theoretically and experimentally, the various mechanisms and their possible interference. More accurate predictions will be calculable once these medium effects are better understood.

6. HEAVY ION COLLISIONS AT RHIC AND LHC: FROM PRE-EQUILIBRIUM TO PARTON PLASMA FORMATION

As mentioned in Chap. 1, one of the major goals of the experimental programs at the future heavy ion colliders, RHIC and LHC, is to detect evidence for the predicted QCD phase transition to a quark gluon plasma. In this context it is of central importance to analyze whether the dissipative dynamics of partons during the early stage of ultra-relativistic heavy ion reactions indeed yields a rapid materialization of the quanta in the central collision region and eventually to the formation of a hot and dense plasma of deconfined quarks and gluons [3, 4, 5]. In other words, the question is whether copious minijet production and parton cascading is a sufficient mechanism to drive the system to kinetic and chemical equilibrium. In order to attack these issues in an explorative way by both experimental and theoretical research, it is necessary to find characteristic signals of quark gluon plasma formation that enable to extract what happens at the very beginning - the "first few fm " [12] - of the nuclear collisions. Numerous theoretical papers addressing the properties and possible unambiguous signatures of this new form of matter have been published (for an overview see Refs. [162, 163, 164]). The most prominent proposed characteristic features [165] are the spectra of photons [166, 168] and dileptons [169, 170, 171, 172, 173, 174] produced during the early stage of the collisions, as well as the yield from pre-equilibrium and plasma of strangeness [176, 177, 83, 178] and charm [179, 178, 180, 181] that should show up in the measurable hadronic fragments.

In the past it was most commonly *assumed* that the quanta which are produced in the central rapidity region by primary parton scatterings quickly reach a state of local thermodynamic equilibrium. In the past, such an assumption was the convenient basis of most theoretical studies addressing the properties of a quark gluon plasma. However, the

validity of this assumption can only be checked by actually performing a dynamical calculation that evolves the initial partons smoothly from their first hard scattering, through a pre-equilibrium state, into a fully equilibrated plasma. Two of the most crucial questions in this context are:

- (i) Does the nuclear collision system have time enough to reach a state of thermal *and* chemical equilibrium with sufficiently large volume and lifetime so that it may leave characteristic marks on the subsequent expansion and cooling?
- (ii) What are the signals that enable one to observe the formation of such a plasma state during the very early stage of the collision, i.e. signatures that survive the complex evolution of comparably long duration from the deconfined plasma phase to the hadronization yielding the final observed particles?

Therefore it is essential to learn about the time scales, energy densities and entropy production associated with the development of the parton matter during the early stage (up to $\approx 3 \text{ fm}$) of heavy ion collisions. In particular, an understanding of the pre-equilibrium dynamics and the characteristic properties of a realistic parton plasma in terms of the gluon and quark composition is inevitable for finding unambiguous signatures that can be identified with the experimental particle detectors.

This Chapter investigates these issues in the context of the dynamical space-time evolution of the parton distributions in AA collisions, mostly for the illustrative case of central $Au + Au$ collisions at RHIC and LHC energies, $\sqrt{s} = 200 \text{ A GeV}$ and $\sqrt{s} = 6000 \text{ GeV}$, respectively. First the properties of thermalization and chemical equilibration will be addressed, and subsequently the consequences and implications for experimentally measurable signatures will be discussed (production of dileptons, strange and charmed particles).

6.1 Characteristics of the space-time evolution of AA collisions

Recall that in the kinetic description of the PCM all the information about the dynamical evolution of the many-parton system is contained in the time dependent single-particle distributions $F_a(p, r)$, eq. (30), for the various parton species a in phase-space. Starting from the initial parton distributions (79) at $t = t_0$, the evolution of the phase-space distributions $F_a(p, r)$ is governed by the transport equation (38), and reflect at any point of time $t > t_0$ the state of the many-parton system in phase-space element around $\vec{r}$ and $\vec{p}$. As explained in Chap. 3, by extracting these distributions, one can calculate the space-time dependent macroscopic quantities (50)-(54), number density $n_a(r)$, energy density $\varepsilon_a(r)$, entropy density $s(r)$, pressure $p(r)$, and even estimate the effective temperature $T(r)$. Employing the method of phase-space discretization (c.f. Sec. 3.4), the parton distributions can be directly obtained at any point of time during a nuclear collision. By evaluating from

those distributions the particle currents energy momentum tensor, etc., along the formulae (43)-(47) in the local rest frame of each phase-space cell, one obtains spatial profiles of e.g. number, energy, and temperature distributions.

It is important to keep in mind that exclusively the *real* particles, being either on mass shell or time-like, i.e. $p^2 = m^2 \geq 0$, must be taken into account when calculating the thermodynamic properties of the system. This includes the initial valence quarks as well as those partons that have been set ‘free’ due to mutual interactions. Only these partons contribute to the entropy production and determine the multiplicities and relative abundances of the final observable hadrons. The *virtual* partons, on the other hand, are understood to be the remaining space-like gluons and sea quarks with $p^2 = q^2 \leq 0$ that represent the color field and the vacuum polarization. These quanta remain pure field excitations whose effect in the process of confining hadronization is to contribute to the mass renormalization of the produced final hadrons.

In Refs. [82, 86, 182] detailed studies of the space-time structure of the parton kinetics in AA collisions were carried out within the PCM. Exemplary examples of how the parton distributions evolve in heavy ion collisions according to this approach are shown in Figs. 48-50, exhibiting qualitatively the characteristics of the partons’ space-time evolution in central $Au + Au$ collisions at RHIC energy $\sqrt{s} = 200$ A GeV. In Fig. 48 the parton distributions are shown at different proper times τ (defined below) along the longitudinal z -axis, and in Fig. 49 along the transverse $r_\perp$ -axis in the total center-of-mass frame of the colliding nuclei. Plotted are the numbers of partons per nuclear mass number A in central $Au + Au$ collisions ($A = 197$) at $\sqrt{s} = 200$ A GeV. The various shaded areas depict the contributions from virtual plus real and from exclusively real gluons, N_g/A , and from quarks plus antiquarks, $(N_q + N_{\bar{q}})/A$, whereas the solid black areas shows the valence quark distribution $(N_q - N_{\bar{q}})/A$, which reflects the baryon number distribution. In Fig. 48 the initial state shows the two strongly Lorentz contracted discs of valence quarks, surrounded by clouds of gluons and sea quarks. During the reaction an increasing number of these latter virtual quanta become real excitations due to multiple parton collisions and in addition new particles are created by gluon bremsstrahlung. By the time $\tau \simeq 2$ fm a rather homogeneous and isotropic plateau has developed, extending over the range $|z| \lesssim 1.5$ fm at $\tau = 2.4$ fm (the end of the calculation). The fast particles recede close to the speed of light as two beam fronts in opposite directions. Fig. 49 demonstrates that the transverse distributions (initially reflecting the Fermi distributions of the nucleons in the incoming nuclei) become increasingly dense, but do not exhibit an appreciable radial expansion during the first 3 fm. Fig. 50 shows the vertex distribution of secondary parton production and indicates the duration of perturbative parton cascade stage, from the actual beginning of the reaction on the parton level, up to the point of time when the partons

stream freely apart — measured in the nuclear CM frame. One sees that it takes a time $\approx 0.7 fm$ after the moment the nuclei began to overlap ($t = 0 fm$) until parton interactions substantially pick up, and the reaction is essentially complete after about $\approx 3 fm$.

For the following it is convenient to use phase-space variables appropriate for a Lorentz invariant description. The *proper time* τ is defined as [39]

$$\tau = \text{sign}(t - t_0(s)) \sqrt{(t - t_0(s))^2 - z^2} , \quad (138)$$

where t is the center-of-mass time and $t_0(s)$ is its value at the moment of maximum overlap of the colliding nuclei, e.g. $t_0 \simeq 0.8 fm$ for $\sqrt{s} = 200 A \text{ GeV}$ (c.f. Fig. 48). Thus, at $z = 0$ in the CM frame (77), the initial nuclear contact occurs at $\tau \simeq -0.8 fm$ and the maximum density is achieved at $\tau \simeq 0$. For the space-time variables one commonly employs the *space-time rapidity* η and the *transverse coordinate* $\vec{r}_\perp$,

$$\eta = \ln \left(\frac{\tilde{t} + z}{\tau} \right) , \quad r_\perp = \sqrt{x^2 + y^2} , \quad (139)$$

where $\tilde{t} \equiv t - t_0 \geq 0$. Thus, $r^\mu = (\tau \cosh \eta, \tau \sinh \eta, \vec{r}_\perp)$. The energy-momentum variables are described in terms of the usual *rapidity* y and the *transverse mass* $m_\perp$,

$$y = \ln \left(\frac{E + p_z}{m_\perp} \right) , \quad m_\perp = \sqrt{p_\perp^2 + m^2} , \quad (140)$$

where $p_\perp = \sqrt{p_x^2 + p_y^2}$, so that $p^\mu = (m_\perp \cosh y, m_\perp \sinh y, \vec{p}_\perp)$. Finally, let me define the *central collision region*, for $\tau \geq 0$, as a cylindrical expanding volume bounded by

$$-1 \leq \eta \leq 1 , \quad 0 \leq r_\perp \leq 5 fm . \quad (141)$$

6.2 Thermalization

The notion *thermal equilibrium* in statistical thermodynamics of many-particle systems is usually defined on account of Boltzmann's H -theorem [80], as the state in which there is no entropy production: if the distribution function tends to a definite limit as time progresses, the state of the system develops into an equilibrium state. In this steady state the entropy of the system attains its maximum value, implying that a necessary condition for equilibrium is that the entropy production vanishes in space-time. This condition, together with the requirement that the equilibrium distribution function must be a solution of the transport equation, uniquely determines its form. These equilibrium distributions are given by eq. (59). An equivalent way of defining thermal equilibrium is the condition that the momentum distribution of the particles becomes isotropic in the rest frame of the

matter, as is explicit in the equilibrium distributions. Therefore the term *kinetic equilibrium* is often used synonym for thermal equilibrium.

On the basis of this understanding, one can analyze the thermodynamic evolution of the parton distributions in nuclear collisions [82, 86, 182]. Figs. 51 and 52 show results from calculations of $Au + Au$ collisions with zero impact parameter at collider energies ranging from $\sqrt{s} = 200$ A GeV to $\sqrt{s} = 6300$ A GeV. They were obtained by discretizing the central collision region (141) into thin slabs of $\Delta z = 0.05$ fm and evaluating the parton distributions as explained in Sec. 3.4, and with these the thermodynamic quantities (50)-(53) in the local rest frame of each slab.

In Fig. 51 the time development of the partons' momentum distributions in the central region (141) of $Au + Au$ collisions at $\sqrt{s} = 200$ A GeV is displayed, individually for gluons and the various quark flavors (scaled by powers of 10). Plotted are the invariant distributions

$$\frac{1}{\bar{n}_a} \frac{d\bar{n}_a}{d^3p}(\tau) = \frac{1}{(2\pi)^3} \frac{1}{\pi R^2 \Delta\eta} \int_{-\Delta\eta/2}^{\Delta\eta/2} d\eta \int_0^R d^2r_\perp F_a(\tau, \eta, r_\perp; y, m_\perp) , \quad (142)$$

versus $p \equiv |\vec{p}|$, which were obtained from the actual parton distributions F_a , averaged over the cylindrical volume defined by (141). The sequence of plots demonstrate that both gluons and quarks indeed rapidly approach kinetic equilibrium in the central region, as is reflected by the increasingly thermal shape $\propto [\exp(E_a/T \pm 1)]^{-1}$. The highly non-thermal initial momentum distribution (79) of the partons in the incoming nuclei first develops a power-law shape characteristic for the harder primary parton collisions and the associated (mini)jet production. With progressing time this QCD power-law tail flattens out and the exponential slope at lower momentum ($p \lesssim 3$ GeV), which is already visible at $\tau = 0.2$ fm, becomes steeper, indicating the decrease in the effective temperature of the system.

Fig. 52 summarizes the thermodynamic properties of the system of partons implied by the time evolution of the partons' phase-space distributions. Shown are the pressure p , the densities n , ε and s , as well as the associated temperature $T \equiv 4/3(\varepsilon/s)$ as functions of the proper time τ (the curves are interpolations of the data points to guide the eye). The time dependence shown in Fig. 52 of the quantities n , p , ε , s and T may be parametrized in terms of negative powers of τ/τ_0 with the parameter $\tau_0 = 0.05$ fm:

$$n(\tau) = 565 \text{ fm}^{-3} \left(\frac{\tau}{\tau_0} \right)^{-0.90} \quad (143)$$

$$p(\tau) = 580 \text{ GeV fm}^{-3} \left(\frac{\tau}{\tau_0} \right)^{-1.25} \quad (144)$$

$$\varepsilon(\tau) = 1300 \text{ GeV fm}^{-3} \left(\frac{\tau}{\tau_0} \right)^{-1.15} \quad (145)$$

$$s(\tau) = 1800 \text{ fm}^{-3} \left(\frac{\tau}{\tau_0} \right)^{-0.85} \quad (146)$$

$$T(\tau) \equiv \frac{4}{3} \frac{\varepsilon(\tau)}{s(\tau)} = 960 \text{ MeV} \left(\frac{\tau}{\tau_0} \right)^{-0.30}. \quad (147)$$

This particular time-dependence remarkably resembles that of an ideal plasma of non-interacting partons in a longitudinally boost invariant scenario of the Bjorken model [100], in which case one has $n(\tau) \propto (\tau/\tau_0)^{-1}$, $p(\tau) \propto (\tau/\tau_0)^{-4/3}$, $\varepsilon(\tau) \propto (\tau/\tau_0)^{-4/3}$, $s(\tau) \propto (\tau/\tau_0)^{-1}$, and $T(\tau) \propto (\tau/\tau_0)^{-1/3}$.

Another good indicator for the degree and the time scale of a thermalization is the *specific entropy*, which is the produced entropy per secondary parton, since as stated above it necessarily must vanish in space-time when the system reaches an equilibrium state. Fig. 53a displays the time development of the specific entropy S/N for various collider energies [82]. The curves show a rapid build-up and relax approximately exponential to reach their final values between 3.9 and 4.3. Comparing these values with $(S/N)_{ideal} \simeq 4$ for an ideal gas of non-interacting massless quarks and gluons, one sees that the deviation between the resulting entropy of the realistic model calculation and the idealized case amounts only to a few percent. The variation of the final S/N values with collider energy is due to a weak temperature dependence of the ratio between entropy and number of partons. Since the temperature of the matter and the effective flavor degrees of freedom in the central region grow slowly with energy [82], an increase of the entropy per particle is natural. In Fig. 53b the relaxation times τ_{relax} associated with the entropy production (the time it takes to reduce the deviation from the equilibrium value by a factor of Euler's constant e) are plotted versus beam energy $\sqrt{s}/A$. The relaxation times are remarkably short, ranging from RHIC to LHC energies between $\simeq 0.4 \text{ fm}$ and $\simeq 0.15 \text{ fm}$, and decrease with growing beam energy roughly as $\tau_{relax}^{-1} \sim \ln(\sqrt{s}/A)$.

To summarize this Section: In heavy ion collisions at collider energies for RHIC and beyond, $\sqrt{s}/A \gtrsim 200 \text{ GeV}$, the highly non-thermal initial parton distributions appear to approach very rapidly a state of thermal equilibrium in the central region (within a fraction of a fm) with very high temperature and very dense particle and energy densities. Although the PCM description includes massive quarks and accounts for interactions among the partons, the system of partons looks effectively like an almost ideal gas.

6.3 Chemical equilibration

Notwithstanding the conclusion that the partons quickly reach a state of thermal equilibrium with isotropic momentum distributions in the central collision region, the question is, do the parton distributions also exhibit the correct (with respect to equilibrium) chemical composition of the matter in terms of quark and gluon admixtures?

As stated in the beginning of Sec. 6.2, when the system reaches a steady state of maximum entropy, then the solutions of the transport equation must take the form of

the equilibrium distributions (59). In this context the term *chemical equilibrium* applies generally to a system of various of distinguishable particle species and means that not only must the phase-space distributions for each species individually have thermal (isotropic) form, but moreover the relative proportions of the number of particles must reach an *equilibrium state of detailed balance* according to their respective degrees of freedom (c.f. discussion in Sec. 3.3). For example, in the simplified case of an ideal gas of massless quarks and gluons, the ratio of their respective number densities [82] would imply $N_{q+\bar{q}}/N_g = 9n_f/16$, i.e. for $n_f = 3$ there would be twice as much quarks plus antiquarks than gluons.

For the case of a closed, spatially bounded system, it is intuitively clear that correct detailed balance of chemical reactions (that change the number of particles) must and will at some point of time establish a chemical equilibrium. However, in nuclear collisions, after the colliding nuclei interpenetrate, the system is not bounded but flows apart, so that the actual question is: do the parton distributions have sufficient time to build up to their equilibrium form as dictated by kinetic theory, or do they approach the free streaming regime before the chemical reactions among quarks and gluons can establish a chemical equilibrium?

To answer this question, one must find a local, time-dependent measure of the degree of chemical equilibration. One way to gain quantitative insight is to compare for each species a the dynamically computed parton number densities n_a and energy densities ε_a (c.f. Sec. 3.2)

$$n_a(r) = \gamma_a \int dQ^2 \int \frac{d^3 p}{(2\pi)^3 2p^0} p^\mu u_\mu F_a(p, r) \quad (148)$$

$$\varepsilon_a(r) = \gamma_a \int dQ^2 \int \frac{d^3 p}{(2\pi)^3 2p^0} p^\mu p^\nu u^\mu u^\nu F_a(r, p) , \quad (149)$$

with the equilibrium densities $n^{(eq)}$ and $\varepsilon^{(eq)}$ for a perfectly equilibrated parton plasma at the *same* temperature,

$$n_a^{(eq)}(T, \mu_a) = \frac{\gamma_a}{2\pi^2} m_a^2 T \sum_{j=1}^{\infty} \frac{(\theta_a)^{j-1}}{j} K_2(j \frac{m_a}{T}) \exp(j \frac{\mu_a}{T}) \equiv \sum_{j=1}^{\infty} n_a^{(eq)}(j) \quad (150)$$

$$\varepsilon_a^{(eq)}(T, \mu_a) = \sum_{a=g,q_f,\bar{q}_f} \sum_{j=1}^{\infty} \frac{n_a^{(eq)}(j)}{j} \left(3T + m_a \frac{K_3(j \frac{m_a}{T})}{K_2(j \frac{m_a}{T})} \right) . \quad (151)$$

As before the factors $\gamma_g = 16$, $\gamma_q = \gamma_{\bar{q}} = 6$ are the products of spin and color degeneracy. In eqs. (148), (149) the four-vector $u^\mu \equiv u^\mu(r)$ is the space-time dependent flow velocity (48) of the matter. In (eqs. 150), (151) the quantum statistical factors are as before $\theta_g = +1$, $\theta_q = \theta_{\bar{q}} = -1$, m_a and μ_a denote the rest mass and the chemical potential, respectively, of parton species a , and the K_i are the usual Bessel functions of order i .

Note that the superscript ‘eq’ in (150), (151) refers to a kinetically equilibrated state of matter, which is not necessarily in chemical equilibrium. In the absence of any net baryon number or electric charge, perfect chemical equilibrium corresponds to $\mu_a = 0$ for all species a individually, since non-vanishing chemical potentials μ_a measure the deviation from chemical equilibrium (c.f. Sec. 3.3).

To make progress in comparing $n_a(\varepsilon_a)$ with $n_a^{(eq)}(\varepsilon_a^{(eq)})$ let me exploit the fact evident in the previous Fig. 49, that transverse expansion effects of the matter can be neglected at least during the first few fm , one can express all the thermodynamic quantities as functions of the proper time τ only, e.g.

$$n_a(r) = n_a(\tau, \vec{r}_\perp) \longrightarrow n_a(\tau), \quad \varepsilon_a(r) = \varepsilon_a(\tau, \vec{r}_\perp) \longrightarrow \varepsilon_a(\tau), \quad (152)$$

and define the ratios

$$\rho_a^{(n)}(\tau) = \frac{n_a(\tau)}{n_a^{(eq)}(T_a, 0)}, \quad \rho_a^{(\varepsilon)}(\tau) = \frac{\varepsilon_a(\tau)}{\varepsilon_a^{(eq)}(T_a, 0)}. \quad (153)$$

Here the chemical potentials μ_a are set to zero, so that the ratios ρ_a measure the deviations from perfect chemical equilibrium.

The *species dependent temperatures* $T_a \equiv T_a(\tau)$ [182] have been introduced to take into account the fact that the momentum distributions of the parton species are rather distinct due to the difference in their masses and the frequency of their interactions. This property reflected by Fig. 54a, in which the time dependence of the average transverse mass $\langle m_{\perp a} \rangle = \langle \sqrt{p_{\perp a}^2 + m_a^2} \rangle$ is shown separately for the gluons and the different quark flavors. Also displayed for comparison is the quantity $\langle m_{\perp}^{(ideal)} \rangle \simeq \langle p_{\perp}^{(ideal)} \rangle = \sqrt{6} T$ which would be the average transverse mass (momentum) of massless partons in an ideal QGP with effective temperature $T \equiv T(\tau)$, eq. (147). Obviously the gluons have a larger $\langle m_{\perp} \rangle$ than $\langle m_{\perp}^{(ideal)} \rangle$, whereas the u , d , and s quarks lie below the ideal curve and are approximately equal in magnitude. The c quarks have the largest $\langle m_{\perp} \rangle$, due to their large mass and the related fact that they are mostly produced in comparably large $p_{\perp}$ parton collisions.

To account for this distinctive property of the parton admixtures, one must replace the effective temperature T of the system as a whole by flavor specific temperatures T_a . To do so, two schemes have been employed, both of which give essentially the same results [82]: The first is a rather nuts and bolts prescription of employing the time-dependent average kinetic energy per particle $\langle E_a^{kin} \rangle \equiv \langle |\vec{p}_a| \rangle = \varepsilon_a/n_a$ as a measure of dissipation, and defining $T_a(\tau) = 1/c_a \langle E_a^{kin} \rangle$ with $c_g = 2.70$, $c_q = c_{\bar{q}} = 3.15$. The second, more founded but also more elaborate, scheme consists of equating the left hand side of eqs. (150) and (151) with the actual parton densities n_a , respectively ε_a , and solving for $T = T_a$ by a numerical root finding algorithm. Astonishingly, the difference between these two methods amounts to

only a few percent. The so extracted temperatures $T_a(\tau)$ are shown in Fig. 54b versus τ for gluons and the different species of quarks plus antiquarks, in comparison to the effective temperature $T(\tau)$, eq. (147), of the parton system as a whole. The essential implication is that the gluon temperature T_g is approximately equal to the overall temperature T , and that quarks are noticeably cooler than the gluons.

To show how this behaviour affects the approach to chemical equilibration, Fig. 55 displays the ratios (153) as functions of time τ for each parton species $a = g, u, d, s, c$ separately. An indication for the establishment of a perfect chemical mixture would be that these ratios tend to unity for gluons and each quark flavor *individually*. It is obvious that the system of partons can *not* establish perfect chemical equilibrium with proper admixtures of gluons and quarks, although, as argued before, it exhibits the thermodynamic behaviour of an ideal parton gas. The gluon density n_g , which rapidly increases and reaches its maximum around $\tau = 0.5 \text{ fm}$, approaches to within 10% of its equilibrium value at about $\tau = 1.2 \text{ fm}$. The dominance of gluon production causes a fast build up of $\rho_g(\tau)$ which overshoots its equilibrium value, but bends over as the phase-space density reaches its maximum and gluon fusion (absorption) processes become substantial. On the other hand, the densities of light quarks plus antiquarks, $n_q + n_{\bar{q}}$, build up only to about 60% of their equilibrium densities. The heavier quarks s and c achieve even less. The general decrease of ρ_a for $\tau \gtrsim 2 \text{ fm}$, is mainly to the diffusion of the quanta as the volume expands and the faster particles escape out of the central region. Analogous conclusions hold for ε_g , $\varepsilon_q + \varepsilon_{\bar{q}}$ and $\varepsilon_q - \varepsilon_{\bar{q}}$.

Two points are important to remark:

- (i) The failure of establishing a chemical equilibrium among the partons is *not* a conceptual defect in the PCM approach. As explained in Secs. 3.3 and 4.2 the model incorporates for each interaction process the reverse process so that detailed balance with respect to the various interaction processes could be established provided the system would have sufficient time. Both the large gluonic component in the initial nuclei and the dominance of gluonic interactions strongly favor the production of gluons rather than quarks. This almost naturally leads to a two-stage equilibration [12], in which the gluons must equilibrate first before the quark densities can build up towards their equilibrium value. In fact, when artificially increasing the quark-gluon coupling by a factor 10, relative to the gluon-gluon coupling, it was found that the production of quarks and antiquarks relative to the gluon yield is accelerated and enhanced such that the particle densities indeed come close to chemical equilibrium, with roughly equal relaxation times for both gluons and quarks. Another check was carried out by enclosing the nuclear collision system in a box in order to prevent the partons from streaming apart. The produced quarks and gluons were reflected from the walls and

eventually established also in this hypothetical case a chemical equilibrium within 5-10 fm (depending on the energy).

- (ii) The property of a decrease of the ratios ρ_a after $\tau \gtrsim 1 fm$ in Fig. 55 does *not* mean that the number of particles decreases. Rather than that the bending over of the curves originates from the fact that the partons have a broad velocity distribution, which causes a dilution of rigidly defined spatial volume of the central collision region (141). With progressing time more and more faster particles escape from this volume so that it is natural that the densities drop. Although this may appear as an artifact due to an arbitrary division into central region and beam fragmentation regions, it is a physical effect of the continuous rapidity distribution of the expanding parton matter, showing that a naive single-fluid expansion is an oversimplified picture.

Independent analyses of parton equilibration properties in heavy ion reactions at collider energies lead to very similar results and confirm that the previously believed standard scenario (see e.g. [100, 183] which assumes perfect chemical equilibration of the partons within a proper time of 1 fm , is certainly too naive. For example, Shuryak [12] found a two-stage equilibration with a fast equilibration of gluons ($\tau_g \approx 0.5 fm$) but a substantially longer equilibration of quarks ($\tau_q \gtrsim 2 fm$). Biro et al. [13] derived and solved a set of rate equations describing the chemical equilibration of partons including medium effects. The results imply again a clear dominance of gluon degrees of freedom over quarks, as well as the lack of chemical equilibration. In Fig. 56 the time evolution of the "fugacities" λ_g and λ_q is shown together with the estimated temperature in $Au + Au$ collisions at RHIC and LHC energies. The fugacities measure the underpopulation of partons in phase-space and reflect the deviation from a perfect chemical equilibrium in which case $\lambda_g = \lambda_q = 1$. Also Xiong and Shuryak [153] investigated the time development of parton equilibration in terms of gluon and quark fugacities. Again, for realistic initial conditions, they obtained the conclusion that the associated time scale is too long for chemical equilibration to be attained.

6.4 The "hot glue scenario"

The previous considerations regarding the parton evolution during the first few fm in ultra-relativistic heavy nucleus-nucleus collisions suggest a drastic modification of the "old" space-time picture as in e.g. the Bjorken model [165, 100]: The kinetics of interacting quarks and gluons during this early stage is governed by very frequent primary semihard scatterings that initiate the development of multiple interrelated cascades accompanied by copious particle production. Both the parton collision processes and the gluon emission processes are the driving forces for a very fast redirection of the large initial longitudinal

momentum in transverse direction and a rapid dissipation and redistribution of the energy in the system. As a consequence the initial partons in the colliding nuclei evolve from their first hard scatterings through a pre-equilibrium stage into a kinetic equilibrium in the central collision region. In other words, the highly non-thermal initial parton distributions develop smoothly into thermal distributions corresponding to very high temperatures and very large energy densities. Based on the investigations of [82, 86, 171, 178], and on other analyses [13, 180, 12, 173, 153, 184], the current understanding of the parton evolution in high energy nuclear collisions during the first 2-3 fm after the nuclear contact at τ_c may be summarized in four stages (here $\tau_c < 0$, with $|\tau_c| \simeq 0.5 - 1 \text{ fm}$, is defined such that $\tau = 0$ is the moment of maximum nuclear overlap [82]):

- 1) $\tau_c \leq \tau \lesssim 0.2 \text{ fm}$: The vehement materialization of initial virtual partons plus associated gluon emission produces most of the entropy and results in a large number of 'few GeV' excitations, which are mostly gluons.
- 2) $0.2 \lesssim \tau \lesssim 0.4 \text{ fm}$: Both the gluons and quarks rapidly reach a local kinetic equilibrium in the central collision region with approximately isotropic momentum distributions in the local restframe of the matter.
- 3) $0.4 \lesssim \tau \lesssim 2 \text{ fm}$: The gluons approach complete equilibration - the saturation of the phase-space density towards its chemical equilibrium value. The phase-space population of quarks evolves significantly slower, mostly by gluon induced processes, and yields at most 1/2 to 2/3 of the quarks' chemical equilibrium densities.
- 4) $\tau \gtrsim 2 \text{ fm}$: The gluon dominated plasma expands and cools without further significant change of its chemical composition.

Thus, the early evolution of these nuclear collisions is governed by the dominant role of gluons, due to their large interaction probability and the large gluonic component in the initial nuclear wavefunctions. This leads to a *hot glue scenario* [12], in which the large number of initially produced energetic partons create a high temperature, high density plasma of predominantly hot gluons and a considerably smaller number of quarks. This scenario is schematically illustrated in Fig. 57, where the different space-time regions are indicated. In the central region the system evolves from initial materialization to a hot, dense parton plasma with subsequent hydrodynamic expansion; in the beam fragmentation region color strings and color ropes may be formed that extend between the wounded nuclei and the surface of the parton plasma [2].

Clearly this optimistic picture is rather qualitative at this point, because many uncertainties and open questions remain to be better understood before a truly quantitative

decription can emerge (I will return to these issues in Chap. 7). However, if such extreme conditions as in this *hot glue scenario* would indeed be established at early times in the central collision region, then this would substantially modify previous expectations for a parton plasma formation, and its experimental signatures should be clearly measurable [12, 82]:

- (i) *Lepton pair and direct photon production:* The spectra of dileptons and direct photons produced during the early stage of the parton cascade evolution should exhibit a rather different behaviour than the spectra obtained in the "old", standard thermal scenario [169]. At early times most quark pairs are produced from 'hot' gluons via $gg \rightarrow q\bar{q}$ and $g^* \rightarrow q\bar{q}$ and therefore should reflect the momentum distribution of the gluons with their high temperature.
- (ii) *Strangeness equilibration:* The enhancement of total strangeness [176] depends on the interrelation of the strange quark mass m_s and the thermal quark mass $m_{th} \sim T$. For high temperatures as $T \gtrsim 300$ MeV, the thermal mass is considerably larger and there should be no essential difference between initially produced u , d and s quarks. If the multiple strange quark pairs, once produced, do not annihilate easily, there should be an enhanced strangeness production observable.
- (iii) *Enhanced production of charm:* Direct charm production in high temperature quark gluon plasmas via $gg \rightarrow c\bar{c}$ is expected to be enhanced significantly during both the pre-equilibrium stage [180] and the QGP evolution [179], for which $c\bar{c}$ production by hot gluons strongly increases with the initial temperature T_i of the thermalized parton matter. For example, the thermal $c\bar{c}$ production rate was estimated [181] for central $Au + Au$ collisions at RHIC to give per collision event about 1 $c\bar{c}$ for $T_i = 400$ MeV, 10 $c\bar{c}$ for $T_i = 550$ MeV and already 100 $c\bar{c}$ for $T_i = 700$ MeV, which clearly exceeds the standard parton model predictions of about 0.8 $c\bar{c}$ without a QGP formation in these collisions [179]. Therefore direct charm production is a very sensitive 'thermometer' that could be an evident signal for a QGP formation.

These issues will be addressed in the following Secs. 6.5 and 6.6.

Aside from these thermodynamic aspects, nuclear and dense medium effects (as partly investigated in Chap. 4) should have significant detectable effects on the final particle spectra and multiplicity distributions in heavy ion collisions. It is a fundamental question on its own right – how is QCD in the hot and ultra-dense nuclear environment modified as compared to QCD in dilute systems or vacuum? This also is of great importance for the future collider experiments at RHIC and LHC, where nuclear effects and dense matter phenomena must clearly deviate naive extrapolations from hadron-hadron collisions.

6.5 Dilepton production

The emission of lepton pairs or photons during a high energy nucleus-nucleus collision has been of great interest for a long time (see e.g. reviews [167, 168]). There are two reasons for this interest. First, the production rates of these electromagnetically interacting particles are strongly increasing functions of the temperature of the excited matter. Therefore, most high energy lepton pairs and photons should be created by parton interactions early in the collision when the energy density, and the energy available in parton scatterings, is the greatest. Second, these particles escape from the system without rescattering and carry information about the high density matter unscathed to the detectors. Conversely, hadrons are not produced until late in the nuclear collision when the energy density is much lower, and their final spectrum is influenced by their last scattering. These two facts make lepton pairs and photons good probes of the early, highly excited pre-equilibrium and plasma phases of the collision. Lepton pairs (synonym dileptons) have the advantage over photons that they have a very useful extra degree of freedom: the invariant mass of dileptons is variable, whereas for photons it is fixed to be zero. In this Section I will concentrate on dilepton emission from cascading partons during the very early stage of nuclear collisions at ultra-relativistic energies; however the following method of analysis is similarly applicable to photon emission.

In the past the usual procedure to quantify dilepton production in heavy ion collisions was to calculate the Drell-Yan contribution and to consider it a background to the yield from an assumed thermalized quark-gluon plasma, the temperature of the latter decreasing with time because of hydrodynamic expansion [11]. Clearly, this is an unsatisfactory state of affairs, because there is an uncertainty in choosing the formation time and temperature of the plasma. The PCM approach on the other hand provides a dynamical description that evolves the initial partons from their first hard scattering, through a pre-equilibrium stage, and into a fully thermalized plasma. Therefore one can study the distribution of radiated lepton pairs as it develops smoothly from the initial Drell-Yan type reactions all the way towards equilibrium radiation. Such an analysis has been carried out in Refs. [171, 172]. For an general overview of dilepton production in nucleus-nucleus collision and recent developments I refer to Refs. [167, 168].

In the PCM the production of lepton pairs has been calculated on the basis of the following elementary partonic processes, which involve the production of a time-like photon that subsequently decays into an l^+l^- pair (l is an electron or muon),

$$q + \bar{q} \rightarrow \gamma^* \rightarrow l^+ + l^- \quad (154)$$

$$q + \bar{q} \rightarrow \gamma^* + g \rightarrow l^+ + l^- + g \quad (155)$$

$$g + q(\bar{q}) \rightarrow \gamma^* + q(\bar{q}) \rightarrow l^+ + l^- + q(\bar{q}) \quad (156)$$

$$g + g \rightarrow \gamma^* + g \rightarrow l^+ + l^- + g \quad (157)$$

$$q(\bar{q}) \rightarrow \gamma^* + q(\bar{q}) \rightarrow l^+ + l^- + q(\bar{q}) . \quad (158)$$

Fig. 58a depicts these processes in terms of Feynman diagrams. This subset of processes is much more general than it seems at first sight, because the structure of arbitrarily complex parton cascades is built from joining elementary $2 \rightarrow 2$, $1 \rightarrow 2$ and $2 \rightarrow 1$ processes. In all these processes the amplitudes are not just the tree level, but are dressed with formfactors corresponding to emission and absorption of soft gluons, as explained in Sec. 4.2. Therefore multiparticle processes such as depicted in Fig. 58b, i.e. higher order processes or processes involving three or more partons in the incoming or outgoing channel are automatically included without overcounting.

The role of the partons' time-like form-factors in this context needs some special attention [174, 175]. Consider a typical parton branching chain initiated by a hard collision, say $g + q \rightarrow g_0 + q_0$, as depicted in Fig. 58c. As discussed in Chap. 2, the probability of the quark q_0 with virtuality Q_0^2 to branch into gluon g_1 and quark q_1 with virtualities Q_1^2 and $Q_1'^2$ is determined by the formfactor of the quark as it effectively describes the gluon substructure, i.e. the number of gluons in the quark q_0 . When one asks for the relative probability that instead of gluon g_1 a photon γ_1 is radiated off q_0 , one has to be aware that the photon does not have a comparable QCD formfactor. This means that the gluon g_1 can itself radiate furtheron, but if instead the γ_1 is emitted, it will not cause any further gluon radiation. Therefore one cannot just count the number of gluon emissions in a cascade and weigh it with the ratio of probabilities for photon to gluon emission, because this would overestimate the photon production rate and in turn the dilepton yield. Rather than that a certain cascade branch must be considered terminated whenever a photon is produced instead of a gluon. As a consequence, the ratio of probabilities for photon emission to gluon bremsstrahlung in general is not a simple function proportional to α_{em}/α_s depending on $Q^2 = M^2$, but involves the form factors for branchings into gluons or $q\bar{q}$ pairs in a non-trivial way. In Ref. [174] a careful analysis of this issue was carried out. with the essential result that this effect is of significance only for dilepton masses $M \gtrsim 8$ GeV, i.e. in the Drell-Yan region.

On account of the processes (154)-(158) the space-time history of dilepton production in a nuclear collision can then be extracted [171] from the time development of the parton phase-space distributions. Since detectors in an experiment register the yield of lepton pairs only in a space-time integrated way, one has to calculate the experimentally observable, space-time integrated number distribution of produced l^+l^- pairs with invariant mass

squared M^2 [171]:

$$\begin{aligned} \frac{dN}{dM^2} = \int \tau d\tau d\eta d^2r_\perp \left\{ \sum_{a,b,X} \int_0^\infty d\hat{s} \left(\frac{d\hat{\sigma}_{ab \rightarrow l^+l^-X}/dM^2}{\sum_{c,d} \hat{\sigma}_{ab \rightarrow cd}} \right) \left(\sum_{c,d} \frac{dn_{ab \rightarrow cd}(\hat{s}; r)}{d\hat{s}} \right) \right. \\ \left. + \sum_{a,Y} \int_0^\infty dq_a^2 \left(\frac{d\hat{\Gamma}_{a \rightarrow l^+l^-Y}/dM^2}{\sum_{c,d} \hat{\Gamma}_{a \rightarrow cd}} \right) \left(\sum_{c,d} \frac{dn_{a \rightarrow cd}(q_a^2; r)}{dq_a^2} \right) \right\}. \quad (159) \end{aligned}$$

where $d^4r = \tau d\tau d\eta d^2r_\perp$, with proper time τ , space-time rapidity η and transverse coordinate $\vec{r}_\perp$ defined in (138) and (139). The dilepton production cross-sections $\hat{\sigma}$ for $ab \rightarrow l^+l^-X$ and widths $\hat{\Gamma}$ for $a \rightarrow l^+l^-Y$ are given in terms of the squared amplitudes for the production of a virtual photon of mass M , weighted with quantum statistical factors involving the phase-space densities $F \equiv F(p, r)$,

$$\begin{aligned} \frac{d\hat{\sigma}_{ab \rightarrow l^+l^-X}}{dM^2} = \frac{\alpha}{48\pi^2 \hat{s}^2 M^2} \int_{\hat{i}_{min}}^{\hat{i}_{max}} d\hat{t} F_a F_b [1 - F_{l^+}] [1 - F_{l^-}] \prod_{k=5}^m [1 + \theta_{x_k} F_{x_k}] \\ \times (2\pi)^4 \delta^4 \left(\hat{s} + \hat{t} + \hat{u} - M^2 - \sum_{k=5}^m m_{x_k}^2 \right) |M_{ab \rightarrow \gamma^* X}(\hat{s}, \hat{t}, M_{\gamma^*}^2)|^2, \quad (160) \end{aligned}$$

$$\begin{aligned} \frac{d\hat{\Gamma}_{a \rightarrow l^+l^-Y}}{dM^2} = \frac{\alpha}{6\pi q_a M^2} \int_{q_{min}^2}^{q_{max}^2} dq^2 F_a [1 - F_{l^+}] [1 - F_{l^-}] \prod_{k=4}^n [1 + \theta_{y_k} F_{y_k}] \\ \times (2\pi)^4 \delta^4 \left(q_a^2 - M^2 - \sum_{k=4}^n q_{y_k}^2 \right) |M_{a \rightarrow \gamma^* Y}(q_a^2, q^2, M_{\gamma^*}^2)|^2. \quad (161) \end{aligned}$$

In (159) $n_{ab \rightarrow cd}$ and $n_{a \rightarrow cd}$ denote the corresponding number of parton interactions that occur at given $\hat{s}$ and space-time point r . The kinematic variables $\hat{s}$, $\hat{t}$ and $\hat{u}$ refer to a pair of colliding partons ab , and q_a^2 is the invariant mass of a virtual parton a . The squared amplitudes for the processes (155)-(157) and for the pure QCD processes $ab \rightarrow cd$ and $a \rightarrow bc$ are evaluated with the standard running strong coupling $\alpha_s(M^2)$, where M^2 sets the scale, chosen as $M^2 = p_\perp^2$, except for $q\bar{q}$ annihilation, for which $M^2 = \hat{s}$, and for gluon bremsstrahlung, where $M^2 = q_a^2$. Furthermore, a K -factor was employed to account for higher order corrections to the lowest order cross-sections in (160). Thus, by summing over the complete space-time history of the partons' evolution in phase-space up to a certain time τ according to eq. (159), one obtains the yield of dileptons as a function of time. This allows to observe how the total lepton pair production develops from the first nuclear contact via pre-equilibrium towards thermalization of the parton system.

As a check of consistency of this method, Fig. 59 shows the dilepton yield in pp collisions as computed in this manner. For masses $M > 6$ GeV the Monte Carlo points agree well with the standard Drell-Yan yield calculated in the usual parton model prescription (see e.g. [46]). For lower masses the PCM results predict slightly higher yield than the Drell-

Yan rate, because even in pp collisions there is a small contribution from multiple parton collisions (c.f. Sec. 5.1, Fig. 34) which show up as the invariant mass gets smaller.

Proceeding to nucleus-nucleus collisions, Fig. 60 shows the total dilepton distribution $dN/dM^2 dy$ at $y = 0$ in $Au+Au$ collisions at RHIC energy (full curve), integrated over space-time up to $\tau = 2 fm$, as well as separate contributions grouped according to the number of parton interactions. The three patterned curves correspond to processes involving exclusively primary partons ('Drell-Yan'), primary plus secondary partons, and only secondary⁴ partons. Below 10 GeV the largest contribution comes from secondary-secondary, followed by primary-secondary. At around 10 GeV all three contributions are approximately equal. Above 10 GeV it may be surmised that Drell Yan is the largest single component. The primary-primary contribution alone should be approximately the same as the total dilepton yield in pp collisions scaled by $A^{4/3}$, because in pp the other two contributions are negligible. This comparison is made in Fig. 61, which proves that the scaling is verified within statistical errors.

Fig. 62 depicts the contributions from the various processes (154)-(158) to the integrated dilepton yield per unit rapidity at $y = 0$. For $M \geq 4$ GeV the annihilation process $q\bar{q} \rightarrow l^+l^-$ dominates, but the process $gq(\bar{q}) \rightarrow l^+l^- q(\bar{q})$ also contributes considerably because of the much larger number of $gq(\bar{q})$ collisions. The low-mass pairs with $M \leq 2$ GeV are mainly from the bremsstrahlung processes $q(\bar{q}) \rightarrow l^+l^- q(\bar{q})$. The process $q\bar{q} \rightarrow l^+l^- g$ contributes about an order of magnitude less than the direct annihilation $q\bar{q} \rightarrow l^+l^-$ and the process $gg \rightarrow l^+l^- g$ is suppressed by 3-4 orders and therefore negligible. Notice that some of the individual yields turn over at smaller mass and actually decrease. This is due to the invariant cut-offs used in the PCM (c.f. Sec. 4.4). Since soft parton interactions are treated phenomenologically, they have not been included in the calculation of the dilepton yields in order not to spoil the results. It must be emphasized that the PCM results cannot, at present be trusted at invariant masses smaller than about 2.5–3 GeV.

It is interesting to compare these results with general considerations as done in Ref. [168]. The PCM predicts for central $Au + Au$ collisions at RHIC energy a rapidity density $dN/dy = 1.7 dN^{ch}/dy \approx 2000$ (c.f. Sec. 5.2, Fig. 37b and Table 4). From this one can estimate an initial temperature $T_0 = 550$ MeV and formation time $\tau_0 = 0.12 fm$ corresponding to a hydrodynamical expansion in the Bjorken model [169], or a free-streaming scenario as considered in [170]. Fig. 63 compares the PCM results for the dilepton yield with the hydrodynamic and free-streaming curves. In the mass range $2 \lesssim M \lesssim 5$ GeV the latter two are very close to the PCM prediction. At high masses $M > 6$ GeV the tail of

⁴Recall that 'primary' refers to those, in some sense pre-existing, partons pulled from the structure functions, whereas 'secondary' partons are those that have scattered at least once or have been newly created during the collision.

the PCM curve is significantly larger than the hydrodynamic and free-streaming curves. The reason for this is to be found in the previous Fig. 51 that shows the time evolution of the parton momentum distributions as they approach thermal equilibrium. Although the distributions become exponential over at least three orders of magnitude, there is always a high momentum tail which takes longer to thermalize. It is this tail which produces the high mass tail in the dilepton spectrum. These power law tails are ultimately due to hard perturbative QCD processes.

In this context it is also elucidating to compare the PCM approach to the *hot glue scenario* [12] mentioned before. Shuryak and Xiong [173] have calculated the dilepton mass spectrum in both the "hot glue scenario" and the "standard scenario", the latter corresponding to the Bjorken hydrodynamic model [100]. The resulting lepton pair yields are shown in Fig. 64 together with the usual Drell-Yan contribution. There is a qualitative agreement with the spectrum of Fig. 63 from the PCM calculation. The large number of energetic, hot gluons in the hot glue scenario gives a much harder energy spectrum of gluons and quarks than the standard scenario, a feature which in turn results in a much enhanced dilepton spectrum at higher masses. However, in Fig. 64 the Drell-Yan contribution wins at masses bigger than about 3.5 GeV, whereas in the PCM the contributions involving secondary partons dominate the Drell-Yan yield even at large masses, as can be seen from Fig. 60.

In conclusion, the computation of the lepton pair yield in central $Au + Au$ collisions at RHIC with the PCM results in a smooth evolution of the l^+l^- mass distribution from Drell-Yan towards quark-gluon plasma, with a clear dominance of pairs from pre-equilibrium and parton plasma over the Drell-Yan yield. Moreover, as analysed in [172], the transverse mass spectrum of dileptons indicates that the specific QCD formfactors of the involved partons result in a characteristic scale-breaking $M_\perp$ -spectrum. This really emphasizes the important role dileptons have in carrying information about the early high energy density parton dynamics to the detectors. Therefore experiments in the central rapidity region are of great importance to test the current understanding of the many-body dynamics of quarks and gluons.

6.6 Strangeness, charm, and bottom production

The properties of strangeness and charm production in high energy heavy ion reactions are similar as dilepton and photon emission, commonly believed to provide characteristic signatures for the formation of a deconfined parton plasma during the early stage of these reactions. Originally this idea was proposed many years ago by Shuryak [165]. Since then numerous investigations have lead to expectations of potentially observable enhanced strange hadron production [176, 177, 83] and J/ψ suppression [185, 186], respectively.

The enhanced abundance of strange particles is expected to be a good indicator for the establishment of a thermalized QGP, because for plasma temperatures $T \gg m_s$ ($m_s \simeq 150$ MeV) there should be no significant difference between u , d , and s quark production. The idea of a suppression of $c\bar{c}$ mesons J/ψ , ψ' , on the other hand, is based on the notion that $c\bar{c}$ pairs are produced mainly via primary hard collisions of energetic gluons during the pre-equilibrium stage up to shortly after the plasma formation (before the initial temperature of the plasma drops below the production threshold), and the J/ψ , ψ' mesons formed from these pairs may subsequently experience deconfinement when traversing the region of the plasma. Additional charm production in the plasma at later times would not occur due to the large charm quark mass ($m_c \simeq 1.3\text{-}1.5$ GeV) and the rapidly dropping temperature. Similar arguments hold for even heavier particles, such as the $b\bar{b}$ meson Υ . However there are other estimates [179, 187] that predict - rather than a suppression - an enhancement of J/ψ production in the plasma, because of the possibility of having several charmed quarks produced independently in close proximity of phase-space. Generally, the various approaches in the literature to calculate the yield strange and charm particles in nucleus-nucleus collisions are plagued by a number of uncertainties which require model dependent phenomenological descriptions. For instance, the assumption of a perfect plasma with a certain initial temperature and formation time, the uncertainty of the charm content in the colliding nuclei and of the value of the c quark mass, and, the lack of knowledge of the mechanisms that affect the conversion of the quarks produced in the plasma at early times into final hadrons at late times [192].

With the PCM it is possible to improve at least some of these shortcomings and to systematically investigate the beam energy and mass dependence of heavy quark production, extract formation times and associated initial plasma temperatures, as well as to separate the individual processes that contribute. In Ref. [178] such an investigation has been presented for strangeness, charm and bottom production in central AA collisions in the energy range between RHIC and LHC. Two different scenarios of heavy ion collisions were compared. The first scenario considers $Au + Au$ collisions *with* plasma formation by realistic simulation of the heavy ion reactions. The second scenario mimics $Au + Au$ collisions *without* plasma formation, on the basis of purely hadronic interactions, by simulating pp collisions and scaling to AA . As I will review now, the results imply a substantial enhancement of strange, charm, as well as bottom particle yields in AA reactions relative to pp collisions. The origin of these findings is again the formation of a high temperature gluon dominated plasma in the central region of $Au + Au$ collisions during the early stage.

The method of extracting the space-time history of the production of heavy quarks (antiquarks) Q ($\bar{Q}$) from the partons' evolution is analogous to the case of dileptons discussed in the previous Section. Each parton cascade is decomposed in a number of elementary

processes that determine the net yield of Q or $\bar{Q}$ through their detailed balance:

- (i) the *flavor creation (annihilation)* processes by which a $Q\bar{Q}$ pair is newly created (annihilated) through quark-antiquark annihilation, gluon fusion, or virtual gluon decay,

$$q + \bar{q} \leftrightarrow Q + \bar{Q} \quad (162)$$

$$g + g \leftrightarrow Q + \bar{Q} \quad (163)$$

$$g^* \leftrightarrow Q + \bar{Q} , \quad (164)$$

where $q = u, d, s, c, b$ and $Q = s, c$ or b .

- (ii) the *flavor excitation* processes in which a Q or a $\bar{Q}$ is knocked out of the sea of one of the incoming nuclei by a collision with a $q, \bar{q}$, or g ,

$$q(\bar{q}) + Q(\bar{Q}) \rightarrow q(\bar{q}) + Q(\bar{Q}) \quad (165)$$

$$g + Q(\bar{Q}) \rightarrow g + Q(\bar{Q}) . \quad (166)$$

Similarly as in Sec. 6.5 the subset of processes (162)-(166) is rather general, as it accounts for the complex space-time structure of multiparticle production from many internested cascades composed of elementary $2 \rightarrow 2$ collisions, $2 \rightarrow 1$ fusion processes and $1 \rightarrow 2$ branchings, plus higher order virtual and real corrections. The decomposition into the elementary processes (162)-(166) is simply a convenient accounting prescription for the different production mechanisms. In fact, all kinds of higher order processes are taken into account. For example, processes like $g + g \rightarrow Q + \bar{Q} + g + g$ or $Q + g + g \rightarrow Q + g$ are automatically included as compositions of the above elementary processes.

The space-time history of Q ($\bar{Q}$) production via (162)-(166) in a nuclear collision can be extracted from the time development of the phase-space distributions $F_a(p, r)$ through the rate equations

$$\partial_\mu n_a^\mu(r) = R_a^{gain}(r) - R_a^{loss}(r) \equiv R_a(r) , \quad (167)$$

where $n_a^\mu(r)$ are the local space-time dependent parton currents given by (43). The quantity

$$\begin{aligned} R_a(r) &= \sum_{processes\ k} \int \frac{d^3p}{(2\pi)^3 E} \left(I_a^{gain(k)}(r, p) - I_a^{loss(k)}(r, p) \right) \\ &= \sum_{processes\ k} \frac{1}{S_a} \int \prod_{i=1}^{n+m} \frac{d^3p_i}{(2\pi)^3 E_i} |\mathcal{M}_{eff}^{(k)}|^2 (2\pi)^4 \delta^4 \left(\sum_{i=1}^n p_i - \sum_{j=n+1}^{m+n} p_j \right) \\ &\quad \times \prod_{i=1}^n F_i(r, p_i) \prod_{j=n+1}^{m+n} (1 + \theta_j F_j(r, p_j)) \end{aligned} \quad (168)$$

is the net gain rate for producing a parton a at $r = (t, \vec{r})$ through any of the considered processes k (c.f. Sec. 4.2).

On account of the processes (162)-(166), the total net production rate for quarks $Q = s, c, b$ (and similar for antiquarks $\bar{Q}$) is

$$R_Q(r) = \sum_{a,b} \left[R_{ab \rightarrow Q\bar{Q}} - R_{Q\bar{Q} \rightarrow ab} \right] + \left[R_{g \rightarrow Q\bar{Q}} - R_{Q\bar{Q} \rightarrow g} \right] + \sum_a R_{aQ(\text{prim}) \rightarrow aQ} \quad , \quad (169)$$

corresponding to the sum of net rates for the flavor creation processes (162)-(164) through $Q\bar{Q}$ production and the flavor excitation processes (165), (166) via scatterings of primary Q out of the coherence of the initial nuclear wavefunctions (indicated by the label '*prim*'). The rates $R_{i \rightarrow f}$ on the right hand side are extracted from the parton cascade simulation [178]. To make connection with experimental quantities, the total restframe rates $R_Q(r)$ and $R_{\bar{Q}}(r)$ must be convoluted over the space-time history of the parton system, since detectors record particle multiplicities accumulated over the whole evolution. Using $d^4r = \tau d\tau d\eta d^2r_\perp$, where as before τ is the proper time, η the space-time rapidity and $\vec{r}_\perp$ the transverse coordinate, the experimentally relevant, space-time integrated number distributions of produced Q and $\bar{Q}$ with respect to their center-of-mass rapidity y are obtained by:

$$\frac{dN_{Q+\bar{Q}}}{dy} = \int \tau d\tau d\eta d^2r_\perp \frac{d}{dy} \left\{ R_Q(r) + R_{\bar{Q}}(r) \right\} \quad . \quad (170)$$

Thus, the Q plus $\bar{Q}$ yield from the parton processes (162)-(166) is accumulated by summing over the complete space-time evolution of the parton system in phase-space up to a certain time τ .

Fig. 65 shows the central rapidity density $dN_{Q+\bar{Q}}/dy|_{y=0}$ versus the normalized beam energy $\sqrt{s}/A$ of produced strange, charm and bottom quarks plus antiquarks ($Q = s, c, b$) in central $Au + Au$ collisions. In Fig. 65a the displayed energy dependence of $dN_{Q+\bar{Q}}/dy$ with plasma formation was obtained on the basis of eq. (170), whereas Fig. 65b shows the purely hadronic production, *without* plasma formation, which was mimicked by simulating pp collisions, and simply scaling the calculated multiplicities to $Au + Au$ [179]. As a consistency check, Fig. 65b also indicates the simple parton model result [179, 105] (thin lines), evaluated for pp collisions and scaled to $Au + Au$, by convoluting the proton structure functions with the relevant flavor production cross-sections. This latter case corresponds to taking into account only primary parton-parton collisions without subsequent cascading and gluon bremsstrahlung. Evidently, in the case of hadronic production, the PCM result generally agrees with the simple parton model estimate, although a steadily growing deviation with $\sqrt{s}/A$ from the simple parton model prediction is obvious. As already observed in the case of dilepton production (Fig. 59), the stronger increase within the PCM arises,

because multiple parton scatterings and $Q\bar{Q}$ production by radiative gluons from parton cascades play an increasingly important role, even in hadron collisions.

Comparing the two scenarios with and without plasma formation, one observes a substantial enhancement of more than one order of magnitude of the quark production for all three flavors s , c and b . Note that this enhancement comes on top of a substantial depletion of partons in the central rapidity region due to nuclear shadowing of the initial gluon and quark distributions and due to the medium effects discussed in Sec. 4.2.

Fig. 66 displays - in correspondence to Fig. 65a - the individual contributions from the processes (162)-(166) to the central rapidity density of strange and charm quarks plus antiquarks as a function of $\sqrt{s}/A$. One can infer some interesting features that elucidate the origin of this intense flavor production: First, the Q ($\bar{Q}$) flavor production is overwhelmingly driven by the gluonic processes $g^* \rightarrow Q\bar{Q}$, $gg \rightarrow Q\bar{Q}$, and $gQ \rightarrow gQ$. Second, among the $2 \rightarrow 2$ processes, the *flavor excitation* process (166), $gQ \rightarrow gQ$, outweighs the *flavor creation* process (163), $gg \rightarrow Q\bar{Q}$, over the entire displayed energy range, although the latter contribution rises steeply with $\sqrt{s}/A$ and becomes comparable in magnitude at and beyond LHC energy. Third, the $1 \rightarrow 2$ *flavor creation* process (164), $g^* \rightarrow Q\bar{Q}$, gives the dominant contribution for large energies ($\sqrt{s}/A \gtrsim 1$ TeV) and is mainly responsible for the drastic multiplicity increase at high energies.

What does this imply now for the potential plasma signatures of strangeness and charm production? The *abundance of strange particles* is a well known dense matter signal, discussed in detail in [176, 177, 83]. In a hot, gluon dominated plasma there should be no significant difference between u , d and s quark production, because the common thermal mass $m_u^{th} = m_d^{th} = m_s^{th} \propto gT$, acting as a production threshold is at high temperatures larger than the current quark masses, $m_q < m_q^{th}$. Indeed, the ratio of produced strange to light quark-antiquark pairs increases from 0.7 (RHIC) to 1 (LHC) as shown in Fig. 67b below. An *enhanced charm production* on the other hand has been advocated to serve as a sensitive thermometer for the initial temperature of QGP [12, 179]. In fact, in the calculations presented here, the majority of charm quarks from $c\bar{c}$ production is due to the large number of energetic gluons with several GeV energy-momentum. These gluons are mostly produced in primary parton collisions at very early times ($\tau \lesssim 0.3$ fm) and provide the main source of $c\bar{c}$ pairs. Therefore the charm quarks should have the momentum distribution of the gluons during the pre-equilibrium stage and upon thermalization, so that they reflect the initial temperature $T_0 \simeq T_g(\tau_0)$ around the plasma formation time τ_0 . The same argument holds also for $b\bar{b}$ creation, however this production rate is by more than an order of magnitude smaller and therefore more difficult to measure with sufficiently good statistics.

In Ref. [178] an independent method to extract the initial plasma temperatures T_0

was employed by comparing the PCM results for $Au + Au$ collisions of Fig. 67a with the multiplicities of produced s and c quarks expected from an ideal QGP. The resulting dependence of T_0 on $dN_{Q+\bar{Q}}/dy$ is plotted in Fig. 67a for $Q = s, c, b$, whereas in Fig. 67b the ratios

$$\rho_{Q+\bar{Q}}(s) = \frac{dN_{Q+\bar{Q}}/dy}{\frac{1}{2}(dN_{u+\bar{u}}/dy + dN_{d+\bar{d}}/dy)} \Big|_{y=0} \quad (171)$$

are shown. Fig. 67 suggests to infer the initial temperature T_0 by simply measuring the total strange or charm yield in the central rapidity region. Although it is to date not completely understood how these absolute multiplicities are reflected in the final hadron spectra, the ratios (171) in Fig. 67b of s, c and b quarks to the light quarks should be equal in magnitude to the measurable ratios of strange, charm and bottom mesons to pions, respectively, because of entropy conservation.

From these considerations, one would expect that the copious production of strange, charm, and bottom quarks during the pre-equilibrium stage of heavy ion collisions should have a significant impact on the production rates of hadrons containing these quarks. In particular the substantial enhancement of the $s\bar{s}$, $c\bar{c}$, and $b\bar{b}$ flavor creation processes (162)-(164) should testify in the production rates of the vector meson resonances ϕ , J/ψ , ψ' , and Υ , and may provide an observable signal for a QGP formation. However, there are a number of uncertainties to be aware of. First, the mechanisms for the formation of these vector mesons from the quark-antiquark pairs are not yet completely understood, and second, the fate of these mesons while traversing the dense nuclear environment before escaping is rather unclear. Both these issues have been addressed for charmonium and bottomonium production within various models [187, 188, 189, 190, 191] with predictions ranging from significant suppression to substantial enhancement of resonance production in a QGP scenario. It is nevertheless of interest to estimate the resonance production rates of ϕ , J/ψ , ψ' , and Υ from the yields of s, c , or b quarks resulting from the PCM calculations. In Ref. [178] a simple coalescence model on the basis of the hadronization scheme described in Sec. 4.3 was pursued to estimate one obtains the the rapidity densities $dN_R/dy|_{y=0}$ of resonances R . The resulting yield of $\phi, J/\psi, \psi'$ and Υ is shown in Fig. 68 as a function of $\sqrt{s}/A$ for central $Au + Au$ collisions *with* plasma formation. In correspondence with Fig. 65a, these particle yields reflect the copious $Q\bar{Q}$ production by the hot and dense parton matter in the central region of these reactions, with the effective initial temperatures T_0 , associated with the beam energy $\sqrt{s}/A$. These estimates are rather optimistic predictions, because it is assumed that all the $Q\bar{Q}$ bound states survive the plasma evolution and the later stages of the nuclear collision. Nevertheless it shows that a considerable increase of resonance production can be expected in heavy ion collisions, if a high density, high temperature environment is created as predicted by the parton cascade simulations [82, 86, 182].

I emphasize that the PCM results discussed here are to be understood as qualitative trends and to be perceived with precaution. Contrary to the previous case of dilepton emission, the calculation quark flavor production involves a number of uncertainties due to the current lack of better experimental knowledge:

- (i) *Intrinsic ('pre-existing') components of gluons and sea quarks:* The magnitude of gluon and s , c sea quark densities in the initial nuclear structure functions sensitively determines the production rates - especially at small x . A large gluon content means a large probability for flavor creation via $gg \rightarrow Q\bar{Q}$ and $gg \rightarrow gg^* \rightarrow gQ\bar{Q}$, whereas the $s\bar{s}$ and $c\bar{c}$ sea quark components determine the amount of flavor excitation through $gQ \rightarrow gQ$ and $qQ \rightarrow qQ$. Although Collins and Tung [193] and more recently Aivazis et al. [194] have shown how to consistently account for flavor excitation processes in perturbative QCD without double counting, it does not resolve the uncertainty of how much intrinsic components there are in nuclei. This requires further experimental information from structure function measurements at small x , which may soon be elucidated by the new HERA data.
- (ii) *The values of m_c , Λ , and structure function dependence:* The exact value of the charm quark mass, choice of renormalization scale Λ and of structure function parametrization are other uncertainties in the game of $Q\bar{Q}$ production, as has recently been investigated in detail by Mangano et al. [195]. Still, the results of the PCM calculations for pp collisions, indicated in Fig. 65b by the thin curves, agree well with the parton model predictions of [195] for charm and bottom production in pp collisions that fit recent experimental data compilations on inclusive heavy flavor production [196] in hadron collisions. For example, the parton cascade model prediction for the total $c\bar{c}$ cross-section at $\sqrt{s}/A = 200$ (6300) GeV is $\sigma_{c\bar{c}} = 0.63(5.9)$ mb with $m_c = 1.35$ GeV and the structure functions of [127]. This may be compared with the corresponding prediction of Mangano et al. [195] $\sigma_{c\bar{c}} \simeq 0.6(4.0)$ mb. Hence, the charm yield calculated with the parton cascade model for pp collisions is relatively accurate within the uncertainties associated with the exact value of m_c , the choice of the structure function parametrization, etc., so that any additional uncertainty in AA collisions is connected with the extrapolation from pp to AA , i.e. the lack of detailed knowledge about nuclear and dense medium effects that are essential in AA reactions but are irrelevant in pp collisions.

To improve the qualitative understanding of the features associated with ultra-dense, high temperature parton matter produced in the central collision region of heavy ion collisions at collider energies more experimental and theoretical information is necessary regarding the heavy flavor sea excitation and glue decay production at RHIC and LHC energies

in order to draw robust conclusions for the experimentally observable heavy flavor yield. In addition, further understanding of a number of issues on both the perturbative evolution of partons and the non-perturbative mechanisms of hadron formation is required.

7. CONCLUSIONS AND OUTLOOK

With this review I intended to give a comprehensive presentation of a QCD based space-time description of high energy nuclear collisions within the the field theoretic parton picture and semiclassical kinetic theory. The central objective was to develop a consistent dynamical formulation for the time evolution of the parton densities in complete phase-space, thereby providing the possibility to study as a function of real time the many particle aspects and the thermodynamic properties of the parton distributions. The underlying philosophy advocated here was rather pragmatic in the sense that emphasis was put on applicability and phenomenology regarding nuclear collisions at the future colliders RHIC and LHC. It condensed in the development of the PCM as a relativistic kinetic formulation for the space-time evolution of nuclear collisions, in which these reactions are described on the microscopic level of parton cascades as sequences of multiple parton-parton collisions and associated parton emission and absorption processes. Further important aspects included are parton shadowing effects in the initial state, the balance between emissive and absorptive processes, dilated formation of gluon radiation, the Landau-Pomeranchuk-Migdal effect, and the properties of soft gluon interference. In addition, a phenomenological hadronization scheme links the perturbative QCD evolution of the parton distributions together with the non-perturbative description of hadron formation. The hadronization is modelled as a recombination of the final state partons at the end of the perturbative phase to form color singlet clusters, followed by the fragmentation of these clusters into observable hadronic states.

It was demonstrated that the PCM sets a consistent framework to simulate and study the time evolution of ultra-relativistic nuclear collisions in complete phase-space, from the first instant of collision to the final particle yield. Although this approach relies on a number of fundamental assumptions and is far from providing an accurate quantitative description, it has the advantage of a wide applicability, which to demonstrate was one of the purposes of the review. It allows to investigate for example,

- (i) general aspects of multiparticle production in hadron-hadron, hadron-nucleus, and nucleus-nucleus collisions,
- (ii) the statistical thermodynamic behavior of off-shell partons and their multiplicative properties, i.e. the evolution of the parton distributions from initial materialization, through pre-equilibrium, towards plasma formation and beyond;

- (iii) properties of characteristic signatures of the early stage of heavy ion collisions to probe quark gluon plasma formation;
- (iv) features of perturbative QCD in hot and dense matter environment.

Although many other issues can be studied in this framework, it is important to keep in mind that the PCM has its clear limitations and cannot (at least at this point) be expected to describe the features of high energy nuclear collisions equally well. In particular, most of the model predictions for heavy ion collisions are to be understood as *qualitative* features associated with ultra-dense, high temperature parton matter produced in the central collision region, and must be viewed as an optimistic scenario that is subject to considerable uncertainties. However, if the underlying description of the partons' kinetics in these reactions is tendentially correct, then the advent of the heavy ion collider experiments at RHIC and LHC will exhibit truly novel physics that has not been accessible before. Particularly the opportunity to create a deconfined macroscopic plasma of predominantly gluons, with temperatures and densities much larger than estimated in the past, is promising to recreate the QCD phase transition of the early universe in the laboratory and to study QCD in a new form of matter.

I hope I have convinced the reader that the main properties of high energy nuclear collisions can be well described on the basis of the semiclassical QCD parton cascade picture. Of course, much more experimental and theoretical study is required before robust conclusions for experimental observables and fundamental theoretical problems can be drawn. The following (incomplete) list of the main problems and open questions connected to the parton cascade approach (c.f. also discussion in Sec. 4.5), may serve as motivating guideline and as a balance to the general optimistic viewpoint of this review.

a) Initial state: The initial parton distributions in nucleus-nucleus collisions are subject to large uncertainty. Particularly the small x behaviour of the gluon structure functions and the question of gluon shadowing [92] in nuclei are of primary importance, since the amount of soft gluons in the initial state sensitively determines the starting conditions for the evolution and of the general reaction dynamics [197]. These gluons densely populate phase-space and due to the large global semihard cross-section they only need to be slightly "tickled" in order to materialize. Thus the gluon content of the nuclei is a large entropy reservoir that is immediately set free because the quantum coherence is so easily destroyed [81]. Even though the recent measurements at HERA [198] of the nucleon structure functions at very small values of x with an extracted gluon component $xg(x) \propto 1/\sqrt{x}$ provide new information, the question of the corresponding form of the structure functions of nuclei composed of many nucleons requires much further investigation [199]. Moreover,

the measured structure functions, which contain solely the single particle aspect of the parton distributions, do not tell us anything about the spatial distribution of partons, or correlations among them - issues which presumably play an important role for large nuclei.

b) Perturbative QCD evolution: Most urgent on the perturbative sector is the medium modification of the parton interaction amplitudes. This open problem is intimately connected to the choice of the model parameters $p_{\perp cut}$ and μ_0 , used to regularize the divergent perturbative QCD parton-parton cross-sections and the parton emission probabilities, respectively. The rather low values of $p_{\perp cut}$ and μ_0 in the model reproduce well a wide range of pp ($p\bar{p}$) collider data, but lead to a huge multiplicity growth of partons when proceeding to heavy ion reactions. Since the parton-parton cross-sections and emission probabilities depend sensitively on $p_{\perp cut}$ and μ_0 , an increase of these cut-off parameters due to dense medium effects would lead to significantly less particle production. Therefore it is important to investigate the color screening length as well as effective medium induced masses of collective gluon and quark modes as a function of space and time. These quantities could then be used as input parameters for the evaluation of medium dependent scattering cross-sections and parton branching probabilities. This would provide a *natural* cut-off of long range interactions in a deconfined QCD plasma, so that no artificial parameters are required to obtain finite interaction amplitudes in a dense medium.

c) Semiclassical approximation of parton transport: The semiclassical kinetic description of the PCM is of course subject to the well known limitations and approximations of the the binary collision approach underlying Boltzmann-type transport equations [79, 80]. Moreover, the use of single particle distributions implies the neglect of e.g. particle correlations, certain aspects of coherent behavior, or effects of long-range color forces that may be subtly latent even at short distances. Specific QCD properties lead to further (yet unsolved) complexities, which have been ignored here, e.g. to the appearance of non-abelian terms in the collision integrals [200], a non-trivial generalization of the Landau-Pomeranchuk-Migdal effect to QCD [201], and more. Some of these problems could be solved by succeeding to establish a rigorous quantum kinetic theory on the basis of Wigner functions rather than classical phase-space densities. By relating the experimentally accessible parton number densities of the nucleon structure functions with the Wigner functions in full phase-space, a connection between the QCD evolution equations in momentum space and the Boltzmann equation in space-time must be established [51]. The semiclassical PCM approach serves here as a pragmatic first step, but ultimately a space-time description well founded on the principles of quantum field theory is desired. Progress in this direction has recently been made in two ways, one starting from a quantum kinetic formulation of QCD

field theory [8], the other one coming from the Schwinger-Keldysh formulation of initial value problems for the exact many-body Schrödinger equation [9].

d) Hadronization: The details of the confinement mechanism are generally difficult to access due to our current lack of non-perturbative QCD dynamics. Although the parton-hadron duality hypothesis [128] and the idea of pre-confinement [52] have been exploited to describe successfully properties of confinement in e^+e^- annihilation, it remains a rather foggy state of affairs in the context of nuclear reactions. It is clear that the conversion of a quark-gluon plasma into hadrons is a much more complicated process than the hadronization of final state partons in free space (as in e^+e^- annihilation), or dilute systems (as in hadron-hadron collisions). In heavy ion collisions the transition is expected to involve a rather long time scale ($\gtrsim 10$ fm) and proceeds as a complex evolution of expansion and cooling of the plasma, perhaps through a mixed parton-hadron phase, until a purely hadronic phase is reached. Several models have been proposed to describe the hadronization of quark-gluon plasmas (see e.g. Ref. [150] and references therein). The phenomenological cluster hadronization scheme employed in the PCM serves at best as a crude prescription that projects the entropy production of the parton evolution onto the final hadron yield. Although a hydrodynamical description may reproduce the gross macroscopic features of hadronic freeze out, ultimately a fully dynamical and microscopic hadronization scheme needs to be developed. For example, a coalescence of partons depending on the local density and temperature, or a nucleation and hadronic bubble formation scenario [14], or an approach based on string cluster formation [202].

In summary: much remains to be done in the nearest future to obtain a more detailed theoretical picture of high energy nuclear collisions, although I believe, already at this point a clear understanding of the fundamental problems on the basis of perturbative QCD and transport theory has been reached. Foremost, we must learn how to utilize medium effects in a dense QCD plasma to eliminate arbitrary cut-offs, so that truly parameter-free calculations become possible. Almost as importantly, we must develop a microscopic theory for the hadronization of a quark-gluon plasma that is based on QCD, doing away with the presently employed QCD-inspired models that have little predictive power. If these two goals can be met, quantitative ab-initio calculations of ultrarelativistic heavy ion collisions will become feasible by the time when RHIC starts operation.

ACKNOWLEDGEMENTS

I am grateful for always inspiring and encouraging collaboration with Berndt Müller, ever since my time at Duke University. Without his enthusiasm this work would have not been possible.

Also I am deeply indebted to J. I. Kapusta, who was a great collaborator during my period at the University of Minnesota, strongly furthering and promoting the developments of the parton cascade approach.

Furthermore I thank all those numerous colleagues that contributed with interesting and fueling conversations and suggestions during the last years: X.-N. Wang, K. Eskola, L. McLerran, E. Shuryak, H. Satz, and many others.

Finally I would like to acknowledge that most of the numerical simulations were carried out at the North Carolina Super Computer Center, and the Minnesota Super Computer Institute, which both provided generous amounts of computer time.

APPENDIX: THE MONTE CARLO PROCEDURE

In the following I will outline the numerical method used in the PCM to solve the transport equation (90), with the collision kernel (91), starting from the initial parton distributions given by eq. (79). For a general overview of Monte Carlo techniques to solve particle transport equations, see e.g. [79, 203, 204]. The PCM Monte Carlo program simulates events of ultra-relativistic hadron-hadron, hadron-nucleus, or nucleus-nucleus collisions on the computer, on the basis of the model framework of Chap. 4. It allows follow the evolution of the parton phase space distributions in real time during the course of a collision in accordance with the concept of independent parton-parton scatterings and connected emission and absorption processes. The Fortran computer program is obtainable from the author.

The probabilistic formulation of parton cascade development (Secs. 2 and 4) provides a suitable basis for a simulation of Markov processes, relying on the property of locality (no memory of the prehistory), because the evolution of parton cascades during a nuclear collision has the structure of independent sequences of elementary interaction processes in between which the partons propagate along classical trajectories. Thus, the path of each parton through space-time has a "zig-zag" shape dictated by their vertices of interaction, and from each vertex a branch extends that incorporates the tree-like structure of further parton interactions.

It is important to realize that the Monte-Carlo procedure is applied on two different levels. First, the propagation of test particles, which are statistically sampled from the

original parton distributions, along classical paths, provides a faithful solution of the transport equation (90) without collision terms. Including the collisions terms, that is allowing for randomly sampled $2 \rightarrow 2$ scatterings and $2 \rightarrow 1$ fusion processes, together with associated $1 \rightarrow 2$ branchings, yields a solution of the full transport equation, when the sampling procedure is repeated sufficiently often to obtain the desired accuracy. In the limit $N \rightarrow \infty$ of N test particles per event, each interacting with a strong coupling constant α_s/N , would give the exact solution of the transport equation. Second, the specific QCD structure of the individual parton interactions is itself sampled by Monte Carlo techniques. As explained in Sec. 4, each cascade sequence has the structure of a Markov process [206], with the interaction probabilities determined by the squared matrix elements that include all leading real and virtual QCD corrections and effectively include the dominant interference terms in a local description. Consequently, the simulation of a nuclear collision, which follows the ensemble of partons in space and time as they cascade and interact through elementary processes, gives the solution to the transport equation.

The simulation of the time-evolution of the parton distributions according to the transport equation [$p \equiv p^\mu = (E, \vec{p})$, $r \equiv r^\mu = (t, \vec{r})$]

$$p^\mu \partial_\mu F_a(p, r) = \left[E \frac{\partial}{\partial t} - \vec{p} \cdot \frac{\partial}{\partial \vec{r}} \right] F_a(p, r) = \sum_{\text{processes } k} I_a^{(k)}(p, r) \quad (A1)$$

is based on the stochastical sampling of the distribution functions $F_a(p, r)$ in coarse grained phase-space, discretized into cells labeled by an index j . The number of partons of type a , $n_a(j)$, in each cell is found at some fixed time in the total center-of-mass frame of the colliding nuclei. Statistics is improved by summing $n_a(j)$ over several collision events of the parton cascade simulation. As explained in Sec. 3.4 the distribution functions $F_a(r, p)$ can be estimated as

$$F_a(p, r) \longrightarrow F_a(j, t) = \frac{1}{\nu_{ev}} \left(\frac{n_a(j, t)}{\Delta \Gamma_a(j)} \right), \quad (A2)$$

where ν_{ev} is the number of collision events accumulated and the phase-space volume of cell j is

$$\Delta \Gamma_a(j) = \gamma_a \int_{\text{cell } j} \frac{d^3 r d^3 p}{(2\pi)^3} = \gamma_a \frac{(\Delta^3 r)_j (\Delta^3 p)_j}{(2\pi)^3} \quad (A3),$$

with $\gamma_g = 16$ and $\gamma_q = \gamma_{\bar{q}} = 6$.

The time evolution of the multi-parton system in phase-space starts from the initial parton distributions of the incoming nuclei. The two nuclear parton clouds approach each other from their initial separation and collide. The evolution of the collision divided into

small discrete time steps ⁵

$$\Delta t = 0.01 \text{ fm} , \quad (A4)$$

chosen such that the structure of parton interaction is completely resolved. In every time step the flavor, 4-momentum and position of each parton is stored, together with the color flow information that connects the parton with its color siblings. These data are continuously updated after each a time step, accounting for the parton interactions that occur. Thus, the change of the parton distributions between time steps t_{i-1} and $t_i = t_{i-1} + \Delta t$ in a given phase-space cell $\Delta\Gamma_a(j)$ with $\vec{r} \in [\vec{r}_j \pm (\Delta r)_j/2]$ and $\vec{p} \in [\vec{p}_j \pm (\Delta p)_j/2]$ corresponding to (A1) is given by

$$F_a(j, t_i) = F_a(j, t_{i-1}) + \sum_{\text{processes } k} E I_a^{(k)}(j, t_{i-1}, t_i) , \quad (A5)$$

i.e. by summing over all partons of type a with momenta and coordinates $(\vec{p}, \vec{r}) \in \Gamma_a(j)$, as they stream freely during Δt , or interact with another according to the space-time dependent collision kernel $\sum_k I_a^{(k)}(j, t_{i-1}, t_i)$. This is the general procedure to simulate the solution of the transport equation. How the specific structure of parton cascades embedded in the collision kernel, is simulated in detail, will be explained in the following.

A.1 The initial state

The initial parton configurations in phase space are obtained from the distribution function (79) for each individual nucleon N_i where the index i numbers the nucleons, being protons or neutrons. The frame of reference is the CM frame of the nuclei, with the kinematics and the notation employed in Sec. 4.

(i) Flavor and momentum distribution of partons

Consider as in Sec. 4 a nucleus-nucleus system of colliding nuclei A and B with the total number of nucleons $N_{nuc} \equiv N_{nuc}^{(A)} + N_{nuc}^{(B)} = \sum_i N_i$. (In case of hadron-hadron or hadron-nucleus collisions one, or both nuclei are simply replaced by a hadron.) For each nucleon N_i individually the flavors and momenta of the confined partons are sampled from the nucleon structure functions $F_a^{N_i}(x, Q^2)$ at the initial resolution scale $Q^2 = Q_0^2$, eq. (78). First the three valence quarks are assigned longitudinal momentum fractions $x = p_z/P$,

⁵Taking $\Delta t = 10^{-2}$ fm implies that the outgoing states c and d in a parton scattering $ab \rightarrow cd$ are not asymptotic states after such a short time, in the sense as used in the parton-parton cross-sections. The question is, how long after a scattering does it take until a parton can be considered an approximate asymptotic state. This is currently described in the PCM in a crude way [82] by taking into account for each interaction the characteristic time scale, given by the uncertainty relation, for the production of the outgoing particles as truly resolvable quanta.

generated according to the proton or neutron structure functions F^{Ni} for the corresponding valence quark configuration uud or udd ,

$$F_v(x, Q_0^2) = F_a^{Ni}(x, Q_0^2) \quad (a = u_v, d_v) . \quad (A6)$$

Next, for the gluons and sea quarks again an individual x -value is sampled, but now from the summed distribution

$$F_{g,s}(x, Q_0^2) = F_g(x, Q_0^2) + \sum_{q=u_s, d_s, s, c, \dots}^{n_f(Q_0^2)} \left[F_q(x, Q_0^2) + F_{\bar{q}}(x, Q_0^2) \right] , \quad (A7)$$

excluding the valence quarks. Here $n_f(Q_0^2)$ is the maximum number of flavors that can be probed at Q_0^2 with $2m_q < Q_0$. The particular flavor $b \equiv g, u_s, d_s, s, \dots$ is then selected from the relative probabilities $F_b/F_{g,s}$. This procedure is repeated until the total nucleon momentum P is exhausted, i.e.

$$\sum_{\kappa}^{n^{Ni}} x_{\kappa} = \sum_{\kappa}^{n^{Ni}} \frac{(p_z)_{\kappa}}{P} = 1 , \quad (A8)$$

[c.f. (82)] from which the effective number of partons in the nucleon $n^{Ni}(Q_0^2, P)$ (84) is obtained.

The nuclear shadowing corrections at small x to the nucleon structure functions in a nucleus as compared to a free nucleon, are then simply incorporated by multiplying F_a^{Ni} with the corresponding shadowing factor $\rho_a^A(x)$, eq. (86).

The transverse momentum $\vec{p}_{\perp}$ of each parton is assigned independent of its flavor according to the Gaussian distribution in (79) with a width fixed by the constraint that the invariant mass of each nucleon's parton composition equals the nucleon mass.

Calculating the partons initial virtualities $q^2 < 0$ and energies $E = \sqrt{p_z^2 + p_{\perp}^2 + m_0^2} + q^2$ completes the construction of the distribution of flavors a and four-momenta p of partons for a given nucleon. The procedure is repeated for all remaining nucleons in the nuclear collision system $A + B$.

Note, that the scale Q_0^2 at which the structure functions are evaluated is initially chosen as a small value ($Q_0 = 1$ GeV), which is continuously updated from the accumulated values of Q^2 , eq. (78), the scales of the primary parton-parton scatterings. In this way it is ensured that the initial parton distributions are sampled at the most appropriate average value of Q^2 (c.f. Tables 1 and 3). It is not necessary to sample the structure functions at the *precise* value of Q^2 , because the initial state radiation allows one to incorporate the Q^2 -evolution of space-like cascades according to the concept of Sec. 4.2.3.

(ii) Spatial distribution of partons

The spatial distribution of partons in a nucleon, eq. (87), is generated as follows. First, the initial separation of the incoming nuclei A and B in the center of mass frame is specified. Then the nucleons are assigned positions randomly throughout the volume of A , respectively B . Next, the partons coordinates are uniformly distributed within a sphere of $R_N^{m,s} = 0.81 \text{ fm}$ around the nucleons positions. Finally the positions of nucleons and their valence quarks are boosted into the center of mass frame occupying a highly contracted region $(\Delta z)_v = 2R_A M_N / P$, whereas the gluons and sea quarks are smeared out over a spatially extended region $(\Delta z)_{g,s} \approx 2R_A$ around the valence quarks. This procedure establishes the property of distributed Lorentz contraction for the partons discussed in Sec. 4.1.1 and yields the fuzzy pancake shape of the incoming nuclei as illustrated in Fig. 19.

(iii) Color structure The last step in the initialization procedure is to provide the initial color flow information. Its purpose is twofold: first, one needs to require the partons of each nucleon to form a color singlet, and second, it allows to keep track of the color flow [30, 69] in parton interactions. Initially, all partons in each nucleon are sequentially connected with color lines to yield a color singlet configuration. This is done such that the gluons and sea quarks stretch color lines between one valence quark on the one end and the remaining two as a diquark on the other end. Each color charge is assigned a value C corresponding to its color multiplet: a quark (antiquark) has $C = +1(-1)$ (triplet (antitriplet)), a gluon has $C = +2$ or $C = -2$ (octet or antioctet) and a diquark (antidiquark) has $C = -1(+1)$ (antitriplet (triplet)). A color singlet, like a nucleon, has $C = 0$. The value of C also specifies the number and direction of color lines from or to a color charge, thus defining the color flow. For example, a quark with $C = +1$ has one outgoing color line, whereas a diquark with $C = -1$ has one incoming color line. A gluon always has two color lines ($C = \pm 2$), one coming in and one going out of the charge. Subject to the requirement that a nucleon is a color singlet ($C = 0$) so that the sum of its partons color values must vanish, the color connection scheme starts with one of the valence quarks (its color line going out) and connects the sea-quark pairs and gluons sequentially along their color lines until the two remaining valence quarks, represented as a diquark, form the end of the color line.

A.2 The space-time development of parton cascades

As mentioned above, once the initialization is complete, the space-time evolution of the collision is now followed in discrete time steps Δt as the partons cascade through coarse grained phase-space subject to $2 \rightarrow 2$ scatterings, $2 \rightarrow 1$ fusions, and $1 \rightarrow 2$ branchings.

(i) Parton-parton scattering processes

The partons propagate along classical trajectories until they scatter, i.e. the path of a parton is given by

$$\vec{r}(t_i) = \vec{r}(t_{i-1}) + \frac{\vec{p}}{E} \Delta t . \quad (A9)$$

To account for mutual interactions, consider a certain phase-space cell. For all partons in this cell during $\Delta t = t_i - t_{i-1}$ it has to be checked if they may scatter during this time step, or if they continue to propagate freely. To determine whether a given parton a will collide with some other parton b within a time step Δt , the total cross-section $\hat{\sigma}_{ab}$ for each possible pair (a, b) is calculated to define an effective area of interaction. This cross-section is evaluated from eq. (97),⁶

$$\begin{aligned} \hat{\sigma}_{ab}(\hat{s}, p_{\perp cut}) &= \sum_{c,d} \left\{ \int_0^{p_{\perp cut}^2} dp_{\perp}^2 \left(\frac{d\hat{\sigma}_{ab \rightarrow cd}^{soft}}{dp_{\perp}^2} \right) + \int_{p_{\perp cut}^2}^{\infty} dp_{\perp}^2 \left(\frac{d\hat{\sigma}_{ab \rightarrow cd}^{hard}}{dp_{\perp}^2} \right) \right\} \\ &\equiv \sum_{c,d} \hat{\sigma}_{ab \rightarrow cd}(\hat{s}, p_{\perp cut}) , \end{aligned} \quad (A10)$$

where the hard and soft elementary cross-sections are specified by (99) and (100). As before, $p_{\perp}^2$ is the relative transverse momentum of the scattering partons, defining the interaction scale $Q^2 \equiv p_{\perp}^2$, and $\hat{s} = (p_a + p_b)^2$, $\hat{t} = (p_a - p_c)^2$, $\hat{u} = (p_a - p_d)^2$ are the Mandelstam variables of the two-parton subsystem, with $\hat{s} + \hat{t} + \hat{u} = m_a^2 + m_b^2 + m_c^2 + m_d^2$, and the sum runs over all possible final states c, d .

The characteristic scale of a scattering $ab \rightarrow cd$, provided it occurs, is given by the specific value of $Q^2(\hat{s}, \hat{t}, \hat{u}) \equiv p_{\perp}^2$ as determined from the differential cross-section for a chosen scattering subprocess (explained below). However, since $\hat{t}$ and $\hat{u}$ and thus $p_{\perp}^2 \approx \hat{t}\hat{u}/\hat{s}$ are not known at this point, the approximate average relation $\overline{Q^2} = \overline{p_{\perp}^2} \approx \hat{s}/3$ is employed here, first for deciding if a hard (soft) scattering occurs $\overline{Q^2} > p_{\perp cut}^2$ ($\overline{Q^2} < p_{\perp cut}^2$), and second, as the argument of the running coupling $\alpha_s(Q^2)$ that enters the matrix elements from which the integrand $d\hat{\sigma}/dp_{\perp}^2$ is obtained.

For each pair of partons (a, b) the cross-section (A10) is used to define an effective area of interaction with radius

$$\hat{b}_{ab}^{max}(p_{\perp cut}^2) = \sqrt{\frac{\hat{\sigma}_{ab}(\hat{s}, p_{\perp cut}^2)}{\pi}} \quad (A11)$$

⁶Recall from Sec. 4.2.2 that the invariant cut-off scale $Q_{cut}^2 \equiv p_{\perp cut}^2$ is introduced such that collisions with $p_{\perp} \geq p_{\perp cut}$ are treated as (semi)hard scatterings within QCD perturbation theory, whereas for $p_{\perp} < p_{\perp cut}$ a soft interaction is assumed to occur.

which sets the maximum allowed impact parameter for the parton pair under consideration. Thus, in a given time interval Δt the two partons can only collide if the distance d_{ab} between them satisfies

$$d_{ab} \leq d_{ab}^{min} = \sqrt{(\hat{b}_{ab}^{max})^2 + (\Delta t)^2} . \quad (A12)$$

If this is the case, it now has to be checked, (a) whether the partons a and b will pass their point of closest approach, and, (b) if this distance of closest approach is smaller than $\hat{b}_{ab}^{max}$. In the center-of-mass frame of the partons a and b , their momenta are $\vec{p}_{cm}$ and $-\vec{p}_{cm}$, respectively, and the distance $\Delta\vec{r} = \vec{r}_a - \vec{r}_b$ in the lab frame becomes in the center-of-mass frame $\Delta\vec{r}_{cm} = (\hat{\gamma} - 1)(\Delta\vec{r} \cdot \vec{\beta}/\beta)\vec{\beta}/\beta + \Delta\vec{r}$, where $\hat{\gamma} = (1 - \beta^2)^{-1/2}$ and $\vec{\beta}$ is the velocity of the center-of-mass of a and b . Thus the two partons will collide within the time interval $\Delta t/2$ and $+\Delta t/2$, if (a)

$$\left| \frac{\Delta\vec{r}_{cm} \cdot \vec{p}_{cm}}{p_{cm}} \right| < \left(\frac{p_{cm}}{E_{1cm}} + \frac{p_{cm}}{E_{2cm}} \right) \frac{\Delta t}{2} , \quad (A13)$$

and if (b) ⁷

$$\hat{b}_{ab} = \sqrt{(\Delta\vec{r}_{cm})^2 - \left(\frac{\Delta\vec{r}_{cm} \cdot \vec{p}_{cm}}{p_{cm}} \right)^2} < \hat{b}_{ab}^{max} . \quad (A14)$$

The maximum allowed impact parameter (A11) depends on $p_{\perp cut}$, however, this sensitivity can be overcome [154] by implementing a dynamical screening of the divergent long wavelength contributions to the interaction probability: All possible collision partners b are considered equal candidates for the actual scattering pair a and b . In case there is more than one collision partner for a , the one with the minimum impact parameter

$$\hat{b}_{ab}^{min} = \min_{b=1,\dots,n} [\hat{b}_{ab}] \quad (A15)$$

is chosen, where n is the number of potential candidates b that satisfy (A13) and (A14). Hence, the parameter $p_{\perp cut}$ is replaced by the dynamical natural cut-off

$$(p_{\perp 0})_{ab} \equiv \frac{1}{\hat{b}_{ab}^{min}} > p_{\perp cut} . \quad (A16)$$

In other words, the long range part of the Coulomb type color field is screened below $(p_{\perp 0})_{ab}$, and each collision pair has its individual value of $(p_{\perp 0})_{ab}$, depending on the local density of surrounding partons.

⁷Actually a more general procedure is realized in the PCM by employing a normalized probability distribution $\Phi(b/b^{max})$ which is a less rigid cut-off than $\Phi(b/b^{max}) = \Theta(1 - b/b^{max})$ which corresponds to eq. (23). A rather diffuse distribution appears to be more appropriate, since the cross-section $\hat{\sigma}_{ab}$ is generally dominated by small-angle scatterings, corresponding to large impact parameters. The particular choice employed is $\Phi(b/b^{max}) = \exp(-b/b^{max})$ which requires, instead of eq. (23), $b < b^{max} \ln(R)$, where R is a uniform random number between 0 and 1.

Once for a parton a the collision partner b is found, then the subprocess is determined. For instance, in the case of two incoming gluons or an incoming quark and antiquark of alike flavor, a number of elementary reactions are possible, $gg \rightarrow gg$, $u\bar{u}$, $d\bar{d}$, ... or $q\bar{q} \rightarrow gg$, $u\bar{u}$, $d\bar{d}$, ..., respectively. Thus the particular subprocess, i.e. the flavors of the outgoing partons c and d , is selected according to the relative magnitude of the cross sections $\hat{\sigma}_{ab \rightarrow cd}/\hat{\sigma}_{ab}$, given by eq. (A10).

Next the scattering angle $\cos \hat{\theta}_{CM} = 1 + 2\hat{s}\hat{t}/[(\hat{s} - (m_a + m_b)^2)(\hat{s} - (m_a - m_b)^2)]$ for the chosen subprocess $ab \rightarrow cd$ is determined by generating the value of $p_\perp^2$ according to the probability distribution

$$\Xi(p_\perp^2) = \frac{1}{\hat{\sigma}_{ab \rightarrow cd}} \int_{(p_{\perp 0}^2)_{ab}}^{p_\perp^2} dp_\perp'^2 \left(\frac{d\hat{\sigma}_{ab \rightarrow cd}}{dp_\perp'^2} \right). \quad (\text{A17})$$

Note that the small-angle scattering is bounded by $\cos \hat{\theta}_{cm} > 1 - 2(p_{\perp 0}^2)_{ab}/\hat{s}$, which is set by the dynamics of the parton system itself and not by an 'external' parameter.

Finally the four-momenta of the outgoing partons and the color flow of the scattering are constructed from standard kinematics (c.f. [126]), and the color flow implied by the specific subprocess is assigned chosen as discussed in detail in Refs. [30, 69].

(ii) Space-like branchings

Recall, that in case one or both of the scattering partons a and b originates as a primary quantum from the initial nuclear parton clouds, the possibility is taken into account that the parton, say a , has participated in a number of space-like branchings before the scattering. As explained in Sec. 4.2.3, the evolution of the initial parton's small space-like virtuality $-q_0^2 \approx Q_0^2$ up to $-q_a^2 \simeq Q_{ab}^2$, the scale of the scattering vertex in the reaction $ab \rightarrow cd$ (see Fig.21), is represented by the Sudakov form-factor for space-like branchings:

$$S_{a_n}(x_{a_n}, \tilde{Q}^2, Q_0^2) = \exp \left\{ - \sum_{a_{n-1}} \int_{Q_0^2}^{\tilde{Q}^2} \int_{\tilde{z}_-(\tilde{Q}')}^{\tilde{z}_+(\tilde{Q}')} d\mathcal{P}_{a_n a_{n-1}}^{(S)}(x_{a_n}, \tilde{z}, \tilde{Q}'^2) \right\}, \quad (\text{A18})$$

where $a \equiv a_n$ and a similar expression holds for parton b . These form factors give the probabilities $P_{a,b}^{(S)} = (1 - S_{a,b})$ that the parton a , respectively parton b , has evolved from the initial point Q_0^2 toward $\tilde{Q}^2 = Q_{ab}^2$ by increasing its virtuality through a sequence of single branching processes. In principle this evolution in its virtuality $\tilde{Q}^2 = |q^2|$ could be carried out starting from the initial $|q_0^2| \approx Q_0^2$ and working upwards to Q_{ab}^2 . This method of evolving the parton virtualities is referred to as *forward evolution scheme* [32, 35]. However there are several problems associated with this approach due to the fact that it is impossible to handle the kinematics of the evolution exactly, since the actual virtualities of the partons cannot be found until after the scattering has occurred. Instead, a *backward evolution*

scheme [30, 31, 34] is used, which is described by the form factor $S_a(x_a, \tilde{Q}^2, Q_0^2)$ with the already determined $\tilde{Q}^2$ of the scattering forming the starting point. The evolution is then reconstructed step by step in falling $\tilde{Q}^2$ sequence back towards the initial $|q_0^2| \approx Q_0^2$.

The algorithm used to generate this backward evolution is based on the one of Bengtsson, Sjöstrand and Zijl [66]. In this approach, a knowledge of $S_a(x_a, \tilde{Q}^2, Q_0^2)$ is sufficient to reconstruct the kinematics. At the branching $a_{n-1} \rightarrow a_n + k_n$ (see Fig. 21) before the scattering that involves the parton $a = a_n$, three quantities have to be determined: the $\tilde{Q}^2$ value of the branching (which also defines the virtuality $q^2 = -\tilde{Q}^2$ of the parton a_n), the flavor of parton a_{n-1} and the splitting variable $\tilde{z} = x_{a_n}/x_{a_{n-1}}$. This information is extracted as follows.

The spacelike branching is sampled from the corresponding branching probability distribution $p_{a_n}^{(S)}(x_{a_n}, \tilde{Q}^2, \tilde{Q}_{max}^2)$ that the parton a_n actually has emerged from the branching $a_{n-1} \rightarrow a_n + k_n$ between $\tilde{Q}^2$ and $\tilde{Q}^2 + d\tilde{Q}^2$,

$$p_{a_n}^{(S)}(x_{a_n}, \tilde{Q}_{max}^2, \tilde{Q}^2) d\tilde{Q}^2 = S_{a_n}(x_{a_n}, \tilde{Q}_{max}^2, \tilde{Q}_{min}^2) \frac{d}{d\tilde{Q}^2} \left(\frac{1}{S_{a_n}(x_{a_n}, \tilde{Q}^2, \tilde{Q}_{min}^2)} \right) d\tilde{Q}^2, \quad (A19)$$

where $\tilde{Q}_{max}^2$ specifies the maximum virtuality of parton a_n which for $a_n = a$ is set by the scale of the scattering vertex Q_{ab}^2 , and $\tilde{Q}_{min}^2 = Q_0^2$ is the lower bound given by the initial resolution scale (78). Solving the equation

$$S_{a_n}(x_{a_n}, \tilde{Q}^2, Q_0^2) = \frac{S_{a_n}(x_{a_n}, \tilde{Q}_{max}^2, Q_0^2)}{R}, \quad (A20)$$

where R is a uniformly distributed random number in $[0, 1]$ determines $\tilde{Q}^2 = |q^2|$. The case that no branching occurs at all, corresponds to $R < S_{a_n}(x_{a_n}, \tilde{Q}^2, Q_0^2)$.

Next, if the splitting occurs the relative probability for the different allowed branching processes $a_{n-1} \rightarrow a_n + k_n$ at given $\tilde{Q}^2$ are given by the integrals

$$\int dz \gamma_{a_{n-1} \rightarrow a_n k_n}(z) \frac{x_{a_{n-1}} F^N(x_{a_{n-1}}, \tilde{Q}^2)}{x_{a_n} F^N(x_{a_n}, \tilde{Q}^2)}, \quad (A21)$$

with the branching functions $\gamma_{a_{n-1} \rightarrow a_n k_n}(z)$ given in eq. (16).

Finally, with $\tilde{Q}^2$ and a_{n-1} known, the probability distribution in $\tilde{z} = x_{a_n}/x_{a_{n-1}}$ is given by the integrand of eq. (A21). The kinematics of the branching can then be uniquely reconstructed and the procedure is repeated in backward direction, by replacing $a_n \rightarrow a_{n-1}$, $x_{a_n} \rightarrow x_{a_{n-1}}$, $\tilde{Q}_{max}^2 = Q_{ab}^2 \rightarrow \tilde{Q}_{max}^2 = \tilde{Q}^2$, and so forth, until the selected virtuality reaches the scale Q_0^2 and the branching stops.

The reconstruction of the branching kinematics is non-trivial. It depends e.g. on the choices of the interpretation of z and of the functional dependence of the evolution variable.

Furthermore the parton k_n in the branching $a_{n-1} \rightarrow a_n + k_n$ is time-like and can itself initiate a sequence of time-like branchings. A detailed discussion of these topics and their physical relevance can be found in e.g. Refs. [66].

(iii) Time-like branchings

The evolution of time-like branchings, i.e. the phenomenon that after the scattering $ab \rightarrow cd$ the outgoing partons c and d may have time-like virtualities $m_{c,d}^2 > 0$ which are successively decreased from $m_{c,d}^2 \simeq Q_{ab}^2$ down to the cutoff μ_0^2 by emitting a number of (also time-like) partons. As explained in Sec. 2.4.4, the probabilities for such subsequent single branchings are determined by the Sudakov form-factor for time-like branchings,

$$T_c(\xi, \xi_{min}) = \exp \left\{ - \int_{\xi_{min}}^{\xi} \sum_f \int_{z-(\xi', E_c)}^{z+(\xi', E_c)} d\mathcal{P}_{cf}^{(T)}(\xi', z) \right\} , \quad (A22)$$

which gives the total branching probability $P_{c,d}^{(T)} = 1 - T_{c,d}$ for partons c and d . The Monte Carlo algorithm in the program that generates such sequences of time-like branchings follows the procedure of Bengtsson and Sjöstrand [70]. Similar to the previous case of space-like branchings, here the form factors $T_{c,d}(Q^2, \mu_0^2)$ form the basis to extract the characteristic quantities of a branching $c \rightarrow fg$ (see Fig. 21), namely, the mass m^2 which defines the point of the branching and determines the virtuality of parton c , (starting from $m^2 \simeq Q_{ab}^2$), the flavors of the daughters f and g and finally the splitting variable $z = E_f/E_c$. The generation procedure works as follows.

From (A22) it follows that the probability distribution $p_c^{(T)}(\xi_{max}, \xi)$ for parton c to actually branch into partons f and g between ξ and $\xi + d\xi$ with $\xi < \xi_{max}$ is given by:

$$p_c^{(T)}(\xi_{max}, \xi) d\xi = T_c(\xi_{max}, \xi_{min}) \frac{d}{d\xi} \left(\frac{1}{T_c(\xi, \xi_{min})} \right) d\xi . \quad (A23)$$

Here ξ_{max} is the value of the angular variable of the vertex at which the parton c was produced and is according to eq. (107) determined by the four-momenta emerging from the vertex. For example, in the case of parton c being produced by the scattering $ab \rightarrow cd$ (as in Fig. 21), one has $\xi = (p_c \cdot p_d)/(E_c E_d)$. The branching is then generated from the distribution $p_c^{(T)}$ by solving for ξ :

$$T_c(\xi, \xi_{min}) = \frac{T_c(\xi_{max}, \xi_{min})}{R} , \quad (A24)$$

with R a random number in $[0,1]$, from which the virtuality M^2 of parton c is obtained. If the virtuality m^2 is less than the cut-off μ_0^2 then the branching does not occur.

If the branching occurs, the second step is to select a branching channel, i.e. the flavor of the daughters f and g from the relative probabilities, given by the z -integrals of the

branching functions $\gamma_{c \rightarrow fg}(z)$,

$$\int dz \gamma_{c \rightarrow c_1 c_2}(z) . \quad (A25)$$

With $m_c^2 = m^2$ and the flavors of f, g determined, the value of the splitting variable z is generated according to the probability distribution given by $\gamma_{c \rightarrow fg}(z)dz$.

Finally, the four-momenta $p_{f,g}^\mu$ and virtualities $m_{f,g}^2 = p_{f,g}^2$ of the daughter partons f and g are then constructed from the chosen values of ξ and z as in Ref. [70]. The partons f and g are subsequently allowed to branch in exactly the same way unless, as stated before, they rescatter or are being absorbed, in which cases the respective branch of the cascade terminates. Furthermore, a perturbative branching process within an individual cascade is forcibly terminated when the virtualities of the partons at a branching vertex fall below the infrared cut-off μ_0 . In this case these partons are put on mass shell with effective masses given by eq. (111) and propagate freely unless they encounter a further collision in which enough energy is transferred to trigger a new cascade.

References

- [1] For a recent overviews, see:
Quark Matter '91, Proceedings of the Ninth International Conference on Ultra-relativistic Nucleus-Nucleus Collisions, Gatlinburg, Tennessee, 1991, edited by T. C. Awes *et al.* [Nucl. Phys. **A544**, 1c (1992)];
Quark Matter '93, Proceedings of the Tenth International Conference on Ultra-relativistic Nucleus-Nucleus Collisions, Borlänge, Sweden, 1993, edited by H. A. Gustafsson *et al.* [Nucl. Phys. **A566**, 1c (1994)].
- [2] B. Müller, *Relativistic heavy ion collisions: "re-creating" the early universe in the laboratory*, in the proceedings of the *Workshop on Pre-equilibrium Parton Dynamics in Heavy Ion Collisions*, edited by X. N. Wang, LBL-Report 34831, Berkeley 1993.
- [3] B. Müller, *The physics of the quark gluon plasma*, Lecture Notes in Physics, Vol. 225 (Springer, Berlin-Heidelberg 1985).
- [4] L. McLerran, Rev. Mod. Phys. **58**, 1020 (1986).
- [5] *Quark Gluon Plasma*, edited by R. C. Hwa (World Scientific, Singapore 1990).
- [6] K. Geiger, Nucl. Phys. **A 566**, 257c (1994).
- [7] H.-T. Elze and U. Heinz, Phys. Rep. **183**, 81 (1989) and references therein;
 S. Mrówczyński and U. Heinz, Ann. Phys. **229**, 1 (1993); *ibid.* **229**, 1 (1994).
- [8] J.-P. Blaizot and E. Iancu, Nucl. Phys. **390**, 589 (1993); Phys. Rev. Lett. **70**, 3376 (1993); preprint SPhT/93-064, Saclay 1993 (to appear in Nucl. Phys. **B**).
- [9] Y. Kluger, J. M. Eisenberg, B. Svetitsky, F. Cooper and E. Mottola, Phys. Rev. Lett. **67**, 2427 (1991); Phys. Rev. **D45**, 4659 (1992); *ibid.* **D48**, 190 (1993).
- [10] L. D. Landau, Izv. Akad. Naul. SSSR, Ser. Fiz. **17**, 51 (1953).
- [11] M. Gyulassy and T. Matsui, Phys. Rev. D **29**, 419 (1984);
 K. Kajantie, R. Raitio, and P. V. Ruuskanen, Nucl. Phys. **B222**, 152 (1983);
 G. Baym, B. L. Fryman, J.-P. Blaizot, M. Soyeur, and W. Czyz, Nucl. Phys. **A407**, 541 (1983);
 H. Gersdorff, L. D. McLerran, M. Kataja, and P. V. Ruuskanen, Phys. Rev. D **34**, 794 (1986);
 N. S. Amelin, E. F. Straubo, L. P. Csernai, V. D. Toneev, K. K. Gudima and D. D. Strottman, Phys. Rev. Lett. **67**, 1523 (1991).

- [12] E. V. Shuryak, Phys. Rev. Lett. **68**, 3270 (1992).
- [13] T. S. Biro, E. van Dorn, M. H. Thoma, B. Müller, and X.-N. Wang, Phys. Rev. **C48**, 1275 (1993).
- [14] L. Csernai and J. I. Kapusta, Phys. Rev. **D46**, 1379 (1992); Phys. Rev. Lett. **69**, 737 (1992);
L. Csernai, G. Kluge, E. E. Zabrodin, and J. I. Kapusta, Z. Phys. **C58**, 453 (1993).
- [15] B. Andersson, G. Gustafson, G. Ingelman and T. Sjöstrand, Phys. Rep. **97**, 33 (1983).
- [16] B. Andersson, G. Gustafson and B. Söderberg, Nucl. Phys. **B264**, 29 (1986).
- [17] L. V. Gribov, E. M. Levin, and M. G. Ryskin, Phys. Rep. **100**, 1 (1983).
- [18] E. M. Levin and M. G. Ryskin, Phys. Rep. **189**, 267 (1990).
- [19] See e.g.: Proceedings of the *Workshop on Pre-equilibrium Parton Dynamics in Heavy Ion Collisions*, edited by X. N. Wang, LBL-Report 34831, Berkeley 1993.
- [20] K. Kajantie, P. V. Landshoff, and J. Lindfors, Phys. Rev. Lett. **59**, 2527 (1987);
K. J. Eskola, K. Kajantie, and J. Lindfors, Nucl. Phys. **B323**, 37 (1989).
- [21] X.-N. Wang, Phys. Rev. **D43**, 104 (1990);
X.-N. Wang and M. Gyulassy, Phys. Rev. Lett. **68**, 1480 (1992).
- [22] A. Selikov and M. Gyulassy, Phys. Lett. **B316**, 373 (1993).
- [23] B. Andersson, G. Gustafson and B. Nilsson-Almqvist, Nucl. Phys. **B281**, 289 (1987)
B. Nilsson-Almqvist and E. Stenlund, Computer Phys. Comm. **43** (1987), 387; B. Lörstad, Int. J. Mod. Phys. **A12**, 2861 (1989).
- [24] M. Gyulassy, in *Proceedings of the Eighth Balaton Conference on Nuclear Physics*, edited by Z. Fodor (KFKI, Budapest, 1987); CERN Report No. CERN-TH-4794/87, 1987 (unpublished).
- [25] K. Werner, Z. Phys. **C42**, 85 (1989); Phys. Rep. **232**, 87 (1993).
- [26] H. Sorge, H. Stöcker, and W. Greiner, Nucl. Phys. **A498**, 567c (1989); Ann. Phys. **192**, 266 (1989).

- [27] N. S. Amelin, K. K. Gudima and V. D. Toneev, *Yad. Fiz.* **51**, 512 (1990); N. S. Amelin, E. F. Staubo, L. P. Csernai, V. D. Toneev, K. K. Gudima and D. Strottman, *Phys. Rev. Lett.* **67**, 1523 (1991).
- [28] D. J. Dean, A. S. Umar, J. S. Wu and M. R. Strayer, *Nucl. Phys.* **A544**, 475c (1992).
- [29] T. Csörgö, J. Zimanyi, J. Bondorf and H. Heiselberg, *Phys. Lett.* **222**, 115 (1989).
- [30] H. U. Bengtsson, *Comp. Phys. Com.* **31**, 323 (1984);
T. Sjöstrand, *Comp. Phys. Com.* **28**, 227 (1983); preprint CERN-TH. 7111/93, 1994
(to appear in *Computer Phys. Comm.*).
- [31] G. Marchesini, B. R. Webber, G. Abbiendi, I. G. Knowles, M. H. Seymour and L. Stanco, *Comp. Phys. Com.* **67**, 465 (1992).
- [32] R. Odrico, *Comp. Phys. Com.* **32**, 139 (1984).
- [33] A. Ali, B. van Eijk and I. ten Have, *Nucl. Phys.* **B292**, 1 (1987).
- [34] R. D. Field and S. Wolfram, *Nucl. Phys.* **B213**, 65 (1983);
T. D. Gottschalk, *Nucl. Phys.* **B214**, 201 (1983); *Nucl. Phys.* **B227**, 413 (1983).
- [35] F. E. Paige and S. D. Protopopescu, in the Proceedings of the summer study of the Physics of the SSC, Snowmass, 1986, edited by R. Donaldson and J. Marx.
- [36] T. Sjöstrand, *Z. Phys.* **C42**, 301 (1989).
- [37] D. Boal, *Phys. Rev.* **C33**, 2206 (1986).
- [38] K. Geiger and B. Müller, *Nucl. Phys.* **B369**, 600 (1992).
- [39] K. Geiger, *Phys. Rev.* **D47**, 133 (1993).
- [40] X. N. Wang, *Phys. Rev.* **D43**, 104 (1991).
- [41] X. N. Wang and M. Gyulassy, *Phys. Rev.* **D44**, 3501 (1991).
- [42] K. Hahn and J. Ranft, *Phys. Rev.* **D41**, 1463 (1991);
P. Aurenche et al., *Phys. Rev.* **D45**, 92 (1992).
- [43] K. Fialkowski and A. Kotanski, *Phys. Lett.* **B107**, 132 (1981);
A. Capella and J. Tran Thanh Van, *Phys. Lett.* **B114**, 450 (1982);
P. Aurenche and F. W. Bopp, *Phys. Lett.* **B114**, 363 (1982);
A. B. Kaidalov, *Phys. Lett.* **B116**, 459 (1982).

- [44] *Handbook of Perturbative QCD*, edited by G. Sterman, CTEQ Collaboration 1993, Fermilab-Pub-93/094.
- [45] *Perturbative Quantum Chromodynamics*, edited by A. H. Mueller, Advanced Series on Directions in High Energy Physics, Vol. 5 (World Scientific, Singapore 1989)
- [46] R. D. Field, *Applications of Perturbative QCD*, Frontiers in Physics **77**, (Addison-Wesley, 1989).
- [47] R. P. Feynman, *Photon Hadron Interactions* (Benjamin Publishers, New York, N.Y., 1972).
- [48] V. N. Gribov, in Proceedings of the *VIII LNPI Winter School*, Leningrad, 1973, Vol. 2.
- [49] J. Kogut and L. Susskind, Phys. Rep. **8**, 75 (1973).
- [50] See e.g.: Yu. L. Dokshitzer, V. A. Khoze, A. H. Müller, and S. I. Troyan, *Basics of Perturbative QCD*, Advanced Series on Directions in High Energy Physics, Vol. 5 (Editions Frontieres, Gif-sur-Yvette Cedex, France, 1991)
- [51] K. Geiger and B. Müller, Phys. Rev. **D50**, (1994);
K. Geiger, Phys. Rev. **D50**, (1994).
- [52] A. Bassetto, M. Ciafaloni and G. Marchesini, Phys. Rep. **100**, 203 (1983).
- [53] J. Kogut and L. Susskind, Phys. Rev. D **9**, 697 (1974); **9**, 3391 (1974).
- [54] V. N. Gribov and L. N. Lipatov, Yad. Fiz. **15**, 781 (1972) [Sov. J. Nucl. Phys. **15**, 438 (1972)]; **15**, 1218 (1972) [**15**, 675 (1972)].
- [55] L. N. Lipatov, Yad. Fiz. **20**, 181 (1974) [Sov. J. Nucl. Phys. **20**, 94 (1975)].
- [56] Yu. L. Dokshitzer, D. I. Dyakonov, and S. I. Troyan, Phys. Rep. **58**, 269 (1980).
- [57] G. Reya, Phys. Rep. **69**, 195 (1981).
- [58] A. H. Mueller, Phys. Rep. **73**, 237 (1981).
- [59] G. Altarelli, Phys. Rep. **81** (1982).
- [60] A. P. Bukhvostov, L. N. Lipatov and N. P. Popov, Sov. J. Nucl. Phys. **20**, 287 (1975).
ibid. **20** 675 (1975).

- [61] Yu. L. Dokshitzer, Sov. Phys. JETP **46**, 641 (1977).
- [62] G. Altarelli and G. Parisi Nucl. Phys. **B126**, 298 (1977).
- [63] J. C. Collins and J. Qiu, Phys. Rev. D **39**, 1398 (1988).
- [64] See e.g.: J. C. Collins, D. E. Soper and G. Sterman, in Ref. [45], p. 1.
- [65] B. R. Webber, Ann. Rev. Nucl. Part. Sci. **36**, 253 (1986).
- [66] T. Sjöstrand, Phys. Lett. **157B**, 321 (1985);
M. Bengtsson, T. Sjöstrand and M. van Zijl, Z. Phys. **C32**, 67 (1986).
- [67] T. D. Gottschalk, Nucl. Phys. **B277**, 100 (1986); and in *From Colliders to Supercolliders*, ed. by V. Barger and F. Halzen, World Scientific, Singapore, 1987), p. 503.
- [68] G. Marchesini and B. R. Webber, Nucl. Phys. **310**, 461 (1988).
- [69] G. Marchesini and B. R. Webber, Nucl. Phys. **B238**, 1 (1984);
B. R. Webber, Nucl. Phys. **B238**, 492 (1984).
- [70] M. Bengtsson and T. Sjöstrand, Nucl. Phys. **B289**, 810 (1987); Z. Phys. **C37**, 465 (1988).
- [71] X. N. Wang and M. Gyulassy, Phys. Rev. **D45**, 844 (1992).
- [72] Proceedings of the *Large Hadron Collider Workshop*, edited by G. Jaroskog and D. Rein, Aachen, 1990, Vol. 2, CERN report 90-10, 1990.
- [73] Yu. L. Dokshitzer and S. I. Troyan, *Proceedings of the XIX Winter School of the LNPI*, Vol. 1, page 144; Leningrad, 1984.
- [74] Yu. L. Dokshitzer and S. I. Troyan, Rev. Mod. Phys. **60**, 373 (1988).
- [75] K. Henzi and P. Valin, Phys. Lett. **B160**, 167 (1985).
- [76] E. A. Uehling and G. E. Uhlenbeck, Phys. Rev. **43**, 552 (1933).
- [77] G. Bertsch, Prog. Part. Nucl. Phys. **4**, 483 (1980);
G. Bertsch and S. Das Gupta, Phys. Rep. **160**, 189 (1988).
- [78] In quantum mechanics the propagation of off-shell particles is exponentially damped according to their virtuality. In a classical visualization of off-shellness as being a property of unstable decaying states, virtual partons propagate as if they were on mass shell, but with the essential difference that their distance of travel is limited due their finite life-time that is associated with their virtuality.

- [79] J. J. Duderstadt and W. R. Martin, *Transport Theory*, Wiley, New York 1979.
- [80] S. de Groot, W. A. van Leuwen and C. G. van Weert, *Relativistic Kinetic Theory*, North Holland, Amsterdam 1980.
- [81] H.-T. Elze, preprint CERN-TH 7131/93, 1994 (unpublished).
- [82] K. Geiger, Phys. Rev. **D46**, 4965 (1992).
- [83] T. Matsui, B. Svetitsky and L. McLerran, Phys. Rev. **D34**, 783 (1986); *ibid.* **D35**, 247 (1986).
- [84] J. Kapusta, Nucl. Phys. **148**, 461 (1979).
- [85] E. V. Shuryak, Phys. Rep. **115**, 151 (1984).
- [86] K. Geiger, Phys. Rev. **D46**, 4986 (1992).
- [87] J. Randrup, Nucl. Phys. **A314**, 429 (1979).
- [88] G. Bertsch and J. Cugnon, Phys. Rev. **C24**, 2514 (1981).
- [89] R. P. Feynman and R. D. Field, Phys. Rev. **D15**, 2590 (1977); Nucl. Phys. **B136**, 1 (1978).
- [90] European Muon Collaboration: J. Ashman *et al.*, Phys. Lett. **B202** 603 (1988); M. Arneodo *et al.*, Phys. Lett. **B211**, 493 (1988).
- [91] S. J. Brodsky and H. J. Lu, Phys. Rev. Lett. **64**, 1364 (1990).
- [92] A. H. Mueller and J. Qui, Nucl. Phys. **B268**, 427 (1986);
J. Qui, Nucl. Phys. **B291**, 746 (1987), 746;
K. J. Eskola, J. Qui and X.-N. Wang, Phys. Rev. Lett. **72**, 36 (1994).
- [93] F. E. Close, J. Qiu and R. G. Roberts, Phys. Rev. **D 40**, 2820 (1989);
A. H. Mueller, Nucl. Phys. **B335**, 115 (1990)
- [94] L. L. Frankfurt and M. I. Strikman, Phys. Rep. **160**, 235 (1988).
- [95] F. Abe *et al.*, Phys. Rev. Lett. **61**, 1819 (1988).
- [96] V. M. Belyaev and B. L. Ioffe, Nucl. Phys. **B313**, 647 (1989);
D. Antreasyan *et al.*, Phys. Rev. Lett. **48**, 302 (1983);
T. D. Gottschalk, CERN Report No. TH.3810-CERN, 1984 (unpublished).

- [97] R. C. Hwa and K. Kajantie, Phys. Rev. Lett. **56**, 696 (1986).
- [98] J. D. Bjorken in *Current Induced Reactions, Proceedings of the International Summer Institute on Theoretical Particle Physics*, Hamburg, 1975, edited by J. Körner et al. (Lecture Notes in Physics, Vol. 56, Springer, New York 1976).
- [99] R. Anishetty, P. Koehler and L. McLerran, Phys. Rev. **D22**, 2793 (1980).
- [100] J. D. Bjorken, Phys. Rev. **D27**, 140 (1983).
- [101] K. Kajantie and L. McLerran, Nucl. Phys. **B214**, 261 (1983).
- [102] A. H. Mueller, Nucl. Phys. **A498**, 41c (1989).
- [103] R. Cutler and D. Sivers, Phys. Rev. **D17**, 196 (1978);
B. L. Combridge, J. Kripfganz and J. Ranft, Phys. Lett. **70B**, 234 (1977).
- [104] P. Nason, S. Dawson and R. K. Ellis, Nucl. Phys. **B303**, 607 (1988); **B327**, 49(1989);
B335, 260 (1990).
- [105] B. L. Combridge, Nucl. Phys. **B151**, 429 (1979).
- [106] B. Svetitsky, Phys. Rev. **D37**, 2484 (1988).
- [107] G. Gustafson, Z. Phys. **C15**, 155 (1982);
H. U. Bengtsson, Comp. Phys. Com. **31**, 323 (1984), 323.
- [108] JADE Collaboration, W. Bartel *et al.*, Phys. Lett. **101B**, 129 (1981); **134B**, 275 (1984); **157B**, 340 (1985);
TPC Collaboration, H. Aihara *et al.*, Z. Phys. **C28**, 31 (1985);
TASSO Collaboration, M. Althoff *et al.*, Z. Phys. **C29**, 29 (1985);
Mark II Collaboration, P. D. Sheldon *et al.*, Phys. Rev. Lett. **57**, 1398 (1986); **134B**, 275 (1984); **157B**, 340 (1985).
- [109] L. D. Landau and I. Ya. Pomeranchuk, Dokl. Akad. Nauk SSSR **92**, 535 (1953); *ibid.*, 735;
A. B. Migdal, Dokl. Akad. Nauk SSSR **96**, 49 (1954).
- [110] N. N. Nikolaev, Sov. Phys. Usp. **24**, 531 (1981);
F. Niedermeyer, Phys. Rev. **D34**, 3494 (1986).
- [111] G. Marchesini and B. R. Webber, Nucl. Phys. **B349**, 617 (1991).

- [112] R. D. Field and S. Wolfram, Nucl. Phys. **B213**, 65 (1983);
T. D. Gottschalk, Nucl. Phys. **B214**, 201 (1983); Nucl. Phys. **B227**, 413 (1983).
- [113] D. Amati, A. Bassetto, M. Ciafaloni, G. Marchesini and G. Veneziano, Nucl. Phys. **B173**, 429 (1980);
G. Marchesini, L. Trentadue and G. Veneziano, Nucl. Phys. **B181**, 335 (1981).
- [114] Note that color neutral systems need not be color singlets: for example, the color octet representation of $SU(3)_c$ contains two color neutral members, determined by the zero eigenvalues of the non-commuting generators τ_3 and τ_8 . In the classical approximation however the final partons are statistically distributed among the possible $SU(3)_c$ representations. For example, a color neutral $q\bar{q}$ pair has a probability $\frac{1}{2}$ to be a color singlet and a probability $\frac{1}{2}$ to be a color octet.
- [115] Another candidate for the measure of the two-particle phase-space separation is $\Delta'_{ij} = \tau_{ij}^\mu p_{ij,\mu}$. This form seems at first sight to be the most natural choice. However, it is absolutely inappropriate, because it correlates the particles positions and momenta. For any two particles 1 and 2 with their relative distance vector $\vec{r}_{12}$ perpendicular to their relative momentum $\vec{p}_{12}$ one has $\Delta'_{12} = 0$, independent of their distance of separation or their relative motion. Consider the extreme example of two particles with equal energy but moving ten miles apart from each other with opposite momenta. Such a configuration certainly does not meet the criteria for a minimal phase-space separation.
- [116] N. Abou-El-Naga, K. Geiger and B. Müller, J. Phys. **G18**, 797 (1992).
- [117] T. K. Gaisser and F. Halzen, Phys. Rev. Lett. **54**, 1754 (1985).
- [118] L. Durand and H. Pi, Phys. Rev. Lett. **58**, 303 (1987); Phys. Rev. **D40**, 1436 (1989).
- [119] M. Block, R. Fletcher, F. Halzen, B. Margolis, and P. Valin, Phys. Rev. **D41**, 978 (1990);
M. M. Block, F. Halzen, and B. Margolis, Phys. Lett. **B252**, 481 (1990).
- [120] See e.g.: M. L. Goldberger and K. M. Watson, *Collision Theory* (Wiley, New York, 1964), p. 330;
H. M. Fried, *Basics of functional methods and eikonal models* (Editions Frontieres, Gif-Sur-Givette, France, 1990).
- [121] M. Block and R. Cahn, in *Physics Simulations at High Energy*, (World Scientific, Singapore, 1987), p. 89;
K. Goulianos, *ibid.*, p. 127.

- [122] J. Pumplin, Phys. Rev. **D8**, 2899 (1973).
- [123] V. P. Sukhatme and F. S. Henyey, Nucl. Phys. **B108**, 317 (1976).
- [124] K. Goulianos, Phys. Rep. **101**, 169 (1983).
- [125] M. Block and R. Cahn, Rev. Mod. Phys. **57**, 563 (1985).
- [126] E. Eichten, I. Hinchliffe, K. Lane and C. Quigg, Rev. Mod. Phys. **56**, 579 (1984).
- [127] M. Glück, E. Reya and A. Vogt, Z. Phys. **C48**, 471 (1990).
- [128] Ya. I. Azimov, Yu. L. Dokshitzer, V. A. Khoze, and S. I. Troyan, Z. Phys. **C27**, 65 (1985).
- [129] S. L. Wu, Phys. Rep. **107**, 59 (1984);
MARK J Collaboration, B. Adeva *et al.*, Phys. Rep. **109**, 131 (1984);
B. Naroska, Phys. Rep. **148**, 67 (1987).
- [130] R. Brandelik *et al.*, Phys. Lett. **86B**, 243 (1979).
- [131] C. Berger *et al.*, Phys. Lett. **81B**, 413 (1979); *ibid.*, **86B**, 413 (1979); *ibid.*, **86B** 418.
- [132] R. Brandelik *et al.*, Phys. Lett. **89B** (1980), 418.
- [133] R. Brandelik *et al.*, Z. Phys. **C4**, 87 (1980).
- [134] R. Brandelik *et al.*, Nucl. Phys. **B148**, 189 (1979);
G. G. Hanson, in *Proceedings of 13th Rencontre de Moriond* (1978), edited by J. Tran Thanh Van;
C. Berger *et al.*, Phys. Lett. **78B**, 176 (1978); *ibid.* **81B** (1979), 410;
W. Bartel *et al.*, Phys. Lett. **88B**, 171 (1979);
R. Brandelik *et al.*, Phys. Lett. **89B**, 418 (1980).
- [135] M. Althoff *et al.*, Z. Phys. **C14**, 5 (1983), and references therein.
- [136] A. Capella, U. Sukhatme and J. Tran Thanh Van, Z. Phys. **C3**, 329 (1980);
J. Ranft, Phys. Lett. **B188**, 379 (1987); Phys. Rev. **D37**, 1842 (1988);
F. E. Paige and S. D. Protopopescu, in *Physics of the Superconducting Super Collider*, Proceedings of the Summer Study, Snowmass, Colorado, 1986, edited by R. Donaldson and J. Marx (APS Division of Particles and Fields, New York 1987);
T. Gaisser and T. Stanev, Phys. Lett. **B219**, 375 (1989).

- [137] UA5 Collaboration, G. J. Alner *et al.*, Phys. Rep. **154**, 247 (1987).
- [138] UA2 Collaboration, M. Banner *et al.*, Phys. Lett. **B122**, 322 (1983).
- [139] UA1 Collaboration, C. Albajar *et al.*, Nucl. Phys. **B309**, 405 (1990).
- [140] F. Abe *et al.*, Phys. Rev. Lett. **61**, 1819 (1988).
- [141] UA5 Collaboration, G. J. Alner *et al.*, Z. Phys. **C33**, 1 (1986).
- [142] F. Abe *et al.*, Phys. Rev. **D41**, 2330 (1990).
- [143] UA5 Collaboration, G. J. Alner *et al.*, Nucl. Phys. **B291**, 445 (1987).
- [144] UA5 Collaboration, R. E. Ansorge *et al.*, Z. Phys. **C43**, 357 (1989).
- [145] W. Thomé *et al.*, Nucl. Phys. **B129**, 365 (1976).
- [146] J. Whitemore *et al.*, Phys. Rep. **10C**, 273 (1974);
W. M. Morse *et al.*, Phys. Rev. **D15**, 66 (1977);
C. P. Ward *et al.*, Nucl. Phys. **B153**, 299 (1979).
- [147] X.-N. Wang, Phys. Rev. **D46**, R1900 (1992).
- [148] K. Geiger, *Nuclear and Medium Effects for Hadron Production in Ultra-relativistic Heavy Ion Collisions*, University of Minnesota UMSI report 92/174 (1992) (unpublished).
- [149] X. N. Wang and M. Gyulassy, Phys. Rev. Lett. **68**, 1480 (1992).
- [150] U. Heinz, K. S. Lee and E. Schedermann in *Quark Gluon Plasma*, edited by R. C. Hwa (World Scientific, Singapore 1990), pp. 471, and references therein.
- [151] J. Zimányi in *Particle Production in Highly Excited Matter*, edited by H. Gutbrod and J. Rafelski (Plenum Press, N.Y., 1993).
- [152] J. Zimányi in M. Gyulassy and M. Pluemer, Phys. Lett. **234B**, 432 (1990).
- [153] L. Xiong and E. Shuryak, Phys. Rev. **C49**, 2203 (1994).
- [154] K. Geiger, Phys. Rev. **C50**, (1994).
- [155] *1989 RHIC Conceptual Design Report*, BNL publication 52195, 1989.

- [156] J. Schukraft, talk given at the *Workshop on a dedicated heavy ion detector for LHC*, CERN, Feb. 1993.
- [157] E. M. Levin and M. G. Ryskin, Nucl. Phys. **B304**, 805 (1989).
- [158] J.-P. Blaizot and A. H. Mueller, Nucl. Phys. **B289**, 847 (1987).
- [159] A. Mueller and J. Qiu, Nucl. Phys. **B286**, 427 (1986).
- [160] See. e.g.: N. A. Krall and A. W. Trivelpiece, *Principles of Plasma Physics* (McGraw-Hill, New York, 1973);
R. Balescu, *Statistical Mechanics of Charged Particles*, (Interscience Publishers, London 1963).
- [161] B. Müller and X.-N. Wang, Phys. Lett. **B283**, 171 (1992);
S. Mrówczyński, Phys. Lett. **B314**, 118 (1993); Phys. Rev. C..., ... (1994).
- [162] Proceedings of the International Workshop on *Quark Gluon Plasma Signatures*, Strasbourg, France, 1991, edited by V. Bernard *et al.* (Editions Frontieres, Gif-sur-Yvette, France, 1991).
- [163] C. P. Singh, Int. J. Mod. Phys. **A 29**, 7185 (1992).
- [164] B. Müller, Nucl. Phys. **544**, 95c (1993).
- [165] E. V. Shuryak, Phys. Lett. **B 78**, 150 (1978); Sov. J. Nucl. Phys. **28**, 408 (1978).
- [166] K. Kajantie and H. Miettinen, Z. Phys. **C 9**, 341 (1981);
F. Halzen and H. C. Liu, Phys. Rev. **D 25**, 1842 (1982);
D. K. Srivastava, B. Sinha, M. Gyulassy and X.-N. Wang, Phys. Lett. **276B** (1991), 285;
J. I. Kapusta, P. Lichard and D. Seibert, Phys. Rev. **D44**, 2774 (1991).
- [167] P. V. Ruuskanen, Nucl. Phys. **A544**, 169c (1992).
- [168] J. I. Kapusta, Nucl. Phys. **A566**, 45c (1994).
- [169] K. Kajantie and H. Miettinen, Z. Phys. **C 9**, 341 (1981);
L. D. McLerran and T. Toimela, Phys. Rev. **D 31**, 545 (1985);
R. C. Hwa and K. Kajantie, Phys. Rev. **D 32**, 1109 (1985);
K. Kajantie, M. Kataja, L. D. McLerran and P. V. Ruuskanen, Phys. Rev. **D 34**, 811 (1986).

- [170] J. I. Kapusta, L. D. McLerran and D. K. Srivastava, Phys. Lett. **B 283**, 145 (1992);
B. Kämpfer and O. P. Pavlenko, Phys. Lett. **B289**, 127 (1992).
- [171] K. Geiger and J. I. Kapusta, Phys. Rev. Lett. **70**, 1290 (1993).
- [172] K. Geiger, Phys. Rev. Lett **71**, 3075 (1993).
- [173] E. Shuryak and L. Xiong, Phys. Rev. Lett. **70**, 2241 (1993).
- [174] X.-N. Wang and K. Eskola, preprints LBL-34409 and LNBL-34664, Berkeley 1993
(to be published).
- [175] X.-N. Wang, *private communication*.
- [176] P. Koch, B. Müller, and J. Rafelski, Phys. Rep. **142**, 167 (1986), and references
therein.
- [177] J. Kapusta and A. Mekjian, Phys. Rev. D **33**, 1304 (1986).
- [178] K. Geiger, Phys. Rev. **D48**, 4129 (1993).
- [179] A. Shor, Phys. Lett. **B 215**, 375 (1988); Phys. Lett. **B 233**, 231 (1989).
- [180] B. Müller and X.-N. Wang, Phys. Rev. Lett. **68**, 2437 (1992);
- [181] J. Alam, S. Raha, and B. Sinha, VECC preprint, Calcutta 1994 (unpublished);
I. Sarcevic and P. Valerio, preprint AZPH-TH/94-13, Tucson (unpublished).
- [182] K. Geiger and J. I. Kapusta, Phys. Rev. **D47**, 4905 (1993).
- [183] H. Satz, preprint CERN. TH-7064/93, 1993 (to appear in Z. Phys. **C**.
- [184] X.-N. Wang and K. Eskola, Phys. Rev. **D49**, 1284 (1994).
- [185] T. Matsui and H. Satz, Phys. Lett. **B178**, 416 (1986).
- [186] J.-P. Blaizot in Ref. [5], pp. 631, and references therein.
- [187] J. Milana, Phys. Rev. Lett. **62**, 2921 (1989).
- [188] J. Cleymans and C. Vanderzande, Phys. Lett. **B147**, 186 (1984);
H. J. Reusch, Z. Phys. **C26**, 105 (1984).

- [189] S. Gavin, M. Gyulassy, and A. Jackson, Phys. Lett. **B208**, 257 (1988);
R. Vogt, M. Prakash, P. Koch, and T. H. Hansson, Phys. Lett. **B208**, 263 (1988);
J.-P. Blaizot and J.-Y. Ollitrault, Phys. Rev. D **39**, 232 (1989).
- [190] S. J. Brodsky and A. H. Mueller, Phys. Lett. **B206**, 685 (1988).
- [191] S. Gupta and H. Satz, Phys. Lett. **B283**, 439 (1992); Z. Phys. **C55**, 391 (1992).
- [192] H. Satz, Nucl. Phys. **A544**, 371c (1992).
- [193] J. C. Collins and W.-K. Tung, Nucl. Phys. **B278**, 934 (1986).
- [194] M. A. G. Aivazis, F. I. Olness and W.-K. Tung, preprint MSU-HEP 93/15, Michigan State University 1993;
M. A. G. Aivazis, J. C. Collins, F. I. Olness and W.-K. Tung, preprint MSU-HEP 93/17.
- [195] M. L. Mangano, P. Nason, and G. Ridolfi, *Fixed Target Hadroproduction of Heavy Quarks*, preprint IFUP-TH-37/92, Istituto Nazionale di Fisica Nucleare (INFN), Pisa, Italy (unpublished).
- [196] J. Appel *et al.*, Ann. Rev. Nucl. Part. Sci. **42**, 367 (1993); L. Rossi, preprint INFN-AE-91-16, Oct. 1991, presented at the 4th Int. Symp. on Heavy Flavor Physics, Orsay, France, June 25-29, 1991 (unpublished); G. Introzzi, E771 Coll., 3rd Topical Seminar on Heavy Flavors, S. Miniato, June 17-21 1991.
- [197] K. Eskola, K. Kajantie and P. V. Ruuskanen, preprint HU-TFT-94-6, Helsinki 1994 (unpublished).
- [198] There is recent experimental evidence from HERA data that at very small $x \lesssim 10^{-3}$ the gluon structure function of a proton exhibits a "Lipaton enhancement" $x\hat{g}(x) \propto x^{-c}$ with $c \approx 1/2$. Nevertheless, for reasons of simplicity mentioned in the text, I will use the approximation of a constant behaviour at small x , which corresponds to $c = 0$.
For recent experimental and phenomenological progress, see e.g.:
A. De Roeck, in the proceedings of the *Durham Workshop on HERA Physics*, Durham, U.K., 1993; Journ. Phys. **G19**, 1549 (1993);
S. R. Magill, in the proceedings of the *Argonne Workshop on Physics at Current Accelerators and the Supercollider*, Argonne, 1993; Journ. Phys. **G19**, 1535 (1993);
G. Wolf, in the proceedings of the *International Workshop on Deep Inelastic Scattering*, Eilat, Israel, 1994.

- [199] L. McLerran and R. Venugopalan, preprints TPI-MINN-93-44/T, TPI-MINN-93-52/T, and TPI-MINN-94-7/T, Minneapolis 1993/94 (unpublished).
- [200] A. V. Selikov, Phys. Lett. **B 268**, 263 (1991).
- [201] X.-N. Wang and M. Guylassy, Columbia University preprint CU-IP-589 (unpublished).
- [202] K. Werner, preprint HD-TV-94-6, Heidelberg 1994 (unpublished).
- [203] M. H. Kalos, *Monte Carlo Methods*, Wiley, New York 1986.
- [204] R. W. Hockney and J. W. Castwood, *Computer Simulations using Particles*, Adam Hilger, Bristol New York 1988.
- [205] *Physics Simulations at High Energy*, edited by V. Barger, T. Gottschalk and F. Halzen World Scientific, Singapore, 1987.
- [206] K. Konishi, A. Ukawa and G. Veneziano, Nucl. Phys. **B157**, 45 (1979); A. Basetto, M. Ciafaloni and G. Marchesini, Nucl. Phys. **B163**, 477 (1980).

TABLE CAPTIONS

Table 1: Dynamical quantities and mean number of parton interactions in simulated $p\bar{p}$ collisions at $\sqrt{s} = 200$ -1800 GeV. The values of Q_0 are obtained from accumulated statistics according to eq. (78), whereas $p_{\perp cut}$, μ_0 and b_{max} are input values.

Table 2: Average multiplicities of partons and resulting hadrons in simulated $p\bar{p}$ collisions at $\sqrt{s} = 200$ -1800 GeV. The partons are distinguished in primary and secondary quanta. The initial parton distributions consist only of primary quanta whereas the final parton configuration shows both the numbers of remaining spectator partons (prim.) and the multiplicities of interacted or produced partons (sec.). The hadron multiplicities and their average $p_{\perp}$ result from the cluster hadronization of these final partons.

Table 3: Model results for dynamical quantities and mean number of parton interactions in central $Au + Au$ collisions at $\sqrt{s} = 200$ A GeV and $\sqrt{s} = 6300$ A GeV. The values of Q_0 are obtained from accumulated statistics according to eq.(78), whereas $p_{\perp cut}$ and μ_0 are input values.

Table 4: Average multiplicities of partons and resulting hadrons in central $Au + Au$ collisions at $\sqrt{s} = 200$ A GeV and $\sqrt{s} = 6300$ A GeV. The partons are distinguished in primary and secondary quanta. The multiplicities of the initial partons were obtained from the nuclear structure functions and reflect the flavor composition of the incoming nuclei. The mean number of final partons refers to the parton distributions just before hadronization. The hadron multiplicities and their average $p_{\perp}$ result from the cluster hadronization of these final state parton configurations.

FIGURE CAPTIONS

Figure 1: Diagram of the longitudinal space-time evolution of a relativistic nucleus-nucleus collision, from the moment of nuclear overlap, via pre-equilibrium and quark gluon plasma to the formation of final state hadrons.

Figure 2: a) *String picture*: primary interactions lead to color flux tubes (strings) which break by $q\bar{q}$ production. b) *Parton approach*: multiple scatterings accompanied by emission and absorption of quarks and gluons are described as internetworked parton cascades.

Figure 3: a) Coherent space-like cascade of partons as part of the wavefunction of a fast moving nucleon. b) The interaction with another particle destroys the coherence of the cascade and sets 'free' the partons of the struck branch of the cascade.

Figure 4: Diagrammatic representation of the Q^2 -evolution of the nucleon structure function as the convolution of a hadronic structure function $h_N^a(x, Q_0^2)$ with a partonic structure function $f_a(x, Q^2, Q_0^2)$.

Figure 5: Schematic picture of the parton evolution in a fast hadron in terms of successive branchings that are ordered as self-enclosed fluctuations.

Figure 6: Probabilistic description in the LLA: a) examples for contributing ladder diagrams; b) typical interference diagrams that are suppressed; c) representation of squared amplitude in terms of elementary branchings in which the non-interacting partons are virtual fluctuations gathered into blobs of self-energy type.

Figure 7: Resolution of the parton substructure of a nucleon when probed at different values of x and Q^2 .

Figure 8: Simplified picture of the probabilistic parton evolution (solid lines are quarks, curly lines are gluons). The x -values are the longitudinal momentum fractions carried by the quarks. [From Ref. [63]].

Figure 9: The three essential elements of the reaction $p + \bar{p} \rightarrow H_1 + H_2 + \dots + \text{anything}$: a) initial parton distributions according to the p and $\bar{p}$ structure functions (SF); b) parton evolution: space-like branchings of incoming partons (SB), hard scattering (HS), time-like branchings of outgoing partons (TS); c) hadronization of final state partons.

Figure 10: Illustration of the characteristics of space-like (left) and time-like (right) branchings. Solid lines are quarks, curly lines are gluons: a) ladder diagram corresponding the squared amplitude of a sequence of space-like and time-like branchings, respectively; b) tree-like structure of subsequent branchings; c) kinematics of a single space-like, respectively time-like branching $a \rightarrow bc$.

Figure 11: Perturbative QCD predictions from numerical calculations of minimum bias pp ($p\bar{p}$) collisions as a function of $\sqrt{s}$ in comparison to experimental data. a) average transverse momentum of charged particles; b) mean charged multiplicity; c) total and inelastic pp ($p\bar{p}$) cross-sections. [From [72] (a,b) and [71] (c).]

Figure 12: *BEL*-effect: the probability distribution $P_{inel}(b)$ for an inelastic pp ($p\bar{p}$) collision versus impact parameter b , for various energies $\sqrt{s}$, shows how the nucleons' appear increasingly blacker, edger, and larger.

Figure 13: Diagrammatic representation of a particular term $I_a^{(k)} = I_a^{(k)gain} - I_a^{(k)loss}$ that contributes to the collision kernel in (38) and changes the parton distribution of species a with momentum p_i due to the process k around $\vec{r}$ at time t .

Figure 14: Absolute error $\Delta(S/N)$ in the entropy per particle due to discretizing phase-space in the case of a spatially uniform system of classical particles. The particles were distributed uniformly with an isotropic momentum distribution $f(\vec{p}) = f_0 \exp[-p/a]$ where $p \equiv |\vec{p}|$ and a a parameter. The cell size in momentum space depends on the resolution parameter $\xi = \Delta p/a$ via Δp and a was chosen to be 1 GeV.

Figure 15: Absolute error in the entropy per particle as a function of phase-space cell size (in units of h^3), resulting from counting statistics with a finite number of particles. The curves correspond to a system with a total number of $N = 200 - 50000$ classical particles with momentum distribution as in Fig. 14, spatially confined in a cylinder of 2 fm in length and 4 fm in radius, i.e. a volume of $\simeq 100 \text{ fm}^3$.

Figure 16: Time evolution of the entropy production of $N = 10000$ non-interacting particles, expanding isotropically from the center of a cylindrical volume $\simeq 100 \text{ fm}^3$. The particles, initially distributed according to the distribution of Fig. 14, progressively populate phase-space with time to finally fill out the volume of the cylinder uniformly.

Figure 17: The three components of the model: (1) the initial state constructed in terms of the parton distribution of the incoming nuclei at a resolution scale Q_0 ; (2) the evolution of parton cascades in phase-space and time within the perturbative QCD domain between Q_0 and the cut-off μ_0 ; (3) the formation of color neutral clusters from the final state partons at the end of the perturbative evolution at μ_0 with subsequent the fragmentation of the clusters into final hadrons at Λ .

Figure 18: The ratio of nuclear structure function to nucleon structure function $\rho^A(x) = A^{-1}F^A(x)/F^N(x)$ as a function of x in the small and medium x region for different nuclear mass number A . The data points are from Ref. [95] and the curves are the parametrization (86). [From Ref. [41]].

Figure 19: Distributed Lorentz contraction: The spatial distribution of the partons in the rest frame of each nucleus becomes, after boosting to the CM frame, a disc of highly Lorentz contracted valence quarks of $(\Delta z)_v$, surrounded by clouds of softer gluons and sea quarks of $(\Delta z)_{g,s}$.

Figure 20: Schematical illustration of a typical parton cascade development initiated by a collision of two partons a and b . The incoming primary parton a evolves through a space-like cascade from the initial resolution scale Q_0^2 up to the scale Q_{ab}^2 at which it collides with parton b . The outgoing partons c and d both initiate time-like cascades which are described as a combination of multiple branchings (parton c), absorptions (parton i) or secondary scatterings (parton e). A parton is not allowed to branch further if its virtuality falls below the invariant mass cut-off μ_0 . In this case, it propagates freely until it interacts again or finally hadronizes.

Figure 21: Diagrams corresponding to the effective matrix elements $\mathcal{M}_{eff}$ of (94)-(96) that contribute to the collision integrals J_{abcd} and K_{abc} . Each of the partons coming into (emerging from) an elementary vertex is dressed by a S (T) formfactor, taking into account higher order perturbative QCD corrections. The S formfactors represent radiative corrections from primary space-like partons having their first collisions, whereas the T formfactors account for multiple emissions and absorptions by time-like secondary partons. Note that there is no S formfactor for the $1 \rightarrow 2$ branchings, because these processes occur only for time-like excited partons originating from previous collisions or decays.

Figure 22: Illustration of the regularization of the infrared divergent parton-parton cross-sections employed in the PCM. The cross-section is divided in a (semi)hard part above $p_{\perp cut}$ and a soft contribution below $p_{\perp cut}$.

Figure 23: Illustration of the cluster formation scheme: First final state gluons are split into collinear quark-antiquark pairs. Then color neutral $q_i \bar{q}_j$ clusters are formed from a quark q_i and an antiquark $\bar{q}_j$ with the smallest separation in phase-space.

Figure 24: Invariant mass spectra of central clusters formed according to the cluster formation scheme in simulations of $p\bar{p}$ collisions at different beam energies. The distribution is strongly damped above $\sim 2\mu_0$ ($\mu_0 = 1$ GeV) and is roughly independent of the beam energy $\sqrt{s}$.

Figure 25: Schematics of the cluster fragmentation: Each cluster is allowed to convert into a single hadron (if the invariant mass is small enough) or to 'decay' isotropically in its rest frame into two hadrons. Very massive clusters are assumed to fission first, until their masses falls below a fission threshold M_{fiss} , before they transform into hadrons.

Figure 26: a) The eikonal QCD-cross section σ_{QCD} and eikonal diffractive cross section σ_0 obtained in Ref. [116] from fits for σ_{ND} and σ_D using eqs. (123) and (124). The cross sections are set equal for $\sqrt{s} > 1.8$ TeV. b) The total, inelastic, nondiffractive, elastic, and diffractive cross sections as obtained from σ_0 and σ_{QCD} on the basis of the eikonal representation. The points correspond to experimental data [124, 125].

Figure 27: The momentum cut-off $p_{\perp cut}$ as obtained in Ref. [116], for EHLQ [126] and GRV [127] structure functions. The points indicate where the detailed numerical analysis was performed. The straight lines represent least-square fits of the form (126).

Figure 28: Transverse momentum distribution of charged particles in simulated e^+e^- -annihilation events at $Q = 15$ GeV and $Q = 29$ GeV. The experimental data are from Ref. [130].

Figure 29: Mean transverse momentum of charged particles in $e^+e^- \rightarrow$ hadrons as a function of center-of-mass energy Q . The data are from Refs. [130, 131].

Figure 30: Overall mean charged multiplicity in e^+e^- hadrons. The data are from Refs. [134].

Figure 31: Charged particle average multiplicities in $e^+e^- \rightarrow$ hadrons: as function of center of mass energy Q . Data are from Ref. [135].

Figure 32: Invariant inclusive cross-sections of a) partons and b) charged hadrons in $p\bar{p}$ collisions at $\sqrt{s} = 200$ -1800 GeV. The histograms are model results and the data are from Refs. [139, 140]. Both the calculation and experimental data refer to the (pseudo)rapidity regions $|\eta| < 2.5$ for $\sqrt{s} = 200, 546$ and 900 GeV, and $|\eta| < 1$ for $\sqrt{s} = 1800$ GeV.

Figure 33: a) Rapidity spectra of partons and b) corresponding pseudorapidity distributions of charged hadrons in $p\bar{p}$ collisions at $\sqrt{s} = 200$ -1800 GeV. The model results are shown as histograms and the data points are from Refs. [141, 142].

Figure 34: Charged particle multiplicity distributions in $p\bar{p}$ collisions at $\sqrt{s} = 200, 546$ and 900 GeV. The data points are from Refs. [143, 144]. The histograms are the calculated distributions. The various patterned curves result from separating the contributions from events with $k = 0, 1, 2$ and ≥ 3 parton-parton interactions.

Figure 35: Central rapidity density at $\eta = 0$ of charged particles in inelastic pp and $p\bar{p}$ collisions versus $\sqrt{s}$. The solid line is the HIJING calculation and the data are from Refs. [141, 145, 146]. The dashed line is for events without jet production in the HIJING simulations. [From Ref. [71]].

Figure 36: The correlation function versus $\Delta\phi$ between two charged particles in the full rapidity range in $p\bar{p}$ collisions at $\sqrt{s} = 1800$ GeV. The dashed line is for all charged

particles, dotted line for particles with transverse momentum $p_\perp > 0.5$ GeV, dash-dotted line for $p_\perp > 1$ GeV, and solid line for $p_\perp > 1.5$ GeV. [From Ref. [147]].

Figure 37: Comparison between the predictions of HIJING and the PCM for the transverse momentum and (pseudo)rapidity spectra of charged hadrons produced in central heavy ion collisions at RHIC and LHC energies. Notice the difference in the central rapidity densities for RHIC (LHC) between HIJING, $dN_{ch}/d\eta|_{\eta=0} \simeq 600$ (1900), and the PCM, $dN_{ch}/d\eta|_{\eta=0} \simeq 1200$ (2400).

Figure 38: Average cumulative number of secondary gluons and quarks per nucleon produced during collisions of $S + S$ and $Au + Au$ at $\sqrt{s} = 200$ A GeV. The curves (a) and (e) refer to the two space-time evolution scenarios explained in the text.

Figure 39: The model predictions (solid histogram) for the pseudorapidity distributions of charged particles in central $Au + Au$ collisions at $\sqrt{s} = 200$ A GeV (left) and $\sqrt{s} = 6300$ A GeV (right). The various histograms show the characteristics of the space-time evolution schemes a) - e) defined in Sec. 4.3: a) ‘naive’ evolution (dotted), b) nuclear shadowing (long dashed), c) parton fusion and absorption (dashed dotted), d) Landau-Pomeranchuk-Migdal effect (short dashed), and e) soft gluon interference (solid). Note that case e) is the full calculation including all these effects.

Figure 40: The model results for the ratio of the inclusive charged hadron spectra of central $Au + Au$ collisions to those of $p + p$ for $\sqrt{s} = 200$ A GeV (dashed histogram) and $\sqrt{s} = 6300$ A GeV (solid histogram). Both the ratio of pseudorapidity distributions (left) and the one of the transverse momentum distributions (right) were calculated with the inclusion of nuclear shadowing, parton fusion and absorption, the Landau-Pomeranchuk effect and soft gluon interference.

Figure 41: ‘Lego’ plot of the transverse energy distribution $dE_\perp/dy d\phi$ of secondary partons versus rapidity y and azimuthal angle ϕ in $p\bar{p}$ (top) and $Au + Au$ collisions (bottom). In $p\bar{p}$ the jets, once produced, propagate freely, whereas in $Au + Au$ the produced jets are quenched and wash out with time.

Figure 42: A considered parton a can interact during a time span Δt with all kinds of partons b , that are contained in the volume $\Delta z \pi R_A^2$ which the parton a traverses within Δt . The density of scatterers F_b determines the magnitude of the interaction probability W_a .

Figure 43: Increase of the interaction probability W_g for gluons to fuse at small rapidities $|y| \leq 1.5$, obtained from pA simulations at $P = 100$ GeV and $P = 3000$ GeV on the basis of eqs. (132) and (133). The time t is measured in the CM_{NN} frame.

Figure 44: Illustration of the prescription for choosing for a given parton a the collision partner b . Of all potential scattering candidates b contained within the effective interaction area $\pi(\hat{b}_{ab}^{max})^2$, given by the cross-section $\hat{\sigma}_{ab}(p_{\perp cut}^2)$, the one with the minimum impact parameter $\hat{b}_{ab}^{min}$ is selected.

Figure 45: Dependence of the average impact parameter $\langle \hat{b} \rangle$ on the mass number A in pA collisions up the first 3 fm after the interpenetration of the proton for the two considered beam momenta $P = 100$ GeV and $P = 3000$ GeV.

Figure 46: Rapidity density dN_{pA}^{ch}/dy of charged hadrons in the central region around $y = 0$ versus mass number A in pA collisions at $P = 100$ and $P = 3000$ GeV. The curves full circles include the effects of parton saturation and dynamical screening, whereas the curves with open circles result from neglecting of these effects.

Figure 47: Nuclear dependence on the mass number A of the calculated average $\langle p_{\perp} \rangle^{ch}$ per secondary charged hadron in pA collisions at $P = 100$ GeV and $P = 3000$ GeV.

Figure 48: Distributions of numbers of partons per nuclear mass number A in central $Au+Au$ collisions ($A = 197$) at $\sqrt{s} = 200$ A GeV in the center-of-mass frame of the colliding nuclei at different times along the longitudinal z -axis. The individual contributions from *virtual + real* (prim.+ sec.) and from exclusively *real* (sec.) gluons and quarks, N_g/A , respectively $(N_q + N_{\bar{q}})/A$, are indicated by the various shaded areas, whereas the solid black areas display the valence quark distribution $(N_q - N_{\bar{q}})/A$, which is proportional to the baryon number distribution.

Figure 49: Distributions of numbers of partons per nuclear mass number A along the transverse $r_{\perp}$ -direction, corresponding to Fig. 1.

Figure 50: Distribution of vertices of the production of secondary partons in longitudinal (z) and transverse ($r_{\perp}$) direction during central $Au + Au$ collisions of The point $t = 0$ fm is the moment the nuclei begin to overlap.

Figure 51: Proper time development of the partons' momentum distributions in the central collision region (141) of $Au + Au$ at $\sqrt{s} = 200$ A GeV from the moment of maximum overlap ($\tau = 0$) until the end of the calculations ($\tau = 2.4$ fm). Shown are the normalized (but scaled) distributions (142) for gluons and the different quark species.

Figure 52: Pressure p , particle density n , energy density ε , entropy density s and associated effective temperature T of the system of partons in the central collision region (15) as a function of proper time τ in $Au + Au$ collisions at $\sqrt{s} = 200$ A GeV and zero impact.

Figure 53: a) Increase with time of the total entropy per particle S/N produced by the secondary partons in the central region of $Au + Au$ collisions with $\sqrt{s} = 200$ A GeV - 6300

A GeV. b) The relaxation times τ_{relax} associated with the entropy production in a) as a function of the beam energy. The points are the values estimated from the approximate exponential relaxation of the specific entropy for the considered beam energies $\sqrt{s} = 200$ A GeV - 6300 A GeV.

Figure 54: a) Average transverse mass $\langle m_{\perp a} \rangle = \langle \sqrt{p_{\perp a}^2 + m_a^2} \rangle$ of the gluons and the different quark flavors as a function of time in central $Au+Au$ collisions at $\sqrt{s} = 200$ A GeV. The quantity $\langle m_{\perp}^{(ideal)} \rangle$ refers to average transverse mass (momentum) of massless partons in an ideal QGP with the same temperature dependence as the actual parton matter. b) The different 'temperatures' for gluons and the quark+antiquark species extracted from the average particle energies in the local restframe of the matter. The dotted curve is the extracted temperature of the parton system as a whole, $T = 4\epsilon/(3s)$.

Figure 55: Time development of the ratios (153) that measure the deviation of the actual particle and energy densities from their equilibrium values for gluons and quarks plus antiquarks in the central region of $Au+Au$ collisions at $\sqrt{s} = 200$ A GeV. For u and d quarks the additional dashed lines represent the contributions of the secondary u , respectively d , quarks alone, i.e. without the already initially present valence quarks (except for those that have scattered at least once).

Figure 56: Time evolution of the temperature T and the fugacities λ_g and λ_q of gluons and quarks in a parton plams created in $Au + Au$ collisions at (a) RHIC energy $\sqrt{s} = 200$ A GeV and (b) LHC energy $\sqrt{s} = 6000$ A GeV. [From Ref. [13]].

Figure 57: Schematic space-time diagram of the parton evolution during the early stage of an ultra-relativistic heavy ion collision. The central collision region around rapidity $Y = 0$ is initially populated by materialization due to primary scatterings and subsequent cascading of partons, and evolves to a thermalized, but not chemically equilibrated, gluon dominated plasma.

Figure 58: a) Feynman diagrams for the lowest order QCD processes to produce an l^+l^- pair through parton-parton collisions (154)-(157), or through bremsstrahlung by a virtual quark or antiquark (158). b) Examples of higher order processes that are effectively accounted for in the present approach by decomposing them into the elementary processes shown in a). c) Difference between gluon and photon emission due to the absence of a QCD formfactor of photons: emitted gluons can radiate themselves further off-spring, whereas a photon emission is not accompanied by additional gluon radiation.

Figure 59: The calculated dilepton yield in pp collisions from the PCM simulations compared to the standard Drell-Yan result.

Figure 60: Invariant mass distributions of dileptons in the central rapidity region in $Au + Au$ collisions at $\sqrt{s} = 200$ A GeV. The bold solid line is the total l^+l^- yield from the parton cascade evolution after $\tau = 2$ fm/c. The three curves below show the separate contributions from ‘Drell-Yan’ processes (primary-primary), from interactions involving primary with secondary partons, and from interactions involving only primary partons.

Figure 61: Scaling the calculated dilepton mass distribution for pp collisions at RHIC energy by $A^{4/3}$ reproduces the primary-primary contribution in central $Au + Au$ collisions.

Figure 62: Individual contributions from the various production processes to the dilepton distribution $dN/dM^2 dy|_{y=0}$ after $\tau = 2$ fm in central $Au + Au$ collisions at $\sqrt{s} = 200$ A GeV.

Figure 63: Comparison of the dilepton mass spectrum from the PCM calculation with the free-streaming and hydrodynamic models discussed in Ref. [168].

Figure 64: The dilepton mass spectrum (from $q\bar{q}$ annihilation only) during $Au + Au$ collisions at RHIC energy, comparing “hot glue”, standard”, and Drell-Yan scenarios. For details, see Ref. [173].

Figure 65: Central rapidity density $dN_{Q+\bar{Q}}/dy|_{y=0}$ versus $\sqrt{s}/A$ of produced strange, charm and bottom quarks plus antiquarks ($Q = s, c, b$) in central $Au + Au$ collisions: (a) *with* plasma formation, as obtained from the parton cascade simulations of $Au + Au$ collisions on the basis of eq. (170), and (b) *without* plasma formation, as calculated from pp collisions and scaled to AA . The thick lines are the parton cascade results, and the thin lines are the corresponding standard parton model predictions without subsequent cascading and gluon bremsstrahlung.

Figure 66: Individual contributions from the processes (162)-(166) to the central rapidity density of (a) strange and (b) charm quarks plus antiquarks as a function of $\sqrt{s}/A$ for the case of $Au + Au$ collisions *with* plasma formation.

Figure 67: Interrelation between initial temperatures T_0 at the time of plasma formation in the central region, and, (a) the rapidity densities $dN_{Q+\bar{Q}}/dy$, as well as (b) the ratios $\rho_{Q+\bar{Q}}$ defined in eq. (171), around $y = 0$.

Figure 68: Rapidity densities $dN_R/dy|_{y=0}$ of the vector meson resonances $R = \phi, J/\psi, \psi', \Upsilon$ as a function of $\sqrt{s}/A$ for central $Au + Au$ collisions *with* plasma formation, as obtained from the quark yields calculated with the parton cascade simulations of $Au + Au$ collisions. For details, see Ref. [178].

Quantity	$\sqrt{s} = 200 \text{ GeV}$	$\sqrt{s} = 546 \text{ GeV}$	$\sqrt{s} = 900 \text{ GeV}$	$\sqrt{s} = 1800 \text{ GeV}$
$Q_0 \text{ (GeV)}$	1.88	2.65	3.42	4.84
$p_{\perp cut} \text{ (GeV)}$	1.50	1.70	2.55	3.45
$\mu_0 \text{ (GeV)}$	1	1	1	1
$b_{max} \text{ (fm)}$	1.30	1.40	1.55	1.95
$\langle n_{2 \rightarrow 2}^{(hard)} \rangle$	2.58	5.35	7.68	9.04
$\langle n_{2 \rightarrow 2}^{(soft)} \rangle$	0.78	1.03	1.86	3.29
$\langle n_{2 \rightarrow 2}^{(prim.coll.)} \rangle$	2.42	3.96	4.94	5.88
$\langle n_{2 \rightarrow 1} \rangle$	0.07	0.15	0.29	0.58
$\langle n_{1 \rightarrow 2}^{(time-like)} \rangle$	3.58	6.85	15.61	24.27
$\langle n_{1 \rightarrow 2}^{(space-like)} \rangle$	0.030	0.029	0.021	0.008
% events with no parton scattering	0.31	0.21	0.16	0.07

Table 1

Quantity	$\sqrt{s} = 200 \text{ GeV}$		$\sqrt{s} = 546 \text{ GeV}$		$\sqrt{s} = 900 \text{ GeV}$		$\sqrt{s} = 1800 \text{ GeV}$	
initial partons	prim.	sec.	prim.	sec.	prim.	sec.	prim.	sec.
$\langle n_{parton} \rangle$	59.11	—	76.10	—	93.70	—	103.09	—
$\langle n_g \rangle$	43.01	—	56.59	—	69.29	—	76.23	—
$\langle n_u + n_{\bar{u}} \rangle$	7.28	—	8.34	—	9.32	—	10.23	—
$\langle n_d + n_{\bar{d}} \rangle$	5.00	—	6.04	—	7.46	—	8.00	—
$\langle n_s + n_{\bar{s}} \rangle$	3.04	—	3.97	—	5.47	—	5.45	—
$\langle n_c + n_{\bar{c}} \rangle$	0.78	—	1.16	—	2.16	—	3.02	—
final partons	prim.	sec.	prim.	sec.	prim.	sec.	prim.	sec.
$\langle n_{parton} \rangle$	56.02	5.52	70.40	10.97	86.88	21.91	93.43	31.60
$\langle n_g \rangle$	40.75	4.41	52.32	8.84	63.29	18.89	68.65	26.58
$\langle n_u + n_{\bar{u}} \rangle$	6.88	0.46	7.38	1.18	8.83	1.41	9.45	1.73
$\langle n_d + n_{\bar{d}} \rangle$	4.74	0.38	5.80	0.50	7.29	0.75	7.50	1.50
$\langle n_s + n_{\bar{s}} \rangle$	2.92	0.20	3.83	0.34	5.42	0.63	5.08	1.28
$\langle n_c + n_{\bar{c}} \rangle$	0.73	0.07	1.07	0.11	2.05	0.23	2.75	0.51
final hadrons								
$\langle n_{ch} \rangle$	20.3		29.9		35.1		42.7	
$\langle n_{ch} \rangle / \langle n_{hadron} \rangle$	0.57		0.57		0.58		0.59	
$\langle p_{\perp ch} \rangle \text{ (GeV)}$	0.39		0.43		0.48		0.51	
$\langle p_{\perp ch}^2 \rangle \text{ (GeV}^2\text{)}$	0.31		0.35		0.47		0.61	

Table 2

Quantity	$\sqrt{s} = 200$ A GeV (RHIC) $\sqrt{s} = 6300$ A GeV (LHC)	
Q_0 (GeV)	2.28	5.23
$p_{\perp cut}$ (GeV)	1.55	3.95
μ_0 (GeV)	1	1
$\langle n_{2 \rightarrow 2}^{(hard)} \rangle$	11026	37980
$\langle n_{2 \rightarrow 2}^{(soft)} \rangle$	2855	10345
$\langle n_{2 \rightarrow 2}^{(prim.coll.)} \rangle$	4674	9249
$\langle n_{2 \rightarrow 1} \rangle$	860	7266
$\langle n_{1 \rightarrow 2}^{(time-like)} \rangle$	7451	24120
$\langle n_{1 \rightarrow 2}^{(space-like)} \rangle$	143	278

Table 3

Quantity	$\sqrt{s} = 200$ A GeV (RHIC) $\sqrt{s} = 6300$ A GeV (LHC)			
initial partons	prim.	sec.	prim.	sec.
$\langle n_{parton} \rangle$	11821	—	20657	—
$\langle n_g \rangle$	8471	—	14825	—
$\langle n_u + n_{\bar{u}} \rangle$	1320	—	2284	—
$\langle n_d + n_{\bar{d}} \rangle$	1214	—	2112	—
$\langle n_s + n_{\bar{s}} \rangle$	666	—	1172	—
$\langle n_c + n_{\bar{c}} \rangle$	150	—	264	—
final partons	prim.	sec.	prim.	sec.
$\langle n_{parton} \rangle$	5687	13638	7590	42454
$\langle n_g \rangle$	4165	10648	5214	33121
$\langle n_u + n_{\bar{u}} \rangle$	566	1146	910	3554
$\langle n_d + n_{\bar{d}} \rangle$	506	1064	822	3318
$\langle n_s + n_{\bar{s}} \rangle$	394	614	524	1930
$\langle n_c + n_{\bar{c}} \rangle$	56	166	120	532
final hadrons				
$\langle n_{ch} \rangle$		6660		21420
$\langle n_{ch} \rangle / \langle n_{hadron} \rangle$		0.58		0.60
$\langle p_{\perp ch} \rangle$ (GeV)		0.39		0.54

Table 4

Fig. 1

a)

b)

Fig. 2

a)

b)

Fig. 3

Fig. 4

Fig. 5

a)

b)

Fig. 6

c)

Fig. 6 cont'd

Fig. 7

Fig. 8

Fig. 9

a)

b)

c)

Fig. 10

a)

b)

Fig. 11

c)

Fig. 11 cont'd

Fig. 12

$$I_{a_i}^{(k)} = I_{a_i}^{(k) \text{ gain}} - I_{a_i}^{(k) \text{ loss}} =$$

Fig. 13

Fig. 14

Fig. 15

Fig. 16

Initial State

SCALE: Q_0^2

Parton Cascade Development

SCALE: $[Q_0^2, \mu_0^2]$

Hadronization

Cluster formation

SCALE: $[\mu_0^2, \Lambda^2]$

Cluster decay

SCALE: Λ^2

Fig. 18

Fig. 19

Fig. 20

Fig. 21

Fig. 22

Fig. 23

Fig. 24

Fig. 25

Fig. 26

Fig. 27

Fig. 28

Fig. 29

Fig. 30

Fig. 31

Fig. 32

Fig. 33

Fig. 34

Fig. 35

Fig. 36

a)

b)

Fig. 37

Fig. 38

Fig. 39

a)

b)

Fig. 40

a) p+p (b=0) at $s^{1/2}=200$ GeV

$\tau = 0.1$ fm

$\tau = 0.4$ fm

b) Au+Au (b=0) at $s^{1/2}=200$ A GeV

$\tau = 0.2$ fm

$\tau = 2.4$ fm

Fig. 41

Fig. 42

Fig. 43

Fig. 44

Fig. 45

Fig. 46

Fig. 47

Fig. 48

Parton distributions (r_T) in Au+Au ($b=0$), $s^{1/2} = 200$ A GeV

Fig. 49

Vertex distribution of parton production

Fig. 50

Fig. 51

Central Au+Au at $s^{1/2} = 200$ A GeV

Fig. 52

Fig. 53

Fig. 54

Fig. 55

Fig. 56

Estimates for RHIC and LHC

	$T(\tau_{th})$ [MeV]	$\varepsilon(\tau_{th})$ [GeV/fm ³]	$T(\tau_g)$ [MeV]	$\varepsilon(\tau_g)$ [GeV/fm ³]
RHIC	700	180	300	15
LHC	950	5400	340	30

Fig. 57

a)

b)

Fig. 58

c)

Fig. 58 cont'd

$p+p \rightarrow \mu^+ \mu^- + X, s^{1/2} = 200 \text{ GeV}$

Fig. 59

Contributions of interaction types

Au+Au ($b=0$), $s^{1/2} = 200$ A GeV

Fig. 60

$A+A \rightarrow \mu^+ \mu^- + X, s^{1/2} = 200 \text{ GeV}$

Fig. 61

Fig. 62

Fig. 63

Fig. 64

Au+Au with QGP Au+Au w/o QGP

Fig. 65

Fig. 66

Fig. 67

Fig. 68