COLLÈGE DE FRANCE
Laboratoire de Physique Corpusculaire

AC 1530

SW 9435

LPC 94 31

MESON RADIATIVE DECAYS AND ANOMALY PHYSICS,
A TEST OF QCD

M. Benayoun, Ph. Leruste, J.L. Narjoux
Collège de France, Paris

L. Montanet
CERN, Genève

Meson Radiative Decays and Anomaly Physics, a Test of QCD

M. Benayoun¹, Ph. Leruste¹, L. Montanet², J.-L. Narjoux¹,

¹ Laboratoire de Physique Corpusculaire, Collège de France IN2P3-CNRS, Paris, France

² CERN, European Organisation for Nuclear Research, Geneva, Switzerland

June 30, 1994

Abstract : We discuss in a wider context the connection of the results previously obtained on the decay processes $\eta/\eta' \rightarrow \pi^+\pi^-\gamma$ with respect to the ξ test of QCD proposed by Chanowitz. We show that one can add consistently to the four Chanowitz equations an equation obtained by Novikov *et al.* Akhoury and Frère which involves J/ψ decays. We show that the formerly defined model M_2 of the ρ mass distribution accommodates easily this new constraint, while model M_1 indicates an inconsistency with η data. We study the consequences of both M_1 and M_2 in the description of $\eta/\eta' \rightarrow \mu^+\mu^-\gamma$ and $\eta/\eta' \rightarrow e^+e^-\gamma$ decays and we find a good agreement with the existing measurements. Consistency of the procedure we propose is illustrated by deriving from $\eta/\eta' \rightarrow \pi^+\pi^-\gamma$ and $\eta/\eta' \rightarrow \mu^+\mu^-\gamma$ decays the branching fractions $\eta/\eta' \rightarrow \gamma\gamma$. We finally examine the consistency of the results obtained using the equations on box and triangle anomalies with the decays $V \rightarrow P\gamma$ and $P \rightarrow V\gamma$, where P and V are the usual pseudoscalar and vector mesons. Using standard SU(3) relations among coupling constants, we find consistency of our previous results with most of these radiative decay modes. The set of meson radiative decays thus addressed consists of 21 processes. As a result of all these checks, we improve the picture provided by these two models, one (M_1) in agreement with QCD, one (M_2) in basic disagreement with it. Ways to go beyond this confirmed ambiguity are proposed.

Keywords : Decay, photon-photon interactions, rho-765 resonances, eta mesons, eta-958 resonances, quarks.

Free keywords : ρ parameters, ρ decay width, box anomaly, triangle anomaly, meson radiative decays, electric charge of quarks.

Πολλὰ μορφὰ τῶν δαιμονίων.
πολλὰ δ' ἄέλπτως κραινόνουσιν θεοί.
καὶ τὰ δοκηθέντ' οὐκ ἔτελέσθη,
τῶν δ' ἄδοκῆτων πόρον ἡῶρε θεός.

Andromache, EURIPIDES

1 Introduction

In a recent paper[1] all available data concerning the decays $\eta/\eta' \rightarrow \pi^+\pi^-\gamma$ have been analyzed in the context of anomaly physics predicted by PCAC[2, 3, 4] and by the Wess-Zumino chiral lagrangian[5, 6, 7]. Beside the classical triangle anomaly[8, 9] responsible for $\pi^0/\eta/\eta'$ decays to two photons, these theories indeed predict the box anomaly[2, 3].

The existence of box anomalies has first been studied by looking for a $\gamma\pi^0\pi^+\pi^-$ vertex in some processes. This search was first successfully performed with the process[10] $e^-\pi^-\pi^0 \rightarrow e^-\pi^-\pi^0$. The same search was also done using the production process of a pion pair in the nuclear Coulomb field[11]. Despite their poor statistics, these two collaborations could conclude to the existence of the π^0 box anomaly, and found consistency with the value of f_π deduced from pion decay (93 MeV) with three colors for quarks. More precisely ref.[11] was able to show that the value found for f_π ($\simeq 84$ MeV) is in fair agreement with the value deduced from π^0 decay.

In ref.[1] (hereafter referred to as paper A) it has been shown that the box anomaly provides a simple and coherent framework to explain the features previously noticed[13]¹ in the $\eta' \rightarrow \pi^+\pi^-\gamma$ decay spectrum. Specific features of η/η' decays allow also for a determination of the pseudoscalar meson nonet parameters by means of the Chanowitz equations[2, 3, 4]. The values found in ref.[1] for the mixing angle θ_{PS} and for f_8 are in agreement with expectations[7, 12]. All these results underline the relevance of the box anomaly in $\eta/\eta'/\pi^0 \rightarrow \pi^+\pi^-\gamma$ processes as expected from chiral theories.

On the other hand, the Chanowitz equations allow for a test of QCD. In these equations a parameter[2, 3, 4], ξ , has to be equal to 1 if QCD holds whereas it can be equal to 2 if quarks carry integral charges (ICQ). Surprisingly, the present experimental data do not favor unambiguously QCD[1], in the sense that $\xi = 1$ is neither the single solution nor the simplest. The ambiguity can be traced back to the interplay of the ρ^0 meson. In paper A, the ρ lineshape was extracted from data[14] on $e^+e^- \rightarrow \pi^+\pi^-$ from threshold to 1 GeV/c. Assuming that this mass interval must be described with only the ρ^0 and ω contributions leads unavoidably to a contradiction with QCD when analyzing η/η' decays, whereas assuming the existence of an additional non-resonant $\gamma\pi^+\pi^-$ coupling allows to reach agreement. Looking at the values obtained for f_8 , f_1 and θ_{PS} does not permit a choice between the two sets of solutions, since in both cases the results are in agreement with all expectations from SU(3), the quark model and

---

¹The full list of references on this subject can be found in ref.[1].

chiral theories. As the mass range used to determine the ρ lineshape from the $e^+e^- \rightarrow \pi^+\pi^-$ data is the same as the one used to study η/η' decays, the validity of the procedure is not affected by a questionable analytic continuation outside the fitted mass range.

On the other hand, the fact that the anomalous and the ρ^0 contributions were considered as independent processes and added up coherently[1] might look arbitrary. Actually, this way to proceed is not new ; it has been proposed long ago[15] to treat the π^0 box anomaly and proved to work[10][11] in the less crucial case of the anomalous $\gamma\pi^0\pi^+\pi^-$ vertex. The same idea has been proposed recently[16] in the context of the $\eta \rightarrow \pi^+\pi^-\gamma$ decay, however with a theoretical shape for the ρ mass distribution ; the partial width deduced in this way agrees well with the corresponding data[12]. In the context of pseudoscalar meson radiative decays into a lepton pair, ref.[17] has shown that such a procedure was self-consistent as long as the resonance contribution goes to zero at zero dilepton mass.

In a recent paper[18] (see also ref.[19] and previous references therein), the question of the η - η' mixing has been revisited within the framework of a chiral theory appreciably different from the standard one[4, 5, 6]. Using as input the $\eta/\eta' \rightarrow \gamma\gamma$ and $J/\psi \rightarrow \gamma$ (η/η') data, the authors of ref.[18] have been able to compute the box anomaly constants². We shall discuss later on their results, however let us notice here that their estimates (and the consequences for QCD) coincide with our previous results[1] in a remarkable way. Since their method is intrinsically not sensitive to any ρ mass contribution, their results support the additive method we have used.

The purpose of this paper is to examine the behaviour of two kinds of ρ^0 models : M_1 which implies $\xi = 1$ (QCD-like) and M_2 (in basic disagreement with QCD since it implies $\xi = 2$) when adding further theoretical constraints and check their consequences on more experimental data.

After having briefly addressed the question of the ρ^0 modeling (section 2), we present updated values for the results obtained in paper A (section 3) and prove their stability when changing the minimization procedures. In section 4, we compare these results to those obtained using in addition to the $\eta/\eta' \rightarrow \gamma\gamma$ formulae, a relation involving J/ψ decays due to Novikov *et al.*[20], Akhoury and Frère[21]. This allows us to show that our treatment of the ρ meson mass distribution is supported by a new independent theoretical constraint and further experimental data on J/ψ decays.

In section 5, we study the consequences of adding this last equation to the set of Chanowitz equations within the fitting procedure ; the agreement with M_2 is found nearly perfect while the agreement with M_1 looks more questionable.

In section 6, we extend our analysis method to the $\eta/\eta' \rightarrow \mu^+\mu^-\gamma$ decays. This allows us to derive from there the branching fractions $\text{Br}(\eta/\eta' \rightarrow \gamma\gamma)$. We also show that our model does not suffer from possible double counting effects.

Using SU(3) relations, we study the consequences of the results obtained with the models M_1 and M_2 on the standard radiative decays $V \rightarrow P\gamma$ and $P \rightarrow V\gamma$, where P and V stand for any of the usual 0^{-+} and 1^{--} light flavour mesons (7 and 8). We also correct here for an error in paper A which affects the connection between the decay process $\rho^0 \rightarrow \eta\gamma$ and $\eta \rightarrow \pi^+\pi^-\gamma$.

---

²In fact, these box anomaly constants contain contributions from triangle anomalies ; we thank V. Petrov for this information. By convention we shall nevertheless continue to call this phase space contribution "box anomaly" as it represents the main part of it.

In section 9, we discuss the relations between the partial results obtained in the previous sections. Section 10 is devoted to a brief summary of our conclusions.

In the appendix, we recall and/or derive model properties of η/η' decays and fix notations of relevance for meson radiative decays.

2 On the Modeling of the ρ^0 Contributions

The $\eta/\eta' \rightarrow \pi^+\pi^-\gamma$ as well as $\eta/\eta' \rightarrow \mu^+\mu^-\gamma$ decays are all heavily dominated by the ρ meson contribution. Since we are interested in determining the numerical value of an expected anomalous term which comes together with the ρ^0 contribution, the precise shape of this last term has to be well defined and justified.

For the time being, the only data sets which allow for an accurate determination of the ρ^0 mass distribution are the measurements[14] of the cross section $e^+e^- \rightarrow \pi^+\pi^-$. In this case, the production mechanism is indeed the best known since it relies on QED and VDM.

In fitting these data[1], two assumptions play a role. The former is whether there exists a non-VDM $\gamma \rightarrow \pi^+\pi^-$ coupling. Such a coupling would change the mass distribution attributed to the ρ^0 meson. In this respect two extreme figures have been presented in paper A with models M₁ and M₂.

The other assumption which could be questioned here is the constancy with respect to the mass of $g_{\rho\gamma}$, the $\gamma - \rho$ transition coupling. We assume this condition[22] ; moreover, the Neutral Detector collaboration[23] recently claimed that there is no evidence for a mass dependence of this parameter. In addition, calculations of finite width corrections[24] to the pion form factor give small contributions ; this may be taken as an indirect support in favor of the constancy of $g_{\rho\gamma}$. Data on $e^+e^- \rightarrow \mu^+\mu^-$ from threshold up to the $\phi(1020)$ mass would be very welcome in order to solve this problem³.

Information about the ρ meson can be obtained from other sources, however none of them can be as conclusive as a simultaneous fit of the $e^+e^- \rightarrow \mu^+\mu^-$ and $e^+e^- \rightarrow \pi^+\pi^-$ cross sections in the ρ mass region. Indeed, information from the $\pi N \rightarrow \pi\pi N$ cross section[25] is hardly model dependent because of the large $a_1(1^{++})$ contribution which takes place besides the one pion exchange mode. It is also to some extent the case of the $J/\psi \rightarrow \pi^+\pi^-\pi^0$ decay mode since the form factor at the $(J/\psi)\rho\pi$ vertex is unknown.

The $\tau^\pm \rightarrow \pi^\pm\pi^0\nu_\tau$ decay is surely useful but will not provide new information as compared to $e^+e^- \rightarrow \pi^+\pi^-$ because of CVC. This has been illustrated by the ARGUS collaboration[26]. Indeed, CVC transfers from e^+e^- annihilation to the τ decay, not only the ρ but also a possible $\gamma\pi\pi$ direct coupling.

We shall use in the following the models for the ρ invariant mass distribution defined in paper A. In this paper, model M₂ was derived assuming that the cross section for $e^+e^- \rightarrow \pi^+\pi^-$ below 1 GeV/c can be completely described assuming only the contributions of the ρ^0 and ω mesons. Model M₁ was derived assuming in addition the existence of a small $\gamma \rightarrow \pi^+\pi^-$ non-VDM contribution. The motivation for this assumption was the high mass value of the ρ^0 meson (780.8 MeV) derived from model M₂. Imposing to the ρ^0 mass the world average value (768.7

---

³It should be noticed that such data have never been published up to now, except for the old result of ACO reproduced in ref.[24] which covers less than a ± 10 MeV interval around the ϕ mass.

MeV), the non-VDM coupling strength had to take the form[1] $A(m^2) = -(c_0 + c_2 m^2)$ in order to get an acceptable χ^2 .

Model M_1 has however the undesirable feature with respect to the theory[27] as well as to the phenomenology[28] to require a mass dependence of the non-VDM contribution in order to recover the usual mass of the ρ meson. Therefore, it is interesting to discuss also the results obtained assuming the existence of a small non-VDM contribution to the pion electromagnetic form factor, but choosing it constant ($A = c_0$) and releasing any condition on the ρ^0 mass. Such a model supposes that the difference observed in fitting the ρ spectrum in e^+e^- annihilation with M_1 and M_2 has to be considered physically meaningful : as discussed in paper A, this is far from being sure. However, as an attempt, this new model (which will be named M_3) is interesting to analyze. The fit of the $e^+e^- \rightarrow \pi^+\pi^-$ data[14] using model M_3 has been done using rel. (A.20). The results are displayed in table 1 where we also recall the results of the fit done with models M_1 and M_2 in paper A. The fit using M_3 has an excellent probability ($\chi^2/dof = 61/77$) and the fitted curve to $e^+e^- \rightarrow \pi^+\pi^-$ (not shown) looks undistinguishable of model M_1 (see fig. (4) in paper A). The $\rho^0 - \omega$ mixing angle φ is in the range expected from theory[24]. The damping power[1] λ , which governs the fall-off of the width $\Gamma_\rho(m)$ as a function of m :

$$\Gamma_\rho(m) = \Gamma_\rho(m_\rho) \left[\frac{q_\pi(m)}{q_\pi(m_\rho)} \right]^3 \left[\frac{m_\rho}{m} \right]^\lambda \quad (1)$$

is found intermediate between models M_1 and M_2 . The constant c_0 is found small as expected from theory[27] but significantly non-zero. Finally the ρ^0 mass is found in the expected range but the width is significantly below the commonly accepted value[12]. There is indeed a correlation between the mass and the width of the ρ meson : within its accepted range (between 768 and 781 MeV), the larger the mass, the larger is the width (between 142 and 153 MeV). To summarize, solution M_3 looks *a priori* acceptable.

We shall not extend our discussion on the ρ meson further on ; most other remarks can be found in paper A. Let us only notice that any model for the ρ meson can be labelled in our approach by its ρ mass. Models M_1 and M_2 are extreme pictures in the sense that the ρ mass cannot be much greater than 780.8 MeV (it is actually bounded by the ω mass) and is unlikely to be significantly below the presently accepted world average value (768.7 MeV). Therefore, by means of models M_1 , M_2 and M_3 , we are performing a kind of rough scanning of the opened possibilities.

3 Update of the Results from $\eta/\eta' \rightarrow \pi^+\pi^-\gamma$

As explained in paper A, no qualitative improvement of the numerical results on η/η' box anomalies can be expected before further and better data on the dipion spectra in the decays $\eta/\eta' \rightarrow \pi^+\pi^-\gamma$ or on the ρ meson mass distribution are available.

Nevertheless, we have been able to improve the algorithms used to fit the available data⁴. The fitting procedure used in paper A was the following : from MINUIT a value was chosen for E_X ($X = \eta, \eta'$)– the box anomaly constants in rel. (A.11)–, then the corresponding value of

---

⁴They represent nine sets of data for the η' meson and only one for the η meson. They are all referred to and discussed in paper A.

F_X —the coupling strength at the $X\rho\gamma$ vertex, see rel. (A.4)— was algebraically derived using the (known) branching fraction $X \rightarrow \pi^+\pi^-\gamma$ and its analytical expression within our model (see paper A for more details) : this procedure was repeated at each step of the minimization procedure up to convergence.

The minimization procedure can be significantly improved by noting (see rel. (A.14) in the appendix) that the suitable variable when fitting the dipion mass spectra is E_X/F_X (or its inverse) as the published experimental spectra are not normalized in absolute magnitude and because we assume the F_X well approximated by constants. Indeed, using this variable the minimization procedure has not to be interrupted at each step to compute F_X . Instead, using the final value of E_X/F_X , the value of F_X is algebraically computed from rel. (A.15), up to an unavoidable sign ambiguity.

This procedure has produced the following results for the η spectrum :

$$\left\{ \begin{array}{ll} E_\eta/F_\eta = -10.64_{-2.07}^{+2.49} & (\text{model } M_1) \\ E_\eta/F_\eta = +13.48_{-3.25}^{+3.80} & (\text{model } M_2) \\ E_\eta/F_\eta = +2.24_{-2.74}^{+3.31} & (\text{model } M_3) \end{array} \right. . \quad (2)$$

The same procedure has been applied to the η' spectra after weighting each spectrum as explained in paper A⁵. The results are :

$$\left\{ \begin{array}{ll} E_{\eta'}/F_{\eta'} = -12.38_{-1.08}^{+1.09} & (\text{model } M_1) \\ E_{\eta'}/F_{\eta'} = -5.03_{-1.08}^{+1.10} & (\text{model } M_2) \\ E_{\eta'}/F_{\eta'} = -7.68_{-1.23}^{+1.24} & (\text{model } M_3) \end{array} \right. . \quad (3)$$

The quality of fits for models M_1 and M_2 is identical to that found in paper A. The χ^2/dof obtained using model M_3 is 9/14 for the η spectrum and 92/102 for the simultaneous fit of the weighted η' spectra.

It is clear from rels. (2) and (3) that the fitting procedure does not provide signs for E_X and F_X separately. Moreover, only the relative sign of E_η/F_η and $E_{\eta'}/F_{\eta'}$ for each of the models M_1 to M_3 is significant since the sign of the vertex function $G_\rho(m)$ (see paper A) associated with the $\rho^0 \rightarrow \pi^+\pi^-$ transition is arbitrarily chosen positive.

Using rel. (A.15), the relevant partial widths[12] and rels. (2) and (3) given above, we can extract F_η and $F_{\eta'}$ for each model. Using rels. (2) and (3), this allows in turn to compute E_η and $E_{\eta'}$ in absolute magnitude for each model. The box anomaly constants are then found to be :

$$\left\{ \begin{array}{ll} E_\eta = -4.39_{-1.00}^{+1.15} , & E_{\eta'} = -5.06_{-0.54}^{+0.53} & (\text{model } M_1) \\ E_\eta = -3.64_{-0.97}^{+1.10} , & E_{\eta'} = -2.17_{-0.49}^{+0.46} & (\text{model } M_2) \\ E_\eta = -0.72_{-0.88}^{+1.06} , & E_{\eta'} = -3.16_{-0.54}^{+0.54} & (\text{model } M_3) \end{array} \right. \quad (4)$$

---

⁵We have determined the weighting factors associated with model M_3 ; they are comparable to the corresponding factors found for models M_1 and M_2 . They are not given here.

and the corresponding values of F_X are :

$$\begin{cases} F_\eta = +0.41 \pm 0.05 & , & F_{\eta'} = +0.41 \pm 0.03 & \text{(model M}_1\text{)} \\ F_\eta = -0.27 \pm 0.03 & , & F_{\eta'} = +0.43 \pm 0.03 & \text{(model M}_2\text{)} \\ F_\eta = -0.32 \pm 0.04 & , & F_{\eta'} = +0.41 \pm 0.03 & \text{(model M}_3\text{)} \end{cases} \quad (5)$$

where the E_X are expressed in units of GeV^{-3} and the F_X in units of GeV^{-1} . The signs quoted here represent a choice which will be justified just below.

The results (4) for models M_1 and M_2 remain stable with respect to paper A. One also observes that model M_3 – which follows from a likely physics in e^+e^- annihilation – has the very peculiar feature that E_η is consistent with zero, while $E_{\eta'}$ is significantly non-zero.

Following paper A, we can solve simultaneously the Chanowitz equations (see rels. (A.16) and (A.17) in the appendix) by fixing the value of ξ at 1 or 2. Then we have four equations and three unknowns after setting $f_\pi = 93$ MeV. Assuming that the signs of B_η and $B_{\eta'}$ (the triangle anomaly constants derived using rel. (A.2)) are negative, the only case where we get good probabilities in solving the Chanowitz equations is when E_η and $E_{\eta'}$ are both negative too. The corresponding results are given in table 2. The other sign assignments for E_X always lead to unacceptable χ^2 whichever is the value chosen for ξ .

For model M_1 or M_2 , the updated results do not substantially differ from those in paper A : the values for f_8 and θ_{PS} are still in fair agreement with expectations from chiral theory estimates[7] or from SU(3) mass formulae. Model M_1 is in agreement with QCD while model M_2 still disagrees with it. Model M_3 exhibits unexpected values for the parameters. Presently, its main interest is to show that a ρ^0 mass in the range $\simeq 772 \div 781$ MeV is inconsistent with QCD.

Model M_3 also proves that the existence of a non-VDM $\gamma\pi^+\pi^-$ coupling is not sufficient to recover an agreement with $\xi = 1$, the most stringent prediction of QCD. One has in addition to accept either that this term is mass dependent (with a ρ mass close to the commonly accepted value 768.7 MeV) or to consider the mass dependent part of the non-VDM coupling as due to higher resonances. We shall return to this question later on.

4 Comparison with other Results and Methods

In a recent paper[18], the problem of the value of ξ and of the anomaly constants E_η and $E_{\eta'}$ has been addressed using a different starting point and new $\eta - \eta'$ mixing schemes[19]. Namely, following previous studies and relying on an unusual lagrangian (see ref.[19] and references quoted therein), Kisselev and Petrov[18] take into account the decay widths $\eta/\eta' \rightarrow \gamma\gamma$, $J/\psi \rightarrow \eta\gamma$ and $J/\psi \rightarrow \eta'\gamma$. In order to perform their analysis, they use new expressions[18, 19] for the triangle anomaly constants B_X (modified forms of rels. (A.16)) and a relation of Novikov *et al.*[20] written in a suitable way by Akhoury and Frère[21] involving J/ψ decays (a version of rel. (A.19) in the appendix modified in agreement with their lagrangian). We recall their main results in the first column in tables 3 and 4 for $\xi = 1$ and $\xi = 2$ respectively. Even fixing the value of ξ , this method does not provide results constrained by a fit as it uses three equations to extract the value of three unknowns. This method may however be preferred to the method used in ref.[29] which uses a value of f_8 deduced from chiral models to get f_1 and θ_{PS} .

The main point concerns the observable quantities E_η and $E_{\eta'}$. In both cases ($\xi = 1$ or 2) the results obtained in ref.[18] agree within errors with the values we have obtained in paper A (see also rels. (4)) for E_η and $E_{\eta'}$. Assuming $\xi = 2$ they get a sign opposite to that coming out when using the Chanowitz equations. They concluded herefrom that $\xi = 2$ is inconsistent with either of their mixing schemes. This is not correct as the fit does not fix the sign of the box anomaly constants, so that one must conclude to an agreement with our results in both cases ($\xi = 1, 2$) for the value of $E_{\eta'}$; the single difference with our approach is that the relative sign between F_η and $F_{\eta'}$ given in rels. (5) is correct in the case of Chanowitz equations (they carry unlike signs) while it should be changed to like signs in the less usual lagrangian approach of Kisselev and Petrov. This sign, which only affects the case $\xi = 2$, is subject to tests with other experimental data, namely radiative decays of light flavour mesons.

We were not aware of the relation of Novikov *et al.*-Akhoury-Frère[20, 21] (hereafter referred to as NAF equation) at the time of paper A. One may indeed solve simultaneously the two Chanowitz equations for the triangle anomalies and the NAF equation in the standard mixing scheme (rels. (A.16) and (A.19) without any modification). In this case we have three equations and three unknowns (ξ is fixed). The pseudoscalar nonet parameters being computed from eqs. (A.16) and (A.19), the (predicted) values of E_η and $E_{\eta'}$ can be derived using eqs. (A.17). All results are displayed in the second column of tables 3 and 4.

The values obtained for $E_{\eta'}$ agree with our fitted values (given in the third data column of tables 3 and 4). Moreover, these equations provide the usual sign as compared to those obtained using the four Chanowitz equations (rels. (A.16) and (A.17)). This conclusion is valid for both ξ . The agreement for E_η is within 1σ when fixing $\xi = 2$, while it is slightly above 2σ when fixing $\xi = 1$.

Several comments can be made on the results presented in tables 3 and 4. For simplicity, we shall limit our comments to the method using the NAF equation and to the method relying on the simultaneous use of the four Chanowitz equations.

i/ The agreement found for both E_η and $E_{\eta'}$ using both values of ξ is of particular interest. Indeed, using only the decay widths $\eta/\eta' \rightarrow \gamma\gamma$ and the radiative J/ψ decays to η/η' , the results (column 2 in tables 3 and 4) are insensitive to any ρ contribution. The agreement between the results obtained in this way and ours (column 3) illustrates that our treatment of the ρ meson in $\eta/\eta' \rightarrow \pi^+\pi^-\gamma$ decays is basically correct.

ii/ Our results rely on nine sets of data for the η' decay (see paper A for references) and on only one set for the η decay[30]. The fair agreement observed between the two methods may be taken as an evidence in favor of the good quality of this unique set of η decay.

iii/ The consistency of the results obtained with these two methods suggests that the NAF equation could be added to the set of the four Chanowitz equations. This would increase the constraints (five equations relating four or three -if we fix ξ - parameters to be fitted). We present the results obtained in this way in the next section.

iv/ This comparison seems to rule out model M_3 since the value for E_η (see rels. (4)) looks inconsistent with Petrov *et al.* and with the NAF equation. Therefore, the correct ρ^0 mass looks to be either close to that of M_1 or close to that of M_2 .

5 Including the NAF Equation

The above results allow us to consider that the Chanowitz and NAF equations are consistent with each other. This assumption underlies previous uses[18, 19, 21] of the relations for $\eta/\eta' \rightarrow \gamma\gamma$ and of the NAF equation, sometimes in a simplified form[31]. Therefore we have used simultaneously eqs. (A.16), (A.17) and (A.19) and replaced their left-hand sides with the world average values[12] of the data on $\eta/\eta' \rightarrow \gamma\gamma$ and $J/\psi \rightarrow (\eta/\eta') \gamma$ decays. As data on the decay processes $\eta/\eta' \rightarrow \pi^+\pi^-\gamma$, we have used the anomalous contributions given in rels. (3) above for each model M_1 , M_2 , M_3 .

Model M_3 is rejected with $\chi^2/dof = 16/1$, pointing to the fact that any model intermediate between M_1 and M_2 looks disfavored. In this case we have not assigned a value to ξ .

Using model M_2 , we obtain a good fit ($\chi^2/dof = 0.87/2$). The mixing angle obtained in this case ($\theta_{PS} = -23.12^\circ$) is in perfect agreement with the value obtained from SU(3) using the linear mass formula[12].

Model M_1 is less favored ($\chi^2/dof = 5.56/2$) than M_2 , but the probability is still acceptable (6%). The mixing angle is found near -16° .

The origin of the problem for M_1 is easy to trace back by looking at table 3 : the discrepancy between the value predicted for E_η and the value obtained from the fit is $\simeq 2.5\sigma$. This points towards an inconsistency with some η data. In order to check this hypothesis, we have removed from the fit the first of the equations (A.17). Then the fit yields as expected a good result ($\chi^2/dof = 0.34/1$) without any substantial change to the parameter values. All these results are collected in table 5.

Having five equations it is now possible to leave also ξ free. One thus finds :

$$\begin{cases} \xi = 1.11^{+0.11}_{-0.09} & \chi^2/dof = 4.09/1 \quad (\text{model } M_1) \\ \xi = 2.06^{+0.37}_{-0.27} & \chi^2/dof = 0.83/1 \quad (\text{model } M_2) \end{cases} \quad (6)$$

Therefore we can conclude to an agreement of our results for E_η and $E_{\eta'}$ obtained using model M_2 with all existing data on the $\eta/\eta' \rightarrow \gamma\gamma$ and $J/\psi \rightarrow (\eta/\eta') \gamma$ decay modes. Instead, model M_1 would point towards a discrepancy with our result on $\eta \rightarrow \pi^+\pi^-\gamma$ decay. As these data are unique[30], no cross check can be performed for the time being on the validity of this result. Let us note however that the value predicted for E_η (see column 2 in table 3) is not unlikely : it is the second solution found by consistency in paper A (see rels. (31) in paper A and the discussion here). This solution has been discarded because of the result obtained with the data set of ref.[30].

6 Fits of $\eta/\eta' \rightarrow \mu^+\mu^-\gamma$ Decay Modes and Spectra

6.1 Motivations and Principle of the Model

The basic idea of the model developed in paper A in order to describe the $\eta/\eta' \rightarrow \pi^+\pi^-\gamma$ decays was to split up the amplitude into the resonant and the non-resonant (anomalous) parts.

The vertex function of the anomalous part was considered to be well approximated in the physical region by its value at the chiral point ; this turns out to identify the anomalous contribution to the amplitude with a pure phase space term (*i.e.* structureless), thus qualitatively different from a pure resonant ρ contribution.

This splitting is well accepted in the case of the box anomaly for π^0 [15] since it provides the expected result[10, 11]. However, in the case of model M_2 which yields a “non-QCD” value for ξ , one may have doubt on the validity of this procedure. However, if the splitting is a correct procedure, it should apply to other decay modes, namely when the triangle anomaly would occur together with the resonant ρ contribution, as in the decays $\eta/\eta' \rightarrow \mu^+\mu^-\gamma$. Moreover, the coupling strengths at the $\eta\rho\gamma$ and $\eta'\rho\gamma$ vertices should be equal to what we have named F_η and $F_{\eta'}$ respectively.

Hence, the processes $\eta/\eta' \rightarrow \mu^+\mu^-\gamma$ can be described with the two contributions depicted in fig. (1). In fig. (1a) (see also section A.1)) the non resonant term contains the triangle anomaly vertex connected with an off-shell photon ; as we have assumed the coefficients E_X nearly constant and approximated them by their value at the chiral point (see rels. (A.17)), consistency implies that the coefficients B_X at the vertex should be treated as constant and approximated by their value at the chiral point (see rels. (A.16)). Moreover, the constants B_X can be deduced from the decay widths $\Gamma(\eta/\eta' \rightarrow \gamma\gamma)$ by means of rel. (A.2).

Fig. (1b) describes the process with an intermediate ρ^0 meson. The coupling strengths at the vertices $X\rho^0\gamma$ should be the same F_X 's as in the processes $\eta/\eta' \rightarrow \pi^+\pi^-\gamma$ (see fig. (2) and especially (2b)). They are listed in rels. (5) according to the choice of the ρ model.

Instead of fixing B_X , we can rather fix F_X . In this case, the results we get from a fit to $\eta/\eta' \rightarrow \mu^+\mu^-\gamma$ allow us to derive the value of B_X and hence to “predict” the values of the triangle anomaly constants (B_η and $B_{\eta'}$) and thus $\Gamma(\eta/\eta' \rightarrow \gamma\gamma)$.

The $\rho \rightarrow \gamma$ transition has to be treated with a constant value $g_{\rho\gamma}$ as we did in paper A to fit the $e^+e^- \rightarrow \pi^+\pi^-$ data. One has however to check for full consistency that the diagram in fig (1b) vanishes at the chiral point[17] to avoid problems with counting twice some contributions[32, 33, 34]. The model we present in the appendix fulfills this condition and corresponds to the models developed in paper A for $\eta/\eta' \rightarrow \pi^+\pi^-\gamma$ decays and for the $e^+e^- \rightarrow \pi^+\pi^-$ annihilation process. A comparison of fig.(1) (which describes the $\eta/\eta' \rightarrow \mu^+\mu^-\gamma$ processes) and fig. (2) (which describes $\eta/\eta' \rightarrow \pi^+\pi^-\gamma$) clearly illustrates this correspondence.

6.2 Data Sets and Analysis Method

The processes $\eta/\eta' \rightarrow \mu^+\mu^-\gamma$ have both been measured only once[35, 36, 37, 38] ; the results can also be found in ref.[34] where they have been widely discussed, mainly in the context of electromagnetic form factor studies.

The statistics for the η decay are reasonable ($\simeq 600$ events), while they are relatively poor for the η' decay ($\simeq 30$ events). The interest in analyzing $\eta/\eta' \rightarrow \mu^+\mu^-\gamma$ rather than $\eta/\eta' \rightarrow e^+e^-\gamma$ is that the former processes are nearly insensitive to radiative corrections[34] and therefore, we can test our model without shadowing effect. The single further hypothesis we have to make is that we can continue analytically our ρ models, which have been fitted between $2m_\pi$ and m_ϕ , to the region $[2m_\mu, 2m_\pi]$. This is not a strong assumption and for what concerns the branching fractions, it is quite safe as the contribution of the $[2m_\mu, 2m_\pi]$ region to the statistics is suppressed by phase space.

Therefore, knowing the partial widths $\Gamma(\eta/\eta' \rightarrow \gamma\gamma)$ [12], the F_η and $F_{\eta'}$ given by rels. (5) and using our ρ^0 models⁶, we can predict all characteristics of the decay modes $\eta/\eta' \rightarrow \mu^+\mu^-\gamma$, without any free parameter.

6.3 Estimation of the $\eta/\eta' \rightarrow \gamma\gamma$ Branching Ratios

A first quality test of models M₁ and M₂ consists in using the branching fractions $\text{Br}(\eta/\eta' \rightarrow \mu^+\mu^-\gamma)$. Using rel. (A.8), they can be expressed as second degree polynomials in B_X . ($X = \eta, \eta'$). Then the solutions to this equation allow to compute the branching fractions $\text{Br}(\eta/\eta' \rightarrow \gamma\gamma)$ by means of rel. (A.2). The value to be used for F_X are those given in rels. (5) together with the corresponding ρ model. The branching ratios $\text{Br}(\eta/\eta' \rightarrow \mu^+\mu^-\gamma)$ being of the order $1 \div 4 \cdot 10^{-4}$, the result is a good test of our method since we shall derive from them large branching fractions ($\simeq 2\%$ and $\simeq 40\%$).

The computation has been done numerically by means of Monte Carlo methods, assuming all input parameters to be random gaussian variables having as mean value the parameter central value and as standard deviation its error. The distributions obtained for the branching fractions are appreciably asymmetric, thus we quote as central values the most probable ones ; the quoted errors have been obtained by fitting separately the upper part and the lower part of these distributions with gaussians. The two possible solutions for B_η (see rel. (A.8)) give two possible values of $\text{Br}(\eta \rightarrow \gamma\gamma)$ distant by less than 1% in absolute magnitude ; we quote the lower result only. We get for $\text{Br}(\eta \rightarrow \gamma\gamma)$ expressed in percents of the full width :

$$\begin{cases} \text{Br}(\eta \rightarrow \gamma\gamma) = (112^{+61}_{-43})\% & (\text{model M}_1) \\ \text{Br}(\eta \rightarrow \gamma\gamma) = (44^{+28}_{-14})\% & (\text{model M}_2) \end{cases} \quad (7)$$

for a reference value of [12] $38.9 \pm 0.5 \%$. Using the decay width [12] of $\rho^0 \rightarrow \eta\gamma$, we can also get another estimate⁷ of F_η using :

$$\Gamma(\rho \rightarrow \eta\gamma) = \frac{1}{96\pi} \left[\frac{m_\rho^2 - m_\eta^2}{m_\eta} \right]^3 |F_\eta|^2. \quad (8)$$

The corresponding $F_\eta = 0.566 \pm 0.053$ (which is independent of any ρ model) gives :

$$\text{Br}(\eta \rightarrow \gamma\gamma) = (206^{+87}_{-69})\%. \quad (9)$$

From these results we conclude that M₂ gives a value of $\text{Br}(\eta \rightarrow \gamma\gamma)$ in good agreement with the reference value [12]. Model M₁ gives too large a central value but is consistent with the PDG value at 1.5σ . Finally the decay rate $\rho \rightarrow \eta\gamma$ gives a value which is too large (at 2σ from the PDG value [12]).

For the branching fraction $\text{Br}(\eta' \rightarrow \gamma\gamma)$, we find :

---

⁶We shall only discuss models M₁ and M₂, leaving aside model M₃ which gives here also intermediate results.

⁷The value quoted for this parameter in paper A (rel. (39)) is wrong by a factor of 2 ; a part of the phase space term was omitted in rel. (38) here. Rel. (8) in this paper is the correct expression.

$$\begin{cases} \text{Br}(\eta' \rightarrow \gamma\gamma) = (2.1_{-0.8}^{+0.9})\% & (\text{model M}_1) \\ \text{Br}(\eta' \rightarrow \gamma\gamma) = (1.9_{-0.5}^{+1.0})\% & (\text{model M}_2) \end{cases} \quad (10)$$

to be compared with the reference value[12] $2.17 \pm 0.17 \%$. Thus for $\eta' \rightarrow \gamma\gamma$ both M_1 and M_2 give an acceptable value.

As a first conclusion from these calculations, one finds that both B_X are consistent with being constants and agree within errors with what can be inferred directly from the η/η' two-photon decays.

Instead of using the values of F_η and $F_{\eta'}$ given by rels. (5), we can assume that the B_X are given by the two-photon decay widths of the η/η' mesons⁸ and consider rels. (A.8) as second degree equations on F_X . Making use of the ρ models M_1 and M_2 , the experimental value of $\text{Br}(\eta \rightarrow \mu^+\mu^-\gamma)$ gives :

$$\begin{cases} F_\eta = -0.224 \pm 0.062 & \text{or} & 0.216 \pm 0.062 & (\text{model M}_1) \\ F_\eta = -0.224 \pm 0.062 & \text{or} & 0.220 \pm 0.062 & (\text{model M}_2) \end{cases} \quad (11)$$

On the other hand, $\text{Br}(\eta' \rightarrow \mu^+\mu^-\gamma)$ gives :

$$\begin{cases} F_{\eta'} = -0.448 \pm 0.098 & \text{or} & 0.404 \pm 0.098 & (\text{model M}_1) \\ F_{\eta'} = -0.491 \pm 0.101 & \text{or} & 0.423 \pm 0.101 & (\text{model M}_2) \end{cases} \quad (12)$$

Rel. (11) and (12) are interesting in several respects :

i/ They show a relative independence with respect to the detailed shape of the ρ mass distribution ; this simply illustrates that the integrals of $1/|D_\rho(m)|^2$ over the physical region for model M_1 and model M_2 are nearly equal.

ii/ The values found here for $F_{\eta'}$ are in agreement with the values obtained from the $\eta' \rightarrow \pi^+\pi^-\gamma$ decay (see rels. (5)), independently of the ρ model applied for the ρ lineshape.

iii/ The values found for F_η are always close to those obtained using model M_2 in $\eta \rightarrow \pi^+\pi^-\gamma$ decay (see rels. (5)). This confirms that M_1 has difficulties to accomodate η data.

The connection between the results in rels. (11) and (12) depends on the model. For model M_2 , F_η and $F_{\eta'}$ must carry unlike signs, while for model M_1 they should carry like signs (see rels. (5)). There are therefore two possibilities for each of models M_1 and M_2 :

$$\begin{cases} F_\eta = -0.224 \pm 0.062 & F_{\eta'} = -0.448 \pm 0.098 & (\text{model M}_1) \\ & \text{or} & \\ F_\eta = +0.216 \pm 0.062 & F_{\eta'} = +0.404 \pm 0.098 & (\text{model M}_1) \end{cases} \quad (13)$$

and :

---

⁸With a negative sign for consistency with the Chanowitz equations.

$$\left\{ \begin{array}{ll} F_\eta = +0.220 \pm 0.062 & F_{\eta'} = -0.491 \pm 0.101 \quad (\text{model M}_2) \\ & \text{or} \\ F_\eta = -0.224 \pm 0.062 & F_{\eta'} = +0.423 \pm 0.101 \quad (\text{model M}_2) \end{array} \right. \quad (14)$$

These sets are, in principle, directly comparable to the corresponding sets in rels. (5). The second solutions given in rels. (13) and (14) are clearly the preferred ones. To summarize these results, we can say that the values for $F_{\eta'}$ are inconclusive since both M_1 and M_2 predict the correct value for this parameter. The values for F_η however favor model M_2 over model M_1 . This discrepancy about model M_1 confirms the problem already mentioned with respect to $\eta \rightarrow \pi^+\pi^-\gamma$ data which arises when one takes into account the NAF equation.

Thus, $\eta \rightarrow \pi^+\pi^-\gamma$ and $\eta \rightarrow \mu^+\mu^-\gamma$ data are conflicting with the NAF equation in case of model M_1 . They are not in the case of model M_2 .

6.4 The η/η' Transition Form Factors

In order to check whether the results about F_η and $F_{\eta'}$ derived from the η/η' branching fractions are not numerical accidents, we have also looked at their transition form factors[34, 35, 36, 37, 38]. As already mentioned, there exists only one measurement for each of the $\eta/\eta' \rightarrow \mu^+\mu^-\gamma$ form factors. The η sample is relatively large ($\simeq 600$ events), the η' sample is much smaller ($\simeq 30$ events).

The model developed in the appendix allows for a one parameter (F_η/B_η) fit to $|\mathcal{F}_\eta(m_{\mu\mu}, 0)|^2$ and a one parameter ($F_{\eta'}/B_{\eta'}$) fit to $|\mathcal{F}_{\eta'}(m_{\mu\mu}, 0)|^2$. However, using rels. (5) and the two photon decay widths of η and η' , we can also fix these parameters and get the functions $|\mathcal{F}_\eta(m_{\mu\mu}, 0)|^2$ and $|\mathcal{F}_{\eta'}(m_{\mu\mu}, 0)|^2$ predicted using the $\eta/\eta' \rightarrow \gamma\gamma$ data and the $\eta/\eta' \rightarrow \pi^+\pi^-\gamma$ results. There only remains a sign ambiguity which originates from the fact that we do not know the relative sign of $g_{\rho\gamma}$ (the $\rho \rightarrow \gamma$ transition amplitude) and $G_\rho(m)$ (the $\rho^0 \rightarrow \pi^+\pi^-$ vertex function) over the physical region ($m \geq 2m_\pi$).

The fit of $|\mathcal{F}_\eta(m_{\mu\mu}, 0)|^2$ has been done using both ρ models and provides the results given in table 6. The quality of fit highly improves if we remove the first two points from the fitting procedure (it becomes $\chi^2/dof = 6.6/6$ for both ρ models). This is not surprising since they are both below the two pion threshold, *i.e.* in a region where our input ρ mass distributions need to be extrapolated. The values expected are also displayed in table 6. This confirms the results already inferred from the branching ratios : model M_2 data are in agreement with the shape of $|\mathcal{F}_\eta(m_{\mu\mu}, 0)|^2$. Table 6 shows that the agreement of model M_1 with expectations is not as good. However, we can find by fit a value of F_η/B_η which allows for a good description of $|\mathcal{F}_\eta(m_{\mu\mu}, 0)|^2$. In figs. (3) and (4), we show the data points together with the fitted function and with the curves obtained by fixing F_η/B_η to its value expected from rels. (2) (up to the quoted sign ambiguity). Even with no free parameter, it is clear that model M_2 gives a suitable description of the data. The continuation from $2m_\pi$ to zero is a consequence of the analytic continuation of the ρ model below the region where it has been fitted ; hence the precise shape displayed in this region is not reliable. It is displayed only in order to show that our fitting functions always go to 1 at $m = 0$. The disagreement for M_1 cannot be attributed to double

counting ; as already noticed in sections 4 and 5, this is due to the difficulty that model M_1 has to accomodate the η data. Moreover, there exists a value of F_η/B_η which provides a correct description of $|\mathcal{F}_\eta(m_{\mu\mu}, 0)|^2$ using the M_1 ρ model.

In figs. (5) and (6) we display the corresponding results for $|\mathcal{F}_{\eta'}(m_{\mu\mu}, 0)|^2$ obtained using models M_1 and M_2 with a fitted parameter and by fixing it to its value expected from rels. (3) (up the above mentioned sign ambiguity). In this case both models agree with the predictions and with the data which are rather unprecise⁹. This could be expected if one remembers that the $F_{\eta'}$ obtained from the branching ratios are consistent for each ρ model and with the $F_{\eta'}$ deduced when fitting the $\eta' \rightarrow \pi^+\pi^-\gamma$ spectra.

It should be noticed that the shape predicted by our models cannot be substantially modified, contrarily to the usual fitting procedure[34]. Having fixed the ρ mass distribution removes to a large extent any flexibility¹⁰.

The most likely way to get steeper shapes - if it were experimentally needed - would be to allow $g_{\rho\gamma}$ to be mass dependent. In this case, however, the whole procedure has to be repeated from the beginning, *i.e.* from fitting the $e^+e^- \rightarrow \pi^+\pi^-$ data[1]. There are however some claims[23] that this is not presently justified by studying several processes in e^+e^- annihilation.

Before closing this section, it is worthwhile to comment once more on the possibility of double counting when looking at fig. (3). If there were some double counting effect when dealing with model M_1 , consistency would imply that the effect would be the same in $|\mathcal{F}_\eta(m_{\mu\mu}, 0)|^2$ and in $|\mathcal{F}_{\eta'}(m_{\mu\mu}, 0)|^2$. Comparing figs. (3) and (5) clearly proves that it is not what happens. Moreover double counting effects should also occur when using model M_2 ; figs. (4) and (6) clearly contradict this possibility.

Therefore, the analysis of $\eta/\eta' \rightarrow \mu^+\mu^-\gamma$ confirms that one can consider the anomaly and the resonant contribution as independent processes provided one does not choose arbitrarily the shape of the resonant contribution. In addition, these data reinforce the conclusion reached using the NAF equation : model M_1 finally has difficulty to accomodate some η data while M_2 is consistent with all the η and η' data examined sofar.

Another remark follows from the analysis of $\eta/\eta' \rightarrow \mu^+\mu^-\gamma$ decays. We have already noted that $g_{\rho\gamma}$ can be mass dependent ; clearly, F_η and $F_{\eta'}$ could share the same property. However, the fact that we recover from $\eta/\eta' \rightarrow \mu^+\mu^-\gamma$ decays the correct triangle anomaly constants B_η and $B_{\eta'}$, suggests that the same procedure provides the correct box anomaly constants E_η and $E_{\eta'}$ from our analysis of $\eta/\eta' \rightarrow \pi^+\pi^-\gamma$ decays. This qualitative argument is confirmed when comparing figs. (1) and (2) and by the remark that the kinematical regions involved for η , η' and ρ^0 are alike in these two cases.

---

⁹The two data sets shown correspond to two different assumptions when subtracting the background[34] to the η' signal. The fitted set is the one represented by star symbols.

¹⁰It should be remarked that the basic idea behind fixing the ρ mass distribution in e^+e^- data was precisely to remove any flexibility to the shape of the ρ . In this way we were able to get the box anomaly constants which otherwise are correlated to the shape of the ρ invariant mass distribution.

7 Predictions of Branching Ratios

Using our results concerning F_η and $F_{\eta'}$, and the branching ratios of $\eta/\eta' \rightarrow \mu^+\mu^-\gamma$, we have obtained “predictions” concerning the branching ratios of $\eta/\eta' \rightarrow \gamma\gamma$ (see rels. (7) and table 7).

Conversely, using F_η , $F_{\eta'}$ and the branching ratios $\eta/\eta' \rightarrow \gamma\gamma$, we can get “predictions” for the branching ratios $\eta/\eta' \rightarrow \mu^+\mu^-\gamma$. The corresponding results are displayed in table 7 together with the reference values[12]. The values predicted with both ρ models for the $\eta' \rightarrow \mu^+\mu^-\gamma$ branching ratios are both in good agreement with the reference value[12]. M_2 looks however more precise ; this is simply a consequence of rels. (12).

Using F_η , we can also evaluate the branching ratio $\rho^0 \rightarrow \eta\gamma$ (see rels. (5) and (8)). The results are given in table 7 and stress that the experimental value reported[12, 23] for $\rho^0 \rightarrow \eta\gamma$ is not consistent with the branching ratios of $\eta \rightarrow \pi^+\pi^-\gamma$ and $\eta \rightarrow \mu^+\mu^-\gamma$. It is too large large by a factor of $2 \div 4$; this is also illustrated by rel. (9).

For the branching ratio of $\eta \rightarrow \mu^+\mu^-\gamma$, both estimations derived from models M_1 and M_2 agree once more with the reference value, however this agreement should not hide that the central values are far from each other in absolute magnitude ; this is a simple consequence of the differences observed for F_η from rels. (13) and (5). Paradoxically, the lack of precision in the estimation is due to the partial width $\eta \rightarrow \gamma\gamma$; due to a larger phase space, $B_{\eta'}$ is indeed known with an accuracy 3 times better than B_η . The errors produced by continuing the ρ^0 mass distribution outside the range where it has been fitted (namely in the $[2m_\mu, 2m_\pi]$ region) is not accounted for in table 7. These errors are likely to be small and affect only the η/η' decays.

Up to possible radiative corrections, the model developed in Appendix A1 clearly applies to $\eta/\eta' \rightarrow e^+e^-\gamma$ decays by changing the muon mass to the electron mass. There is only one rather unprecise measurement of the η branching ratio[12] while nothing is available for the η' decay. In this case however, the error produced by continuing the ρ mass distribution down to nearly zero is surely much more important than for the muon final state. Proceeding as for $\eta/\eta' \rightarrow \mu^+\mu^-\gamma$, we obtain for $\text{Br}(\eta/\eta' \rightarrow e^+e^-\gamma)$ the values given in table 7. Comparison with existing data is only possible for η and both M_1 and M_2 provide a good agreement ; we give also the predictions for the η' decay. The errors quoted here are only those due to the errors on the B_X and F_X .

Leaving aside a detailed comparison of models M_1 and M_2 predictions with actual data, one should stress that both models give quite reasonable predictions concerning η and η' decays. Since these predictions follow from the splitting assumption, we consider that the observed qualitative and quantitative agreement with measurements supports the method.

The parameters F_η and $F_{\eta'}$ used here and in paper A are identical to the usual $g_{\rho\eta\gamma}$ and $g_{\rho\eta'\gamma}$ couplings of SU(3)[24, 39] (see section 7 in the appendix). Therefore, using SU(3) symmetry relations between coupling constants, one can relate these parameters with some other radiative decays. More precisely, using F_η , $F_{\eta'}$ and our results on the pseudoscalar mixing angle θ_{PS} (see table 5), we can calculate the coupling constant[39] g . This provides predictions on the decays

$\rho \rightarrow \pi\gamma$ and $K^* \rightarrow K\gamma$ since they only depend on g (see appendix A6).

If SU(3) were not broken, one would expect strictly $\Gamma(\rho^0 \rightarrow \pi^0\gamma) = \Gamma(\rho^\pm \rightarrow \pi^\pm\gamma)$ (see rels. (A.21) and (A.26) in the appendix). Experimentally the ratio[12] of these two partial widths is 1.76 ± 0.49 . On the other hand, one also expects strictly $\Gamma(K^{*0} \rightarrow K^0\gamma) = 4\Gamma(K^{*\pm} \rightarrow K^\pm\gamma)$, while experimental data[12, 40] give a ratio for these decay modes of 2.28 ± 0.29 instead of $\simeq 4$. Therefore, one cannot expect that estimates performed using rels. (A.24) and (A.26) can provide a simultaneous agreement for both ρ and both K^* radiative decay branching ratios.

Our results together with the reference values[12] are displayed in table 7. In the case of $\rho \rightarrow \pi\gamma$ decays, M_2 and is in better agreement with the PDG values[12] than M_1 which is nevertheless consistent with both experimental measurements at the 2σ level.

The situation for K^* mesons looks more intriguing : model M_1 is in good agreement with K^{*0} data but does not agree with the $K^{*\pm}$ data, while model M_2 is in agreement with the $K^{*\pm}$ measurements and disagrees at the 2.2σ level with the K^{*0} data[40].

The result on $\rho \rightarrow \eta\gamma$ is indeed not conclusive as the experimental branching ratio[12, 23] is inconsistent with $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow \mu^+\mu^-\gamma$. It may well be that this discrepancy follows from experimental errors or inadequate modeling ; however, one cannot exclude another possibility. Indeed, if F_η were ρ mass dependent, the prediction for $\rho \rightarrow \eta\gamma$ could be different from estimations done using $\eta \rightarrow \rho\gamma$. Only new measurements and/or modeling can answer the question.

Therefore, this check does not fully permit to discriminate between M_1 and M_2 , even if M_2 is in better agreement with the data. Nevertheless, we can readily conclude that the values of F_η and $F_{\eta'}$ deduced from our treatment of the anomalies give predictions for radiative decay branching ratios within the expected order of magnitude. Knowing that M_2 is basically inconsistent with QCD, this feature has to be underlined.

We have presented in section 4, the model results of Kisselev and Petrov[18]. Fixing $\xi = 1$, we have shown that they agree with the model M_1 results on anomalies ; consequently F_η and $F_{\eta'}$ have to be taken with like signs as we did above (the corresponding results are given in table 7) and this leads to the conclusions already presented for model M_1 . However, in the case $\xi = 2$, they differ from the M_2 standard results because F_η and $F_{\eta'}$, instead of being unlike sign, should carry like signs (see the discussion in sections 4 and 5). We have examined the consequences of this change on the decay modes already quoted : for all ρ and K^* radiative decays this change of sign provides systematically too small branching fractions (at least by a factor of 5).

The results described in this section show that the relative sign of F_η and $F_{\eta'}$ predicted by our ρ models (see rels. (5)) is an observable quantity : this relative sign has to be positive in the case of model M_1 , while a negative sign in the case of model M_2 is preferred by the data. This rules out the mass mixing scheme of Kisselev and Petrov[18, 19] and favors the standard approach if $\xi = 2$.

8 Further Predictions on Radiative Decays

The results discussed above lead us to attempt a global fit of the data on radiative decays together with our own results which summarize the properties of $\eta/\eta' \rightarrow \gamma\gamma$, $\eta/\eta' \rightarrow \pi^+\pi^-\gamma$, $\eta/\eta' \rightarrow \mu^+\mu^-\gamma$ and $J/\psi \rightarrow (\eta/\eta')\gamma$. As far as we know, such an attempt has never been done.

It involves *a priori* 19 measured decay modes and includes all the corresponding branching ratios and some mass spectra.

Let us notice that our results on anomalies, provides the mixing angle θ_{PS} , F_η and $F_{\eta'}$ for each ρ model, *i.e.* the values of three parameters (g , g_1 , θ_{PS}) of the SU(3) formulation[39] : then, from additional radiative decays, we can expect in addition to determine g_1 and θ_V .

We have already seen that the radiative decays exhibit inconsistencies relative to each other when assuming exact SU(3) symmetry. In order to trace back the origin of possible further inconsistencies when merging up these data and the results we have obtained from other decays, it is useful to analyze first separately the $V \rightarrow P\gamma$ and $P \rightarrow V\gamma$ decays by themselves.

8.1 Preliminary Analysis

We have performed the analysis of the eleven decay modes (see table 8 for the list) $V \rightarrow P\gamma$ and $P \rightarrow V\gamma$ using the method[39] given in the appendix A7. We summarize here only the final conclusions. For obvious reasons, we have removed from this set the decay modes $\eta/\eta' \rightarrow \pi^+\pi^-\gamma$. Even if they are dominated by $\eta/\eta' \rightarrow \rho^0\gamma$, there is a non-resonant contribution (the anomaly) which is not accounted for in the usual approaches[29, 23].

We have first attempted a five parameter ($g, g_1, g_1', \theta_V, \theta_{PS}$) fit to the whole set of radiative decay modes. Within the formalism of ref.[39], we have found that the decay process $K^{*0} \rightarrow K^0\gamma$ always degrades the quality of the fit, whereas $K^{*\pm} \rightarrow K^\pm\gamma$ is perfectly consistent with the other decay modes. Therefore we remove $K^{*0} \rightarrow K^0\gamma$ from the set under fit.

Using the remaining set of ten radiative decay modes, we have performed a five parameter fit. In this case we get :

$$\left\{ \begin{array}{lcl} g & = & 0.721 \pm 0.018 \text{ GeV}^{-1} \\ g_1 & = & 0.645 \pm 0.020 \text{ GeV}^{-1} \\ g_1' & = & 0.760 \pm 0.028 \text{ GeV}^{-1} \\ \theta_V & = & (34.75 \pm 0.90)^\circ \\ \theta_{PS} & = & -(24.71 \pm 1.14)^\circ \end{array} \right. \quad (15)$$

with $\chi^2/dof = 14/5$. The branching fraction of $\rho^0 \rightarrow \eta\gamma$ is relatively close to that predicted using model M₁ (see table 7). However, the value of θ_{PS} is not as expected. In this case, consistency between the standard radiative decays and the anomaly physics (see tables 3 and 5) would have provided $\theta_{PS} \simeq -16^\circ$.

After several attempts we have found that removing in addition the process $\phi \rightarrow \eta\gamma$ allows for a more suitable fit :

$$\begin{cases} g = & 0.739 \pm 0.019 \text{ GeV}^{-1} \\ g_1 = & 0.672 \pm 0.023 \text{ GeV}^{-1} \\ g'_1 = & 0.637 \pm 0.060 \text{ GeV}^{-1} \\ \theta_V = & (34.42 \pm 1.06)^\circ \\ \theta_{PS} = & -(13.46^{+5.60}_{-5.40})^\circ \end{cases} \quad (16)$$

with an improved quality $\chi^2/dof = 6/4$. Removing any other process does not produce such an improvement. We thus find that two experimental results have to be discarded to reach an agreement with the anomaly physics results. For the remaining processes, it is not necessary to invoke SU(3) symmetry breaking¹¹. It remains to understand the origin of the disagreement of $K^{*0} \rightarrow K^0 \gamma$ with the other radiative decay modes and that of $\phi \rightarrow \eta \gamma$ with the results coming from $\eta/\eta' \rightarrow \gamma \gamma$ and $J/\psi \rightarrow (\eta/\eta') \gamma$ decays.

Moreover, we have obtained a good fit assuming nonet symmetry in the vector sector (*i.e.* $g \equiv g_1$ in rels. (A.24) and (A.25)). In this case we get :

$$g \equiv g_1 = 0.713 \pm 0.016 \text{ GeV}^{-1} \quad g'_1 = 0.624 \pm 0.027 \text{ GeV}^{-1} \quad \theta_V = (38.61 \pm 0.18)^\circ \quad (17)$$

with $\chi^2/dof = 8/5$ and $\theta_{PS} = (-16.24 \pm 1.16)^\circ$, in good agreement with expectations in this case. In particular, it is in agreement with a recent experimental result[41] $\theta_{PS} = (-17.3 \pm 1.8)^\circ$ which has been obtained from a systematic comparison of η and η' production in $p\bar{p}$ annihilation at rest. Using rels. (A.24) to (A.26) it is possible to reconstruct all branching ratios and the values of F_η and $F_{\eta'}$. The corresponding values are given in table 8. As expected all branching ratios are correctly reproduced except for $K^{*0} \rightarrow K^0 \gamma$ and $\phi \rightarrow \eta \gamma$ which are predicted around twice the measurements[12] also given in table 8. The value found for $F_{\eta'}$ guarantees a good description of $\eta' \rightarrow \pi^+ \pi^- \gamma$ and $\eta' \rightarrow \mu^+ \mu^- \gamma$ decays (compare with rels. (5) and (12)). The value found for F_η , however, is a little too large to provide a good agreement with $\eta \rightarrow \pi^+ \pi^- \gamma$ data and provides a poor agreement with $\eta \rightarrow \mu^+ \mu^- \gamma$ whichever model is used for the ρ meson.

The results given here and in the first data column of table 8 are the reference values to be used to study the consequences of models M₁ and M₂ on radiative decay modes. They have been obtained by means of rels. (17). In this respect, one should note that eqs. (A.24) to (A.26) together with the corresponding experimental data[12] allow also for surprising solutions ; for instance, a good fit is obtained ($\chi^2/dof = 6.5/4$) with $\theta_V = 32.8^\circ \pm 1.5^\circ$ but also $\theta_{PS} = -86.2^\circ \pm 9^\circ$. The reconstructed decay rates are nevertheless in good agreement with the data (comparable to those displayed in the first column of table 8). This kind of strange solution (η is nearly SU(3) singlet and η' nearly pure octet) is in complete contradiction with all expectations (including those of anomaly physics) and will not be discussed here any longer.

8.2 Radiative Decays and Anomaly Results

The consequences of our analysis of $\eta/\eta' \rightarrow \gamma \gamma$, $\eta/\eta' \rightarrow \pi^+ \pi^- \gamma$, $\eta/\eta' \rightarrow \mu^+ \mu^- \gamma$ and $J/\psi \rightarrow (\eta/\eta') \gamma$ (*i.e.* eight processes) can be summarized by the values obtained for F_η , $F_{\eta'}$ and θ_{PS}

---

¹¹This does not mean, of course, that such a breaking is not present, but only that we cannot detect it.

(see rels. (5) and table 5) for each ρ model (M_1 and M_2).

Following the preliminary analysis (8.1), we remove from the sample to be fitted the $K^{*0} \rightarrow K^0 \gamma$ and $\phi \rightarrow \eta \gamma$ decay modes which exhibit inconsistencies with respect to the other decay modes. From the analysis presented in 6.3, we are also led to remove $\rho^0 \rightarrow \eta \gamma$ which exhibits inconsistencies with $\eta \rightarrow \mu^+ \mu^- \gamma$ and $\eta \rightarrow \gamma \gamma$ (see rel. (9)). We have also found that this decay mode is inconsistent with M_1 at the 2σ level and with M_2 at the 3σ level (see table 7). Therefore, we shall keep as input the eight remaining radiative decays and our results on : F_η , $F_{\eta'}$, θ_{PS} .

When discussing model M_1 , we thus have :

$$F_\eta = (0.41 \pm 0.05) \text{ GeV}^{-1} \quad F_{\eta'} = (0.41 \pm 0.03) \text{ GeV}^{-1} \quad \theta_{PS} = (-15.38 \pm 1.20)^\circ \quad , \quad (18)$$

and, when discussing model M_2 , we use :

$$F_\eta = (-0.27 \pm 0.03) \text{ GeV}^{-1} \quad F_{\eta'} = (0.43 \pm 0.03) \text{ GeV}^{-1} \quad \theta_{PS} = (-23.12 \pm 1.23)^\circ \quad , \quad (19)$$

Whichever is the model to be used (M_1 or M_2), we find that a good fit is achieved assuming nonet symmetry for vector mesons ($g \equiv g_1$) and we get no substantial improvement by releasing this assumption. This is interesting since it shows that the additional set of radiative decays mainly determine θ_V , all other parameters being (nearly) fixed by the eight processes mentioned above ($\eta/\eta' \rightarrow \gamma \gamma$, $\eta/\eta' \rightarrow \pi^+ \pi^- \gamma$, $\eta/\eta' \rightarrow \mu^+ \mu^- \gamma$ and $J/\psi \rightarrow (\eta/\eta') \gamma$).

Using model M_1 the result of the fit is :

$$\left\{ \begin{array}{l} g \equiv g_1 = 0.704 \pm 0.016 \text{ GeV}^{-1} \\ g_1' = 0.608 \pm 0.027 \text{ GeV}^{-1} \\ \theta_V = (38.65 \pm 0.18)^\circ \\ \theta_{PS} = -(15.58 \pm 1.15)^\circ \end{array} \right. \quad (20)$$

with $\chi^2/dof = 14/7$. The values inferred for the branching ratios are given in the second data column of table 8. Clearly, the input value of F_η (see rel. (18)) is not large enough to accomodate easily the other decay modes and the fit nearly restores the value associated with the $\rho^0 \rightarrow \eta \gamma$ decay[12] (compare F_η fitted values in the first two data columns of table 8). Otherwise, all predicted branching fractions are consistent with measurements[12], except for $\phi \rightarrow \eta \gamma$ and $K^{*0} \rightarrow K^0 \gamma$ as expected.

It should be noticed that model M_1 nearly agrees with complete nonet symmetry ($g \equiv g_1 \equiv g_1'$), the single decay mode which deteriorates the χ^2 when fitting under this assumption is $\omega \rightarrow \eta \gamma$. This decay mode is found by fit at $\simeq 10^{-3}$, *i.e.* nearly twice the value predicted assuming only $g \equiv g_1 (\simeq 0.62 \cdot 10^{-3})$, but in agreement with the recent measurement of GAMS[43].

Using model M_2 we get :

$$\left\{ \begin{array}{ll} g \equiv g_1 = & 0.714 \pm 0.016 \text{ GeV}^{-1} \\ g_1' = & -0.337 \pm 0.029 \text{ GeV}^{-1} \\ \theta_V = & (31.91 \pm 0.18)^\circ \\ \theta_{PS} = & -(23.03 \pm 1.21)^\circ \end{array} \right. \quad (21)$$

with a good fit ($\chi^2/dof = 6.3/5$). The values of branching ratios which can be inferred from rels. (21) are given in the third data column of table 8. They closely reproduce the values obtained for F_η and $F_{\eta'}$ using exclusively the $\eta/\eta' \rightarrow \pi^+\pi^-\gamma$ decays (see rels. (5)) and θ_{PS} (see table 5) is found at the expected value. Therefore we find consistency of the results coming from anomaly physics with eight out of eleven radiative decays. The prediction for $\rho^0 \rightarrow \eta\gamma$ (resp. $\phi \rightarrow \eta\gamma$) is found a factor of 4 below (resp. above) the reported measurement[12, 23]. The prediction for $K^{*0} \rightarrow K^0\gamma$ is also too large with respect to the measurement[40] by $\simeq 50\%$.

It is interesting to remark that the value for θ_V obtained with M_1 is close to the value predicted from quadratic mass formulae[12]. On the contrary, the value obtained for θ_V using model M_2 is somehow unusual ; it should be noticed that it coincides with a recent prediction of Dillon and Morpurgo[42] following from an analysis of $g_{\rho\gamma}$ and $g_{\omega\gamma}$. This analysis is based on a general parametrization of physical quantities which exploits the $SU(3)$ flavor structure of QCD.

Therefore we find that M_1 gives results close to those obtained using only a consistent set of radiative decays (all but the $K^{*0} \rightarrow K^0\gamma$ and $\phi \rightarrow \eta\gamma$ decays) ; however the global fit fails to provide a good value for F_η and hence to describe the η decays.

On the other hand, model M_2 gives a good description of the eight processes used to establish our anomaly results and eight additional radiative decays of light flavour mesons. Only three decays remain outside this scheme : $K^{*0} \rightarrow K^0\gamma$, $\phi \rightarrow \eta\gamma$ and $\rho^0 \rightarrow \eta\gamma$.

We note again that the disagreement observed for $K^{*0} \rightarrow K^0\gamma$ cannot be attributed to our models ; it is intrinsic to the set of equations (A.24) and (A.25). The disagreement observed for $\phi \rightarrow \eta\gamma$ is already apparent when using only the set of equations ((A.24)-(A.25)) (see 8.1). Moreover, both M_1 and M_2 disagree with the measurement[23] of $\rho^0 \rightarrow \eta\gamma$.

8.3 Comments on the results

By adding to the set of decays used to establish our anomaly results (and the value of ξ) most of the radiative decay modes, we find that the consistency of both models M_1 and M_2 improves.

Concerning M_1 (and hence a picture in agreement with QCD) we find that the main problem is the consistency with the existing $\eta \rightarrow \pi^+\pi^-\gamma$ and $\eta \rightarrow \mu^+\mu^-\gamma$ data. This could indicate a theoretical inconsistency in adding the NAF equation to the set of Chanowitz equations. This inconsistency is however confirmed by the global fit with the standard radiative decays.

Concerning M_2 (and hence a picture in basic disagreement with QCD), we find full consistency when considering most, *i.e.* eight of the eleven measured radiative decays of the type $V(PS) \rightarrow PS(V)\gamma$ and assuming exact $SU(3)$ symmetry.

The additional results on radiative decays suppose that exact SU(3) symmetry holds in the form displayed by eqs. (A.24) and (A.25) : in this case, we have shown that nonet symmetry holds for the coupling constants in the vector meson sector. Then, g , g_1 and θ_{PS} are practically determined by our results on F_η , $F_{\eta'}$ and θ_{PS} from anomaly analysis. As in this case $g_1 \equiv g$, the additional set of decays mainly helps to determine θ_V . In other words we fit the standard set of radiative decay modes with only one parameter (θ_V), all other parameters being determined by an external set of data.

The origin of the disagreements observed between some measurements and the predictions is not easy to trace back. Clearly, the K^{*0} branching ratio cannot be accommodated ; whether this is due to a failure of exact SU(3) symmetry or to the measurement[40] is an open question. If we can rely on SU(3) symmetry, the observed departures from predictions of $\phi \rightarrow \eta\gamma$ and $\rho^0 \rightarrow \eta\gamma$ become significant. It should be noticed that a large value for the branching ratio $\phi \rightarrow \eta\gamma$ has also been predicted by O'Donnell[39] and Oshima[44]¹².

The disagreement for $\rho^0 \rightarrow \eta\gamma$, with respect to $\eta \rightarrow \rho^0\gamma$ (*i.e.* the value of F_η) might be attributed to a mass dependence of F_η . In this case however, the agreement reached with M₂ would be hard to understand.

Finally we have shown that $\phi \rightarrow \eta\gamma$ generates a surprising modification of θ_{PS} (see 8.1). It would be interesting to confirm that this is due to symmetry breaking effects.

The disagreement observed for $\phi \rightarrow \eta\gamma$ and $\rho^0 \rightarrow \eta\gamma$ is particularly surprising as these radiative decays have been measured[23] using the annihilation channel $e^+e^- \rightarrow \eta\gamma$ which involves diagrams identical to those describing the decay $\eta \rightarrow \mu^+\mu^-\gamma$ (one has just to invoke crossing and change muons to electrons).

This leads to the remark that the measurements of $\phi \rightarrow \eta\gamma$ and $\rho^0 \rightarrow \eta\gamma$ are not model independent, while to some extent $\omega \rightarrow \eta\gamma$ is. Indeed the numerical value of the ρ^0 branching ratio is influenced by the Breit Wigner chosen to describe its mass spectrum. As the ϕ is relatively far on the tail of the ρ distribution its description could be sensitive to this mass distribution ; as the ω meson stays on the top of the ρ , its dependence on the ρ distribution should be weaker. Moreover the process $\eta \rightarrow \gamma\gamma$ implies the existence of a non-VDM contribution to $e^+e^- \rightarrow \eta\gamma$ of a fixed strength (B_η). Therefore, one cannot fully rely on the present results obtained[23] assuming full VDM when analyzing this annihilation process¹³. It would be interesting to analyze the $e^+e^- \rightarrow \eta\gamma$ and $e^+e^- \rightarrow \pi^0\gamma$ data using motivated shapes for the ρ meson (models M₁ and M₂) and introducing the non-VDM contributions.

If the model developed in ref.[39] and reproduced in the appendix is suitable to describe radiative decays, up to limited symmetry breaking effects, a test of M₁ versus M₂ (and hence of QCD) is possible. All predictions using only the standard radiative decay modes[23, 39] and model M₁ yield for Br($\phi \rightarrow \eta'\gamma$) a value of $1 \div 2 \cdot 10^{-4}$, slightly below the present upper limit[23] ($4.1 \cdot 10^{-4}$). Model M₂ predicts this branching ratio at the level of 10^{-6} . Even if SU(3) symmetry breaking effects are large, it looks unlikely that they could be responsible for a factor of about 100 which would smear out the difference between M₁ and M₂ predictions. Therefore

¹²Using SU(3) symmetry in another manner ref.[23] claims however lower values.

¹³Likewise arguments can be asserted with the decay modes $\rho/\omega/\phi \rightarrow \pi^0\gamma$ in $e^+e^- \rightarrow \pi^0\gamma$ because of the ρ mass distribution assumed and because of the existence of the decay mode $\pi^0 \rightarrow \gamma\gamma$, which implies a non-VDM contribution to the annihilation process.

η -factories[45, 46] will soon provide interesting informations on this issue¹⁴.

9 General Discussion

In this paper, we have addressed several topics in connection with anomaly physics and the ξ test of QCD formerly proposed by Chanowitz[2, 3, 4]. We now summarize all the results we have obtained.

9.1 Independence of Anomalous and Resonant Contributions

The main tool in treating the decay spectra of $\eta/\eta' \rightarrow \pi^+\pi^-\gamma$ in paper A was the assumption that the resonant (mainly ρ^0) contribution and the anomalous contribution are independent processes which have to be added coherently. As stressed in the introduction, this assumption is not new[15] and has proved to work for the π^0 box anomaly[10, 11]. It has even been used recently to describe the η decay[16]. The consequence of such an assumption for QCD appears however crucial enough[1] to call for an unambiguous test.

We have found that the $\eta/\eta' \rightarrow \mu^+\mu^-\gamma$ processes allow for such an unambiguous test. For the first time the decay widths $\Gamma(\eta/\eta' \rightarrow \gamma\gamma)$ have been derived from the $\eta/\eta' \rightarrow \pi^+\pi^-\gamma$ and $\eta/\eta' \rightarrow \mu^+\mu^-\gamma$ processes. This analysis leads to a good agreement with the experimental values (see table 7). The precision of model M_2 predictions is striking ; model M_1 is not as good but is not ruled out.

Conversely, knowing F_η , $F_{\eta'}$ from $\eta/\eta' \rightarrow \pi^+\pi^-\gamma$ and B_η , $B_{\eta'}$ from $\eta/\eta' \rightarrow \gamma\gamma$, we can make definite predictions about the η/η' form factors in the $\eta/\eta' \rightarrow \mu^+\mu^-\gamma$ decays. This provides a second test of our model and shows that M_2 gives a prediction of both form factors in agreement with the present data[35, 36, 37, 38]. Model M_1 succeeds in predicting the shape of the η' form factor but fails to predict the η form factor (see figs. (3) and (5)). Therefore, in order to test M_1 versus M_2 (and hence QCD), it is important to get better data on the η and η' electromagnetic form factors. Interesting informations can be expected from the η -factory planned to start data taking by the end of 1995 in Saclay[47].

9.2 The ρ Models and the VDM Assumption

To perform the analysis discussed above, we have had to analyze spectra dominated by a ρ contribution. As widely discussed in paper A and in this article, the box anomaly constants $E_{\eta,\eta'}$ are sensitive to the shape of the ρ mass distribution¹⁵ and their precise numerical values are highly correlated with it.

We have chosen to derive the ρ mass distribution from the $e^+e^- \rightarrow \pi^+\pi^-$ cross section[14]. Here arises a theoretical ambiguity connected with the possible existence of a non-VDM $\gamma \rightarrow \pi^+\pi^-$ coupling considered recently in the theory of effective lagrangians[27]. We thus define model M_2 which is 100% VDM and model M_1 which assumes the existence of a small (mass

¹⁴In the $e^+e^- \rightarrow \eta'\gamma$ process, according to our approach, one would have to account for the ϕ as intermediate state, but also for the ρ^0 and a photon because of the triangle anomaly process $\eta' \rightarrow \gamma\gamma$.

¹⁵If we plan to get the triangle anomaly constants $B_{\eta,\eta'}$ from the decays $\eta/\eta' \rightarrow \mu^+\mu^-\gamma$ instead of $\eta/\eta' \rightarrow \gamma\gamma$, we would get the same problem and need the same solution.

dependent) non-VDM contribution. The former gives a mass for the ρ^0 meson close to the ω mass ($\simeq 781$ MeV), the latter is built up in order to give the ρ mass its world average value[12]. We have attempted here to introduce an intermediate solution (M_3) but with little success, pointing to the fact that if the correct ρ model is not exactly M_1 or M_2 , it is probably close to one of them.

Model M_2 is simple : it assumes that no (or no important) non-VDM coupling $\gamma \rightarrow \pi^+\pi^-$ exists. The problem already noticed with M_1 is related to the fact that the non-VDM coupling is mass dependent. However, this result could as well be an effective result obtained by mixing a constant non-VDM coupling to the tail of higher resonances contributing to $e^+e^- \rightarrow \pi^+\pi^-$. Possible candidates exists (they are given in paper A) but have seemingly too small couplings[48] to $\pi^+\pi^-$.

Anyhow, the existence of a non-VDM coupling in e^+e^- annihilation remains an open question which cannot be answered with the presently available data. As explained in **2**, the best way to answer this question would be a simultaneous fit of $e^+e^- \rightarrow \pi^+\pi^-$ and $e^+e^- \rightarrow \mu^+\mu^-$ from threshold to about 1 GeV. In this last process no non-VDM contribution is expected to occur, except for the standard QED background. This would also give an answer to the question of a possible mass dependence of $g_{\rho\gamma}$ (the transition amplitude $\gamma \rightarrow \rho^0$).

9.3 Adding the NAF Equation to the Chanowitz Equations

Following ref.[18, 19], we have studied the consequences of performing a simultaneous fit of the (four) Chanowitz equations together with the equation originating from the work of Novikov[20] *et al.* under the form given by Akhoury and Frère[21]. In this way, we have increased the constraints on f_1 , f_8 , θ_{PS} with respect to the experimental data.

Again, model M_2 provides a good fit and yields a pseudoscalar mixing angle at the value given by the SU(3) linear mass formulae (see table 5), f_8 being in agreement with the chiral model predictions[7] (see table 5).

Model M_1 provides a fit with a pseudoscalar mixing angle in agreement with the recent CRISTAL BARREL result[41]. However, it has some difficulty to accommodate the existing $\eta \rightarrow \pi^+\pi^-\gamma$ data.

In both cases, including the NAF equation improves the accuracy of the results obtained using only the Chanowitz equations (compare M_1 and M_2 results in tables 5 and 2).

In addition, using the NAF equation together with both equations for the η and η' triangle anomalies, makes it possible (assuming $\xi = 1$ or $\xi = 2$) to solve (not fit) for f_1 , f_8 , θ_{PS} and then, by means of the equations (A.17), to predict the values of both E_η and $E_{\eta'}$. This method is intrinsically insensitive to any choice of the ρ mass distribution, and therefore its predictions are of particular interest as our results rely on specific ρ distributions. The results are given in tables 3 and 4. In the case $\xi = 2$ both predictions for E_η and $E_{\eta'}$ are in good agreement with our M_2 results (compare columns 2 and 3 in table 4), pointing to the fact that our modeling of the ρ is suitable. When $\xi = 1$, table 3 shows that $E_{\eta'}$ is correctly predicted (columns 2 and 3), the prediction for E_η being also consistent with the result obtained with the single existing η data set[30] but not as well (compare however this prediction with rel. (33) in paper A). This points once more towards a difficulty of model M_1 to accommodate η data which certainly need

confirmation

To summarize, we consider that the predictions obtained by this standard method support our treatment of the ρ meson. Moreover, it allows to increase from 4 to 5 the number of equations relating f_1 , f_8 , θ_{PS} and ξ . In this way, we are able to confirm the values of ξ associated with each of the models M_1 and M_2 .

9.4 Anomaly Physics and SU(3) Symmetry

From the analysis of the data available on $\eta/\eta' \rightarrow \gamma\gamma$, $\eta/\eta' \rightarrow \pi^+\pi^-\gamma$ and $J/\psi \rightarrow (\eta/\eta')\gamma$, the main results obtained in this work are the values of f_1 , f_8 , θ_{PS} and ξ .

In addition, we also get for each ρ model the coupling strengths F_X ($X = \eta, \eta'$) at the $X\rho^0\gamma$ vertex. Using SU(3) relations[39], it is possible to relate them to all couplings of the kind $V \rightarrow P\gamma$ and $P \rightarrow V\gamma$, where P and V are light flavour mesons.

We have therefore studied the consistency of our results with eleven additional processes (assuming SU(3) is not too badly broken). Assuming in addition nonet symmetry for vector meson, we have shown that model M_1 predicts correctly eight of these decays, but in a way which does not accomodate easily the η radiative decays to two pions or two muons. Under the same assumption, model M_2 does not meet these difficulties.

Among the decay modes which remain outside model M_2 predictions, one finds the K^{*0} meson radiative width[40] which anyhow is difficult to reconcile with exact SU(3) symmetry, and the $\phi \rightarrow \eta\gamma$ decay which is found two to three times the accepted measurement value[23, 12]. This is however not specific of our approach (see the first column in table 8), and, moreover, this measurement contradicts the results obtained for θ_{PS} using the anomaly equations (see the discussion in 8.1).

Finally model M_2 has also some difficulty to accommodate the measured value of $\rho^0 \rightarrow \eta\gamma$. This could indicate a significant mass dependence of F_η , but the analysis of the data on $e^+e^- \rightarrow \eta\gamma$ may have to be reconsidered (see 8.3). Indeed, this process is basically the same as $\eta \rightarrow \mu^+\mu^-\gamma$ which is successfully described using M_2 .

Finally, assuming SU(3) symmetry, model M_1 predicts a branching fraction $\simeq 1 \div 2 \cdot 10^{-4}$ for the $\phi \rightarrow \eta'\gamma$ decay mode in agreement with several approaches[23, 39, 44]. Instead, model M_2 predicts this branching fraction at a much lower level $\simeq 10^{-6}$.

10 Conclusion

In this paper, we have made an attempt to present a coherent picture of all the data available in relation with anomaly physics. We have shown that an equation due to Novikov *et al.*, Akhoury and Frère can be consistently added to the four Chanowitz equations. Within this constrained scheme, we have shown that model M_1 and model M_2 yield better results, showing that the method proposed in ref.[1] and extended here is suitable.

The introduction of this additional equation also shows that our treatment of the ρ mass distribution is in agreement with data on J/ψ decays to $\eta\gamma$ and $\eta'\gamma$. Moreover, the successful reconstruction of $\Gamma(\eta/\eta' \rightarrow \gamma\gamma)$ from the decay processes $\eta/\eta' \rightarrow \pi^+\pi^-\gamma$ and $\eta/\eta' \rightarrow \mu^+\mu^-\gamma$ confirms that splitting up amplitudes into anomalous and resonant contributions is methodologically correct. All this confirms the results on box anomalies and pseudoscalar nonet parameters

given in paper A ; their values are improved in the present paper.

Finally, we have shown that the results explained by models M_1 and M_2 cover $\eta/\eta' \rightarrow \gamma\gamma$, $J/\psi \rightarrow (\eta/\eta')\gamma$, $\eta/\eta' \rightarrow \pi^+\pi^-\gamma$, $\eta/\eta' \rightarrow \mu^+\mu^-\gamma$, $\eta/\eta' \rightarrow e^+e^-\gamma$ and most of the radiative decay modes $V \rightarrow P\gamma$ or $P \rightarrow V\gamma$. This is the largest set of radiative decays (21) addressed in a unified scheme ; it turns out that M_1 and M_2 are able to provide such a scheme. Knowing that M_2 is in basic disagreement with QCD, this result is already remarkable.

We have shown the relevance of several radiative decay and annihilation processes for QCD (e^+e^- to $\pi^+\pi^-$, to $\mu^+\mu^-$, to $\eta\gamma$ or to $\pi^0\gamma$) in relation with the existence of anomalies. Most of this physics is under the scope of several existing detectors and facilities like ϕ -factories or machines planned to start soon (the η -factory in Saclay). The only hypothetical facility is that needed to make suitably the η' physics ; it could be a goal for a τ - e factory[49].

Even if model M_2 behaves in a better way, it is clear that a final answer supposes a better understanding of the ρ^0 physics. The ρ^0 looks as important for testing QCD as the Z^0 was crucial to test the Standard Model. The main question addressed for this purpose is twofold : to which extent $g_{\rho\gamma}$ can be considered as constant? what is the magnitude of a possible non-VDM $\gamma \rightarrow \pi^+\pi^-$ transition? $e^+e^- \rightarrow \mu^+\mu^-$ data should provide an answer. We have also seen that η spectra offer the interesting possibility to check M_1 versus M_2 as these models exhibit different predictions for some of these spectra. Testing the order of magnitude of the $\phi \rightarrow \eta'\gamma$ branching ratio would also provide a valuable contribution. However establishing with a good accuracy the value of ξ supposes a better knowledge of η' physics.

On another hand, the surprising consistency of the NAF equation with the Chanowitz equations call for a search of other relations which could strengthen the constraints on the experimental data. Further predictions like those of Dillon and Morpurgo go the same way.

It happens that this physics is of particular relevance to perform and achieve the test of QCD formerly proposed by Chanowitz. As follows from the discussion just above, several processes converge to check whether $\xi = 1$ as predicted by QCD. Using the available data (which call for improvements), the possibility that $\xi = 2$ does not look especially unlikely. Whatever such a result could actually mean, this possibility is important enough to call for better data in meson spectroscopy.

Acknowledgements

One of us (M.B.) acknowledges many fruitful discussions or mails with V. Petrov (Serpukhov) and J.-M. Frère (Brussels) on anomaly physics, with V. Chernyak (Novossibirsk) about J/ψ decays in the QCD framework, with F. Renard (Montpellier) about SU(3) and radiative decays of mesons and with C. Rybicki (Krakow) for informations about $\pi N \rightarrow \pi\pi N$ data. We also acknowledge M. Froissart (Collège de France, Paris) for useful discussions, comments and reading the manuscript.

Appendix A : Models and Relations in η/η' Decays

We give here a description of the model used to describe the $\eta/\eta' \rightarrow \mu^+\mu^-\gamma$ decays. For the sake of completeness, we also remind some of the formulae which describe the $\eta/\eta' \rightarrow \pi^+\pi^-\gamma$ decays developed in ref.[1], but using precisely the same rules as those used to model the dimuon radiative decays of η/η' mesons. It is however meaningless to copy fully the appendices from paper A, therefore one has to refer to it for the informations not recalled here.

A1 Model for $\eta/\eta' \rightarrow \mu^+\mu^-\gamma$ Decays

The model we propose is based on the existence of the two independent contributions depicted in fig. (1). In figure (1a) we have a term which only proceeds from the combination of the standard triangle anomaly for η/η' (denoted X in the following) and quantum electrodynamics. The triangle anomaly vertex is written :

$$M(X \rightarrow \gamma\gamma) = B_X(q \cdot k) \epsilon_{\mu\nu\rho\sigma} k^\mu q^\nu \varepsilon_i^\rho(k) \varepsilon_j^\sigma(q) \quad , \quad X = \eta, \eta' \quad , \quad (\text{A.1})$$

where the ε are the two photon polarization vectors and i, j are helicity indices, assuming we stay in the X rest frame. Finally q and k are the photon 4-momenta. We shall approximate the functions B_X by their values $B_X(0)$ for on-shell photons :

$$\Gamma(X \rightarrow \gamma\gamma) = \frac{M_X^3}{64\pi} |B_X(0)|^2 \quad , \quad X = \eta, \eta' \quad , \quad (\text{A.2})$$

where the X meson 4-momentum is P and $P^2 = M_X^2$. Denoting by $m^2 = q^2$ the dimuon invariant squared mass, $e = \sqrt{4\pi\alpha_{em}}$ the muon electric charge, the amplitude for the process in figure (1a) can be written :

$$T_i = -\frac{eB}{m^2} \epsilon_{\mu\nu\rho\sigma} k^\mu q^\nu \varepsilon_i^\rho(k) [\bar{v}(q_2) \gamma_\sigma u(q_1)] \quad . \quad (\text{A.3})$$

There are basically three ingredients which arise with the contribution depicted in figure (1b). We have first the vertex function describing the $X \rightarrow \rho^0\gamma$ transition :

$$V_{ij}^1 = F_X \epsilon_{\mu\nu\rho\sigma} q^\mu k^\nu \varepsilon_i^\rho(k) \varepsilon_j^\sigma(q) \quad , \quad X = \eta, \eta' \quad , \quad (\text{A.4})$$

where the vertex function F_X is approximated by a constant as in ref.[1], k and q are respectively the photon and ρ^0 meson 4-momenta and the ε 's are the corresponding polarization vectors. The second ingredient is the (VDM) transition amplitude from ρ^0 to photon :

$$V_{ij}^2 = e g_{\rho\gamma} \varepsilon_i^{\star\alpha}(q) \varepsilon_j^\alpha(q) \quad , \quad (\text{A.5})$$

where α denotes a Lorentz index¹⁶. The quantity $g_{\rho\gamma}$ is supposed to be given by the e^+e^- decay width of the ρ^0 meson in the standard way (see rel.(A.5) in ref.[1] for instance). It is moreover assumed to be real and positive. The last ingredient is the standard QED vertex already used to derive rel. (A.3) above :

---

¹⁶We assume implicitly summation over repeated greek (*i.e.* Lorentz) and latin (*i.e.* helicity) indices throughout this paper.

$$V_3 = -ie [\bar{v}(q_2) \gamma_\sigma u(q_1)] \varepsilon_i^{\sigma\gamma}(q) \quad , \quad (\text{A.6})$$

where $q = q_1 + q_2$. With these Feynman rules, the photon propagator is simply i/m^2 , while the ρ propagator is written $i/D_\rho(m)$. For this last quantity we shall use expression (A.3) from ref.[1] with the parameters obtained in fitting the $e^+e^- \rightarrow \pi^+\pi^-$ cross section[14].

Using these rules, it is a simple matter to compute the diagram in figure (1b) :

$$T_2 = -\frac{ie^2 F_X}{m^2 D_\rho(m)} g_{\rho\gamma} [\bar{v}(q_2) \gamma_\sigma u(q_1)] \varepsilon_{\mu\nu\rho\sigma} k^\mu q^\nu \varepsilon_i^\rho(k) \quad . \quad (\text{A.7})$$

Using elementary rules and integrating over angles, we can deduce the width of the $\eta/\eta' \rightarrow \mu^+\mu^-\gamma$ decay modes :

$$\frac{d\Gamma}{dm} = \frac{\alpha_{em}}{48\pi^2} \frac{\sqrt{m^2 - 4\mu^2}}{M_X^3 m^4} (m^2 + 2\mu^2) (M_X^2 - m^2)^3 \left| B_X + \frac{ie g_{\rho\gamma} F_X}{D_\rho(m)} \right|^2 \quad . \quad (\text{A.8})$$

where μ is the muon mass.

Using rel. (A.2) above, rel. (A.8) can be cast in a more usual form[34] allowing the introduction of the η/η' transition form factors $\mathcal{F}_X(m^2, 0)$:

$$\frac{1}{\Gamma(X \rightarrow \gamma\gamma)} \frac{d\Gamma}{dm} = \frac{4\alpha_{em}}{3\pi} \frac{\sqrt{m^2 - 4\mu^2}}{m^2} \left(1 + \frac{2\mu^2}{m^2} \right) \left(1 - \frac{m^2}{M_X^2} \right)^3 \left| \mathcal{F}_X(m^2, 0) \right|^2 \quad . \quad (\text{A.9})$$

with :

$$\left| \mathcal{F}_X(m^2, 0) \right|^2 = \left| 1 + ie g_{\rho\gamma} \frac{F_X}{B_X} \frac{1}{D_\rho(m)} \right|^2 \quad . \quad (\text{A.10})$$

In analyzing the experimental data on η/η' conversion decays into muons[35][36][37][38] reproduced in ref.[34], we shall have to use expressions (A.8) or (A.9) when dealing with the branching fractions and rel. (A.10) to study the form factor.

Strictly speaking, we would have to add in rels. (A.8) and (A.10) the contributions of the ω and ϕ mesons. This would introduce here contributions $g_{\omega\gamma} F_X(\omega)/D_\omega(m)$ and $g_{\phi\gamma} F_X(\phi)/D_\phi(m)$ besides the term $g_{\rho\gamma} F_X(\rho)/D_\rho(m)$ already accounted for¹⁷. However, since the mass spectrum is cut at M_η in one case, it is useless to go beyond the ρ contribution. In the case of the η' meson, it is also useless to account for the ϕ contribution because this meson is outside the available phase space. Moreover, the accuracy of the data does not allow to feel any ω contribution and therefore we shall also neglect it. In other words, we work under the assumption of ρ dominance.

---

¹⁷Of course, what is called F_X is actually $F_X(\rho)$; since F_X is the coupling constant of X ($= \eta, \eta'$) to a photon and a vector meson V , it should be understood as $F_X(V)$ whenever needed. We shall use this notation each time it is free from ambiguities.

A2 Reminder of the Model for $\eta/\eta' \rightarrow \pi^+\pi^-\gamma$ Decays.

In ref.[1], we have developed a model to describe the decays $\eta/\eta' \rightarrow \pi^+\pi^-\gamma$. For the sake of completeness and in order to keep consistency with the previous model, we remind here the main formulae. The contributions taken into account are depicted in figure (2). More precisely, the anomalous contribution of fig. (2a) is :

$$T'_i = E_X \epsilon_{\mu\nu\rho\sigma} \varepsilon_i^\mu(\gamma) q_1^\nu q_2^\rho k^\sigma \quad (X = \eta, \eta'), \quad , \quad (\text{A . 11})$$

where q_1 and q_2 are respectively the π^+ and π^- 4-momenta. All other notations have been defined just above. E_X is assumed to be constant and is approximated by its value at the chiral point. We define :

$$T_j = iG_\rho \varepsilon_\mu^*(q_1 - q_2)^\mu \quad , \quad (\text{A . 12})$$

the amplitude which describes the ρ^0 decay to $\pi^+\pi^-$; its expression¹⁸ in terms of the $\Gamma(\rho^0 \rightarrow \pi^+\pi^-)$ width can be found in ref.[1]. Using rel. (A.4) above, the contribution depicted in fig. (2b) can be written :

$$T_i = -\frac{2G_\rho F_X}{D_\rho(m)} \epsilon_{\mu\nu\rho\sigma} \varepsilon_i^\mu(\gamma) q_1^\nu q_2^\rho k^\sigma \quad . \quad (\text{A . 13})$$

and the decay width to $\pi^+\pi^-\gamma$ is :

$$\frac{d\Gamma}{dm} = \frac{1}{48\pi^3} \left| \frac{2G_\rho F_X}{D_\rho(m)} + E_X \right|^2 k_\gamma^3 q_\pi^3 \quad , \quad (\text{A . 14})$$

with $k_\gamma = (M_X^2 - m^2)/2M_X$ and $q_\pi = 1/2\sqrt{m^2 - 4m_\pi^2}$. The function $G_\rho(m)$ is deduced from fitting[1] the e^+e^- data[14][1].

As soon as a value of E_X/F_X is obtained from a minimization procedure the value of F_X is simply given by :

$$\frac{1}{F_X^2} = \frac{1}{\Gamma(X \rightarrow \pi^+\pi^-\gamma)} \frac{1}{48\pi^3} \int_{2m_\pi}^{M_X} \left| \frac{2G_\rho}{D_\rho(m)} + \frac{E_X}{F_X} \right|^2 k_\gamma^3 q_\pi^3 dm \quad . \quad (\text{A . 15})$$

The expressions for the constants E_X and B_X at the chiral point are recalled below.

As in the previous model, we should have to account for a possible ω contribution in (A.14) in the case of η' . However, a direct inspection of the experimental spectra reproduced in ref.[1], as well as the study in ref.[13] clearly show that such a complication is presently useless. Another possible contribution would be $\eta' \rightarrow \rho^\pm \pi^\mp$ with a subsequent decay $\rho^\pm \rightarrow \pi^\pm \gamma$; it is easy to estimate its magnitude and convince oneself that it is insignificant at the present level of accuracy of the data¹⁹.

¹⁸Notice the additional factor i in expression (A.12) here compared with rel. (B.2) in ref.[1]. It is due to the fact that, following the usual custom, we define now the ρ propagator by $i/D_\rho(m)$ instead of $1/D_\rho(m)$. Actually to get a consistent picture, each coupling involving an odd number of momenta has to carry a factor i .

¹⁹Its intensity would not exceed that of an ω contribution but spread out onto the whole $\pi^+\pi^-$ mass spectrum.

A3 Problems of Signs

As it can be seen from rels. (A.8) and (A.10), we have access experimentally only to the *relative* signs of the B_X and F_X constants. However, the signs of the ratios B_η/F_η and $B_{\eta'}/F_{\eta'}$ can be meaningfully compared²⁰. The situation is completely symmetric for E_η/F_η and $E_{\eta'}/F_{\eta'}$ from rels. (A.14) ; the reason now is that the sign in rel. (A.12) is arbitrary and therefore the coefficients in rels. (A.11) and (A.13) can carry like signs or opposite signs as well. However, here again one can assume safely that the relative sign of E_η/F_η and $E_{\eta'}/F_{\eta'}$ is significant.

The signs of the F_X have been fixed in ref.[1], because self-consistency of the Chanowitz equations (see below) implies that the four quantities B_X and E_X ($X = \eta, \eta'$) are likely to be negative. In the scheme developed in ref.[18], the corresponding sequence of signs is always negative (as for the Chanowitz equations) when $\xi = 1$ (fractionally charged quarks), but not necessarily when $\xi = 2$ (case of integral-charge quarks). This remark is of concern in connexion with possible SU(3) predictions concerning the relative signs of F_η and $F_{\eta'}$.

A4 The Chanowitz Equations

The functions E_X and B_X introduced above are generally approximated by their values at the chiral point. In the form proposed by Chanowitz[2][3][4], they are written :

$$\begin{cases} B_\eta(0) &= -\frac{\alpha_{cm}}{\pi\sqrt{3}} \left[\frac{\cos\theta_{PS}}{f_8} - 2\sqrt{2}\xi \frac{\sin\theta_{PS}}{f_1} \right] \\ B_{\eta'}(0) &= -\frac{\alpha_{cm}}{\pi\sqrt{3}} \left[\frac{\sin\theta_{PS}}{f_8} + 2\sqrt{2}\xi \frac{\cos\theta_{PS}}{f_1} \right] \end{cases} \quad (\text{A.16})$$

and :

$$\begin{cases} E_\eta(0) &= -\frac{\epsilon}{4\pi^2\sqrt{3}} \frac{1}{f_\pi^2} \left[\frac{\cos\theta_{PS}}{f_8} - \sqrt{2} \frac{\sin\theta_{PS}}{f_1} \right] \\ E_{\eta'}(0) &= -\frac{\epsilon}{4\pi^2\sqrt{3}} \frac{1}{f_\pi^2} \left[\frac{\sin\theta_{PS}}{f_8} + \sqrt{2} \frac{\cos\theta_{PS}}{f_1} \right] \end{cases} \quad (\text{A.17})$$

These equations connect a phase space term contribution with the pseudoscalar meson parameters : the singlet and octet coupling constants, and the mixing angle.

The parameter ξ which occurs in rels. (A.16) is connected with the underlying theory of strong interactions. If QCD holds, we have necessarily $\xi = 1$. In most realizations of integral-quark charge models (ICQ) $\xi = 2$. This allows for a test of QCD in its most stringent prediction. Even if motivated by ICQ models, equations (A.16) and (A.17) are interesting since they allow for something else than QCD.

Setting $\beta = \epsilon f_\pi^2$, eqs. (A.16) and (A.17), give :

$$\xi = \frac{1}{2\beta} \frac{B_\eta^2 + B_{\eta'}^2 - \beta(B_\eta E_\eta + B_{\eta'} E_{\eta'})}{B_{\eta'} E_{\eta'} + B_\eta E_\eta - \beta(E_\eta^2 + E_{\eta'}^2)}. \quad (\text{A.18})$$

---

²⁰Since the sign of $g_{\rho\gamma}$ is not fixed by theory (SU(3) or even VDM for instance), the sign of each of these ratios is not not known. Referring to the fits done in ref.[1] on e^+e^- data, one could possibly assert that the quantities $g_{V\gamma}G_V(m)$ are of like signs for $V = \omega$ and $V = \rho$.

This equation replaces rel. (34) of paper A which has been wrongly copied : the numerical values of rel. (35) in paper A and the subsequent discussion are however correct.

A5 A Relation using J/ψ Decays

Using a relation of Novikov[20], Akhoury and Frère[21] have derived a useful relation :

$$\frac{\Gamma(J/\psi \rightarrow \eta' \gamma)}{\Gamma(J/\psi \rightarrow \eta \gamma)} = \left[\frac{M_{J/\psi}^2 - M_{\eta'}^2}{M_{J/\psi}^2 - M_{\eta}^2} \right]^3 \left[\frac{M_{\eta'}}{M_{\eta}} \right]^4 \left[\frac{\sqrt{2}/f_8 \cos \theta_{PS} + 1/f_1 \sin \theta_{PS}}{1/f_1 \cos \theta_{PS} - \sqrt{2}/f_8 \sin \theta_{PS}} \right]^2 \quad (\text{A . 19})$$

where the numerical factor is simply the ratio of the phase space terms in the J/ψ radiative decays to η' and η . In connection with the ξ test of QCD this relation has not to be changed if ξ is allowed to get values different of 1. This is simply connected with the fact that these processes involve a single photon.

A6 Description of $e^+e^- \rightarrow \pi^+\pi^-$ Data

The cross section $e^+e^- \rightarrow \pi^+\pi^-$ is fitted using the expression :

$$\sigma(m) = \frac{12\pi}{m^4} \left| A \frac{\alpha_{em}}{3} \left[\frac{2q_\pi^3(m)}{m} \right]^{1/2} + \frac{m_\rho^2 \sqrt{\Gamma_\rho(e^+e^-)\Gamma_\rho(m)}}{D_\rho(m)} + e^{i\varphi} \frac{m_\omega^2 \sqrt{\Gamma_\omega(e^+e^-)\Gamma_\omega(m)}}{D_\omega(m)} \right|^2 \quad (\text{A . 20})$$

This expression[1] contains the usual resonant contributions beside a possible non-VDM contribution (*i.e.* the first term). q_π is the pion 3-momentum and φ is the $\rho^0 - \omega$ mixing angle. The other quantities of concern are :

$$\begin{cases} D_\rho(m) = (m_\rho^2 - m^2) - im_\rho \Gamma_\rho(m) \\ D_\omega(m) = (m_\omega^2 - m^2) - im_\omega \Gamma_\omega(m_\omega) \end{cases} \quad (\text{A . 21})$$

and the (varying) widths are :

$$\begin{cases} \Gamma_\rho(m) = \Gamma_\rho(m_\rho) \left[\frac{q_\pi(m)}{q_\pi(m_\rho)} \right]^3 \left[\frac{m_\rho}{m} \right]^\lambda \\ \Gamma_\omega(m) = Br_\omega(\pi^+\pi^-) \Gamma_\omega(m_\omega) \left[\frac{q_\pi(m)}{q_\pi(m_\omega)} \right]^3 \frac{m_\omega}{m} \end{cases}, \quad (\text{A . 22})$$

where λ is a parameter to be fitted. As the ω is narrow a corresponding λ parameter is useless[1]. It is also the reason why one chooses the tabulated fixed width in the second expression (A.21).

A7 SU(3) Predictions for Radiative Decays

One generally assumes that the photon has no (flavour) SU(3) singlet component[24, 39] when relating with each other all amplitudes of the type $V \rightarrow P\gamma$ and $P \rightarrow V\gamma$, where V and

P denote respectively any (light flavour) vector and pseudoscalar meson. The vertex function for such processes can be written :

$$T = g_{VP\gamma} \epsilon^{\mu\nu\rho\sigma} q_\mu k_\nu \varepsilon_\rho(V) \varepsilon_\sigma(\gamma) \quad , \quad (\text{A.23})$$

where q and k are the 4-momenta of respectively the V meson and the photon. The ε are the corresponding polarization 4-vectors.

The coupling strengths $g_{VP\gamma}$ depend on the two mixing angles θ_V and θ_{PS} and on three independent couplings constants : one describing the coupling $g_{V8P8\gamma} \equiv g$, the second describing the coupling $g_{V1P8\gamma} \equiv g_1$ and finally $g_{V8P1\gamma} \equiv g_1'$. The assumption of no SU(3) singlet component for the photon means that $g_{V1P1\gamma} \equiv 0$.

Traditionally[39] one prefers using α_V and α_P to express these couplings ($\alpha_{V,P} = \theta_{V,PS} - \arctan 1/\sqrt{2}$).

Comparing rel. (A.23) with rel. (A.4) clearly show that $F_\eta \equiv g_{\rho\eta\gamma}$ and $F_{\eta'} \equiv g_{\rho\eta'\gamma}$, up to a possible global sign.

We take the following relationships from ref.[39] :

$$\left\{ \begin{array}{l} g_{\rho^0\pi^0\gamma} = \frac{1}{3}g \\ g_{\rho^\pm\pi^\pm\gamma} = \frac{1}{3}g \\ g_{K^{*0}K^0\gamma} = -\frac{2}{3}g \\ g_{K^{*\pm}K^\pm\gamma} = \frac{1}{3}g \\ g_{\rho^0\eta\gamma} = \frac{1}{3} \left[\sqrt{2}(g - g_1') \cos \alpha_P - (2g_1' + g) \sin \alpha_P \right] \\ g_{\rho^0\eta'\gamma} = \frac{1}{3} \left[\sqrt{2}(g - g_1') \sin \alpha_P + (2g_1' + g) \cos \alpha_P \right] \\ g_{\omega\pi^0\gamma} = \frac{1}{3} \left[\sqrt{2}(g - g_1) \sin \alpha_V + (2g_1 + g) \cos \alpha_V \right] \\ g_{\phi\pi^0\gamma} = \frac{1}{3} \left[\sqrt{2}(g - g_1) \cos \alpha_V - (2g_1 + g) \sin \alpha_V \right] \end{array} \right. \quad (\text{A.24})$$

and also :

$$\left\{ \begin{array}{l}
g_{\omega\eta\gamma} = \frac{1}{9} \cos \alpha_V \sin \alpha_P [g - 2g_1' - 2g_1] - \frac{2}{9} \sin \alpha_V \cos \alpha_P [g + g_1' + g_1] \\
\quad + \frac{\sqrt{2}}{9} \sin \alpha_V \sin \alpha_P [g - 2g_1' + g_1] + \frac{\sqrt{2}}{9} \cos \alpha_V \cos \alpha_P [-g - g_1' + 2g_1] \\
g_{\phi\eta\gamma} = -\frac{1}{9} \sin \alpha_V \sin \alpha_P [g - 2g_1' - 2g_1] - \frac{2}{9} \cos \alpha_V \cos \alpha_P [g + g_1' + g_1] \\
\quad + \frac{\sqrt{2}}{9} \cos \alpha_V \sin \alpha_P [g - 2g_1' + g_1] - \frac{\sqrt{2}}{9} \sin \alpha_V \cos \alpha_P [-g - g_1' + 2g_1] \\
g_{\omega\eta'\gamma} = -\frac{1}{9} \cos \alpha_V \cos \alpha_P [g - 2g_1' - 2g_1] - \frac{2}{9} \sin \alpha_V \sin \alpha_P [g + g_1' + g_1] \\
\quad - \frac{\sqrt{2}}{9} \sin \alpha_V \cos \alpha_P [g - 2g_1' + g_1] + \frac{\sqrt{2}}{9} \cos \alpha_V \sin \alpha_P [-g - g_1' + 2g_1] \\
g_{\phi\eta'\gamma} = \frac{1}{9} \sin \alpha_V \cos \alpha_P [g - 2g_1' - 2g_1] - \frac{2}{9} \cos \alpha_V \sin \alpha_P [g + g_1' + g_1] \\
\quad - \frac{\sqrt{2}}{9} \cos \alpha_V \cos \alpha_P [g - 2g_1' + g_1] - \frac{\sqrt{2}}{9} \sin \alpha_V \sin \alpha_P [-g - g_1' + 2g_1]
\end{array} \right. \quad (\text{A . 25})$$

We have corrected the last term in the second relation (A.25) for an error in ref.[39]. Using these coupling constants, the decays widths can be written :

$$\left\{ \begin{array}{l}
\Gamma(V \rightarrow P\gamma) = \frac{1}{96\pi} \left[\frac{m_V^2 - m_P^2}{m_V} \right]^3 |g_{VP\gamma}|^2 \\
\Gamma(P \rightarrow V\gamma) = \frac{1}{32\pi} \left[\frac{m_P^2 - m_V^2}{m_P} \right]^3 |g_{VP\gamma}|^2
\end{array} \right. \quad (\text{A . 26})$$

Assuming full nonet symmetry amounts to set $g = g_1' = g_1$ in rels.(A.24) and (A.25). Since SU(3) is much more badly broken in the pseudoscalar sector than in the vector sector (if we refer to meson masses), nonet symmetry could be better justified for vector mesons only ; in this case one could attempt to see whether $g = g_1$ would not be consistent with the data on radiative decays of light flavour mesons.

Figures

Figure 1: Diagrams of the $\eta/\eta' \rightarrow \mu^+\mu^-\gamma$ decay processes. In (a) the contribution induced by the triangle anomaly ; in (b) the contribution involving the ρ^0 with its subsequent decay to a muon pair.

Figure 2: Diagrams of the $\eta/\eta' \rightarrow \pi^+\pi^-\gamma$ decay processes. In (a) the contribution induced by the box anomaly ; in (b) the contribution involving the ρ^0 which further decays to a pion pair.

Figure 3: The η transition form factor $\eta \rightarrow \mu^+\mu^-\gamma$ described using model M₁ for the ρ^0 . In full line, the fit ; in dashed and dashed-dotted lines the two predictions using the results of the fit of $\eta \rightarrow \pi^+\pi^-\gamma$ (see text). The location of the two pion threshold is indicated.

Figure 4: The η transition form factor $\eta \rightarrow \mu^+ \mu^- \gamma$ described using model M₂ for the ρ^0 . In full line, the fit ; in dashed and dashed-dotted lines the two predictions using the results of the fit of $\eta \rightarrow \pi^+ \pi^- \gamma$ (see text). The location of the two pion threshold is indicated.

Figure 5: The η' transition form factor $\eta' \rightarrow \mu^+\mu^-\gamma$ described using model M_1 for the ρ^0 . In full line, the fit ; in dashed and dashed-dotted lines the two predictions using the results of the fit of $\eta' \rightarrow \pi^+\pi^-\gamma$ (see text). The location of the two pion threshold is indicated.

Figure 6: The η' transition form factor $\eta' \rightarrow \mu^+\mu^-\gamma$ described using model M₂ for the ρ^0 . In full line, the fit ; in dashed and dashed-dotted lines the two predictions using the results of the fit of $\eta' \rightarrow \pi^+\pi^-\gamma$ (see text). The location of the two pion threshold is indicated.

Tables

	Model M ₁	Model M ₂	Model M ₃
m_ρ (MeV)	768.7 ± 0.7	$780.8^{+0.5}_{-0.4}$	772.8 ± 1.4
Γ_ρ (MeV)	$142.8^{+1.5}_{-2.4}$	$153. \pm 2.$	$143.4^{+2.2}_{-2.1}$
λ	1.748 ± 0.079	0.659 ± 0.046	1.268 ± 0.100
φ (degrees)	111.4 ± 3.2	94.3 ± 2.9	108.5 ± 4.8
c_0	0.61 ± 0.06	0.	0.26 ± 0.04
c_2 (GeV ⁻²)	-0.28 ± 0.07	0.	0.
χ^2/dof	61/77	90/78	61/77

Table 1 : Parameter values in the fit of the $e^+e^- \rightarrow \pi^+\pi^-$ cross section ; numerical values written boldface are fixed (input) values. The first two parameters are the mass and width of the ρ^0 meson, λ is the damping power affecting the ρ^0 high-mass tail and φ is the $\rho^0 - \omega$ mixing angle. The last two lines give the parameter values of the non-VDM $\gamma \rightarrow \pi^+\pi^-$ coupling $A(m) = c_0 + c_2m^2$.

	Model M ₁	Model M ₂	Model M ₃
ξ	1	2	2
χ^2/dof	1.9/1	0.06/1	1.3/1
$\frac{f_\pi}{f_1}$	0.96 ± 0.03	0.48 ± 0.02	0.49 ± 0.01
$\frac{f_\pi}{f_8}$	$0.34^{+0.29}_{-0.26}$	$0.42^{+0.22}_{-0.20}$	$-0.15^{+0.17}_{-0.16}$
θ_{PS}	$-29.33^\circ \begin{smallmatrix} +6.62^\circ \\ -6.05^\circ \end{smallmatrix}$	$-27.68^\circ \begin{smallmatrix} +5.24^\circ \\ -4.71^\circ \end{smallmatrix}$	$-39.34^\circ \begin{smallmatrix} +3.97^\circ \\ -3.74^\circ \end{smallmatrix}$
χ^2/dof (converse ξ)	27/1	46/1	37/1

Table 2 : Values of the pseudoscalar nonet parameters obtained by fitting the η and η' mass spectra using the quoted ρ^0 model.

$\xi = 1$	Kisselev–Petrov (current mixing)	Chanowitz–NAF	Model M ₁
$\frac{f_\pi}{f_1}$	$1.12^{+0.09}_{-0.10}$	0.93 ± 0.02	0.96 ± 0.03
$\frac{f_\pi}{f_8}$	$1.19^{+0.11}_{-0.13}$	0.87 ± 0.05	$0.34^{+0.29}_{-0.26}$
θ_{PS}	$-19.65^\circ \pm^{+2.19^\circ}_{-2.15^\circ}$	$-18.85^\circ \pm 1.17^\circ$	$-29.33^\circ \pm^{+6.62^\circ}_{-6.05^\circ}$
E_η	$-7.37^{+0.71}_{-0.77}$	-6.84 ± 0.31	$-4.39^{+1.00}_{-1.15}$
$E_{\eta'}$	$-4.14^{+0.48}_{-0.53}$	-5.32 ± 0.18	$-5.06^{+0.54}_{-0.53}$

Table 3 : Case $\xi = 1$; comparison of our results obtained using model M₁ with the corresponding results of Kisselev–Petrov[18] and that obtained using the Novikov[20] equation in the standard way.

$\xi = 2$	Kisselev–Petrov (mass mixing)	Chanowitz–NAF	Model M ₂
$\frac{f_\pi}{f_1}$	$0.88^{+0.09}_{-0.10}$	0.48 ± 0.01	0.48 ± 0.02
$\frac{f_\pi}{f_8}$	$1.39^{+0.06}_{-0.08}$	0.57 ± 0.04	$0.42^{+0.22}_{-0.20}$
θ_{PS}	$-19.55^\circ \pm^{+1.83^\circ}_{-1.86^\circ}$	$-25.22^\circ \pm 1.16^\circ$	$-27.68^\circ \pm^{+5.24^\circ}_{-4.71^\circ}$
E_η	$-7.27^{+0.76}_{-0.83}$	-4.41 ± 0.23	$-3.64^{+0.97}_{-1.10}$
$E_{\eta'}$	$+2.47^{+0.81}_{-0.74}$	-2.05 ± 0.13	$\pm(-2.17^{+0.49}_{-0.46})$

Table 4 : Case $\xi = 2$; comparison of our results obtained using model M₂ with the corresponding results of Kisselev–Petrov[18] and that obtained using the Novikov[20] equation in the standard way.

	Model M ₁		Model M ₂
	four equations	five equations	five equations
ξ	1	1	2
χ^2/dof	0.34/1	5.56/2	0.87/2
$\frac{f_\pi}{f_1}$	0.92 ± 0.02	0.91 ± 0.02	0.48 ± 0.01
$\frac{f_\pi}{f_8}$	0.93 ± 0.06	0.90 ± 0.05	0.61 ± 0.04
θ_{PS}	$-16.86^\circ \pm 1.20^\circ$	$-16.38^\circ \pm 1.20^\circ$	$-23.12^\circ \pm 1.23^\circ$

Table 5 : Values of the pseudoscalar nonet parameters obtained using three of the Chanowitz equations and the NAF equation with model M₁ (first data column) ; values obtained using the four Chanowitz equations and the NAF equation with both model M₁ and model M₂ results (second and third data columns).

	Model M ₁		Model M ₂	
	predicted	fitted	predicted	fitted
$\frac{F_\eta}{B_\eta}$	$\pm(-17.30 \pm 3.49)$	-10.92 ± 0.70	$\pm(-11.39 \pm 2.22)$	-11.22 ± 0.73
χ^2/dof		14.4/8		15.8/8
$\frac{F_{\eta'}}{B_{\eta'}}$	$\pm(-12.77 \pm 1.00)$	$-9.74^{+2.41}_{-1.97}$	$\pm(+13.40 \pm 1.01)$	$11.80^{+2.07}_{-2.58}$
χ^2/dof		4.4/4		5.8/4

Table 6 : Fits of $\eta/\eta' \rightarrow \mu^+\mu^-\gamma$ transition form factors using models M₁ and M₂ for the ρ invariant mass distribution. We also give the value of the parameter expected from $\eta/\eta' \rightarrow \gamma\gamma$ and from $\eta/\eta' \rightarrow \pi^+\pi^-\gamma$.

	Model M ₁	Model M ₂	PDG
$\eta \rightarrow \gamma\gamma$ (units 10^{-2})	112^{+61}_{-43}	44^{+28}_{-14}	39.8 ± 0.5
$\eta' \rightarrow \gamma\gamma$ (units 10^{-2})	$2.1^{+0.9}_{-0.8}$	$1.9^{+1.0}_{-0.5}$	2.17 ± 0.17
$\eta \rightarrow \mu^+\mu^-\gamma$ (units 10^{-4})	5.56 ± 1.18	3.59 ± 0.84	3.10 ± 0.40
$\eta' \rightarrow \mu^+\mu^-\gamma$ (units 10^{-4})	1.12 ± 0.14	1.13 ± 0.14	1.06 ± 0.27
$\eta \rightarrow e^+e^-\gamma$ (units 10^{-3})	8.03 ± 2.24	8.71 ± 2.28	5.00 ± 1.20
$\eta' \rightarrow e^+e^-\gamma$ (units 10^{-4})	5.30 ± 0.58	6.65 ± 0.75	no data
$\rho^0 \rightarrow \eta\gamma$ (units 10^{-4})	2.18 ± 0.53	1.00 ± 0.22	3.80 ± 0.70
$\rho^0 \rightarrow \pi^0\gamma$ (units 10^{-4})	2.44 ± 0.87	5.23 ± 0.79	7.90 ± 2.00
$\rho^\pm \rightarrow \pi^\pm\gamma$ (units 10^{-4})	2.46 ± 0.87	5.28 ± 0.80	4.50 ± 0.50
$K^{*0} \rightarrow K^0\gamma$ (units 10^{-3})	1.70 ± 0.61	3.64 ± 0.55	2.30 ± 0.20
$K^{*\pm} \rightarrow K^\pm\gamma$ (units 10^{-3})	0.43 ± 0.15	0.92 ± 0.14	1.01 ± 0.09

Table 7 : Values of branching fractions predicted using the results of models M₁ and M₂ and the corresponding PDG values. The errors on $e^+e^-\gamma$ decay fractions of the η/η' mesons predicted using models M₁ and M₂ do not take into account the uncertainty in continuing the ρ^0 mass distribution down to $m = 1$ MeV. See text for further comments.

	Internal fit	Model M ₁	Model M ₂	PDG
F_η (units GeV^{-1})	0.538 ± 0.014	0.525 ± 0.016	-0.271 ± 0.015	
$F_{\eta'}$ (units GeV^{-1})	0.374 ± 0.24	0.369 ± 0.030	0.414 ± 0.022	
$\rho^0 \rightarrow \pi^0 \gamma$ (units 10^{-4})	5.08 ± 0.24	4.95 ± 0.24	5.09 ± 0.24	7.90 ± 2.00
$\rho^\pm \rightarrow \pi^\pm \gamma$ (units 10^{-4})				4.50 ± 0.50
$\rho^0 \rightarrow \eta \gamma$ (units 10^{-4})	3.41 ± 0.18	3.25 ± 0.17	0.87 ± 0.10	3.80 ± 0.70
$K^{*0} \rightarrow K^0 \gamma$ (units 10^{-3})	3.54 ± 0.17	3.45 ± 0.17	3.55 ± 0.17	2.30 ± 0.20
$K^{*\pm} \rightarrow K^\pm \gamma$ (units 10^{-3})	0.89 ± 0.04	0.87 ± 0.04	0.89 ± 0.04	1.01 ± 0.09
$\omega \rightarrow \pi^0 \gamma$ (units 10^{-2})	8.71 ± 0.32	8.49 ± 0.31	8.73 ± 0.32	8.50 ± 0.50
$\omega \rightarrow \eta \gamma$ (units 10^{-4})	6.50 ± 0.60	6.14 ± 0.58	3.27 ± 0.44	$4.70^{+2.20}_{-1.80}$
$\phi \rightarrow \pi^0 \gamma$ (units 10^{-3})	1.40 ± 0.73	1.40 ± 0.73	1.40 ± 0.70	1.31 ± 0.13
$\phi \rightarrow \eta \gamma$ (units 10^{-2})	2.84 ± 0.18	2.85 ± 0.18	4.62 ± 0.18	1.28 ± 0.06
$\phi \rightarrow \eta' \gamma$ (units 10^{-6})	$141. \pm 10.$	$132. \pm 10.$	1.09 ± 0.91	≤ 410
$\eta' \rightarrow \omega \gamma$ (units 10^{-2})	3.42 ± 0.41	3.32 ± 0.40	3.20 ± 0.33	3.00 ± 0.30

Table 8 : Results from the fits and the PDG reference values. The first data column gives the internal fit to the radiative decays only. The second (third) data column gives the results obtained by adding the results of the other η and η' decays using model M₁ (M₂) results.

References

- [1] M. Benayoun *et al.* Zeit. für Phys. **C58** (1993) 31.
- [2] M.S. Chanowitz, Phys. Rev. Lett. **35** (1975) 977.
- [3] M.S. Chanowitz, Phys. Rev. Lett. **44** (1980) 59.
- [4] M.S. Chanowitz, in “Proceedings of the VIth International Workshop on Photon–Photon Collisions 1984”, R.L. Lander Editor, World Scientific, Singapore 1985.
- [5] J. Wess and B. Zumino, Phys. Lett. **B37** (1971) 95.
- [6] J. F. Donoghue, B. R. Holstein and Y. R. Lin, Phys. Rev. Lett. **55** (1985) 2766.
- [7] J. Gasser and H. Leutwyler, Nucl. Phys **B250** (1985) 465.
- [8] L. Adler, Phys. Rev. **117** (1969) 2426.
- [9] J.S. Bell and R. Jackiw, Nuov. Ciment. **A60** (1969) 47.
- [10] S.R. Amendolia *et al.* Phys. Lett. **B155** (1985) 457.
- [11] Yu. M. Antipov *et al.* Phys. Rev. Lett. **56** (1986) 796.
- [12] Review of Particle Properties Phys. Rev. **D45** (1992) S1.
- [13] S.I. Bityukov *et al.*, Zeit. für Phys. **C50** (1991) 451.
- [14] L. M. Barkov *et al.*, Nucl. Phys. **B256** (1985) 365.
- [15] M.V. Terentev, Phys. Lett. **B38** (1972) 419.
- [16] M.K. Volkov and A.A. Osipov, Sov. Journ. Nucl. Phys. **55** (1992) 107 ; Yad. Fiz. **55** (1992) 191.
- [17] A.I. Ivanov and V.M. Shekhter, Sov. Journ. of Nucl. Phys. **32** (1980) 410 ; Yad. Fiz. **32** (1980) 796.
- [18] A.V. Kisselev and V.A. Petrov, preprint CERN–TH 7184–94, Geneva, jan. 94.
- [19] A.V. Kisselev and V.A. Petrov, Zeit. für Phys. **C58** (1993) 595.
- [20] V.A. Novikov *et al.* Nucl. Phys. **B165** (1980) 55.
- [21] R. Akhoury and J.–M. Frère, Phys. Lett. **B220** (1989) 258.
- [22] R. P. Feynman, “Photon–Hadron Interactions”, W.A. Benjamin and Sons, Reading, 1972.
- [23] S.I. Dolinsky *et al.* Pys. Rep. **202** (1991) 99.
- [24] J. P. Perez–Y–Jorba and F. M. Renard, Phys. Rep. **31** (1977) 1.

- [25] V. Chabaud *et al.*, Nucl. Phys. **B223** (1983) 1.
- [26] H. Albrecht *et al.*, 'Measurement of the decay $\tau^- \rightarrow \rho^- \nu_\tau$ ', Desy preprint 92-082, Hamburg June 1992.
- [27] M. Bando, T. Kugo, S. Uehara, K. Yamawaki and T. Yanagida, Phys. Rev. Lett. **54** (1985) 1215.
- [28] H.-J. Behrend *et al.*, CELLO Collaboration, Zeit. für Phys. **C56** (1992) 381.
- [29] S. Kawabata in "Proceedings of the Joint International Lepton-Photon symposium and Europhysics Conference on High Energy Physics", Geneva 1991, vol.2, p 53, S. Hegarty, K. Potter and E. Quercigh editors, World Scientific, Singapore 1992.
- [30] J. G. Layter *et al.*, Phys. Rev. **D7** (1973) 2565.
- [31] F. J. Gilman and R. Kauffman, Phys. Rev. **D36** (1987) 2761.
- [32] A. Bramon and E. Masso, Phys. Lett. **B104** (1981) 311.
- [33] Ll. Amettler *et al.*, Nucl. Phys. **B228** (1983) 301.
- [34] L.G. Landsberg, Phys. Rep. **128** (1985) 301.
- [35] V.A. Viktorov *et al.* Sov. Journ of Nucl. Phys. **32** (1980) 516 ; Yad. Fiz. **32** (1980) 998.
- [36] R.I. Djhelyadin *et al.* Phys. Lett. **B94** (1980) 548.
- [37] R.I. Djhelyadin *et al.* Phys. Lett. **B88** (1980) 379.
- [38] V.A. Viktorov *et al.* Sov. Journ of Nucl. Phys. **32** (1980) 520 ; Yad. Fiz. **32** (1980) 1005.
- [39] P. O'Donnell, Rev. of Mod. Phys. **53** (1981) 673.
- [40] D. Carlsmith *et al.* Phys. Rev. Lett. **56** (1986) 18.
- [41] C. Amsler *et al.* Phys. Lett. **B294** (1992) 451.
- [42] G. Dillon and G. Morpurgo, "QCD parametrization of the $\rho\gamma$, $\omega\gamma$ and $\phi\gamma$ couplings : Why $f_{\rho\gamma} : f_{\omega\gamma} = 3 : 1$ in spite of flavor breaking", Genova, preprint INFN/GE-34118, SW 9417, April 7, 1994.
- [43] D. Alde *et al.* Zeit. für Phys. **C61**, (1994) 35.
- [44] T. Oshima, Phys. Rev. **D22** (1980) 707.
- [45] A. Bramon *et al.*, in "The DAΦNE Physics Handbook", vol. 2, L. Maiani, G. Pancheri, N. Paver (eds), p305, Frascati: INFN 1992.
- [46] R.R. Akhmetshin *et al.* "Preliminary results from the CMD-2 Detector at c.m. energy 1 GeV", Novossibirsk preprint 1993.

- [47] “Rare decays of the eta meson”, Laboratoire National Saturne, Proposal 284, november 1993. B. Mayer, in “Future directions in particle and nuclear Physics at multiGeV hadron beam facilities”, BNL report 52-389, 1993.
- [48] D. Aston *et al.* Preprint SLAC-PUB-5006 Stanford, january 1994, submitted to Zeit. für Phys. C.
- [49] M. Benayoun *et al.* in “Proceedings of the Marbella Workshop on the tau-charm factory”, june 1993, to be published.