AB

EUROPEAN ORGANIZATION FOR NUCLEAR RESEARCH

CERN LIBRARIES, GENEVA

CERN-PPE/94-149 /
September 13th, 1994

CERN-PPE-94-149

see 9442

OBSERVATION OF THE $\omega \rightarrow \pi^0 \pi^0 \gamma$ DECAY

D. Alde³, F.G. Binon^{2*}, M. Boutemeur^{5**}, C. Bricman^{2**}, S.V. Donskov¹,
 M. Gouanère⁴, A.V. Inyakin¹, V.A. Kachanov¹, G.V. Khaustov¹, E.A. Knapp³,
 A.A. Kondashov¹, A.V. Kulik¹, G.L. Landsberg^{1***}, A.A. Lednev¹, V.A. Lishin¹,
 J.P. Peigneux⁴, S.A. Polovnikov¹, V.A. Polyakov¹, M. Poulet⁴, Yu.D. Prokoshkin¹,
 S.A. Sadovsky¹, V.D. Samoylenko¹, P.M. Shagin¹, A.V. Shtannikov¹, A.V. Singovsky¹,
 J.P. Stroot^{2**} and V.P. Sugonyaev¹

GAMS Collaboration : IHEP¹-IISN²-LANL³-LAPP⁴
 (Joint CERN - IHEP experiment)

Abstract

The rare radiative decay $\omega \rightarrow \pi^0 \pi^0 \gamma$ has been detected with the GAMS-2000 multi-photon spectrometer at the IHEP 70 GeV accelerator. The $\omega \rightarrow \pi^0 \gamma$ decay events have been used for normalization. The measured branching ratio is $BR(\omega \rightarrow \pi^0 \pi^0 \gamma) = (7.2 \pm 2.6) \cdot 10^{-5}$.

(Submitted to Physics Letters B)

-
- 1) Institute for High Energy Physics, Protvino 142284, Russia.
 - 2) Institut Interuniversitaire des Sciences Nucléaires, B-1050 Brussels, Belgium.
 - 3) Los Alamos National Laboratory, Los Alamos, NM 87544, USA.
 - 4) Laboratoire d'Annecy de Physique des Particules, F-74019 Annecy-le-Vieux, France.
 - 5) Université de Montréal, Laboratoire de Physique Nucléaire, H3C, 3J7 Montréal, Canada.
 - *) Mailing address : Université Libre de Bruxelles, CP 229, B-1050 Brussels, Belgium.
 - **) Mailing address : CERN, CH-1211 Geneva 23, Switzerland.
 - ***) Mailing address : SUNY at Stony Brook, NY 11794-3800, USA.

The first observation of the rare radiative decay of the ω meson

$$\omega \rightarrow \pi^0 \pi^0 \gamma, \quad (1)$$

and the measurement of its branching ratio are presented here by the GAMS Collaboration. The decays of light vector mesons (ρ , ω , ϕ) into a pair of neutral pseudoscalar mesons and a photon have been discussed for more than three decades [1-4]. None had previously been observed experimentally [5]. In the chiral perturbation models which are currently used for evaluation of these processes, the decay (1) is described by the vector meson dominance model with a relatively small inner kaon loop contribution [4].

The data were collected at the IHEP 70 GeV proton accelerator during three high-statistics runs, one in 1984 (I), and two in 1987 (II, III) [6, 7]. Five photons of the decay (1) were detected with the GAMS-2000 multiphoton spectrometer.

The charge-exchange reaction

$$\pi^- p \rightarrow \omega n \quad (2)$$

at the π^- beam momentum of 38 GeV/c has been used as a source of monoenergetic omegas. The detailed description of the apparatus, the event reconstruction procedure, and the data analysis algorithms is given elsewhere [6, 8].

The target-to-GAMS distance differed during the three runs, resulting in different spectrometer acceptance and mass resolution [6, 7]. The data were thus analysed separately for each run.

The $\pi^0 \pi^0 \gamma$ candidates have been selected out of a grand total of 10^8 charge-exchange events by requiring five photons in the final state with the GAMS reconstruction program. A series of cuts is then applied to suppress both instrumental and physical backgrounds (see Refs. [6, 7] for details). The contribution of one-pion exchange processes is lowered through a cut of small momentum transfer squared events ($-t > 0.1$ (GeV/c)²) in order to enhance the ω signal [6].

The major physical background, which originates from copiously produced $3\pi^0$ and $\eta\pi^0\pi^0$ systems with one undetected photon, gives no enhancement in the ω mass region. It is strongly suppressed by the above cuts.

The $\omega \rightarrow \pi^0 \eta \gamma$ decay could fake the process (1), but its branching ratio is expected to be two orders of magnitude lower [4].

The $K^{*0}(892) \rightarrow K_s^0 \pi^0 \rightarrow 3\pi^0 \rightarrow 6\gamma$ decay might contribute to the mass spectrum around the ω mass when the K_s^0 decays into two neutral pions at a certain distance from the spectrometer target, with one photon being lost (K^{*0} is produced in a pair with a hyperon, a small fraction of which might escape detection by the target guard system [7]). A Monte Carlo study shows that after all cuts the contribution of this process to the measured $\pi^0 \pi^0 \gamma$ mass spectrum is negligible.

Further selection of the reaction

$$\pi^- p \rightarrow \pi^0 \pi^0 \gamma n \quad (3)$$

events is based on a kinematical 3C fit (masses of both π^0 's and of the recoil neutron being fixed). The χ^2 values are computed for all possible combinations in which the invariant masses of two γ -pairs, composed of different photons, fall into the π^0 mass range (60 - 210 MeV). Only the events with $\chi^2 < 9$ (CL > 97%) are used in further analysis.

Both the raw $\pi^0 \pi^0 \gamma$ mass spectrum and the efficiency-corrected spectrum exhibit a narrow peak in the ω mass region. The spectra are fitted with a sum of a Gaussian and a polynomial background.

Fig. 1 shows the spectrum summed over runs I and II which have similar mass resolution. The mass of the peak in this figure is measured to be 785 ± 3 MeV and its width is dominated by the instrumental GAMS resolution. The similar peak obtained in run III is 20% wider (better acceptance and signal-to-background ratio but lower mass resolution in this run).

The total number of the $\omega \rightarrow \pi^0 \pi^0 \gamma$ events is 40 ± 14 . The branching ratio of this rare ω decay mode has been evaluated through normalization with the abundant $\omega \rightarrow \pi^0 \gamma$ decay events that have been simultaneously measured in the present experiment ($1.5 \cdot 10^5$ virtually background-free events):

$$\frac{\text{BR}(\omega \rightarrow \pi^0 \pi^0 \gamma)}{\text{BR}(\omega \rightarrow \pi^0 \gamma)} = (8.5 \pm 2.9) \cdot 10^{-4}. \quad (4)$$

With $\text{BR}(\omega \rightarrow \pi^0 \gamma) = (8.5 \pm 0.5) \cdot 10^{-2}$ [5] this gives :

$$\text{BR}(\omega \rightarrow \pi^0 \pi^0 \gamma) = (7.2 \pm 2.6) \cdot 10^{-5}. \quad (5)$$

Theoretical estimates of this value vary from $3 \cdot 10^{-5}$ to $9 \cdot 10^{-5}$ [2-4].

A search for several other rare ω meson decays was carried out along with this measurement. The following upper limits (at 90% CL) were obtained for the C-violating decays :

$$\text{BR}(\omega \rightarrow 3\pi^0) < 1.5 \cdot 10^{-3}, \quad (6)$$

$$\text{BR}(\omega \rightarrow \eta \pi^0) < 1 \cdot 10^{-3}; \quad (7)$$

and for the direct 3- γ decay :

$$\text{BR}(\omega \rightarrow 3\gamma) < 8 \cdot 10^{-4}. \quad (8)$$

This work was supported, in part, by the Soros Foundation grant awarded by the American Physical Society.

References

- [1] P. Singer, Phys. Rev. 128 (1962) 2789; 130 (1963) 2441.
- [2] S. Fajfer, R.J. Oakes, Phys. Rev. D42 (1990) 2392.
- [3] A. Bramon et al., Phys. Lett. B289 (1992) 97.
- [4] A. Bramon et al., The DAΦNE Physics Handbook, Frascati, 1992, p. 467.
- [5] Particle Data Group, Phys. Rev. D45 (1992) VII.15.
- [6] D. Alde et al., Yad. Fiz. 56 (1993) 137; Z. Phys. C61 (1994) 35.
- [7] D. Alde et al., Yad. Fiz. 54 (1991) 745; Phys. Lett. B284 (1992) 455.
- [8] F.G. Binon et al., Nucl. Instr. Meth. A248 (1986) 86; in : Cherenkov detectors and their applications in science and techniques. Ed. Nauka, Moscow, 1990, p. 149.

Figure caption

Fig. 1. The $\pi^0\pi^0\gamma$ mass spectrum in the reaction (3) corrected for efficiency (the combined result of runs I and II). The curve shows the fit result.