J7A

**X-ray Emission from the Historical Supernovae in the
Spiral Galaxy NGC 6946: SN1980K and SN1968D
Recovered?**

Eric M. Schlegel

8u 8445

Laboratory for High Energy
Astrophysics

SCAN-9411090

CERN LIBRARIES, GENEVA

**High Energy Astrophysics Science
Archive Research Center**

**NASA Goddard Space Flight Center
Greenbelt, MD 20771**

X-ray Emission from the Historical Supernovae in the Spiral Galaxy NGC 6946: SN1980K and SN1968D Recovered?

Eric M. Schlegel¹

Code 668, NASA-Goddard Space Flight Center, Greenbelt, MD 20771

E-mail: eric@heasfs.gsfc.nasa.gov (I)

Received _____; accepted _____

¹Research Scientist, Universities Space Research Association

ABSTRACT

This paper describes a study of the *ROSAT* PSPC observation of the spiral galaxy NGC 6946 to search for X-ray emission from the six historical supernovae that have occurred in that galaxy. At the age of these supernovae, all could be emitting X-rays due to circumstellar interactions. SN1980K may be present based upon the hardness ratio and the coincidence of an X-ray source with the optical position. Its luminosity agrees with a recent prediction by Chevalier & Fransson. The situation for SN1968D, apparently recovered in VLA data, is less clear due to source confusion, but suggests SN1968D has not been recovered. Only upper limits are available for the remaining four supernovae that have occurred in this galaxy.

Subject headings: supernovae: individual: SN1917A, SN1939C, SN1948B, SN1968D, SN1969P, SN1980K; X-rays: stars

Submitted:

1. Introduction

X-ray emission may be detected from supernovae by at least three possible processes during their evolution (c. f., Schlegel 1994a). First, there is a prompt thermal burst due to shock breakout. If this burst is sufficiently hot, X-rays may be produced. Second, Compton scattered γ -rays from the radioactive decay of nuclides synthesized in the explosion produce X-rays at least down to the photoelectric absorption cutoff at ~ 16 -20keV. Third, the interaction of a shock with circumstellar matter generates X-ray emission as well as radio emission (Chevalier 1990 and references therein). Two possible X-ray emitting regions are present in the circumstellar interaction model: at the outgoing shock where X-rays (up to ~ 100 keV) are generated; at the reverse shock softer X-rays (~ 1 keV) are produced.

Observations of supernovae have yielded five detections to date (for a review, see Schlegel 1994a). Only two supernovae have been seen at early times (< 50 days old), both almost certainly due to circumstellar interaction: SN1980K (Canizares et al. 1982) and SN1993J (Zimmermann et al. 1994). SN1994I in NGC 5194 may be a promising addition to this list based upon its early radio detection (Rupen et al. 1994). Three supernovae have been seen at later times: SN1987A, due to Compton-scattered γ -rays (Kumagai et al. 1989); SN1986J (Bregman & Pildis 1992) and SN1978K (Ryder et al. 1993) due to circumstellar interactions. Two of these sources (SN1986J, SN1978K) are still under observation as they are extremely X-ray luminous (10^{39-40} ergs s^{-1}) in the 0.5-2.0keV band. The X-ray light curve of SN1978K is under construction and will be the first late-time X-ray light curve of a supernova (Schlegel et al. 1994). Other detections of historical supernovae may also be reported soon. For example, Pietsch (1992) notes that some X-ray emission is observed "near" the sites of SN1970G and SN1951H in M101, although the details have not yet been published.

At some point, however, a supernova becomes a supernova remnant. Is the transition

observable? Supernova remnants (SNRs) are known X-ray sources with luminosities generally in the range 10^{34-36} ergs s^{-1} (c.f. Seward 1989). A few SNRs are much brighter with luminosities approaching 10^{38-40} ergs s^{-1} . This list of luminous SNRs includes the SNR in NGC 4449 (Blair et al. 1983), N49 in the LMC (Vancura et al. 1992), and the SNR in NGC 6946 (Schlegel 1994b). Is there a connection between these luminous SNRs and luminous supernovae such as SN1986J and SN1978K? What is the nature of the transition?

Two approaches exist to observe the transition or to understand whether a connection exists. First, one can follow a single supernova (or a small set) from supernova to remnant. If one chooses wisely, one may live sufficiently long to see this transition. The second approach examines as many supernovae as possible. The supernovae will exist in different stages of their evolution, but the observer may at least see if any have made the transition and exhibit properties similar to known SNRs. Two galaxies exist that are ideally suited for such a study: NGC 5236 (M83) and NGC 6946. Both galaxies have six historical supernovae each spread over ~ 80 years. Each galaxy also happens to be the parent of one of the relatively bright supernovae that have occurred in the last decade or so: SN1983N in NGC 5236 and SN1980K in NGC 6946.

I describe here the search for X-ray emission from the historical supernovae in NGC 6946 using *ROSAT* PSPC data. The paper is organized as follows. Section 2 describes the observations and analysis. Section 3 details the SN1980K observation, Section 4 describes SN1968D, and Section 5 describes the upper limits on the remaining four supernovae. A brief summary ends the paper. In this paper, I use a value of 5.1Mpc (deVaucouleurs 1979) for the distance to NGC 6946.

2. Observations

The *ROSAT* PSPC observed NGC 6946 on 1992 June 16-21, obtaining a total exposure time of 36713 s scattered over 14 observation intervals. The mean count rate in each observation interval was calculated, as well as the mean of these means. Observation intervals in which an interval's mean count rate was more than 3σ from the total mean count rate were screened out. A total of 70 s failed this criterion, leaving a total exposure time of 36643 s. The background light curve showed a relatively constant rate of ~ 5 counts s^{-1} for each observation interval. No evidence for short- or long-term enhancements, as described in Snowden et al. (1994), was seen. No systematic effects were seen in the master veto rate (Plucinsky et al. 1993; Snowden et al. 1992). More complete details of the data reduction are in Schlegel (1994c).

The positions of the historical supernovae in NGC 6946 are known to rather high accuracy (Table 1). The values were converted from 1950 coordinates to J2000. These coordinates must then be "corrected" to the PSPC image coordinates. The positional accuracy of the PSPC was tested by comparing the optical positions of three sources within 20' of NGC 6946 with their corresponding X-ray positions. Each of the 3 sources had at least several tens of counts, so the position centroids should be accurate to $\sim 8''$ - $10''$ (Briel et al. 1988). Two of the sources are known *Einstein* sources (Harris et al. 1990; Hertz & Grindlay 1988), so the X-ray and optical correspondence is known. The resulting, average RA offset (optical - X-ray) was $+1^{\circ}.6$ and the resulting Dec offset was $-6.4''$. These values are typical of the boresight shifts of *ROSAT* (Kürster & Hasinger 1993). The corrections were then applied to the coordinates of the historical supernovae to obtain "raw" coordinates.

The raw coordinates were used to place the extraction apertures in their proper location on the PSPC image. Figure 1 shows the apertures on a background-subtracted, smoothed version (described in Schlegel 1994c). The apertures are all $24''$ in size, which is comparable to the overall FWHM of the PSPC point spread function, and which samples $\sim 90\%$ of the point spread function at 1keV (Hasinger et al. 1993). The counts were extracted

within the aperture and treated as “source” + background counts. The background was determined from a nearby, source-free region. Most of the historical supernovae happened to have occurred at a sizable distance from the nucleus, hence, most are in relatively source-free regions. SN1968D requires additional work as it lies atop the diffuse emission from the hot interstellar medium of NGC 6946 (Schlegel 1994c). For this source, the background extraction region was also taken from a region apparently free of point sources, but encircling a region in which the emission comes from the hot interstellar medium of NGC 6946. That location lies just north of the site of SN1968D. As all of the sources are essentially on-axis (the largest offset is SN1980K at $\sim 5'$), vignetting corrections, while applied, are essentially negligible.

Upper limits were computed using the Bayesian prescription of Kraft, Burrows, & Nousek (1991). The Bayesian approach assumes the measured data constrain the experimental hypothesis by applying *a priori* knowledge of the system being measured (for example, that the flux is never less than zero). The Kraft et al. approach is an approximation to the proper formulation. They assume the background count rate is infinitely well-measured. For a large field-of-view instrument such as the *ROSAT* PSPC, this is a reasonable approximation as a relatively large extraction region for the background may be defined. The required inputs for the Bayesian upper limit are the observed counts for the source plus background and the mean background counts. All upper limits were calculated for the 99% confidence level. The counts were then converted to fluxes by assuming a thermal bremsstrahlung spectrum of temperature 0.5 keV absorbed by the known galactic column to NGC 6946 ($N_H \sim 5.0 \times 10^{21} \text{ cm}^{-2}$ (Burstein & Heiles 1984)). A temperature of 0.5 keV was chosen as representative of the total radiative cooling function of shocked gas (e.g., Raymond, Cox, & Smith 1976) in which free-free and line emission are important based upon observations of SN1978K (Ryder et al. 1993) and SN1986J (Bregman & Pildis 1992). Both of these X-ray bright supernovae are of comparable age to SN1980K, so may

be in a similar evolutionary phase. The temperature range for shocked gas under the above conditions is $\sim 5 \times 10^5$ °K to $\sim 4 \times 10^7$ °K. Below ~ 3 keV, the X-rays produced by the reverse shock dominate the spectrum; their temperature is expected to be ~ 0.9 keV (Chevalier & Fransson 1994). As it turns out, the calculated fluxes are relatively insensitive to the exact value of the temperature or the exact nature of the spectrum, whether bremsstrahlung or a Raymond plasma. To obtain a flux, the normalization was increased or decreased until the reported counts in the simulated spectrum matched the number of counts from the upper limit calculation. Fluxes and luminosities were then calculated for the 0.5-2.0 keV band and at 2.0 keV. All of these entries are listed in Table 1.

3. SN1980K

SN1980K is among the best-studied supernovae. Of the six historical supernovae in NGC 6946, it was the brightest by ~ 2 magnitudes. It lies well away from the nucleus ($\sim 325''$) to the southeast. SN1980K was a Type II-Linear supernova. SN1980K is special for two reasons. First, until the X-ray detection of SN1993J in M81 (Zimmermann et al. 1994 and references therein), it was the only supernova to have been seen near optical maximum. Second, it has been one of a handful of strong radio supernovae the observations of which have led to considerable understanding of the circumstellar interaction mechanism (Weiler et al. 1992). It was detected in the radio at 6 cm ~ 35 days after the optical maximum (Weiler et al. 1986). Its radio light curve was monitored until 1990 December when it faded below the VLA detection limit (Weiler et al. 1992). The 10-year radio light curves show that the Chevalier circumstellar interaction shell model (Chevalier 1990 and references therein) provides a good description of the data. That model predicts the radio light curve should follow the relation

$$S(mJy) = K_1 \left[\frac{\nu}{5GHz} \right]^\alpha \left[\frac{t - t_0}{1day} \right]^\beta e^{-\tau}, \tau = K_2 \left[\frac{\nu}{5GHz} \right]^{-2.1} \left[\frac{t - t_0}{1day} \right]^\delta, \quad (1)$$

with $\delta = \alpha - \beta - 3$ (Chevalier 1984). Given that the radio light curve represents the interaction of the shock with circumstellar matter, X-ray emission is also expected. An *Einstein* IPC observation of SN1980K detected an $\sim 5\sigma$ X-ray source at the supernova's location ~ 35 days after the explosion (Canizares et al. 1982). A subsequent observation at ~ 50 days after maximum showed a $\sim 2\sigma$ source at that location. The resulting decline by a factor of ~ 2 in ~ 15 days was consistent with the expectations of the model. Chevalier & Fransson (1994) predict SN1980K to be a $\sim 2 \times 10^{38}$ ergs s $^{-1}$ source in the 0.2-4keV X-ray band at an age of 10 years. The predicted temperature is ~ 0.9 keV.

The prediction of Chevalier & Fransson can be tested as the prediction was made for an X-ray band largely covered by *ROSAT*. Chevalier & Fransson adopted a distance of 7.5Mpc to NGC 6946. With their predicted luminosity, this gives a flux of $\sim 3 \times 10^{-14}$ ergs s $^{-1}$ cm $^{-2}$. The prediction was given in the 0.2-4keV band, essentially that of the *Einstein* IPC. I simulated a bremsstrahlung spectrum and a Raymond-Smith thermal plasma spectrum, both at the predicted temperature and absorbed by a column of $\sim 3\text{--}5 \times 10^{21}$ cm $^{-2}$. I adjusted the model normalizations until the flux in the 0.2-4.0keV band matched the predicted flux. The flux in the 0.2-2.0keV band was calculated, yielding a value of $\sim 2 \times 10^{-14}$ ergs cm $^{-2}$ s $^{-1}$. This flux can be expected to generate ~ 29 counts in the PSPC for a $\sim 5\sigma$ detection, assuming no contribution to the background from NGC 6946 itself. The PSPC observation of NGC 6946 occurred at a supernova age of ~ 11.6 years. As the predicted evolution is relatively slow at an age of ~ 10 years, the expected luminosity should not differ significantly from the above prediction.

The position of SN1980K agrees quite well with a “lump” of X-ray emission. As the supernova occurred $> \sim 5'$ from the nucleus, the quantity of diffuse emission present at that radius from the galaxy itself is small (Schlegel 1994c), so source confusion is reduced. The positional difference between SN1980K and the nearby “lump” is $\sim 8''$ ($\Delta\alpha = 1''.1$; $\Delta\delta = 1''.2$). The number of net counts in the lump is 18.4, corresponding to a 99% upper limit on the

flux of $\sim 7 \times 10^{-15}$ ergs s $^{-1}$ cm $^{-2}$. The hardness ratio is +0.36. The number of counts at the position of SN1980K is 20.5 with a hardness ratio of +0.49. The hardness ratio is defined as $\frac{H-S}{H+S}$, where S = counts in channels 20-90, ~ 0.2 -0.9keV; and H = counts in channels 91-210, ~ 0.9 -2.1keV. The error range on the extracted counts for SN1980K and the lump easily cover this range of hardness. While 20.5 counts barely represents a detection ($\sim 3.5\sigma$), some consideration should be given to whether the "detection" is consistent with SN1980K.

A hardness ratio of $\sim +0.4$ seems low for a source expected to be relatively hard (kT ~ 0.9 ; Chevalier & Fransson 1994). A brief digression is necessary, however, to test one's intuition on this point. A simulation is necessary to investigate the range of hardness ratios possible for various combinations of temperature and column. A simulated spectrum, assuming a bremsstrahlung continuum or a Raymond thermal plasma model at various temperatures and absorbed by several different values for the galactic column to the galaxy, yielded the results presented in Tables 2 and 3. The observed values of $\sim +0.4$ correspond to very high temperatures at low columns or low temperatures with a high column. Unfortunately, because there are so few observed counts, the range of possible hardness ratios is large. The known column to NGC 6946, even though it is not known precisely, does eliminate, however, the highest and lowest entries in Table 2 for the bremsstrahlung spectrum; the Raymond spectrum is less constrained. A reasonable match to the observed hardness may be made if the column is closer to 3×10^{21} cm $^{-2}$ than to 5×10^{21} cm $^{-2}$; the temperature then falls in the range 0.7-0.9keV regardless of the continuum model. Such a value for the column is within the errors determined from the spectral fits to the very luminous supernova remnant in NGC 6946 (Schlegel 1994b). The lower column also corresponds more closely with the E(B-V) value to SN1980K (0.37, Buta 1982). The hardness ratio thus lends additional evidence that SN1980K has been detected.

The difference between the predicted X-ray flux and the upper limit for SN1980K is about a factor of 1.5-2, well within the errors. Chevalier & Fransson (1994) show the

luminosity is related to $\frac{\dot{M}v_w^{-1}v_s^3}{(n-3)^3}$, where $\dot{M}$ is the mass loss rate, v_w is the wind velocity, v_s is the shock velocity, and n is the power law index. $\dot{M}v_w^{-1}$ is measured by the $H\alpha$ emission line to be ~ 3 . A slightly lower value of v_s by ~ 2 would decrease the expected luminosity; so also would a small change in the index. Several of the parameters are sensitive to n ; observations to date have not placed overly restrictive limits on its value. Hence, the difference between prediction and observation is even more negligible at this stage of our understanding.

Of greater concern is the potential for source confusion. The probability of a source of the "correct" flux falling in a given detection cell of size $30''$ by $30''$ is $\sim 4 \times 10^{-3}$, using the log N-log S relation of Hasinger, Schmidt, & Trümper (1991). The probability is actually somewhat higher since SN1980K lies near a galaxy which locally may not follow the log N-log S relation. Few other types of sources are capable of mimicking SN1980K. Stars, cataclysmic variables, and stellar coronal sources are insufficiently luminous by orders of magnitude. X-ray binaries have the correct luminosity but are usually much harder. Perhaps the only type of source that could mimic SN1980K is another SNR. Examination of the optical recovery image of SN1980K (Fesen & Becker 1990) shows no apparent SNR within at least $10\text{-}20''$ of the position of SN1980K.

On the basis of the available evidence, I tentatively conclude that SN1980K has been detected. Certainty, however, will require additional observations to establish definitively whether SN1980K is present. For example, detecting X-ray emission lines typical of a supernova remnant would establish the case. Unfortunately, it is likely that SN1980K is fading in the X-ray band as it already has in the radio (Weiler et al. 1992), so additional observations may not be possible until a more sensitive instrument is available.

The possible detection of SN1980K may be placed in a larger context. Weiler et al. (1986) compiled the 6cm fluxes of supernovae and selected remnants and plotted them against the log of the age of the supernova. The result was a 6cm radio "evolution" curve

for the ensemble of supernovae then known. A start at a similar figure has been made in Figure 3; it shows a 2keV evolution curve for SN1980K and a few well-known supernova remnants of known age. A more complete version is in Schlegel (1994a). The *Einstein* points represent a fit to the spectra presented in Canizares et al. (1982). The model used for the fitting was the thermal bremsstrahlung model described above, using $kT \sim 0.5$ keV and $N_H \sim 3 \times 10^{21} \text{ cm}^{-2}$. The flux was calculated from the model, but is relatively insensitive to the choice of model. The figure clearly shows gaps exist at the few year scale and from ~ 10 years to ~ 300 years. The SN1980K observations are known to lie near the limit of detection; none of the well-known supernova remnants would be visible were they to have occurred in a nearby galaxy such as NGC 6946. The SN1980K detection, if real, shows a luminosity approximately 100 times larger than the ensemble of known remnants. It is likely that an evolution plot such as Figure 3 will reveal phases of evolution from supernova to remnant once we are able to place more observations on the figure. The observed phases can then be used to test the theoretical models.

4. SN1968D

SN1968D occurred closest to the nucleus, $\sim 49''$ northeast of the galaxy's optical center on 1968 February 29. It was identified as a Type II supernova. In the PSPC data, the supernova lies $\sim 30''$ from a relatively soft source labeled source S4 in Schlegel (1994c). The fitted temperature of S4 of $\sim 0.2-0.5$ keV, while not precisely measured, is consistent with a young supernova remnant (among other possibilities, such as stellar coronal sources). The hardness ratio for S4 is $\sim +0.1$. The local background is very high here due to the diffuse emission of the galaxy, so the X-ray "upper limit" is unrestrictive. Source confusion is a larger problem. Given that S4 has ~ 200 counts, its position is likely to be accurate to $\sim 8-10''$ (Briel et al. 1988). A tentative detection of SN1968D in data from the VLA has

been made recently (Van Dyk et al. 1994) The position of the radio source is 20:34:58.4, +60:09:32.4 (J2000) which corresponds to the optical position to $\sim 1''$. Based upon the available data, it is not clear whether the X-ray source corresponds to the radio source. Attempts are being made to identify the optical counterpart of source S4. If these succeed, the identification of the X-ray emission in this region should then be possible. Note, however, that the coordinate shift necessary to register the optical and X-ray images increases the separation between the potential X-ray counterpart and the optical position of SN1968D. This suggests the X-ray source is not the counterpart to SN1968D.

5. Upper Limits on the Remaining Supernovae

Upper limits are reported for the remaining four supernovae. In this section each supernova is discussed in turn along with a brief summary of its optical behavior.

5.1. SN1917A

SN1917A was discovered on 1917 July 19. It lies $\sim 111''$ from the nucleus in the region of low X-ray surface brightness that exists south of the nucleus (Schlegel 1994c). Based upon a photographic spectrum by Ritchey (1917), SN1917A is a Type II supernova. The X-ray emission at its location is completely consistent with the X-ray background local to the NGC 6946 region.

5.2. SN1939C

SN1939C was discovered on 1939 July 17 by F. Zwicky and identified by him as a Type I supernova. The supernova lies $\sim 216''$ from the nucleus on the outer arm west of

the nucleus. Little else is apparently known about this object. The X-ray emission from its location is also consistent with the X-ray background local to NGC 6946. If SN1939C is a Type Ia supernova (no hydrogen and believed to be from a low-mass progenitor), then the upper limit is relatively unrestrictive. Type Ia supernovae are not expected to be X-ray emitters in the few decades after the detonation due to the low density of the circumstellar matter. At outburst, they may be X-ray emitters but, to date, only upper limits exist (Schlegel 1994a; Schlegel & Petre 1993). The X-ray behavior of a Type Ib supernova (no hydrogen, but believed to be from a massive progenitor) should follow that of a Type II supernova (Chevalier 1984).

5.3. SN1948B

SN1948B was discovered on 1948 June 15 at a position $\sim 230''$ east-northeast of the nucleus on the outer limits of the prominent north arm. Based upon the spectrum (Mayall 1948) and the light curve (Barbon, Ciatti, & Rosino 1979; Patat et al. 1993), the supernova is a Type II-Plateau object. X-ray emission from the location of SN1948B is consistent with the X-ray emission of the spiral arm; that emission could be diffuse emission from a hot interstellar medium or the summed emission of unresolved point sources. The first explanation appears to be the most likely at this point (Schlegel 1994c).

5.4. SN1969P

SN1969P occurred on 1969 December 11. The type of the supernova was never determined, apparently, because no optical follow-up was ever done. The object lies $\sim 180''$ south of the nucleus in a region of low X-ray surface brightness. The X-ray upper limit is unrestrictive if the supernova is a Type Ia.

6. Summary Discussion

Estimates may be made for the average mass loss over the evolutionary life of these historical supernovae. Chevalier (1984) showed that an estimate of $\dot{M}$ may be made from $L = 10^{40} A^2 (\frac{t}{10 \text{ days}})^{-1}$, where A is defined by $\frac{\dot{M}}{v_w} = 10^{-6} A M_{\odot} \text{ yr}^{-1} (\text{km s}^{-1})^{-1}$ with v_w the wind speed, usually assumed to be 10 km s^{-1} . This relation applies to SN Ia only, as it assumes a density profile with a power law index of 7. A similar relation is available for SN II (e.g., Fransson, Lundqvist, & Chevalier 1994). Since the upper limits on the luminosities are all approximately $2 \times 10^{37} \text{ ergs s}^{-1}$ and the ages of the historical supernovae fall in the range of few 10^3 to few 10^4 days, the values of $\dot{M}$ are all approximately $1\text{--}2 \times 10^{-8} M_{\odot} \text{ yr}^{-1}$ for the SN Ia and $\sim 5 \times 10^{-6}$ to 5×10^{-7} for the SN II. These are very small values, and clearly represent a mass loss averaged over the full life of these evolving remnants. The possible detections of SN1980K and SN1968D clearly indicate that an average mass loss does not reflect the conditions surrounding these historical supernovae. Furthermore, a high mass loss rate cannot be maintained without losing a substantial fraction of the progenitor. One thus expects the X-ray and radio flux to reflect the duration of mass loss; such behavior was observed in the optical [O III] line emission of SN1957D (Long et al. 1992).

In summary, a 36ksec *ROSAT* PSPC observation of the spiral galaxy NGC 6946 has likely detected SN1980K, may have detected SN1968D, and places upper limits on the X-ray emission of the remaining four historical supernovae known. Observations are starting to close the gap in our understanding of the evolution of a supernova to a remnant.

7. Acknowledgments

I thank the referee, Dr. Roger Chevalier, for helpful comments that improved this paper and for a reminder about SN1957D. This research was supported by a grant from the

US *ROSAT* Guest Observer Program funded by NASA. The NASA Extragalactic Database (NED) and the SIMBAD database, operated by the CDS, Strasbourg, France, were used in this work.

REFERENCES

- Barbon, R., Ciatti, F., & Rosino, L. 1979, *A&A*, 72, 287
- Blair, W., Kirshner, R., & Winkler, P. F. 1983, *ApJ*, 272, 84
- Bregman, J. & Pildis, R. 1992, *ApJ*, 398, L107
- Briel, U., Pfeffermann, E., Hartner, G., & Hasinger, G. 1988, *X-ray Instrumentation in Astronomy II (Proc SPIE)*, 982, 401
- Burstein, D. & Heiles, C. 1984, *ApJS*, 54, 33
- Buta, R. 1982, *PASP*, 94, 578
- Canizares, C., Kriss, G., & Feigelson, E. 1982, *ApJ*, 253, L17
- Chevalier, R. 1990 in *Supernovae*, ed. A. Petschek, (Berlin: Springer-Verlag), 91
- Chevalier, R. 1984, *ApJ*, 285, L63.
- Chevalier, R. & Fransson, C. 1994, *ApJ*, 420, 268
- Chugai, N. 1993, *ApJ*, 414, L101
- deVaucouleurs, G. 1979, *ApJ*, 227, 729
- Fesen, R. & Becker, R. H. 1990, *ApJ*, 351, 437
- Fransson, C., Lundqvist, P., & Chevalier, R. 1994, *ApJ* preprint
- Harris, D. et al. 1990, *The Einstein Observatory Catalog of IPC X-ray Sources*, (Cambridge: Smithsonian Astrophysical Observatory)
- Hasinger, G., Turner, T. J., George, I., & Boese, G. 1992, *NASA-OGIP Calibration Memo* 92-001
- Hasinger, G., Schmidt, M., & Trümper, J. 1991, *A&A*, 246, L2
- Hertz, P. & Grindlay, J. 1988, *AJ*, 96, 233

- Kraft, R., Burrows, D., & Nousek, J. 1991, *ApJ*, 374, 344
- Kumagai, S., Shigeyama, T., Nomoto, K., Itoh, M., Nishimura, J., & Tsuruta, S. 1989, *ApJ*, 345, 412
- Kürster, M. & Hasinger, G. 1993, Internal MPE memo, 1993 Mar 10
- Long, K., Winkler, P. F., & Blair, W. P. 1992, *ApJ*, 395, 632
- Mayall, N. 1948, *PASP*, 60, 266
- Patat, F., Barbon, R., Capellaro, E., & Turatto, M. 1993, *A&AS*, 98, 443
- Pietsch, W. 1992 in *Physics of Nearby Galaxies: Nature or Nurture?*, ed. T. Thuan et al. (France: Editions Frontieres), 67
- Plucinsky, P., Snowden, S., Briel, U., Hasinger, G., & Pfeffermann, E. 1993, *ApJ*, 418, 519
- Raymond, J., Cox, D., & Smith, B. 1976, *ApJ*, 204, 290
- Richmond, G. 1917, *Harvard Coll. Obs. Bull.*, 641, July 28
- Rupen, M., Sramek, R., VanDyk, S., Weiler, K., & Panagia, N. 1994, *IAU Circular* 5963
- Ryder, S., Staveley-Smith, L., Dopita, M., Petre, R., Colbert, E., Malin, D., and Schlegel, Eric M. 1993, *ApJ*, 416, 167
- Schlegel, Eric M. 1994a, in preparation for *RepProgPhys*
- Schlegel, Eric M. 1994b, *ApJ*, 424, L99
- Schlegel, Eric M. 1994c, *ApJ*, 4xx, 000 (20 October)
- Schlegel, Eric M. & Petre, Robert 1993, *ApJ*, 412, L29
- Schlegel, Eric M., Colbert, E., & Petre, R. 1994, in preparation
- Seward, F. 1989, *SpSciRev*, 49, 385
- Snowden, S., McCammon, D., Burrows, D., & Mendenhall, J. 1994, *ApJ*, 424, 714
- Snowden, S., Plucinsky, P., Briel, U., Hasinger, G., & Pfeffermann, E. 1992, *ApJ*, 393, 819

- Vancura, O., Blair, W., Long, K., & Raymond, J. 1992, ApJ, 394, 158
- Weiler, K., Van Dyk, S., Panagia, N., & Sramek, R. 1992, ApJ, 398, 248
- Weiler, K., Sramek, R., Panagia, N., van der Hulst, J., & Salvati, M. 1986, ApJ, 301, 790
- Zimmermann, H. et al. 1994, Nature, 367, 621

TABLE 1. Upper Limits on the Historical Supernovae

SN	RA (2000)	Date of	Mag	Source	Bkgrd	99% UL	Flux	
Name	Dec (2000)	opt. max	at max	Cts ^a	Cts ^a	Counts	Limit ^b	L_X^c
SN1980K ^d	20:35:30.3	1980 Oct 31	V11.4	31	10.5±1.9	37.2	1.0(-14)	3.1(37)
	+60:06:23.6							
SN1969P	20:34:51.3	1969 Dec 11	PG13.9	19	11.3±4.4	20.6	8.3(-15)	2.6(37)
	+60:06:14.9							
SN1968D	20:34:58.4	1968 Feb 29	PG13.5	96	71.9±9.5	50.6	2.0(-14)	6.2(37)
	+60:09:32.4							
SN1948B	20:35:21.5	1948 Jun 15	PG14.9	37	34.0±6.9	20.7	8.3(-15)	2.6(37)
	+60:10:16.6							
SN1939C	20:34:23.0	1939 Jul 17	PG13.0	24	23.1±5.8	16.2	6.5(-15)	2.0(37)
	+60:09:37.3							
SN1917A	20:34:46.9	1917 Jul 19	PG14.6	31	32.2±6.7	16.6	6.7(-15)	2.2(37)
	+60:07:29.7							

Notes to Table 1

^aBoth “source” and background counts were extracted in the energy range 0.2-2.0keV.

^bUnits are $\text{ergs s}^{-1} \text{ cm}^{-2}$. The flux upper limit comes from the spectrum with the equivalent number of counts. The energy range is 0.5-2.0keV.

^cUnits are ergs s^{-1} in the 0.5-2.0keV range. The assumed distance is 5.1Mpc (de Vaucouleurs 1979).

^dThe “upper limit” is included here for completeness, or if the reader does not believe the evidence for the detection of SN1980K.

TABLE 2. Hardness Ratios from Simulated Spectra: Bremsstrahlung^a

N_H (cm ⁻²)	kT (keV)									
	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9	1.0	1.5
1x10 ²¹	-0.82	-0.59	-0.38	-0.23	-0.12	-0.04	+0.03	+0.08	+0.12	+0.25
3x10 ²¹	-0.51	-0.13	+0.12	+0.36	+0.26	+0.43	+0.48	+0.51	+0.54	+0.63
5x10 ²¹	-0.16	+0.25	+0.46	+0.57	+0.64	+0.68	+0.71	+0.73	+0.75	+0.80
7x10 ²¹	+0.16	+0.52	+0.67	+0.75	+0.79	+0.82	+0.84	+0.85	+0.86	+0.89

Notes to Table 2

^aThe spectral model was a thermal bremsstrahlung of the tabulated temperature absorbed by the tabulated galactic column.

TABLE 3. Hardness Ratios from Simulated Spectra: Raymond Plasma^a

N_H (cm ⁻²)	kT (keV)									
	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9	1.0	1.5
1x10 ²¹	-0.86	-0.68	-0.55	-0.43	-0.27	-0.04	+0.15	+0.41	+0.46	+0.53
3x10 ²¹	-0.71	-0.48	-0.35	-0.22	-0.06	+0.17	+0.37	+0.61	+0.67	+0.78
5x10 ²¹	-0.50	-0.26	-0.13	-0.01	+0.15	+0.36	+0.53	+0.74	+0.79	+0.89
7x10 ²¹	-0.26	-0.02	+0.09	+0.20	+0.35	+0.53	+0.67	+0.83	+0.87	+0.94

Notes to Table 3

^aThe spectral model was a Raymond-Smith thermal plasma of the tabulated temperature absorbed by the tabulated galactic column.

Figure Caption

Figure 1 - A background-subtracted, smoothed image of NGC 6946 in the *ROSAT* band (0.2-2.0keV). North is up and east is left. The positions of the historical supernovae are indicated by circles 24" in diameter. This image emphasizes the diffuse background present in the galaxy.

Figure 3 - The 2keV light curve of SN1980K and a few supernova remnants. A more complete version of this plot is in Schlegel (1994a). The SN1980K points labeled "E" are from re-fits to the *Einstein* observations described in Canizares et al. (1982). The SN1980K point labeled "R" is the *ROSAT* PSPC observation.