BB

8x8446

LABORATOIRE DE PHYSIQUE NUCLEAIRE

UNITE ASSOCIEE AU CNRS/IN2P3 (UA 57)

2, rue de la Houssinière 44072 NANTES CEDEX 03

IN2P3

SCAN-941196

CERN LIBRARIES, GENEVA

When and how are fragments formed in heavy ion collisions ?

C.O. Dorso and J. Aichelin

Rapport Interne LPN - 94-09

UNIVERSITE DE NANTES

When and how are fragments formed in heavy ion collisions?

C.O.Dorso

Universidad de Buenos Aires

Departamento de Física, FCEN-UBA, Pab.1, Ciudad Universitaria

1428 Buenos Aires, Argentina

J.Aichelin

Laboratoire de Physique Nucléaire, CNRS/IN2P3 Université de Nantes,

2, Rue de la Houssinière, 44072 Nantes Cedex 03, France

Abstract : Using the Quantum Molecular Dynamics (QMD) approach for the simulation of nucleus nucleus collisions we compare the fragmentation pattern obtained with the standard minimum spanning tree(MST) procedure with that obtained by the Early Cluster Recognition Algorithm (ECRA) formalism. We find, that the ECRA algorithm gives the same clusters as the MST but is able to identify them much earlier, already shortly after the initial phase of violent collisions between projectile and target nucleons has passed. These findings rule out any coalescence description of fragment formation. We study the time evolution of the fragmentation pattern, the time correlation of the fragment constituent nucleons and the role of collisions in the early formation of clusters.

1 Introduction

The study of fragmentation processes in highly excited finite systems is a topic of current studies in different branches of physics. Examples of such processes are: atomization of pressurized fluids flowing through nozzles, lithium jets heated by neutrons in inertial confinement fusion reactors and nuclear collisions at intermediate energies. The analysis of them has strongly benefited from microscopic simulations. For the case of nuclear collisions at intermediate energies many dynamical microscopical models have been devised in order to study this process. They can be roughly classified in two main groups. Lets first mention those based on different realizations of the numerical simulation of one-body quantum kinetic equation (BUU [1], VUU [2], Landau-Vlassov[3]) in which the time evolution of the one body phase space density is followed in time. A second group are n - body molecular dynamics calculations in which the time evolution of each individual nucleon is followed (QCNM[6], QMD[4], FMD[5]). It is immediate to see that for this second group information of correlations at all levels in phase space is accessible at all times.

One of the main unsolved problems is the dynamics of the formation of fragments. Most of the calculations devoted to this topic, both for many body and one body formalisms, have focused on the late stages of the evolution. In microscopic models individual nucleons experience pair wise interactions consisting in a short range attraction and a finite range repulsion so that the final configurations are formed by free nucleons plus bounded sets of nucleons. After sufficient time has elapsed this mixture is sufficiently dilute, the individual components are well separated in space and the constituents of each aggregate are close both in q and in p space. Then the cluster structure can be simply obtained by using a minimum spanning tree recognition algorithm in q -space with an arbitrary clusterization radius. For one body approaches the fragments are built through a mapping on n -body configurations in q -space and a subsequent application of percolation like recognition algorithm. This procedure, however, is not justified, because in a calculation of a one body phase space distribution

any fluctuation is artificial being a consequence of the numerical noise only. In these calculations, however, fluctuations are the reason why one obtains different size fragments.

Here we employ the microscopic model known as Quantum molecular dynamics(QMD) for the modelation of the nucleus nucleus collision and fragments are recognized via a search of the most bound density fluctuations in phase space (ECRA) which allows us to get information about the cluster structure of the systems at all times. The fragments found with ECRA are compared with those obtained by the minimum spanning tree procedure at the end of the calculation. We find that in the dilute limit, i.e. at the end of the calculation, both cluster search routines give almost identical results. ECRA, however, detects the fragments much earlier, already when the cluster nucleons are still embedded in a surrounding nucleon gas. It allows for the first time to investigate when fragments have been formed and which mechanism drives their formation.

In section 2 we briefly review the main characteristics of the employed formalisms and the observables to be computed, in section 3 the results of the calculations are shown. Finally we present a discussion of the results

2 QMD and ECRA formalisms

The QMD model is a n body theory which simulates heavy ion reactions between 30 MeV/N and 1 GeV/N on a event by event basis. Each nucleon is represented by a coherent state of the form ($\hbar, c = 1$)

$$\phi_{\alpha}(x_1, t) = \left(\frac{2L}{\pi}\right)^{3/4} e^{-(x_1 - x_{\alpha}(t))^2 L} e^{i x_1 p_{\alpha}(t)} e^{-i p_{\alpha}^2(t) t / 2m} \quad (1)$$

Thus the wave function has two parameters x_{α}, p_{α} , L is fixed. The total n body wave function we assume to be the direct product of coherent states.

$$\phi = \phi_{\alpha}(x_1, x_{\alpha}, p_{\alpha}, t) \phi_{\beta}(x_2, x_{\beta}, p_{\beta}, t) \cdots \quad (2)$$

thus antisymmetrization is neglected. The initial values of the parameters are chosen in that way that the ensemble of $A_T + A_P$ nucleons gives a proper density distribution as well as a proper momentum distribution of the projectile and target nucleus. The time evolution of the system is

calculated by means of a generalized variational principle : we start out from the action

$$S = \int_{t_1}^{t_2} \mathcal{L}[\phi, \phi^*] dt \quad (3)$$

with the Lagrange functional $\mathcal{L}$

$$\mathcal{L} = \left(\phi \left| i\hbar \frac{d}{dt} - H \right| \phi \right) \quad (4)$$

where the total time derivative includes the derivation with respect to the parameters. The evolution is obtained by the requirement that the action is stationary under the allowed variation of the wave function.

If the true solution of the Schrödinger equation is contained in the restricted set of wave functions $\phi_\alpha(x_1, x_\alpha, p_\alpha)$ this variation of the action gives the exact solution of the Schrödinger equation. If the parameter space is too restricted, we obtain that wave function in the restricted parameter space which comes closest to the solution of the Schrödinger equation.

For the coherent states and an Hamiltonian of the form $H = \sum_i T_i + \frac{1}{2} \sum_{ij} V_{ij}$ (T_i = kinetic energy, V_{ij} = potential energy) the Lagrangian and the variation can be easily calculated and we obtain:

$$\mathcal{L} = \sum_{\alpha} -\dot{x}_{\alpha} p_{\alpha} + \sum_{\beta} \langle V_{\alpha\beta} \rangle + \frac{3}{8Lm} \quad (5)$$

$$\dot{\bar{x}}_{\alpha} = \frac{p_{\alpha}}{m} + \nabla_{p_{\alpha}} \sum_{\beta} \langle V_{\alpha\beta} \rangle \quad (6)$$

$$\dot{p}_{\alpha} = -\nabla_{x_{\alpha}} \sum_{\beta} \langle V_{\alpha\beta} \rangle \quad (7)$$

with $\bar{x}_{\alpha} = x_{\alpha} + \frac{p_{\alpha}}{m}t$. These are the time evolution equations which are solved numerically. The interaction V_{ij} is the real part of the Brückner g - matrix. If the energy is sufficiently high, the g - matrix becomes complex and the imaginary part acts like a cross section. For detail we have to refer to [4].

We apply a minimum spanning tree procedure to identify fragments. Two nucleons are considered as part of a fragment if another nucleons is closer than 3fm (standard) or 6 fm. Of course with this procedure the

clusters can only be identified at a very late time, when the average nuclear density is quite low.

The calculation of the fragments spectra in the ECRA formalism is based on the following definition of clusters:

given a set of particles whose dynamics is generated by a Hamiltonian of the form

$$H = \sum_i \frac{p_i^2}{2m} + V$$

According to [6] a cluster is defined as that set of particles that satisfies the following condition

$$\forall i \in C, \quad e_i = T_{iCM} + V_i < 0 \quad (8)$$

where T_{iCM} is the kinetic energy of particle i calculated in the C.M. of the cluster. And V_i is the interaction energy between particle i and all other particles belonging to the same cluster C ; i.e. all particles must be bound. For a system of particles we define the cluster structure is a partition such that for each subset of the partition of the system, condition (8) is satisfied. Because this condition can be satisfied by more than one partition we introduce the following extra condition: cluster structure is that partition that maximizes the total binding energy of the system when each subset is considered as non interacting with the others. As can be easily seen this is a very complicated problem because the binding energy is calculated in center of momentum of the cluster, as a consequence the problem is highly self consistent. To solve this problem in ref[6] an algorithm in the spirit of simulated annealing was developed. It is clear that this cluster definition is free from arbitrary "clusterization parameters", on the other side the method used to determine the most bound density fluctuations is statistical in nature and although we can be confident that we will be quite close to the minimum, we cannot be sure. It should be emphasized at this point that it is possible to apply this method at any point in the evolution of the system. At the beginning, i.e. shortly after the interaction between projectile and target has set in, this algorithm recognizes the "most bound fragmentation structure" in phase space. During this stage particles are close together in q space but the p distribution shows very strong fluctuations, and no

cluster structure can be recognized by standard configuration (percolation like) cluster recognition algorithms. At late times, when the fragments and nucleons are well separated ECRA should yield the same fragmentation pattern like a minimum spanning tree procedure.

3 Calculations

We performed 30 collision Ca + Ca at 80 MeV/A and zero impact parameter in the QMD approach. The time evolution of the system were analyzed using the ECRA algorithm as well as a simple minimum spanning tree formalism with clusterization radius of 3 and 6 fm. In figure 1a) we show the time evolution of the average total multiplicity for $A > 2$, in 1b) the population of different fragment types as a function of time and in 1c) the average maximum fragment. In all cases full lines denote the result of the ECRA analysis. In the ECRA algorithm after 50 fm/c the final distribution is obtained, long before in the minimum spanning tree approach (dashed lines).

Moreover it is important to note that when configurations are analyzed in q-space, like MST, there will always be a stage, during the initial violent part of the collision, in which all particles are part of a unique fragment. If the analysis is performed in phase space, the clusters which are identified during this stage reflect the process of destruction of initial correlations in phase space and creation of new ones. This change of the correlations is caused by collisions.

We can define two time scales,

τ_e , given by that time at which the cluster spectra given by the ECRA stops varying and

τ_c , given by the time at which the same behavior is found when the minimum spanning tree recognition algorithm is used.

We see that τ_e is about 40 - 50 fm/c. At that time the system is still quite dense and the fragment nucleons are surrounded by others finally not belonging to the cluster. This point can be seen in figure 2) where snapshots at the relevant times are displayed. The difference between τ_e and τ_c shows that fragments are much earlier formed than they can be recognized as clusters in coordinate space.

Once this general behavior has been established we focus on the time development of the microscopic structure of the clusters. Let $n_1 \dots n_N$ the nucleons which belong to the cluster C at the time t . $b_C(t) = 0.5 * n_N(n_N - 1)$ is then the number of pairs of nucleons in cluster C at that time. At the time Δt later, some of the nucleons may have left the clusters and are part of another cluster or singles. Let N_{CA} be the number of nucleons which have been in the cluster C at time t and are at $t + \Delta t$ in the cluster A . We define $a(t + \Delta t) = \sum_A 0.5 * (N_{CA}(N_{CA} - 1))$

Now we are able to introduce the persistence coefficient (P)

$$P(t + \frac{\Delta t}{2}) = \langle \sum [a(t + \Delta t)/b(t)] \rangle$$

where $\langle \dots \rangle$ denotes the average over the ensemble of nucleus nucleus collisions. $P(t)$ will be equal to 1 if all particles remain together and 0 if the clusters break up completely. It measures the tendency of the members of a given cluster to remain together. The average value of P when we remove one particle from clusters of mass in the range $3 \leq A \leq 14$ is $\langle P_{-1} \rangle = (1/12) \sum_{n=3,14} \left[\frac{0.5(n-1)(n-2)}{0.5n(n-1)} \right] \sim 0.7$

In Fig.3) we show the time evolution of the persistence coefficient (full line) for fragments of mass $3 \leq A \leq 14$. From it we can conclude that the constituent nucleons are strongly correlated in time and that after 45 fm/c the microscopic structure of the fragments is essentially fixed.[7]

This behavior has already been found when analyzing the fragmentation of spherical systems excited via a monopole vibration, but in this case the fragmenting system is excited via the simulation of the complete collisional process and as a consequence the excitation and the initial fluctuations in phase space are fixed totally by the dynamics of the microscopic model. In this way we introduce a third time scale (τ_p), the time at which the persistence coefficient attains a value above 0.8.

Between the different time scales the following relation holds

$$\tau_e \leq \tau_p \ll \tau_c$$

In order to investigate the role played by nucleon nucleon collisions, we also display in Fig.3) the collision rate (properly rescaled to ease the comparison between the two curves) averaged over medium mass fragments. Comparison between the persistence coefficient and the collision rate

is particularly enlightening. The collision rate attains its maximum value for a time of 30 fm/c (τ_{cm}) and reaches its asymptotic plateau at approximately 50 fm/c (when P surpasses 0.7 value). From this the following view of the fragment formation is apparent : We have a first stage in which the system is highly collisional, i.e. nucleon nucleon collisions scatter the nucleons into different phase space regions and fluctuations begin to develop in phase space ($t \leq \tau_{cm}$). At the second stage the collision rate begins to diminish and the persistence coefficient increases while the curves in Fig 1) begin to reach their asymptotic value ($\tau_{cm} \leq t \leq \tau_p$) it is obvious that the effect of collisions has been that of "aligning" particles in momentum space building the so called MBDF. After this point collisions are not dominant any more, the persistence coefficient is almost constant, as well as the bin population, total multiplicity, etc. In this stage the evolution of the system is dominated by the expansion and after sufficient time the system becomes dilute enough to be viewed as a mixture of free nucleon plus small and medium mass aggregates, recognizable by a simple coalescence model. This free nucleons and aggregates turn out to be the spatially separated MBDF.

4 Discussion

In view of the results presented in the previous section it is clear that the process of fragment formation has the following characteristics : i) final fragments result from the time evolution of the initial most bound density fluctuations, ii) the MBDF are formed during the short time during which the evolution of the system is dominated by collisions. iii) It takes place in phase space and iv) is a many particle correlation effect. v) It takes place in a time scale much smaller than the one associated with spatial separation of fragments $\tau_p \ll \tau_c$. vi) The microscopic structure of the MBDF remains pretty constant after a time $\tau_e \ll \tau_c$, i.e. when collisions stop being dominant. v) The following relation holds between the three characteristic time scales $\tau_e \leq \tau_p \ll \tau_c$. vii) non equilibrium effects seem to play an essential role. Certainly these results rule out any coalescence type fragmentation model or those in which only the one body phase space density is considered. Fragments are clearly based on fluctuations and

a complete description of the n- body phase space is required for their modeling.

One of us (C.O.D.) gratefully acknowledge the Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET) and Universite de Nantes for partial financial support.

References

- [1] J. Aichelin and G. Bertsch, Phys. Rev. C31 (1985) 1730 W. Cassing et al., Phys. Rep. 188 (1990) 361,
- [2] H. Kruse, B. Jacak and H. Stöcker, Phys. Rev. Lett. 54 (1985) 289 H. Stöcker and W. Greiner, Phys. Rep. 137 (1986) 277. A simplified PC version of this program is available in K. Langanke, J.A. Maruhn and S.E. Koonin, Computational Nuclear Physics II, Springer-Verlag, Heidelberg
- [3] F. Sebillie et al., Nucl. Phys. A501 (1989) 137
- [4] J. Aichelin, Phys. Rep. 202 (1991) 233
- [5] H. Feldmeier, Nucl. Phys. A515 (1990) 147
- [6] C.O.Dorso and J.Randrup, Phys.Lett.B 301 (1993) 328
- [7] C.O.Dorso and P.E.Balonga, Phys.Rev.C (in press)

Figure Captions

Figure 1

a) The maximum fragment as a function of time, according to ECRA formalism (full lines), percolation like algorithm with $R_c = 6$ fm (dot-dashed) and $R_c = 3$ fm (dashed); b) the population of mass bins : $\alpha) 3 \leq A \leq 5$ and $\beta) 7 \leq A \leq 9$, in both cases ECRA results are denoted by full lines and percolation like algorithm with $R_c = 3$ fm by dashed lines; c) The average total multiplicity as a function of time(same convention as in a))

Figure 2

x-y projections of snapshots of one collision simulation at times a) 0 fm/c, b) 40 fm/c, c) 60 fm/c, d) 80 fm/c. The particles are represented by circles of 3 fm in the corresponding scale

Figure 3

The persistence coefficient (full line) and the rescaled average collisional rate(dashed line)as a function of time, dashed horizontal lines denote the value 1 (upper limit for the P coefficient) and value 0.7 (lower limit of strong correlation. vertical dashed line marks the time 45 fm/c where P has already surpasses the value 0.7.

Fig 1

Fig 2

Fig 3