114

**Finding Curves on General Spaces through Quantitative Topology,
with Applications for Sobolev and Poincaré Inequalities**

Stephen SEMMES

5w3545

Institut des Hautes Etudes Scientifiques
35, route de Chartres
91440 – Bures-sur-Yvette (France)

Août 1995

IHES/M/95/77

Finding Curves on General Spaces through Quantitative Topology, with Applications for Sobolev and Poincaré Inequalities

Stephen Semmes

Abstract. In many metric spaces one can connect an arbitrary pair of points with a curve of finite length, but in Euclidean spaces one can connect a pair of points with a lot of rectifiable curves, curves that are well distributed across a region. In the present paper we give geometric criteria on a metric space under which we can find similar families of curves. We shall find these curves by solving first a “dual” problem of building Lipschitz maps from our metric space into a sphere with good properties. These families of curves can be used to control the values of a function in terms of its gradient (suitably interpreted on a general metric space), and to derive Sobolev and Poincaré inequalities.

Contents

1. Introduction

Part I. Constructions of Lipschitz Mappings

2. Basic facts about metric spaces
3. Compactness and convergence of metric spaces
4. Topological dimension and polyhedra
5. Constructing continuous mappings into spaces with controlled contractability function
6. Topological dimension with bounds
7. Mappings into polyhedra with Lipschitz bounds
8. Mappings into spheres with nonzero degree and Lipschitz bounds
9. Bounds for the degrees of Lipschitz mappings
10. Lipschitz bounds for homotopies of Lipschitz mappings
11. Two summarizing theorems

Part II. Applications of Lipschitz Mappings

12. Two useful reformulations
13. Some topological lemmas
14. The coarea estimate
15. Finding curves inside compact sets with finite length
16. The proofs of Theorems 12.1 and 12.5
17. Some observations about differential forms
18. Nontrivial Lipschitz mappings on “codimension-1” sets in Euclidean spaces

Appendices

- A. Review of degree theory and homology
- B. Cleaner statements about Sobolev and Poincaré inequalities
- C. Regular metric spaces

References

1. Introduction.

Given a pair of points $x, y \in \mathbf{R}^n$, we can join them with line segment whose length is $|x - y|$. This line segment is the unique curve between x and y of minimal length. There are many other curves with approximately minimal length. Given any $z \in \mathbf{R}^n$ with $|z - (x + y)/2| \leq |x - y|/10$, we can combine the line segments which join x to z and z to y to get another curve from x to y which is not too long. The family of all these curves fills up a region in $\mathbf{R}^n$ which looks like a cone near x and y and which is fat in the middle. The mass of these curves is distributed well across this region. The masses concentrate at x and y , but that is unavoidable.

Under what conditions on a metric space can we connect a given pair of points by a similar family of rectifiable curves?

We shall give here some sufficient conditions, in Theorems 1.9 and 1.11.

The existence of nice families of curves in a metric space is closely related to inequalities for functions in terms of their gradients, as in Sobolev and Poincaré inequalities. (Theorems 1.22 and 1.27, Appendix B.)

The existence of these families of curves is also related to the problem of building families of homotopically nontrivial Lipschitz mappings into spheres which are concentrated in prescribed balls, with bounds on their Lipschitz norms. (Theorems 1.29 and 1.30, Sections 11 and 12.)

Lipschitz mappings of this type give rise to differential forms with nice properties, and which can also be used to control a function in terms of its gradient. (Section 17.)

These themes have familiar counterparts in topology. Given a space X , mappings from X into the k -sphere $\mathbf{S}^k$ are related to k -dimensional homology and cohomology on X . If X is something like an m -dimensional manifold, then mappings from X into $\mathbf{S}^k$ also give rise to $(m - k)$ -dimensional homology classes in X , coming from the fibers of the mapping.

The conditions on metric spaces that we shall consider are close to conditions used in topological finiteness theorems, as in [Fe1], [Fe2], [GP], [GPW], [P1], [P2].

For the record, a metric space is a nonempty set M equipped with a function $d(x, y)$ defined on $M \times M$ such that $d(x, y)$ is symmetric and nonnegative, vanishes exactly on the diagonal, and satisfies the triangle inequality. We write $d_M(x, y)$ when we want to be explicit about the metric space whose metric is under consideration. All metric spaces in this paper will be assumed to be separable. Subsets of Euclidean spaces will be assumed to be equipped with the induced Euclidean metric unless something to the contrary is explicitly stated. Given a set $A \subseteq M$ we write $\text{diam } A$ for its diameter, which we take to be 0 when A is empty and ∞ when A is unbounded. We write $B(x, r)$ for the open ball in M with center x and radius r , we write $\overline{B}(x, r)$ for the closed ball, and we may write $B_M(x, r)$ when we want to make the metric space M explicit. If B is a ball in M and $k > 0$, then we write kB for the ball with the same center and k times the radius. The diameter of a set $E \subseteq M$ is denoted by $\text{diam } E$, or by $\text{diam}_M E$ when we want to make the dependence on M explicit.

A map $f : (M, d(\cdot, \cdot)) \rightarrow (M', d'(\cdot, \cdot))$ between metric spaces is *Lipschitz* if there is a

The author is supported by the U.S. National Science Foundation.

constant C such that

$$(1.1) \quad d'(f(x), f(y)) \leq C d(x, y)$$

for all $x, y \in M$. We say that f is C -Lipschitz if we want to make the constant explicit.

The following technical definition makes precise the idea of a family of curves in a metric space, and the manner in which a measure on the metric space can control the distribution of the masses of the curves.

Definition 1.2. Let $(M, d(x, y))$ be a metric space, and let (S, λ) be a measure space. (In this paper S will typically be a standard sphere with its usual Lebesgue measure.) A **family of curves in M with parameter space S** consists of a function $L : S \rightarrow [0, \infty)$ on S and a collection of 1-Lipschitz mappings $\gamma_p : [0, L(p)] \rightarrow M$, $p \in S$. We make no assumptions about the dependence of $L(p)$ or γ_p on p , and this part of the definition does not involve the measure λ on S . If ν is a Borel measure on M , we say that this family of curves in M is **controlled in measure by ν** if for each Borel set $A \subseteq M$ there is a measurable function $a : S \rightarrow [0, \infty]$ such that

$$(1.3) \quad |\gamma_p^{-1}(A)| \leq a(p) \quad \text{for all } p \in S,$$

where $|\cdot|$ denotes one-dimensional Lebesgue measure, and

$$(1.4) \quad \int_S a(p) d\lambda(p) \leq \nu(A).$$

In the case of $\mathbf{R}^n$ we can connect any given pair of points x and y by a family of curves which is controlled by a measure ν that is absolutely continuous with respect to Lebesgue measure. The density $f(u)$ of ν will blow up at the points x and y , and the singularity will be like $|u - x|^{-n+1}$ near x , and similarly near y . The family of paths described at the beginning of this introduction has this property.

We would like to know when we can find similar families of curves in other metric spaces, with good bounds on ν . This does not happen in general. For instance, if one takes a Riemannian manifold with a large balloon that has a small neck, and if one tries to connect a point on the balloon to a point away from the balloon, then the measure ν will have to concentrate on the neck of the balloon. There are interesting fractals in which every pair of points can be connected by a curve which is not too long, but such that there are not so many curves with this property. A particularly interesting example is given by the Sierpinski carpets, in which there will be a lot of such curves, but not enough to get a family of curves controlled by a measure ν which is absolutely continuous with respect to the natural background measure. (See [BB] for a discussion of some interesting properties of Sierpinski carpets.)

We are going to give some criteria under which one can find nice families of rectifiable curves in a metric space, and with good bounds. To do this we need some definitions.

Definition 1.5. We call $(M, U, d(x, y))$ an **admissible triple** if $(M, d(x, y))$ is a metric space and U is an open subset of M with the property that $U_1 = \{x \in M : \text{dist}(x, U) < \text{diam } U\}$

has compact closure in M . (We shall always use U_1 to denote the neighborhood of U just defined.)

This definition of U_1 was chosen so that it would be scale-invariant, and we shall be careful to make our other definitions scale-invariant as well. It is sometimes simpler to just assume that $\text{diam } U = 1$.

Definition 1.6. Let $(M, U, d(x, y))$ be an admissible triple, and let $N : (0, 1] \rightarrow (0, \infty)$ be a function such that $N(t) = O(t^{-k})$ as $t \rightarrow 0$ for some finite k . We say that $N(t)$ is a *covering function* for $(M, U, d(x, y))$ if for each $t \in (0, 1]$ we can cover $\overline{U}_1$ (as in Definition 1.5) with at most $N(t)$ balls of radius $t \text{diam } U$.

The requirement in Definition 1.5 that $\overline{U}_1$ be compact implies that there is automatically a covering function, except for the condition that $N(t) = O(t^{-k})$ as $t \rightarrow 0$. By fixing a covering function we get some basic control on the complexity of our metric space. This will be important for us because we shall be using some compactness arguments (in the spirit of [Gr1], [Gr2], and [P2]).

The assumption that $N(t) = O(t^{-k})$ will be needed only to permit us to use Theorem 6.5 below. See Remark 6.11.

Definition 1.7. Let $(M, U, d(x, y))$ be an admissible triple, and let $\rho : (0, 1] \rightarrow (0, \infty)$ be a nondecreasing function such that $\rho(t) \geq t$ for all t and $\lim_{t \rightarrow 0} \rho(t) = 0$. We say that ρ is a *contractability function* for $(M, U, d(x, y))$ if each ball $B(x, r \text{diam } U)$, $x \in \overline{U}_1$, $0 < r < 1$, can be contracted to a point in $B(x, \rho(r) \text{diam } U)$. (That is, there should exist a one-parameter family of continuous mappings from $B(x, r \text{diam } U)$ into $B(x, \rho(r) \text{diam } U)$ which begins with the identity and ends with a constant mapping.)

If M is a topological manifold, for instance, or a finite polyhedron, then there will automatically be a contractability function for $(M, U, d(x, y))$. We want to give it a name because our estimates will depend on it.

See [P2] for information about how bounds on contractability functions can be used.

A bound on the contractability function is a strong condition, and it would be nice to be able to weaken it in a significant manner in the results below. See Remark 6.10 for a discussion of how it is used in this paper, and how it might be weakened or replaced with other conditions. One important point is that we would have less need for a bound on the contractability function if we made stronger assumptions about the size of the space (as measured by a covering function).

The quantitative content of the assumptions that we shall be making in this paper will be in terms of covering and contractability functions. We shall also need some non-quantitative topological assumptions. For ease of reference we state separately the package of assumptions that we shall use.

Standard Assumptions 1.8. $(M, U, d(x, y))$ is an admissible triple with covering function $N(\cdot)$ and contractability function $\rho(\cdot)$. M is an oriented topological manifold of dimension n , and U is connected.

The assumption of orientability here is not serious, but it will be convenient. The requirement that M be a topological manifold is not so serious either. We shall really

only need M to have topological dimension n and to have a reasonable version of degree theory for maps from M into the standard n -sphere $\mathbf{S}^n$. We shall be more precise about this point in Remark A.35.

When thinking about the manifold assumption in 1.8 in the context of the theorems below, one should keep in mind that one gets to choose M and U . One can try to choose them to avoid singularities, e.g., if one is working on polyhedron.

The $n = 1$ case of the definitions and results in this paper is somewhat degenerate and not terribly interesting. The reader is probably better off forgetting about it.

If M is as in Standard Assumptions 1.8, then the topological dimension n of M is controlled by the covering function $N(\cdot)$ (i.e., $n \leq k$, where k is as in Definition 1.5). See Theorem 12 in [P2], and also Corollary 4.7 below.

Conditions like the ones in Standard Assumptions 1.8 have been much studied, especially in connection with topological finiteness theorems. See [Fe1], [Fe2], [GP], [GPW], [P1], [P2].

Let H^n denote n -dimensional Hausdorff measure (whose definition is recalled in (2.14)). (Do not confuse H^n with cohomology.) We shall use the notation $H^n|_E$ for the restriction of H^n to the set E .

Theorem 1.9. *Suppose that $(M, U, d(x, y))$ is as in Standard Assumptions 1.8. Suppose also that $H^n(U_1) < \infty$. Let B_0 and B_1 be two balls contained in U with radius $\delta \operatorname{diam} U$, where δ is some positive number.*

Then there is a family of curves in M with parameter space $\mathbf{S}^n$ such that all the curves are contained in U_1 , they all start in $\overline{B_0}$ and end in $\overline{B_1}$, and they are controlled in measure by $\nu = C_0 (\operatorname{diam} U_1)^{-n+1} H^n|_{U_1}$. Here C_0 depends only on n , δ , and the functions $N(\cdot)$ and $\rho(\cdot)$ from Standard Assumptions 1.8. Note that we are implicitly using Lebesgue measure on our parameter space $\mathbf{S}^n$.

The case where $M = \mathbf{R}^n$ provides a nice exercise for understanding the definitions.

Standard Assumptions 1.8 do not imply anything about the behavior of H^n on M . Think about snowflakes, like $M = \mathbf{R}^n$ equipped with the metric $|x - y|^s$, $0 < s < 1$.

The reader might want to spend a moment contemplating the geometric meaning of the conclusions of Theorem 1.9. It is one thing to say that a pair of balls can be connected by a curve which is not too long; the existence of a family of curves as in the theorem is something more. It allows for an interesting interaction with the measure theory on M , for instance.

If we want to connect a given pair of points with a nice family of curves, then we need to make stronger assumptions.

Standard Assumptions 1.10. Let $(M, U, d(\cdot, \cdot))$, $N(\cdot)$, and $\rho(\cdot)$ be as in Standard Assumptions 1.8. Let $x_0, x_1 \in U$ be given, with $B(x_0, \delta \operatorname{diam} U), B(x_1, \delta \operatorname{diam} U) \subseteq U$ for some $\delta \in (0, 1)$. We require also that $(M, B(x_i, s \delta \operatorname{diam} U), (s \delta)^{-1} d(\cdot, \cdot))$ have covering function $N(\cdot)$ and contractability function $\rho(\cdot)$ for each $s \in (0, 1]$ and $i = 0, 1$ (in the sense of Definitions 1.6 and 1.7, with U replaced by $B(x_i, s \delta \operatorname{diam} U)$).

In other words we are asking for uniform and scale invariant bounds on the covering and contractability properties of M near x and y . (These conditions are slightly messy, and the reader might prefer the cruder assumptions in Lemma 1.16 below.)

Theorem 1.11. *Suppose that $(M, U, d(x, y))$, x_0, x_1 , and δ are as in Standard Assumptions 1.10. Define a measure ν on M by $\nu = h \cdot H^n|_{U_1}$, where the density h is defined by*

$$(1.12) \quad h(y) = (d(y, x_0)^{-n+1} + d(y, x_1)^{-n+1}).$$

(That is, ν is a measure on the H^n -measurable subsets of M , say.) Assume also that $\nu(U_1) < \infty$.

Then there is a family of curves in M with parameter space S^n such that all the curves are contained in U_1 , they all start at x_0 and end at x_1 , and they are controlled in measure by $C_1 \nu$, where C_1 depends only on n, δ , and the functions $N(\cdot)$ and $\rho(\cdot)$ from Standard Assumptions 1.10. Note that we are implicitly using Lebesgue measure on our parameter space S^n .

In this case we also have that

$$(1.13) \quad x_0 \text{ and } x_1 \text{ can be connected by a curve in } U_1 \text{ of length } \leq C_1 |S^n|^{-1} \int_{U_1} \nu,$$

where $|S^n|$ denotes the measure of the standard n -sphere.

Again, if $M = \mathbf{R}^n$, then the conclusions of this theorem can be verified directly.

In order to digest Standard Assumptions 1.10 it is helpful to consider a cruder version of it. First we record a couple of definitions.

Definition 1.14. A metric space $(M, d(x, y))$ is said to be *doubling* if there is a constant C_2 such that every ball in M can be covered by at most C_2 balls of half the radius.

Notice that the doubling condition can be iterated easily. If $(M, d(x, y))$ is as in Definition 1.14 and B is a ball in M of radius r , then B can be covered by C_2^l balls of radius $2^{-l}r$. Thus the doubling condition gives rise to covering conditions as in Definition 1.6.

Notice also that if $(M, d(x, y))$ is doubling and complete, then every closed and bounded subset of M is compact. This follows from standard facts about metric spaces.

Definition 1.15. A metric space $(M, d(x, y))$ is said to be *locally linearly contractable* if there is a constant C_3 such that if $x \in M$ and $0 < r \leq C_3^{-1} \text{diam } M$, then $B(x, r)$ can be contracted to a point inside $B(x, C_3 r)$.

Lemma 1.16. *Suppose that $(M, d(x, y))$ is a complete metric space which is doubling and locally linearly contractable. Suppose also that M is a connected orientable topological manifold of dimension n . Then there exist a constant $\delta > 0$ and functions $N(\cdot)$ and $\rho(\cdot)$ – depending only on the constants C_2 and C_3 from the definitions of the doubling condition and local linear contractability – with the property that if $x_0, x_1 \in M$ are given, then there is an open set U in M such that Standard Assumptions 1.10 hold for these choices of $x_0, x_1, U, \delta, N(\cdot)$, and $\rho(\cdot)$, and such that $\text{diam } U \leq C d(x, y)$, where C depends only on the local linear contractability constant for M .*

To prove this we first make an auxiliary observation.

Sublemma 1.17. *If $(M, d(x, y))$ is connected and locally linearly contractable, then for each subset E of M we can find a connected open subset U of M such that $E \subseteq U$ and*

$\text{diam } U \leq C' \text{diam } E$, where C' depends only on the local linear contractability constant for M .

To prove this we may as well assume that $\text{diam } E$ is small compared to $\text{diam } M$, because otherwise we can take $U = M$. When $\text{diam } E$ is small we take U to be the set of points in M which can be connected to a point in E by a curve of diameter $< k \text{diam } E$, where k is a large constant. The local linear contractability condition implies that U is open and contains E when k is large enough, and we have that $\text{diam } U \leq (2k + 1) \text{diam } E$ by definition. This proves Sublemma 1.17.

Suppose now that $(M, d(x, y))$ is as in Lemma 1.16, and let $x_0, x_1 \in M$ be given. Choose U as in Lemma 1.17, with $E = B(x_0, d(x_0, x_1)) \cup B(x_1, d(x_0, x_1))$. Then all the requirements of Standard Assumptions 1.10 are met, with suitable choices of δ , $N(\cdot)$, and $\rho(\cdot)$. Specifically, we can take $N(t)$ to be a constant times a power of t , and $\rho(t)$ will be linear. In order to get that U_1 has compact closure in M , as in Definition 1.5, we use also the fact that all closed and bounded subsets of M are compact (since M is doubling and complete). This proves Lemma 1.16.

The families of curves obtained in Theorems 1.9 and 1.11 and Lemma 1.16 can be used to control the values of a function on the metric space in terms of its “gradient”. To formulate this precisely we need a generalization of the gradient which makes sense for metric spaces. In the next definition we follow Hajlasz, Heinonen, and Koskela.

Definition 1.18. Let $(M, d(x, y))$ be a metric space, and let f and g be two Borel measurable functions defined on an open set Ω in M , with f real-valued and g taking values in $[0, \infty]$. We say that g is a *generalized gradient* of f if

$$(1.19) \quad |f(\gamma(a)) - f(\gamma(b))| \leq \int_a^b g(\gamma(t)) dt$$

whenever $a, b \in \mathbf{R}$ and $\gamma : [a, b] \rightarrow \Omega$ is 1-Lipschitz (so that $d(\gamma(s), \gamma(t)) \leq |s - t|$ for all $s, t \in [a, b]$).

To illustrate this definition consider the following special case.

Lemma 1.20. Let $(M, d(x, y))$ be a metric space, let Ω be an open subset of M , and suppose that f is a real-valued Lipschitz function on Ω . Define g on Ω by

$$(1.21) \quad g(x) = \liminf_{r \rightarrow 0} r^{-1} \sup_{y \in B(x, r)} |f(y) - f(x)|.$$

Then g is a generalized gradient of f on Ω .

Indeed, if $\gamma(t)$ is as in Definition 1.18, then $h(t) = f(\gamma(t))$ is a Lipschitz function on $[a, b]$, and $|h'(t)| \leq g(\gamma(t))$ whenever $h'(t)$ is defined (which is almost everywhere). Thus we get (1.19), since h is Lipschitz and hence absolutely continuous. This proves Lemma 1.20.

The definition of a generalized gradient permits us to reinterpret Theorem 1.11 as follows.

Theorem 1.22. *Same notations and assumptions as in Theorem 1.11. If f and g are Borel measurable functions on U_1 and g is a generalized gradient of f , then*

$$(1.23) \quad |f(x_0) - f(x_1)| \leq C_1 \int_{U_1} \left(\frac{1}{d(y, x_0)^{n-1}} + \frac{1}{d(y, x_1)^{n-1}} \right) g(y) dH^n(y),$$

where C_1 is as in Theorem 1.11.

To prove this we need only combine the definitions and theorems. Since we are making the same assumptions as in Theorem 1.11, we get the same conclusions, which is to say that we get a collection of 1-Lipschitz mappings $\gamma_p : [0, L(p)] \rightarrow M$, $p \in \mathbf{S}^n$, which connect x_0 to x_1 and satisfy other properties as in Definition 1.2 and Theorem 1.11. For each $p \in \mathbf{S}^n$ we have that

$$(1.24) \quad |f(x_0) - f(x_1)| \leq \int_0^{L(p)} g(\gamma(t)) dt,$$

because of (1.19). To get (1.23) we average this over $p \in \mathbf{S}^n$. More precisely, there is a Borel measurable function $G : \mathbf{S}^n \rightarrow [0, \infty]$ such that

$$(1.25) \quad \int_0^{L(p)} g(\gamma(t)) dt \leq G(p) \quad \text{for all } p \in \mathbf{S}^n, \text{ and}$$

$$(1.26) \quad \int_{\mathbf{S}^n} G(p) dp \leq C_1 \int_{U_1} (d(y, x_0)^{-n+1} + d(y, x_1)^{-n+1}) g(y) dH^n(y).$$

(The dp on the left side of (1.26) denotes Lebesgue measure on $\mathbf{S}^n$.) This Borel function $G(p)$ plays the same role as the function $a(p)$ in Definition 1.2, except that it is associated to the Borel function g rather than a Borel set A . This extension of Definition 1.2 poses no difficulty, it is simply a question of approximating g by simple functions. We are also using Theorem 1.11 here to know what to put on the right side of (1.26), i.e., what measure ν to use. Once we have (1.25) and (1.26) we can derive (1.23) from (1.24). This proves Theorem 1.22.

There is also a version of this for Theorem 1.9.

Theorem 1.27. *Under the same conditions as in Theorem 1.9, if f and g are Borel measurable functions on U_1 and g is a generalized gradient of f , and if f is constant on each of B_0 and B_1 , say $f = \beta_i$ on B_i , then*

$$(1.28) \quad |\beta_0 - \beta_1| \leq C_0 (\text{diam } U_1)^{-n+1} \int_{U_1} g(y) dH^n(y),$$

where C_0 is as in Theorem 1.9.

This can be derived from the definitions in the same manner as for Theorem 1.22.

Estimates like (1.22) permit one to reduce questions of Sobolev and Poincaré inequalities to problems about integral operators, basically problems in measure theory and classical analysis. If we assume that $(M, d(x, y))$ is locally linearly contractable, for instance,

and also that the mass of M is evenly distributed, then one has practically the same kind of Sobolev and Poincaré inequalities for $(M, d(x, y))$ as for $\mathbf{R}^n$. See Appendix B, especially Theorem B.10. Gromov has indicated another approach to Sobolev inequalities on such spaces, based on the methods of [Gr3]. See Remark B.20.

See Section 17 for another approach to (1.23) and (1.28), based on differential forms.

To prove Theorems 1.9 and 1.11 we shall not construct the required families of curves directly. We shall construct some Lipschitz mappings into spheres, and we shall find our curves in the fibers of these mappings. This argument uses ideas similar to those in [DS1].

We now state the basic results about Lipschitz mappings into spheres that we shall prove. These statements will involve the notion of the degree of a continuous mapping from an oriented topological manifold of dimension n into $\mathbf{S}^n$. See Appendix A at the end of this paper for a review of topological degree theory. An important point to remember is that the degree of such a mapping exists when the mapping is constant off a compact set.

Theorem 1.29. *Suppose that $(M, U, d(x, y))$ is as in Standard Assumptions 1.8. Fix a point $\sigma \in \mathbf{S}^n$.*

(a) *Let an open ball B in U with radius $\delta \operatorname{diam} U$ be given, where δ is some positive number. Then we can find a Lipschitz map $f : M \rightarrow \mathbf{S}^n$ with Lipschitz constant $\leq C(\delta)(\operatorname{diam} U)^{-1}$ which takes only the value σ on $M \setminus B$ and which has nonzero degree. (In particular f cannot be homotoped to a constant through continuous mappings of M into $\mathbf{S}^n$ which take only the value σ on the complement of a compact subset of M .) This constant $C(\delta)$ depends on the functions $N(\cdot)$ and $\rho(\cdot)$ from Standard Assumptions 1.8 as well as n and δ (but nothing else). (For this part we do not need the assumption of connectedness of U in Standard Assumptions 1.8.)*

(b) *Suppose that $g : M \rightarrow \mathbf{S}^n$ takes only the value σ on $M \setminus U$ and has Lipschitz norm $\leq k(\operatorname{diam} U)^{-1}$ for some $k \geq 1$. Then the degree D of g satisfies $|D| \leq C(k)$, where $C(k)$ depends on the functions $N(\cdot)$ and $\rho(\cdot)$ from Standard Assumptions 1.8 as well as n and k (but nothing else).*

Theorem 1.30. *Suppose that $(M, U, d(x, y))$ is as in Standard Assumptions 1.8. Fix a point $\sigma \in \mathbf{S}^n$, and set $r = \operatorname{diam} U$. Suppose that $f_0, f_1 : M \rightarrow \mathbf{S}^n$ take only the value σ on $M \setminus U$, are Lipschitz with norm $\leq L r^{-1}$ for some constant L , and have the same degree. Then there is a mapping $F : M \times [0, r] \rightarrow \mathbf{S}^n$ such that*

$$(1.31) \quad F(x, 0) = f_0(x) \text{ and } F(x, r) = f_1(x) \text{ for all } x \in M,$$

$$(1.32) \quad F(x, t) = \sigma \text{ when } x \in M \setminus U_1 \text{ and } t \in [0, r], \text{ and}$$

$$(1.33) \quad F \text{ is Lipschitz with norm } \leq C(L) r^{-1}.$$

Here $C(L)$ depends on the functions $N(\cdot)$ and $\rho(\cdot)$ from Standard Assumptions 1.8 as well as n and L (but nothing else).

The main point in Theorems 1.29 and 1.30 is to get the Lipschitz bounds and the bound on the degree in part (b) of Theorem 1.29. Without these bounds these results would not have real content beyond standard topology.

In Theorem 1.29 (a) we shall need to have a controlled covering function $N(\cdot)$ only in order to control the “topological dimension function” (Definition 6.1). We shall spell out this point in detail in Remark 8.24. This is an important point because it is clear that a bound on the covering function is not really a natural assumption to make in order to guarantee the existence of a Lipschitz mapping to the sphere with bounded norm and nonzero degree. The covering function could be very large, the manifold could be very large, and that could cause no problem for the map to the sphere, it would simply have to squeeze points together, and there is nothing wrong with that. By contrast the bounds in Theorems 1.29 (b) and 1.30 rely on the controlled covering function in a much more substantial way.

When we prove Theorem 1.30 in Section 10 we shall derive it from a more general result. If one assumes that f_0 and f_1 are homotopic through mappings which take only the value σ on $M \setminus U$ – but without a bound on the modulus of continuity of the homotopy – then the analogue of Theorem 1.30 can be proved without the requirement that M be an oriented topological manifold, or that U be connected. This also works for much more general target spaces than $\mathbf{S}^n$.

One should keep in mind that the topological assumptions in these two theorems are stronger than needed. The assumption that M be a connected oriented topological manifold is needed only for the familiar facts about continuous mappings from M into $\mathbf{S}^n$ (in terms of homotopy and degree). This will be spelled out in Appendix A. (See Remark A.35.) The contractability assumptions are also not optimal, but there the story is more complicated. See Remark 6.10.

A large part of this paper is aimed at the proofs of Theorems 1.29 and 1.30, and we shall deal with them first, in Part I. This will require a certain amount of preliminary material about metric spaces and topological dimension, and the actual proofs are given in Sections 8, 9, and 10. We shall derive Theorems 1.9 and 1.11 from Theorems 1.29 and 1.30 in Part II, with the transition between them being spelled out in Sections 11 and 12.

In Section 17 we explain another way of getting information from the Lipschitz maps that we construct, in terms of differential forms. These differential forms provide an alternate approach to the inequalities (1.23) and (1.28).

In Section 18 we give a very different criterion for constructing nontrivial Lipschitz mappings on a subset of a Euclidean space. This criterion involves the geometry of the complement of the set, rather than the intrinsic geometry. The method of Section 18 provides an alternate approach to some Sobolev inequalities given in [Se1] and Section 6 of [DS2].

In Appendices B and C we consider metric spaces which satisfy scale-invariant and uniform versions of our assumptions. These conditions are cruder but the results are simpler and cleaner.

See the Table of Contents for more information about the structure of this paper.

I have the impression that this paper shares with fiber bundles the feature of being locally trivial. Globally the structure is more amusing, some dualities converting the pieces into different conclusions. Lipschitz mappings to curves to Poincaré inequalities. In many places arguments are spelled out more for the purpose of accessibility, bridging cultural gaps, than for novelty.

In this paper we are giving much attention to properties of controlled geometry and topology on bounded pieces of a metric space. Similar issues arise in studying the asymptotic properties of spaces at infinity. That is, instead of considering some property on a piece of a metric space with a uniform bound, and asking whether it implies some other property with a uniform bound, one can look at the large-scale behavior of a metric space at infinity, and the way in which various asymptotic properties are related. For instance, instead of asking that a piece of a metric space have a uniformly bounded contractability function, as in Definition 1.7, one can consider the property that an unbounded metric space M be *uniformly contractable*, which means that for every $r > 0$ there is an $R > 0$ such that every ball in M of radius r can be contracted inside a ball of radius R . Unlike Definition 1.7, we do not ask that $R \rightarrow 0$ when $r \rightarrow 0$; the point is to control the behavior at infinity, for large r , in a uniform way. Instead of asking for a controlled map from a space into a sphere as in Theorem 1.29 (a), in the asymptotic context one asks for the existence of a proper Lipschitz map from our metric space M – a topological manifold of dimension n , say – into $\mathbf{R}^n$ which is Lipschitz and has nonzero degree. Such an M is sometimes said to be (*rationally*) *hyper-Euclidean (or hyperspherical)*. A question of current interest is to know under what conditions a metric space which is uniformly contractable is also hyper-Euclidean. References about this point include [CGM], [DFW], [FH], [Gr4] (0.3 II'), [Gr5] (especially the beginning of § 4 and § 9 $\frac{1}{4}$).

For these considerations of the large-scale geometry a typical example of a metric space of interest would be hyperbolic space. This is an important point to keep in mind, because this paper normally relies on bounds on the covering function which are incompatible with that kind of geometry, that kind of growth at infinity.

Other references that deal with controlled geometry and topology at infinity include [BW], [FeW], and [FHP]. From the perspective of an analyst, the topologists' story of the bounded category is quite intriguing.

See [Gr4] for another discussion of asymptotic properties of metric spaces.

I would like to thank T. Coughlon, M. Gromov, J. Heinonen, P. Koskela, and D. Sullivan for their comments and suggestions. I am especially grateful to Gromov for his criticisms of an earlier version of this paper. I am not convinced however that my attempts at improvement have been entirely successful. I am not at all certain that the results given here have found their proper formulation. I am troubled by the strength of the assumptions, I am suspicious that I am missing key points that may reveal themselves in time and show that certain pieces of this story are unnaturally crude and clumsy. But this seems a reasonable time to tell the story, and at the large scale the story is coherent and nice, even if the pillars on which it stands are merely logs dragged in from Le Bois Marie which might eventually be replaced by stones sculpted with greater skill.

Part I. Constructions of Lipschitz Mappings.

2. Basic facts about metric spaces.

Throughout this section we let $(M, d(x, y))$ be any metric space. For the sake of completeness we shall state and prove some simple facts about extensions of Lipschitz functions and approximations of continuous functions by Lipschitz functions.

If E is a subset of M , then $f : E \rightarrow \mathbf{R}$ is said to be Lipschitz if there is a constant $C > 0$ so that

$$(2.1) \quad |f(x) - f(y)| \leq C d(x, y) \quad \text{for all } x, y \in E.$$

We shall sometimes say that f is C -Lipschitz to make the constant explicit. Remember always that if $x_0 \in M$ then $f(x) = d(x, x_0)$ is 1-Lipschitz, because of the triangle inequality.

Lemma 2.2. *Let $\mathcal{F}$ be a family of real-valued C -Lipschitz functions on a metric space M . If $F(x) = \inf_{f \in \mathcal{F}} f(x)$ is finite at one point, then it is finite on all of M , and it is C -Lipschitz. The same conclusion holds for the supremum.*

This is a well known and straightforward consequence of the definition.

Lemma 2.3. *If E is a nonempty subset of a metric space M and $g : E \rightarrow \mathbf{R}$ is C -Lipschitz, then there is an extension of g to a C -Lipschitz function on all of M .*

This is also well-known. Given $e \in E$ we define $f_e : M \rightarrow \mathbf{R}$ by $f_e(x) = g(e) + C d(x, e)$. Then each f_e is C -Lipschitz, $g(x) \leq f_e(x)$ for all $x, e \in E$, since g is assumed to be C -Lipschitz, and $g(e) = f_e(e)$ for all $e \in E$. Set $G(x) = \inf_{f \in \mathcal{F}} f(x)$. Then $g = G$ on E , and hence G is finite everywhere and C -Lipschitz by Lemma 2.2. This proves Lemma 2.3.

Next we record the fact that uniformly continuous functions can be approximated uniformly by Lipschitz functions, but with bounds that suit our purposes.

Lemma 2.4. *Given a bounded function $f : M \rightarrow \mathbf{R}$ and $j > 0$ set*

$$(2.5) \quad f_j(x) = \inf \{ f(y) + j d(x, y) : y \in M \}.$$

Then

$$(2.6) \quad \inf_M f \leq f_j(x) \leq f(x) \quad \text{for all } x \in M.$$

each f_j is j -Lipschitz, and f_j converges to f uniformly as $j \rightarrow \infty$ when f is uniformly continuous. More precisely we have the bound

$$(2.7) \quad \sup_M |f_j - f| \leq \phi(2j^{-1} \sup_M |f|),$$

where ϕ is the modulus of continuity of f ,

$$(2.8) \quad \phi(t) = \sup \{ |f(x) - f(y)| : x, y \in M, d(x, y) \leq t \}.$$

This is very simple. The bounds (2.6) follow immediately from the definition (2.5) of f_j , and the j -Lipschitzness of f_j is implied by Lemma 2.2. To prove (2.7) we observe that

$$(2.9) \quad f_j(x) = \inf \{ f(y) + j d(x, y) : y \in M, j d(x, y) \leq 2 \sup_M |f| \}.$$

since we may ignore the y 's in the infimum in (2.5) which satisfy $f(y) + j d(x, y) > f(x)$. This implies that

$$(2.10) \quad f_j(x) \geq \inf \{ f(y) : y \in M, j d(x, y) \leq 2 \sup_M |f| \}.$$

Since we already know that $f_j \leq f$ we get (2.7) easily. Of course the statement about uniform convergence is an immediate consequence of (2.7). This proves Lemma 2.4.

Notice that (2.9) implies that

$$(2.11) \quad f_j(x) = f(x) \quad \text{when } f(y) = f(x) \text{ for all } y \in B(x, 2j^{-1} \sup_M |f|).$$

Lemma 2.12. *Suppose that $f : M \rightarrow \mathbf{S}^n$ is a continuous map into the standard n -sphere which takes only the value σ , $\sigma \in \mathbf{S}^n$, off a compact subset K of M . Let $\epsilon > 0$ be given, and set $K_\epsilon = \{x \in M : \text{dist}(x, K) < \epsilon\}$. Then we can approximate f uniformly by Lipschitz maps into $\mathbf{S}^n$ which equal σ off of K_ϵ . These approximations have the property that they are homotopic to f through mappings which equal σ off K_ϵ .*

This is an easy consequence of Lemma 2.4. One can check that f is uniformly continuous using the standard argument, since we are assuming that $f \equiv \sigma$ off a compact set. We can approximate f by f_j 's as in Lemma 2.4, or rather by applying Lemma 2.4 to each component of f separately (viewing f as a map into $\mathbf{R}^{n+1}$). These mappings f_j are each Lipschitz and they converge uniformly to f (since f is uniformly continuous), but we should replace them by the mappings $\tilde{f}_j = f_j |f_j|^{-1}$ in order to get $\mathbf{S}^n$ -valued mappings. We can do this when j is large enough because f_j will then satisfy $|f_j| \geq \frac{1}{2}$ on all of M . Also, $\tilde{f}_j(x) = f(x)$ when $\text{dist}(x, K) \geq 2j^{-1}$, because of (2.9). To get the last statement, about homotopies, observe that if $F : M \rightarrow \mathbf{S}^n$ is continuous and satisfies $\sup_M |F - f| \leq \frac{1}{2}$, then $(tf + (1-t)F)|tf + (1-t)F|^{-1}$ provides a homotopy from F to f , and it is constant in t at any point x where $F(x) = f(x)$. This proves Lemma 2.12.

We shall use this kind of argument for building homotopies several times in this paper, but we shall often have to be more careful about the estimates.

Let us recall now the definition of Hausdorff measures. Let $s > 0$ be given. Given $E \subseteq M$ and $\delta > 0$ set

$$(2.13) \quad H_\delta^s(E) = \inf \left\{ \sum (\text{diam } E_j)^s : \{E_j\}_j \text{ is any sequence of subsets of } M \right. \\ \left. \text{such that } E \subseteq \bigcup_j E_j \text{ and } \text{diam } E_j < \delta \text{ for all } j \right\}.$$

We can require that the E_j 's be closed subsets of M without affecting $H_\delta^s(E)$. The true Hausdorff measure H^s is defined by

$$(2.14) \quad H^s(E) = \lim_{\delta \rightarrow 0} H_\delta^s(E).$$

This limit exists because $H_\delta^s(E)$ increases as $\delta \rightarrow 0$. ($H^s(E) = \infty$ is allowed.) It is well known that the restriction of H^s to the σ -algebra of Borel sets in M is a countably additive measure (while H^s is countably subadditive in all subsets of M).

Note that H^n agrees with Lebesgue measure on $\mathbf{R}^n$, except perhaps for a normalization. It is easy to check directly that H^1 equals Lebesgue measure on $\mathbf{R}$, i.e., there is no subtlety about normalizations when $n = 1$. When $s = 0$ Hausdorff measure reduces to counting measure.

Lemma 2.15. *If $(M, d(x, y))$ and $(M', d'(x, y))$ are metric spaces, $f : M \rightarrow M'$ is C -Lipschitz, and $E \subseteq M$, then $H^s(f(E)) \leq C^s H^s(E)$.*

This inequality represents a mild abuse of notation, since the two versions of H^s live on different spaces.

The lemma follows directly from the definitions.

Definition 2.16. A subset E of M has finite Hausdorff dimension if $H^s(E) < \infty$ for some s , and the infimum of all such s is called the Hausdorff dimension of E .

Lemma 2.17. *Let X be a metric space, and let $N : (0, 1] \rightarrow (0, \infty)$ be a function with $N(t) = O(t^{-\mu})$ as $t \rightarrow 0$ for some $\mu \geq 0$. Suppose that X can be covered by at most $N(\epsilon)$ closed balls of radius ϵ for each $0 < \epsilon < 1$. Then X has Hausdorff dimension $\leq \mu$.*

Indeed, in this case one can check that $H^\mu(X) < \infty$ directly from the definitions.

3. Compactness and convergence of metric spaces.

We shall use compactness arguments heavily in this paper, and we shall use Gromov's notion of convergence of compact metric spaces, as in [Gr1], [Gr2]. The paper [P2] provides a nice reference for these topics. See also Section 1 of [BS].

We begin with the relevant definitions.

Definition 3.1. Let $\{M_i\}$ be an infinite sequence of compact metric spaces. (For the simplicity of notation we do not write explicitly their different metrics, but the reader should not forget them.) We say that $\{M_i\}$ converges to a compact metric space M if there exist a sequence of metric spaces $\{Z_i\}$ and sequences of isometric embeddings $\phi_i : M_i \rightarrow Z_i$, $\psi_i : M \rightarrow Z_i$ so that the Hausdorff distance between $\phi_i(M_i)$ and $\psi_i(M)$ inside Z_i converges to 0. This means that there should exist a sequence $\{\epsilon_i\}$ of positive numbers such that $\epsilon_i \rightarrow 0$ as $i \rightarrow \infty$ and

$$(3.2) \quad \begin{aligned} \sup_{x \in \phi_i(M_i)} \text{dist}_{Z_i}(x, \psi_i(M)) &< \epsilon_i \quad \text{and} \\ \sup_{y \in \psi_i(M)} \text{dist}_{Z_i}(y, \phi_i(M_i)) &< \epsilon_i \end{aligned}$$

for all i .

We could require that all the Z_i 's be the same here, but we shall not need this additional structure, and it is a little simpler to do without it. We could also formulate this notion of convergence of metric spaces in terms of the "Gromov-Hausdorff distance" between them, but we shall not bother with that either.

This notion of convergence leads to a corresponding notion of compactness, and the “Gromov Compactness Theorem” provides nice characterizations of compact and precompact families of metric spaces. (See [P2].) The following criterion for compactness will be sufficient for our purposes.

Theorem 3.3. *Let $\{M_i\}$ be an infinite sequence of compact metric spaces. Suppose that $\{M_i\}$ is “uniformly compact”, in the sense that $\sup_i \text{diam}_i M_i < \infty$ and there is a single function $N : (0, 1] \rightarrow (0, \infty)$ such that for each $\epsilon \in (0, 1]$ we can cover each M_i with at most $N(\epsilon)$ closed balls in M_i of radius ϵ . Then there is a subsequence of $\{M_i\}$ which converges to a compact metric space M in the sense of Definition 3.1.*

See [P2] for a proof. Let us review a few of the basic ideas. In order to understand how the compactness is obtained it is helpful to think first about the case where all the M_i ’s have a bounded number of elements. This case comes down to the compactness of closed and bounded intervals in $\mathbf{R}$, but still one has to be a little careful about the definitions. In general the M_i ’s are uniformly approximable by finite metric spaces with boundedly many elements, because of the uniform covering function $N(\cdot)$ in Theorem 3.3, and this permits one to work down to the finite case. See [P2] for details.

Lemma 3.4. *Let $\{M_i\}$ be an infinite sequence of compact metric spaces which converges to a compact metric space M in the sense of Definition 3.1. Then $\text{diam } M = \lim_{i \rightarrow \infty} \text{diam } M_i$. If there is a function $N : (0, 1] \rightarrow (0, \infty)$ such that for each $\epsilon \in (0, 1]$ we can cover each M_i with at most $N(\epsilon)$ closed balls in M_i of radius ϵ , then M enjoys the same property (with the same function N).*

This is an easy consequence of the definition of the convergence. (To be honest, it is slightly easier to prove the statement about covering functions if we cut ourselves a little slack for the covering function for the limit, e.g., if we settled for $N(\epsilon/2)$ instead of $N(\epsilon)$. This cruder statement would suffice for our purposes.)

Definition 3.5. A collection $\mathcal{F}$ of compact metric spaces will be called precompact if every sequence in $\mathcal{F}$ has a subsequence which converges in the sense of Definition 3.1.

Remark 3.6. We shall be using compactness arguments several times in this paper, and as a result we shall sometimes make statements of the form “Let $\mathcal{F}$ be the collection of all metric spaces with the properties ...” Strictly speaking, this will not always make sense, because the properties may be so general that $\mathcal{F}$ would not be a set. However, this is not at all a serious issue. As stated at the beginning of the introduction, we shall assume that all of our metric spaces are separable. For these compactness arguments we do not really care about the metric spaces themselves, only their equivalence classes up to isometry. With these two restrictions we can easily restrict our set-theoretic universe to avoid the potential problems of setness. For the simplicity of exposition we shall ignore this point from now on.

4. Topological dimension and polyhedra.

Definition 4.1. Let X be a compact metric space and let $\{U^i\}$ be a finite covering of X by open sets. We say that $\{U^i\}$ has *order* $\leq n$ if no point in X lies in more than $n + 1$ of the U^i 's. The *mesh* of the covering is the maximum of the diameters of the U^i 's.

Definition 4.2. Let X be a compact metric space and let $n \geq 0$ be given. We say that X has *topological dimension* $\leq n$ if for each $\epsilon > 0$ there is a finite covering of X by open sets with mesh $< \epsilon$ and order $\leq n$.

This is not the standard definition for the dimension, but it is equivalent. (See the corollary on p.67 of [HW].) One normally speaks of the topological dimension of a space, rather than the topological dimension being less than or equal to something, and one normally allows noncompact spaces as well, but for our purposes these restrictions are sufficient and simpler.

It is an amusing exercise to check that the unit cube in $\mathbf{R}^n$ has topological dimension $\leq n$. At any rate it is well known that a compact subset of a topological manifold of dimension n has dimension $\leq n$. (This is immediate if one uses the definition of topological dimension given in [HW].)

The notion of topological dimension is beautiful and fascinating. It imposes nontrivial restrictions on the structure of a space while letting other aspects of the structure run wild. In Section 6 we shall consider a more quantitative version of the statement that “ X has topological dimension $\leq n$ ”, in which the shape of the covering sets is also controlled.

Let us recall some basic facts about the topological dimension.

Lemma 4.3. *Suppose that X is a compact metric space with topological dimension $\leq n$ and that X_0 is a compact subset of X . Then X_0 also has topological dimension $\leq n$.*

This is a trivial exercise. The next results are more interesting.

Definition 4.4. If Y and Z are metric spaces, $g : Y \rightarrow Z$ is a continuous mapping, and $\epsilon > 0$, then we say that g is an ϵ -mapping if $g^{-1}(z)$ has diameter $< \epsilon$ for every z in the image of g in Z .

Theorem 4.5. *If Y is a compact metric space, then Y has topological dimension $\leq n$ if and only if for each $\epsilon > 0$ there is a continuous ϵ -mapping from Y into a finite polyhedron of dimension $\leq n$.*

This theorem is given on p.72 of [HW]. Of course the polyhedron is permitted to depend on ϵ .

Theorem 4.6. *Let X be a compact metric space. Then X has topological dimension $\leq n$ if it has Hausdorff dimension $< n + 1$.*

See p.107 of [HW].

Corollary 4.7. *Let X be a compact metric space and $N : (0, 1] \rightarrow (0, \infty)$ a function such that $N(t) = O(t^{-\mu})$ as $t \rightarrow 0$. Suppose that X can be covered by at most $N(\epsilon)$ closed balls of radius ϵ for each $0 < \epsilon < 1$. Then X has topological dimension $\leq \mu$.*

This follows by combining Theorem 4.6 with Lemma 2.17.

There is a nice way to take a finite covering of a space and associate to it a mapping of the space into a polyhedron. This construction can be used to prove the “only if” part of Theorem 4.5, and we shall use it a lot in this paper, and so we describe it in some detail.

Let us begin with a more formal discussion of polyhedra. We shall restrict ourselves to finite polyhedra, and it will be convenient to always have them living inside some finite dimensional Euclidean space (instead of considering abstract polyhedra).

Definition 4.8. A k -*simplex* in $\mathbf{R}^m$ is a closed subset of $\mathbf{R}^m$ which is the convex hull of $k + 1$ affinely independent points (i.e., they should not all lie on a $(k - 1)$ -plane). These points are called the *vertices* of the simplex. If τ is a simplex, then the *faces* of τ are the simplices generated by the various subsets of the vertices of τ . If τ is a simplex, then the *interior* of τ means τ with all its faces removed. We may sometimes call this the “geometric interior” of τ , since it is usually not the interior of τ as a subset of $\mathbf{R}^m$. A *finite polyhedron* is a subset P of some $\mathbf{R}^m$ which is the union of some finite set $\mathcal{S}(P)$ of simplices in $\mathbf{R}^m$. For us a polyhedron is not simply a subset of $\mathbf{R}^m$, but it is “marked” in the sense that we distinguish this finite collection $\mathcal{S}(P)$ of constituent simplices from all the other simplices which are contained in P set-theoretically. We also require that if a simplex τ is contained in $\mathcal{S}(P)$ then all its faces are contained in $\mathcal{S}(P)$ too, and that every pair of elements of $\mathcal{S}(P)$ have disjoint interiors. We say that the finite polyhedron P has dimension $\leq n$ if all the simplices in $\mathcal{S}(P)$ are k -simplices with $k \leq n$.

This definition differs in both terminology and details from its counterpart on p.66 of [HW], but the differences are minor and not significant for the statement of Theorem 4.5 above.

Let X be a compact metric space, and suppose that $U^1, \dots, U^m$ is a finite covering of X by nonempty open sets. There is a standard procedure for associating to this covering a finite polyhedron $P \subseteq \mathbf{R}^m$. Let $\epsilon^1, \dots, \epsilon^m$ denote the standard basis vectors in $\mathbf{R}^m$. Let $\mathcal{S}$ denote the collection of nonempty subsets S of $\{1, \dots, m\}$ such that $\bigcap_{i \in S} U^i \neq \emptyset$. Given $S \in \mathcal{S}$, let $\tau(S)$ denote the simplex which is the convex hull of the ϵ^i 's, $i \in S$. Let P denote the polyhedron in $\mathbf{R}^m$ which is the union of all the $\tau(S)$'s, with $\mathcal{S}(P) = \{\tau(S) : S \in \mathcal{S}\}$. (These notations will be used many times in this paper.)

Lemma 4.9. *Notations and assumptions as in the preceding paragraph. If the covering $U^1, \dots, U^m$ of X has order $\leq n$ (as in Definition 4.1), then the corresponding polyhedron P has dimension $\leq n$.*

This is an immediate consequence of the definitions.

Next we build a mapping from X into P . We begin with a standard definition.

Definition 4.10. Let X be a compact metric space, and let $U^1, \dots, U^m$ be a finite covering of X by nonempty open subsets of X . A family of functions $\phi^1, \dots, \phi^m$ on X is called a *partition of unity subordinate to $U^1, \dots, U^m$* if each ϕ^i is a continuous function which takes values in $[0, 1]$ and vanishes on $X \setminus U^i$, and if $\sum_i \phi^i \equiv 1$ on X .

It is a standard exercise that there is always a partition of unity which is subordinate to a given covering of X . In later sections we shall have to be a little careful about the way that we choose our partitions of unity, in order to get Lipschitz estimates.

Lemma 4.11. *Let X be a compact metric space, let $U^1, \dots, U^m$ be a finite covering of X by nonempty open subsets of X , let $\phi^1, \dots, \phi^m$ be a partition of unity subordinate to $U^1, \dots, U^m$, and let P , etc., be defined as in the paragraph which precedes Lemma 4.9. Define $f : X \rightarrow \mathbf{R}^m$ by $f = \sum_{i=1}^m \epsilon^i \phi^i$. Then f is a continuous mapping which takes values in P , and $\text{diam } f^{-1}(\tau)$ is at most twice the mesh of $U^1, \dots, U^m$ for each $\tau \in S(P)$.*

This is straightforward. Let $x \in X$ be given, and let $S(x)$ denote the set of $i \in \{1, \dots, m\}$ such that $\phi^i(x) \neq 0$. Then $x \in U^i$ for each $i \in S(x)$, and so $S(x) \in \mathcal{S}$. Clearly $f(x) \in \tau(S(x))$, whence $f(x) \in P$.

Conversely, fix an $S \in \mathcal{S}$, and consider $f^{-1}(\tau(S))$. Set $U(S) = \bigcup_{i \in S} U^i$. If $f(x) \in \tau(S)$, then $\phi^i(x) = 0$ when $i \notin S$, and hence $S(x) \subseteq S$. This means that $x \in U(S)$. On the other hand $\text{diam } U(S)$ is at most twice the mesh of $U^1, \dots, U^m$, as one can check from the definitions (using $\bigcap_{i \in S} U^i \neq \emptyset$). This proves Lemma 4.11.

Remark 4.12. In this paper we shall often be concerned with controlling the geometry of the finite polyhedra that we are using, and not just their dimension. In some cases we shall have assumptions which ensure that the finite polyhedra being used range through only a finite collection of finite polyhedra, so that their complexity is controlled. In any case the construction given just before Lemma 4.9 produces polyhedra of a somewhat special or normalized nature, in that all the distinguished simplices are convex hulls of basis vectors of the ambient Euclidean space. This means that the distinguished simplices all look like standard simplices, and it also prevents them from intersecting at small angles. In other words, while the complexity of these polyhedra may deteriorate in that the number of distinguished simplices may be enormous, this construction does not permit the individual distinguished simplices to degenerate, nor does it permit them to meet at a small angle.

5. Constructing continuous mappings into spaces with controlled contractibility function.

In this section we provide some technical results which enable us to construct continuous mappings with prescribed behavior into spaces which have controlled local contractability properties. We shall be able to approximately specify the values of these continuous mappings, but we shall not get any bounds on their moduli of continuity.

The results in this section are covered already in [P1], [P2], except for minor technical differences. For the sake of completeness we explain the proofs, modulo some imprecisions that hopefully make it easier to read. In order to simplify the arguments we shall accept conclusions that are not as strong as possible. See the Main Obstruction Result in [P2] for a sharper formulation.

In our setting we shall not assume that a space has good contractability properties, but rather that a subset of a metric space has good contractability properties relative to the ambient space. Let us formalize this with a definition.

Definition 5.1. Let M be a metric space, let Z be a subset of M , and let $\rho : (0, 1] \rightarrow (0, \infty)$ be a nondecreasing function such that $\rho(r) \geq r$ for all r and $\lim_{r \rightarrow 0} \rho(r) = 0$. We say that ρ is a *contractability function* for the pair (M, Z) if each ball $B(x, r)$ with $x \in Z$ and $0 < r < 1$ can be contracted to a point inside $B(x, \rho(r))$.

In other words, M is locally contractable near elements of Z in a way which is controlled by $\rho(\cdot)$, but we do not say anything about what happens away from Z . Unfortunately [P1], [P2] deal only with the case of spaces which are themselves locally contractable (i.e., $Z = M$), but this is not a significant issue.

Roughly speaking, local contractability of a space implies that maps into the space can be manipulated topologically in almost the same manner as if the space were locally linear. This principle is illustrated by the following observation. Suppose that M , Z , and ρ are as in Definition 5.1. Let τ be a simplex and let $\theta : \partial\tau \rightarrow Z$ be a continuous mapping. If $\text{diam } \theta(\partial\tau)$ is very small, then we can extend θ to a continuous mapping $\Theta : \tau \rightarrow M$ in such a way that $\text{diam } \Theta(\tau)$ is also small. We can choose Θ so that $\text{diam } \Theta(\tau) \leq 2\rho(2\text{diam } \theta(\partial\tau))$. The existence of this extension and the bound on the diameter follow from Definition 5.1. Keep in mind that $\rho(\epsilon) \rightarrow 0$ as $\epsilon \rightarrow 0$.

We shall use this argument repeatedly in this section. We shall start with compact metric spaces, approximate them by polyhedra, and then build mappings on these polyhedra by induction on skeleta, using the preceding argument at each step. The crucial point will be that the estimates will depend only on $\rho(\cdot)$ and not on the specific mappings involved.

We want to construct continuous mappings into a metric space given approximately continuous data. We begin with a definition (taken from [P2]).

Definition 5.2. A mapping $f : X \rightarrow Y$ between compact metric spaces is said to be ϵ -continuous if there is a $\delta > 0$ such that $d_X(x_1, x_2) \leq \delta$ implies that $d_Y(f(x_1), f(x_2)) \leq \epsilon$ for all $x_1, x_2 \in X$.

The point here is that we ask for the existence of such a δ for only the one choice of ϵ .

Notation 5.3. If A is a nonempty subset of some metric space M and $t > 0$ is given, then we write A_t for the set of points in M which are at distance $< t$ from A .

The next result says that an ϵ -continuous mapping may be perturbed slightly to get a continuous mapping if we control the local contractability properties of the target.

Proposition 5.4. *Suppose that X is a compact metric space with topological dimension $\leq k$, M is another metric space (not necessarily compact), Z is a subset of M , and $\rho : (0, 1] \rightarrow (0, \infty)$ is a contractability function for (M, Z) . Then for every $\eta > 0$ and $0 \leq s < t \leq 1$ there is an $\epsilon > 0$ so that if $f : X \rightarrow Z_s$ is ϵ -continuous, then we can find a continuous mapping $g : X \rightarrow Z_t$ such that $d_M(f(x), g(x)) \leq \eta$ for all $x \in X$. Here ϵ depends on k, η, s, t , and $\rho(\cdot)$, but not on the specific choices of X, M, Z , or f .*

It is important here that t be larger than s , so that Z_t is larger than Z_s by a definite amount. We shall need this extra bit of room to obtain g from f .

Let X, M, η, s, t , etc. be given as in the proposition. Let $\epsilon > 0$ be small, to be chosen later, and let $f : X \rightarrow Z_s$ be an ϵ -continuous mapping. Let δ be as in Definition 5.2. We want to show that we can find a continuous mapping $g : X \rightarrow Z_t$ as in the proposition if ϵ is small enough. Note that our smallness condition on ϵ may not depend on δ , and that δ depends on our choice of f , which ϵ is not permitted to do.

We need to approximate X by a polyhedron of dimension $\leq n$. (If X is already a polyhedron of dimension $\leq n$, then the argument can be simplified.) Let $U^1, \dots, U^m$ be a

covering of X by finitely many nonempty open sets with order $\leq k$ and mesh $< \delta/2$, as in Definition 4.2. Let $P, \mathcal{S}(P)$ be the polyhedron associated to this covering in the manner described just before Lemma 4.9. We shall use the notation from that discussion freely now. Let $\phi^1, \dots, \phi^m$ be a partition of unity subordinate to $U^1, \dots, U^m$, and let $h : X \rightarrow P$ denote the mapping given by $h = \sum_i \epsilon^i \phi^i$, as in Lemma 4.11. We shall obtain g as a composition of h and another mapping.

Choose and fix a point x^i in each U^i . Note that the ϵ^i 's are precisely the vertices of P .

Lemma 5.5. *If ϵ is small enough (depending only on the parameters allowed in the proposition), then there is a continuous mapping $F : P \rightarrow Z_t$ such that $F(\epsilon^i) = f(x^i)$ for each $i, 1 \leq i \leq m$, and $\text{diam}_M F(\tau) < \eta/2$ for each simplex $\tau \in \mathcal{S}(P)$.*

This follows by induction on skeleta. Let $\mathcal{S}$ be as defined just before Lemma 4.9, and let $\mathcal{S}_j$ be the collection of $S \in \mathcal{S}$ such that S has at most $j+1$ elements, $0 \leq j \leq k$. Let P_j be the polyhedron obtained by taking the union of $\tau(S)$, $S \in \mathcal{S}_j$. Thus P_0 is the set of vertices of P ($= \epsilon^1, \dots, \epsilon^m$), P_1 is the set of vertices and edges, and in general P_j is the union of all the (distinguished) simplices in P of dimension $\leq j$. Set $s_j = s + j(t-s)/k$, so that $s = s_0 < s_1 < \dots < s_k = t$, and set $Z(j) = Z_{s_j}$, $0 \leq j \leq k$. We want to define $F_j : P_j \rightarrow Z(j)$ recursively.

We define $F_0 : P_0 \rightarrow Z(0) = Z_s$ by $F_0(\epsilon^i) = f(x^i)$ for each $i, 1 \leq i \leq m$. If $F_j : P_j \rightarrow Z(j)$ has been defined for some $j < k$ then we want to extend it to $F_{j+1} : P_{j+1} \rightarrow Z(j+1)$ in the following manner. By definitions, $P_{j+1} \setminus P_j$ consists of the interiors of the $(j+1)$ -simplices $\tau(S)$ with $S \in \mathcal{S}$, S having $j+1$ elements. All the faces of such a $\tau(S)$ are of the form $\tau(T)$, $T \in \mathcal{S}_j$, and so F_j is already defined on all the faces of $\tau(S)$. We want to use our local contractibility assumption on (M, Z_1) to extend F_j to these $(j+1)$ -simplices (in the manner described just after Definition 5.1) to get our mapping $F_{j+1} : P_{j+1} \rightarrow Z(j+1)$. To do this we need to be more careful about estimates.

If $S \in \mathcal{S}$, then

$$(5.6) \quad d_M(f(x^i), f(x^l)) \leq \epsilon \quad \text{when } i, l \in S.$$

Indeed, in this case we have that $U^i \cap U^l \neq \emptyset$, and hence $d_X(x^i, x^l) < \delta$, since our covering $U^1, \dots, U^m$ has mesh less than $\delta/2$. Thus (5.6) follows from the ϵ -continuity of f .

Suppose that $F_j : P_j \rightarrow Z(j)$ has already been defined, and that $\text{diam}_M F_j(\tau(S)) \leq \epsilon_j$ for all $S \in \mathcal{S}_j$, where ϵ_j is very small. (When $j = 0$ we use (5.6).) If ϵ_j is small enough, then we can define $F_{j+1} : P_{j+1} \rightarrow Z(j+1)$ as above, by filling it in on the various $(j+1)$ -simplices in P_{j+1} using the local contractibility property of (M, Z_1) . These fillings can be performed independently of each other. Keep in mind that F_{j+1} is supposed to take its values in $Z(j+1)$, for which we need ϵ_j to be quite small. (It is here that we use the fact that $Z(j+1)$ is larger than $Z(j)$ by a definite amount.) If we do this filling in an approximately optimal manner, then we get that $\text{diam}_M F_{j+1}(\tau(S)) \leq \epsilon_{j+1}$ for all $S \in \mathcal{S}_{j+1}$, where ϵ_{j+1} can be controlled in terms of ϵ_j using $\rho(\cdot)$ (without regard to the specific choices of f or F_j), and where $\epsilon_{j+1} \rightarrow 0$ as $\epsilon_j \rightarrow 0$. This permits us to continue the induction for as long as the ϵ_j 's remain sufficiently small. Because we are only interested in $j \leq k$ with k fixed we can ensure that all of the ϵ_j 's are sufficiently small by taking ϵ

small enough. We can compute a bound for how small ϵ should be in terms of k, s, t , and $\rho(\cdot)$, without regard to the specific choices of f or the F_j 's.

We repeat this construction until we get to $j = k$, and then we stop and set $F = F_k$.

We also want to have that $\text{diam}_M F(\tau(S)) \leq \eta/2$ for all $S \in \mathcal{S}$. We can ensure that this holds by requiring that ϵ be sufficiently small, depending on η, k , and $\rho(\cdot)$, since ϵ controls the ϵ_j 's.

This completes the proof of Lemma 5.5, modulo some imprecision which is not hard to clarify. More detailed arguments of this nature can be found in [P1], [P2], with more precise computations.

Now let us finish the proof of Proposition 5.4 using Lemma 5.5. Set $g = F \circ h$. Thus g maps X into Z_t and is continuous. It remains to check that $d_M(f(x), g(x)) \leq \eta$ for all $x \in X$. Fix x , and choose $S \in \mathcal{S}$ so that $h(x) \in \tau(S)$. Choose also an i so that $\phi^i(x) > 0$, and note that $i \in S$, $c^i \in \tau(S)$. Lemma 5.5 says that $\text{diam}_M F(\tau(S)) \leq \eta/2$. Hence $d_X(F(c^i), F(h(x))) \leq \eta/2$. We also have that $F(c^i) = f(x^i)$, and so

$$(5.7) \quad \begin{aligned} d_M(f(x), g(x)) &= d_M(f(x), F(h(x))) \\ &\leq d_M(f(x), f(x^i)) + d_M(f(x^i), F(h(x))) \leq \epsilon + \eta/2. \end{aligned}$$

For the last inequality we have used the assumption that f is ϵ -continuous. Specifically, we have that x and x^i both lie in U^i , whence $d_X(x, x^i) < \delta$ (since our covering $U^1, \dots, U^m$ has mesh $< \delta/2$), and therefore $d_M(f(x), f(x^i)) \leq \epsilon$. From (5.7) we conclude that $d_M(f(x), g(x)) \leq \eta$ if ϵ is small enough.

This completes the proof of Proposition 5.4.

Next we are interested in taking two mappings which are close to each other and showing that they can be homotoped into each other.

Proposition 5.8. *Suppose that X is a compact metric space with topological dimension $\leq k$, M is another metric space (not necessarily compact), Z is a subset of M , and that $\rho : (0, 1] \rightarrow (0, \infty)$ is a contractibility function for (M, Z_1) . For every $\eta > 0$ and $0 \leq s < t \leq 1$ there is an $\epsilon > 0$ so that if $f_0, f_1 : X \rightarrow Z_s$ are continuous mappings with*

$$(5.9) \quad d_M(f_0(x), f_1(x)) \leq \epsilon \quad \text{for all } x \in X,$$

then we can find a continuous mapping $H : X \times [0, 1] \rightarrow Z_t$ such that

$$(5.10) \quad H(x, 0) = f_0(x) \text{ and } H(x, 1) = f_1(x) \quad \text{for all } x \in X, \quad \text{and}$$

$$(5.11) \quad d_M(f_0(x), H(x, \tau)) \leq \eta \quad \text{for all } x \in X \text{ and } \tau \in [0, 1].$$

Here ϵ depends on k, η, s, t , and $\rho(\cdot)$, but not on the specific choices of X, M, Z, f_0 , or f_1 .

The proof of this is much like the proof of Proposition 5.4, except that an infinite construction is required. Again, this is basically contained in the Main Obstruction Result in [P2], with only the difference that our target is not itself locally contractible but rather satisfies a local contractibility condition relative to a larger space.

Let X , M , η , s , t , etc. be given as in the proposition. Let $\epsilon > 0$ be small, to be chosen later, and suppose that $f_0, f_1 : X \rightarrow Z_s$ are continuous mappings which satisfy $d_M(f_0(x), f_1(x)) \leq \epsilon$ for all $x \in X$. We want to build a homotopy H between f_0 and f_1 as in the proposition. The source of the additional complication here is that we really want to specify the values of H on $X \times \{0, 1\}$, and not just specify its approximate behavior. This makes it necessary for us to employ an infinite covering of $X \times (0, 1)$.

Lemma 5.12. *Let $\delta > 0$ be given. We can find a countable covering $\{V^\alpha\}_{\alpha \in A}$ of $X \times (0, 1)$ by nonempty open sets in $X \times (0, 1)$ in such a way that*

$$(5.13) \quad \{V^\alpha\}_{\alpha \in A} \text{ has order } \leq 2k + 1, \text{ i.e., each element of } X \times (0, 1)$$

lies in at most $2k + 2$ of the V^α 's,

$$(5.14) \quad \text{each } V^\alpha \text{ has compact closure in } X \times (0, 1)$$

and intersects only finitely many other V^β 's, and

$$(5.15) \quad \text{diam } V^\alpha \leq \delta \inf\{t(1-t) : (x, t) \in V^\alpha\} \text{ for all } \alpha \in A.$$

(This last condition implies that all the V^α 's have small diameter in $X \times (0, 1)$, and that their diameters shrink to 0 as the V^α 's approach either $X \times \{0\}$ or $X \times \{1\}$.) There is also a partition of unity on $X \times (0, 1)$ which is subordinate to this covering.

The $2k + 1$ in (5.13) should be offensive to some readers, because it really ought to be a $k + 1$. However, $2k + 1$ is fine for our purposes, and it makes the lemma more trivial.

The lemma is extremely easy to prove. Think first about the case where the X is not present, and we are just covering $(0, 1)$. The content of the lemma is that we can cover $(0, 1)$ by tiny open intervals in such a way that no point in $(0, 1)$ lies in more than two of these intervals and the lengths of these intervals goes to 0 as we approach 0 or 1. This is easy to arrange. One can even write down formulae for the intervals. To cover $X \times (0, 1)$ we take one of these little intervals in $(0, 1)$, call it I , and we associate to it a covering of X by finitely many nonempty open sets with order $\leq k$ and with very small mesh, as in Definition 4.2. We then take the Cartesian product of these open subsets of X with I , and the totality of all subsets of $X \times (0, 1)$ obtained in this manner (with I ranging through our initial covering of $(0, 1)$) is our covering of $X \times (0, 1)$. This construction gives (5.13) and (5.14) automatically, and it is easy to arrange (5.15) by choosing the initial covering of $(0, 1)$ and all the subsequent auxiliary coverings of X to be sufficiently fine.

The partition of unity on $X \times (0, 1)$ subordinate to this covering is also easy to obtain. The covering of $(0, 1)$ by tiny intervals certainly has a partition of unity, for which one can even write down formulas. Each of the auxiliary (finite) open coverings on X has a partition of unity subordinate to it too. The required partition of unity on $X \times (0, 1)$ can be produced by taking products.

This proves Lemma 5.12, modulo simple details.

Let us now use Lemma 5.12 to get Proposition 5.8. Let $\delta > 0$ be small enough so that

$$(5.16) \quad d_M(f_l(x), f_l(y)) \leq \epsilon \quad \text{when} \quad d_X(x, y) \leq \delta, \quad l = 0, 1.$$

We can do this because f_0 and f_1 are uniformly continuous, since X is compact. Thus δ depends on f_0 and f_1 , but we do not mind. Let $\{V^\alpha\}_{\alpha \in A}$ be a covering of $X \times (0, 1)$ as in Lemma 5.12, using this choice of δ .

The first thing that we have to do is to construct a polyhedron P associated to this covering. This polyhedron should be constructed in essentially the same manner as described before Lemma 4.9, but now we have to permit it to have infinitely many simplices. Let us work in the (infinite-dimensional) vector space $\mathbf{R}^A$ of real-valued functions on the index set A of our covering $\{V^\alpha\}_{\alpha \in A}$ which vanish except at finitely many points. Let e^α denote the standard basis vector associated to $\alpha \in A$, i.e., the function which satisfies $e^\alpha(\alpha) = 1$, $e^\alpha(\beta) = 0$ for all other $\beta \in A$. Let $\mathcal{S}$ denote the collection of nonempty finite subsets S of A such that $\bigcap_{\alpha \in S} V^\alpha \neq \emptyset$. Thus any $S \in \mathcal{S}$ has at most $2k + 2$ elements, by (5.13). Given $S \in \mathcal{S}$ let $\tau(S)$ denote the simplex in $\mathbf{R}^A$ which is the convex hull of $e^\alpha, \alpha \in S$. We take P to be the union of all the $\tau(S)$'s, $S \in \mathcal{S}$. This is now an infinite polyhedron, which is slightly a nuisance, but it has the nice property that it is locally finite, i.e., no simplex in P touches more than finitely many other simplices. This is not difficult to check, using (5.14). (Note that if two of these simplices intersect, then their intersection must contain a vertex, i.e., an e^α . Also, each vertex e^α lies in only finitely many of these simplices.) Let $\mathcal{S}_j$ denote the set of $S \in \mathcal{S}$ with at most $j + 1$ elements, $0 \leq j \leq 2k + 1$, and set $P_j = \bigcup \{\tau(S) : S \in \mathcal{S}_j\}$. Thus $P = P_{2k+1}$.

Let $\{\phi^\alpha\}_{\alpha \in A}$ be a partition of unity on $X \times (0, 1)$ which is subordinate to $\{V^\alpha\}_{\alpha \in A}$ (whose existence is promised by Lemma 5.12). Define $\Pi : X \times (0, 1) \rightarrow \mathbf{R}^A$ by

$$(5.17) \quad \Pi(x) = \sum_{\alpha \in A} \epsilon^\alpha \phi^\alpha(x).$$

As in Lemma 4.11, Π actually maps $X \times (0, 1)$ into P . Given $S \in \mathcal{S}$ set $V(S) = \bigcup_{\alpha \in S} V^\alpha$. Then

$$(5.18) \quad \Pi^{-1}(\tau(S)) \subseteq V(S) \quad \text{for all } S \in \mathcal{S},$$

just as in the proof of Lemma 4.11.

The homotopy H required in Proposition 5.8 will be defined on $X \times (0, 1)$ by $H = F \circ \Pi$, where $F : P \rightarrow Z_t$ is a map that we need to construct. We shall construct F in stages.

Define s_j for $j = 0, 1, \dots, 2k + 1$ by $s_j = s + j(t - s)/(2k + 1)$, so that $s = s_0 < s_1 < \dots < s_{2k+1} = t$, and set $Z(j) = Z_{s_j}$. We want to define mappings $F_j : P_j \rightarrow Z(j)$, $0 \leq j \leq 2k + 1$, in such a way that each F_j is continuous on all the simplices in P_j and each F_{j+1} is an extension of F_j .

Let us first define F_0 . By definitions, $P_0 = \{e^\alpha : \alpha \in A\}$. Choose, for each $\alpha \in A$, a point $(x^\alpha, t^\alpha) \in V^\alpha$. Define F_0 by

$$(5.19) \quad \begin{aligned} F_0(e^\alpha) &= f_0(x^\alpha) & \text{when } t^\alpha \in (0, 1/2] \\ &= f_1(x^\alpha) & \text{when } t^\alpha \in (1/2, 1). \end{aligned}$$

Let us check that F_0 does not oscillate too much, in the sense that

$$(5.20) \quad \text{diam}_M F_0(\tau(S) \cap P_0) \leq 2\epsilon \quad \text{for all } S \in \mathcal{S}.$$

Let $S \in \mathcal{S}$ be given, and let $\alpha, \beta \in S$ be arbitrary. This amounts to choosing arbitrary elements e^α and e^β of $\tau(S) \cap P_0$. We have that $V^\alpha \cap V^\beta \neq \emptyset$, by definition of $\mathcal{S}$. This implies that

$$(5.21) \quad d_X(x^\alpha, x^\beta) \leq \text{diam}_X V^\alpha + \text{diam}_X V^\beta \leq \delta,$$

where the last inequality comes from (5.15). Using (5.16) we get that

$$(5.22) \quad d_M(f_l(x^\alpha), f_l(x^\beta)) \leq \epsilon, \quad l = 0, 1.$$

This implies that

$$(5.23) \quad d_M(F_0(x^\alpha), F_0(x^\beta)) \leq 2\epsilon,$$

because of the definition (5.19) of F_0 and our assumption (5.9). This proves (5.20).

We want to construct the F_j 's for $j > 0$ recursively. Suppose that $F_j : P_j \rightarrow Z(j)$ has been defined already, and that

$$(5.24) \quad \text{diam}_M F_j(\tau(S)) \leq \epsilon_j \quad \text{when } S \in \mathcal{S} \text{ has exactly } j+1 \text{ elements,}$$

where ϵ_j is some small number. If ϵ_j is small enough, then we can extend F_j to a mapping $F_{j+1} : P_{j+1} \rightarrow Z(j+1)$ in the obvious way. Indeed, P_{j+1} is obtained from P_j by adding to P_j the $(j+1)$ -simplices, or, in other words, the simplices $\tau(S)$ such that $S \in \mathcal{S}$ has exactly $j+2$ elements. Fix such an S , and notice that all of the faces of $\tau(S)$ are among the simplices in P_j . (This statement uses the easy observation that if $S \in \mathcal{S}$, then each nonempty subset of S also lies in $\mathcal{S}$.) Thus F_j is already defined on the faces of $\tau(S)$, and we want to extend it to all of $\tau(S)$. We can do this if ϵ_j is small enough, because of our contractability assumption for (M, Z_1) . That is, (5.24) will ensure that $\text{diam}_M F_j(\partial\tau(S))$ will have small diameter in $Z(j)$, and this implies that we can fill it in inside $Z(j+1)$ if ϵ_j is small enough, as discussed just after Definition 5.1. (When $j = 0$ one should use (5.20) instead of (5.24).) By doing this for each of the $\tau(S)$'s we get $F_{j+1} : P_{j+1} \rightarrow Z(j+1)$.

When we define F_{j+1} in this manner, we should be careful to choose each of these fillings on the $(j+1)$ -simplices so that their diameters are approximately as small as possible (to within a factor of 2, say). This will give

$$(5.25) \quad \text{diam}_M F_{j+1}(\tau(S)) \leq \epsilon_{j+1} \quad \text{when } S \in \mathcal{S} \text{ has exactly } j+2 \text{ elements,}$$

where ϵ_{j+1} can be controlled in terms of ϵ_j using $\rho(\cdot)$, and in such a way that $\epsilon_{j+1} \rightarrow 0$ when $\epsilon_j \rightarrow 0$. We can repeat the construction if ϵ_{j+1} is small enough. This will happen automatically if ϵ_j was sufficiently small.

The conclusion of this induction is that we can construct all the mappings $F_j : P_j \rightarrow Z(j)$ with $0 < j \leq 2k+1$ if ϵ is small enough. That is, we can ensure that all the ϵ_j 's are sufficiently small by taking ϵ small enough, because of our inductive control on the ϵ_j 's, and because we only need to deal with a bounded number of j 's.

Set $F = F_{2k+1}$. Define $H : X \rightarrow [0, 1]$ by setting $H = F \circ \Pi$ on $X \times (0, 1)$ and using (5.10) on $X \times \{0, 1\}$.

Lemma 5.26. *H is continuous on $X \times [0, 1]$.*

The continuity of H on $X \times (0, 1)$ follows from the continuity of F and Π . Strictly speaking this statement does not make sense, because we have not defined a topology on our infinite polyhedron P , but in this case we only need the obvious topology on the finite

parts of P . That is, if we restrict ourself to a compact subset of $X \times (0, 1)$, then Π takes values in a finite polyhedron contained in P (because of (5.14)), and the continuity issues become especially transparent. (We could also have defined a topology on P , but this is not necessary.)

The continuity of H on $X \times \{0, 1\}$ is slightly more tricky. The main point is that

$$(5.27) \quad \begin{aligned} &\text{for any (small) } \theta > 0 \text{ we have that } \text{diam}_M F(\tau) < \theta \\ &\text{for all but finitely many (distinguished) simplices } \tau \text{ in } P. \end{aligned}$$

To prove (5.27) we make an induction on skeleta as before. We start with an analogue of (5.20). For each $\theta_0 > 0$ we have that

$$(5.28) \quad \text{diam}_M F_0(\tau \cap P_0) < \theta_0$$

for all but finitely many simplices τ . This is an easy consequence of the definition (5.19) of F_0 , the continuity of f_0 and f_1 , and (5.15). In ordinary words, the main point is that the simplices τ come from elements of our covering $\{V^\alpha\}$ of $X \times (0, 1)$, and all but finitely many of the τ 's are associated to V^α 's which live near "the boundary" $X \times \{0, 1\}$ and which therefore have very small diameter, by (5.15). This permits us to derive (5.28) from the (uniform) continuity of f_0 and f_1 . Next we need a version of (5.24) in this context. Given $j \leq 2k + 1$ and $\theta_j > 0$, we have that

$$(5.29) \quad \text{diam}_M F_j(\tau \cap P_j) < \theta_j$$

for all but finitely many simplices τ . This is proved by induction on j , using the same type of argument as we used to go from (5.24) to (5.25). That is, we obtained F_{j+1} from F_j by taking each $(j+1)$ -simplex σ in P , observing that F_j was already defined on $\partial\sigma$, and then extending this map on $\partial\sigma$ to all of σ using the local contractibility of (M, Z_1) . If we know (5.29) for j (use (5.28) for $j = 0$), then $\text{diam}_M F_j(\partial\tau)$ is as small as we like for all but finitely many τ 's, and the local contractibility condition implies that $\text{diam}_M F_{j+1}(\tau)$ is as small as we like for all but finitely many τ 's. (For this point it is important that we chose our fillings in such a way that $\text{diam}_M F_{j+1}(\tau)$ is always approximately as small as it can be. Of course we are using here the argument described just after Definition 5.1.) Thus we can get (5.29) for all $j \leq 2k + 1$, by induction. This implies (5.27).

Now that we have (5.27) let us check that H is continuous on $X \times \{0, 1\}$. Fix $x \in X$ and let us check the continuity of H at $(x, 0)$, say. Let $(y, t) \in X \times (0, 1)$ be given, with (y, t) close to $(x, 0)$. Choose an $\alpha \in A$ so that $\phi^\alpha(y, t) > 0$, where ϕ^α comes from our partition of unity (as in (5.17)). Let S denote the set of $\beta \in A$ such that $(y, t) \in V^\beta$. Then $\alpha \in S$, $S \in \mathcal{S}$, and $\Pi(y, t) \in \tau(S)$, by definitions. Using (5.27) we get that $\text{diam}_M F(\tau(S))$ is as small as we like when t is sufficiently small. Since ϵ^α is a vertex of $\tau(S)$ (by definitions), we conclude that $d_M(H(y, t), F(\epsilon^\alpha)) = d_M(F(\Pi(y, t)), F(\epsilon^\alpha))$ is as small as we like if t is small enough. On the other hand we have that $F(\epsilon^\alpha) = F_0(\epsilon^\alpha) = f_0(x^\alpha)$, by (5.19), and so we can make $d_M(H(y, t), f_0(x^\alpha))$ as small as we like by taking t small enough. Since (x^α, t^α) and (y, t) both lie in V^α we have that $d_X(x^\alpha, y) \leq \text{diam } V^\alpha \leq t$, because of (5.15). We are assuming that f_0 is continuous, and hence uniformly continuous, since X is compact.

This means that $d_M(f_0(x^\alpha), f_0(y))$ is as small as we like when t is sufficiently small, and hence $d_M(H(y, t), f_0(y))$ is as small as we like. Using again the uniform continuity of f_0 we obtain that $d_M(H(y, t), f_0(x))$ is small (since (y, t) is assumed to be close to $(x, 0)$). This implies that H is continuous at $(x, 0)$, which is what we wanted. (Of course the argument for $(x, 1)$ is the same.)

This proves Lemma 5.26.

It remains to show that (5.11) holds (if ϵ is small enough, depending on η). Let $x \in X$ and $t \in [0, 1]$ be given. We may as well assume that $t \in (0, 1)$, since otherwise (5.11) follows from (5.10) and (5.9). As in the preceding argument, we fix $\alpha \in A$ so that $\phi^\alpha(x, t) > 0$, and we let S denote the set of $\beta \in A$ such that $(x, t) \in V^\beta$. We get that $\alpha \in S$, $S \in \mathcal{S}$, $\Pi(x, t) \in \tau(S)$, and e^α is a vertex of $\tau(S)$. If $(x^\alpha, t^\alpha) \in V^\alpha$ is as in (5.19), then $d_X(x, x^\alpha) \leq \text{diam } V^\alpha \leq \delta$, by (5.15). Thus (5.16) implies that $d_M(f_0(x), f_0(x^\alpha)) \leq \epsilon$, and (5.9) yields $d_M(f_0(x), f_1(x^\alpha)) \leq 2\epsilon$. Therefore $d_M(f_0(x), F_0(e^\alpha)) \leq 2\epsilon$, by (5.19). On the other hand we have that $d_M(F(\Pi(x, t)), F_0(e^\alpha)) \leq \text{diam}_M F(\tau(S))$ is small in a way which is controlled by ϵ , because of (5.24) with $j = 2k + 1$ and the fact that the ϵ_j 's are controlled in terms of ϵ . In particular $d_M(F(\Pi(x, t)), F_0(e^\alpha)) \leq \eta/2$ if ϵ is small enough. Thus $d_M(F(\Pi(x, t)), f_0(x)) \leq \eta/2 + 2\epsilon \leq \eta$ if ϵ is small enough. This says exactly that $d_M(H(x, t), f_0(x)) \leq \eta$, which is what we wanted.

This completes the proof of Proposition 5.8. For more details or precise estimates see the Main Obstruction Result and its proof in [P2].

Note that there is more care taken in [P2] about exactly what kind of contractability assumptions are needed to make this argument work. For instance, we do not really need to be able to contract whole balls in Definition 5.1, it is enough to be able to contract mappings from spheres of certain dimensions into M . We shall ignore this point, but one should not forget it. Also, [P1] and [P2] are more precise about the estimates, which means the relationship between η and ϵ in Propositions 5.4 and 5.8.

6. Topological dimension with bounds.

We are going to need a quantitative version of topological dimension (Definition 4.2). Recall Notation 5.3, which we shall use frequently in this section.

Definition 6.1. Let X be a compact metric space, let k be a nonnegative integer, and let $\delta_k(\epsilon)$ be a positive function on $(0, 1]$ which satisfies $\delta_k(\epsilon) \leq \epsilon$ for each ϵ . We say that X has topological dimension function δ_k (with dimension k) if for each $\epsilon > 0$ we can cover X by finitely many nonempty subsets A^i such that $\text{diam } A^i < \epsilon$ for each i and no point in X lies in more than $k + 1$ of the A_k^i 's, where $\delta = \delta_k(\epsilon)$ and A_k^i is as in Notation 5.3.

This definition should be compared with the notion of asymptotic dimension in §1 F of [Gr4], which controls the asymptotic behavior of an unbounded metric space at infinity. See also [Y] in connection with this notion.

Lemma 6.2. *If X is a compact metric space, then X has topological dimension $\leq k$ if and only if there is a function $\delta_k : (0, 1] \rightarrow (0, 1]$ such that X has topological dimension function δ_k (with dimension k).*

This is not hard to derive from Definitions 4.2 and 6.1. (Exercise. The main point is that if a compact set X is covered by a collection of open subsets, then there is a $\delta > 0$ so that for each $x \in X$ the ball $B(x, \delta)$ is wholly contained in one of the elements of the covering. This implies that the compact set is also covered by open sets that are slightly smaller, by the amount δ .)

It is useful to have a bound on the topological dimension of a metric space in order to build mappings from that metric space into other spaces. We saw examples of this phenomenon in the preceding section. In this paper we shall need to make some constructions of mappings with controlled moduli of continuity, and for this it will be necessary to have control on the topological dimension functions. In particular we shall need to have criteria for the existence of a uniform choice of a topological dimension function for a family of compact metric spaces.

Proposition 6.3. *Let $\mathcal{F}$ be a precompact family of compact metric spaces (Definition 3.5) and let k be a nonnegative integer. Then there is a (single) function $\delta_k : (0, 1] \rightarrow (0, 1]$ such that every $X \in \mathcal{F}$ has topological dimension function δ_k (with dimension k) if every $X \in \mathcal{F}$ has topological dimension $\leq k$ and if every compact metric space which arises as the limit of a sequence in $\mathcal{F}$ (as in Definition 3.1) has topological dimension $\leq k$.*

The converse is true too (and easy to check), but we shall not need it.

To prove this we use the obvious argument. Suppose that $\mathcal{F}$ and k satisfy the conditions in the proposition, but that there is no single topological dimension function δ_k that works for all $X \in \mathcal{F}$. This means that there is an $\epsilon > 0$ with the property that for each positive integer j we can find an $X(j) \in \mathcal{F}$ which cannot be covered by finitely many sets $A^i \subseteq X(j)$ such that $\text{diam } A^i < \epsilon$ for each i and no element of $X(j)$ lies in A^i for more than $k + 1$ choices of i . The precompactness of $\mathcal{F}$ implies that there is a subsequence of $\{X(j)\}$ which converges in the sense of Definition 3.1 to a compact metric space X , and our hypotheses imply that X has topological dimension $\leq k$. Hence there is a $\delta > 0$ and a finite family of subsets E^i of X such that $X \subseteq \bigcup_i E^i$ and no element of X lies in E^i for more than $k + 1$ choices of i . It is easy to use these E^i 's to induce similar coverings of $X(j)$'s for arbitrarily large values of j (but with δ replaced by $\delta/2$ for the $X(j)$'s). This contradicts our assumption about the $X(j)$'s, and the proposition follows.

Proposition 6.4. *Let $\mathcal{F}$ be a family of compact metric spaces with bounded diameter with the property that there is a function $N : (0, 1] \rightarrow (0, \infty)$ such that for every $0 < \epsilon \leq 1$ and every $X \in \mathcal{F}$ we can cover X by at most $N(\epsilon)$ closed balls in X of radius ϵ . Assume also that $N(t) = O(t^{-\mu})$ for some $\mu \geq 0$, and that k is an integer with $\mu < k + 1$. Then there is a (single) function $\delta_k : (0, 1] \rightarrow (0, 1]$ such that every $X \in \mathcal{F}$ has topological dimension function δ_k (with dimension k).*

Notice first that every $X \in \mathcal{F}$ has Hausdorff dimension $\leq \mu$, by Lemma 2.17. This implies that the topological dimension of each $X \in \mathcal{F}$ is $\leq k$, by Theorem 4.6.

Now suppose that X is a compact metric space which arises as the limit of a sequence in $\mathcal{F}$. Then $N(\cdot)$ also controls the covering properties of X , by Lemma 3.4. Therefore X has Hausdorff dimension $\leq \mu$, and hence topological dimension $\leq k$ also. Proposition 6.4 now follows from Proposition 6.3.

Theorem 6.5. *Fix functions $N : (0, 1] \rightarrow (0, \infty)$ and $\rho : (0, 1] \rightarrow (0, \infty)$ such that $N(t) = O(t^{-k})$ for some $k \geq 0$ and $\rho(t) \geq t$ for all t . Fix a nonnegative integer n too. Let $\mathcal{A}$ denote the collection of all admissible triples $(M, U, d(x, y))$ (Definition 1.5) with $\text{diam } U = 1$, covering function $N(\cdot)$, contractability function $\rho(\cdot)$ (as in Definitions 1.6 and 1.7), and such that $\bar{U}_1$ has topological dimension $\leq n$. For each such triple $(M, U, d(x, y))$ and any $t > 0$ define $U_t \subseteq M$ by $U_t = \{x \in M : \text{dist}(x, U) < t\}$. Define $\mathcal{F}_t$ to be the family of metric spaces of the form $(\bar{U}_t, d(x, y))$ with $(M, U, d(x, y)) \in \mathcal{A}$. Then for each $t \in (0, 1)$ there is function $\delta_n : (0, 1] \rightarrow (0, 1]$ (depending on t) such that every $X \in \mathcal{F}$ has topological dimension function δ_n (with dimension n). (Of course this function δ_n depends on k, N , and ρ too.)*

Observe that there is no point in taking $n > k$, we can always replace it with $\min(n, k)$, because of Proposition 6.4.

Thus we are gaining uniform control on the topological dimension functions from our uniform assumptions on the covering and contractability functions and from the requirement that our metric spaces have topological dimension at most n . Note that k is allowed to be as large as it wants, independently of n .

The proof of this theorem is based on the same argument as the one at the end of Section 4 of [GP]. Suppose that we are given our initial objects as in the theorem, and fix $t \in (0, 1)$. Theorem 3.3 (and Definition 1.5) imply that $\mathcal{F}_t$ is a precompact family of compact metric spaces. In view of Proposition 6.3 it suffices to show that if $(X, D(\cdot, \cdot))$ is a compact metric space which arises as the limit of a sequence $\{X^j\}$ in $\mathcal{F}_t$, then X has topological dimension $\leq n$.

Notice that X has topological dimension $\leq k$. Indeed, its covering properties are controlled by the function $N(\cdot)$, by Lemma 3.4, and our assumption on N implies that X has Hausdorff dimension $\leq k$, by Lemma 2.17. Thus X has topological dimension $\leq k$ by Theorem 4.6.

Let $\{(M^j, U^j, d_j(\cdot, \cdot))\}$ be the sequence of elements in $\mathcal{A}$ which corresponds to $\{X^j\}$.

Lemma 6.6. *There is a sequence of mappings $f_j : X \rightarrow X^j$, not necessarily continuous, such that*

$$(6.7) \quad \lim_{j \rightarrow \infty} \sup\{|d_j(f_j(x), f_j(y)) - D(x, y)| : x, y \in X\} = 0.$$

This is an easy consequence of the convergence of X^j to X . (See Definition 3.1.)

Lemma 6.8. *There is a sequence of continuous mappings $g_j : X \rightarrow U_1^j$ such that*

$$(6.9) \quad \lim_{j \rightarrow \infty} \sup\{|d_j(g_j(x), g_j(y)) - D(x, y)| : x, y \in X\} = 0.$$

This follows from Lemma 6.6 and Proposition 5.4. That is, the mappings f_j in Lemma 6.6 are ϵ -continuous with choices of ϵ which tend to 0 as $j \rightarrow \infty$, and Proposition 5.4 permits us to modify them slightly to get g_j 's which are continuous. Note that we are allowing the g_j 's to take values in slightly larger spaces than the f_j 's, which is important

for Proposition 5.4. Also, the fact that our targets depend on j causes no problems, because we have uniform estimates.

To finish the proof of Theorem 6.5 we need to show that X has topological dimension $\leq n$. According to Theorem 4.5 it is enough to show that for each $\epsilon > 0$ there is a finite polyhedron $P(\epsilon)$ of dimension $\leq n$ and an ϵ -mapping (Definition 4.4) from X into $P(\epsilon)$. Theorem 4.5 and our assumptions ensure that there are $\epsilon/2$ -mappings from the U_1^j 's into finite polyhedra of dimension $\leq n$. We can compose these mappings with the mappings provided by Lemma 6.8 (for sufficiently large j) to get ϵ -mappings from X into polyhedra of dimension $\leq n$. Thus X has topological dimension $\leq n$.

This completes the proof of Theorem 6.5.

Remark 6.10. In Standard Assumptions 1.8 and 1.10 we are requiring a bound on the local contractability properties of our spaces, with the bound being given by the contractability function $\rho(\cdot)$. This is a strong assumption, and it is worthwhile to try to understand how it is being used, and how it can be weakened. One very important way in which we use this bound is in applying Theorem 6.5 in order to get a bound on the topological dimension function δ_n . However, one could also control the topological dimension function δ_n through other means, e.g., in terms of bounds on the geometric “size” of the space, as in Proposition 6.4. (See also the second paragraph of Remark B.21.)

This bound on the topological dimension function will be very important for us. We saw in Section 4 how a compact metric space X with topological dimension $\leq n$ can be approximated by a finite polyhedron P of dimension $\leq n$, in the sense that we can build a continuous map $f : X \rightarrow P$ for which the inverse images of the simplices in P have small diameter. We shall see in the next section that a bound on the topological dimension function will permit us to do this in such a way that we can control the Lipschitz norm of f . This fact will be crucial for us; it will be the source of our constructions of special Lipschitz mappings with bounds.

We shall also use our bound on the contractability function in very different way. If our space X is well approximated by a polyhedron P in the manner just described, with a map $f : X \rightarrow P$ as above, then the bound on local contractability permits us to build a mapping $g : P \rightarrow X$ (or into a slightly larger space) which is approximately the inverse of f , and in such a way that $g \circ f$ is homotopic to the identity (or to an inclusion). These approximate inverses and homotopies are obtained using Propositions 5.4 and 5.8. However, we never really need such strong approximate inverses, at the level of mappings and homotopies, but rather only approximate inverses at the level of n -dimensional (relative) homology and cohomology groups. This is a much weaker requirement.

More precisely, in the proofs of Theorems 1.29 (a) and 1.30 in Sections 8 and 10 we shall be faced with precisely this point of approximating our space by a finite polyhedron and approximately inverting the mapping f . The arguments will not be written in such a manner as to maximize the relevance of homology or cohomology over actual mappings – the arguments become messier that way – but this point will be in the background, and we shall make comments about it. (See Remarks 8.26 and 10.52.) The proof of Theorem 1.29 (b) in Section 9 comes down exactly to showing a correspondence between relative homology classes associated to the orientations of our spaces (as in Theorem A.17). This will be a little different from the situation just described, because it will involve maps

between two of our spaces rather than a map from a given space into an approximating polyhedron.

We shall also use the local contractability assumption in a third and less exciting way. In Theorem 1.29 (a) we need to assume that U be connected, and this requirement propagates to the need for making other connected sets in other places. Specifically, it is necessary to do this in Sublemmas 9.38 and 9.39, Lemma 11.10, and Sublemma 1.17. For this purpose we do not need a bound on local contractability, we just need to know that any given pair of points can be put inside a connected set of controlled diameter.

Remark 6.11. Our assumption that $N(t) = O(t^{-k})$ in Theorem 6.5 is needed only to ensure that the limiting space X in the proof has finite topological dimension. We assumed that $N(t) = O(t^{-k})$ in Definition 1.6 and elsewhere in this paper only for this step.

7. Mappings into polyhedra with Lipschitz bounds.

Theorem 4.5 says that a compact metric space with topological dimension $\leq n$ admits continuous mappings into polyhedra with dimension $\leq n$ and with small fibers. We are going to need a quantitative version of this statement, in which we control the complexity of the polyhedron and the Lipschitz norm of the mappings.

Theorem 7.1. *Let a positive integer n and functions $N, \delta_n : (0, 1] \rightarrow [0, \infty)$ be given, and let $\mathcal{F}$ denote the collection of compact metric spaces X such that X can be covered by at most $N(\epsilon)$ closed balls in X of radius ϵ for each $0 < \epsilon \leq 1$ and such that X has topological dimension function δ_n (Definition 6.1) with dimension n . Then for each $0 < \epsilon \leq 1$ there exists an integer l , a finite collection $\mathcal{P}$ of finite polyhedra in $\mathbf{R}^l$ all with dimension $\leq n$, and a constant C – all depending on $n, \epsilon, N(\cdot)$, and $\delta_n(\cdot)$ but not on anything else – such that for each $X \in \mathcal{F}$ there is a polyhedron $P \in \mathcal{P}$ and a C -Lipschitz map $f : X \rightarrow P$ such that if τ is a (distinguished) simplex in P , then $\text{diam}_X f^{-1}(\tau) \leq \epsilon$.*

For the record, we are using the metric on P induced from the Euclidean metric on $\mathbf{R}^l$ here.

Let $X \in \mathcal{F}$ and $0 < \epsilon \leq 1$ be given, and set $\delta = \delta_n(\epsilon/6)$. Let $\{A^i\}_{i=1}^m$ be as in Definition 6.1, but with ϵ replaced with $\epsilon/6$, so that $X \subseteq \bigcup_i A^i$, $\text{diam } A^i < \epsilon/6$ for each i , and no element of X lies in more than $n+1$ of the A_δ^i 's, where A_δ^i is defined as in Notation 5.3.

Let us bound the number m of A^i 's. By assumption we can cover X by at most $N(\delta/2)$ closed balls of radius $\delta/2$. If B is any ball in X of radius $\delta/2$, then there are not more than $n+1$ of the A^i 's which intersect B , since no point in X lies in more than $n+1$ of the A_δ^i 's. Thus we conclude that $m \leq (n+1) \cdot N(\delta/2)$. Note that this bound does not depend on X .

Set $U^i = A_\delta^i$, so that $U^1, \dots, U^m$ is a covering of X with order $\leq n+1$. Let P be the polyhedron in $\mathbf{R}^m$ as constructed just before Lemma 4.9, so that P has dimension $\leq n$. We shall use the same notations ($e^i, S, \mathcal{S}, \tau(S)$) as before.

It is clear that the set $\mathcal{P}$ of polyhedra constructed from metric spaces in $\mathcal{F}$ in this manner is finite, and that we can consider them as all living in some fixed $\mathbf{R}^l$ (that does not depend on X).

It remains to construct our mapping f . We shall do this in the same manner as in Lemma 4.11, except that we must choose carefully our partition of unity in order to get Lipschitz bounds. Define auxiliary mappings $\psi^i : M \rightarrow [0, 1]$ by

$$(7.2) \quad \psi^i(p) = \max(1 - \delta^{-1} \text{dist}(x, A^i), 0) \quad \text{for } x \in X.$$

Thus ψ^i is Lipschitz with norm $\leq \delta^{-1}$, $\psi^i \equiv 1$ on A^i , and $\psi^i \equiv 0$ on $X \setminus U^i$. To convert this into a partition of unity we use the usual recipe. Observe that $1 \leq \sum_j \psi^j \leq n+1$ on X . Set

$$(7.3) \quad \phi^i = \psi^i \left(\sum_j \psi^j \right)^{-1}.$$

Thus the ϕ^i 's satisfy $0 \leq \phi^i \leq 1$ on X , $\sum \phi^i \equiv 1$ on X , $\phi^i \equiv 0$ on $X \setminus U^i$, and each ϕ^i is $C(n, \delta)$ -Lipschitz. The Lipschitz condition comes from the Lipschitz condition on the ψ^i 's, and is not hard to verify by direct computation.

Define $f : X \rightarrow \mathbf{R}^m$ by

$$(7.4) \quad f(x) = \sum_{i=1}^m \epsilon^i \phi^i(x) \quad \text{for } x \in X.$$

This is the same formula as in Lemma 4.11, the only difference is that we have chosen a specific partition of unity $\{\phi^i\}$. Thus the conclusions of Lemma 4.11 are still valid in this case. Specifically, f takes values in P , and $\text{diam } f^{-1}(\tau)$ is at most twice the mesh of the covering $U^1, \dots, U^m$ of X for all the (distinguished) simplices τ in P . In our case the mesh of $U^1, \dots, U^m$ is $\leq \epsilon/2$, because of the way that $\{A^i\}$ and δ were chosen, and so we get that $\text{diam } f^{-1}(\tau) \leq \epsilon$ for all simplices τ in P .

It is easy to see that f is Lipschitz with bounded norm, since the ϕ^i 's are. This completes the proof of Theorem 7.1.

Remark 7.5. In Theorem 7.1 we needed to have a controlled covering function $N(\cdot)$ for $X \in \mathcal{F}$ only to ensure that the collection $\mathcal{P}$ of polyhedra would be finite, and to get a bound on the dimension l of the Euclidean spaces in which they live. If we drop the assumption of a uniform covering function $N(\cdot)$, then we can still get a finite polyhedron P as in Theorem 7.1 and a Lipschitz mapping f which enjoys the same estimates as before. In particular, the bound on the Lipschitz norm of f does not depend on $N(\cdot)$, as one can easily check. (When one goes to estimate $|f(x) - f(y)|$ for any pair of points $x, y \in X$, one need not deal with more than $2(n+1)$ choices of i .) In order for this Lipschitz bound to be meaningful one should remember how the polyhedron P is constructed; if we do not impose any conditions on P , then the Lipschitz bound is meaningless, because we could simply rescale the polyhedron. In fact we get our Lipschitz bound while taking P to be of a very special form, as in Remark 4.12.. We obtained P from the construction which precedes Lemma 4.9, and that construction produces a polyhedron P in some Euclidean space in which every distinguished simplex in P is the convex hull of standard basis vectors of the Euclidean space. In other words, all the vertices are standard basis vectors. With this restriction

the Lipschitz bound for f as a mapping into P has content, because the distinguished simplices in P are not permitted to become small or to degenerate. Although we permit the complexity of P to be unbounded, in terms of the number of these distinguished simplices, we do not permit the distinguished simplices themselves to degenerate.

8. Mappings into spheres with nonzero degree and Lipschitz bounds.

In this section we shall prove part (a) of Theorem 1.29. This means that we want to construct a Lipschitz map from a given space into a sphere in such a way that the map is constant outside of a given ball, it has nonzero degree, and its Lipschitz norm is bounded. Let $(M, U, d(x, y)), \sigma, B$, and δ be given as in Theorem 1.29. We also have the functions $N(\cdot)$ and $\rho(\cdot)$ coming from Standard Assumptions 1.8. We shall use heavily the results and definitions in Appendix A concerning degree theory and homology. Recall that M is required to be an oriented topological manifold of dimension n in Standard Assumptions 1.8, so that we can talk about the degree of mappings from M into $\mathbf{S}^n$ which are constant off a compact set.

We may also assume that $\text{diam } U = 1$, since otherwise we can rescale the metric (and because the statement of Theorem 1.29 scales properly).

Let us write $U(t)$ for $\bar{U}_t$, where U_t is as in Notation 5.3.

As mentioned in the introduction, we could drop the assumption of a fixed covering function $N(\cdot)$ from Theorem 1.29 (a) if we are willing to assume control on the topological dimension function. We shall spell this point out in Remark 8.24, but the reader may wish to keep this point in mind while going through the arguments that follow.

Remark. In the next three lemmas we do not need to assume as much structure as in Theorem 1.29 (a). We do not need to assume that M is an oriented topological manifold. We shall need this fact in Section 10, and so we spell it out: in Lemmas 8.1, 8.2, and 8.3 it suffices to assume that $(M, U, d(x, y))$ is an admissible triple (Definition 1.5) with covering function $N(\cdot)$ (Definition 1.6), contractability function $\rho(\cdot)$ (Definition 1.7), that $\bar{U}_1$ has topological dimension $\leq n$ (Definition 4.2), and that $\text{diam } U = 1$. Keep in mind that we are requiring in Definition 1.6 that the covering function $N(\cdot)$ satisfy $N(t) = O(t^{-k})$ for some $k < \infty$.

Lemma 8.1. *There is a function $\delta_n : (0, 1] \rightarrow (0, 1]$ so that $U(1/2)$ has topological dimension function $\delta_n(\cdot)$ with dimension n . Here $\delta_n(\cdot)$ depends on $N(\cdot)$, and $\rho(\cdot)$, but nothing else.*

This follows from Theorem 6.5. The reader will please forgive the confusing use of $\delta_n(\cdot)$ in this lemma and will hopefully not confuse it with the parameter $\delta > 0$ from Theorem 1.29 (a).

Lemma 8.2. *For each $\epsilon > 0$ there is a finite polyhedron P with dimension $\leq n$ and a C -Lipschitz mapping $f : U(1/2) \rightarrow P$ such that $\text{diam } f^{-1}(\tau) \leq \epsilon$ for all (distinguished) simplices τ in P . Here C depends only on $n, \epsilon, N(\cdot)$, and $\rho(\cdot)$, and P is an element of a finite collection of finite polyhedra in some $\mathbf{R}^l$ which depends only on the same parameters.*

This follows from Theorem 7.1 and Lemma 8.1.

This notation is slightly bad. In Theorem 1.29 (a) we called the map that we are seeking f , and that map is not the mapping f in the lemma. For more global reasons it seems better to stick to this notation, and to leave unnamed the mapping that we are trying to construct.

Lemma 8.3. *For each $\theta > 0$ there is a finite polyhedron P with dimension $\leq n$, a C -Lipschitz mapping $f : U(1/2) \rightarrow P$, a continuous mapping $g : P \rightarrow U(3/4)$, and a continuous mapping $H : U(1/2) \times [0, 1] \rightarrow U(4/5)$ with the following properties:*

$$(8.4) \quad \text{diam } f^{-1}(\tau) \leq \theta \quad \text{for all (distinguished) simplices } \tau \text{ in } P;$$

$$(8.5) \quad \text{diam } g(\tau) \leq \theta \quad \text{for all (distinguished) simplices } \tau \text{ in } P;$$

$$(8.6) \quad d_M(g(f(x)), x) \leq \theta \quad \text{for all } x \in U(1/2);$$

$$(8.7) \quad H(x, 0) = x \text{ and } H(x, 1) = g(f(x)) \text{ for all } x \in U(1/2);$$

$$(8.8) \quad d_M(x, H(x, t)) \leq \theta \quad \text{for all } x \in U(1/2) \text{ and } t \in [0, 1].$$

Here C depends only on $n, \theta, N(\cdot)$, and $\rho(\cdot)$, and P is an element of a finite collection of finite polyhedra in some $\mathbf{R}^l$ which depends only on the same parameters.

Note that we do not get any bounds on the moduli of continuity of g or H .

Let $\theta > 0$ be given, and let $\alpha, \beta > 0$ be small, to be chosen later. We shall let β depend on θ and α depend on β and θ , and we require that $\alpha < \beta/10 < \theta/10$. Let P and f be as in Lemma 8.2, with $\epsilon = \alpha/2$, so that

$$(8.9) \quad \text{diam } f^{-1}(\tau) \leq \frac{\alpha}{2}$$

for all distinguished simplices τ in P . Let $g_0 : P \rightarrow U(1/2)$ be a mapping, possibly discontinuous, such that

$$(8.10) \quad g_0(p) \in f^{-1}(p) \quad \text{for each } p \in P.$$

We get from (8.9) that

$$(8.11) \quad \text{diam } g_0(\tau) \leq \frac{\alpha}{2}$$

for all (distinguished) simplices τ in P . One can check that this implies that g_0 is α -continuous (Definition 5.2). (If two points in P are sufficiently close to each other, then they lie in simplices which touch.) This permits us to apply Proposition 5.4 with f taken to be g_0 , $X = P$, $k = n$, $Z = U$, $s = 1/2$, $t = 3/4$, $\eta = \beta/2$, and $\epsilon = \alpha$. We obtain that there is a continuous mapping $g : P \rightarrow U(3/4)$ such that

$$(8.12) \quad d_M(g_0(p), g(p)) \leq \beta/2 \quad \text{for all } p \in P.$$

if α is small enough, depending on β . This implies that

$$(8.13) \quad \text{diam } g(\tau) \leq \alpha/2 + \beta$$

for all distinguished simplices τ in P , because of (8.11). We also get that

$$(8.14) \quad d_M(g(f(x)), x) \leq \alpha/2 + \beta/2 \leq \beta$$

for all $x \in U(1/2)$, by (8.10), (8.9), and (8.12).

(Note that the proof of Proposition 5.4 could be simplified somewhat for this application. The previous use of coverings and partitions of unity are not needed in this case because $X = P$ is already a polyhedron, one can proceed directly to the induction on skeleta.)

In order to get our homotopy H we apply Proposition 5.8 with $f_0 =$ the identity, $f_1 = g \circ f$, $X = U(1/2)$, $k = n$, $Z = U$, $s = 3/4$, $t = 4/5$, $\eta = \theta$, and $\epsilon = \beta$. We conclude that if β is small enough then there is a mapping H as in Lemma 8.3, with (8.7) and (8.8) corresponding to (5.10) and (5.11), and with the condition (5.9) coming from (8.14).

These choices of f , g , and H satisfy the conclusions of Lemma 8.3. Specifically, (8.4) comes from (8.9), (8.5) from (8.13), (8.6) from (8.14), (8.7) and (8.8) from (5.10) and (5.11). Let us be a little more careful about the way that α and β are chosen. We start with θ , and then we choose β so that Proposition 5.8 works with $\eta = \theta$ and $\epsilon = \beta$. We then choose α so that Proposition 5.4 works with $\eta = \beta/2$ and $\epsilon = \alpha$. Although we have to apply these two propositions in the opposite order, we really have to choose β before α , and we can do this, because of the correct dependence on parameters in Propositions 5.4 and 5.8.

This completes the proof of Lemma 8.3.

Recall that $\sigma \in \mathbf{S}^n$, B , and $\delta > 0$ are as in Theorem 1.29 (a). Set $\theta = \min(\delta/10, 10^{-2})$, and let P , f , g , and H be as in Lemma 8.3 and fixed for the rest of the section. We want to build a mapping from M into $\mathbf{S}^n$ which takes only the value σ outside B , which has nonzero degree, and which has controlled Lipschitz norm. The idea is to build this mapping out of f , since we already control the Lipschitz norm of f . To make this work we need to know that f does not collapse too much of the topological structure of M , and this is made precise by the properties of g and H in Lemma 8.3.

We shall start with a continuous mapping into the sphere with nonzero degree but no Lipschitz bound and make a series of modifications to it. (A slightly different approach, in which one works with homology more directly, is described in Remark 8.26.)

Lemma 8.15. *Let P_{n-1} denote the union of the distinguished simplices in P of dimension $\leq n-1$. There is a continuous mapping $\phi : P \rightarrow \mathbf{S}^n$ which satisfies the following properties: (i) ϕ takes only the value σ on a neighborhood of $P_{n-1} \cup f(U(1/2) \setminus \frac{1}{2}B)$; (ii) if we define $\psi : M \rightarrow \mathbf{S}^n$ by the recipe*

$$(8.16) \quad \begin{aligned} \psi &= \phi \circ f && \text{on } U(1/2) \\ &= \sigma && \text{on } M \setminus U(1/2). \end{aligned}$$

then ψ is continuous and has nonzero degree.

Remember from our hypotheses in Theorem 1.29 (a) that $B \subseteq U$.

We shall build this mapping in stages, beginning with just some continuous mapping on M with nonzero degree.

Let $v_0 : M \rightarrow \mathbf{S}^n$ be a continuous mapping which takes only the value σ on $M \setminus B_0$, $B_0 = \frac{1}{10}B$, and which has nonzero degree. Such a mapping exists, as in Lemma A.27, because M is an oriented topological manifold of dimension n .

Define $\psi_1 : M \rightarrow \mathbf{S}^n$ by $\psi_1 = v_0 \circ g \circ f$ on $U(1/2)$, $\psi_1 \equiv \sigma$ on $M \setminus U(1/2)$. Then ψ_1 takes only the value σ on $M \setminus B_1$, $B_1 = \frac{2}{10}B$, because of (8.6) and the fact that $\psi_0 \equiv \sigma$ on $M \setminus B_0$. (Remember also that $\theta \leq \delta/10$.) In particular ψ_1 is continuous on M , even across the boundary of $U(1/2)$. Let us check that ψ_1 has nonzero degree, using the fact that the degree is preserved by suitable homotopies (Lemma A.21).

Define $\Psi : M \times [0, 1] \rightarrow \mathbf{S}^n$ by $\Psi(x, t) = v_0(H(x, t))$ when $x \in U(1/2)$ and $\Psi(x, t) = \sigma$ when $x \in M \setminus U(1/2)$. We have in fact that $\Psi(x, t) = \sigma$ when $x \in M \setminus B_1$, since $H(x, t) \in M \setminus B_0$ when $x \in M \setminus B_1$, by (8.8). This implies also that Ψ is continuous across $\partial U(1/2)$. From (8.7) we get that $\Psi(x, 0) = v_0(x)$ and $\Psi(x, 1) = \psi_1(x)$ for all $x \in M$. Thus Ψ defines a homotopy between v_0 and ψ_1 which takes only the value σ off a compact subset of M , and we conclude that v_0 and ψ_1 have the same degree (Lemma A.21). Therefore ψ_1 has nonzero degree.

Our first approximation to ϕ is the mapping $\phi_1 : P \rightarrow \mathbf{S}^n$ defined by $\phi_1 = \psi_0 \circ g$. This continuous mapping takes only the value σ on $P \setminus g^{-1}(B_0)$. Let K denote the closure of $g^{-1}(B_0)$. We have that $K \cap f(U(1/2) \setminus \frac{1}{2}B) = \emptyset$, because of (8.6) and our choice of θ . Note that K and $f(U(1/2) \setminus \frac{1}{2}B)$ are both compact subsets of P . We need to modify ϕ_1 to get a mapping which takes only the value σ on P_{n-1} .

Given any $\epsilon > 0$ we can apply Lemma 2.12 to get a Lipschitz mapping $\phi_2 : P \rightarrow \mathbf{S}^n$ which takes only the value σ on $P \setminus K_\epsilon$ and which is homotopic to ϕ_1 through mappings which take only the value σ on $P \setminus K_\epsilon$. We do not control the Lipschitz norm of ϕ_2 here, but we do not mind, we care only that it is finite. We choose ϵ so small, once and for all, so that K_ϵ is disjoint from a neighborhood of $f(U(1/2) \setminus \frac{1}{2}B)$. Thus ϕ_2 takes only the value σ on a neighborhood of $f(U(1/2) \setminus \frac{1}{2}B)$, and ϕ_2 is homotopic to ϕ_1 through mappings which take only the value σ on a neighborhood of $f(U(1/2) \setminus \frac{1}{2}B)$. This mapping ϕ_2 is our second approximation to ϕ .

Because P_{n-1} is a finite union of simplices of dimension $\leq n-1$ and ϕ_2 is Lipschitz we conclude that $\phi_2(P_{n-1})$ has n -dimensional measure 0 in $\mathbf{S}^n$. Hence there is a point $\xi \neq \sigma$ in the complement of $Z = \phi_2(P_{n-1})$ in $\mathbf{S}^n$. Note that Z is a compact subset of $\mathbf{S}^n$, so that ξ lies in the interior of $\mathbf{S}^n \setminus Z$.

Let u be a map on $\mathbf{S}^n$ which takes the value σ on a neighborhood of $\sigma \cup Z$ and which is homotopic to the identity through maps which send σ to itself. It is easy to construct such a mapping. (Lemma A.26 addresses a similar point.)

Set $\phi = u \circ \phi_2$. Thus ϕ is continuous. The definition of u ensures that ϕ takes only the value σ on a neighborhood of P_{n-1} . We also have that ϕ takes only the value σ on a neighborhood of $f(U(1/2) \setminus \frac{1}{2}B)$, because we know that this is true for ϕ_2 .

If we define v as in (8.16) then v is continuous because ϕ takes only the value σ on $f(U(1/2) \setminus \frac{1}{2}B)$. It remains to show that v has nonzero degree.

Define $\psi_2 : M \rightarrow \mathbf{S}^n$ by $\psi_2 = \phi_2 \circ f$ on $U(1/2)$ and $\psi_2 \equiv \sigma$ on $M \setminus U(1/2)$. This mapping is continuous, and it is homotopic to ψ_1 through continuous mappings from M into $\mathbf{S}^n$ which take only the value σ on $M \setminus \frac{1}{2}B$. The existence of this homotopy follows from the homotopy between ϕ_1 and ϕ_2 through mappings which take only the value σ on a

neighborhood of $f(U(1/2) \setminus \frac{1}{2}B)$ which was obtained above. Thus ψ_2 has the same degree as ψ_1 , and it is nonzero in particular. We have that $\psi = u \circ \psi_2$, by definitions, and this permits us to conclude that ψ has the same nonzero degree. (See Lemmas A.20 and A.21.)

This completes the proof of Lemma 8.15.

In order to finish the proof of Theorem 1.29 (a) we need to replace the mapping ψ from Lemma 8.15 with one which has a good Lipschitz bound. We shall do this by expressing ψ in terms of building blocks for which we have better estimates.

Let $\mathcal{Q}$ denote the (finite) collection of n -dimensional distinguished simplices Q in P which intersect $f(\frac{1}{2}B)$. Note that $\mathcal{Q}$ has only a bounded number of elements, because P can be one of only finitely many finite polyhedra, as in Lemma 8.3. We want to build some auxiliary mappings from these simplices into $\mathbf{S}^n$, in such a way that their boundaries get mapped to σ , but we need to choose these mappings carefully.

For each $Q \in \mathcal{Q}$ and $0 < \epsilon < 1$ let $Q_{1-\epsilon}$ denote the m -dimensional (closed) simplex contained in Q which has the same center but which has been shrunk by a factor of $1 - \epsilon$. For each $\epsilon \leq \frac{1}{2}$ let H'_Q be a mapping from Q to $\mathbf{S}^n$ which sends $Q \setminus Q_{1-\epsilon}$ to σ and which maps $Q_{1-\epsilon} \setminus \partial Q_{1-\epsilon}$ homeomorphically onto $\mathbf{S}^n \setminus \{\sigma\}$. Such mappings exist and can be constructed explicitly, and we can do this in such a way that each H'_Q has Lipschitz norm $\leq C$, where C does not depend on ϵ . (C does depend on the finite collection of finite polyhedra mentioned in Lemma 8.3.) Set $H_Q = H'_Q$, where ϵ is chosen small enough so that the restriction of ϕ to Q can be written as $\Phi_Q \circ H_Q$ for some continuous mapping $\Phi_Q : \mathbf{S}^n \rightarrow \mathbf{S}^n$ which takes σ to itself, where ϕ is as in Lemma 8.15. This is possible because ϕ takes the value σ on a neighborhood of P_{n-1} . We extend each H_Q to all of P by setting it to be equal to σ on the complement of Q . These extensions also have uniformly bounded Lipschitz norms, because P ranges through a finite collection of finite polyhedra.

Lemma 8.17. *Define $J_Q : M \rightarrow \mathbf{S}^n$ for each $Q \in \mathcal{Q}$ by*

$$(8.18) \quad \begin{aligned} J_Q &= H_Q \circ f && \text{on } U(1/2) \\ &= \sigma && \text{on } M \setminus U(1/2). \end{aligned}$$

Then each J_Q takes only the value σ on $M \setminus B$, is Lipschitz with bounded norm (with the bound depending on the same parameters as in Lemma 8.3), and there is at least one $Q \in \mathcal{Q}$ such that J_Q has nonzero degree.

Let us begin with some auxiliary notation. For each $Q \in \mathcal{Q}$ set $V(Q) = f^{-1}(Q \setminus \partial Q)$. Let us check that

$$(8.19) \quad V(Q) \subseteq B \quad \text{for all } Q \in \mathcal{Q}.$$

Indeed, the definition of $\mathcal{Q}$ ensures that Q intersects $f(\frac{1}{2}B)$ when $Q \in \mathcal{Q}$, and so (8.19) follows from (8.4) and our choice of θ .

A useful consequence of (8.19) is that $V(Q)$ is an open subset of M . It is automatically a relatively open subset of $U(1/2)$, since $f : U(1/2) \rightarrow P$ is continuous, but it is also open in M because (8.19) ensures that $V(Q)$ stays away from $\partial U(1/2)$. (Remember that $B \subseteq U$, by our hypotheses from Theorem 1.29 (a).)

Another consequence of (8.19) is that each J_Q , $Q \in \mathcal{Q}$, takes only the value σ on $M \setminus B$. This follows from the definitions, which imply that

$$(8.20) \quad J_Q \equiv \sigma \quad \text{on } M \setminus V(Q).$$

Because $B \subseteq U$ we have that J_Q is continuous (the only possible danger being near $\partial U(1/2)$). It is easy to see that J_Q is also Lipschitz with bounded norm, because of the corresponding bounds for H_Q and f . (See Lemma 8.3.)

It remains to show that some J_Q has nonzero degree. We shall derive this from the corresponding property of the mapping ψ in Lemma 8.15 using some properties of the degree discussed in Appendix A. More precisely we shall decompose ψ into pieces associated to the elements of $\mathcal{Q}$.

Set $V = \bigcup_{Q \in \mathcal{Q}} V(Q)$. Thus V is an open subset of M .

Sublemma 8.21. *ψ and each J_Q , $Q \in \mathcal{Q}$, take only the value σ on $M \setminus V$. For each $T \in \mathcal{Q}$ there is a compact subset $V_0(T)$ of $V(T)$ such that ψ and each J_Q , $Q \in \mathcal{Q}$, take only the value σ on $V(T) \setminus V_0(T)$.*

Notice first that if W is a neighborhood of ∂Q , $Q \in \mathcal{Q}$, then $f^{-1}(Q \setminus W)$ is a compact subset of M , since the domain $U(1/2)$ of f is compact. For our purposes it is better to say that $f^{-1}(Q \setminus W)$ is a compact subset of the open set $V(Q)$.

The definitions of J_Q and H_Q imply that $J_Q \equiv \sigma$ off a compact subset of $V(Q)$. This comes down to the fact that $H_Q \equiv \sigma$ on $P \setminus Q$ and also on a neighborhood of ∂Q . Thus the conclusions of the sublemma hold for the J_Q 's, and so we need only concern ourselves with ψ .

We know from the definition (8.16) of ψ that it takes only the value σ on $M \setminus U(1/2)$ and that $\psi = \phi \circ f$ on $U(1/2)$. Note that $f(\frac{1}{2}B)$ is covered by the Q 's in $\mathcal{Q}$, by definition of $\mathcal{Q}$. It is not difficult to check that $f(U(1/2) \setminus V) \subseteq f(U(1/2) \setminus \frac{1}{2}B) \cup P_{n-1}$, using the definitions of V and $V(Q)$. This implies that $\psi \equiv \sigma$ on $U(1/2) \setminus V$, because of Lemma 8.15 (i). Therefore $\psi \equiv \sigma$ on $M \setminus V$.

Fix $T \in \mathcal{Q}$. From Lemma 8.15 we know that $\phi \equiv \sigma$ on a neighborhood of ∂T , because $\partial T \subseteq P_{n-1}$. This implies that the restriction of ψ to $V(T)$ takes only the value σ on the complement of a compact subset of $V(T)$ (using also (8.16) and (8.19)).

This completes the proof of Sublemma 8.21.

Set $V_0 = \bigcup_{Q \in \mathcal{Q}} V_0(Q)$, where $V_0(Q)$ is as in Sublemma 8.21. Thus V_0 is a compact subset of V , and ψ and each J_Q , $Q \in \mathcal{Q}$, take only the value σ on $M \setminus V_0$. This implies that the degree of the restriction of ψ to V is well-defined and equals the degree of ψ as a mapping defined on all of M . (See Proposition A.24.)

Let ψ^Q denote the restriction of ψ to $V(Q)$, $Q \in \mathcal{Q}$. Thus ψ^Q takes only the value σ on the complement of $V_0(Q)$, and so its degree is well-defined. Since $V = \bigcup_{Q \in \mathcal{Q}} V(Q)$ we get that

$$(8.22) \quad \text{degree } \psi = \sum_{Q \in \mathcal{Q}} \text{degree } \psi^Q.$$

(See Lemma A.25.) Since the degree of ψ is nonzero, we conclude that there is at least one $Q \in \mathcal{Q}$ such that ψ^Q has nonzero degree. Fix such a $Q \in \mathcal{Q}$.

Let J'_Q denote the restriction of J_Q to $V(Q)$.

Sublemma 8.23. *degree $\psi^Q = \text{degree } \Phi_Q \cdot \text{degree } J'_Q$, where $\Phi_Q : S^n \rightarrow S^n$ is as defined just before Lemma 8.17.*

Indeed, the definition of ψ^Q , Φ_Q , and J_Q imply that $\psi^Q = \Phi_Q \circ J'_Q$. Thus Sublemma 8.23 is just the usual composition rule for degrees (Lemma A.20).

Sublemma 8.23 implies that the degree of J'_Q is nonzero, since we chose Q so that ψ^Q has nonzero degree.

As a mapping on all of M , $J_Q \equiv \sigma$ off the compact subset $V_0(Q)$ of $V(Q)$. Thus Proposition A.24 implies that J_Q and J'_Q have the same degree. Therefore the degree of J_Q is nonzero, which is what we wanted. This completes the proof of Lemma 8.17.

The proof of Theorem 1.29 (a) is now complete as well. We just take a Q for which J_Q has nonzero degree and use that for our mapping.

Remark 8.24. As indicated in the introduction, Theorem 1.29 (a) remains true if we do not ask for a controlled covering function $N(\cdot)$, but we ask instead for a controlled topological dimension function $\delta_n(\cdot)$. To see this we simply go through the various steps of the preceding arguments. This change in hypothesis amounts to making the conclusions of Lemma 8.1 part of the assumptions, so that is fine. Lemma 8.2 continues to hold if we replace our assumption of a fixed covering function with the requirement that the topological dimension function be controlled, except that we lose the conclusion that our polyhedron lies in a finite collection of polyhedra, or even that it lies in a Euclidean space of bounded dimension. This variant of Lemma 8.2 follows from Remark 7.5. Note that, as in Remark 7.5, we do retain some control on the geometry of the polyhedron P , namely that it lies in some Euclidean space and that the vertices of all of its distinguished simplices are all standard basis vectors for the ambient Euclidean space. Thus the distinguished simplices in P are not permitted to degenerate, even if we do not control their number. Lemma 8.3 also admits a similar variation, that is, we may trade a bound on the covering function for a bound on the topological dimension function at the expense of losing control on the number of distinguished simplices in the polyhedron P (but not on the geometry of these simplices). The main point behind this assertion is that Propositions 5.4 and 5.8 did not require bounds on the covering function, just on the topological dimension and the contractability function. Next, Lemma 8.15 is undisturbed by our change in hypotheses, because it only needs P to be a finite polyhedron of dimension $\leq n$. Note that there are no real bounds in the conclusion of Lemma 8.15, except for the control on the set where ϕ is permitted to take a value different from σ .

In the remainder of the proof of Theorem 1.29 (a), beginning just after the proof of Lemma 8.15, we said that we were using the fact that P ranges through a finite collection of finite polyhedra, but in fact it is enough to have the control on the geometry of P that we have here. All we really need is that the mappings H_Q have bounded Lipschitz norm, and this we can certainly get. When we define the H_Q 's first on the Q 's we get a uniform Lipschitz bound simply because all the Q 's look like the standard n -dimensional simplex. When we extend the H_Q 's to all of P as above we have to be careful not to destroy our uniform Lipschitz bound, and this depends also on the way that the Q 's sit inside P , i.e., they never meet at a small angle. To be precise, if τ and τ' are two distinct n -dimensional

distinguished simplices in P , and if $x \in \tau$, then

$$(8.25) \quad \text{dist}(x, \tau') \geq \text{dist}(x, \partial\tau').$$

As always we are using here the Euclidean metric from the ambient Euclidean space in which P is constructed. This inequality is very easy to verify, using only the fact that τ and τ' are convex hulls of basis vectors of the ambient Euclidean space. It is easy to check that this inequality does indeed imply a uniform Lipschitz condition for the extended H_Q 's (since H_Q is defined so that it takes only the value σ on ∂Q and on $P \setminus Q$). Once we have this the rest of the argument is the same, with no further mention of the complexity of P . The main point is to get that one of the J_Q 's has nonzero degree, for which the complexity of P is irrelevant. The bound on the Lipschitz norm of J_Q then follows from the corresponding bounds for f and H_Q .

This completes the discussion of the modification of the proof of the variant of Theorem 1.29 (a) in which the bound of the covering function is traded for a bound on the topological dimension function.

Remark 8.26. The starting point of the preceding construction was the mapping $\psi_0 : M \rightarrow \mathbf{S}^n$ introduced at the beginning of the proof of Lemma 8.15. This map had the properties of being continuous, taking only the value σ on the complement of $B_0 = \frac{1}{10}B$, and having nonzero degree, and our construction replaced it with a mapping which enjoyed a Lipschitz bound. There is a nice coherence to this argument, starting with a mapping and replacing it with a better one, but there is a minor variant of this approach which has some advantages. In this variant one would forget about ψ_0 and work directly with the n -dimensional relative homology classes on M associated to its orientation, as in Theorem A.17. One would use Lemma 8.3 to show that the Lipschitz mapping $f : U(1/2) \rightarrow P$ given there is nondegenerate in the sense that it induces injections between suitable n -dimensional relative homology groups. One could then prove Lemma 8.17 by analyzing the relevant relative homology groups of the polyhedron P in a standard way. (It is a CW-complex, so that one can use the injectivity of the homomorphism j_n in Theorem 4.2 on p.233 of [Ma].)

This second approach is not enormously different from the other one, but it is better for reducing our reliance on bounds on local contractability, as discussed in Remark 6.10. More precisely, the polyhedron P and the mapping f in Lemma 8.3 came from Lemma 8.2, and they required the bound on the local contractability function only through the bound on the topological dimension function in Lemma 8.1. The contractability function entered into that bound through Theorem 6.5, but there are other reasonable ways in which one can bound the topological dimension function, as discussed in Remark 6.10. By contrast, the approximate inverse g to f given in Lemma 8.3 and the homotopy H given there were obtained through Propositions 5.4 and 5.8 and used the contractability function in a more direct and brutal way. The nice feature of this second approach to the construction above is that it makes it more clear that we do not really need an approximate inverse to f as a mapping, we simply need to know that it is nondegenerate on n -dimensional homology.

9. Bounds for the degrees of Lipschitz mappings.

In this section we prove Theorem 1.29 (b). This means that we want to show that the degree of a Lipschitz mapping from a certain space into a sphere can be bounded in terms of the Lipschitz norm (and other quantities related to the underlying space).

Fix a dimension n , a covering function $N(\cdot)$, and a contractability function $\rho(\cdot)$ (as in Definitions 1.6 and 1.7). Let $\mathcal{F}$ denote the family of triples $(M, U, d(x, y))$ which satisfy Standard Assumptions 1.8 with these choices of $N(\cdot)$ and $\rho(\cdot)$ and with the additional normalization $\text{diam } U = 1$.

Fix $\sigma \in \mathcal{S}^n$ and $k \geq 1$. Let $\mathcal{G}$ denote the family of quadruples $(M, U, d(x, y), g)$ where $(M, U, d(x, y)) \in \mathcal{F}$ and $g : M \rightarrow \mathcal{S}^n$ takes only the value σ on $M \setminus U$ and is k -Lipschitz. Theorem 1.29 (b) is equivalent to saying that if $(M, U, d(x, y), g) \in \mathcal{G}$, then the degree of g is uniformly bounded. (The normalization $\text{diam } U = 1$ causes no trouble here, because one can always rescale the metric to get it.)

Let us reformulate this assertion in a more convenient way.

Theorem 9.1. *Notations and assumptions as above. There is a finite subset $\mathcal{G}_0$ of $\mathcal{G}$ with the property that if $(M, U, d(x, y), g) \in \mathcal{G}$, then there is a $(M', U', d'(x, y), g') \in \mathcal{G}_0$ such that g and g' either have the same degree or their degrees are negatives of each other.*

Theorem 1.29(b) will follow once we prove this.

To prove Theorem 9.1 we shall basically use compactness, although the argument is not truly nonconstructive. We are going to show that every element of $\mathcal{G}$ is “close” (in a technical sense) to an element of a finite subset $\mathcal{G}'$ of $\mathcal{G}$ (Lemma 9.7), and then we are going to show two elements of $\mathcal{G}$ which are close in this way have mappings into $\mathcal{S}^n$ with either the same degrees or degree of opposite sign (Proposition 9.15).

If $(M, U, d(x, y), g) \in \mathcal{G}$, then we let U_t be as in Notation 5.3, and we write $U(t)$ for $\overline{U}_t$.

Definition 9.2. Let $\epsilon > 0$ be given. We say that $(M, U, d(x, y), g), (M', U', d'(x, y), g') \in \mathcal{G}$ are ϵ -close if there exist mappings $f : U(1) \rightarrow U'(1)$, $f' : U'(1) \rightarrow U(1)$ with the following properties:

$$(9.3) \quad f(U) \subseteq U', f'(U') \subseteq U;$$

$$(9.4) \quad d(f' \circ f(x), x) < \epsilon \quad \text{and} \quad d'(f \circ f'(x'), x') < \epsilon \quad \text{for all } x \in U(1), x' \in U'(1);$$

$$(9.5) \quad |d'(f(x), f(y)) - d(x, y)| < \epsilon \quad \text{and} \quad |d(f'(x'), f'(y')) - d'(x', y')| < \epsilon \quad \text{for all } x, y \in U(1), x', y' \in U'(1);$$

$$(9.6) \quad |g' \circ f(x) - g(x)| < \epsilon \quad \text{and} \quad |g \circ f'(x') - g'(x')| < \epsilon \quad \text{for all } x \in U(1), x' \in U'(1).$$

These mappings are not required to be continuous.

In other words, f and f' are almost isometries which are almost inverses of each other and under which g and g' almost correspond.

Lemma 9.7. *For each $\epsilon > 0$ there is a finite subset $\mathcal{G}_\epsilon$ of $\mathcal{G}$ such that every element $(M, U, d(x, y), g)$ of $\mathcal{G}$ is ϵ -close to some $(M', U', d'(x, y), g') \in \mathcal{G}_\epsilon$.*

To prove this we use a straightforward compactness argument. We first choose some discrete models for the elements of $\mathcal{G}$. Set $\delta = 10^{-2}k^{-1}\epsilon$. Given $(M, U, d(x, y), g) \in \mathcal{G}$ let $x_1, \dots, x_j$ be a maximal set of points in U such that

$$(9.8) \quad d(x_a, x_b) > \delta \text{ when } a \neq b.$$

Let $x_1, \dots, x_l$, $l \geq j$, be an extension of $x_1, \dots, x_j$ to a maximal subset of $U(1)$ such that (9.8) holds. Remember that $U(1)$ is compact, and so there is no problem finding finite sets with these properties.

Since $N(\cdot)$ is a covering function for $(M, U, d(x, y))$ (Definition 1.6) we have that $l \leq N(\delta/2)$. Also,

$$(9.9) \quad U \subseteq \bigcup_{i=1}^j \overline{B}_M(x_i, \delta) \quad \text{and} \quad U(1) \subseteq \bigcup_{i=1}^l \overline{B}_M(x_i, \delta).$$

by maximality.

Our discrete model for $(M, U, d(x, y), g)$ is given by the data

$$(9.10) \quad (j, l, \{x_i\}_{i=1}^l, \{d(x_a, x_b) : 1 \leq a, b \leq l\}, \{g(x_a) : 1 \leq a \leq l\}).$$

Note that the x_a 's live in M and the $g(x_a)$'s live in $\mathbf{S}^n$. We choose such a discrete model for each $(M, U, d(x, y), g) \in \mathcal{G}$. We do not care how the specific model is chosen, just that one is associated to each element of $\mathcal{G}$ in this manner.

Let us say that $(M, U, d(x, y), g), (M', U', d'(x, y), g') \in \mathcal{G}$ are *discretely close* if their discrete models

$$(9.11) \quad (j, l, \{x_i\}_{i=1}^l, \{d(x_a, x_b) : 1 \leq a, b \leq l\}, \{g(x_a) : 1 \leq a \leq l\}) \quad \text{and} \\ (j', l', \{x'_i\}_{i=1}^{l'}, \{d'(x'_a, x'_b) : 1 \leq a, b \leq l'\}, \{g'(x'_a) : 1 \leq a \leq l'\})$$

satisfy $j' = j, l' = l$,

$$(9.12) \quad |d'(x'_a, x'_b) - d(x_a, x_b)| < \delta, \quad 1 \leq a, b \leq l, \text{ and}$$

$$(9.13) \quad |g(x_a) - g'(x'_a)| < \delta, \quad 1 \leq a \leq l.$$

Sublemma 9.14. *There is a finite subset $\mathcal{G}_\epsilon$ of $\mathcal{G}$ such that every element of $\mathcal{G}$ is discretely close to an element of $\mathcal{G}_\epsilon$.*

This follows from an easy compactness argument. Keep in mind that we have a uniform bound on the l 's and the numbers $d(x_a, x_b)$ in our discrete model, and of course we are also using the compactness of $\mathbf{S}^n$.

Lemma 9.7 is an easy consequence of Sublemma 9.14. Let $\mathcal{G}_\epsilon$ be as in Sublemma 9.14, let $(M, U, d(x, y), g) \in \mathcal{G}$ be given, and choose $(M', U', d'(x, y), g') \in \mathcal{G}_\epsilon$ so that it is discretely close to $(M, U, d(x, y), g)$, with the discrete models given as in (9.11). We need to prove that $(M, U, d(x, y), g)$ and $(M', U', d'(x, y), g')$ are ϵ -close, as in Definition 9.2. We define maps $f : U(1) \rightarrow U'(1)$, $f' : U'(1) \rightarrow U(1)$ as follows. First we set $f(x_i) = x'_i$ and

$f'(x'_i) = x_i$ for all $1 \leq i \leq l$. Next, if $x \in U$, then we use (9.9) to find an $i, 1 \leq i \leq j$, such that $x \in \overline{B}_M(x_i, \delta)$, and we set $f(x) = x'_i$. This choice of i need not be unique, but we do not mind, we simply take one. (Remember that f need not be continuous.) We define f' on U' in the same way. Next we define f on $U(1) \setminus U$. Given $x \in U(1) \setminus U$ we choose an $i, 1 \leq i \leq l$, so that $x \in \overline{B}_M(x_i, \delta)$ (as in (9.9)), and we set $f(x) = x'_i$. Again this choice need not be unique but we do not mind. We extend f' to $U'(1)$ in the same manner. It is not hard to check that (9.3) - (9.6) hold, using (9.9), (9.12), (9.13), our choice of δ , and the fact that g and g' are both Lipschitz with norm $\leq k$. This proves Lemma 9.7.

Proposition 9.15. *If $(M, U, d(x, y), g), (M', U', d'(x, y), g') \in \mathcal{G}$ are ϵ -close for a sufficiently small ϵ (which does not depend on $(M, U, d(x, y), g)$ or $(M', U', d'(x, y), g')$), then g and g' either have the same degree or their degrees are negatives of each other.*

Theorem 9.1 will follow from Lemma 9.7 once we prove this.

Let $\epsilon > 0$ be small, to be chosen later. We may as well assume that $\epsilon < 10^{-3}$, for instance. Let $(M, U, d(x, y), g), (M', U', d'(x, y), g') \in \mathcal{G}$ be given, and assume that they are ϵ -close. Let $f : U(1) \rightarrow U'(1), f' : U'(1) \rightarrow U(1)$ be as in Definition 9.2. We shall use Proposition 5.4 to modify these mappings to get continuous ones, and then we shall use these continuous mappings and Proposition 5.8 to relate g and g' .

We shall need to use the sets $U(t)$ and $U'(t)$ for various values of t between 0 and 1, and we shall be choosing values of t like .5 and .6. Notice that

$$(9.16) \quad f(U(t)) \subseteq U'(t + \epsilon) \quad \text{and} \quad f'(U'(t)) \subseteq U(t + \epsilon)$$

when $0 \leq t \leq 1$, by (9.5), (9.3), and the definition of $U(t), U'(t)$.

Lemma 9.17. *If $\eta_1 > 0$ is given and ϵ is small enough, depending on η_1 , then we can find continuous mappings $\phi : U(.5) \rightarrow U'(.6)$ and $\phi' : U'(.5) \rightarrow U(.6)$ such that*

$$(9.18) \quad d_{M'}(\phi(x), f(x)) < \eta_1 \quad \text{for all } x \in U(.5), \text{ and}$$

$$(9.19) \quad d_M(\phi'(x'), f'(x')) < \eta_1 \quad \text{for all } x' \in U'(.5).$$

To see this we note that f and f' are 2ϵ -continuous, in the sense of Definition 5.2, because of (9.5). The mappings ϕ, ϕ' are obtained from f, f' as in Proposition 5.4, modulo many changes in notation. More precisely, to get ϕ we apply Proposition 5.4 with $X = U(.5)$, $k = n$, M replaced with M' , $Z = U$, $s = .51$, and $t = .6$. We can do this because (9.16) ensures that f maps X into Z_s with these choices if ϵ is small enough. (It is for this reason that we need to take $s = .51 > .5$.) We take ϕ to be the mapping provided by Proposition 5.4, which was called g there. The mapping ϕ' is obtained in the same way. This proves Lemma 9.17.

From now on we let ϕ and ϕ' be as in Lemma 9.17, with η_1 small and to be specified. We require that η_1 at least be $< 10^{-3}$, say.

Define $h : M \rightarrow S^n$ by

$$(9.20) \quad \begin{aligned} h(x) &= g'(\phi(x)) && \text{when } x \in U(.5), \\ &= \sigma && \text{when } x \in M \setminus U(.5). \end{aligned}$$

This mapping will provide a bridge for us between g and g' .

Lemma 9.21. *If $k\eta_1 + \epsilon < 10^{-3}$ then h is continuous and has the same degree as g .*

Recall that k is an upper bound for the Lipschitz norms of the g 's. (See the definition of $\mathcal{G}$ at the beginning of this section.)

To see that h is continuous it is sufficient to know that

$$(9.22) \quad h(x) = \sigma \quad \text{when } x \in M \setminus U(.2).$$

The definition of $\mathcal{G}$ ensures that $g' \equiv \sigma$ on $M' \setminus U'$, and so (9.22) reduces to

$$(9.23) \quad \phi(U(.5) \setminus U(.2)) \subseteq M' \setminus U'.$$

To prove this inclusion it is enough to show that $f(U(.1) \setminus U(.2)) \subseteq M' \setminus U'(.1)$, because of (9.18) (and $\eta_1 < 10^{-3}$). To get this inclusion for f we use (9.16) to get $f'(U'(.1)) \subseteq U(.11)$, since $\epsilon < 10^{-3}$. This implies that $f(U(.1) \setminus U(.2)) \subseteq M' \setminus U'(.1)$, because $f' \circ f$ is almost the identity, in the sense of (9.4) (and using $\epsilon < 10^{-3}$ again). Thus we get (9.23), and hence (9.22) and the continuity of h .

Next we check that

$$(9.24) \quad |h(x) - g(x)| < k\eta_1 + \epsilon \quad \text{when } x \in M.$$

When $x \in U(.5)$ we have that

$$(9.25) \quad |h(x) - g(x)| = |g'(\phi(x)) - g(x)| \leq |g'(\phi(x)) - g'(f(x))| + |g'(f(x)) - g(x)|.$$

Thus (9.24) follows from our Lipschitz bound on g' , (9.18), and (9.6) when $x \in U(.5)$. When $x \in M \setminus U(.5)$ we have that $h(x) = g(x) = \sigma$, and (9.24) is trivial.

Our hypothesis in Lemma 9.21 yields

$$(9.26) \quad |h - g| < 10^{-3} \quad \text{on } M.$$

This implies that g and h are homotopic. Specifically, define $H : M \times [0, 1] \rightarrow \mathbf{S}^n$ by

$$(9.27) \quad H(x, t) = \frac{th(x) + (1-t)g(x)}{|th(x) + (1-t)g(x)|}.$$

This mapping is continuous, because (9.26) ensures that the denominator never vanishes. We have that $H(x, 0) = g(x)$ and $H(x, 1) = h(x)$ for all $x \in M$, and $H(x, t) = \sigma$ for all $x \in M \setminus U(.2)$ and $t \in [0, 1]$, because $h(x) = g(x) = \sigma$ for the same choices of x . Thus H provides a homotopy between g and h which takes only the value σ on $M \setminus U(.2)$, and this implies that g and h have the same degree. (See Lemma A.21.)

This proves Lemma 9.21.

Lemma 9.28. *If ϵ and η_1 are small enough, then the degrees of h and g' are either equal or negatives of each other.*

Proposition 9.15 will follow once we prove this.

The idea behind this lemma is that the mappings ϕ and ϕ' are almost inverses of each other, so that they are almost like homeomorphisms, or at least like homotopy equivalences. We are going to need to show that the fundamental classes of M and M' – the top-dimensional homology classes determined by the orientations of M and M' – correspond to each other under ϕ and ϕ' , except possibly for a change in sign. We have to be a little careful, because the mappings and homology classes with which we shall be working are not defined on all of M and M' . We shall have to be careful about which sets are mapped where by ϕ and ϕ' . It will be important for us to work with connected subsets of M and M' , to ensure that the relevant homology groups are infinite cyclic. (See Lemma A.28.) It is for this reason that we need to assume that U and U' are connected, as in Standard Assumptions 1.8. We like infinite cyclic groups because their generators are uniquely determined up to sign, and it is this uniqueness which will give Lemma 9.28.

Notice that

$$(9.29) \quad \phi(U(t)) \subseteq U'(t + \epsilon + \eta_1) \quad \text{and} \quad \phi'(U'(t)) \subseteq U(t + \epsilon + \eta_1)$$

when $0 \leq t \leq .5$, by (9.16), (9.18), and (9.19). Assume that $\eta_1 < 10^{-3}$. This permits us to define $\psi : U(.4) \rightarrow U(.6)$ and $\psi' : U'(.4) \rightarrow U'(.6)$ by

$$(9.30) \quad \psi = \phi' \circ \phi \quad \text{and} \quad \psi' = \phi \circ \phi'.$$

These mappings ψ, ψ' are close to the identity, in the sense that

$$(9.31) \quad d_M(\psi(x), x) < 2\epsilon + 2\eta_1 \quad \text{for all } x \in U(.4),$$

$$(9.32) \quad d_{M'}(\psi'(x'), x') < 2\epsilon + 2\eta_1 \quad \text{for all } x' \in U'(.4).$$

These estimates follow from (9.18), (9.19), (9.4), and (9.5).

Next we homotop ψ and ψ' to the identity in a controlled way. This will provide another sense in which ϕ and ϕ' are approximately inverses of each other, and it will be used to analyze the induced maps on homology.

Sublemma 9.33. *Let $\eta_2 > 0$ be given. If $\eta_1 + \epsilon$ is sufficiently small, then there exist continuous mappings $\Psi : U(.4) \times [0, 1] \rightarrow M$ and $\Psi' : U'(.4) \times [0, 1] \rightarrow M'$ such that*

$$(9.34) \quad \Psi(x, 0) = x \quad \text{and} \quad \Psi(x, 1) = \psi(x) \quad \text{for all } x \in U(.4),$$

$$(9.35) \quad \Psi'(x', 0) = x' \quad \text{and} \quad \Psi'(x', 1) = \psi'(x') \quad \text{for all } x' \in U'(.4),$$

$$(9.36) \quad d_M(\Psi(x, t), x) \leq \eta_2 \quad \text{for all } x \in U(.4) \text{ and } t \in [0, 1], \text{ and}$$

$$(9.37) \quad d_{M'}(\Psi'(x', t), x') \leq \eta_2 \quad \text{for all } x' \in U'(.4) \text{ and } t \in [0, 1].$$

This follows from Proposition 5.8. Specifically, in order to obtain Ψ we take $X = U(.4)$, $k = n$, $f_0 = \text{the identity}$, $f_1 = \psi$, $Z = U$, $s = .61$, $t = 1$. The ϵ and η in Proposition 5.8 should be replaced by $2\epsilon + 2\eta_1$ and η_2 , respectively. The mapping Ψ corresponds to H in Proposition 5.8, and Ψ' can be obtained in the same way.

We shall use these homotopies to control the effect of ψ and ψ' on homology, but before we do that we need to construct some auxiliary compact *connected* subsets of M and M' . We begin with a preliminary fact.

Sublemma 9.38. *For each $\alpha \in (0, 1)$ (think of α as being small) there is a $\beta = \beta(\alpha) > 0$ and a compact connected set $J \subseteq M$ such that $U(\beta) \subseteq J \subseteq U(\alpha)$. Here β depends only on α and our contractability function $\rho(\cdot)$ from the beginning of this section, and the analogous statement holds for M' instead of M (with the same β).*

Given α as in the sublemma, take J to be the closure of the points in M which can be connected to U by a curve whose diameter is $< \alpha$. Thus $J \subseteq U(\alpha)$, by definition. We also have that J is compact and connected, since U is connected (as in Standard Assumptions 1.8). Our local contractability assumptions (coming from Standard Assumptions 1.8 and Definition 1.7) imply that $U(\beta) \subseteq J$ for a choice of β which depends only on α and $\rho(\cdot)$. This proves Sublemma 9.38.

Sublemma 9.39. *There exist compact connected sets $K \subseteq U(.1)$ and $K', L' \subseteq U'(.1)$ which satisfy the following inclusions, at least if η_2, η_1 , and ϵ are small enough (in a way which can be computed just in terms of $\rho(\cdot)$). (Think of L' as being the largest of $\{K, K', L'\}$, with K the second largest and K' the smallest. This is made precise by (9.46)-(9.48) below, and it is helpful in understanding (9.42)-(9.44).)*

$$(9.40) \quad U(2\epsilon + \eta_1) \subseteq K, \quad U'(0) \subseteq K' \subseteq L'$$

$$(9.41) \quad \phi(U(2\epsilon + \eta_1)) \subseteq K'$$

$$(9.42) \quad \phi'(U'(.5) \setminus L') \subseteq U(.6) \setminus K$$

$$(9.43) \quad \phi(U(.5) \setminus K) \subseteq U'(.6) \setminus K'$$

$$(9.44) \quad \Psi'(\{U'(.4) \setminus L'\} \times [0, 1]) \subseteq M' \setminus K'$$

Let $\beta(\alpha)$ be the function given in Sublemma 9.38, and notice that $\beta(\alpha) \leq \alpha$. Choose $\alpha_1, \alpha_2 > 0$ so that $\alpha_2 < \alpha_1 < .1$.

$$(9.45) \quad \alpha_1 < \beta(.1)/2 \quad \text{and} \quad \alpha_2 < \beta(\alpha_1)/2.$$

(The smallness requirements for η_2, η_1 , and ϵ can be computed just in terms of α_1 and α_2 .) Choose K, K', L' so that they are compact connected sets which satisfy

$$(9.46) \quad U'(\beta(\alpha_2)) \subseteq K' \subseteq U'(\alpha_2),$$

$$(9.47) \quad U(\beta(\alpha_1)) \subseteq K \subseteq U(\alpha_1),$$

$$(9.48) \quad U'(\beta(.1)) \subseteq L' \subseteq U'(.1).$$

We can do this because of Sublemma 9.38, applied to M and M' with α taken to be α_2, α_1 , and $.1$.

If η_2, η_1 , and ϵ are small enough, then (9.40)-(9.44) are true. This is clear for (9.40), using (9.46)-(9.48). The main point for (9.44) is (9.37), but we also use (9.48), (9.46), and

(9.45). This gives (9.44) if η_2 is small enough. To get (9.41) we observe that $\phi(U(2\epsilon + \eta_1)) \subseteq U'(3\epsilon + 2\eta_1)$, because of (9.29). This implies (9.41) if ϵ and η_1 are small enough, because of (9.46). For (9.42) and (9.43) we need another piece of information, namely

$$(9.49) \quad \text{dist}_{M'}(f(x), U') \geq \text{dist}_M(x, U) - 2\epsilon \quad \text{for all } x \in U(1), \text{ and}$$

$$(9.50) \quad \text{dist}_M(f'(x'), U) \geq \text{dist}_{M'}(x', U') - 2\epsilon \quad \text{for all } x' \in U'(1).$$

Let us check (9.49). The main point is that f' is almost an inverse of f , because of (9.4). Suppose that $x \in U(1)$ but that $\text{dist}_{M'}(f(x), U') < \text{dist}_M(x, U) - 2\epsilon$. Notice that

$$(9.51) \quad \text{dist}_M(f'(f(x)), U) \leq \text{dist}_{M'}(f(x), U') + \epsilon,$$

because of (9.5) and (9.3). Thus $\text{dist}_M(f'(f(x)), U) < \text{dist}_M(x, U) - \epsilon$, by our assumption. This inequality implies that $\text{dist}_M(x, U) < \text{dist}_M(x, U)$, because of (9.4). This is impossible, and (9.49) follows. The proof of (9.50) is the same. Once we have (9.49) and (9.50) we can get versions of them for ϕ and ϕ' as well, to wit,

$$(9.52) \quad \text{dist}_{M'}(\phi(x), U') \geq \text{dist}_M(x, U) - 2\epsilon - \eta_1 \quad \text{for all } x \in U(.5), \text{ and}$$

$$(9.53) \quad \text{dist}_M(\phi'(x'), U) \geq \text{dist}_{M'}(x', U') - 2\epsilon - \eta_1 \quad \text{for all } x' \in U'(.5),$$

because of (9.18) and (9.19). These estimates permit us to derive (9.42) and (9.43) from (9.45)-(9.48) when ϵ and η_1 are small enough. This completes the proof of Sublemma 9.39.

From now on we assume that η_2, η_1 , and ϵ are sufficiently small for the preceding results to hold, and we let K, K' , and L' be as in Sublemma 9.39.

(Let us be a little more explicit about the way in which η_2, η_1 , and ϵ are chosen. We first choose η_2 in a way which is small enough for the requirements of Sublemma 9.39. Sublemma 9.33 then imposes a condition on the size of $\eta_1 + \epsilon$, but we can also take them to be smaller still. Next we choose η_1 small enough so that Sublemma 9.39 holds and so that Lemma 9.21 can hold. Lemma 9.17 then forces us to take ϵ small enough depending on this choice of η_1 , but again we can take it smaller still to ensure that the requirements of Lemma 9.21 and Sublemma 9.39 are met. These choices are consistent with each other because our choice of η_2 does not depend on η_1 or ϵ , nor does our choice of η_1 depend on ϵ . This is because the requirements of Lemma 9.12 and Sublemma 9.39 are “absolute” requirements which do not depend on the specific choices of these parameters, only on the parameters of our data.)

We are still working towards proving Lemma 9.28. The point now is to understand the effect of our various maps on homology. We shall use basic definitions and results from homology theory, as summarized in Appendix A.

Recall that M and M' are oriented topological manifolds of dimension n , as in Standard Assumptions 1.8. Let their orientations be denoted by $\{\lambda_p\}_{p \in M}$ and $\{\lambda'_{p'}\}_{p' \in M'}$, in accordance with Definition A.15. Given compact sets $J \subseteq M$ or $J' \subseteq M'$, there are relative homology classes $\lambda_J \in H_n(M, M \setminus J)$ and $\lambda'_{J'} \in H_n(M', M' \setminus J')$ which are determined by the orientations. (See Theorem A.17.)

We care about these relative homology classes because we can use them to compute the degrees of h and g' . Recall that g' takes only the value σ on $M' \setminus U'$. Let us check that

$$(9.54) \quad h \equiv \sigma \quad \text{on } M \setminus K.$$

where h is defined in (9.20). Since g' takes only the value σ on $M' \setminus U'$ we need only know that $\phi(U(.5) \setminus K) \subseteq M' \setminus U'$, and this follows from (9.43). Thus we have (9.54).

Let d denote the degree of h and let d' denote the degree of g' . Following Definition A.19 we express these degrees in terms of homology. Let ω be the standard generator of $H_n(\mathbf{S}^n)$, and let $\tilde{\omega}$ be its image in $H_n(\mathbf{S}^n, \{\sigma\})$. ($\tilde{\omega}$ is called ω' in Definition A.19, but that is not such good notation here.) We have that

$$(9.55) \quad h_*(\lambda_K) = d \cdot \tilde{\omega} \quad \text{and} \quad g'_*(\lambda'_{K'}) = d' \cdot \tilde{\omega},$$

where $h_* : H_n(M, M \setminus K) \rightarrow H_n(\mathbf{S}^n, \{\sigma\})$ and $g'_* : H_n(M', M' \setminus K') \rightarrow H_n(\mathbf{S}^n, \{\sigma\})$ are the mappings induced on homology by h and g' . (Remember that $U' \subseteq K'$, by (9.40), so that g' takes only the value σ on $M' \setminus K'$. Remember also that the degree does not depend on the choice of compact set — in these cases K and K' — so long as the mapping is constant off the compact set.)

Roughly speaking, (9.20) says that h is the composition of g' and ϕ . This is not quite true, because h is defined on all of M and ϕ is not, but it is true near K , which is good enough for the computation of homology that we need to make. The point now is to understand what ϕ does to homology well enough to conclude that $d = \pm d'$.

Sublemma 9.56. *The relative homology groups $H_n(M, M \setminus K)$, $H_n(M', M' \setminus K')$, and $H_n(M', M' \setminus L')$ are all infinite cyclic groups, and the homology classes λ_K , $\lambda'_{K'}$, and $\lambda_{L'}$ are generators for these groups (respectively).*

This follows from the connectedness of the compact sets K , K' , and L' . (See Lemma A.28.)

We need to sort out how our various mappings induce homomorphisms between homology groups. Recall that if X and Y are topological spaces, $A \subseteq X$, $B \subseteq Y$, $f : X \rightarrow Y$ is continuous, and $f(A) \subseteq B$ (i.e., $f : (X, A) \rightarrow (Y, B)$), then f induces a homomorphism $f_* : H_n(X, A) \rightarrow H_n(Y, B)$. This notation is slightly bad, because f_* does not reflect the dependence on (X, A) , (Y, B) , and there are often many reasonable choices for these pairs. However, the various choices are usually related in a simple way. In our case we shall suffer small confusions by needing to use different pairs of spaces at different times, but it all washes out.

A priori our maps ϕ' , ϕ , and ψ' are maps as follows:

$$(9.57) \quad \phi' : U'(.5) \rightarrow U(.6), \quad \phi : U(.5) \rightarrow U(.6), \quad \psi' : U'(.4) \rightarrow U'(.6).$$

See Lemma 9.17 and (9.30). We can also restrict their domains, or view them as mappings between pairs of spaces in various ways. For our purposes it will be convenient to view them as follows:

$$(9.58) \quad \phi' : (U'(.4), U'(.4) \setminus L') \rightarrow (U(.5), U(.5) \setminus K),$$

$$(9.59) \quad \phi : (U(.5), U(.5) \setminus K) \rightarrow (M', M' \setminus K'),$$

$$(9.60) \quad \psi' : (U'(.4), U'(.4) \setminus L') \rightarrow (M', M' \setminus K').$$

In (9.58) we are saying that $\phi'(U'(.4)) \subseteq U(.5)$ and that $\phi'(U'(.4) \setminus L') \subseteq U(.5) \setminus K$. The first inclusion follows from (9.29), while the second comes from (9.42). Similarly (9.59) follows

from (9.43), and (9.60) is a special case of (9.44) (by (9.35)). With these interpretations of ϕ' , ϕ , and ψ' we have that $\psi' = \phi \circ \phi'$, i.e., this composition is compatible with our choices of source and targets.

Corresponding to (9.58)-(9.60) we have the induced homomorphisms

$$(9.61) \quad \phi'_* : H_n(U'(.4), U'(.4) \setminus L') \rightarrow H_n(U(.5), U(.5) \setminus K),$$

$$(9.62) \quad \phi_* : H_n(U(.5), U(.5) \setminus K) \rightarrow H_n(M', M' \setminus K'),$$

$$(9.63) \quad \psi'_* : H_n(U'(.4), U'(.4) \setminus L') \rightarrow H_n(M', M' \setminus K')$$

on homology. By functoriality we have that

$$(9.64) \quad \psi'_* = \phi_* \circ \phi'_*.$$

These homomorphisms live on inconvenient spaces, and so we want to make some identifications. The “excision theorem” (Theorem A.10) implies that the homomorphisms

$$(9.65) \quad H_n(U'(.4), U'(.4) \setminus L') \rightarrow H_n(M', M' \setminus L')$$

$$(9.66) \quad H_n(U(.5), U(.5) \setminus K) \rightarrow H_n(M, M \setminus K)$$

induced by the obvious inclusions are isomorphisms. This application of excision is permitted because $L' \subseteq U'(.1)$ and $K \subseteq U(.1)$ (Sublemma 9.39), so that the closure of $M' \setminus U(.4)$ is contained in the interior of $M' \setminus L'$ and the closure of $M \setminus U(.5)$ is contained in the interior of $M \setminus K$. One can think of the pairs of relative homology groups in (9.65) and (9.66) as being identified via the inclusion isomorphisms. Using these isomorphisms we can convert the homomorphisms in (9.61)-(9.63) into

$$(9.67) \quad \tilde{\phi}'_* : H_n(M', M' \setminus L') \rightarrow H_n(M, M \setminus K),$$

$$(9.68) \quad \tilde{\phi}_* : H_n(M, M \setminus K) \rightarrow H_n(M', M' \setminus K'),$$

$$(9.69) \quad \tilde{\psi}'_* : H_n(M', M' \setminus L') \rightarrow H_n(M', M' \setminus K').$$

Sublemma 9.70. $\tilde{\psi}'_* = \tilde{\phi}_* \circ \tilde{\phi}'_*.$

This follows from chasing definitions.

Let us now use (9.55) to relate the degrees of h and g' .

Sublemma 9.71. $h_*(\lambda_K) = g'_*(\tilde{\phi}_*(\lambda_K)).$

This is almost just the definition (9.20) of h and functoriality, but we should be a little careful. As in (9.55), the induced mappings h_* and g'_* are, more specifically,

$$(9.72) \quad \begin{aligned} h_* &: H_n(M, M \setminus K) \rightarrow H_n(\mathbf{S}^n, \{\sigma\}) \\ g'_* &: H_n(M', M' \setminus K') \rightarrow H_n(\mathbf{S}^n, \{\sigma\}). \end{aligned}$$

We know from the excision theorem that the homomorphism (9.66) induced by inclusion is an isomorphism, and hence there is a relative homology class $\lambda \in H_n(U(.5), U(.5) \setminus K)$

whose image in $H_n(M, M \setminus K)$ under the homomorphism induced by inclusion is λ_K . Let h_*^1 denote the homomorphism

$$(9.73) \quad h_*^1 : H_n(U(.5), U(.5) \setminus K) \rightarrow H_n(\mathbf{S}^n, \{\sigma\})$$

induced by h . We have that

$$(9.74) \quad h_*^1(\lambda) = h_*(\lambda_K),$$

because of the usual compatibility properties of homology. (This comes down to the statement that $h : (U(.5), U(.5) \setminus K) \rightarrow (\mathbf{S}^n, \{\sigma\})$ is the composition of $h : (M, M \setminus K) \rightarrow (\mathbf{S}^n, \{\sigma\})$ with the inclusion $(U(.5), U(.5) \setminus K) \rightarrow (M, M \setminus K)$.) On the other hand, h , as a map from $(U(.5), U(.5) \setminus K)$ into $(\mathbf{S}^n, \{\sigma\})$, is just the composition of ϕ as in (9.59) with $g' : (M', M' \setminus K') \rightarrow H_n(\mathbf{S}^n, \{\sigma\})$. Thus

$$(9.75) \quad h_*^1(\lambda) = g'_*(\phi_*(\lambda)).$$

We also have that $\tilde{\phi}_*(\lambda_K) = \phi_*(\lambda)$, by definitions. This proves Sublemma 9.71.

To finish the proof of Lemma 9.28 it suffices to show the following.

Sublemma 9.76. *$\tilde{\phi}_*(\lambda_K)$ is either $\lambda'_{K'}$ or $-\lambda'_{K'}$.*

To prove this we need to look again at the mappings (9.67)-(9.69).

Let j denote the obvious inclusion from $(U'(.4), U'(.4) \setminus L')$ into $(M', M' \setminus K')$, and let $j_* : H_n(U'(.4), U'(.4) \setminus L') \rightarrow H_n(M', M' \setminus K')$ denote the induced homomorphism on homology.

Claim 9.77. $\psi'_* = j_*$

Indeed, as mappings from $(U'(.4), U'(.4) \setminus L')$ into $(M', M' \setminus K')$, ψ and j are homotopic, with the homotopy being provided by Ψ . This follows from Sublemma 9.33, especially (9.35), and from (9.44). The existence of this homotopy implies that the induced mappings on homology are the same. (See Appendix A below, and also p.173-174 of [Ma].) This proves the claim.

Claim 9.78. j_* is an isomorphism.

We can write j as the composition of the inclusions $(U'(.4), U'(.4) \setminus L') \rightarrow (M', M' \setminus L')$ and $(M', M' \setminus L') \rightarrow (M', M' \setminus K')$, and j_* is the composition of the two induced maps on homology

$$(9.79) \quad H_n(U'(.4), U'(.4) \setminus L') \rightarrow H_n(M', M' \setminus L') \quad \text{and}$$

$$(9.80) \quad H_n(M', M' \setminus L') \rightarrow H_n(M', M' \setminus K').$$

The homomorphism in (9.79) is an isomorphism, because of excision (Theorem A.10) and the fact that $L' \subseteq U'(.1)$ (Sublemma 9.39). The homomorphism in (9.80) sends $\lambda'_{L'}$ to $\lambda'_{K'}$. (This follows from the uniqueness statement in Theorem A.17.) The two relative homology groups in (9.80) are both infinite cyclic groups which are generated by $\lambda'_{L'}$ and

$\lambda'_{K'}$, respectively, by Sublemma 9.56. This implies that (9.80) is also an isomorphism, and the claim follows.

Claim 9.81. $\tilde{\psi}'_{\star}$ is an isomorphism.

Indeed, $\psi'_{\star}$ is an isomorphism by Claims 9.77 and 9.78, and $\tilde{\psi}'_{\star}$ is defined by $\psi'_{\star}$ through the isomorphism (9.65).

Claim 9.82. $\tilde{\phi}_{\star}$ is an isomorphism.

Consider first the following statement: if the composition of two group homomorphisms on the group of integers $\mathbf{Z}$ is an isomorphism, then each of the two homomorphisms is an isomorphism. This is an elementary exercise. Claim 9.82 follows from this fact, because of Sublemma 9.70 and Claim 9.81, and because Sublemma 9.56 ensures that the relevant homology groups are isomorphic to $\mathbf{Z}$.

It is easy to derive Sublemma 9.76 from Sublemma 9.56 and Claim 9.82.

Lemma 9.28 now follows from (9.55) and Sublemmas 9.71 and 9.76. Proposition 9.15 follows in turn from Lemma 9.28, and Theorem 9.1 follows from Proposition 9.15.

This completes the proof of Theorem 1.29 (b).

10. Lipschitz bounds for homotopies of Lipschitz mappings.

We shall prove Theorem 1.30 in this section. We first formulate a variant of it which will be more convenient to prove. This variant holds under much more general conditions, and so the statement will be more complicated than we need for Theorem 1.30. We are going to both weaken the assumptions that we make on M and permit more general target spaces than $\mathbf{S}^n$. We shall not maximize the generality of the targets that we allow, but we shall permit some pretty general targets.

Definition 10.1 A space E is called an *ENR*, or *Euclidean Neighborhood Retract*, if it is a closed set inside some Euclidean space $\mathbf{R}^k$ and if there is an open set $W \subseteq \mathbf{R}^k$ with $E \subseteq W$ and a continuous mapping $R : W \rightarrow E$ which is a retraction, i.e., $R(x) = x$ for all $x \in E$. In topology one might think of a space as being an ENR if it is homeomorphic to such a set, but we shall consider E to be an actual set in $\mathbf{R}^k$ which is equipped with the induced Euclidean metric. We shall call E a *Lipschitz ENR* if the retraction R can be taken to be Lipschitz. Of course a sphere $\mathbf{S}^n$ satisfies these conditions, as do finite polyhedra, etc. We shall sometimes call $\{W, R\}$ the *entourage* of E .

We shall typically assume that E is compact, so that there is an $\epsilon > 0$ such that

$$(10.2) \quad x \in W \quad \text{whenever} \quad \text{dist}(x, E) \leq 10\epsilon.$$

Theorem 10.3. *Let E be a compact ENR, and fix a point $\sigma \in E$. Let $\nu : [0, \infty) \rightarrow [0, \infty)$ be a function which is continuous at the origin and satisfies $\nu(0) = 0$.*

Suppose that $(M, U, d(x, y))$ is an admissible triple (Definition 1.5) with covering function $N(\cdot)$ (Definition 1.6) and contractability function $\rho(\cdot)$ (Definition 1.7). (As in Definition 1.6, $N(\cdot)$ should satisfy $N(t) = O(t^{-k})$ for some k .) Assume also that $\text{diam}_M U = 1$.

Let $f_0, f_1 : M \rightarrow E$ be continuous mappings which take only the value σ on $M \setminus U$, are continuous with modulus of continuity ν (so that $|f_i(y) - f_i(x)| \leq \nu(d(x, y))$ for all $x, y \in M$), and are homotopic through continuous mappings from M into E which take only the value σ on $M \setminus U$. (We do not require that this homotopy satisfy any bounds on its modulus of continuity.)

Then there is a homotopy between f_0 and f_1 with controlled modulus of continuity. That is, there is function $\omega : [0, \infty) \rightarrow [0, \infty)$ and a mapping $F : M \times [0, 1] \rightarrow E$ such that $\omega(0) = 0$, ω is continuous at the origin,

$$(10.4) \quad F(x, 0) = f_0(x) \text{ and } F(x, 1) = f_1(x) \text{ for all } x \in M,$$

$$(10.5) \quad F(x, t) = \sigma \text{ when } x \in M \setminus U_1 \text{ and } t \in [0, 1], \text{ and}$$

$$(10.6) \quad |F(x, t) - F(y, s)| \leq \omega(d(x, y) + |s - t|) \\ \text{for all } x, y \in M, s, t \in [0, 1].$$

This function ω depends on E (and its entourage), ν , $N(\cdot)$, and $\rho(\cdot)$, but not on anything else.

What happens if we try to apply Theorem 10.3 under the assumptions of Theorem 1.30, with $E = \mathbf{S}^n$? Of course $\mathbf{S}^n$ is a compact ENR, so that is fine. The assumptions on $(M, U, d(x, y))$ in Theorem 10.3 hold under Standard Assumptions 1.8. We have normalized the diameter of U to be 1 here, but that is not a restriction, because Theorem 1.30 was formulated to be scale-invariant. In Theorem 1.30 we required f_0 and f_1 to be Lipschitz with norm $\leq Lr^{-1}$, which just means that we can take $\nu(t) = Lt$ (since we have made the normalization $r = \text{diam } U = 1$). We have assumed here that f_0 and f_1 are homotopic through mappings constant off U , while in Theorem 1.30 we assumed only that f_0 and f_1 had the same degree. The latter implies the former, by a well-known result in topology (Proposition A.29). This step uses the extra structure assumed in Theorem 1.30, to wit, the connectedness of U and the requirement that M be an oriented topological manifold (from Standard Assumptions 1.8), and the fact that the target space is $\mathbf{S}^n$. The remaining difference is that Theorem 1.30 asks for a Lipschitz bound on the homotopy F , while Theorem 10.3 provides only a bound on the modulus of continuity. Let us first see how to pass from (10.6) to a Lipschitz bound, at least in the case where $E = \mathbf{S}^n$, and then deal with Theorem 10.3 afterwards.

Lemma 10.7. *Let E be a Lipschitz ENR which is also compact, let $(M, d(x, y))$ be a metric space, and let V be an open subset of M . Suppose that $f_0, f_1 : M \rightarrow E$ are L -Lipschitz mappings for some constant L , and suppose that there exist a function $\omega : [0, \infty) \rightarrow [0, \infty)$ and a mapping $F : M \times [0, 1] \rightarrow E$ such that $\omega(0) = 0$, ω is continuous at the origin, and (10.4)-(10.6) hold, with U_1 replaced by V in (10.5). Then there is a mapping $F' : M \times [0, 1] \rightarrow E$ such that (10.4) and (10.5) hold, with U_1 replaced by V in (10.5), and such that F' is $C(L)$ -Lipschitz. Here $C(L)$ depends on E (and its entourage), L , and ω , but nothing else.*

Note that we ask only that M be a metric space here, and nothing like extra conditions in Theorem 10.3. We switch from U_1 to V here to make this point more clear. (On the

other hand, if we did make extra assumptions on M , then we could weaken the hypotheses on E .)

Let us prove Lemma 10.7. Let V, f_0, f_1, F , etc., be given, as above.

Let W and $R : W \rightarrow E$ be as in Definition 10.1, with R Lipschitz in this case. Let $\epsilon > 0$ be as in (10.2).

We want to approximate F by a Lipschitz function, and so we use Lemma 2.4, but applied to the metric space $M \times [0, 1]$ (with the metric $D((x, s), (y, t)) = d(x, y) + |s - t|$) and the function F . More precisely, we apply Lemma 2.4 to each component of F separately, with F viewed as map into $\mathbf{R}^k$. For the function ϕ in (2.8) (with f, M replaced with $F, M \times [0, 1]$) we can simply take ω . Lemma 2.4 provides us with approximations $F_j : M \times [0, 1] \rightarrow \mathbf{R}^k$ to F which are κj -Lipschitz and which satisfy

$$(10.8) \quad \sup_{M \times [0, 1]} |F_j - F| \leq \kappa \omega(2j^{-1}).$$

These extra factors of κ come from the fact that we are applying Lemma 2.4 to the κ components of F separately. The estimate (10.8) is just (2.7) rewritten in the current notation. Choose j to be the smallest positive integer such that $\kappa \omega(2j^{-1}) \leq \epsilon$, and let it be fixed from now on. Thus j is bounded by a constant that depends only on ϵ, ω , and κ , and this F_j has bounded Lipschitz norm. Let us rewrite (10.8) as

$$(10.9) \quad \sup_{M \times [0, 1]} |F_j - F| \leq \epsilon$$

for this choice of j .

There are several problems with F_j . It does not necessarily take values in E , and we do not automatically have that $F_j(x, t) = \sigma$ when $x \in M \setminus V$. It also does not reduce to f_0 and f_1 when $t = 0, 1$. To take care of the first problems we shall compose F_j with some auxiliary mappings. Basically we just want to compose it with our retraction R , but we need to do a little more to fix it on $M \setminus V$.

Let $\beta : \mathbf{R}^k \rightarrow \mathbf{R}^k$ be a Lipschitz mapping such that $\beta(x) = x$ when $x \notin B(\sigma, 2\epsilon)$, $\beta(B(\sigma, 2\epsilon)) \subseteq B(\sigma, 2\epsilon)$, $\beta(x) = \sigma$ when $x \in \overline{B}(\sigma, \epsilon)$. This is easy to construct. Of course β depends only on κ and ϵ . Note that $|\beta(x) - x| \leq 4\epsilon$ for all $x \in \mathbf{R}^k$.

Define $G : M \times [0, 1] \rightarrow E$ by $G = R \circ \beta \circ F_j$. We must first check that this is well-defined, i.e., that $\beta \circ F_j$ takes values in W . Let us show that

$$(10.10) \quad \text{dist}(\beta(F_j(x, t)), E) \leq 2\epsilon \quad \text{for all } (x, t) \in E \times [0, 1].$$

This is automatic when $F_j(x, t) \in B(\sigma, 2\epsilon)$, since $\beta(F_j(x, t)) \in B(\sigma, 2\epsilon)$ in that case, and $\sigma \in E$. If $F_j(x, t) \notin B(\sigma, 2\epsilon)$, then $\beta(F_j(x, t)) = F_j(x, t)$, and (10.10) follows from (10.9) and the fact that F takes values in E . Thus we have (10.10). It follows from (10.2) that $\beta \circ F_j$ takes values in W , so that G is well-defined.

Notice that G is Lipschitz with bounded norm, since the same is true of F_j , and since R and β are just fixed Lipschitz mappings (which depend only on E and its entourage).

Next let us verify that

$$(10.11) \quad G(x, t) = \sigma \quad \text{when } x \in M \setminus V \text{ and } t \in [0, 1].$$

If $x \in M \setminus V$ and $t \in [0, 1]$, then $|F_j(x, t) - \sigma| \leq \epsilon$, because of (10.9) and our assumption that F satisfy (10.5) with U_1 replaced with V . Hence

$$(10.12) \quad \beta(F_j(x, t)) = \sigma,$$

by definition of β . Therefore $G(x, t) = R(\beta(F_j(x, t))) = R(\sigma) = \sigma$ (since R equals the identity on E). This implies (10.11).

Unfortunately G does not reduce to f_0 and f_1 at $t = 0, 1$, and so we must do some more to fix that. Define $f_{0,j}$ and $f_{1,j}$ by $f_{0,j}(x) = F_j(x, 0)$, $f_{1,j}(x) = F_j(x, 1)$. Define $G_i : M \times [0, 1] \rightarrow E$ for $i = 0, 1$ by

$$(10.13) \quad G_i(x, t) = R((1-t)f_i(x) + t\beta(f_{i,j}(x))).$$

Let us first check that this makes sense. For each $x \in M$ we have that

$$(10.14) \quad \begin{aligned} |f_i(x) - \beta(f_{i,j}(x))| &\leq |f_i(x) - f_{i,j}(x)| + |f_{i,j}(x) - \beta(f_{i,j}(x))| \\ &\leq \epsilon + 4\epsilon = 5\epsilon. \end{aligned}$$

In the second inequality we used (10.9) and the definition of β . This implies that the distance between $f_i(x)$ and $(1-t)f_i(x) + t\beta(f_{i,j}(x))$ is $\leq 5\epsilon$ when $0 \leq t \leq 1$. Therefore the distance from $(1-t)f_i(x) + t\beta(f_{i,j}(x))$ to E is $\leq 5\epsilon$, since f_i takes values in E , and $(1-t)f_i(x) + t\beta(f_{i,j}(x))$ lies in the domain W of R , by (10.2). Thus (10.13) is well defined.

We also have that

$$(10.15) \quad G_i(x, t) = \sigma \quad \text{when } x \in M \setminus V \text{ and } t \in [0, 1].$$

$i = 0, 1$. Indeed, if $x \in M \setminus V$, then $\beta(f_{i,j}(x)) = \sigma$, by (10.12), and of course $f_i(x) = \sigma$ by assumptions. Thus (10.15) reduces to $R(\sigma) = \sigma$, just as for (10.11). This proves (10.15).

The G_i 's are Lipschitz with bounded norm. This follows from (10.13), since we have Lipschitz bounds for f_i and $f_{i,j}$, and since R and β are Lipschitz.

Define $g_0, g_1 : M \rightarrow E$ by $g_0(x) = G(x, 0)$, $g_1(x) = G(x, 1)$. Thus G is a homotopy between g_0 and g_1 , and G_i is a homotopy between f_i and g_i , $i = 0, 1$, because (10.13) reduces to $R \circ \beta \circ f_{i,j}$ when $t = i$, $i = 0, 1$. Let F' be the homotopy between f_0 and f_1 obtained by combining G_0 , G , and G_1 in the obvious manner. Then F' has bounded Lipschitz norm, because of the corresponding bounds for G and G_i . We also have that $F'(x, t) = \sigma$ when $x \in M \setminus V$ and $t \in [0, 1]$, because of (10.11) and (10.15). Thus F' has all the required properties.

This completes the proof of Lemma 10.7.

This also finishes the proof of the assertion that Theorem 1.30 will follow from Theorem 10.3.

To prove Theorem 10.3 we first establish a weaker result in which the bounds depend on the given metric space. We shall reduce to that case afterwards, by approximating our space M by a polyhedron that comes from a finite collection of polyhedra.

Lemma 10.16. *Let E be a compact ENR, and fix a point $\sigma \in E$. Let $\nu : [0, \infty) \rightarrow [0, \infty)$ be a function which is continuous at the origin and satisfies $\nu(0) = 0$.*

Let $(X, d(x, y))$ be a compact metric space, and let K be a closed subset of X . Suppose that $f_0, f_1 : X \rightarrow E$ is a pair of continuous mappings with modulus of continuity ν (so that $|f_i(x) - f_i(y)| \leq \nu(d(x, y))$ for all $x, y \in M$) which take only the value σ on K and which are homotopic to each other through mappings which take only the value σ on K . (We do not assume a bound on the modulus of continuity of the homotopy.)

Then we can find a homotopy between f_0 and f_1 which has a controlled modulus of continuity. That is, there exists a function $\omega(t) : [0, \infty) \rightarrow [0, \infty)$, depending only on E (and its entourage), X , K , and ν (and not on f_0 or f_1), and a mapping $F : X \times [0, 1] \rightarrow E$ such that $\omega(0) = 0$, $\omega(t)$ is continuous at the origin,

$$(10.17) \quad F(x, t) = \sigma \text{ when } x \in K \text{ and } t \in [0, 1],$$

and the analogues of (10.4) and (10.6) hold with M replaced by X .

This lemma would be automatic if we were willing to permit ω to depend on f_0 and f_1 , because continuous mappings on compact metric spaces are uniformly continuous. We could simply take F to be the homotopy provided by the hypotheses, and we would take ω to be its modulus of continuity (as in (2.8)). To prove the lemma we shall reduce to the case where we are allowed to have ω depend on f_0 and f_1 by using the Arzela-Ascoli theorem to reduce the problem to only finitely many choices of f_0 and f_1 .

Let $\mathcal{F}$ denote the family of continuous mappings from X to E with modulus of continuity ν which take only the value σ on K . Choose $\epsilon > 0$ so that (10.2) holds. The Arzela-Ascoli theorem implies that there is a finite subset $\mathcal{G}$ of $\mathcal{F}$ such that

$$(10.18) \quad \text{for each } f \in \mathcal{F} \text{ there is a } g \in \mathcal{G} \text{ with } \sup_X |f - g| \leq \epsilon.$$

(Note that we are using the compactness of X and E here.) If $f \in \mathcal{F}$ and $g \in \mathcal{G}$ are as in (10.18), then we claim that there is a homotopy $H : X \times [0, 1] \rightarrow E$ between them which has a controlled modulus of continuity (as a function on $X \times [0, 1]$) and which satisfies (10.17) with F replaced by H . To see this we take

$$H(x, t) = R((1 - t)f(x) + tg(x)),$$

where R is the retraction promised in Definition 10.1. This is well defined, because (10.18) and (10.2) ensure that $(1 - t)f(x) + tg(x)$ takes values in W when $x \in X$ and $0 \leq t \leq 1$ (since f and g take values in E). Clearly H defines a homotopy between f and g , and we need to check that it has a controlled modulus of continuity. Observe that $(1 - t)f(x) + tg(x)$ takes values in the compact subset $\hat{E} = \{u \in \mathbf{R}^n : \text{dist}(u, E) \leq \epsilon\}$ of W . Since R is continuous it is uniformly continuous on $\hat{E}$, and therefore we can control the modulus of continuity of H in terms of ν and the modulus of continuity of R on $\hat{E}$ (which does not depend on f or g). Thus H has all the required properties.

If $g_0, g_1 \in \mathcal{G}$ are homotopic through mappings which take the value σ on K , then we can find a homotopy between them of the same type which satisfies the analogue of (10.6) with a choice of ω that does not depend on g_0 or g_1 . This is true because $\mathcal{G}$ is finite, so that this assertion is practically the same as the case where ω is permitted to depend on

g_0 and g_1 . It just comes down to the fact that any homotopy is uniformly continuous, by compactness.

Suppose now that we are given any $f_0, f_1 \in \mathcal{F}$ which are homotopic through mappings which take only the value σ on K . We want to find a homotopy F between them as in Lemma 10.16, with a uniform choice of ω . To do this we first choose $g_0, g_1 \in \mathcal{G}$ corresponding to f_0, f_1 as in (10.18). We can homotop f_0 to g_0 and f_1 to g_1 with homotopies H_0 and H_1 that have controlled moduli of continuity, as noted above. This implies that g_0 and g_1 are homotopic, and hence we can find a homotopy G between them which satisfies (10.6) with a uniform choice of ω , by the argument in the preceding paragraph. All these homotopies take only the value σ on K . We can combine these three homotopies H_0, G, H_1 to get a homotopy F of f_0 and f_1 with the required properties. In particular the combined homotopy has a controlled modulus of continuity. This proves Lemma 10.16.

The basic plan for proving Theorem 10.3 is to reduce the problem from M to a finite polyhedron, and in fact one of only finitely many finite polyhedra. This will permit us to use Lemma 10.16. In order to make this reduction we first need to approximate our metric spaces by finite polyhedra. The next result contains the information that we need. (The reader may wish to review Definition 4.8 for terminology pertaining to finite polyhedra.)

Lemma 10.19. *Suppose that $(M, U, d(x, y))$ is as in Theorem 10.3. For each $\theta > 0$ there exists an integer l , a finite polyhedron P in $\mathbf{R}^l$, a C -Lipschitz mapping $\phi : U(1/2) \rightarrow P$, and a continuous mapping $\psi : P \rightarrow U(3/4)$ with the following properties:*

$$(10.20) \quad \text{diam } \phi^{-1}(\tau) \leq \theta \quad \text{for all (distinguished) simplices } \tau \text{ in } P;$$

$$(10.21) \quad \text{diam } \psi(\tau) \leq \theta \quad \text{for all (distinguished) simplices } \tau \text{ in } P;$$

$$(10.22) \quad d_M(\psi(\phi(x)), x) \leq \theta \quad \text{for all } x \in U(1/2).$$

Here l and C depend only on θ , the covering function $N(\cdot)$, and the contractability function $\rho(\cdot)$ from the hypotheses of Theorem 10.3, and not on anything else. The polyhedron P can be selected from a finite collection $\mathcal{P}$ of finite polyhedra in $\mathbf{R}^l$, where $\mathcal{P}$ depends only on θ , $N(\cdot)$, and $\rho(\cdot)$.

Note that we are not asserting any bound for the modulus of continuity of ψ .

This is just a portion of Lemma 8.3 rewritten for the present circumstances, and with f and g in Lemma 8.3 relabelled as ϕ and ψ . See the remark which precedes Lemma 8.1 for a discussion of the hypotheses of Lemma 8.3, but notice that there is a small subtlety here concerning the hypotheses. In Lemma 8.3 we had a special dimension n in play, and it was important that the polyhedron P have dimension $\leq n$. In the present circumstances there is no such special dimension. However we still need to have at least a bound on the topological dimension of $U(1)$ in order to apply Lemma 8.3. We get this from the requirement that $N(t) = O(t^{-k})$ as $t \rightarrow 0$, as in Corollary 4.7. Thus we get a bound on the topological dimension that depends on $N(\cdot)$, and that is fine. (Note that Lemma 8.3 used Lemma 8.1, which itself used Theorem 6.5. In this case, since we do not have a special dimension that we care about, we can use Proposition 6.4 instead of Theorem 6.5.)

Lemma 8.3 also provides a homotopy H between $\psi \circ \phi$ and the identity. It turns out that we shall not need this homotopy here, because of our bounds on the moduli of continuity of f_0 and f_1 . The proof of Lemma 8.3 is simpler without the homotopy.

Let us now begin the proof of Theorem 10.3 in earnest. Let $E, (M, U, d(x, y)), \sigma, f_0, f_1$, and ν be as given in Theorem 10.3. Let $\mathbf{R}^k, W$, and R be as in Definition 10.1, and let $\epsilon > 0$ be as in (10.2). Choose θ so that

$$(10.23) \quad \theta < 10^{-2} \quad \text{and} \quad \sup_{0 \leq t \leq \theta} \nu(t) \leq \frac{\epsilon}{2}$$

(and so that θ depends only on ν and ϵ). Let l, P, ϕ, ψ , and $\mathcal{P}$ be as in Lemma 10.19, corresponding to this choice of θ . Recall from Definition 4.8 that $\mathcal{S}(P)$ denotes the collection of distinguished simplices in P . Define $K \subseteq P$ by

$$(10.24) \quad K = \bigcup \{ \tau \in \mathcal{S}(P) : v(\tau) \cap U = \emptyset \}.$$

In the next lemma we obtain models for f_0 and f_1 on P . We shall prove Theorem 10.3 by working with these models and then going back to the f_i 's and M .

Lemma 10.25. *There exist continuous mappings $g_0, g_1 : P \rightarrow E$ and a function $\mu : [0, \infty) \rightarrow [0, \infty)$ such that the following statements are true.*

- (a) μ is continuous at 0, $\mu(0) = 0$, and μ depends only on E (and its entourage), ν , the covering function $N(\cdot)$, and the contractability function $\rho(\cdot)$.
- (b) $g_i \equiv \sigma$ on K , $i = 0, 1$.
- (c) g_i is homotopic to $f_i \circ \psi$ through continuous mappings from P into E which take only the value σ on K .
- (d) $|g_i(p) - g_i(q)| \leq \mu(d(p, q))$ for all $p, q \in P$.
- (e) If we define $\tilde{g}_i : M \rightarrow E$ by

$$(10.26) \quad \begin{aligned} \tilde{g}_i(x) &= g_i(\phi(x)) && \text{when } x \in U(1/2) \\ &= \sigma && \text{when } x \in M \setminus U(1/2), \end{aligned}$$

$i = 0, 1$, then $\tilde{g}_i$ is continuous. (Recall the definition of $U(t)$, given shortly before Lemma 8.1.)

(f) There are continuous mappings $H_i : M \times [0, 1] \rightarrow E$, $i = 0, 1$, which provide homotopies between f_i and $\tilde{g}_i$ with bounded modulus of continuity. Specifically, $H_i(x, 0) = f_i(x)$ and $H_i(x, 1) = \tilde{g}_i(x)$ for all $x \in M$, $H_i(x, t) = \sigma$ when $x \in M \setminus U(3\theta)$ and $0 \leq t \leq 1$, and

$$(10.27) \quad \begin{aligned} |H_i(x, t) - H_i(y, s)| &\leq \mu(d_M(x, y) + |s - t|) \\ &\text{for all } x, y \in M, s, t \in [0, 1]. \end{aligned}$$

$i = 0, 1$.

To prove this we define first an auxiliary mapping $h_i : P \rightarrow \mathbf{R}^k$ for $i = 0, 1$ by setting $h_i(p) = f_i(\psi(p))$ when p is a vertex of P and by extending h_i to the various distinguished simplices τ of P by requiring h_i to be affine on each such τ .

Let us check that

$$(10.28) \quad h_i \equiv \sigma \quad \text{on } K.$$

If $\tau \in \mathcal{S}(P)$ is one of the distinguished simplices of P , and if $\tau \subseteq K$, then $\psi(p) \in M \setminus U$ for each vertex p of τ . This means that $h_i(p) = f_i(\psi(p)) = \sigma$ for all these vertices. The definition of h_i implies that $h_i \equiv \sigma$ on such a τ , and (10.28) follows.

Next let us check that

$$(10.29) \quad |h_i(p) - f_i(\psi(p))| \leq \sup_{0 \leq t \leq \theta} \nu(t) \leq \frac{\epsilon}{2} \quad \text{for all } p \in P.$$

Fix $p \in P$ and let τ be a distinguished simplex in P which contains p . Let $V(\tau)$ denote the set of vertices of τ . Thus $h_i(p)$ is a convex combination of the points $h_i(q)$, $q \in V(\tau)$, and $h_i(q) = f_i(\psi(q))$ for each $q \in V(\tau)$, by definition of h_i . Also, $d(\psi(p), \psi(q)) \leq \theta$ for each $q \in V(\tau)$, by (10.21), and hence

$$(10.30) \quad |f_i(\psi(p)) - f_i(\psi(q))| \leq \nu(d(\psi(p), \psi(q))) \leq \sup_{0 \leq t \leq \theta} \nu(t) \leq \frac{\epsilon}{2}$$

for each $q \in V(\tau)$, since f_i has modulus of continuity ν , by assumption. These facts combine to give (10.29).

Set $\widehat{E} = \{u \in \mathbf{R}^k : \text{dist}(u, E) \leq \epsilon\}$. This is a compact subset of W , since E is compact, and because of (10.2).

Notice that

$$(10.31) \quad h_i(p) \in \widehat{E} \quad \text{and} \quad (1-t)h_i(p) + tf_i(\psi(p)) \in \widehat{E} \quad \text{for all } p \in P, t \in [0, 1].$$

This follows from (10.29), since f_i takes values in E .

Define $g_i : P \rightarrow E$ by $g_i = R \circ h_i$. This is well defined and continuous because of (10.31). Part (b) of Lemma 10.25 follows from (10.28) and the fact that R equals the identity on E and $R(\sigma) = \sigma$ in particular.

To get part (c) of Lemma 10.25 we simply take $R((1-t)h_i(p) + tf_i(\psi(p)))$ for the required homotopy. This is defined and continuous because of (10.31). If $p \in K$, then $h_i(p) = \sigma$, by (10.28), and $f_i(\psi(p)) = \sigma$ because $\psi(p) \in M \setminus U$, by the definition (10.24) of K . This and the fact that $R(\sigma) = \sigma$ imply that our homotopy takes only the value σ on K , as desired.

Let us now prove (d). Observe first that $h_i : P \rightarrow \mathbf{R}^k$ is Lipschitz with bounded norm (depending only on the allowed objects). This is because h_i is bounded (since it takes values in the convex hull of the compact set E) and because it is affine on each distinguished simplex τ of P . Remember also that P is a finite polyhedron, so that there are only finitely many distinguished simplices. This implies that h_i is Lipschitz with norm bounded by a constant that depends only on the polyhedron P , and we know from Lemma 10.19 that there are only finitely many possibilities for P . Since g_i is simply $R \circ h_i$, we get a uniform bound for the modulus of continuity for g_i from the Lipschitz bound for h_i . More precisely, h_i takes values in $\widehat{E}$, and R is uniformly continuous on $\widehat{E}$ since it is compact. Thus our bound on the modulus of continuity of the g_i 's depends on the modulus of continuity of R on $\widehat{E}$, but that is fine.

To prove (e) it suffices to show that

$$(10.32) \quad \tilde{g}_i(x) = \sigma \quad \text{when } x \in M \setminus U(3\theta), 0 \leq t \leq 1.$$

Let us check that

$$(10.33) \quad \phi(x) \in K \quad \text{when } x \in U(1/2) \setminus U(3\theta).$$

Suppose that this is not true. Let $x \in U(1/2) \setminus U(3\theta)$ be given, and let τ be a distinguished simplex in P such that $\phi(x) \in \tau$. If $\phi(x) \notin K$ then we have that $\psi(\tau) \cap U \neq \emptyset$, because of (10.24). Since $\phi(x) \in \tau$ we get that $\text{dist}_M(\psi(\phi(x)), U) \leq \theta$, by (10.21). This implies that $\text{dist}_M(x, U) \leq 2\theta$, because of (10.22). This contradicts our choice of x , and (10.33) follows. This implies (10.32), because of part (b) of Lemma 10.25. The continuity of $\tilde{g}_i$ follows from (10.32), and this is part (e) of Lemma 10.25.

Before we start to prove part (f) of Lemma 10.25 define $\tilde{h}_i : M \rightarrow E$ by

$$(10.34) \quad \begin{aligned} \tilde{h}_i(x) &= h_i(\phi(x)) && \text{when } x \in U(1/2), \\ &= \sigma && \text{when } x \in M \setminus U(1/2), \end{aligned}$$

$i = 0, 1$. As for $\tilde{g}_i$ we have that

$$(10.35) \quad \tilde{h}_i(x) = \sigma \quad \text{when } x \in M \setminus U(3\theta), \quad 0 \leq t \leq 1,$$

because of (10.33) and (10.28). Thus $\tilde{h}_i$ is continuous.

To prove part (f) of Lemma 10.25 we want to take

$$(10.36) \quad H_i(x, t) = R((1-t)f_i(x) + t\tilde{h}_i(x)).$$

We need to show that this makes sense, etc.

Notice first that

$$(10.37) \quad (1-t)f_i(x) + t\tilde{h}_i(x) = \sigma \quad \text{when } x \in M \setminus U(3\theta), \quad 0 \leq t \leq 1.$$

Indeed, $\tilde{h}_i(x) = \sigma$ when $x \in M \setminus U(3\theta)$, because of (10.34) and (10.35), and $f_i(x) = \sigma$ when $x \in M \setminus U$ by our original assumptions on f_i . This yields (10.37).

Next let us check that

$$(10.38) \quad |f_i(x) - \tilde{h}_i(x)| \leq \epsilon \quad \text{when } x \in M.$$

We may restrict our attention to $x \in U(1/2)$, since $f_i(x) = \tilde{h}_i(x) = \sigma$ when $x \in M \setminus U(1/2)$. If $x \in U(1/2)$, then

$$(10.39) \quad \begin{aligned} |f_i(x) - \tilde{h}_i(x)| &= |f_i(x) - h_i(\phi(x))| \\ &\leq |f_i(x) - f_i(\psi(\phi(x)))| + |f_i(\psi(\phi(x))) - h_i(\phi(x))|. \end{aligned}$$

We already know that $|f_i(\psi(\phi(x))) - h_i(\phi(x))| \leq \epsilon/2$, because of (10.29). As for the first term on the right, we have that

$$(10.40) \quad |f_i(x) - f_i(\psi(\phi(x)))| \leq \nu(d_M(x, \psi(\phi(x)))) \leq \sup_{0 \leq t \leq \theta} \nu(t) \leq \frac{\epsilon}{2},$$

because of our assumption that f_i has modulus of continuity ν , and because of (10.22). Combining these estimates we get (10.38), as desired.

From (10.38) we obtain that

$$(10.41) \quad (1-t)f_i(x) + t\tilde{h}_i(x) \in \widehat{E} \quad \text{when } x \in M, 0 \leq t \leq 1.$$

That is, $(1-t)f_i(x) + t\tilde{h}_i(x)$ is at distance $\leq \epsilon$ from $f_i(x) \in E$, and so (10.41) follows from the definition of $\widehat{E}$. From (10.41) we obtain that $H_i(x, t)$ as defined in (10.33) does make sense and is continuous, because of (10.2).

Observe that

$$(10.42) \quad H_i(x, t) = \sigma \quad \text{when } x \in M \setminus U(3\theta), 0 \leq t \leq 1.$$

This follows from (10.37) and the fact that $R(\sigma) = \sigma$.

We have that $H_i(x, 0) = f_i(x)$ for all $x \in M$ because of the definition (10.36) of H_i and the fact that R equals the identity on E , and hence on the image of f_i . We also want to know that $H_i(x, 1) = \tilde{g}_i(x)$ for all $x \in M$. When $x \in M \setminus U(1/2)$ this is true because both sides equal σ , by (10.42) and (10.26). When $x \in U(1/2)$ we use the definition (10.36) of H_i and the definition (10.34) of $\tilde{h}_i$ to get that $H_i(x, 1) = R(\tilde{h}_i(x)) = R(h_i(\phi(x)))$. This is the same as $g_i(\phi(x)) = \tilde{g}_i(x)$, by the definition of g_i (just after (10.31)) and (10.26).

It remains to show that the modulus of continuity of H_i is uniformly bounded. The argument is practically the same as for g_i . The definition of H_i involves R only on the compact set $\widehat{E}$, and R is uniformly continuous on that set, with a modulus of continuity that only depends on E and its entourage. We are assuming a bound on the modulus of continuity of f_i (to wit, ν), and we have already observed in the proof of part (d) that the h_i 's are Lipschitz with bounded norm. For the present argument we need to know that the $\tilde{h}_i$'s are Lipschitz with uniformly bounded norm. This is true on $U(1/2)$ because of the definition (10.34) of $\tilde{h}_i$ and the Lipschitz bounds for h_i and ϕ (from Lemma 10.19). The Lipschitz bound for $\tilde{h}_i$ on all of M follows from (10.35) and the fact that θ is small.

This completes the proof of Lemma 10.25. (To be truly thorough we ought to go back and write down the μ that we get, but we won't bother with that.)

From now on we let $g_0, g_1 : P \rightarrow E$ be as in Lemma 10.25.

Lemma 10.43. *g_0 and g_1 are homotopic through continuous mappings from P into E which take only the value σ on K .*

We already know that g_i is homotopic to $f_i \circ \psi$ through continuous mappings from P into E which take only the value σ on K , by Lemma 10.25 (c). Thus it suffices to know that $f_0 \circ \psi$ and $f_1 \circ \psi$ are homotopic through continuous mappings from P into E which take only the value σ on K . By hypothesis $f_0, f_1 : M \rightarrow E$ are homotopic through mappings which take only the value σ on $M \setminus U$. This implies what we want, since $\psi(K) \subseteq M \setminus U$ by (10.24). This proves Lemma 10.43.

Lemma 10.44. *There is a homotopy between g_0 and g_1 which takes only the value σ on K and which also satisfies a uniform bound on its modulus of continuity. That is, there exist a function $\omega_0(t) : [0, \infty) \rightarrow [0, \infty)$ - depending only on E (and its entourage), ν , the covering*

function $N(\cdot)$, and the contractability function $\rho(\cdot)$ – and a mapping $G : P \times [0, 1] \rightarrow E$ such that $\omega_0(0) = 0$, $\omega_0(t)$ is continuous at the origin,

$$(10.45) \quad G(p, 0) = g_0(p) \text{ and } G(p, 1) = g_1(p) \text{ for all } p \in P,$$

$$(10.46) \quad G(p, t) = \sigma \text{ when } p \in K \text{ and } t \in [0, 1], \text{ and}$$

$$(10.47) \quad |G(p, t) - G(q, s)| \leq \omega_0(d_P(p, q) + |s - t|) \\ \text{for all } p, q \in P, s, t \in [0, 1].$$

To prove this we apply Lemma 10.16 with X replaced by P and f_i replaced by g_i . Note that Lemma 10.25 supplies the bounds for the moduli of continuity of g_0 and g_1 that we need to apply Lemma 10.16, while Lemma 10.43 supplies the initial homotopy. From Lemma 10.16 we get a homotopy G as above, except that the modulus of continuity depends on P and K . However, there are only finitely many possibilities for P and K (see Lemma 10.19 and (10.24)), and so we can get a uniform choice of ω_0 that works for all of them. (There are only finitely many possibilities for K because it is a union of distinguished simplices in P , and there are only finitely many of those.) This proves Lemma 10.44.

From now on we let G be as in Lemma 10.44.

Lemma 10.44 provides us with a homotopy between g_0 and g_1 which is like the one that we want for the conclusions of Theorem 10.3. It remains to convert it into a suitable homotopy between f_0 and f_1 . We start by pulling G back to M .

Define $\tilde{G} : M \times [0, 1] \rightarrow E$ by

$$(10.48) \quad \begin{aligned} \tilde{G}(x, t) &= G(\phi(x), t) && \text{when } x \in U(1/2), 0 \leq t \leq 1, \\ &= \sigma && \text{when } x \in M \setminus U(1/2), 0 \leq t \leq 1. \end{aligned}$$

Lemma 10.49. $\tilde{G}$ is continuous, and

$$(10.50) \quad \tilde{G}(x, t) = \sigma \quad \text{when } x \in M \setminus U(3\theta), 0 \leq t \leq 1.$$

In fact, there is a function $\omega_1(t) : [0, \infty) \rightarrow [0, \infty)$, depending only on E (and its entourage), ν , the covering function $N(\cdot)$, and the contractability function $\rho(\cdot)$, such that $\omega_0(0) = 0$, $\omega_1(t)$ is continuous at the origin, and

$$(10.51) \quad |\tilde{G}(p, t) - \tilde{G}(q, s)| \leq \omega_1(d_P(p, q) + |s - t|) \\ \text{for all } p, q \in P, s, t \in [0, 1].$$

Let us first check (10.50). This is practically the same as (10.32). As in (10.33), we have that $\phi(x) \in K$ when $x \in U(1/2) \setminus U(3\theta)$. This implies (10.50), because of (10.48) and (10.46).

The continuity of $\tilde{G}$ follows immediately from (10.50). To get the bound on the modulus of continuity we use the corresponding bound (10.47) for G and the fact that ϕ is Lipschitz

with a bounded norm. (See Lemma 10.19.) These bounds imply a controlled modulus of continuity for G when we restrict ourselves to $x \in U(1/2)$, and then (10.50) permits us to deal with all $x \in M$ at once.

This proves Lemma 10.49.

We can now finish the proof of Theorem 10.3. We need to build a homotopy $F : M \times [0, 1] \rightarrow E$ between f_0 and f_1 which takes only the value σ on $M \setminus U_1$ and which has a uniformly bounded modulus of continuity. To do this we combine the homotopies H_0 , $\tilde{G}$, and H_1 in the obvious manner. That is, H_0 provides a homotopy between f_0 and $\tilde{g}_0$, $\tilde{G}$ provides a homotopy between $\tilde{g}_0$ and $\tilde{g}_1$ (by (10.48), (10.26), and (10.45)), and H_1 gives a homotopy from $\tilde{g}_1$ to f_1 . This gives a homotopy F between f_0 and f_1 , i.e., F satisfies (10.4). Each of these intermediate homotopies take only the value σ when $x \in M \setminus U(3\theta)$, because of Lemma 10.25 (f) and (10.50), and so the same is true of F . This implies (10.5). The uniform modulus of continuity condition (10.6) follows from the corresponding bounds for H_i and $\tilde{G}$ in Lemmas 10.25 (f) and 10.49.

This completes the proof of Theorem 10.3.

As discussed in the beginning of this section, Theorem 1.30 follows from Theorem 10.3 and Lemma 10.7, and so the proof of Theorem 1.30 is complete now as well.

Remark 10.52. In passing from Theorem 1.30 to Theorem 10.3 we strengthened the result in some ways, because we asked for much less information about our source and target spaces. We could also try to strengthen it in a very different way. We could require, as in Theorem 1.30, that our target space be $\mathbf{S}^n$ and our metric space M have topological dimension $\leq n$, and then try to use this extra information to weaken our other assumptions. In particular we would like to weaken the assumption of a bound on the contractability function $\rho(\cdot)$, as in Remark 6.10.

Without these additional conditions Theorem 10.3 lives in the world of homotopy, and the required bound on the contractability function is reasonable. However, if we do assume that our target space is $\mathbf{S}^n$ and our metric space M has topological dimension $\leq n$, then homotopy classes of maps from M into $\mathbf{S}^n$ which are constant on the complement of a compact set can be seen in terms of cohomology. (See Theorem VIII 2 on p.149 of [HW], although that deals with slightly different circumstances.) This makes them more flexible than homotopy classes are in general.

We used the bound on the local contractability function $\rho(\cdot)$ in the proof of Theorem 10.3 only in Lemma 10.19, to get the polyhedron P and the mappings $\phi : U(1/2) \rightarrow P$ and $\psi : P \rightarrow U(3/4)$. Remember that Lemma 10.19 was really just a reformulation of part of Lemma 8.3. Let us think about P , ϕ , and ψ , how exactly we got them and how they were used.

The polyhedron P and the mapping ϕ were indispensable for our argument, they gave us our uniform bounds. P and ϕ came originally from Theorem 7.1 (via Lemmas 8.3 and 8.2), which required a bound on the covering function $N(\cdot)$ and on a topological dimension function. In Lemma 10.19 above there was no special dimension n , and so we could just use the requirement that $N(t) = O(t^{-k})$ as $t \rightarrow 0$ for some k to get a bound on the topological dimension function for the dimension k , as in Proposition 6.4 or Theorem 6.5 (with the aid of Corollary 4.7). If we want to try to get some advantage from the cohomological analysis of homotopy classes when M has topological dimension $\leq n$ and

the target space is $\mathbf{S}^n$, then we do have a special dimension n again, and we need to choose P so that it has dimension $\leq n$. This means that we have to be able to control a topological dimension function with dimension n . In Lemma 8.3 we had a bound on the topological dimension function from Lemma 8.1 and Theorem 6.5, and it relied on our bound for the local contractability function. However, this is the only way in which F and f depend on the local contractability function, and there are other reasonable ways to control the topological dimension function without it, as discussed in Remark 6.10.

What about the mapping ψ in Lemma 10.19? It also came from Lemma 8.3, and it was obtained through Proposition 5.4, which required a bound on the contractability function $\rho(\cdot)$. More precisely, we needed to bound the local contractability function in order to get a mapping ψ which was continuous. If we are willing to forgo the continuity of ψ , and ask only for the bounds (10.21) and (10.22), then we can get ψ for free, and by a much simpler argument.

Of course we needed ψ to be continuous in the argument above, but how exactly did we use the continuity of ψ ? We used ψ to define K in (10.24), but this does not use the continuity of ψ . We used ψ in the definition of h_i in the beginning of the proof of Lemma 10.25. This use of ψ is basically trivial, in that we needed ψ to be defined only on the vertices of P , for which the continuity of ψ is irrelevant. The definition of g_i just below (10.31) required ψ only through h_i , and hence it did not use the continuity of ψ either. The continuity properties of h_i and g_i did not depend at all on the continuity of ψ .

Part (c) of Lemma 10.25 does depend on the continuity of ψ . Otherwise we would not know that $f_i \circ \psi$ is continuous, and the homotopy condition would not make sense. Lemma 10.43 also required the continuity of ψ , both in order to use Lemma 10.25 (c) and again in its own proof. We did not use the continuity of ψ again after that, and Lemma 10.25 (c) was used only to obtain Lemma 10.43.

Thus the continuity of ψ was needed only to obtain Lemma 10.43, i.e., to show that g_0 was homotopic to g_1 through maps on P that take only the value σ on K . If we are now assuming that M and P have topological dimension $\leq n$ and that our target space is $\mathbf{S}^n$, then this statement is really a statement about cohomology. As such it ought to require less than the continuous approximate inverse ψ of ϕ provided by Lemma 10.19 in order to be derived from the corresponding homotopy between f_0 and f_1 on M that was part of our original assumptions. (Actually, we should also modify (10.24) to make K a little smaller. This is not a serious issue, we have lots of room, because θ is small.) Remember from Lemma 10.25 (f) that $\tilde{g}_i$ and f_i are homotopic, with suitable bounds. Thus $\tilde{g}_0$ and $\tilde{g}_1$ are homotopic, and we simply need to be able to conclude from this that g_0 and g_1 are homotopic, with restrictions in both cases about where the mappings must take the value σ . This needs less “invertibility” from ϕ than the existence of a continuous choice of ψ .

11. Two summarizing theorems.

The proofs of Theorems 1.29 and 1.30 were completed in Sections 8, 9, and 10, and in Part II below we shall derive Theorems 1.9 and 1.11 from them. In this section we give some consequences of Theorems 1.29 and 1.30 which will be more convenient for the proofs of Theorems 1.9 and 1.11.

Theorem 11.1. *Suppose that $(M, U, d(x, y))$ is as in Standard Assumptions 1.8. Fix a point $\sigma \in \mathbf{S}^n$, and set $\text{diam}_M U = r$. Let B_0 and B_1 be two balls contained in U with radius δr , where δ is some positive number. Then there is a $C' r^{-1}$ -Lipschitz mapping $F : M \times [0, r] \rightarrow \mathbf{S}^n$ such that*

$$(11.2) \quad F(x, t) = \sigma \text{ when } x \in M \setminus U_1 \text{ and } 0 \leq t \leq r,$$

$$(11.3) \quad F(x, t) = \sigma \text{ when } t = 0 \text{ and } x \in M \setminus B_0, \text{ and when } t = r \text{ and } x \in M \setminus B_1,$$

$$(11.4) \quad \text{each } f_t = F(\cdot, t) : M \rightarrow \mathbf{S}^n, 0 \leq t \leq r, \text{ has nonzero degree and cannot be homotoped to a constant through maps that equal } \sigma \text{ on } M \setminus U_1.$$

Here U_1 is as in Definition 1.5, and C' depends only on n, δ , and the functions $N(\cdot)$ and $\rho(\cdot)$ from Standard Assumptions 1.8.

Theorem 11.1 follows easily by combining our earlier results. Let $(M, U, d(x, y))$, σ , B_0 , and B_1 be given as above. We start by applying Theorem 1.29 (a) to get two mappings $f_0, f_1 : M \rightarrow \mathbf{S}^n$ with $f_i \equiv \sigma$ on $M \setminus B_i$ such that each f_i is Lipschitz with norm $\leq C' r^{-1}$ and has nonzero degree. Theorem 1.29 (b) implies that the degrees of these mappings are uniformly bounded, and so we can reduce to the case where they have the same degree by composing them with auxiliary (smooth) mappings on $\mathbf{S}^n$. (This uses basic facts about the degree, as in Lemmas A.21 and A.26. Note that we need only a finite family of auxiliary mappings on $\mathbf{S}^n$ here — which does not depend on the particular choice of f_i — because of the bound on the degrees, and so this composition will not destroy our Lipschitz bounds.) Once we have f_0 and f_1 we can get F as above by applying Theorem 1.30. The Lipschitz bound for F comes from (1.33), the properties (11.2) and (11.3) come from (1.32) and (1.31), and (11.4) follows from the fact that f_0 and f_1 have nonzero degree and are therefore homotopically nontrivial (as in Lemma A.21).

Theorem 11.5. *Suppose that $(M, U, d(x, y))$, x_0, x_1 , and δ are as in Standard Assumptions 1.10. Fix a point $\sigma \in \mathbf{S}^n$, and set $\text{diam}_M U = R$. Then there is a mapping $F : M \times (0, R) \rightarrow \mathbf{S}^n$ with the following properties:*

$$(11.6) \quad F \text{ is Lipschitz with norm } \leq C' \epsilon^{-1} \text{ on } M \times (\epsilon, R - \epsilon) \text{ when } 0 < \epsilon < R;$$

$$(11.7) \quad F(z, t) = \sigma \text{ when } z \in M \setminus U_1, 0 < t < R,$$

$$\text{and when } d(z, x_0) > C't, \text{ and when } d(z, x_1) > C'(R - t);$$

$$(11.8) \quad \text{each } f_t = F(\cdot, t) : M \rightarrow \mathbf{S}^n, 0 < t < R, \text{ has nonzero degree and cannot be homotoped to a constant through maps that all equal } \sigma \text{ on } M \setminus U_1.$$

Here U_1 is as in Definition 1.5, and C' depends only on n, δ , and the functions $N(\cdot)$ and $\rho(\cdot)$ from Standard Assumptions 1.10.

To prove this we shall combine sequences of mappings which are obtained as in Theorem 11.1.

The assumptions and conclusions of Theorem 11.5 are scale-invariant, and so we assume from now on the normalization

$$(11.9) \quad R = \text{diam } U = 1.$$

We first address a technical point.

Lemma 11.10. *Same assumptions as in Theorem 11.5. Given $s \in (0, 1]$ and $i = 0, 1$, let $U(x_i, s)$ denote the connected component of $B(x_i, s\delta)$ which contains x_i . Then there is a small constant $a \in (0, 1/10)$ such that*

$$(11.11) \quad B(x_i, a s \delta) \subseteq U(x_i, s) \subseteq B(x_i, s \delta),$$

$$(11.12) \quad a s \delta \leq \text{diam } U(x_i, s) \leq 2 s \delta,$$

and such that $(M, U(x_i, s), (s\delta)^{-1} d(\cdot, \cdot))$ satisfies Standard Assumptions 1.8 (with the same $N(\cdot)$ and $\rho(\cdot)$ from our original assumptions in Theorem 11.5). This constant a depends only on $\rho(\cdot)$.

The second inclusion in (11.11) is automatic. The first holds if a is small enough because $(M, B(x_i, s\delta), (s\delta)^{-1} d(\cdot, \cdot))$ has contractability function $\rho(\cdot)$, as in Standard Assumptions 1.10. Choose and fix such an $a \in (0, 1/10)$.

The second inequality in (11.12) follows immediately from the second inclusion in (11.11). The first inequality almost follows from the first inclusion in (11.11), but there is a small issue. We need to know that $\{z \in M : d(z, x_i) = r\} \neq \emptyset$ when $0 < r < a s \delta$. Actually this is true for all $0 < r < \frac{1}{2}$, which includes the preceding range by our assumptions on a , s , and δ . To see this we observe that $\{z \in M : d(z, x_i) = r\} = \emptyset$ implies that $U \subseteq B(x_i, r)$, since U is connected and contains x_i by assumption. Thus $\{z \in M : d(z, x_i) = r\} = \emptyset$ implies that $\text{diam } U \leq 2r$, which contradicts (11.9) when $r < \frac{1}{2}$. Therefore $\{z \in M : d(z, x_i) = r\} \neq \emptyset$ when $0 < r < \frac{1}{2}$, and we can derive the first inequality in (11.12) from the first inclusion in (11.11).

Next let us check that $(M, U(x_i, s), (s\delta)^{-1} d(\cdot, \cdot))$ is an admissible triple (in the sense of Definition 1.5). This is a matter of chasing definitions. Let $U_1(x_i, s)$ be the analogue of U_1 for $(M, U(x_i, s), (s\delta)^{-1} d(\cdot, \cdot))$, so that $U_1(x_i, s)$ is the set of points in M whose $(s\delta)^{-1} d(\cdot, \cdot)$ -distance from $U(x_i, s)$ is less than the $(s\delta)^{-1} d(\cdot, \cdot)$ -diameter of $U(x_i, s)$. Thus

$$(11.13) \quad U_1(x_i, s) = \{z \in M : \text{dist}(z, U(x_i, s)) < \text{diam } U(x_i, s)\},$$

where these distances and diameters are taken with respect to $d(\cdot, \cdot)$. Observe that $U(x_i, s) \subseteq U$, because of (11.11) and $B(x_i, \delta) \subseteq U$ (by Standard Assumptions 1.10). Hence $\text{diam } U(x_i, s) \leq \text{diam } U$, and so $U_1(x_i, s) \subseteq U_1$. This implies that $U_1(x_i, s)$ has compact closure in M , since the same is assumed to be true of U_1 . Therefore $(M, U(x_i, s), (s\delta)^{-1} d(\cdot, \cdot))$ is an admissible triple.

Thus $(M, U(x_i, s), (s\delta)^{-1} d(\cdot, \cdot))$ satisfies the first part of Standard Assumptions 1.8. The connectedness of $U(x_i, s)$ is automatic from its definition, and M is an oriented topological manifold by our hypotheses. It remains to check that $(M, U(x_i, s), (s\delta)^{-1} d(\cdot, \cdot))$ has covering function $N(\cdot)$ and contractability function $\rho(\cdot)$. This follows immediately from (11.11) and Standard Assumptions 1.10.

This completes the proof of Lemma 11.10.

From now on we let a be as in Lemma 11.10.

The next lemma provides us with the “slices” that we need for Theorem 11.5.

Lemma 11.14. *Same assumptions as in Theorem 11.5. Given $s \in (0, a]$ and $i = 0, 1$ there is a mapping $f^{i,s} : M \rightarrow \mathbf{S}^n$ such that*

$$(11.15) \quad f^{i,s} \text{ is } C(s\delta)^{-1}\text{-Lipschitz.}$$

$$(11.16) \quad f^{i,s} \equiv \sigma \text{ on } M \setminus B(x_i, s\delta),$$

$$(11.17) \quad f^{i,s} \text{ has nonzero degree on } M.$$

In fact we may choose the $f^{i,s}$'s so that they all have the same degree. The constant C here depends only on n and the functions $N(\cdot)$ and $\rho(\cdot)$ from Standard Assumptions 1.10.

Let us set aside the issue of making the degrees be equal for the moment and just look for mappings which satisfy (11.15) - (11.17) and which have bounded degree. To get $f^{i,s}$ we apply Theorem 1.29 with $(M, U, d(\cdot, \cdot))$ replaced with $(M, U(x_i, a^{-1}s), (a^{-1}s\delta)^{-1}d(\cdot, \cdot))$ and B taken to be $B(x_i, s\delta)$. We may apply Theorem 1.29 with these choices because of Lemma 11.10 (with s replaced with $a^{-1}s$). For this application of Theorem 1.29 we choose the δ in Theorem 1.29 to be $s\delta(\text{diam } U(x_i, a^{-1}s))^{-1}$ (and not the δ from the hypotheses of Theorem 11.5), and this number lies in $[a/2, 1]$ because of (11.12). The mapping $f^{i,s}$ provided by Theorem 1.29 is Lipschitz with respect to $(a^{-1}s\delta)^{-1}d(\cdot, \cdot)$ with norm at most a bounded constant times the reciprocal of the $(a^{-1}s\delta)^{-1}d(\cdot, \cdot)$ -diameter of $U(x_i, a^{-1}s)$. This is the same as saying that $f^{i,s}$ is Lipschitz with respect to $d(\cdot, \cdot)$ with norm at most a constant times the reciprocal of the $d(\cdot, \cdot)$ -diameter of $U(x_i, a^{-1}s)$. This implies (11.5), because of (11.12). We also get uniform bounds on the degrees of these mappings from Theorem 1.29 (b). Of course we get (11.16) and (11.17) from Theorem 1.29 (a) too.

Once we have mappings $f^{i,s}$ which satisfy (11.15)-(11.17) and which have bounded degree, we can modify them so that they all have the same degree by composing them with auxiliary (smooth) mappings on $\mathbf{S}^n$ (using Lemmas A.21 and A.26). Our bounds on the degrees ensure that we only need finitely many auxiliary mappings, and so these modifications will not destroy our Lipschitz bounds. This proves Lemma 11.14.

From now on we let $f^{i,s} : M \rightarrow \mathbf{S}^n$ be as in Lemma 11.14.

Next we fill in the slices.

Lemma 11.18. *Same assumptions as in Theorem 11.5. Given $s \in (0, a]$ and $i = 0, 1$ there is a mapping $F^{i,s} : M \times [s/3, s] \rightarrow \mathbf{S}^n$ such that*

$$(11.19) \quad F^{i,s}(\cdot, s) = f^{i,s} \text{ and } F^{i,s}(\cdot, s/3) = f^{i,t}, \quad t = s/3.$$

$$(11.20) \quad F^{i,s} \text{ is } C s^{-1}\text{-Lipschitz.}$$

$$(11.21) \quad F^{i,s}(z, \tau) = \sigma \text{ when } z \in M \setminus (B(x_i, 3a^{-1}s\delta) \cap U_1) \text{ and } \tau \in [s/3, s].$$

Also, there is a mapping $F^a : M \times [\frac{1}{3}, \frac{2}{3}] \rightarrow \mathbf{S}^n$ such that

$$(11.22) \quad F^a(\cdot, \frac{1}{3}) = f^{0,a} \text{ and } F^a(\cdot, \frac{2}{3}) = f^{1,a},$$

$$(11.23) \quad F^a \text{ is } C\text{-Lipschitz.}$$

$$(11.24) \quad F^a(z, \tau) = \sigma \text{ when } z \in M \setminus U_1 \text{ and } \tau \in [\frac{1}{3}, \frac{2}{3}].$$

This constant C depends only on n , δ , and the functions $N(\cdot)$ and $\rho(\cdot)$ from Standard Assumptions 1.10.

These odd-looking parameter intervals $[s/3, s]$ and $[\frac{1}{3}, \frac{2}{3}]$ are chosen for their convenience for the application, nothing more. The 3 in the $B(x_i, 3a^{-1}s\delta)$ has nothing to do with the other 3's.

To prove Lemma 11.18 we simply need to apply Theorem 1.30 and unwind the definitions. Let us begin with the mapping F^a in the second half of the lemma. For this we can apply Theorem 1.30 directly, with $f_0 = f^{0,a}$, $f_1 = f^{1,a}$, and the replacement of $[0, 1]$ with $[\frac{1}{3}, \frac{2}{3}]$.

Now consider $F^{i,s}$, as in the first part of the lemma. For this we apply Theorem 1.30 with $(M, U, d(\cdot, \cdot))$ replaced with $(M, U(x_i, a^{-1}s), (a^{-1}s\delta)^{-1}d(\cdot, \cdot))$ and $f_0 = f^{i,s}$, $f_1 = f^{i,t}$, $t = s/3$. Both of these mappings take only the value σ outside $U(x_i, a^{-1}s)$, because of (11.16) and (11.11). This enables us to apply Theorem 1.30 in this way, and we get a mapping that we shall denote by $G^{i,s} : M \times [0, r] \rightarrow \mathbf{S}^n$, where r is the diameter of $U(x_i, a^{-1}s)$ with respect to the metric $(a^{-1}s\delta)^{-1}d(\cdot, \cdot)$, as in Theorem 1.30.

We need to be a little careful in deriving information from Theorem 1.30, because we are changing our metric here. Let us compute r , the diameter of $U(x_i, a^{-1}s)$ with respect to the metric $(a^{-1}s\delta)^{-1}d(\cdot, \cdot)$. We have that

$$(11.25) \quad r = (a^{-1}s\delta)^{-1} \text{diam} U(x_i, a^{-1}s) \quad \text{and} \quad a \leq r \leq 2,$$

because of (11.12). In other words r is comparable in size to 1, modulo quantities that we can bound.

Set $F^{i,s}(z, \tau) = G^{i,s}(z, A(\tau))$, where $A : \mathbf{R} \rightarrow \mathbf{R}$ is the affine function such that $A(s/3) = 0$ and $A(s) = r$. Thus $F^{i,s} : M \times [s/3, s] \rightarrow \mathbf{S}^n$ satisfies (11.19), because of (1.31).

Next, (1.32) translates into

$$(11.26) \quad F^{i,s}(z, \tau) = \sigma \quad \text{when } z \in M \setminus U_1(x_i, a^{-1}s) \text{ and } \tau \in [s/3, s],$$

where $U_1(x_i, a^{-1}s)$ is the analogue of U_1 for $(M, U(x_i, a^{-1}s), (a^{-1}s\delta)^{-1}d(\cdot, \cdot))$. Equivalently, $U_1(x_i, a^{-1}s)$ is given by (11.13). We have that

$$(11.27) \quad U_1(x_i, a^{-1}s) \subseteq B(x_i, 3a^{-1}s\delta),$$

because of (11.13) and (11.12). We also have that $U_1(x_i, a^{-1}s) \subseteq U_1$, as observed just after (11.13). These observations together with (11.26) yield (11.21).

It remains to verify (11.20). We know from (1.33) and (11.25) that $G^{i,s} : M \times [0, r] \rightarrow \mathbf{S}^n$ is Lipschitz with bounded norm, at least if we use the metric $(a^{-1}s\delta)^{-1}d(\cdot, \cdot)$ on M and the Euclidean metric on $[0, r]$. This is the same as saying that $G^{i,s}$ is Lipschitz with bounded norm if we use the metric $s^{-1}d(\cdot, \cdot)$ on M and the Euclidean metric on $[0, r]$, because our constants are allowed to depend on a and δ . This implies that $F^{i,s}$ is Cs^{-1} -Lipschitz on $M \times [s/3, s]$, where we use the metric $d(\cdot, \cdot)$ on M and the Euclidean metric on $[s/3, s]$, because of the definition of $F^{i,s}$ in terms of $G^{i,s}$. This is the same as (11.20).

This completes the proof of Lemma 11.18.

To prove Theorem 11.5 we combine the mappings from Lemma 11.18. Remember our normalization (11.9). Define $F : M \times (0, 1) \rightarrow \mathbf{S}^n$ by
(11.28)

$$\begin{aligned} F(z, t) &= F^a(z, t) && \text{when } t \in [\frac{1}{3}, \frac{2}{3}] \\ &= F^{0,s}(z, t) && \text{when } t \in [3^{-j-1}, 3^{-j}], \text{ with } s = a 3^{-j+1}, j = 1, 2, 3, \dots \\ &= F^{1,s}(z, 1-t) && \text{when } t \in [1 - 3^{-j}, 1 - 3^{-j-1}], \text{ with } s = a 3^{-j+1}, j = 1, 2, 3, \dots \end{aligned}$$

These formulas are compatible with each other because of (11.22) and (11.19). This compatibility implies that F is continuous.

The Lipschitz bound (11.6) follows from (11.23) and (11.20). To see this it is helpful to observe that it is enough to prove Lipschitz bounds for $F(z, t)$ in z and t separately. The Lipschitz condition in z follows directly from (11.23) and (11.20), while for the Lipschitz bound in t one should also notice that Lipschitz mappings splice together nicely. That is, if g is a Lipschitz mapping on $[a, b] \subseteq \mathbf{R}$, and if h is a Lipschitz mapping on $[b, c]$, and if $g(b) = h(b)$, then the combination of g and h on $[a, c]$ is Lipschitz, with norm at most the maximum of the Lipschitz norms of g and h .

The localization condition (11.7) follows from (11.21) and (11.24).

The topological nontriviality condition (11.8) follows from (11.17) (and Lemma A.21) when $t = 3^{-j}$ or $t = 1 - 3^{-j}$ for some positive integer j . It then follows for all t because F itself provides a homotopy between the f_t 's which takes only the value σ off U_1 .

This completes the proof of Theorem 11.5.

Part II. Applications of Lipschitz Mappings.

12. Two useful reformulations.

Our next main goal is to prove Theorems 1.9 and 1.11. These two results can each be split naturally into two parts, with Theorems 11.1 and 11.5 providing the first parts. In this section we state a pair of theorems which correspond to the second parts.

The conclusions of these theorems involve Definition 1.2, which the reader may wish to review. Recall that $H^n(\cdot)$ denotes n -dimensional Hausdorff measure, and that $H^n|_E$ denotes the restriction of H^n to the set E .

Theorem 12.1. *Let $(M, d(x, y))$ be a metric space, and let V be an open subset of M with compact closure such that $H^n(V) < \infty$. Set $T = \text{diam } V$, and fix a point $\sigma \in \mathbf{S}^n$. Let B_0 and B_1 be a pair of open balls contained in V , and let k be a positive number. Assume that there is a kT^{-1} -Lipschitz mapping $F : M \times [0, T] \rightarrow \mathbf{S}^n$ such that*

$$(12.2) \quad F(x, t) = \sigma \text{ when } x \in M \setminus V \text{ and } 0 \leq t \leq T.$$

$$(12.3) \quad F(x, t) = \sigma \text{ when } t = 0 \text{ and } x \in M \setminus B_0, \text{ and when } t = T \text{ and } x \in M \setminus B_1,$$

$$(12.4) \quad \text{each } f_t = F(\cdot, t) : M \rightarrow \mathbf{S}^n, 0 \leq t \leq T, \text{ cannot be homotoped to a constant through maps that equal } \sigma \text{ on } M \setminus V.$$

Then there is a family of curves in M with parameter space $\mathbf{S}^n$ such that all the curves are contained in V , they all start in $\overline{B}_0$ and end in $\overline{B}_1$, and they are controlled in measure by $\nu = C_0 (\text{diam } V)^{-n+1} H^n \llcorner V$. Here C_0 depends only on n and k . Note that we are implicitly using Lebesgue measure on our parameter space $\mathbf{S}^n$.

Theorem 1.9 will follow once we prove Theorem 12.1, because Theorem 11.1 implies that the hypotheses of Theorem 12.1 hold under the assumptions of Theorem 1.9. (The V in Theorem 12.1 corresponds to the U_1 in Theorems 1.9 and 11.1. Note that $\text{diam } U \leq \text{diam } U_1 \leq 3 \text{diam } U$ when U and U_1 are as in Definition 1.5.)

Theorem 12.5. *Let $(M, d(x, y))$ be a metric space, and let V be an open subset of M with compact closure. Set $T = \text{diam } V$, and fix a point $\sigma \in \mathbf{S}^n$. Let x_0 and x_1 be a pair of points in V , and let k be a positive number. Assume that there is a mapping $F : M \times (0, T) \rightarrow \mathbf{S}^n$ with the following properties:*

$$(12.6) \quad F \text{ is Lipschitz with norm } \leq k\epsilon^{-1} \text{ on } M \times (\epsilon, T - \epsilon) \text{ when } 0 < \epsilon < T;$$

$$(12.7) \quad F(z, t) = \sigma \text{ when } z \in M \setminus V, 0 < t < T.$$

$$\text{and when } d(z, x_0) > kt, \text{ and when } d(z, x_1) > k(T - t);$$

$$(12.8) \quad \text{each } f_t = F(\cdot, t) : M \rightarrow \mathbf{S}^n, 0 < t < T, \text{ cannot be homotoped to a constant through maps that equal } \sigma \text{ on } M \setminus V.$$

Define a measure ν on M by $\nu = h \cdot H^n \llcorner U_1$, where the density h is defined by

$$(12.9) \quad h(y) = (d(y, x_0)^{-n+1} + d(y, x_1)^{-n+1}).$$

(That is, ν is a measure on the H^n -measurable subsets of M , say.) Assume also that $\nu(U_1) < \infty$.

Then there is a family of curves in M with parameter space $\mathbf{S}^n$ such that all the curves are contained in V , they all start at x_0 and end at x_1 , and they are controlled in measure by $C_1 \nu$. Here C_1 depends only on n and k . Note that we are implicitly using Lebesgue measure on our parameter space $\mathbf{S}^n$ (as in Definition 1.2).

In this case we also have that

$$(12.10) \quad x_0 \text{ and } x_1 \text{ can be connected by a curve in } V \text{ of length } \leq C_1 |\mathbf{S}^n|^{-1} \int_V \nu,$$

where $|\mathbf{S}^n|$ denotes the measure of the standard n -sphere.

Theorem 1.11 will follow once we prove Theorem 12.5, because Theorem 11.5 implies that the hypotheses of Theorem 12.5 hold under the assumptions of Theorem 1.11. (Again, the V in Theorem 12.5 corresponds to U_1 in Theorems 1.11 and 11.5.)

Thus to prove Theorems 1.9 and 1.11 we need only prove Theorems 12.1 and 12.5. These theorems are not very novel, and similar issues arose in [DS1]. Their proofs will involve only standard ingredients from topology and geometric measure theory. The more serious point is that the hypotheses of these theorems can be verified in interesting situations.

At this stage, where we are ready to assume the existence of Lipschitz mappings with various properties, the covering and contractability functions in Definitions 1.6 and 1.7

will no longer play a role, nor will Standard Assumptions 1.8 or 1.10, nor will topological dimension or topological manifolds or homology.

In both of Theorems 12.1 and 12.5 we assume the existence of Lipschitz mappings which satisfy certain bounds and topological nontriviality properties, and we conclude that there are many paths in our metric space with certain properties. We shall find these paths in the fibers of the mappings. The next three sections provide the tools for doing this.

13. Some topological lemmas.

As indicated above, we are interested in finding curves in the fibers of certain mappings. As a first step we show in this section that these fibers have the right connectivity properties.

Definition 13.1. If $(M, d(x, y))$ is a metric space and Γ is a closed subset of M , then we say that Γ **connects** two subsets A_0 and A_1 of M if we cannot find disjoint closed subsets E_0, E_1 of M such that $A_0 \subseteq M \setminus E_1$, $A_1 \subseteq M \setminus E_0$, and $\Gamma = E_0 \cup E_1$.

We allow E_0 or E_1 to be empty here. By taking $E_0 = \emptyset$ and $E_1 = \Gamma$, or vice-versa, we see that Γ must intersect each of A_0 and A_1 if it is to connect them.

Notice that if Γ connects A_0 and A_1 , then any closed set which contains Γ also connects A_0 and A_1 .

Lemma 13.2. *Let $(M, d(x, y))$ be a metric space, and let $n \in \mathbf{Z}_+$ and $\sigma \in \mathbf{S}^n$ be given. Let V be a subset of M , let A_0 and A_1 be two subsets of M , and let T be a positive real number. Suppose that $F : M \times [0, T] \rightarrow \mathbf{S}^n$ is a continuous mapping such that*

$$(13.3) \quad F(x, t) = \sigma \text{ when } x \in M \setminus V \text{ and } 0 \leq t \leq T.$$

$$(13.4) \quad F(x, t) = \sigma \text{ when } t = 0 \text{ and } x \in M \setminus A_0, \text{ and when } t = T \text{ and } x \in M \setminus A_1,$$

$$(13.5) \quad \text{each } f_t = F(\cdot, t) : M \rightarrow \mathbf{S}^n, \ 0 \leq t \leq T, \text{ cannot be homotoped to a constant through maps that equal } \sigma \text{ on } M \setminus V.$$

Fix $p \in \mathbf{S}^n$, $p \neq \sigma$, and set $\Gamma(p) = \{z \in M : F(z, t) = p \text{ for some } t \in [0, T]\}$. Then $\Gamma(p)$ is a closed subset of M which connects A_0 and A_1 , and $\Gamma(p) \subseteq V$.

Let $p \in \mathbf{S}^n \setminus \{\sigma\}$ be given. That $\Gamma(p)$ is closed is not hard to check, using the continuity of F and the compactness of $[0, T]$. We also have $\Gamma(p) \subseteq V$ automatically from (13.3).

It remains to show that $\Gamma(p)$ connects A_0 and A_1 . Suppose, to the contrary, that this is not the case. This means that we can find disjoint closed subsets E_0, E_1 of M such that $A_0 \subseteq M \setminus E_1$, $A_1 \subseteq M \setminus E_0$, and $\Gamma(p) = E_0 \cup E_1$.

Let $h : M \rightarrow [0, T]$ be a continuous function on M such that $h \equiv T$ on E_0 and $h \equiv 0$ on E_1 . For instance, take h to be defined by

$$(13.6) \quad h(z) = T \frac{\text{dist}(z, E_1)}{\text{dist}(z, E_0) + \text{dist}(z, E_1)}.$$

This is well defined and continuous because E_0 and E_1 are closed and disjoint, so that the denominator never vanishes. Define $g : M \rightarrow \mathbf{S}^n$ by $g(z) = F(z, h(z))$. Then $g \equiv \sigma$ on

$M \setminus V$, because of (13.3). In fact, if we set $g_t(z) = F(z, th(z))$, then we get a homotopy between g and f_0 (as defined in (13.5)) through maps which take only the value σ on $M \setminus V$. Therefore g cannot be homotoped to a constant through maps which take only the value σ on $M \setminus V$, because of (13.5).

Let us check now that g never takes the value p . If $z \notin \Gamma(p)$ then certainly $g(z) \neq p$, by definition of $\Gamma(p)$. So suppose that $z \in \Gamma(p)$, so that z either lies in E_0 or E_1 . If $z \in E_0$, then $h(z) = T$, and $g(z) = F(z, T) \neq p$ because $z \notin A_1$ and because of (13.4). Similarly, if $z \in E_1$, then $h(z) = 0$, $g(z) = F(z, 0)$, and this cannot be p because $z \in E_1 \subseteq M \setminus A_0$. Thus g never takes the value p .

Since g never takes the value p it is easy to deform it to a constant through maps which take the value σ whenever g does. Indeed, we can deform the identity mapping on $\mathbf{S}^n \setminus \{p\}$ to a constant through maps which always take the value σ at σ . (This is equivalent to saying that we can deform the identity mapping on the unit ball in $\mathbf{R}^n$ to a constant through mappings which all take the origin to itself. We can do this just by multiplying by t .) By composing with this deformation on $\mathbf{S}^n \setminus \{p\}$ we get a homotopy of g to a constant through mappings which take only the value σ on $M \setminus V$. We already saw that this is impossible, because of (13.5). This gives a contradiction, and Lemma 13.2 follows.

Lemma 13.2 will be used in the proof of Theorem 12.1, and we need a slightly different version for Theorem 12.5.

Lemma 13.7. *Let $(M, d(x, y))$ be a metric space, and let $n \in \mathbf{Z}_+$, and $\sigma \in \mathbf{S}^n$ be given. Let V be a subset of M and let x_0 and x_1 be points in V . Suppose that $F : M \times (0, T) \rightarrow \mathbf{S}^n$ is a continuous mapping which satisfies (12.7) (for some $k > 0$) and (12.8). Fix $p \in \mathbf{S}^n$, $p \neq \sigma$, and set $\Gamma(p) = \{z \in M : F(z, t) = p \text{ for some } t \in (0, T)\} \cup \{x_0, x_1\}$. Then $\Gamma(p)$ is a closed subset of M which connects $\{x_0\}$ and $\{x_1\}$ and which is contained in V .*

That $\Gamma(p) \subseteq V$ follows from (12.7) and our other assumptions.

Let us check that $\Gamma(p)$ is closed. Given $\epsilon > 0$, let $\Gamma_\epsilon(p)$ be defined in the same manner as $\Gamma(p)$ was, except that we restrict ourselves to $t \in [\epsilon, T - \epsilon]$. It is not hard to show that $\Gamma_\epsilon(p)$ is closed for each $\epsilon > 0$, because of the compactness of the interval $[\epsilon, T - \epsilon]$. (This is the same as the corresponding step for Lemma 13.2.) The closedness of $\Gamma(p)$ follows from this observation and (12.7), which controls the behavior of $F(z, t)$ as $t \rightarrow 0$ or $t \rightarrow T$.

It remains to check that $\Gamma(p)$ connects $\{x_0\}$ and $\{x_1\}$. Suppose that this were not true. This would mean that there exist disjoint closed subsets E_0 and E_1 of M such that $x_0 \notin E_1$, $x_1 \notin E_0$, and $\Gamma(p) = E_0 \cup E_1$. Since E_0 and E_1 are closed we get that $\overline{B}(x_0, k\epsilon) \subseteq M \setminus E_1$ and $\overline{B}(x_1, k\epsilon) \subseteq M \setminus E_0$ for some small $\epsilon > 0$. This means that $\Gamma(p)$ does not connect $\overline{B}(x_0, k\epsilon)$ and $\overline{B}(x_1, k\epsilon)$. However, Lemma 13.2 implies that $\Gamma(p)$ does connect $\overline{B}(x_0, k\epsilon)$ and $\overline{B}(x_1, k\epsilon)$. That is, we apply Lemma 13.2 to the restriction of F to $M \times [\epsilon, T - \epsilon]$, with $A_i = \overline{B}(x_i, k\epsilon)$. The assumptions of Lemma 13.7 imply that this restriction satisfies the hypotheses of Lemma 13.2, and so we are able to apply Lemma 13.2 here. The set $\Gamma(p)$ associated to this restriction of F is not the same as the one that we are using, but it is contained in our $\Gamma(p)$, and so we obtain that our $\Gamma(p)$ connects $\overline{B}(x_0, k\epsilon)$ and $\overline{B}(x_1, k\epsilon)$. This gives a contradiction, and we conclude that $\Gamma(p)$ connects $\{x_0\}$ and $\{x_1\}$, as desired.

This completes the proof of Lemma 13.7.

Lemma 13.8. *Let $(M, d(x, y))$ be a metric space, and suppose that Γ is a compact subset of*

M which connects two closed subsets A_0 and A_1 of M . Then there is a compact connected subset Γ' of Γ which intersects both A_0 and A_1 .

A kind reader has pointed out that this is a restatement of the fact that quasicomponents are the same as components in a compact Hausdorff space, at least in the case where A_0 and A_1 are singletons. (See p.47 of [HY].)

To prove the lemma we use the Brouwer reduction method. (See p.161 of [HW].)

Since Γ is compact there is a sequence of open balls $\{B_i\}_{i=0}^\infty$ which induce a basis for the topology of Γ . That is, if $u \in \Gamma$ and $r > 0$, then there is a B_i such that $u \in B_i \subseteq \mathcal{B}(u, r)$. For given u and r there must be infinitely many such choices of i , because we can shrink r as much as we like.

Define a nonincreasing sequence $\{\Gamma_i\}_{i=0}^\infty$ of closed subsets of Γ as follows. Set $\Gamma_0 = \Gamma$. If Γ_i has been chosen, then we take $\Gamma_{i+1} = \Gamma_i \setminus B_i$ if $\Gamma_i \setminus B_i$ connects A_0 and A_1 , and we set $\Gamma_{i+1} = \Gamma_i$ otherwise. By continuing this process indefinitely we obtain our sequence $\{\Gamma_i\}_{i=0}^\infty$. Each Γ_i connects A_0 and A_1 . Set $\Gamma' = \bigcap \Gamma_i$. This set is automatically compact, and it intersects both of A_0 and A_1 , since each Γ_i does (and since A_0 and A_1 are closed).

Suppose that Γ' is not connected, so that it can be written as the union of two disjoint nonempty closed subsets H_+ and H_- . We want to get a contradiction. The idea for this is that our construction of Γ' should make it minimal, while the decomposition of Γ' into $H_+ \cup H_-$ should enable us to violate this minimality. Let U_+ and U_- be disjoint open subsets of M such that $H_+ \subseteq U_+$, $H_- \subseteq U_-$. (They can be produced using a continuous function which equals 0 on H_+ and 1 on H_- , as in (13.6).) There is an i such that $\Gamma_i \subseteq U_+ \cup U_-$; otherwise $\{M \setminus \Gamma_i\}_{i=0}^\infty$ together with U_+ and U_- would provide an open cover of Γ without a finite subcover. (Remember that $\{\Gamma_i\}$ is nonincreasing by construction.) Given $j \geq i$ set $\Gamma_j^+ = \Gamma_j \cap U_+ = \Gamma_j \setminus U_-$ and $\Gamma_j^- = \Gamma_j \cap U_- = \Gamma_j \setminus U_+$, so that Γ_j^+ and Γ_j^- are disjoint compact sets whose union is Γ_j .

Claim 13.9. For each $j \geq i$ we have that either Γ_j^+ connects A_0 and A_1 or Γ_j^- connects A_0 and A_1 .

If the claim were not true, then we could decompose Γ_j^+ into the union of two disjoint closed sets, one of which is disjoint from A_0 and the other of which is disjoint from A_1 , and we could do the same for Γ_j^- . These decompositions could then be combined to provide a similar decomposition for Γ_j , which would contradict the fact that Γ_j connects A_0 and A_1 . This proves the claim.

Choose $j \geq i$ so that $B_j \cap H_+ \neq \emptyset$ and $B_j \subset U_+$. We can do this, since $\{B_i\}$ provides a basis for the topology of M . Since $H_+ \subseteq \Gamma'$ we conclude that $\Gamma_j \setminus B_j$ does not connect A_0 and A_1 . (Otherwise, our construction would give $\Gamma_{j+1} = \Gamma_j \setminus B_j$, and Γ' would not be able to contain all of H_+ .) Since $B_j \subset U_+$ we conclude that Γ_j^- does not connect A_0 and A_1 . (Otherwise $\Gamma_j \setminus B_j \supseteq \Gamma_j^-$ would connect A_0 and A_1 .)

Choose $k > j$ so that $B_k \cap H_- \neq \emptyset$ and $B_k \subset U_-$. The same argument as the one that we just used implies that Γ_k^+ does not connect A_0 and A_1 . However, we also have that Γ_k^- does not connect A_0 and A_1 , because $\Gamma_k^- \subseteq \Gamma_j^-$. This contradicts Claim 13.9.

Thus Γ' must be connected. This completes the proof of Lemma 13.8.

14. The coarea estimate.

The main result of this section is a version of the coarea formula. This will give us control on the 1-dimensional Hausdorff measures of the preimages of points of the mappings F in Theorems 12.1 and 12.5. The traditional version for maps between Euclidean spaces is given in Theorem 3.2.12 on p.249 of [Fd]. We shall need a much less precise result, but on more general spaces. We could use Theorem 2.10.25 on p.188 of [Fd], but the present situation can be handled by a simpler argument which we include here for the sake of completeness.

Proposition 14.1. *Let $(N, D(x, y))$ be a metric space, let l be a nonnegative integer, and suppose that $\psi : N \rightarrow \mathbf{S}^n$ is K -Lipschitz. Given any subset A of N there is a Borel measurable function $h : \mathbf{S}^n \rightarrow [0, \infty]$ such that*

$$(14.2) \quad H^l(A \cap \psi^{-1}(y)) \leq h(y) \quad \text{for all } y \in \mathbf{S}^n, \text{ and}$$

$$(14.3) \quad \int_{\mathbf{R}^n} h(y) dy \leq C(n) K^n H^{n+l}(A),$$

where $C(n)$ depends only on n .

Here H^l and H^{n+l} denote l -dimensional and $(n + l)$ -dimensional Hausdorff measure (see (2.14)), respectively, and dy denotes n -dimensional Lebesgue measure on $\mathbf{S}^n$.

Let us prove the proposition. Let N, ψ, l , and A be as above. We may as well assume that $H^{n+l}(A) < \infty$. Let $\delta > 0$ be given, arbitrarily small. We shall first try to prove (14.3) with H^l replaced by H_δ^l (as in (2.13)). Basically we shall simply start with the definition of $H^{n+l}(A)$ and then move sums around.

By definition of $H^{n+l}(A)$ we can cover A with a sequence of sets $\{E_j\}$ in N such that $\text{diam } E_j < \delta$ and

$$(14.4) \quad \sum_j (\text{diam } E_j)^{n+l} \leq H^{n+l}(A) + \delta.$$

Let F_j denote the closure of $\psi(E_j)$, so that

$$(14.5) \quad \text{diam } F_j \leq K \text{diam } E_j.$$

Define $h_\delta(y)$ by

$$(14.6) \quad h_\delta(y) = \sum_j (\text{diam } E_j)^l \mathbf{1}_{F_j}(y),$$

where $\mathbf{1}_{F_j}$ denotes the characteristic function of F_j . The definition of H_δ^l in (4.1) implies that

$$(14.7) \quad H_\delta^l(A \cap \psi^{-1}(y)) \leq h_\delta(y)$$

for all $y \in \mathbf{S}^n$. That is, we can cover $A \cap \psi^{-1}(y)$ by the E_j 's for which $y \in F_j$, so that the sum in (14.6) gives the correct sum for $H_\delta^l(A \cap \psi^{-1}(y))$.

The Lebesgue measure of F_j in $\mathbf{S}^n$ is $\leq C(n)(\text{diam } F_j)^n$. Hence

$$(14.8) \quad \begin{aligned} \int_{\mathbf{S}^n} h_\delta(y) dy &\leq \sum_j (\text{diam } E_j)^l C(n) (\text{diam } F_j)^n \\ &\leq C(n) K^n \sum_j (\text{diam } E_j)^{n+l} \leq C(n) K^n (H^{n+l}(A) + \delta). \end{aligned}$$

This is a good approximation to (14.3).

Set $h(y) = \liminf_{k \rightarrow \infty} h_{\frac{1}{k}}(y)$. This is a Borel measurable function, and we have that

$$(14.9) \quad H^l(A \cap v^{-1}(y)) \leq h(y)$$

for all $y \in \mathbf{S}^n$, because of (14.7) and (2.14). (Remember that the limit in (2.14) exists, because of monotonicity.) Fatou's lemma and (14.8) imply that

$$(14.10) \quad \int_{\mathbf{S}^n} h(y) dy \leq C(n) K^n H^{n+l}(A).$$

This proves Proposition 14.1.

The next lemma provides a version of the coarea estimate that will be more convenient for us to use.

Lemma 14.11. *Let $(M, d(x, y))$ be a metric space and let k, R be nonnegative real numbers. Suppose that $F : M \times [0, R] \rightarrow \mathbf{S}^n$ is kR^{-1} -Lipschitz. Set $\Phi(p) = \{z \in M : F(z, t) = p \text{ for some } t \in [0, R]\}$ for $p \in \mathbf{S}^n$. Given any subset A of M there is a Borel measurable function $h : \mathbf{S}^n \rightarrow [0, \infty]$ such that*

$$(14.12) \quad H^1(A \cap \Phi(p)) \leq h(p) \quad \text{for all } p \in \mathbf{S}^n, \text{ and}$$

$$(14.13) \quad \int_{\mathbf{S}^n} h(p) dp \leq C(n) k^n R^{-n+1} H^n(A)$$

where $C(n)$ depends only on n .

To see this we apply Proposition 14.1, with $N = M \times [0, R]$, A replaced with $A \times [0, R]$, and $l = 1$. We get (14.12) from (14.2) because

$$(14.14) \quad H^1(A \cap \Phi(p)) \leq H^1(\{A \times [0, R]\} \cap F^{-1}(p))$$

(since $A \cap \Phi(p)$ is just the image of $\{A \times [0, R]\} \cap F^{-1}(p)$ under the obvious projection from $M \times [0, R] \rightarrow M$). We get (14.13) from (14.3), with the main point being that we picked up an extra factor of R from the transition from $H^{n+1}(A \times [0, R])$ to $H^n(A)$. This proves Lemma 14.11.

We shall use Lemma 14.11 in the proof of Theorem 12.1, but for Theorem 12.5 we shall need a slightly more complicated result.

Lemma 14.15. *Let $(M, d(x, y))$ be a metric space, let V be an open subset of M , and fix a point $\sigma \in \mathbf{S}^n$. Let x_0 and x_1 be a pair of points in V , and let k, T be nonnegative*

real numbers. Suppose that $F : M \times (0, T) \rightarrow \mathbf{S}^n$ satisfies (12.6) and (12.7), and set $\Phi(p) = \{z \in M : F(z, t) = p \text{ for some } t \in (0, T)\}$ for $p \in \mathbf{S}^n$. Let A be an H^n -measurable subset of M (e.g., a Borel set). Then there is a Borel measurable function $a(p) : \mathbf{S}^n \rightarrow [0, \infty]$ such that

$$(14.16) \quad H^1(A \cap \Phi(p)) \leq a(p) \quad \text{for all } p \in \mathbf{S}^n, \text{ and}$$

$$(14.17) \quad \int_{\mathbf{S}^n} a(p) dp \leq C \int_{A \cap V \cap \overline{B}_M(x_0, kT)} \left\{ \frac{1}{d(x_0, z)^{n-1}} + \frac{1}{d(x_1, z)^{n-1}} \right\} dH^n(z),$$

where C depends only on n and k .

The assumption of measurability of A is not serious, it just permits us to avoid interpreting the right side of (14.17) as an outer integral.

Observe that we may ignore $p = \sigma$ in the integral on the left side of (14.17). This will permit us to ignore large portions of M , because of (12.7). In particular we may simply assume that $V = M$, since $F(z, t) = \sigma$ when $z \in M \setminus V$, by (12.7). Thus we shall forget about V from now on. Similarly, $F(z, t) = \sigma$ for all t when $z \in M \setminus \overline{B}(x_0, kT)$, by (12.7), and so we may assume that $M \setminus \overline{B}(x_0, kT)$ is empty; we can just throw it away. The bottom line is that it suffices to prove (14.17) with the integral on the right side taken over all of A .

We shall prove Lemma 14.15 by applying Lemma 14.11 to the restrictions of F to various pieces of $M \times (0, T)$ and then summing the pieces up again.

Let x, y, T , etc., be as in the lemma. Define intervals I_j^- and I_j^+ by

$$(14.18) \quad I_j^- = [2^{-j-1}T, 2^{-j}T] \quad \text{and} \quad I_j^+ = [(1 - 2^{-j})T, (1 - 2^{-j-1})T],$$

$j = 1, 2, 3, \dots$. Thus $(0, T) = \bigcup_j (I_j^- \cup I_j^+)$. Let $F_j^\pm$ denote the restriction of F to $M \times I_j^\pm$, and set $\Phi_j^\pm(p) = \{z \in M : F(z, t) = p \text{ for some } t \in I_j^\pm\}$. We have that

$$(14.19) \quad \Phi(p) = \bigcup_{j=1}^{\infty} \{\Phi_j^-(p) \cup \Phi_j^+(p)\}$$

by the definition of $\Phi(p)$ (in Lemma 14.15).

We want to apply Lemma 14.11 to each $F_j^\pm$ separately. Let us first observe that

$$(14.20) \quad F_j^\pm \text{ is Lipschitz with norm } \leq k 2^j T^{-1},$$

because of (12.6). Set $B_j^- = \overline{B}(x_0, k 2^{-j}T)$ and $B_j^+ = \overline{B}(x_1, k 2^{-j}T)$. Using (12.7) we get that

$$(14.21) \quad F_j^\pm \equiv \sigma \quad \text{on } (M \setminus B_j^\pm) \times I_j^\pm.$$

We now apply Lemma 14.11 to $F_j^\pm$, with $M = B_j^\pm$, $[0, R]$ replaced with $I_j^\pm$, and A replaced with $(A \cap B_j^\pm) \times I_j^\pm$. We obtain Borel measurable functions $h_j^\pm : \mathbf{S}^n \rightarrow [0, \infty]$ such that

$$(14.22) \quad H^1(A \cap B_j^\pm \cap \Phi_j^\pm(p)) \leq h_j^\pm(p) \quad \text{for all } p \in \mathbf{S}^n, \text{ and}$$

$$(14.23) \quad \int_{\mathbf{S}^n} h_j^\pm(p) dp \leq C(n) k^n 2^{j(n-1)} T^{-n+1} H^n(A \cap B_j^\pm).$$

Notice that

$$(14.24) \quad H^1(A \cap \Phi_j^\pm(p)) = H^1(A \cap \Phi_j^\pm(p) \cap B_j^\pm) \quad \text{for all } p \in \mathbf{S}^n \setminus \{\sigma\},$$

because of (14.21). Thus (14.22) is really an estimate for $H^1(A \cap \Phi_j^\pm(p))$, except when $p = \sigma$.

Define $a(p) : \mathbf{S}^n \rightarrow [0, \infty]$ by

$$(14.25) \quad \begin{aligned} a(p) &= \sum_{j=1}^{\infty} \{h_j^-(p) + h_j^+(p)\} && \text{when } p \in \mathbf{S}^n \setminus \{\sigma\}, \\ &= \infty && \text{when } p = \sigma. \end{aligned}$$

This is a Borel measurable function, because the $h_j^\pm(p)$'s are. The estimate (14.16) follows from (14.22), (14.24), and (14.19). To get (14.17) we observe that

$$(14.26) \quad \sum_{j=1}^{\infty} 2^{j(n-1)} T^{-n+1} \{\mathbf{1}_{B_j^-(z)} + \mathbf{1}_{B_j^+(z)}\} \leq C \left\{ \frac{1}{d(x_0, z)^{n-1}} + \frac{1}{d(x_1, z)^{n-1}} \right\},$$

where $\mathbf{1}_X$ denotes the characteristic function of X , and C depends on k and n . This is not hard to check, using the definition of $B_j^\pm$ and summing a geometric series. Once we have this inequality the estimate (14.17) follows from (14.23) and the reductions made at the beginning of the proof (to the effect that we only need to prove (14.17) with the integral on the right taken over A itself).

This completes the proof of Lemma 14.15.

15. Finding curves inside compact sets with finite length.

The main result in this section provides a criterion for finding the rectifiable curves in our metric space that will be used in the proof of Theorems 12.1 and 12.5.

Proposition 15.1. *Let $(M, d(x, y))$ be a metric space, and let $x, y \in M$ be given, $x \neq y$. Suppose that $\Gamma \subseteq M$ is a compact connected set with $H^1(\Gamma) < \infty$ (see (2.14)) and $x, y \in \Gamma$. Then there is an $L > 0$, $L \leq H^1(\Gamma)$, and an injective 1-Lipschitz mapping $\gamma : [0, L] \rightarrow M$ such that $\gamma(t) \in \Gamma$ for all t , $\gamma(0) = x$, $\gamma(L) = y$, and $H^1(\gamma(E)) = |E|$ for all Lebesgue measurable sets $E \subseteq [0, L]$, where $|E|$ denotes the 1-dimensional Lebesgue measure of E .*

This result is basically well known, but we give the argument for the sake of completeness, and to ensure that the reader has no doubts about its validity in general metric spaces.

For the record, note that H^1 and Lebesgue measure coincide on $\mathbf{R}$. This is easy to check directly, without any covering lemmas. (This could even be the definition of Lebesgue measure on $\mathbf{R}$, depending on the reader's preferences.)

Let M, x, y , and Γ be as in the proposition, and fixed from now on. Let us first establish a simpler result.

Lemma 15.2. *There is a 1-Lipschitz mapping $f : [0, H^1(\Gamma)] \rightarrow \Gamma$ such that $f(0) = x$ and $f(H^1(\Gamma)) = y$.*

In order to obtain this mapping f we shall first build some discrete versions of it and then take a limit using compactness.

Let $\delta > 0$ be small and given. Let $\mathcal{U}$ be a collection of relatively open subsets of Γ such that

$$(15.3) \quad \Gamma = \bigcup_{U \in \mathcal{U}} U, \text{ diam } U < \delta \text{ for all } U \in \mathcal{U}, \text{ and } \sum_{U \in \mathcal{U}} \text{diam } U < H^1(\Gamma) + \delta.$$

The existence of such a family follows from the definition of H^1 in (2.14), except for the requirement that the U 's be relatively open. This is easily accomplished by enlarging slightly the E_j 's given in (2.13), with the cost absorbed in the δ at the end of (15.3). We may also require that $\mathcal{U}$ be finite, since Γ is compact.

Let us call a finite sequence $\{U_i\}_{i=1}^k$ of elements of $\mathcal{U}$ a *chain* if $U_i \cap U_{i+1} \neq \emptyset$ for all i , $1 \leq i < k$. A chain $\{U_i\}_{i=1}^k$ *connects x to y* if $x \in U_1$ and $y \in U_k$. A chain $\{U_i\}_{i=1}^k$ is *reduced* if no element of $\mathcal{U}$ appears in $\{U_i\}$ more than once.

Sublemma 15.4. *There exists a reduced chain $\{U_i\}_{i=1}^k$ of elements of $\mathcal{U}$ which connects x to y .*

Let X denote the set of elements of Γ which can be connected to x by a chain in $\mathcal{U}$. If $U \in \mathcal{U}$ intersects X , then $U \subseteq X$, and similarly if $U \in \mathcal{U}$ intersects $\Gamma \setminus X$, then $U \subseteq \Gamma \setminus X$. It follows that X and $\Gamma \setminus X$ are both relatively open in Γ , and hence $X = \Gamma$, since Γ is connected. This implies that x can be connected to y by a chain. To get a reduced chain one need only remove any unnecessary subchains. This proves Sublemma 15.4.

Let $\{U_i\}_{i=1}^k$ be as in Sublemma 15.4, and let us use it to define an approximately Lipschitz map from $[0, 1] \rightarrow \Gamma$ which almost connects x to y . Choose points z_i , $1 \leq i < k$, such that $z_i \in U_i \cap U_{i+1} \subseteq \Gamma$, and set $z_k = y$. Except for these conditions the z_i 's could be arbitrary. Set $A_j = \sum_{i=1}^j \text{diam } U_i$ and $a_j = A_j A_k^{-1}$, $1 \leq j \leq k$, and set $a_0 = 0$. Thus $A_k \leq H^1(\Gamma) + \delta$, by (15.3), the a_j 's are increasing, and $a_k = 1$. Define $g_\delta : [0, 1] \rightarrow \Gamma$ by

$$(15.5) \quad g_\delta(t) = z_i \text{ when } t \in [a_{i-1}, a_i) \text{ and } 1 < i < k, \quad g_\delta(t) = z_k = y \text{ when } t \in [a_{k-1}, 1].$$

Note that

$$(15.6) \quad g_\delta(1) = y \quad \text{and} \quad d(g_\delta(0), x) = d(z_1, x) \leq \text{diam } U_1 \leq \delta,$$

since $U_1 \in \mathcal{U}$ (so that (15.3) applies).

Sublemma 15.7. *$d(g_\delta(s), g_\delta(t)) \leq (H^1(\Gamma) + \delta)|s - t| + \delta$ when $s, t \in [0, 1]$.*

Let $s, t \in [0, 1]$ be given, and suppose that $s \in [a_{i-1}, a_i)$ and $t \in [a_{j-1}, a_j)$ for some $1 \leq i, j \leq k$. (If i or j equals k , then we should allow the right-hand endpoint $a_k = 1$ in

the corresponding range for s or t .) We have that $g_\delta(s) = g_\delta(t)$ when $i = j$, and so we may assume that $i < j$. Using the definitions and the triangle inequality we get that

$$\begin{aligned}
 (15.8) \quad d(g_\delta(s), g_\delta(t)) &= d(z_i, z_j) \leq \sum_{l=i}^{j-1} d(z_l, z_{l+1}) \leq \sum_{l=i}^{j-1} \text{diam } U_{l+1} \\
 &\leq (A_{j-1} - A_i) + \text{diam } U_j \\
 &\leq (a_{j-1} - a_i)A_k + \delta \leq |s - t|(H^1(\Gamma) + \delta) + \delta.
 \end{aligned}$$

This implies Sublemma 15.7.

The preceding discussion works for all $\delta > 0$, and so we can think of the g_δ 's as a family of functions for which we can choose the parameter δ . The point now is to take a sequence of δ 's tending to 0 such that the corresponding g_δ 's converge uniformly to a function $g : [0, 1] \rightarrow \Gamma$. Such sequences exist, because of the Arzela-Ascoli argument, using also Sublemma 15.7 and the compactness of Γ . (One can find first a sequence of g_δ 's which converges pointwise on a countable dense subset of $[0, 1]$, using the compactness of Γ and a Cantor diagonalization argument. Then one uses the bounds in Sublemma 15.7 to extend the convergence of the g_δ 's to all of $[0, 1]$, and to get uniform convergence.) It is easy to check that g is Lipschitz with norm $\leq H^1(\Gamma)$, and that $g(0) = x$, $g(1) = y$ (by (15.6)). Lemma 15.2 follows by taking $f(t) = g(H^1(\Gamma)^{-1}t)$.

Lemma 15.9. *There is an $L > 0$ and a 1-Lipschitz map $\gamma : [0, L] \rightarrow \Gamma$ such that $\gamma(0) = x$, $\gamma(L) = y$, and L is minimal (no such map exists for a smaller L).*

This lemma is equivalent to the assertion that there exists a Lipschitz map $h : [0, 1] \rightarrow \Gamma$ with $h(0) = x$, $h(1) = y$, and with minimal Lipschitz norm. The existence of such a Lipschitz map with minimal norm is an easy consequence of Lemma 15.2 (which ensures that the set of competitors is nonempty) and the Arzela-Ascoli theorem. This proves Lemma 15.9.

From now on we let γ and L be as in Lemma 15.9. Notice that $L \leq H^1(\Gamma)$, because of Lemma 15.2. The minimality of L implies that γ is one-to-one. (Otherwise γ would have a loop which we could cut out, thereby reducing L .) We want to use the minimality of γ to prove that $H^1(\gamma(E)) = |E|$ for all Lebesgue measurable sets $E \subseteq [0, L]$, as in Proposition 15.1.

Lemma 15.10. $H^1(\gamma([0, L])) = L$.

Of course we have that $H^1(\gamma([0, L])) \leq H^1([0, L]) = L$, since γ is 1-Lipschitz. Suppose that $H^1(\gamma([0, L])) < L$. Then we could apply Lemma 15.2 with Γ replaced with $\gamma([0, L])$ (which is compact and connected and has finite H^1 -measure) to get a 1-Lipschitz map $F : [0, H^1(\gamma([0, L]))] \rightarrow \gamma([0, L])$ which takes the values x, y at the endpoints. This would violate the minimality of L , since $H^1(\gamma([0, L])) < L$ and $\gamma([0, L]) \subseteq \Gamma$. This proves Lemma 15.10.

Lemma 15.11. $H^1(\gamma(E)) = |E|$ for all Lebesgue measurable sets $E \subseteq [0, L]$.

Again $H^1(\gamma(E)) \leq |E|$ is automatic since γ is 1-Lipschitz, and so we need only show that $H^1(\gamma(E)) \geq |E|$. Set $E' = [0, L] \setminus E$, so that $H^1(\gamma(E')) \leq |E'|$ also. Thus

$$(15.12) \quad H^1(\gamma([0, L])) \leq H^1(\gamma(E)) + H^1(\gamma(E')) \leq |E| + |E'| = L.$$

All of these inequalities must be equalities because of Lemma 15.10, and we conclude that $H^1(\gamma(E)) = |E|$. This proves Lemma 15.11.

Thus γ and L have all the required properties. The proof of Proposition 15.1 is now complete.

16. The proofs of Theorems 12.1 and 12.5.

Theorems 12.1 and 12.5 follow easily from the results of the previous three sections, but let us check this carefully. We begin with Theorem 12.1.

Let $M, V, T = \text{diam } V, F, \sigma, B_0, B_1$, and k be as in Theorem 12.1. Given $p \in \mathbf{S}^n$ set $\Gamma(p) = \{z \in M : F(z, t) = p \text{ for some } t \in [0, T]\}$. We shall restrict ourselves to $p \neq \sigma$, so that $\Gamma(p)$ is closed and contained in V , as in Lemma 13.2. This implies that $\Gamma(p)$ is compact, since we are assuming in Theorem 12.1 that V has compact closure. Lemmas 13.2 and 13.8 imply that there is a compact connected set $\Gamma'(p) \subseteq \Gamma(p)$ which intersects $\overline{B}_0$ and $\overline{B}_1$.

Lemma 14.11 applies in this situation. Given any subset A of M we can find a Borel measurable function $h : \mathbf{S}^n \rightarrow [0, \infty]$ such that

$$(16.1) \quad H^1(A \cap \Gamma(p)) \leq h(p) \quad \text{for all } p \in \mathbf{S}^n, \text{ and}$$

$$(16.2) \quad \int_{\mathbf{S}^n} h(p) dp \leq C(n) k^n T^{-n+1} H^n(A)$$

where $C(n)$ depends only on n .

Lemma 16.3. $H^1(\Gamma(p)) < \infty$ for almost all $p \in \mathbf{S}^n$.

Indeed, take $A = V$ above. We are assuming that $H^n(V) < \infty$, and so Lemma 16.3 follows from (16.2) and (16.1) with $A = V$ (and the fact that $\Gamma(p) \subseteq V$ for all $p \neq \sigma$).

Lemma 16.4. *For almost all $p \in \mathbf{S}^n$ there is an $L(p) > 0$, $L(p) \leq H^1(\Gamma(p))$, and an injective 1-Lipschitz mapping $\gamma_p : [0, L(p)] \rightarrow M$ such that $\gamma_p(t) \in \Gamma(p)$ for all t , $\gamma_p(0) \in \overline{B}_0$, $\gamma_p(L(p)) \in \overline{B}_1$, and $H^1(\gamma_p(E)) = |E|$ for all Lebesgue measurable sets $E \subseteq [0, L(p)]$, where $|E|$ denotes the usual Lebesgue measure on $\mathbf{R}$.*

This follows from Proposition 15.1. That is, for each $p \neq \sigma$ we know already that there is a compact connected set $\Gamma'(p) \subseteq \Gamma(p)$ which intersects $\overline{B}_0$ and $\overline{B}_1$. Lemma 16.3 implies that $H^1(\Gamma'(p)) < \infty$ for almost all p , and for these p 's we can apply Proposition 15.1 with $\Gamma = \Gamma'(p)$ to get the curve γ_p . This proves Lemma 16.4.

The conclusion of Theorem 12.1 now follows by chasing definitions. Lemma 16.4 provides us with a family of curves $\gamma_p : [0, L(p)] \rightarrow M$ in M with parameter space $\mathbf{S}^n$, as in Definition 1.2, except that this family is defined only for almost all $p \in \mathbf{S}^n$, and not all p . This is of course not a real problem, for the bad p 's we just take $\gamma_p : [0, L(p)] \rightarrow M$ to be the same as for one of the good p 's. (One should include σ among the bad p 's.) These curves all lie in V because the $\Gamma(p)$'s are contained in V . The curves all start in $\overline{B}_0$ and end in $\overline{B}_1$ because of Lemma 16.4. This family of curves is controlled in measure by $C(n) k^n (\text{diam } V)^{-n+1} H^n(A)$ because of (16.1) and (16.2). Actually, (16.1) is not literally the same as (1.3) (with $a(p) = h(p)$), but Lemma 16.4 implies that (16.1) implies

(1.3). (Note that Definition 1.2 deals only with Borel sets A , which permits us to avoid measurability problems.)

This completes the proof of Theorem 12.1.

The proof of Theorem 12.5 is similar, but there are some minor changes required. Let $M, V, T = \text{diam } V, F, \sigma, x_0, x_1$, and k be as in Theorem 12.1. Given $p \in \mathbf{S}^n \setminus \{\sigma\}$ let $\Gamma(p)$ be as in Lemma 13.7. Thus $\Gamma(p)$ is a closed subset of M which connects x_0 and x_1 and which is contained in V . It is also compact, since we are assuming that V has compact closure in M . Lemma 13.8 implies that there is a compact connected subset $\Gamma'(p)$ of $\Gamma(p)$ which contains both x_0 and x_1 .

Let us apply Lemma 14.15. Given an H^n -measurable set $A \subseteq M$ we get a Borel measurable function $a(p) : \mathbf{S}^n \rightarrow [0, \infty]$ such that

$$(16.5) \quad H^1(A \cap \Gamma(p)) \leq a(p) \quad \text{for all } p \in \mathbf{S}^n, \text{ and}$$

$$(16.6) \quad \int_{\mathbf{S}^n} a(p) dp \leq C \int_{A \cap V \cap \overline{B}_{M(x_0, kT)}} \left\{ \frac{1}{d(x_0, z)^{n-1}} + \frac{1}{d(x_1, z)^{n-1}} \right\} dH^n(z),$$

where C depends only on n and k . Note that $\Phi(p)$ in Lemma 14.15 differs from $\Gamma(p)$ by $\{x_0, x_1\}$, but that this does not affect (16.5).

Lemma 16.7. $H^1(\Gamma(p)) < \infty$ for almost all $p \in \mathbf{S}^n$.

This follows from (16.5) and (16.6), with $A = V$. Remember that we are assuming in Theorem 12.5 that the right side of (16.6) is finite when $A = V$. Remember also that $\Gamma(p) \subseteq V$ when $p \neq \sigma$.

Lemma 16.8. *For almost all $p \in \mathbf{S}^n$ there is an $L(p) > 0$, $L(p) \leq H^1(\Gamma(p))$, and an injective 1-Lipschitz mapping $\gamma_p : [0, L(p)] \rightarrow M$ such that $\gamma_p(t) \in \Gamma(p)$ for all t , $\gamma_p(0) = x_0$, $\gamma_p(L(p)) = x_1$, and $H^1(\gamma_p(E)) = |E|$ for all Lebesgue measurable sets $E \subseteq [0, L(p)]$, where $|E|$ denotes the usual Lebesgue measure on $\mathbf{R}$.*

This follows from Proposition 15.1 and Lemma 16.7 in the same way as for Lemma 16.4.

Theorem 12.5 follows from Lemma 16.8, (16.5), and (16.6) in practically the same manner as before, with only cosmetic changes. One should also notice that (12.10) follows automatically from the first part of the conclusions of Theorem 12.5.

This completes the proof of Theorem 12.5.

17. Some observations about differential forms.

Theorems 1.9 and 1.11 assert the existence of some nice families of paths on a metric space under certain circumstances. We obtained these curves from special Lipschitz mappings, as in Theorems 11.1, 11.5, 12.1, 12.5. We also saw in Theorems 1.22 and 1.27 how these curves could be used to get estimates for a function in terms of its gradient. In this section we present another variation on these themes. We shall show how Lipschitz mappings as in Theorems 11.1 and 11.5 give rise to differential forms with nice properties.

In order to avoid technical issues we shall assume a priori that our metric space is actually a smooth manifold, with a Riemannian metric that is “locally” compatible with

the original distance function. We could do with much less, but the main points will be clear and simpler in this case.

Definition 17.1. Let $(M, d(x, y))$ be a metric space. We say that M is *almost Riemannian* if M is a smooth manifold and if there is a smooth Riemannian metric η on M and a constant $C_0 > 0$ such that each $x \in M$ has a neighborhood $U(x)$ with the property that

$$(17.2) \quad C_0^{-1} d(y, z) \leq D_\eta(y, z) \leq C_0 d(y, z) \quad \text{for all } y, z \in U(x).$$

Here $D_\eta(y, z)$ denotes the Riemannian geodesic distance on M determined by the Riemannian metric η .

Of course this condition holds when $d(x, y)$ is itself the Riemannian distance $D_\eta(x, y)$, but it is better to allow this extra generality. For instance, if M is a smooth submanifold of some $\mathbf{R}^N$, and if we take $d(x, y)$ to be the Euclidean metric $|x - y|$, then $(M, d(x, y))$ is automatically almost Riemannian with respect to the Riemannian metric inherited from $\mathbf{R}^N$, and with C_0 taken to be any constant larger than 1. This assertion can be verified directly from the definitions. The intrinsic Riemannian distance on such a submanifold could still be very different from the Euclidean distance $|x - y|$ at large scales.

Note that n -dimensional Hausdorff measure on M (with respect to $d(x, y)$) and the Riemannian volume on M are each bounded by a constant multiple of the other when M is almost Riemannian, where the constant multiples can be chosen to depend only on n and the constant C_0 in (17.2).

Theorem 17.3. *Let $(M, d(x, y))$ be an oriented almost Riemannian metric space, and suppose that $(M, d(x, y))$ satisfies the same assumptions as in Theorem 12.1 (with $V, T, \sigma, B_0, B_1, k$, and F as before). Assume also that the f_t 's in (12.4) have nonzero degree D . Then there exist bounded measurable differential forms θ_0, θ_T of degree n and Λ of degree $n - 1$ on M with the following properties:*

$$(17.4) \quad d\Lambda = \theta_T - \theta_0;$$

$$(17.5) \quad \theta_{iT} = 0 \text{ a.e. on } M \setminus B_i, \quad i = 0, 1;$$

$$(17.6) \quad \|\theta_{iT}\|_\infty \leq C k^n T^{-n} \quad i = 0, 1;$$

$$(17.7) \quad \Lambda = 0 \text{ a.e. on } M \setminus V;$$

$$(17.8) \quad \|\Lambda\|_\infty \leq C k^n T^{-n+1};$$

$$(17.9) \quad \int_M \theta_i = D, \quad i = 0, 1.$$

This constant C depends on n and the constant C_0 from Definition 17.1 but not on anything else.

Remember from Theorem 11.1 that our criterion for the hypotheses of Theorem 12.1 to hold also implies that the mappings f_t in (12.4) have nonzero degree.

Note that the powers of T on the right sides of (17.6) and (17.8) scale correctly.

Our differential forms θ_{iT}, Λ are only bounded and measurable because we are only requiring F to be Lipschitz. If we smoothed up F then they would be smooth. If M were only a Lipschitz manifold then we would have to settle for bounded measurable forms.

It makes sense to talk about differential forms on M which are bounded and measurable, or even just locally integrable. It also makes sense to take their exterior derivative in the weak sense of currents (by integrating by parts, as in distribution theory), and (17.4) should be interpreted in that way. Note that we are implicitly using the Riemannian metric from Definition 17.1 when we take the L^∞ norms of θ_{iT} or Λ .

Theorem 17.3 can be used to give another approach to the estimate (1.28) (by integrating by parts).

Before we prove Theorem 17.3 let us state its counterpart for Theorem 12.5.

Theorem 17.10. *Let $(M, d(x, y))$ be an oriented almost Riemannian metric space, and suppose that $(M, d(x, y))$ satisfies the same assumptions as in Theorem 12.5 (with $V, T, \sigma, x_0, x_1, k$, and F as before). Assume also that the f_i 's in (12.8) have nonzero degree D . Then there is a locally integrable differential form Λ of degree $n - 1$ on M with the following properties:*

$$(17.11) \quad d\Lambda = (\delta_{x_1} - \delta_{x_0}) D;$$

$$(17.12) \quad \Lambda = 0 \text{ a.e. on } M \setminus V;$$

$$(17.13) \quad |\Lambda(y)| \leq C_1 (d(y, x_0)^{-n+1} + d(y, x_1)^{-n+1}) \quad \text{for almost all } y \in M.$$

This constant C_1 depends on n, k , and the constant C_0 from Definition 17.1 but not on anything else.

Again, the nonzero degree assumption is reasonable, in view of Theorem 11.5.

In (17.11) we are using δ_x to denote the Dirac mass at $x \in M$. More precisely this is the Dirac current, and it should be viewed as a differential form of degree n . In other words, (17.11) really means that

$$(17.14) \quad \int_M du \wedge \Lambda = -(u(x_1) - u(x_0)) D$$

when u is a C^1 function on M (for instance).

Notice that the estimate (17.13) implies that Λ is locally integrable on M , because of our a priori assumption that M be almost Riemannian.

Theorem 17.10 provides another approach to the estimate (1.23).

Before we begin the proof of these theorems in earnest we should deal with a technical point about regularization of Lipschitz mappings.

Lemma 17.15. *Let N be a smooth Riemannian manifold, and let $g : N \rightarrow \mathbf{R}$ be a Lipschitz function which vanishes off a compact subset K of N . Then there is a sequence of smooth functions $g_j : N \rightarrow \mathbf{R}$ which vanish off of any prescribed neighborhood of K in N such that the g_j 's converge uniformly to g , they have uniformly bounded Lipschitz norm, and such that $dg_j \rightarrow dg$ pointwise almost everywhere on N .*

"Lipschitz" here can be defined in terms of the Riemannian metric, but it does not really matter, since we are working with a compactly supported function on a smooth manifold. We do not care here about constants or norms, just the quality of the approximation.

The “ dg ” here denotes the differential of g . Recall that real-valued Lipschitz functions on $\mathbf{R}^n$ are differentiable almost everywhere. The same is true for Lipschitz functions on smooth manifolds, one simply has to interpret the differential differently.

To prove the lemma we may reduce to the case where g is supported in a coordinate chart, by using a partition of unity to break up g into localized pieces. This permits us to reduce to the case of Lipschitz mappings on Euclidean spaces. On Euclidean spaces one can simply regularize by convolving with an approximation to the identity. This kind of regularization procedure is used in the proof of Proposition 1 on p.122 of [St1].

The details of this argument are standard, and we omit them. This proves Lemma 17.15.

Lemma 17.16. *Let N be a smooth Riemannian manifold, and let $G : N \rightarrow \mathbf{S}^n$ be a Lipschitz function which takes only the value σ on the complement of a compact subset K of N . Then there is a sequence of smooth mappings $G_j : N \rightarrow \mathbf{S}^n$ such that each G_j takes only the value σ on the complement of any prescribed neighborhood of K in N , the G_j ’s converge uniformly to G , they have uniformly bounded Lipschitz norm, and $dG_j \rightarrow dG$ pointwise almost everywhere on N .*

This statement about $dG_j \rightarrow dG$ should be treated with a little care. The differential of a map from N to $\mathbf{S}^n$ is a linear map between the tangent spaces, and so it actually depends also on the value of the mapping at the given point. This was not an issue for real-valued maps, and it is really only a minor issue here. The simplest way to deal with it in the present circumstances is to think of G and the G_j ’s as mappings into $\mathbf{R}^{n+1}$, where it is again not an issue. One could make other reasonable formulations but they all amount to the same thing.

To prove Lemma 17.16 we use Lemma 17.15. The latter implies the existence of a sequence of mappings $H_j : N \rightarrow \mathbf{R}^{n+1}$ such that each H_j takes only the value σ on the complement of any prescribed neighborhood of K in N , the H_j ’s converge uniformly to G , they have uniformly bounded Lipschitz norm, and $dH_j \rightarrow dG$ pointwise almost everywhere on N , as maps from N into $\mathbf{R}^{n+1}$. We may as well assume that $\|H_j - G\|_\infty < 1/2$ for all j , because of uniform convergence. Set $G_j = H_j |H_j|^{-1}$. These functions take only the value σ off the same prescribed but arbitrary neighborhood of K , they converge uniformly to G , and they have uniformly bounded Lipschitz norm. It is not hard to check that $dG_j \rightarrow dG$ pointwise almost everywhere on N , using the corresponding fact for the H_j ’s and the chain rule. This proves Lemma 17.16.

Lemma 17.16 is useful because it permits us to reduce certain computations for Lipschitz mappings to the case of smooth mappings. The next lemma provides an example of this.

Let Ω denote the standard volume form on $\mathbf{S}^n$.

Lemma 17.17. *Let N be an oriented smooth Riemannian manifold of dimension n , and let $G : N \rightarrow \mathbf{S}^n$ be a Lipschitz function which takes only the value σ on the complement of a compact subset K of N . Then the pull-back $G^*(\Omega)$ of Ω to N via G is a bounded measurable n -form on N , and*

$$(17.18) \quad \int_N G^*(\Omega) = \text{the degree of } G.$$

Note that $G^*(\Omega)$ makes sense as a bounded measurable n -form as soon as $G : N \rightarrow \mathbf{S}^n$ is Lipschitz, just by using the existence of dG pointwise a.e.

Suppose first that G is smooth. Then (17.18) is practically the definition of the degree, and it is sometimes taken to be. (See [N].) In our case we are using the homological definition of the degree, but it is a standard fact that they are the same. (It is also not very difficult to prove.)

Thus we take (17.18) as given for smooth mappings. For Lipschitz mappings in general we approximate by smooth mappings G_j as in Lemma 17.16. If j is large enough so that $\|G_j - G\|_\infty < 1/2$, then G_j and G are homotopic through maps which take the value σ whenever G_j and G do. (The homotopy is simply obtained by projecting $(1-t)G_j + tG$ onto $\mathbf{S}^n$ inside $\mathbf{R}^{n+1}$.) This means that G_j and G have the same degree when j is large enough (Lemma A.21). Let us check that

$$(17.19) \quad \lim_{j \rightarrow \infty} \int_N G_j^*(\Omega) = \int_N G^*(\Omega).$$

We want to use the Dominated Convergence Theorem. Notice first that $G^*(\Omega)$ and the $G_j^*(\Omega)$'s all vanish off a compact subset of N , because G and the G_j 's take only the value σ on the complement of a compact subset of N . (For this we should be careful to choose the neighborhood of K in Lemma 17.16 to have compact closure.) We also know that the $G_j^*(\Omega)$'s are all uniformly bounded, because of the uniform bound on the Lipschitz norm of the G_j 's, and the definition of the pull-backs. The remaining point is that $G_j^*(\Omega) \rightarrow G^*(\Omega)$ pointwise a.e. on N . This is not hard to verify, using both the uniform convergence of the G_j 's to G and the pointwise a.e. convergence of the differentials. We omit the details. We conclude from dominated convergence that (17.19) holds, and (17.18) follows.

This completes the proof of Lemma 17.17.

Let us now prove Theorem 17.3. Let M, V, F , etc. be as in the theorem. Let f_t be as in (12.4), and set

$$(17.20) \quad \theta_t = f_t^*(\Omega), \quad 0 \leq t \leq T.$$

As in Lemma 17.17, these make sense as bounded measurable n -forms on M . We also get that

$$(17.21) \quad \int_M \theta_t = D$$

for all $t \in [0, T]$, where D is as in Theorem 17.3, because of Theorem 17.17. This gives (17.9).

Let us check (17.5) and (17.6). For (17.5) we observe that $f_{iT} \equiv \sigma$ on $M \setminus B_i$, $i = 0, 1$, because of (12.3). This implies that $df_{iT} = 0$ a.e. on $M \setminus B_i$, $i = 0, 1$. (This assertion comes down to the fact that if $g : \mathbf{R}^n \rightarrow \mathbf{R}$ is Lipschitz and is constant on some measurable set E , then the gradient of g vanishes almost everywhere on E . If one really wants to check this carefully, it is helpful to use the fact that almost every point of E is a point of density

of E , and the Lipschitz condition.) Thus we get (17.5). As for the bound (17.6) one uses the definition of the pull-back and the Lipschitz condition on F and hence f_t . More precisely, our Lipschitz condition is initially given in terms of the metric $d(x, y)$, but then our almost Riemannian assumption implies that the differential of each f_t is bounded by $C k T^{-1}$, where the norm of the differential is computed in terms of the Riemannian metric provided by Definition 17.1. It is easy to derive (17.6) from this bound.

To define the $(n-1)$ -form Λ we first define some auxiliary $(n-1)$ -forms λ_t . In the end we shall take

$$(17.22) \quad \Lambda = \int_0^T \lambda_t dt.$$

Thus we would like to choose λ_t so that $d\lambda_t$ is the t -derivative of θ_t . There is a well-known formula for doing this, but we shall simply write everything down.

The forms λ_t will only be bounded and measurable, and we only need to worry about defining them almost everywhere. Fix $z \in M$ and $t \in (0, T)$, and let us try to define λ_t at z . We need to assume that the differential of f_t exists at z , and that the derivative of $f_t(z)$ in t exists. These derivatives will exist for almost all (z, t) , because F is Lipschitz. Set $p = f_t(z)$, and let us write Ω_p for the volume form on the tangent space $T_p \mathbf{S}^n$ of $\mathbf{S}^n$ at p . (This is slightly bad notation, because the subscript t in λ_t plays a completely different role from the subscript p in Ω_p , and so the reader should be vigilant to avoid confusion.) Let $v \in T_p \mathbf{S}^n$ denote the tangent vector at p which represents the derivative of $f_t(z)$ in t at our fixed choice of t . Let $\omega_{z,t}$ denote the alternating multilinear $(n-1)$ -form on $T_p \mathbf{S}^n$ obtained by taking the interior product of Ω_p with v . (See p.61 of [Wa] for a discussion of interior multiplication.) We take λ_t at z to be the alternating $(n-1)$ -form on $T_z M$ obtained by pulling $\omega_{z,t}$ back using (the differential of) f_t . This is the form that we want. Note that λ_t is bounded and measurable, since F is Lipschitz.

Lemma 17.23. $d\{\int_0^T \lambda_t dt\} = \theta_T - \theta_0$, in the sense of currents.

(Keep in mind that the integral on the left defines a bounded measurable $(n-1)$ -form on M .)

To prove this we may as well assume that F is smooth. Indeed, if F is not smooth, but merely Lipschitz, then we can approximate it by smooth mappings F_j , as in Lemma 17.16. The main point is that the analogues of λ_t and θ_t for F_j converge to λ_t and θ_t in the L^1 topology. This can be proved using the Dominated Convergence Theorem. Specifically, all of these forms are uniformly bounded and vanish off a compact set, and one can use the pointwise convergence of the differentials provided by Lemma 17.16 to get pointwise convergence of these forms here. We encountered a similar point in the proof of Lemma 17.17. The details are straightforward but slightly messy, and we omit them. The equation in Lemma 17.23 is stable under L^1 convergence, because we are interpreting it in the sense of currents. Thus it suffices to prove Lemma 17.23 under the assumption that F is smooth.

To prove Lemma 17.23 in the smooth case we need a more convenient formula for λ_t . Let us start with the temporary lie that f_t takes values in $\mathbf{R}^n$, and not $\mathbf{S}^n$, and that Ω is the standard volume form on $\mathbf{R}^n$. That is, $\Omega = dw_1 \wedge dw_2 \wedge \cdots \wedge dw_n$, where $\{w_i\}$ are the standard coordinates for $\mathbf{R}^n$. Let us write f_t^i for the i^{th} coordinate of f_t , and let $\dot{f}_t$ and $\dot{f}_t^i$

denote the t derivative of f_t and f'_t . Then df_t^i is a 1-form on M for each i , and θ_t becomes $df_t^1 \wedge df_t^2 \wedge \cdots \wedge df_t^n$, since θ_t is the pull-back of Ω by f_t . In this situation we have that

$$(17.24) \quad \lambda_t = \sum_{j=1}^n (-1)^{j+1} \dot{f}_t^j df_t^1 \wedge \cdots \wedge \widehat{df_t^j} \wedge \cdots \wedge df_t^n.$$

where we follow the custom of putting a hat over the factor that should be omitted from the wedge product. This is simply an explicit realization of the interior product mentioned above.

This temporary lie is not really a lie, it is the truth in a different coordinate system. That is, given any point $q \in \mathcal{S}^n$, we can find a neighborhood U of q in $\mathbf{S}^n$ and a diffeomorphism ϕ from U onto an open set in $\mathbf{R}^n$ such that Ω corresponds to the standard volume form on $\mathbf{R}^n$ under ϕ . To do this we start with any diffeomorphism ψ from a neighborhood V of q in $\mathbf{S}^n$ onto an open subset of $\mathbf{R}^n$. This permits us to reduce to the problem of making a local change of variables on $\mathbf{R}^n$ which transforms a given nonvanishing volume form in the neighborhood of $\psi(q)$ to the standard volume form. This is the same as saying that if we have a smooth nonvanishing function J in a neighborhood of $\psi(q)$ in $\mathbf{R}^n$, then we can find a (diffeomorphic) change of variables on a neighborhood of $\psi(q)$ whose jacobian is J . This can be accomplished with a mapping $\Psi(w)$ which changes only the w_1 coordinate of w , and one can easily write down a formula for it (by integrating J in the first variable).

Thus there are local coordinates on $\mathbf{S}^n$ which permit us to reduce (locally) to the situation of the paragraph of (17.24). If (U, ϕ) is such a good coordinate system for an open subset U of $\mathbf{S}^n$ (so that Ω corresponds to the standard volume form on $\mathbf{R}^n$ under ϕ), then (17.24) is a correct formula for λ_t on $f_t^{-1}(U)$ if we take f_t^i to be the i^{th} coordinate of $\phi \circ f_t$. This assertion can be derived easily from the definitions of interior multiplication and the pull-back of a form under a differentiable mapping, using also the chain rule. The advantage of our original definition of λ_t is that it makes sense directly, without any auxiliary choices, while this description using good local coordinates on $\mathbf{S}^n$ makes it easier for us to compute.

If we use good coordinates as just described then we get that

$$(17.25) \quad d\lambda_t = \sum_{j=1}^n df_t^1 \wedge \cdots \wedge d\dot{f}_t^j \wedge \cdots \wedge df_t^n,$$

by definition of the exterior derivative. In this formula only the j^{th} factor is being differentiated in t (i.e., has a dot). The right hand side is plainly the derivative in t of $df_t^1 \wedge \cdots \wedge df_t^n$. Thus we get that

$$(17.26) \quad d\lambda_t = \frac{d}{dt} \theta_t$$

in these local coordinates. This formula holds everywhere on M , since it does not matter where on $\mathbf{S}^n$ we are. (We have good local coordinates everywhere on $\mathbf{S}^n$.)

Of course (17.26) implies Lemma 17.23, just by integrating. This proves Lemma 17.23 in the case where F is smooth, and therefore in general, as we saw before.

Define Λ as in (17.22), so that (17.4) follows from Lemma 17.23. It remains to verify (17.7) and (17.8). Let us check that

$$(17.27) \quad \lambda_t = 0 \text{ a.e. on } M \setminus V$$

$$(17.28) \quad \|\lambda_t\|_\infty \leq C k^n T^{-n}$$

for $0 \leq t \leq T$. For (17.27) we use the fact that $F(z, t) = \sigma$ for all t when $z \in M \setminus V$, which implies that the first derivatives of F vanish for almost such (z, t) . This implies (17.27) by definition of λ_t . It is also easy to derive (17.28) from the definition of λ_t and our assumption that F is $k T^{-1}$ -Lipschitz. Keep in mind that the norm in (17.28) uses the Riemannian metric in Definition 17.1, and that our almost-Riemannian assumption implies that the Riemannian norm of the differential of F is bounded by $C k T^{-1}$. (Note that the derivatives of F in both the z and t variables arise in the definition of λ_t .) This proves (17.27) and (17.28). (Notice that they are both compatible with (17.24) as well.) Once we have (17.27) and (17.28) we can derive (17.7) and (17.8) directly from (17.22), just by integrating.

This completes the proof of Theorem 17.3.

Now let us prove Theorem 17.10. Let M, V, F, x_0, x_1 , etc. be as given in the statement of Theorem 17.10. Define θ_t as in (17.20), but only for $0 < t < T$ now. We still get (17.21) for this range of t 's, for the same reason as before.

Lemma 17.29.

$$\lim_{t \rightarrow iT} \theta_t = \delta_x, D, \quad i = 0, 1,$$

in the sense of currents.

For simplicity of notation let us prove this only when $i = 0$. We need three facts. The first is that

$$(17.30) \quad \theta_t = 0 \text{ a.e. on } M \setminus \overline{B}(x_0, kt).$$

This follows from (12.7), which implies that $f_t \equiv \sigma$ on $M \setminus \overline{B}(x_0, kt)$, so that its differential vanishes at almost all points in this set. The second fact is that

$$(17.31) \quad \|\theta_t\|_\infty \leq C k^n t^{-n}.$$

This follows from (12.6), which implies that f_t is Lipschitz with norm $\leq k t^{-1}$, and the definition of the pull-back. We are also using the assumption that M is almost Riemannian here, to pass from the Lipschitz condition with respect to $d(x, y)$ to bounds on the Riemannian norm of the differential. The third fact is that (17.21) holds. Once we have these three facts it is not hard to prove the lemma for $i = 0$. This comes down to the statement that

$$(17.32) \quad \lim_{t \rightarrow iT} \int_M \phi \theta_t = \phi(x_0) D$$

for all continuous functions ϕ on M , and the proof of this is a standard exercise. The proof of the $i = 1$ case is the same, just with more complicated notation, and the lemma follows.

We can define λ_t for $0 < t < T$ exactly as before (just after (17.22)). This gives a bounded measurable $(n-1)$ -form for each such t , but the L^∞ norm of λ_t remains bounded only when we restrict ourselves to a compact subinterval of $(0, T)$.

Lemma 17.33. *If $0 < s < t < T$, then*

$$(17.34) \quad d\left\{\int_s^t \lambda_t dt\right\} = \theta_t - \theta_s$$

in the sense of currents.

This is the same as Lemma 17.23, with only trivial changes.

We want to define Λ as in (17.22), but we have to be careful to make certain that this makes sense (that the integral converges). For this we need some simple estimates.

Lemma 17.35. *We have that*

$$(17.36) \quad \lambda_t = 0 \text{ a.e. on } M \setminus (V \cap \overline{B}(x_0, kt)) \quad \text{and} \quad \|\lambda_t\|_\infty \leq C k^n t^{-n}$$

when $0 < t \leq T/2$, and

$$(17.37) \quad \lambda_t = 0 \text{ a.e. on } M \setminus (V \cap \overline{B}(x_1, k(T-t))) \quad \text{and} \quad \|\lambda_t\|_\infty \leq C k^n (T-t)^{-n}$$

when $T/2 \leq t < T$.

This is proved in essentially the same manner as (17.27) and (17.28). That is, we use (12.7) to conclude that F is constant on certain sets, and therefore that its first derivatives vanish almost everywhere on these sets, and we use the Lipschitz bound (12.6) and the almost-Riemannian assumption to get the bounds on $\|\lambda_t\|_\infty$. This proves Lemma 17.35.

One consequence of Lemma 17.35 is that

$$(17.38) \quad \|\lambda_t\|_1 = O(1) \quad \text{as } t \rightarrow 0 \text{ or } t \rightarrow T.$$

We don't have a uniform bound on $\|\lambda_t\|_1$ in terms of constants that we control, but we do get boundedness of these L^1 norms because of our a priori almost-Riemannian assumption (which ensures that M is smooth in particular). This implies that (17.22) does define a measurable $(n-1)$ -form on M with finite L^1 norm. This form Λ satisfies (17.11) because of (17.34), Lemma 17.29, and a limiting argument. We also get (17.12) from Lemma 17.35. The derivation of (17.13) from (17.36) and (17.37) is a straightforward exercise in calculus.

This completes the proof of Theorem 17.10.

18. Nontrivial Lipschitz mappings on “codimension-1” sets in Euclidean spaces.

In Theorems 11.1 and 11.5 we saw how one can build interesting Lipschitz mappings on metric spaces under certain conditions on the spaces. In this section we shall show how similar mappings can be constructed on a subset of a Euclidean space under geometric conditions on its complement.

Let n be a fixed positive integer. We shall be working with subsets of $\mathbf{R}^{n+1}$, which we always view as being equipped with the Euclidean metric.

Definition 18.1. Let M be a closed subset of $\mathbf{R}^{n+1}$, and let x_0 and x_1 be two elements of M . We say that M *has a nice complement relative to x_0 and x_1* (with constant C) if there are two distinct components Ω_+, Ω_- of $\mathbf{R}^{n+1} \setminus M$ and a pair of C -Lipschitz mappings $\gamma_+ : [0, |x_1 - x_0|] \rightarrow \Omega_+$, $\gamma_- : [0, |x_1 - x_0|] \rightarrow \Omega_-$ such that $\gamma_{\pm}(0) = x_0$, $\gamma_{\pm}(|x_0 - x_1|) = x_1$, $\gamma_+(t) \in \Omega_+$ and $\gamma_-(t) \in \Omega_-$ when $0 < t < |x_0 - x_1|$, and

$$(18.2) \quad \text{dist}(\gamma_{\pm}(t), M) \geq C^{-1} \min\{t, |x_0 - x_1| - t\} \quad \text{when } 0 < t < |x_0 - x_1|.$$

This condition means that M should not look like a cusp near x_0 or x_1 , that its complementary domains should not have unavoidable narrow passages between x_0 and x_1 , etc. This condition holds for all $x_0, x_1 \in M$ when the complementary domains are “NTA (nontangentially accessible) domains” or “uniform domains”.

Theorem 18.3. *Suppose that M is a closed subset of $\mathbf{R}^{n+1}$ with a nice complement relative to x_0 and x_1 , and set $T = |x_1 - x_0|$. Fix a point $\sigma \in \mathbf{S}^n$. Then there is a constant k and a mapping $F : M \times (0, T) \rightarrow \mathbf{S}^n$ with the following properties:*

$$(18.4) \quad F \text{ is Lipschitz with norm } \leq k\epsilon^{-1} \text{ on } M \times (\epsilon, T - \epsilon) \text{ when } 0 < \epsilon < T;$$

$$(18.5) \quad F(z, t) = \sigma \text{ when either } d(z, x_0) > kt \text{ or } d(z, x_1) > k(T - t);$$

$$(18.6) \quad \text{each } f_t = F(\cdot, t) : M \rightarrow \mathbf{S}^n, \ 0 < t < T, \text{ cannot be homotoped to a constant through maps that all equal } \sigma \text{ on the complement of some compact subset of } M.$$

Here k depends only on n and the constant in Definition 18.1.

Under suitable topological conditions on M we can also get the mappings f_t in (18.6) to have nonzero degree. For instance we can do this if M is a smooth oriented n -dimensional embedded submanifold of $\mathbf{R}^{n+1}$. In that case the f_t 's are very easy to understand when t is close to 0 or to T , because they will be concentrated in small balls in which M is very flat.

The conclusion of this theorem is practically the same as the conclusion of Theorem 11.5. The hypotheses are very different. The assumptions of Theorem 18.3 are much simpler than those of Theorem 11.5, and in some ways they are much weaker. However the hypotheses of Theorem 11.5 have the advantage of being intrinsic, requiring no embedding into a Euclidean space nor information about any “complement”.

The main ingredient in the proof of Theorem 18.3 is the following classical result in topology. Given $p \in \mathbf{R}^{n+1}$, define $\pi_p : \mathbf{R}^{n+1} \setminus \{p\} \rightarrow \mathbf{S}^n$ by

$$(18.7) \quad \pi_p(z) = \frac{z - p}{|z - p|}.$$

Theorem 18.8. *Let K be a compact subset of $\mathbf{R}^{n+1}$. If $p, q \in \mathbf{R}^{n+1} \setminus K$ lie in different components of $\mathbf{R}^{n+1} \setminus K$, then the restrictions of π_p and π_q to K are not homotopic to each other.*

This is one-half of Theorem VI.10 on p.97 of [HW]. For our purposes it is not in the most convenient form, and we shall modify it slightly.

Lemma 18.9. *Let K be a compact subset of $\mathbf{R}^{n+1}$. If $p \in \mathbf{R}^{n+1} \setminus K$ lies in a bounded component of the complement of K , then the restriction of π_p to K is not homotopically trivial.*

This is just a special case of Theorem 18.8. The point is simply that the restriction of π_q to K is homotopically trivial when q lies outside of a ball that contains K .

Lemma 18.10. *Set $\epsilon_+ = (1, 0, \dots, 0)$ and $\epsilon_- = (-1, 0, \dots, 0)$, and let $\sigma \in \mathbf{S}^n$ be given. There is a smooth mapping $\Sigma : \mathbf{R}^{n+1} \setminus \{\epsilon_+, \epsilon_-\} \rightarrow \mathbf{S}^n$ with $\Sigma(z) = \sigma$ when $|z| \geq 10$ which satisfies the following property. If E is a closed subset of $\mathbf{R}^{n+1} \setminus \{\epsilon_+, \epsilon_-\}$ which separates ϵ_+ from ϵ_- (i.e., if ϵ_+ and ϵ_- lie in different components of the complement of E), then the restriction of Σ to E cannot be deformed to a constant through mappings from E into $\mathbf{S}^n$ which equal σ off a compact set. (This last part is relevant only when E is unbounded.)*

Let $\widehat{\mathbf{R}}^{n+1} = \mathbf{R}^{n+1} \cup \{\infty\}$ denote the one-point compactification of $\mathbf{R}^{n+1}$, which we can identify with $\mathbf{S}^{n+1}$. Let P denote the $z_1 = 0$ hyperplane in $\mathbf{R}^{n+1}$, and set $\widehat{P} = P \cup \{\infty\}$. Let $h : \widehat{\mathbf{R}}^{n+1} \rightarrow \widehat{\mathbf{R}}^{n+1}$ be a Möbius transformation such that $h(\epsilon_+) = \infty$, $h(\epsilon_-) = 0$, $h(\widehat{P}) = \mathbf{S}^n$, and $h(\infty) = \sigma$. We are going to use h to reduce to the situation of Lemma 18.9, with $p = 0$.

Let $\Pi : \mathbf{R}^{n+1} \setminus \{0\} \rightarrow \mathbf{S}^n$ be a smooth mapping which is homotopic to π_0 (from (18.7)) and which equals σ on $h(\mathbf{R}^{n+1} \setminus B(0, 10))$ (= a reasonably small ball around σ). (One can realize Π as the composition of π_0 with a map from $\mathbf{S}^n$ to itself which is homotopic to the identity.) We take $\Sigma = \Pi \circ h$, so that $\Sigma(z) = \sigma$ when $|z| \geq 10$.

Let E be as in the lemma, and set $\widehat{E} = E \cup \{\infty\}$ when E is unbounded and $\widehat{E} = E$ when E is bounded. Then $h(\widehat{E})$ is a compact subset of $\widehat{\mathbf{R}}^{n+1}$, and 0 lies in a bounded component of $\widehat{\mathbf{R}}^{n+1} \setminus \widehat{E}$. Hence the restriction of π_0 to $h(\widehat{E})$ is not homotopically trivial, by Lemma 18.9. The restriction of $\Pi \circ h$ to $\widehat{E}$ is also not homotopically trivial, since Π is homotopic to π_0 and h is a homeomorphism. Therefore the restriction of Σ to E has the required homotopic nontriviality property. This proves Lemma 18.10.

Next we address a small technical point.

Lemma 18.11. *In Definition 18.1 we may assume without loss of generality that $\gamma_{\pm}$ are smooth on $(0, |x_0 - x_1|)$, at least if we permit ourselves to increase C slightly.*

This follows from a standard approximation argument. That is, we can approximate $\gamma_{\pm}$ in Definition 18.1 by smooth mappings on $(0, |x_0 - x_1|)$ without disturbing the other properties and estimates in Definition 18.1 too much. We omit the details. We could also get precise bounds on the higher derivatives of a modified $\gamma_{\pm}$, but we shall not need that.

Let us now begin the proof of Theorem 18.3. Let $x_0, x_1 \in M$ and $\gamma_{\pm} : [0, T] \rightarrow \mathbf{R}^{n+1}$ be as in Definition 18.1, but with the additional assumption that γ_+ and γ_- are smooth, as in Lemma 18.11. Set $T = |x_0 - x_1|$, and let $\sigma \in \mathbf{S}^n$ be given. We shall change variables in order to renormalize our picture into the form required for Lemma 18.10. We shall then use the mapping Σ from Lemma 18.10 to build the desired mapping F .

Set $c(t) = (\gamma_+(t) + \gamma_-(t))/2$ and $d(t) = |\gamma_+(t) - \gamma_-(t)|/2$ for $t \in [0, T]$, and observe that

$$(18.12) \quad C^{-1} \min\{t, T-t\} \leq d(t) \leq C \min\{t, T-t\}$$

for some C and all t , because of the conditions in Definition 18.1. (That is, we use (18.2) to get the lower bound in (18.12), and we use the Lipschitz conditions on $\gamma_{\pm}$ to get the upper bounds.) We would like to straighten our picture out by mapping $c(t)$ to the origin and $\gamma_{\pm}(t)$ to the x_1 -axis in a nice way for each t . The next lemma will enable us to do that.

Lemma 18.13. *There is a smooth family of rotations $\rho_t : \mathbf{R}^{n+1} \rightarrow \mathbf{R}^{n+1}$, $t \in (0, T)$, such that*

$$(18.14) \quad \rho_t(\gamma_{\pm}(t) - c(t)) = (\pm d(t), 0, \dots, 0) \quad \text{and}$$

$$(18.15) \quad \left| \frac{d}{dt} \rho_t \right| \leq C \min\{t, T-t\}^{-1}$$

for all $t \in (0, T)$. Here C depends only on n and the constant from Definition 18.1, and we are implicitly using the standard metric on the rotation group in (18.15).

It is easy to find rotations which satisfy (18.14), but we have to be a little careful to get (18.15). We shall do this “infinitesimally” first, and then we shall obtain ρ_t by integrating an ordinary differential equation.

Infinitesimal deformations of rotations can be represented in terms of antisymmetric linear transformations, and our first task is to write down the relevant antisymmetric linear transformations. Set $v(t) = (\gamma_+(t) - c(t))/d(t)$ when $0 < t < T$, and note that $|v(t)| = 1$ for all t , because of the definition of $c(t)$ and $d(t)$. The smoothness of $\gamma_{\pm}$ on $(0, T)$ implies that $v(t)$ is also smooth. Let $\dot{v}(t)$ denote the derivative of $v(t)$ in t . Notice that $\langle \dot{v}(t), v(t) \rangle = 0$ for all t , since $|v(t)| \equiv 1$, where $\langle \cdot, \cdot \rangle$ denotes the standard inner product on $\mathbf{R}^{n+1}$. Let A_t denote the linear transformation on $\mathbf{R}^{n+1}$ given by

$$(18.16) \quad A_t(w) = -\dot{v}(t)\langle w, v(t) \rangle + v(t)\langle w, \dot{v}(t) \rangle.$$

Thus $A_t(v(t)) = -\dot{v}(t)$ and A_t is antisymmetric for all t .

Fix a rotation ρ which satisfies (18.14) at $t = T/2$. By well-known results there is a smooth family of rotations ρ_t , $t \in (0, T)$, which satisfies $\dot{\rho}_t = \rho_t \circ A_t$ for all t and $\rho_t = \rho$ when $t = T/2$. These ρ_t 's are rotations for all t because the A_t 's are antisymmetric. (It is a little easier to verify first that the transposes of the ρ_t 's are rotations.) The differential equation defining ρ_t implies that

$$(18.17) \quad \frac{d}{dt} \{ \rho_t(v(t)) \} = \dot{\rho}_t(v(t)) + \rho_t(\dot{v}(t)) = \rho_t(A_t(v(t))) + \rho_t(\dot{v}(t)) = 0,$$

since $A_t(v(t)) = -\dot{v}(t)$. Thus $\rho_t(v(t))$ is constant, and the fact that (18.14) holds when $t = T/2$ implies that $\rho_t(v(t)) = (1, 0, \dots, 0)$ for all t . This implies (18.14) for all t , because of the definitions and a little algebra.

As for (18.15), we have that $|\frac{d}{dt}\rho_t| \leq 2|\dot{v}(t)|$, and it is easy to check that $|\dot{v}(t)| \leq C d(t)^{-1}$ for all t . This last comes down to the fact that $\gamma_{\pm}$ are Lipschitz with bounded norm, by Definition 18.1, and the definition of $v(t)$ and $d(t)$. Combining these inequalities with (18.12) we get (18.15). This proves Lemma 18.13.

We want to use Lemma 18.13 to produce the mapping F required in Theorem 18.3. Let Σ be as in Lemma 18.10, and let ρ_t be as in Lemma 18.13. Define $F : M \times (0, T) \rightarrow \mathbf{S}^n$ by

$$(18.18) \quad F(z, t) = \Sigma\left(\frac{\rho_t(z - c(t))}{d(t)}\right).$$

In order to verify that this has the desired properties we first establish some preliminary information. Observe that

$$(18.19) \quad \left|\frac{\rho_t(z - c(t))}{d(t)} - c_{\pm}\right| = \frac{|z - \gamma_{\pm}(t)|}{d(t)}.$$

This uses (18.14) and the fact that the ρ_t 's are rotations. Next, there is a constant C_0 such that

$$(18.20) \quad \left|\frac{\rho_t(z - c(t))}{d(t)} - c_{\pm}\right| \geq C_0^{-1}$$

when $z \in M$ and $t \in (0, T)$. This follows from (18.19), using (18.2) and (18.12).

Let us prove that F satisfies (18.5). The main point is that $\Sigma(\zeta) = \sigma$ when $|\zeta| \geq 10$, as in Lemma 18.10. We also need some small computations with distances. Suppose that $|z - x_0| > 2Kt$, where $K > 1$ is some constant. If K is large enough, then $|z - \gamma_{\pm}(t)| > Kt$, because $|\gamma_{\pm}(t) - x_0| \leq Ct$, by the properties of $\gamma_{\pm}$ in Definition 18.1. This implies that

$$(18.21) \quad \left|\frac{\rho_t(z - c(t))}{d(t)} - c_{\pm}\right| \geq 10$$

if K is large enough, because of (18.19) and (18.12). Thus $|z - x_0| > 2Kt$ implies that $F(z, t) = \sigma$ if K is chosen large enough, since $\Sigma(\zeta) = \sigma$ when $|\zeta| \geq 10$. A similar argument shows that $F(z, t) = \sigma$ when $|z - x_1| > 2K(T - t)$. This proves (18.5).

Now let us check the Lipschitz property for $F(z, t)$ required in (18.4). The first point is that $\Sigma(\zeta)$ is Lipschitz on $\{\zeta \in \mathbf{R}^{n+1} : |\zeta - c_{\pm}| \geq C_0^{-1}\}$, since Σ is smooth away from $c_{\pm}$ and constant off a compact set. These are the only ζ 's which matter in the definition (18.18) of F , because of (18.20). The Lipschitz norm for Σ on this set does not depend on anything important (like x_0 or x_1). The second point is that we don't need to worry about the (z, t) 's such that $|z - x_0| > 10Kt$ or $|z - x_1| > 10K(T - t)$, because of the preceding observation that F takes only the value σ on or near such points. Let us check that the remaining pairs (z, t) satisfy

$$(18.22) \quad |z - c(t)| \leq C d(t)$$

for some constant $C = C(K)$. Indeed, $|c(t) - x_0| \leq C t$ and $|c(t) - x_1| \leq C(T - t)$, because of the definition of $c(t)$ and the properties of $\gamma_{\pm}$ in Definition 18.1. These inequalities and our assumptions on z imply that $|z - c(t)| \leq C \min(t, T - t)$. Therefore (18.22) holds, because of (18.12).

The conclusion of these two observations is that to prove the Lipschitz property for $F(z, t)$ required in (18.4) it is enough to prove the same kind of Lipschitz estimate for $\rho_t(z - c(t))/d(t)$, and only for pairs (z, t) which satisfy a bound like (18.22). This Lipschitz estimate for $\rho_t(z - c(t))/d(t)$ can be derived from various estimates established already, namely (18.15), the fact that $c(t)$ and $d(t)$ are Lipschitz with bounded norm (because $\gamma_{\pm}$ are Lipschitz with bounded norm), the lower bound for $d(t)$ in (18.12), (18.22), and the fact that the ρ_t 's are all rotations.

It remains to prove the homotopic nontriviality property in (18.6). For this we employ the corresponding assertion in Lemma 18.10. Let E_t denote the subset of $\mathbf{R}^{n+1}$ which is the image of M under the mapping $z \rightarrow \rho_t(z - c(t))/d(t)$. This is a closed subset of $\mathbf{R}^{n+1}$, and E_t separates c_+ from c_- for each t because $\gamma_+(t)$ and $\gamma_-(t)$ lie in different components of $\mathbf{R}^{n+1} \setminus M$ for all $t \in (0, T)$, as in Definition 18.1. Thus the restriction of Σ to E_t (for any fixed t) cannot be deformed to a constant through continuous maps from E_t into $\mathbf{S}^n$ which equal σ off a compact set, because of Lemma 18.10. This implies the homotopic nontriviality property in (18.6), by the definition (18.18) of F .

This completes the proof of Theorem 18.3.

Remark 18.23. In Theorem 18.3 we gave another criterion for the conclusions of Theorem 11.5 to hold, and for most of the hypotheses of Theorem 12.5 to hold. If we add suitable assumptions on the n -dimensional Hausdorff measure of M , then we can combine Theorems 18.3 and 12.5 to get the existence of a nice family of curves which connect x_0 to x_1 in M . We can then derive the analogue of Theorem 1.22 in this case. If M is smooth oriented n -dimensional embedded submanifold of $\mathbf{R}^{n+1}$, then we can apply Theorem 17.10. (Note that M is automatically almost Riemannian (Definition 17.1) in this case, with constant C_0 as close to 1 as we like. Also, if M is smooth, then the mapping F defined in (18.18) is also smooth, because of the smoothness of Σ and $\gamma_{\pm}$.) If we assume that M has a nice complement with respect to all pairs of points x_0 and x_1 in M , and if we make suitable assumptions on the Hausdorff measure, then we can obtain Sobolev and Poincaré inequalities on M . (See Appendix B, especially Theorem B.16.) Similar results were obtained in [Se1] and Section 6 [DS2] by different methods (although there remains a certain similarity). The approach given here is more direct and provides an explanation for the slightly miraculous arguments in [Se1] and [DS2].

Appendices.

A. Review of degree theory and homology.

Let M be an oriented topological manifold of dimension n and let f be a continuous map from M into the n -sphere $\mathbf{S}^n$. The degree of f , which measures the number of times that the image of f wraps around $\mathbf{S}^n$ (counted with orientations), plays an important role in this paper, and we review some of its main features in this appendix.

In order for the degree of f to be well defined we need to assume some compactness, i.e., that M is compact, or that f is constant off a compact set.

Suppose for the moment that M is compact and that M and f are smooth. Let z be a point in $\mathbf{S}^n$ such that z is not the image of a critical point of f and z has only finitely many preimages in M . Almost every point in $\mathbf{S}^n$ has these properties. To each preimage $x \in M$ of z one can associate a choice of $+1$ and -1 , depending on whether the differential of f at x preserves or reverses orientations. The degree of f for the value z is defined to be the sum of all these ± 1 's. If there are no preimages then one takes the degree to be 0. It is a great fact that this value of the degree does not depend on the choice of z , and that it remains unchanged by homotopies of f .

See [Mi] and [N] for more information about the degree of a mapping in the smooth category. Unfortunately this approach does not work when M is simply a topological manifold. The easiest way to define the degree in the topological category is to use homology, which we review in this appendix.

All homology groups in this appendix will be taken with respect to the integers. We shall follow [Ma]. As in Chapter VII of [Ma], given a topological space X we let $H_n(X)$ denote the n -dimensional homology group of X , and given a topological subspace A of X we denote by $H_n(X, A)$ the relative homology group of the pair (X, A) . Thus $H_n(X)$ is the abelian group of n -cycles in X (singular n -chains which lie in the kernel of the boundary operator) modulo the subgroup of cycles which are boundaries (i.e., which lie in the image of the boundary operator). Roughly speaking, the relative homology of pair (X, A) is defined in the same way as the homology of X except that the parts of the singular chains (formal sums of continuous maps from cubes into X) living in A are thrown to the garbage (when defining cycles and boundaries). If $A = \emptyset$, then $H_n(X, A) = H_n(X)$. The $n = 0$ case behaves a little differently from the others, in such a way that it is often preferable to replace $H_0(X)$ with the "reduced homology". We shall ignore this technical point because we shall not need it, but unfortunately it will make some of the statements that follow more complicated in low dimensions.

Our main concern here will be the n -dimensional homology of spaces of dimension n and the fundamental class. In order to deal with noncompact spaces, and to make certain cutting-and-pasting arguments, it is convenient to use the relative homology groups too.

Examples A.1. (a) $H_n(\mathbf{R}^n) = 0$.

(b) $H_n(\mathbf{S}^n) = \mathbf{Z}$.

(c) If B^n denotes the standard closed unit ball in $\mathbf{R}^n$ (so that $\partial B^n = \mathbf{S}^{n-1}$), then $H_n(B^n) = 0$ but $H_n(B^n, \mathbf{S}^{n-1}) = \mathbf{Z}$. Roughly speaking, the relative homology in this case has the effect of collapsing the boundary to a point, in such a way that B^n is transformed into a sphere.

(d) $H_n(\mathbf{R}^n, \mathbf{R}^n \setminus \{0\}) = \mathbf{Z}$.

(e) If p is an element of $\mathbf{S}^n$, then $H_n(\mathbf{S}^n, \{p\}) = \mathbf{Z}$.

These examples are quite well known (see [Ma], especially p.186-188), but we shall take a closer look at them in the course of this review of basic properties of the homology groups.

If X and Y are topological spaces and $f : X \rightarrow Y$ is a continuous mapping then f induces a group homomorphism $f_* : H_n(X) \rightarrow H_n(Y)$ on the homology groups. The

identity mapping on X induces the identity mapping on $H_n(X)$. If $g : Y \rightarrow Z$ is another continuous mapping, then $(f \circ g)_* = f_* \circ g_*$. In particular homeomorphisms between topological spaces induce isomorphisms between homology groups. If f_0 and f_1 are continuous maps from X to Y which are homotopic, then $(f_0)_* = (f_1)_*$. (See p.163-168 of [Ma].)

Note that constant maps induce the trivial homomorphism (with image = $\{0\}$) in homology with positive dimension. The homotopy property then implies that contractible spaces (like balls and Euclidean spaces) have trivial homology in positive dimensions.

There are analogous statements for relative homology and pairs of spaces. By a *pair of spaces* (X, A) we mean a topological space X and a subspace A . Suppose that (X, A) and (Y, B) are two pair of spaces and $f : X \rightarrow Y$ is continuous. If also $f(A) \subseteq B$, then we say that f maps (X, A) into (Y, B) , and we write $f : (X, A) \rightarrow (Y, B)$. In this case f induces a homomorphism $f_* : H_n(X, A) \rightarrow H_n(Y, B)$. (Beware of this notation, since we are also using f_* to denote the induced homomorphism between $H_n(X)$ and $H_n(Y)$.) If $f(X) \subseteq B$, then f_* is identically zero, because of the way that the relative homology group $H_n(Y, B)$ is defined (it throws the pieces that live wholly in B to the garbage). If $g : (Y, B) \rightarrow (Z, C)$ is also continuous, then $(f \circ g)_* = f_* \circ g_*$. If $f_0, f_1 : (X, A) \rightarrow (Y, B)$ are two continuous mappings which are homotopic as maps from (X, A) into (Y, B) (so that the intermediate maps in the homotopy also send A to B), then we have again the equality $(f_0)_* = (f_1)_*$ between the induced homomorphisms on the relative homology groups. (See p.173-174 of [Ma].)

In short, mappings between topological spaces induce mappings between the homology groups, with the expected functorial properties. Let us consider some concrete examples. The inclusion map $\mathbf{S}^{n-1} \rightarrow \mathbf{R}^n \setminus \{0\}$ induces a homomorphism $H_j(\mathbf{S}^{n-1}) \rightarrow H_j(\mathbf{R}^n \setminus \{0\})$ for each j . Similarly, the usual radial projection $\mathbf{R}^n \setminus \{0\} \rightarrow \mathbf{S}^{n-1}$ induces a homomorphism $H_j(\mathbf{R}^n \setminus \{0\}) \rightarrow H_j(\mathbf{S}^{n-1})$. These two homomorphisms are inverses of each other. The composition of the inclusion with the radial projection is just the identity on $\mathbf{S}^{n-1}$, and so the composition of the induced homomorphisms on homology also equals the identity. The composition of the radial projection with the inclusion of the sphere (in the other order) gives a mapping from $\mathbf{R}^n \setminus \{0\}$ into itself which is not the identity – it looks like the radial projection again – but it is homotopic to the identity, and so the induced homomorphism on homology is again the identity. This proves that our original induced homomorphisms on homology are inverses of each other.

For an example for relative homology, let B, B' be (standard) balls in $\mathbf{R}^n$ with $B' \subseteq B$, and consider the inclusion $(\mathbf{R}^n, \mathbf{R}^n \setminus B) \rightarrow (\mathbf{R}^n, \mathbf{R}^n \setminus B')$. This induces a homomorphism

$$(A.2) \quad H_j(\mathbf{R}^n, \mathbf{R}^n \setminus B) \rightarrow H_j(\mathbf{R}^n, \mathbf{R}^n \setminus B')$$

on the relative homology groups. This homomorphism is actually an isomorphism. Indeed, the pairs $(\mathbf{R}^n, \mathbf{R}^n \setminus B)$ and $(\mathbf{R}^n, \mathbf{R}^n \setminus B')$ are homeomorphic, and one can check that the inclusion is homotopic to such a homeomorphism as a map between pairs. Suppose now that p is an element in B , and consider the homomorphism

$$(A.3) \quad H_j(\mathbf{R}^n, \mathbf{R}^n \setminus B) \rightarrow H_j(\mathbf{R}^n, \mathbf{R}^n \setminus \{p\})$$

induced by the obvious inclusion. This is an isomorphism too, but it does not quite follow from the facts mentioned above. Using the definition of the relative homology

groups it is easy to check that if $\beta \in H_j(\mathbf{R}^n, \mathbf{R}^n \setminus \{p\})$, then β is the image of some α in some $H_j(\mathbf{R}^n, \mathbf{R}^n \setminus B')$ (where B' is a sufficiently small ball that contains p) under the homomorphism induced by inclusion. This really is just a question of chasing definitions, and it is true at the level of chains. The point is that, for a given β , we only have to deal with finitely many singular j -cubes, each of which either touches p or does not, and the finitely many which do not must also lie in the complement of some small ball B' which contains p . Once we have our α in some $H_j(\mathbf{R}^n, \mathbf{R}^n \setminus B')$ we know that it really has to come from an element of $H_j(\mathbf{R}^n, \mathbf{R}^n \setminus B)$, since we know already that (A.2) is an isomorphism. This proves the surjectivity of (A.3). As for injectivity, a similar argument implies that if an element α in $H_j(\mathbf{R}^n, \mathbf{R}^n \setminus B)$ should become zero when mapped into $H_j(\mathbf{R}^n, \mathbf{R}^n \setminus \{p\})$ – i.e., that it could be represented as a boundary of a chain in $(\mathbf{R}^n, \mathbf{R}^n \setminus \{p\})$ – then it would have to do that already in $H_j(\mathbf{R}^n, \mathbf{R}^n \setminus B')$ for some small ball B' containing p (because of the definition of $(\mathbf{R}^n, \mathbf{R}^n \setminus \{p\})$, as before). This is impossible, because of (A.2). This proves that (A.3) is an isomorphism. (We shall encounter another version of this isomorphism in (A.7) below.)

The isomorphism in (A.3) will be useful for us because it provides a way to understand $H_j(\mathbf{R}^n, \mathbf{R}^n \setminus \{p\})$ as we move p .

Given any pair of spaces (X, A) the inclusion from A into X induces a homomorphism from $H_n(A)$ into $H_n(X)$. We also have a homomorphism from $H_n(X)$ into $H_n(X, A)$. This homomorphism comes directly from the definition of $H_n(X, A)$, but one can also see it as the homomorphism induced by the identity as a mapping from $(X, \emptyset)$ into (X, A) . It turns out that there is a canonical but slightly less trivial homomorphism from $H_n(X, A)$ into $H_{n-1}(A)$ which makes the sequence

$$(A.4) \quad H_n(A) \rightarrow H_n(X) \rightarrow H_n(X, A) \rightarrow H_{n-1}(A) \rightarrow H_{n-1}(X)$$

exact. (See Theorem A.1 on p.171 of [Ma].) To be honest, this is not correct when $n = 1$, although there is a natural way to fix it, and one also has to make some decisions for $n = 0$. We shall not go into these matters because $n > 1$ in all the cases that we really care about in this paper.

The mapping from $H_n(X, A)$ into $H_{n-1}(A)$ in (A.4) is not very difficult to describe. Any homology class in $H_n(X, A)$ is represented by some singular n -chain ξ in X . Consider the boundary β of this chain. Then β lives in A ; this comes from the definition of $H_n(X, A)$. This class is closed, since it is a boundary, and so it defines an element of $H_{n-1}(A)$. This element of $H_{n-1}(A)$ turns out to be independent of the representative ξ of our original homology class in $H_n(X, A)$. Note that β need not be a boundary as a chain in A , even though it is a boundary as a chain in X , which is why it can represent a nontrivial element of $H_{n-1}(A)$.

As a simple illustration of this exact sequence, note that if p is an element of $\mathbf{S}^n$, then the induced homomorphism

$$(A.5) \quad H_n(\mathbf{S}^n) \rightarrow H_n(\mathbf{S}^n, \{p\})$$

is an isomorphism. This comes from the exactness of (A.4) and the fact that $H_n(\{p\}) = H_{n-1}(\{p\}) = 0$ when $n > 1$. When $n = 1$ one has to modify (A.4), but (A.5) remains an isomorphism. When $n = 0$ (A.5) is not an isomorphism.

Similarly, if q is any element of $\mathbf{S}^n$, then the homomorphism

$$(A.6) \quad H_n(\mathbf{S}^n) \rightarrow H_n(\mathbf{S}^n, \mathbf{S}^n \setminus \{q\})$$

induced by inclusion is an isomorphism when $n > 1$, because $\mathbf{S}^n \setminus \{q\}$ is a topological n -ball and $H_n(\mathbf{S}^n \setminus \{q\}) = H_{n-1}(\mathbf{S}^n \setminus \{q\}) = 0$. When $n = 1$ (A.6) is still an isomorphism but one must modify the argument, and (A.6) is not an isomorphism when $n = 0$. By combining (A.5) and (A.6) we get that the homomorphism

$$(A.7) \quad H_n(\mathbf{S}^n, \{p\}) \rightarrow H_n(\mathbf{S}^n, \mathbf{S}^n \setminus \{q\})$$

induced by inclusion is an isomorphism when $n > 1$ (and also when $n = 1$), as long as p and q are distinct points in $\mathbf{S}^n$.

Another simple example is provided by B_n and $\mathbf{S}^{n-1}$ as in (A.1) (c) above. In this case we get that the natural homomorphism

$$(A.8) \quad H_n(B_n, \mathbf{S}^{n-1}) \rightarrow H_{n-1}(\mathbf{S}^{n-1})$$

in the exact sequence is an isomorphism when $n > 1$, since $H_n(B_n) = H_{n-1}(B_n) = 0$. (Note that we do not get an isomorphism in (A.8) when $n = 1$.) Similarly, the homomorphism

$$(A.9) \quad H_n(\mathbf{R}^n, \mathbf{R}^n \setminus \{0\}) \rightarrow H_{n-1}(\mathbf{R}^n \setminus \{0\})$$

from the exact sequence is an isomorphism when $n > 1$ because $H_n(\mathbf{R}^n) = H_{n-1}(\mathbf{R}^n) = 0$. (Again the $n = 1$ version of this is wrong). Because $\mathbf{R}^n \setminus \{0\}$ is homotopy equivalent to $\mathbf{S}^{n-1}$ we have that $H_{n-1}(\mathbf{R}^n \setminus \{0\})$ is isomorphic to $H_{n-1}(\mathbf{S}^{n-1})$.

The next result describes the *excision* property of homology.

Theorem A.10. *Let (X, A) be a pair of spaces, and let W be a subset of A such that $\overline{W}$ is contained in the interior of A . Consider the inclusion map $(X \setminus W, A \setminus W) \rightarrow (X, A)$ and the induced homomorphism*

$$(A.11) \quad H_n(X \setminus W, A \setminus W) \rightarrow H_n(X, A).$$

This induced homomorphism is an isomorphism.

This is Theorem 6.2 on p.175 in [Ma]. If one thinks of $H_n(X, A)$ as the homology of X with everything in A thrown to the garbage, then the excision property says that it does not matter whether one cuts out W before or after A is thrown to the garbage when W lives well inside A .

Here is a nice dumb example to illustrate excision. If V is any subset of $\mathbf{R}^n$ which contains the origin 0 in its interior, then the natural inclusion from $(\mathbf{R}^n, \mathbf{R}^n \setminus \{0\})$ to $(V, V \setminus \{0\})$ induces an isomorphism

$$(A.12) \quad H_n(\mathbf{R}^n, \mathbf{R}^n \setminus \{0\}) \rightarrow H_n(V, V \setminus \{0\})$$

between the corresponding relative homology groups. For a slightly fancier example, let M be a topological manifold of dimension n and let p be any point in M . Then

$$(A.13) \quad H_n(M, M \setminus \{p\}) \text{ is isomorphic to } H_n(\mathbf{R}^n, \mathbf{R}^n \setminus \{0\}).$$

Indeed, one can first use excision to say that $H_n(M, M \setminus \{p\})$ is isomorphic to $H_n(U, U \setminus \{p\})$ for any neighborhood U of p in M . Since M is a manifold we can take U to be a topological ball and then reduce to (A.12).

Using some of the isomorphisms above we get that

$$(A.14) \quad H_n(\mathbf{S}^n) \text{ is isomorphic to } H_{n-1}(\mathbf{S}^{n-1})$$

when $n > 1$. Specifically, we use (A.6), (A.13) (with $M = \mathbf{S}^n$), and (A.9) to reduce this assertion to the observation made above that $H_{n-1}(\mathbf{R}^n \setminus \{0\})$ is isomorphic to $H_{n-1}(\mathbf{S}^{n-1})$. Once we have (A.14) we can reduce (A.1) (b) to the $n = 1$ case. Similar methods permit the $n = 1$ case to be reduced to an $n = 0$ problem. Note that the rest of the examples in (A.1) can be derived from (A.1) (b) and the other facts mentioned above, except for some minor changes when $n = 1$.

Before we get to degree theory we need to say a little about the homology of topological manifolds, beginning with the homological definition of an orientation.

Definition A.15. Let M be a topological manifold of dimension n , so that $H_n(M, M \setminus \{p\})$ is isomorphic to $\mathbf{Z}$ for every $p \in M$, because of (A.13) and (A.1) (d). An *orientation* for M is a choice of generator $\lambda_p \in H_n(M, M \setminus \{p\})$ for each $p \in M$ with the following continuity property: for each $p \in M$ there is a neighborhood V of p and an element λ_V of $H_n(M, M \setminus V)$ so that for every $x \in V$ we have that λ_x is the image of λ_V under the homomorphism $H_n(M, M \setminus V) \rightarrow H_n(M, M \setminus \{x\})$ induced by inclusion.

This is the same as Definition 2.1 on p.351 of [Ma]. It is easy to see that if M is oriented in the usual sense (in terms of coordinate charts) then one gets this kind of orientation as well. This statement is a little bit specious because it presumes that one has already a good definition for orientation-preserving homeomorphisms, while in fact homology arguably provides the best way to define the notion of orientation-preserving homeomorphisms. Of course this is much simpler when M is smooth.

Note that an orientation on M induces an orientation on any open subset U of M (which can be viewed as a topological manifold in its own right). This uses the excision principle (Theorem A.10). Also, given an orientation $\{\lambda_p\}$ on M we have that $\{-\lambda_p\}$ is also an orientation for M .

The continuity condition in Definition A.15 is illustrated well by the isomorphism (A.3) for $M = \mathbf{R}^n$. Using excision one can produce similar isomorphisms on any manifold. Let us use this fact to prove that orientations enjoy the appropriate uniqueness properties.

Lemma A.16. *If M is a connected topological manifold, then two orientations which agree at a point agree everywhere. If the two orientations are different then they are negatives of each other.*

At any given point, two orientations either agree or are negatives of each other, because there are only two generators for $\mathbf{Z}$ (+1 and -1). To prove the lemma it suffices to show

that the subset of M where two orientations agree is always an open set. (The set where they disagree is the same as the set where one orientation equals the negative of the other and hence will also be open.) Let $\{\lambda_p\}$ and $\{\lambda'_p\}$ be two orientations of M which agree at the point x . Then there exist neighborhoods V and V' of x and homology classes $\lambda_V \in H_n(M, M \setminus V)$ and $\lambda'_{V'} \in H_n(M, M \setminus V')$ such that λ_p is the image of λ_V in $H_n(M, M \setminus \{p\})$ (under the homomorphism induced by inclusion) when $p \in V$ and similarly for V' . We may shrink V and V' without destroying these properties (we simply have to map the old λ_V into the new relative homology group using inclusion, etc.). This observation permits us to reduce to the case where $V = V'$ and V is a topological ball whose closure is contained in another topological ball in M in a standard way. Using excision (Theorem A.10) we can reduce further to the case where $M = \mathbf{R}^n$ and V is a standard ball in $\mathbf{R}^n$. The isomorphism (A.3) then implies that $\lambda_V = \lambda'_{V'}$, and hence the two orientations have to agree on all of V . Thus the set where the two orientations agree is open, as desired. This proves Lemma A.16.

An orientation on a manifold determines some other, more global, homology classes.

Theorem A.17. *Suppose that M is an n -dimensional topological manifold with orientation $\{\lambda_x\}$ (as defined above). Then for each compact set $K \subseteq M$ there is a unique homology class $\lambda_K \in H_n(M, M \setminus K)$ such that, for every $x \in K$, we have that the image of λ_K under the homomorphism $H_n(M, M \setminus K) \rightarrow H_n(M, M \setminus \{x\})$ induced by inclusion equals λ_x . In particular, if M is compact then there is a homology class $\lambda_M \in H_n(M)$ (the fundamental class) with this property (but with $H_n(M, M \setminus K)$ replaced with $H_n(M)$).*

This is Theorem 2.2 on p.353 of [Ma].

To understand what this means one should start with the case $M = \mathbf{R}^n$. Remember that homology classes are always represented by formal sums of mappings of cubes into our space. There is a lot of algebra around, but it always comes down to cubes. In the case of $\mathbf{R}^n$ we can simply put our given compact set into one big cube, and one gets the homology class λ_K from that cube. The sphere is slightly more complicated; one has to use two cubes. Theorem A.17 says that on any oriented topological manifold one can do something similar with a finite number of cubes (for a given compact subset).

Let us now define the degree of a mapping into the sphere. Fix the dimension n .

Notation A.18. Let ω be the generator of $H_n(\mathbf{S}^n) \simeq \mathbf{Z}$ which corresponds to the usual orientation on $\mathbf{S}^n$. That is, ω induces an orientation $\{\omega_p\}$ of $\mathbf{S}^n$, by taking $\omega_p \in H_n(\mathbf{S}^n, \mathbf{S}^n \setminus \{p\})$ to be the image of ω under the homomorphism on homology induced by inclusion, and we want this induced orientation to be the standard one. (ω_p really is a generator of $H_n(\mathbf{S}^n, \mathbf{S}^n \setminus \{p\})$, as required in Definition A.15, because (A.6) is always an isomorphism.) Equivalently, ω is the fundamental class corresponding to the usual orientation of $\mathbf{S}^n$ (as in Theorem A.17). (The fact that (A.6) is an isomorphism ensures that this fundamental class is a generator of $H_n(\mathbf{S}^n)$.) We shall always use this orientation $\{\omega_p\}$ for $\mathbf{S}^n$, no matter whether $\mathbf{S}^n$ is being used as the domain of a given mapping, the range, or both, unless the contrary is explicitly stated.

This choice of ω can also be formulated as a choice of an isomorphism between $H_n(\mathbf{S}^n)$ and the group of integers $\mathbf{Z}$.

Definition A.19. Let M is a topological manifold of dimension n with orientation $\{\lambda_x\}$ as above, and let $f : M \rightarrow \mathbf{S}^n$ be a continuous mapping. If M is compact then the *degree* of f is defined to be the integer d such that $f_*(\lambda_M) = d \cdot \omega \in H_n(\mathbf{S}^n)$. If M is not compact then assume that f takes a single value p on the complement of a compact subset K of M , so that $f : (M, M \setminus K) \rightarrow (\mathbf{S}^n, \{p\})$. In this case the degree of f is defined to be the integer d such that $f_*(\lambda_K) = d \cdot \omega' \in H_n(\mathbf{S}^n, \{p\})$, where ω' denotes the image of ω in $H_n(\mathbf{S}^n, \{p\})$ under the natural inclusion map. (Note that ω' generates $H_n(\mathbf{S}^n, \{p\})$, since (A.5) is an isomorphism.) Here λ_M, λ_K are as in Theorem A.17.

This definition of the degree does not depend on the choice of K when M is not compact, because $f_*(\lambda_K) \in H_n(\mathbf{S}^n, \{p\})$ does not depend on K , as one can check from the properties of the induced maps on homology. (Note that the f in $f_*(\lambda_K)$ does depend on K , because we are viewing it as map from the pair $(M, M \setminus K)$ into $(\mathbf{S}^n, \{p\})$, and the first pair depends on K . A precise explanation as to why the compatibilities work out correctly would be more difficult to read than to derive for oneself, and so we omit it.) Also, the definition of the degree in the noncompact case also works in the compact case, because (A.5) is an isomorphism.

We can extend the definition of the degree, if we wish, to continuous mappings $f : M \rightarrow \mathbf{S}^n$ which omit some fixed value $q \in \mathbf{S}^n$ on the complement of a compact subset of M (rather than actually being constant on such a region). This follows from the isomorphism (A.7). The various properties of the degree stated below have counterparts for this extension. We won't need this extension directly, but it makes some of the stability properties of the degree more transparent even for mappings which are constant off a compact set.

For the record, note that the identity mapping from $\mathbf{S}^n$ to itself has degree 1, and that constant mappings have degree zero. A mapping from $\mathbf{S}^n$ into itself which simply reflects points about some hyperplane has degree -1 . (See (g) on p.189 of [Ma].)

Let us now record a few facts about the degree.

Lemma A.20. *Let M be an oriented topological manifold of dimension n . If $f : M \rightarrow \mathbf{S}^n$ and $g : \mathbf{S}^n \rightarrow \mathbf{S}^n$ are continuous, and if f is constant off a compact subset of M , then the degree of $g \circ f$ equals the product of the degrees of f and g .*

This follows easily from the definition. Note that we are assuming that the sphere $\mathbf{S}^n$ is given its standard orientation when we use it as the domain of g (as in Notation A.18).

Lemma A.21. *Suppose that M is an oriented topological manifold of dimension n and that $f_0, f_1 : M \rightarrow \mathbf{S}^n$ are continuous mappings which are constant off a compact subset of M and which are homotopic through mappings which are each constant off a (fixed) compact set. Then the degrees of f_0 and f_1 are equal.*

Let f_t , $0 \leq t \leq 1$, be the assumed homotopy of between f_0 and f_1 , and let K be a compact subset of M such that each the f_t is constant off of K . Let $p(t)$ denote the value of f_t on $M \setminus K$. We do not need to require that $p(t)$ be constant, but let us assume for the moment that this is true. In this case the existence of the homotopy implies that the induced mappings $(f_t)_* : H_n(M, M \setminus K) \rightarrow H_n(\mathbf{S}^n, \{p\})$, $p = p(t)$, are constant (Theorem 6.1 on p.174 of [Ma]). This implies that the degrees of f_t are all the same. If $p(t)$ is not constant then we can reduce to the case where it is constant by composing the f_t 's

with a family of rotations, but we can also argue as follows. Suppose first that there is a $q \in \mathbf{S}^n$ such that $p(t) \neq q$ for all t . In this case we have that the induced mappings $(f_t)_* : H_n(M, M \setminus K) \rightarrow H_n(\mathbf{S}^n, \mathbf{S}^n \setminus \{q\})$ are all equal, and we can recover the constancy of the degree from the isomorphism (A.7). In general even if $p(t)$ is some kind of Peano curve it will have the property that it omits a point $q \in \mathbf{S}^n$ on a neighborhood about any $t_0 \in [0, 1]$, and so the preceding argument applies to tell us that the degree of f_t is locally constant in t , which implies that it is constant. This proves Lemma A.21.

Our next goal is to understand how the degree of a mapping can be localized (under certain conditions).

Lemma A.22. *Let M be a topological manifold of dimension n with orientation $\{\lambda_y\}$. Suppose that $f : M \rightarrow \mathbf{S}^n$ is a continuous mapping which is constant off a compact set, and also that there is a point $q \in \mathbf{S}^n$ such that q has only one preimage $x \in M$. Then the degree d is determined by the equation $f_*(\lambda_x) = d \cdot \omega_q$, where f_* denotes the induced homomorphism from $H_n(M, M \setminus \{x\})$ into $H_n(\mathbf{S}^n, \mathbf{S}^n \setminus \{q\})$, and ω_q is as in Notation A.18.*

This follows from the usual compatibility properties of the homomorphisms on homology induced by mappings between spaces. More precisely, if M is compact, then the degree is defined by $f_*(\lambda_M) = d \cdot \omega$, where now $f_* : H_n(M) \rightarrow H_n(\mathbf{S}^n)$. This implies the formula in the lemma because of the definition of ω_q (in Notation A.18) and the fact that λ_x is the image of λ_M in $H_n(M, M \setminus \{x\})$ under the homomorphism induced by inclusion. (We are also using the compatibility of compositions of mappings between topological spaces with the various induced homomorphisms on homology.) The noncompact case is similar. (It uses also the trivial observation that if $p \in \mathbf{S}^n$, $p \neq q$, and ω' is the image of ω in $H_n(\mathbf{S}^n, \{p\})$ under the natural inclusion map, then ω_q is the image of ω' under the homomorphism (A.7) induced by inclusion.) This proves Lemma A.22.

Lemma A.23. *If M is an oriented topological manifold of dimension n and $f : M \rightarrow \mathbf{S}^n$ is a continuous mapping which is constant off a compact set and omits some value $q \in \mathbf{S}^n$, then f has degree 0.*

This can be proved in approximately the same way as Lemma A.22, or it can be derived from Lemma A.21. (That is, if f omits a point in $\mathbf{S}^n$, then it can be homotoped to a constant.)

The next result is a version of the excision principle for the degree.

Proposition A.24. *Suppose that M is an oriented topological manifold of dimension n , U is an open subset of M , K is a compact subset of U , and $f : M \rightarrow \mathbf{S}^n$ is continuous. Suppose also that f is constant on $M \setminus K$. Then the degree of f as a mapping on M equals the degree of f as a mapping on U (where U is viewed as a topological manifold in its own right, with the orientation inherited from M).*

This is an easy consequence of the definitions and the excision principle Theorem A.10. More precisely, the excision principle implies that the natural map from $H_n(U, U \setminus \{x\})$ to $H_n(M, M \setminus \{x\})$ induced by inclusion is an isomorphism for every $x \in U$. (In fact one uses this isomorphism to make sense of the orientation on U inherited from M .) The uniqueness part of Theorem A.17 implies that the relative homology class $\lambda_K \in H_n(M, M \setminus K)$ corresponds under the obvious inclusion map to its counterpart for U . (That is, the former

is the image of the latter under the homomorphism $H_n(U, U \setminus K) \rightarrow H_n(M, M \setminus K)$ induced by inclusion.) The equality of the degrees follows from this correspondence and ridiculous nonsense (we factor $f : (U, U \setminus K) \rightarrow (\mathbf{S}^n, \dots)$ into the inclusion $\iota : (U, U \setminus K) \rightarrow (M, M \setminus K)$ and $f : (M, M \setminus K) \rightarrow (\mathbf{S}^n, \dots)$, and then use the corresponding maps on homology). This proves Proposition A.24.

Next we consider the situation where the manifold and mapping can be decomposed into a finite number of separate pieces.

Lemma A.25. *Suppose that M is an oriented topological manifold of dimension n which is a finite union of disjoint open subsets U_i , and that $f : M \rightarrow \mathbf{S}^n$ is a continuous map which is constant off of a compact subset of each U_i . Then the degree of f as a map on M equals the sum of the degrees of the restrictions of f to the various U_i 's (when viewed as topological manifolds in their own right, with the inherited orientations).*

This is a trivial consequence of the definition of the homology groups, which ensures that homology and relative homology groups on M split into direct sums of homology and relative homology groups on the U_i 's in the obvious way.

In practice we can sometimes compute the degree of a mapping by using Proposition A.24 and Lemma A.25 to reduce to a finite number of pieces, each of which satisfies the hypotheses of Lemma A.22.

It is a pleasant exercise to compare these various properties of the degree with the discussion of the degree of smooth mappings at the beginning of this appendix. That is, if one takes that definition of the degree then many of the results just stated are very clear (like Proposition A.24 and Lemma A.25). Conversely, one can use these results to check that that definition of the degree for smooth mappings agrees with the homological definition.

The next two lemmas provide useful building blocks for constructing maps with good properties.

Lemma A.26. *Suppose that $p \in \mathbf{S}^n$ and V is a neighborhood of p in $\mathbf{S}^n$. Then there is a smooth mapping from $\mathbf{S}^n$ into itself which is homotopic to the identity, equals the identity on a neighborhood of p , sends no other point to p besides p , and sends $\mathbf{S}^n \setminus V$ to $-p$. We can even choose this homotopy so that all the intermediate maps send $-p$ to itself. There also exist smooth mappings from $\mathbf{S}^n$ into itself with any prescribed integer degree. We can choose these mappings so that they fix some prescribed point $\sigma \in \mathbf{S}^n$.*

This is an easy exercise. We may assume that V is a round neighborhood (the intersection of $\mathbf{S}^n$ with a ball centered at p), because we can always shrink it if necessary. We can build the mapping to be a homeomorphism from V onto $\mathbf{S}^n \setminus \{-p\}$ which is obtained "radial" expansion along the great circles on $\mathbf{S}^n$ which emanate from p , and we can do this expansion of V in such a way that the mapping is homotopic to the identity in the required manner. We can also keep everything fixed on a neighborhood of p . Note that this mapping has degree 1, since it is homotopic to the identity, and also because of Lemma A.22. We can get a mapping of degree -1 by composing the preceding map with a reflection across a hyperplane. (Such a reflection has degree -1 , as in (g) on p.189 of [Ma].)

To get mappings of any integer degree one builds mappings which are constant off a finite union of disjoint closed balls and which agree with a mapping of the type just

constructed (or its reflection) on each little ball. The degree of such a map reduces to the sum of the individual degrees by Proposition A.24 and Lemma A.25, and hence it can be arranged to be any given integer. One can also do this in such a way that a given point σ is fixed. This proves Lemma A.26.

Lemma A.27. *Suppose that M is an oriented topological manifold of dimension n , and let U be any nonempty open subset of M . Let $\sigma \in \mathbf{S}^n$ be given. Then there is a continuous mapping $f : M \rightarrow \mathbf{S}^n$ which takes only the value σ outside U and which has degree 1.*

By shrinking U if necessary we may assume that it is a topological ball whose closure in M is a compact set. Let $f : M \rightarrow \mathbf{S}^n$ be a continuous mapping which takes the value σ on $M \setminus U$ and which restricts to a homeomorphism from U onto $\mathbf{S}^n \setminus \{\sigma\}$. Given any $x \in U$, we have that the homomorphism $H_n(U, U \setminus \{x\}) \rightarrow H_n(\mathbf{S}^n \setminus \{\sigma\}, \mathbf{S}^n \setminus \{\sigma, f(x)\})$ induced by f is an isomorphism, because f restricts to a homeomorphism from $(U, U \setminus \{x\})$ onto $(\mathbf{S}^n \setminus \{\sigma\}, \mathbf{S}^n \setminus \{\sigma, f(x)\})$. Using excision (Theorem A.10) twice we conclude that the homomorphism $H_n(M, M \setminus \{x\}) \rightarrow H_n(\mathbf{S}^n, \mathbf{S}^n \setminus \{f(x)\})$ induced by f is an isomorphism. Lemma A.22 now implies that f has degree equal to $+1$ or -1 . If it is the latter then we can compose it with a reflection on the sphere to get a mapping of degree 1. This proves Lemma A.27.

The remaining facts that we shall need about homology and degree concern connected sets.

Lemma A.28. *Suppose that M is an n -dimensional topological manifold with orientation $\{\lambda_x\}$. If K is a compact connected nonempty subset of M , then $H_n(M, M \setminus K)$ is an infinite cyclic group, and the homology class $\lambda_K \in H_n(M, M \setminus K)$ from Theorem A.17 is a generator for $H_n(M, M \setminus K)$.*

We know that $H_n(M, M \setminus K)$ is nonzero because it contains λ_K , and so it is infinite cyclic by Exercise 2.5 (b) on p.357 of [Ma]. The second part of that exercise says that for each $x \in K$ the homomorphism from $H_n(M, M \setminus K)$ into $H_n(M, M \setminus \{x\})$ induced by inclusion is injective. Since this homomorphism sends λ_K to λ_x and the latter generates $H_n(M, M \setminus \{x\})$, we conclude that λ_K generates $H_n(M, M \setminus K)$.

The next result provides a converse to Lemma A.21.

Proposition A.29. *Let M be an oriented topological manifold of dimension n , and let U be a connected open subset of M whose closure in M is compact. Let $\sigma \in \mathbf{S}^n$ be given, and let $f_1, f_2 : M \rightarrow \mathbf{S}^n$ be continuous mappings which each take only the value σ on $M \setminus U$. If f_1 and f_2 have the same degree, then they are homotopic through continuous mappings from M into $\mathbf{S}^n$ which all equal σ on $M \setminus U$.*

This is well-known, but we include a proof for the sake of completeness. For this proof we shall assume that there is a metric $d(x, y)$ on U which defines the same topology on U as the one inherited from M and for which the n -dimensional Hausdorff measure of U with respect to this metric is finite. (See (2.14) for the definition of Hausdorff measure.) This is not truly an additional assumption, but it makes the argument simpler. The main purpose of including a proof of Proposition A.29 is to provide the reader with an argument that does not require much topological machinery, and for this purpose the extra assumption is not too onerous.

From now on we let M, U, σ, f_1 , and f_2 be as in the proposition.

Lemma A.30. *Proposition A.29 is true when U is homeomorphic to the unit ball in $\mathbf{R}^n$.*

Let $h : M \rightarrow \mathbf{S}^n$ be a continuous mapping which sends $M \setminus U$ to σ and which restricts to be a homeomorphism from U onto $\mathbf{S}^n \setminus \{\sigma\}$. We can factor f_1, f_2 by h , i.e., we can find continuous mappings $f'_1, f'_2 : \mathbf{S}^n \rightarrow \mathbf{S}^n$ which send σ to itself and satisfy $f_i = f'_i \circ h$. Lemma A.20 implies that the degree of f_i equals the degree of f'_i times the degree of h , $i = 1, 2$. The degree of h is either 1 or -1 , because of Lemma A.22 and the fact that h is a homeomorphism on U . (Specifically, for each $x \in U$, the homomorphism $H_n(U, U \setminus \{x\}) \rightarrow H_n(\mathbf{S}^n \setminus \{\sigma\}, \mathbf{S}^n \setminus \{\sigma, h(x)\})$ induced by h is an isomorphism, because $h : (U, U \setminus \{x\}) \rightarrow (\mathbf{S}^n \setminus \{\sigma\}, \mathbf{S}^n \setminus \{\sigma, h(x)\})$ is a homeomorphism. The excision principle (Theorem A.10) can be used to show the homomorphism $H_n(M, M \setminus \{x\}) \rightarrow H_n(\mathbf{S}^n, \mathbf{S}^n \setminus \{h(x)\})$ induced by h is also an isomorphism. The statement about the degree of h now follows from Lemma A.22.) Thus we conclude that these auxiliary mappings f'_1, f'_2 have the same degree.

This passage from f_i to f'_i amounts to reducing Lemma A.30 to the case of the sphere. In the case of the sphere Lemma A.30 is equivalent to the statement that the homotopy class of a continuous mapping on the sphere (with basepoint σ) is determined by its degree. This is well known; it is mentioned in (a) on p.189 of [Ma], it is stated and the proof is discussed in Lemma 17.10.2 on p.215 of [BT], and the original paper is apparently [Wh]. (In some formulations the basepoint is dropped, but this is not a serious issue.) This proves Lemma A.30.

Lemma A.31. *In order to prove Proposition A.29 we may assume that f_1 and f_2 take only the value σ off a compact subset of U , and that $f_i^{-1}(p)$ is finite for $i = 1, 2$ and some $p \in \mathbf{S}^n \setminus \{\sigma\}$.*

Let θ be a map from $\mathbf{S}^n$ to itself which maps a neighborhood of σ onto σ and which is homotopic to the identity through maps which send σ to itself. Such a mapping exists, by Lemma A.26. We can reduce to the case where f_1 and f_2 take only the value σ off a compact subset of U by replacing them with $\theta \circ f_1$ and $\theta \circ f_2$ if necessary. This uses Lemma A.21. (We are also using here the fact that a closed subset of M which does not intersect $M \setminus U$ must be compact, since the closure of U in M is compact.)

Next, we may also assume that f_1 and f_2 are Lipschitz with respect to our metric $d(x, y)$, because we can use Lemma 2.12 to deform them to Lipschitz mappings through mappings which take only the value σ off a compact subset of U . The Lipschitzness of f_1, f_2 and our assumption of finite n -dimensional Hausdorff measure imply that $f_i^{-1}(p)$ is finite for almost all $p \in \mathbf{S}^n$, by a well-known fact. (See Theorem 2.10.25 on p.188 of [Fd] or Proposition 14.1 above (with $l = 0$).) This proves Lemma A.31.

Lemma A.32. *Suppose that $f : M \rightarrow \mathbf{S}^n$ is a continuous mapping which takes only the value σ on $M \setminus U$ and that $f^{-1}(p)$ is finite for some $p \in \mathbf{S}^n$. Let V be any nonempty open subset of U . Then there is a continuous map $\tilde{f} : M \rightarrow \mathbf{S}^n$ which takes only the value σ on $M \setminus V$ and which is homotopic to f through mappings which take only the value σ on $M \setminus U$.*

Proposition A.29 will follow once we have this. Indeed, we first use Lemma A.31 to reduce to the case where f_1 and f_2 satisfy the hypotheses of Lemma A.32. We then apply

Lemma A.32 with V taken to be a topological ball. This permits us to reduce Proposition A.29 to the case where U is a topological ball, and this case is covered by Lemma A.30.

Let us now prove Lemma A.32. Let f and p be as above.

Given a mapping from M into the sphere $\mathbf{S}^n$, let us call the closure of the set where it does not take the value σ the *support* of the mapping.

If W is an open subset of M which contains $f^{-1}(p)$, then we can deform f to a continuous mapping which is supported on W and which takes the value p at exactly the same points that f does. To see this we observe first that our assumptions imply that $f(M \setminus W)$ is a compact subset of the sphere which does not contain p . Hence there is a mapping $\theta : \mathbf{S}^n \rightarrow \mathbf{S}^n$ which sends $f(M \setminus W)$ to σ and which is homotopic to the identity through maps which fix the point σ . We may also require that p be the only point sent to p by θ . Such a mapping is easy to construct, as in Lemma A.26.

Thus we may deform f inside of its homotopy class in such a way that we shrink the support of f to an arbitrarily small neighborhood of $f^{-1}(p)$ while keeping the preimages of p fixed. Let us call this “the shrinking argument”.

Sublemma A.33. *Notation and assumptions as above. Suppose that $x \in f^{-1}(p)$, $y \in M \setminus f^{-1}(p)$, and γ is a curve in M which connects x to y but does not touch any element of $f^{-1}(p)$ besides x . Then we can homotop f to a continuous mapping $g : M \rightarrow \mathbf{S}^n$ in such a way that $g^{-1}(p) = (f^{-1}(p) \setminus \{x\}) \cup \{y\}$, and so that the intermediate mappings in the homotopy are all supported in the union of the support of f and any prescribed neighborhood of γ (no matter how small).*

In other words, we can move points in $f^{-1}(p)$ around, so long as we move them continuously and we don't let them collide with each other.

To prove this consider first the case where $f^{-1}(p)$ contains only the one element x . In fact, let us assume for the moment that $M = \mathbf{R}^n$. Think of f (symbolically) as being some kind of bump on $\mathbf{R}^n$, and the lemma says that we can slide that bump along γ . On $\mathbf{R}^n$ we can do this sliding in the simplest possible way, by just translating x along γ and translating f along with it. We can control the support of this sliding by applying the shrinking argument at the beginning to make sure that f is supported in an arbitrarily small neighborhood of x . If we do this then all the intermediate maps obtained by sliding f along γ will be supported in an arbitrarily small neighborhood of γ .

Next we continue to assume that $f^{-1}(p)$ contains only the one element x , but now we let M be any topological manifold. If γ lies in a single coordinate patch of M , then we can reduce to the case where $M = \mathbf{R}^n$ using these coordinates. (We also use the shrinking argument, if necessary, to get the support of f to lie within this coordinate patch.) In general γ can be decomposed into a finite collection of arcs, each of which lies in a coordinate patch. We can apply the same argument to each arc separately, using the shrinking argument if necessary at the end of any intermediate arc to squeeze the support of the map into the coordinate patch of the next arc. We only have to do this a finite number of times, and so we can get the deformation promised in Sublemma A.33 in the case where x is the only element of $f^{-1}(p)$. In particular we can keep the support of the deformation of f as close to γ as we like, except at the beginning where the support of our original f is simply given.

The general case follows from the same argument as when $f^{-1}(p)$ contains only x . We first apply the shrinking argument to reduce to the case where the support of f is contained in a very small neighborhood of $f^{-1}(p)$. Specifically, we can reduce to the case where γ does not touch the pieces of the support of f near the elements of $f^{-1}(p) \setminus \{x\}$, and where the part of the support of f near x lies in an arbitrarily small neighborhood of x . We then apply the method of the previous case to slide the part of f supported near x along the curve γ , in such a way that the part of f that is sliding along γ is supported in an arbitrarily small neighborhood of our curve γ , and the parts of f which live near the elements of $f^{-1}(p) \setminus \{x\}$ are not changed or affected. This deformation gives the mapping g required in Sublemma A.33.

Although this argument is more descriptive than precise, it can be made precise with a lot of notation which is not pleasant to read, and so we leave the details to the reader. This concludes the proof of Sublemma A.33.

Sublemma A.33 is the main ingredient in the proof of Lemma A.32. There is one remaining point, which is the following.

Sublemma A.34. *Let $x, y \in M$ be given, $x \neq y$, and let F be any finite subset of M which does not contain x or y . Then there is a curve γ in U which connects x to y and does not touch any element of F .*

We are already assuming that U is open and connected, so that it is automatically arcwise connected. We simply need to make the arcs avoid the finite set F . This is easy to accomplish. Given $u \in F$ let $D(u)$ be a very small topological ball in M which contains u and whose boundary is a topological sphere. We can do this because M is a topological manifold. We can choose these topological balls so that they have disjoint closures which are disjoint from x and y . Let γ_0 be a curve in U which connects x to y . If γ_0 wanders into some $D(u)$ then we take the first and last points in γ_0 which lie in $D(u)$ and replace the arc which connects them with an arc in $\partial D(u)$. If we do this for each $u \in F$, then in the end we get a curve γ with the correct properties. This proves Sublemma A.34.

We can now finish the proof of Lemma A.32. We can use the shrinking argument to reduce to the case where f is supported in a small neighborhood of the finite set $f^{-1}(p)$. Sublemma A.33 permits us to deform f so that $f^{-1}(p)$ is moved to any other subset of U with the same cardinality, using the curves provided by Sublemma A.34. In particular we can deform f so that $f^{-1}(p)$ is moved into V . Sublemmas A.33 and A.34 permit us to make this deformation while keeping the supports of the intermediate mappings inside U , and we can use the shrinking argument to ensure that our final map is supported in V . This proves Lemma A.32.

The proof of Proposition A.29 is now complete.

Remark A.35. In many of the results in this paper we have assumed that our underlying topological space M is an oriented topological manifold of dimension n . This assumption is much stronger than we really need. Basically we just need for M to have topological dimension n and that a suitable version of degree theory should work for M . This means that M should admit a system of homology classes $\lambda_K \in H_n(M, M \setminus K)$ which satisfy the obvious compatibility conditions under the homomorphisms induced by inclusions, that the conclusion of Lemma A.27 should hold, with the degree of mappings into $\mathbf{S}^n$ being defined

using the homology classes λ_K , and that the conclusions of Lemma A.28 and Proposition A.29 should also hold. (In the case of Lemma A.27 it would be enough to simply know that the mappings have nonzero degree instead of all having degree 1. It would be even better to forget about Lemma A.27 and follow the method of Remark 8.26 instead.)

B. Cleaner statements about Sobolev and Poincaré inequalities.

We saw in Theorems 1.22 and 1.27 how the constructions in this paper lead to estimates for a function in terms of its gradient. Conclusions of this type are simpler and easier to obtain under stronger “uniform” assumptions about our underlying metric space. In this appendix we shall make such assumptions and discuss the Sobolev and Poincaré inequalities that follow from them. The reader should keep in mind that we are aiming here more for simplicity than sharp or optimally general results, and that we are throwing away some of the delicacy and flexibility of the constructions.

We shall discuss in Remark B.20 a different approach due to Gromov for obtaining Sobolev inequalities.

Let us think for a moment about the kind of assumptions that were made in Section 1. They basically came in three types. We assumed some non-quantitative conditions on the topology of our metric space, i.e., we required that it be an oriented topological manifold. This assumption could be weakened considerably, as in Remark A.35 and the comments that followed Standard Assumptions 1.8, but for this appendix we shall not worry about that. The second kind of assumption controlled the size of our metric space. In this category one should include bounds on the covering function (Definition 1.6) and assumptions on the Hausdorff measure of our metric space. The third kind of assumption entailed quantitative bounds on the contractability function (Definition 1.7) of our space.

What are uniform versions of these assumptions? The first assumption, the non-quantitative restrictions on the topology, we shall simply leave alone. For the quantitative condition that we imposed on the topology, namely bounds on the contractability function, the uniform version that we shall use is the local linear contractability condition in Definition 1.15. (Note that Definition 1.15 does not require any bounds on the moduli of continuity on the topological contractions given there.) To control the size of our metric space we shall use (Ahlfors) regularity.

Definition B.1. A metric space $(M, d(x, y))$ is said to be *regular* with dimension $n > 0$ if it is complete and if there is a Borel measure μ on M and a constant $C_1 > 0$ such that

$$(B.2) \quad C_1^{-1} r^n \leq \mu(B(x, r)) \leq C_1 r^n$$

for all $x \in M$ and $0 < r \leq \text{diam } M$.

This can be viewed as a uniform and scale-invariant version of the requirement that M have Hausdorff dimension equal to n . If a metric space is regular with dimension n , then it does have Hausdorff dimension equal to n , and it satisfies (B.2) is satisfied with μ taken to be n -dimensional Hausdorff measure (but perhaps with a different constant). In fact regularity implies that μ and the Hausdorff measure are each bounded by a constant times the other. See Lemma B.5 below.

Examples B.3. (a) $M = \mathbf{R}^n$ is regular with dimension n and μ taken to be Lebesgue measure.

(b) Let us say that $(M, d(x, y), \mu)$ is Riemannian if M is a smooth Riemannian manifold and $d(x, y)$ and μ are the Riemannian metric and volume, respectively. If $(M, d(x, y), \mu)$ is Riemannian and M is compact then M is regular, but with a constant that depends on M and its Riemannian metric. The inequality (B.2) will hold asymptotically as $r \rightarrow 0$ with a constant C_1 that depends only on the dimension, but for r 's of definite size (B.2) is not automatic.

(c) The usual self-similar fractals, like Cantor sets, snowflake curves, trees, and Sierpinski carpets, are all regular, with self-similar measures μ which are usually obvious. Self-similar Cantor sets can have any dimension in $(0, \infty)$.

(d) The product of two regular metric spaces is also regular, at least if they are both bounded or both unbounded. The regularity constant C_1 will be large if the diameters of the two spaces are very different.

(e) (The snowflake functor.) If $(M, d(x, y))$ is a regular metric space with dimension n and $0 < s < 1$, then $(M, d(x, y)^s)$ is a regular metric space of dimension $\frac{n}{s}$.

(f) The Heisenberg group, equipped with its Carnot metric, is regular. For this one takes μ to be Lebesgue measure, and the regularity dimension is 1 larger than the topological dimension.

(g) If $\{S_j\}$ is a sequence of n -dimensional spheres in some fixed Euclidean space, with the radius of each S_j equal to 2^{-j} , and if S_j intersects S_{j+1} for each j , then the union of all the S_j 's together with their limit point is regular with dimension n (when equipped with the Euclidean metric).

Regular metric spaces are special cases of "spaces of homogeneous type" as in [CW1], [CW2]. Many aspects of real analysis work on them, e.g., covering lemmas, Lebesgue points, integral operators, Hardy spaces and BMO. Nothing like Sobolev embeddings are true in general for regular metric spaces, because they can be Cantor sets or snowflakes.

Let us think about which of the examples of regular metric spaces above are also locally linearly contractable. Euclidean spaces and compact smooth manifolds are locally linearly contractable. Cantor sets and Sierpinski carpets are definitely not. The standard self-similar fractal trees and snowflake curves are locally linearly contractable, as is the Heisenberg example (1.6) (f). (Thus there are "fractals" which are both regular and locally linearly contractable.) Local linear contractability is undisturbed by the snowflake functor (1.6) (e). It does prevent some bad behavior, like cusps, bubbles, long thin tubes, and tiny handles. Example (B.3) (g) is not locally linearly contractable.

The next two lemmas contain some basic facts about regular metric spaces.

Lemma B.4. *Suppose that $(M, d(x, y), \mu)$ is a regular metric space of dimension n . Then there is a constant C so that for any ball B in M with radius $R > 0$ and any $r \in (0, R)$ there is a covering of B by a family of at most $C(R/r)^n$ balls in M of radius r . In particular it is doubling (Definition 1.14), and closed and bounded subsets of M are compact.*

Lemma B.5. *If $(M, d(x, y), \mu)$ is a regular metric space of dimension n , then there is a constant $C > 0$ so that $C^{-1}\mu(E) \leq H^n(E) \leq C\mu(E)$ for all Borel sets $E \subseteq M$.*

Remember that H^n denotes n -dimensional Hausdorff measure, defined in (2.14).

We postpone the proofs of these two lemmas until the beginning of Appendix C (Lemmas C.1 and C.3).

Let us now formulate a version of Theorem 1.11 under these stronger assumptions.

Theorem B.6. *Assume that*

- (B.7) *$(M, d(x, y), \mu)$ is a regular metric space of dimension n (Definition B.1) which is also locally linearly contractable (Definition 1.15) and a connected orientable topological manifold of dimension n .*

(a) *There is a constant C , depending only on n and the constants in regularity and local linear contractability conditions, such that the following is true. For each $x_0, x_1 \in M$ there is a family of curves in M with parameter space $\mathbf{S}^n$ (see Definition 1.2) such that all the curves are contained in $B = B_M(x_0, C d(x_0, x_1))$, they all start at x_0 and end at x_1 , and they are controlled in measure by $\nu = h \cdot \mu|_B$, where the density h is defined by*

$$(B.8) \quad h(y) = C' (d(y, x_0)^{-n+1} + d(y, x_1)^{-n+1}).$$

Note that we are implicitly using Lebesgue measure on our parameter space $\mathbf{S}^n$.

(b) *There is a constant C' , depending only on n and the constants in regularity and local linear contractability conditions, such that*

$$(B.9) \quad \begin{aligned} &\text{any pair of points } x_0, x_1 \in M \text{ can be connected} \\ &\text{by a curve in } M \text{ of length } \leq C' d(x_0, x_1). \end{aligned}$$

To see this we use Lemmas B.4 and B.5 to conclude that M is doubling and that n -dimensional Hausdorff measure on M is comparable in size to the measure μ given by the regularity assumption. This permits us to derive part (a) of the theorem from Theorem 1.11 and Lemma 1.16. That is, Lemma 1.16 permits us to convert most of the hypotheses of Theorem B.6 into a form that can be used by Theorem 1.11. One still needs to check the assumption $\nu(U_1) < \infty$ stated just after (1.12), but this follows from the assumption of regularity, as we shall see in Lemma C.7 in the next appendix. Part (b) of the theorem follows from (a) and Lemma C.7 below.

Note that the local linear contractability condition and the connectedness of M automatically imply that any pair of points x_0, x_1 in M can be connected by a curve whose diameter is at most $C d(x_0, x_1)$. Controlling the length is another matter. Snowflake curves, for instance, satisfy the local linear contractability condition but do not contain any nontrivial rectifiable curves.

The next theorem provides Sobolev and Poincaré inequalities under the assumptions (B.7). We shall formulate these results in terms of the notion of generalized gradient given in Definition 1.18.

Theorem B.10. *Suppose that $(M, d(x, y), \mu)$, n are as in (B.7). Then functions on M can be controlled by their gradients in the following ways. Suppose that u and g are Borel*

measurable functions on an open set Ω in M , with u real-valued and g taking values in $[0, \infty]$, and suppose that g is a generalized gradient of u .

(a) Assume that $\Omega = M$, and that either M is unbounded and $u(z) \rightarrow 0$ as $z \rightarrow \infty$ in M , or that M is bounded and $\int_M u(z) d\mu(z) = 0$. Then

$$(B.11) \quad |u(x)| \leq C \int_M \frac{1}{d(x, y)^{n-1}} g(y) d\mu(y) \quad \text{for all } x \in M.$$

In this case we also have the Sobolev inequality $\|u\|_q \leq C(p) \|g\|_p$ when $1 \leq p < n$, $\frac{1}{q} = \frac{1}{p} - \frac{1}{n}$. (These L^p norms are taken with respect to the measure μ .)

(b) There is a constant $L_1 > 1$ so that if $B(w, L_1 r) \subseteq \Omega$ then we have the Poincaré inequality

$$(B.12) \quad r^{-2n} \int_B \int_B |u(x) - u(y)| d\mu(x) d\mu(y) \leq L_1 r^{-n} \int_{B(w, L_1 r)} r g(z) d\mu(z),$$

where $B = B(w, r)$.

(c) There is a constant L_2 so that if $x, y \in \Omega$ and $B(x, L_2 |x - y|) \subseteq \Omega$ then

$$(B.13) \quad |u(x) - u(y)| \leq L_2 \int_{B(x, L_2 |x - y|)} \left\{ \frac{1}{d(x, z)^{n-1}} + \frac{1}{d(y, z)^{n-1}} \right\} g(z) d\mu(z).$$

In this case we also have

$$(B.14) \quad |u(x) - u(y)| \leq C(p) L_2 |x - y|^{1 - \frac{n}{p}} \|g\|_p$$

when $n < p \leq \infty$.

These constants depend only on n and the constants in the local linear contractability and regularity conditions (Definitions 1.15 and B.1), and also on p where appropriate.

This theorem says that a lot of the standard results in analysis on Euclidean spaces extend to metric spaces which satisfy (B.7). It is pretty pointless when the dimension $n = 1$ (in which case M would be bilipschitz equivalent to a line or a circle, as one can check using “arclength” parameterizations).

Note that a standard reason for Sobolev inequalities not to hold on a space is that the space has a big bubble with a small neck. This means that one can have a function whose derivative is concentrated on the small neck even though the function is large (and constant) on the big bubble. In our case it is the local linear contractability condition which prevents this from happening.

Part (b) of the theorem is not an automatic consequence of (a), because it holds for all balls. We shall derive both (a) and (b) from (c).

In Theorem B.10 we could also give other standard variants of the Sobolev and Poincaré inequalities, but we shall not bother with that. See [HaK] for the more interesting issue of knowing when there are inequalities like (B.12) in which one can use the same ball on both sides of the inequality.

Before we explain why Theorem B.10 is true let us give a reformulation of it in which the role of the hypotheses is made more precise.

Theorem B.15. *Suppose that $(M, d(x, y), \mu)$ is an n -dimensional regular metric space, and assume that part (a) of the conclusions of Theorem B.6 holds. Then $(M, d(x, y), \mu)$ satisfies the conclusions of Theorem B.10, with constants that depend only on the constants in our assumptions and on the exponent p where appropriate.*

Thus the topological parts of the hypotheses (B.7) of Theorem B.10 are needed only for getting the families of curves in part (a) of Theorem B.6. Once we have these families of curves we do not need the topology any more, only the regularity.

Of course Theorem B.10 will follow once we prove Theorem B.15, because of Theorem B.6.

So why is Theorem B.15 true? It is not difficult to derive (B.13) from the existence of curve families as in part (a) of Theorem B.6, because of the definition of the generalized gradient. The argument is the same as the one used to derive Theorem 1.22 from Theorem 1.11 and Definition 1.2. Once we have (B.13) the rest of the conclusions of Theorem B.10 can be derived from it, using only the assumption that $(M, d(x, y), \mu)$ be an n -dimensional regular metric space. This assertion follows from standard arguments, and we include it in Proposition C.6 below for the sake of completeness. This proves Theorem B.15, modulo Proposition C.6.

For the record, let us state another variant of Theorem B.10.

Theorem B.16. *Suppose that $(M, d(x, y), \mu)$ is an n -dimensional regular metric space, and fix a point $\sigma \in \mathbf{S}^n$. Assume that there is a constant $k > 0$ so that for each pair of points x_0, x_1 we can find a mapping $F : M \times (0, T) \rightarrow \mathbf{S}^n$, $T = d(x_0, x_1)$, with the following properties:*

(B.17) *F is Lipschitz with norm $\leq k \epsilon^{-1}$ on $M \times (\epsilon, T - \epsilon)$ when $0 < \epsilon < T$;*

(B.18) *$F(z, t) = \sigma$ when either $d(z, x_0) > kt$ or $d(z, x_1) > k(T - t)$;*

(B.19) *each $f_t = F(\cdot, t) : M \rightarrow \mathbf{S}^n$, $0 < t < T$, cannot be homotoped to a constant through maps that all equal σ on the complement of some compact subset of M .*

Then $(M, d(x, y), \mu)$ satisfies the conclusions of Theorem B.10, with constants that depend only on k , n , and the regularity constant for M , and also on the exponent p where appropriate.

This follows from Theorems B.15 and 12.5. That is, Theorem 12.5 ensures the existence of families of curves as in part (a) of Theorem B.6 under the hypotheses of Theorem B.16, and then Theorem B.15 permits us to conclude. We are also using Lemma C.7 here to know that the assumption $\nu(U_1) < \infty$ in Theorem 12.5 (just below (12.9)) is valid.

A nice point about Theorem B.16 is that Theorem 18.3 provides another criterion for its assumptions to hold (that is, a criterion very different from (B.7)).

In summary, Theorems B.10, B.15, and B.16 provide criteria for Sobolev and Poincaré inequalities to hold on a regular metric space.

Remark B.20. Let us briefly sketch a different approach to Sobolev inequalities on metric spaces which satisfy (B.7) which is due to Gromov and which is based on [Gr3]. We shall not try to obtain Sobolev inequalities directly, we shall look for isoperimetric inequalities instead. That is, we want to show that if we are given a bounded region in M , then the Hausdorff measure of its boundary will control either the measure of the original region or the measure of the complement of the region inside M . (We need to allow this alternative only when M is compact, otherwise the Hausdorff measure of the boundary should control the measure of the region.)

Let us first deal with a minor topological issue. For the isoperimetric inequality it is better to think of a region and its boundary geometrically, as sets. For the argument that follows we want to think homologically, in terms of chains and cycles. Thus we want to give ourselves a (bounded) region in M and think of its boundary initially as a set, but then as a cycle of dimension $n - 1$. (The reader is invited to make whatever a priori smoothness or simplicial assumptions desired, to keep to the main ideas.) We then want to try to fill that cycle by an n -dimensional chain in M which is not too large. The small technical point is that for our purposes it suffices to control the measure of the support of this chain rather than the mass of the chain which counts multiplicities. For indeed the support of this chain will contain our original region if M is unbounded, and if M is bounded then the support of the chain will contain either the original region or its complement in M . (The sum of this n -chain and the one determined by our original region has no boundary, which means that either it vanishes identically, or M is compact and the sum is a nonzero multiple of the fundamental class of M , in which case the complement of our original region is contained in the support of the new n -chain.)

The point of departure in Gromov's approach is the well-known isometric embedding of M into the Banach space of bounded functions on M . A wonderful fact in [Gr3] is that one can always fill a cycle in a Banach space with control on the mass of the filling, i.e., with a bound of the type that one would expect from the isoperimetric inequality. If one is working inside a finite-dimensional Euclidean space, then the existence of such a controlled filling is an old result of Federer and Fleming. However their result gives an estimate that deteriorates when the ambient dimension goes to infinity. Michael and Simon showed that in fact one can do the filling with bounds that do not depend on the ambient dimension, and then Gromov proves in [Gr3] that one can do the filling in Banach spaces (and not just Hilbert spaces).

Thus one embeds M isometrically into $L^\infty(M)$, one takes a region in M , one considers the $(n - 1)$ -dimensional cycle determined by its boundary, and one can fill this cycle inside $L^\infty(M)$ in the manner just described. That is, one can fill it inside $L^\infty(M)$, with a good bound on the mass of the filling. We have to try to push the filling back inside M without losing control on the mass.

One of the main points is to use a "minimal filling", as in Section 4.3.C on p.36ff of [Gr3]. The words "minimal filling" have to be taken with some care, as in the remark on p.38 of [Gr3]. Minimal fillings have the advantage that they do not have long fingers stupidly sticking out, because of a lower bound on the mass of the intersection of the filling with a ball centered on the filling and which does not reach the boundary, as in Theorem 4.3.C" on p.37 of [Gr3].

The next step is to cover the filling by balls centered on the filling and whose radius is comparable to the distance to the boundary. More precisely, one should be a little careful and use a covering lemma to make a covering with the property that the sum of the n^{th} powers of the radii of the balls is bounded by a constant times the mass of the filling. (For instance, one can choose the covering so that the balls become disjoint after shrinking them by a bounded amount.) One can then push this filling into M , in such a way that the resulting n -chain in M is supported in a union of balls whose radii admit a similar estimate. This pushing can be accomplished using the local linear contractability property of M . The *linear* contractability is very important here; when we go to push forward part of the chain that lives in one of these balls, we need to know that the image lies in a ball of comparable size. We do not have any control on these pushings-forward, in terms of a Lipschitz constant or anything like that, but we don't need any control of that nature, we only need to control the measure of the support of the total push-forward. This is provided by the Ahlfors regularity of our metric space. Specifically, the measure of the support of the n -chain that we have constructed in M is bounded by a constant times the sum of the radii of the balls to the n^{th} power, which is controlled in turn by the mass of the filling in the ambient Banach space.

This rough sketch lacks many details, but these are the main ingredients.

One should compare the logic of this argument to the methods of this paper. In some ways Gromov's approach is clearly superior, it does a much better job of getting to the geometric heart of Sobolev inequalities. If M happens to be a subset of some finite-dimensional Euclidean space the filling issues become more simple and more classical, while the methods of this paper do not become especially simpler. In any case the filling issues are completely separate from the assumptions on M , which is very nice. The assumptions on M are employed in a much less focussed manner in this paper, it is harder to understand exactly what is needed and why. On the other hand the methods of this paper do provide other types of information, and the circumstances under which they can be used are not the same as in Gromov's approach. It just happens that in this special case of many assumptions the two methods provide similar information.

Remark B.21. The assumption of regularity in this appendix is quite strong in comparison to the assumptions that we made in Section 1. If one is ready to accept it then one should take a closer look at our earlier arguments to see how they can be strengthened or simplified. In this regard there is a particular point which stands out. In Theorem 6.5 we went to some trouble to show that the spaces that we were considering not only have topological dimension $\leq n$, but they enjoy this property with a certain uniform bound. This uniform bound used a compactness argument and the bounds on the contractability function. If we are willing to make an assumption like regularity, then we can get this kind of topological dimension condition with a bound more directly. We could use Proposition 6.4, for example. See Remark 6.10 for further discussion of how the assumption of a bound on the contractability function might be weakened in these circumstances.

For a regular metric space there is a more direct approach to topological dimension conditions with bounds than Proposition 6.4. The main ingredient is a theorem of Assouad, to the effect that if $(M, d(x, y))$ is a metric space which is doubling, then for each $s \in (0, 1)$ there is a bilipschitz embedding of $(M, d(x, y)^s)$ into some $\mathbf{R}^N$. (See Proposition 2.6 on

p.436 of [A3], and also [A1] and [A2].) It would be nicer if one could take $s = 1$ here, but this is unfortunately not possible. One can take s to be arbitrarily close to 1, however. If $(M, d(x, y))$ is a regular metric space of dimension n , then $(M, d(x, y)^s)$ is a regular metric space of dimension n/s , and we can make this close to n by taking s close to 1. Once we have $(M, d(x, y)^s)$ embedded in some $\mathbf{R}^N$ we can build mappings from it into polyhedra with bounds using well-known geometric arguments (retracting onto skeleta, using the Ahlfors regularity of the set to be able to avoid the singularities of the retractions).

For the record:

Definition B.22. Let $(M, d(x, y))$ and $(M', d'(v, w))$ be metric spaces. A mapping $f : M \rightarrow M'$ is *bilipschitz* if there is a constant C such that

$$(B.23) \quad C^{-1} d(x, y) \leq d'(f(x), f(y)) \leq C d(x, y)$$

for all $x, y \in M$. We sometimes say that f is C -bilipschitz to make the constant explicit. Note that we do not require that f be surjective. If there is a bilipschitz bijection between two metric spaces, then we say that these metric spaces are *bilipschitz equivalent*.

Let us now think more carefully about what the conditions (B.7) mean, and compare them with more familiar conditions concerning the existence of good parameterizations of a metric space $(M, d(x, y))$.

If M is bilipschitz equivalent to $\mathbf{R}^n$ then it satisfies (B.7). In this case the conclusions of Theorems B.6 and B.10 are not very interesting, because they can all be reduced to familiar facts about $\mathbf{R}^n$ using the bilipschitz parameterization.

Suppose now that $(M, d(x, y), \mu)$ is regular of dimension n and that it is quasisymmetrically equivalent to $\mathbf{R}^n$. This means that there is a mapping f from $\mathbf{R}^n$ onto M and a homeomorphism $\eta : [0, \infty) \rightarrow [0, \infty)$ such that

$$(B.24) \quad |x - a| \leq t |x - b| \text{ implies } d(f(x), f(a)) \leq \eta(t) d(f(x), f(b))$$

for all $t > 0$ and $x, a, b \in \mathbf{R}^n$. In other words, f should distort *relative* distances by only a bounded amount, i.e., if a is much closer to x than b is, then the analogous statement is true for $f(a), f(x)$, and $f(b)$. A priori this does not prevent $|f(x) - f(a)|$ from being wildly different from $|x - a|$, but it turns out that the regularity of M and an argument of Gehring [Ge1] imply that the pull-back of the measure μ on M via f is absolutely continuous with respect to Lebesgue measure on $\mathbf{R}^n$, and that the density satisfies some scale-invariant L^p bounds with $p > 1$. (See [DS1] and [Se2].) In this case the conclusions of Theorems B.6 and B.10 are not so trivial and they are equivalent to some results in [DS1]. (See also [Se6].)

It is not very easy to tell when a metric space is bilipschitz or quasisymmetrically equivalent to some $\mathbf{R}^n$ though.

There are also examples of metric spaces which satisfy (B.7) but which do not admit such good parameterizations, even locally. For instance, there are finite polyhedra which are topological manifolds but which do not admit bilipschitz local coordinates. These examples arise from the work of Edwards and Cannon ([E], [C1], [C2]) on double suspensions

of homology spheres. See also [SS]. These polyhedra satisfy (B.7) as soon as they are connected and orientable. Although these polyhedra are very strange in terms of local coordinates, it is easy to verify directly that they satisfy the conclusions of Theorems B.6 and B.10 (without using the complicated arguments given here). For this we do not need the polyhedra to be topological manifolds; connected finite polyhedra are fine in general so long as one avoids simple pathologies, like a pair of planes which meet in only a point.

As another example, take M to be $\mathbf{R}^3$ equipped with the following metric. Take a self-similar Antoine's necklace A in $\mathbf{R}^3$, as in [Mo], and deform the (Riemannian) Euclidean metric conformally by the factor $\text{dist}(x, A)^5$ (say). One gets in this way a Riemannian metric which degenerates on A , but it turns out that the associated distance function does not degenerate, because of the disconnectedness of A . One can show that $\mathbf{R}^3$ equipped with this new distance function satisfies (B.7), and that this metric space is not bilipschitz equivalent to $\mathbf{R}^3$. There are also no local bilipschitz coordinates for this space near A . See [Se4] for details.

There is another example in [Se4] of a conformal deformation of a Euclidean space with the property that the resulting metric space satisfies (B.7) but is not bilipschitz equivalent to any subset of any Euclidean space (of finite dimension).

Another interesting class of examples can be produced using some famous examples of "decompositions" in topology (as in Section 9 of [Da]). These examples are obtained from procedures which have a kind of topological self-similarity, but one can make them geometrically self-similar in a natural way. The simplest example of this type is based on the construction of the Whitehead continuum (see Example 7 on p.68 of [Da], or Section 4 on p.81 of [K]). Bing's dogbone space (Example 4 on p.64 of [Da]) and Bing doubling (Example 1 on p.62 of [Da]) give rise to other interesting examples. Using these constructions one can build metric spaces (subsets of Euclidean spaces even) which satisfy (B.7) but which do not admit quasisymmetric local coordinates, and which have other interesting properties as well. See [Se5].

It is natural to wonder whether the nonsmoothable compact topological 4-manifolds constructed by Freedman (see [FQ]), like the simply connected E_8 manifold, admit metrics which satisfy the properties in (B.7). In other words, does there exist a topological obstruction which prevents (B.7) from being realized? In dimensions less than 4 every topological manifold is smoothable, while in dimensions larger than 4 every topological manifold admits a (unique) Lipschitz structure, by Sullivan's theorem [Su]. This Lipschitz structure gives rise to a metric which satisfies (B.7) (as long as the manifold is compact, connected, and orientable). Sullivan's theorem fails in dimension 4, because [DoS] implies that many of the topological manifolds constructed by Freedman (like the simply connected E_8) do not admit Lipschitz structures. Note that a connected topological 4-manifold is smoothable away from a single point, by a theorem of Quinn (see §8.2 of [FQ]), but not much is known, by way of quantitative estimates, about what will happen at the singularity.

Let us contemplate for a moment the relationship between analysis and topology, between the validity of basic analytic statements like Sobolev inequalities and the existence of homeomorphic coordinates which satisfy good estimates. If one has a manifold with bilipschitz coordinates, for instance, then one can transfer many analytical results (like Sobolev and Poincaré inequalities) from Euclidean space to the manifold. The point of

Theorem B.10 is that one can also get some analytical results without good local coordinates. In most of the examples above the obstruction to the existence of good local coordinates is somehow related to π_1 . I do not mean the fundamental group of the whole space, but rather π_1 of pieces of the space. For example, if one takes the double suspension of a non-simply connected 3-sphere, as in the work of Edwards and Cannon, then one gets a finite polyhedron which is homeomorphic to a sphere, but there is a polygonal curve in the polyhedron whose complement is not simply connected, and that prevents the double suspension from being bilipschitz equivalent to a sphere. Part of the point of Theorem 1.16 is that there is a lot of analysis which is not sensitive to problems with π_1 of this type. Gauge theory in dimension 4 has the important feature that it is more sensitive to π_1 phenomena.

It is very important that we require that the Hausdorff and topological dimensions be the same in (B.7). Except for this requirement snowflakes could satisfy (B.7), but they would not satisfy the conclusions of Theorem B.10. By contrast the Heisenberg group with its Carnot metric has the very interesting properties that it satisfies (B.7) except for the equality of the Hausdorff and topological dimensions, and that it also satisfies (B.9) and suitable versions of the Poincaré and Sobolev inequalities and the rest. (See [Fo], [FS1], [FS2], [Je], [P], [St2] for information about analysis on the Heisenberg group.)

C. Regular metric spaces.

Recall that the notion of a regular metric space was defined in Definition B.1. In this appendix we prove some facts about regular metric spaces which were needed in Appendix B. The main result is that the conclusions of Theorem B.10 hold for any regular metric space as soon as (B.13) is established (Proposition C.6).

The material in this appendix is quite standard, but we include it for the sake of completeness and clarity.

Lemma C.1. *Suppose that $(M, d(x, y), \mu)$ is a regular metric space of dimension n . Then there is a constant C so that for any ball B in M with radius $R > 0$ and any $r \in (0, R)$ there is a covering of B by a family of at most $C(R/r)^n$ balls in M of radius r . In particular it is doubling (Definition 1.14), and closed and bounded subsets of M are compact.*

This is the same as Lemma B.4, whose proof was postponed before.

Let M be as in the lemma, let B be a ball in M of radius R , and let $r \in (0, R)$ be given. We may as well assume that $R \leq \text{diam } M$. Let A be any subset of B such that $d(x, y) \geq r$ whenever x and y are distinct points in A . Then the balls $B(x, r/2)$, $x \in A$, are pairwise disjoint, and so

$$(C.2) \quad \sum_{x \in A} \mu(B(x, r/2)) \leq \mu(2B) \leq C R^n.$$

On the other hand $\mu(B(x, r/2)) \geq C^{-1} r^n$ for each x , and so we conclude that A has at most $C(R/r)^n$ elements. If we choose A so that it has a maximal number of elements, then every point in B is contained in the union of the balls $B(x, r)$, $x \in A$, by maximality. This implies the covering property asserted in the lemma, and the doubling condition follows.

Since M is complete (see Definition B.1) and bounded sets can be covered by finite families of balls of arbitrarily small radius, we conclude that closed and bounded subsets of M are (complete and totally bounded and hence) compact. This proves the lemma.

Lemma C.3. *If $(M, d(x, y), \mu)$ is a regular metric space of dimension n , then there is a constant $C > 0$ so that $C^{-1}\mu(E) \leq H^n(E) \leq C\mu(E)$ for all Borel sets $E \subseteq M$.*

This is the same as Lemma B.5, whose proof was postponed before. Remember that H^n denotes n -dimensional Hausdorff measure, defined in (2.14).

Let M and μ be as in the lemma. The estimate $\mu \leq C H^n$ follows immediately from the definition of H^n and the fact that $\mu(A) \leq C(\text{diam } A)^n$ for any measurable (in this case closed) set A in M . The reverse inequality is a bit more involved. Notice first that $H^n(B) \leq CR^n$ whenever B is a ball with radius R in M , because Lemma C.1 implies that $H_\delta^n(B) \leq CR^n$ for all $\delta > 0$. Next let us check that $H^n(U) \leq C\mu(U)$ when U is an open subset of M . We may as well assume that $\mu(U) < \infty$, so that U does not contain balls with unbounded radii. Since U is open we can realize it as a union of open balls. Using a covering argument as on p.9 of [St1] (which works for any metric space) we can find a sequence of disjoint balls $\{B_i\}$ contained in U such that either $U \subseteq \bigcup_i 5B_i$ or the radii of these balls do not tend to 0. The latter alternative is prevented by the assumption that $\mu(U) < \infty$ and the regularity of M . Thus we get

$$H^n(U) \leq \sum_i H^n(5B_i) \leq \sum_i C\mu(5B_i) \leq C' \sum_i \mu(B_i) \leq C' \mu(U).$$

This gives the desired inequality when U is an open set. For the general case we use the fact that μ must be outer regular in the classical sense in measure theory, i.e.,

$$(C.4) \quad \mu(E) = \inf\{\mu(U) : E \subseteq U \text{ and } U \text{ is open}\}$$

for all Borel sets E . This fact is well known, and it follows from Theorem 2.18 in [R]. (This theorem requires that M be locally compact and that every open set in M can be realized as a countable union of compact subsets. In our case both of these conditions hold, because of Lemma C.1.) This proves Lemma C.3.

Many familiar results from real analysis on Euclidean spaces are also true on regular metric spaces, or even on the “spaces of homogeneous type” studied in [CW1], [CW2]. The existence of Lebesgue points is an example of this principle.

Lemma C.5. *If $(M, d(x, y), \mu)$ is a regular metric space of dimension n , and f is a locally integrable function on M (with respect to μ), then*

$$\lim_{r \rightarrow 0} r^{-n} \int_{B(r, x)} |f(y) - f(x)| d\mu(y) = 0$$

for μ -almost all $x \in M$.

As explained in [CW2], the point here is that our modest geometric assumptions are sufficient to ensure that the usual proof on Euclidean spaces goes through. For instance,

the covering lemma on p.9 of [St1] goes through in this context because it uses only the fact that $\mathbf{R}^n$ is a metric space with a measure such that the mass of the ball is a power of the radius. (In fact it is sufficient to know that the mass of $2B$ is at most a constant times the mass of B for every ball B .) Once one has this covering lemma one gets the estimates on the maximal function as in Theorem 1 on p.5 of [St1] in exactly the same way as on Euclidean space. The derivation of Corollary 1 on p.5 of [St1] (the existence almost everywhere of the pointwise averages) and the argument to get from there to the existence of Lebesgue points on p.11 of [St1] also use no special structure of Euclidean space. This proves Lemma C.5.

Proposition C.6. *Let $(M, d(x, y), \mu)$ be a regular metric space of dimension n . If the estimate (B.13) holds under the conditions of part (c) of Theorem B.10, then the rest of the conclusions of Theorem B.10 also hold.*

To prove this we again mimic classical arguments on $\mathbf{R}^n$.

Lemma C.7. *Let $(M, d(x, y), \mu)$ be a regular metric space of dimension n , and let $s < n$ be given. Then for each $x \in M$ and $R > 0$ we have that*

$$(C.8) \quad \int_{B(x, R)} \frac{1}{d(x, y)^s} d\mu(y) \leq C(s) R^{n-s}.$$

To see this we note that

$$(C.9) \quad \int_{B(x, 2r) \setminus B(x, r)} \frac{1}{d(x, y)^s} d\mu(y) \leq C(s) r^{n-s}$$

for all $r > 0$, by our regularity condition (B.2). It is easy to derive (C.8) from (C.9) by summing a geometric series. This proves Lemma C.7.

Lemma C.10. *Let $(M, d(x, y), \mu)$ be a regular metric space of dimension n . Suppose that the estimate (B.13) in part (c) of Theorem B.10 has been established. Then the Poincaré inequality (B.12) and the bound (B.14) follow automatically. The estimate (B.11) is also valid for Borel measurable functions u, g such that (u is real-valued, g takes values in $[0, \infty]$, g is the generalized gradient of u , and) one of the following conditions holds: (i) M is bounded and $\int_M u(x) d\mu(x) = 0$; (ii) M is unbounded and $u(x) \rightarrow 0$ as $x \rightarrow \infty$ in M ; (iii) $\mu(\{x \in M : u(x) \neq 0\}) < \frac{1}{2} \mu(M)$. (When M is unbounded this last assumption means that $\mu(\{x \in M : u(x) \neq 0\}) < \infty$.)*

In order to obtain (B.12) from (B.13) it is enough to use Fubini's theorem and (C.8). It is not difficult to derive (B.14) from (B.13), Hölder's inequality, and (C.8). Now let us check (B.11) in the various cases. In case (i) we average (B.13) over y and use the assumption that u has integral zero to get (B.11). (We also use Lemma C.7 here to control the integral in y of $d(y, z)^{-n+1}$.) When (ii) holds we average (B.13) over $y \in B(x, 2R) \setminus B(x, R)$ and then send $R \rightarrow \infty$ to obtain (B.11). The average of the right side of (B.13) behaves properly for this computation, because the average of $d(y, z)^{-n+1}$ over $y \in B(x, 2R) \setminus B(x, R)$ is $\leq C R^{-n+1} \leq C d(x, z)^{-n+1}$ when $z \in B(x, 2L_2 R)$. The third case is handled by the

same kind of averaging arguments, except that one takes care to average only over y 's such that $u(y) = 0$. This proves Lemma C.10.

To finish the proof of Proposition C.6 it remains to establish the Sobolev inequalities in Theorem B.10 (a).

Lemma C.11. *Let $(M, d(x, y), \mu)$ be a regular metric space of dimension n , and define the linear operator I_1 on functions on M by*

$$(C.12) \quad I_1(g)(x) = \int_M \frac{1}{d(x, y)^{n-1}} g(y) d\mu(y).$$

Suppose that $1 \leq p < n$ and q is defined by $\frac{1}{q} = \frac{1}{p} - \frac{1}{n}$. If $g \in L^p(\mu)$, then the integral in (C.12) converges absolutely for almost all $x \in M$. When $p > 1$ I_1 defines a bounded linear operator from $L^p(\mu)$ into $L^q(\mu)$. When $p = 1$ I_1 is weak-type $(1, q)$, which means that there is a constant $C > 0$ so that

$$(C.13) \quad \mu(\{x \in M : |I_1(g)(x)| > \lambda\}) \leq C \lambda^{-q} \|g\|_1^q$$

for all $g \in L^1(\mu)$. The constants in these bounds depend only on the constants coming from the regularity of the metric space M (as in Definition B.1).

If $M = \mathbf{R}^n$ then this is a classical result which is the same as Theorem 1 on p.119 of [St1]. Exactly the same proof as given in [St1] works in this more abstract setting, modulo cosmetic changes which stem from the fact that I_1 is not a convolution operator in the present situation. In outline the argument is as follows. One shows first that I_1 is weak-type (p, q) when $1 \leq p < n$. This is sufficient, because the Marcinkiewicz interpolation theorem then implies the boundedness of the operator when $p > 1$. To get the weak-type result one splits I_1 into the near and far-away parts, which correspond to the portions of the kernel which live near or far from the diagonal. One uses L^1 estimates for the near part and L^∞ estimates for the far-away part. The precise computations are as in [St1], but the loss of the convolution structure here means only that one has to use the Minkowski integral inequalities instead of Hausdorff-Young. We leave the details to the reader.

Lemma C.11 implies that the Sobolev inequality promised in part (a) of Theorem B.10 is an automatic consequence of (B.11) when $p > 1$. However, it is well known and easy to show that I_1 does not map $L^p(\mu)$ into $L^q(\mu)$ when $p = 1$. To finish the proof of Proposition C.6 we need the following.

Proposition C.14. *The $p = 1$ case of the Sobolev inequality asserted in Theorem B.10 (a) holds under the same assumptions as in Proposition C.6.*

The main point of the proof of this proposition is that the true relationship between a function and its generalized gradient is much more local than in (B.11), and this extra degree of localization permits us to win something in the estimates. More precisely, we shall break our functions u and g into pieces, in such a way that each piece of u is approximately a multiple of a characteristic function, and then we shall apply (C.13) to each of these various pieces separately and combine the results. The argument will be very similar to the well-known derivation of the $p = 1$ Sobolev inequality from the isoperimetric inequality, but for technical reasons it will be formulated a bit differently.

Let u, g be given, as in part (b) of Theorem B.10. Thus u and g are Borel measurable functions on M , g is a generalized gradient of u , and either

$$(C.15) \quad \text{diam } M = \infty \text{ and } u(x) \rightarrow 0 \text{ as } x \rightarrow \infty \text{ in } M, \text{ or}$$

$$(C.16) \quad \text{diam } M < \infty \text{ and } \int_M u(x) d\mu(x) = 0.$$

Because of Lemma C.10 we know the Poincaré inequality (B.12) in this case, and so

$$(C.17) \quad \int_M |u(x)| d\mu(x) \leq C \int_M \text{diam } M g(z) d\mu(z)$$

when M is bounded. Thus we may replace (C.16) with the assumption that (C.17) hold when M is bounded. We can also reduce to the case where u is nonnegative. Indeed, if we replace u by $|u|$, then g is automatically a generalized gradient of $|u|$, and (C.15) and (C.17) remain undisturbed by this substitution. Hence we may assume that u is nonnegative.

The proof of Proposition C.14 that follows will be clearer in the special case where u is continuous. Let us make this auxiliary assumption for the time being and then get rid of it afterwards with an approximation argument.

For each integer j let u_j be the function defined by

$$(C.18) \quad \begin{aligned} u_j(x) &= 0 && \text{when } u(x) \leq 2^j \\ &= u(x) - 2^j && \text{when } 2^j < u(x) < 2^{j+1} \\ &= 2^j && \text{when } 2^{j+1} \leq u(x). \end{aligned}$$

Set $F_j = \{x \in M : 2^j \leq u(x) \leq 2^{j+1}\}$, and (for technical reasons) let V be any open subset of M which contains F_j . We shall eventually choose V so that it shrinks down to F_j , in the sense that $\mu(V \setminus F_j)$ goes to 0. Define g_j by $g_j(x) = g(x)$ on V , $g_j(x) = 0$ on $M \setminus V$.

Lemma C.19. *g_j is a generalized gradient of u_j .*

We can derive this from Definition 1.18. Let γ be a curve in M , as in Definition 1.18, and let us check that the analogue of (1.19) holds for u_j and g_j . If γ is contained in V , then the desired inequality for u_j and g_j is equivalent to the original version for u and g , and so there is nothing to do. If γ is contained in $M \setminus F_j$, then u_j is constant along γ (since u is continuous), and the analogue of (1.19) in this case is vacuous. Now suppose that γ is any curve in M . We can decompose γ into finitely many subarcs, each of which is contained in either V or $M \setminus F_j$. This statement reduces to the fact that if an interval $[a, b]$ in $\mathbf{R}$ is covered by two open sets, then it can be broken up into finitely many closed intervals, each of which is contained in one of the covering open sets. (Note that $M \setminus F_j$ is open, because u is continuous.) Once we have this decomposition, the analogue of (1.19) in the general case follows from the preceding special cases. This proves Lemma C.19.

The plan now is to get a good estimate for u_j using (B.11) and the $p = 1$ part of Lemma C.11. From now on we let A denote the average of u on M when M is bounded, and we set $A = 0$ when M is unbounded. Notice that

$$(C.20) \quad \mu(\{x \in M : u(x) > 2A\}) < \frac{1}{2} \mu(M)$$

when M is bounded, as in Telebychev's inequality. Hence

$$(C.21) \quad \mu(\{x \in M : u_j(x) > 0\}) < \frac{1}{2} \mu(M)$$

when M is bounded and $2^j \geq 2A$.

Let us check that

$$(C.22) \quad u_j(x) \leq C \int_M \frac{1}{d(x, y)^{n-1}} g_j(y) d\mu(y) \leq C I_1(g_j)(x)$$

for all $x \in M$ when $2^j > 2A$. This application of (B.11) holds because of Lemma C.10 applied to u_j and g_j . Specifically, we are using cases (ii) and (iii) of the last part of Lemma C.10, and we are using the fact that either (C.15) or (C.21) holds. Here C depends only on the constants that arise in the hypotheses of Proposition C.6.

Let q be associated to $p = 1$ as in Theorem B.10 (a) and Lemma C.11. Using (C.13) we obtain

$$(C.23) \quad \mu(\{x \in M : u(x) > 2^{j+1}\}) = \mu(\{x \in M : u_j(x) > 2^j\}) \leq C 2^{-jq} \left(\int_M g_j(y) d\mu(y) \right)^q$$

when $2^j > 2A$. This implies that

$$(C.24) \quad \mu(\{x \in M : u(x) > 2^{j+1}\}) \leq C 2^{-jq} \left\{ \int_{F_j} g(y) d\mu(y) \right\}^q$$

when $2^j > 2A$. Indeed, (C.24) would be identical to (C.23) if we integrated over V instead of F_j , but we can replace V with F_j since V was taken to be an arbitrary open superset of F_j . (We may as well assume here that g is integrable, since otherwise there is nothing to prove.)

We want to control the L^q norm of u using (C.24). Set $G = \{x \in M : u(x) \leq 8A\}$, so that

$$(C.25) \quad M \subseteq G \cup \left(\bigcup_{2^j > 2A} \{x \in M : 2^{j+1} < u(x) \leq 2^{j+2}\} \right).$$

Then

$$(C.26) \quad \begin{aligned} \int_M u(x)^q d\mu(x) &\leq (8A)^q \mu(G) + \sum_{2^j > 2A} 2^{(j+2)q} \mu(\{x \in M : 2^{j+1} < u(x) \leq 2^{j+2}\}) \\ &\leq (8A)^q \mu(G) + C \sum_{2^j > 2A} \left\{ \int_{F_j} g(y) d\mu(y) \right\}^q, \end{aligned}$$

by (C.24). We interpret the $(8A)^q \mu(G)$ term to be 0 in the case where M is unbounded (and $A = 0$, $\mu(G) = \infty$). The well known inequality

$$(C.27) \quad \left\{ \sum_i a_i \right\}^s \leq \sum_i a_i^s$$

for nonnegative numbers a_i and $0 \leq s \leq 1$ implies that

$$(C.28) \quad \|u\|_q \leq 8A \mu(G)^{\frac{1}{q}} + C \sum_j \int_{F_j} g(y) d\mu(y).$$

This is pretty much what we want. No point in M belongs to more than 2 of the F_j 's, and so the sum on the right is bounded by a constant times the L^1 norm of g . Therefore

$$(C.29) \quad \|u\|_q \leq 8A \mu(G)^{\frac{1}{q}} + C \int_M g(y) d\mu(y).$$

When M is unbounded this is exactly what we want, because the first term on the right is 0. When M is bounded the first term on the right is bounded by a constant times the L^1 norm of g , because the definition of q implies that $\mu(G)^{\frac{1}{q}} \leq \mu(M)^{\frac{n-1}{n}} \leq C \mu(M) (\text{diam } M)^{-1}$ (since M is regular), and because the average A of u on M is controlled by (C.17). Altogether we get that $\|u\|_q \leq C \|g\|_1$.

This proves Proposition C.14 when u is continuous. To complete the proof we need to use an approximation argument to get rid of this assumption. The basic idea is to approximate u by functions which have been smoothed out on small balls of a fixed size. In order to do this we need to have a convenient covering of M by balls of a fixed size, and we need to construct some partitions of unity.

Let $t > 0$ be given, with $t \leq \text{diam } M$. We shall be interested in approximating u by a nicer function at the scale of t .

Lemma C.30. *There is a family of balls $\{B_i\}$ in M , all with radius t , such that $M \subseteq \bigcup_i B_i$ and such that for each $k > 1$ and every ball B_i there are at most $C(k)$ balls B_j with the property that kB_j intersects kB_i . Here $C(k)$ depends also on the doubling constant of M (which is controlled by the regularity constant and the dimension, as in Lemma C.1), but not on t .*

Note that there will be only finitely many B_i 's when M is bounded, and infinitely many when M is unbounded.

Let us call a subset A of M *separated* if $d(x, y) \geq t$ whenever x and y are distinct elements of A . To prove the lemma we shall need a maximal separated subset A of M . In order to produce this maximal set A we start with a sequence $\{K_l\}$ of compact subsets of M which exhaust M . Let A_1 be a maximal separated subset of K_1 . Every separated subset of K_1 has a bounded number of elements (with the bound depending only on t and K_1), and so one can simply take A_1 to be a separated subset of K_1 with a maximal number of elements. If A_j has been chosen for some $j \geq 1$ take A_{j+1} to be a maximal subset of K_{j+1} which contains A_j and which is separated. In this way we can choose A_j for all j . The union A of the A_j 's is then a maximal separated subset of M .

Let $\{B_i\}$ be the family of balls centered on the elements of A and having radius t . The maximality of A ensures that $\{B_i\}$ covers M . We also have that the balls $\frac{1}{2}B_i$ are pairwise disjoint, since A is separated. To get the bounded overlap property in Lemma C.30 we observe that $B_j \subseteq (2k+1)B_i$ when kB_j intersects kB_i . The doubling condition on M ensures that $(2k+1)B_i$ can be covered by at most $C(k)$ balls of radius $t/10$, and

each such ball can contain the center of at most one B_j , since the balls $\frac{1}{2}B_j$ are pairwise disjoint. Thus there are at most $C(k)$ balls B_j such that kB_j intersects kB_i (for fixed i). This proves Lemma C.30.

From now on (in this section) we let the B_i 's be as in Lemma C.30.

Lemma C.31. (Partitions of unity.) *There is a family of functions $\{\phi_i\}$ on M such that $\sum_i \phi_i \equiv 1$ on M and, for each i , $0 \leq \phi_i \leq 1$, $\phi_i \equiv 0$ on $M \setminus 2B_i$, and ϕ_i is Lipschitz with norm $\leq Ct^{-1}$, where C depends only on the doubling constant of M .*

We first choose auxiliary functions $\tilde{\phi}_i$ which are nonnegative, vanish off $2B_i$, equal 1 on B_i , and have Lipschitz norm bounded by Ct^{-1} . Indeed, if x_i is the center of B_i , then we can choose $\tilde{\phi}_i$ to be of the form $\tilde{\phi}_i(x) = f(d(x, x_i))$ for a suitable function $f: \mathbf{R} \rightarrow \mathbf{R}$. (Exercise.) If we define the ϕ_i 's by

$$(C.32) \quad \phi_i = \tilde{\phi}_i \left\{ \sum_j \tilde{\phi}_j \right\}^{-1},$$

then the ϕ_i 's have the required properties. This is not hard to verify. (Note that there are only finitely many $\tilde{\phi}_j$'s which are nonzero at any given point in M , because of the bounded overlap condition in Lemma C.30. Thus there is no funny business in the sum.) This proves Lemma C.31.

We are now ready to make our approximation of u . As before we assume that u and g are Borel-measurable functions, with g a generalized gradient of u , and that either (C.15) or (C.17) holds. For each B_i let $u(B_i)$ denote the average of u over $2B_i$, and define u_t by

$$(C.33) \quad u_t = \sum_i \phi_i u(B_i),$$

where $\{\phi_i\}$ is as in Lemma C.31. This is our approximation to u which is supposed to be "smooth" at the scale of t . It is certainly Lipschitz on any compact set, since the ϕ_i 's are Lipschitz.

Let us estimate $|u - u_t|$. We have that

$$(C.34) \quad u - u_t = \sum_i \phi_i \cdot (u - u(B_i))$$

by definitions. Hence

$$(C.35) \quad \begin{aligned} \int_M |u - u_t| d\mu &\leq \sum_i \int_{2B_i} |u - u(B_i)| \\ &\leq \sum_i L_1 \int_{2L_1 B_i} t g d\mu, \end{aligned}$$

because we are assuming that we already know the Poincaré inequality (B.12). (See Lemma C.10.) The bounded overlap property in Lemma C.30 now implies that

$$(C.36) \quad \|u - u_t\|_1 \leq C(L_1) t \|g\|_1.$$

It is important that this constant $C(L_1)$ does not depend on t .

Next we should provide generalized gradients for the u_t 's. Define functions g_t by

$$(C.37) \quad g_t(x) = t^{-1} \max\{|u(B_i) - u(B_j)| : 2B_i \text{ and } 2B_j \text{ touch } B(x, t)\}.$$

This function is Borel measurable. Let us check that

$$(C.38) \quad |u_t(y) - u_t(z)| \leq C |y - z| g_t(x) \quad \text{when } y, z \in B(x, t).$$

By definitions we have that $u_t(y) - u_t(z) = \sum_i (\phi_i(y) - \phi_i(z)) u(B_i)$. Fix a j so that $x \in B_j$. Then

$$(C.39) \quad u_t(y) - u_t(z) = \sum_i (\phi_i(y) - \phi_i(z)) (u(B_i) - u(B_j)),$$

since $\sum_i \phi_i \equiv 1$. It is not hard to derive (C.38) from (C.39) using the bound on the Lipschitz norms of the ϕ_i 's provided by Lemma C.31 and the fact that only a bounded number of terms in the sum are nonzero (by Lemmas C.30 and C.31).

Clearly g_t is a generalized gradient for u_t (Definition 1.18), since (C.38) holds. It remains to show that

$$(C.40) \quad \|g_t\|_1 \leq C \|g\|_1,$$

where C does not depend on t . Let us first check that

$$(C.41) \quad \sup_{x \in B_l} g_t \leq C t^{-1} t^{-2n} \int_{6B_l} \int_{6B_l} |u(y) - u(z)| d\mu(y) d\mu(z)$$

for any B_l . Let $x \in B_l$ be given. If $2B_i$ and $2B_j$ touch $B(x, t)$, then $2B_i \cup 2B_j \subseteq 6B_l$, and $t^{-1}|u(B_i) - u(B_j)|$ is bounded by the right side of (C.41), since $u(B_i)$ is the average of u over $2B_i$. This implies (C.41), because of the definition (C.37) of g_t . Our Poincaré inequality (B.12) implies that

$$(C.42) \quad \sup_{x \in B_l} g_t \leq C t^{-n} \int_{6L_1 B_l} g d\mu.$$

Hence

$$(C.43) \quad \int_{B_l} g_t d\mu \leq C \int_{6L_1 B_l} g d\mu.$$

To get (C.40) we need only sum this in l and use the fact that the balls $6L_1 B_l$ have bounded overlap, by Lemma C.30.

Observe that the u_t 's satisfy (C.15) when M is unbounded, by their definition (C.33), and that they satisfy (C.17) when M is bounded (and $t \leq \text{diam } M$), because of (C.17) applied to u itself and (C.36). Because the u_t 's are also continuous we may apply our previous argument to them to get that $\|u_t\|_q \leq C \|g_t\|_1$ for all t , where q is the usual Sobolev

exponent, and where C does not depend on t . From (C.40) we obtain that $\|u_t\|_q \leq C\|g\|_1$ for all t , and (C.36) permits us to conclude that $\|u\|_q \leq C\|g\|_1$. (We are using a variant of Fatou's lemma here.) This completes the proof of Proposition C.14.

Now that Proposition C.14 is established the proof of Proposition C.6 is also complete.

References

- [A1] P. Assouad, *Espaces Métriques, Plongements, Facteurs*, Thèse de Doctorat (January, 1977), Université de Paris XI, 91405 Orsay Cedex, France.
- [A2] P. Assouad, *Étude d'une dimension métrique liée à la possibilité de plongement dans $\mathbf{R}^n$* , C. R. Acad. Sci. Paris **288** (1979), 731-734.
- [A3] P. Assouad, *Plongements Lipschitziens dans $\mathbf{R}^n$* , Bull. Soc. Math. France **111** (1983), 429-448.
- [BB] M. Barlow and R. Bass, *Coupling and Harnack inequalities for Sierpinski carpets*, Bull. Amer. Math. Soc. **29** (1993), 208-212.
- [BS] M. Bridson and G. Swarup, *On Hausdorff-Gromov convergence and a theorem of Paulin*, L'Enseignement Mathématique **40** (1994), 267-289.
- [BT] R. Bott and L. Tu, *Differential Forms in Algebraic Topology*, Graduate Texts in Mathematics **82**, Springer-Verlag, 1982.
- [BW] J. Block and S. Weinberger, *Large scale homology theories and geometry*, preprint.
- [C1] J. Cannon, *The characterization of topological manifolds of dimension $n \geq 5$* , Proceedings I.C.M. (Helsinki, 1978), 449-454.
- [C2] J. Cannon, *Shrinking cell-like decompositions of manifolds: codimension 3*, Ann. Math. (2) **110** (1979), 83-112.
- [CGM] A. Connes, M. Gromov, and H. Moscovici, *Group cohomology with Lipschitz control and higher signatures*, Geometric and Functional Analysis **3** (1993), 1 - 78.
- [CW1] R. Coifman and G. Weiss, *Analyse Harmonique Non-commutative sur Certains Espaces Homogènes*, Lecture Notes in Math. **242** (1971), Springer-Verlag.
- [CW2] R. Coifman and G. Weiss, *Extensions of Hardy spaces and their use in analysis*, Bull. Amer. Math. Soc. **83** (1977), 569-645.
- [Da] R. Daverman, *Decompositions of Manifolds*, Academic Press, 1986.
- [DS1] G. David and S. Semmes, *Strong A_∞ -weights, Sobolev inequalities, and quasiconformal mappings*, in *Analysis and Partial Differential Equations*, edited by C. Sadosky, Lecture Notes in Pure and Applied Mathematics **122**, Marcel Dekker 1990.
- [DS2] G. David and S. Semmes, *Quantitative rectifiability and Lipschitz mappings*, Transactions Amer. Math. Soc. **337** (1993), 855-889.
- [DoS] S. Donaldson and D. Sullivan, *Quasiconformal 4-manifolds*, Acta Math. **163** (1989), 181-252.
- [DFW] A. Dranishnikov, S. Ferry, and S. Weinberger, *Large Riemannian manifolds which are flexible*, preprint.
- [E] R. Edwards, *The topology of manifolds and cell-like maps*, Proceedings I.C.M. (Helsinki, 1978), 111-127.
- [FH] F. T. Farrell and W. C. Hsiang, *On Novikov's conjecture for non-positively curved manifolds, I*, Ann. Math. (2) **113** (1981), 199-209.

- [Fd] H. Federer, *Geometric Measure Theory*. Springer-Verlag, 1969.
- [Fe1] S. Ferry, *Counting simple-homotopy types of topological manifolds*, preprint.
- [Fe2] S. Ferry, *Topological finiteness theorems for manifolds in Gromov-Hausdorff space*, Duke Math. J. **74** (1994), 95-106.
- [FeW] S. Ferry and S. Weinberger, *A course approach to the Novikov conjecture*, preprint.
- [FHP] S. Ferry, I. Hambleton, and E. K. Pedersen, *A survey of bounded surgery theory and applications*, preprint.
- [Fo] G. Folland, *Subelliptic estimates and function spaces on nilpotent Lie groups*, Arkiv f. Mat. **13**(1975), 161-207.
- [FS1] G. Folland and E. Stein, *Estimates for the ∂_b -complex and analysis on the Heisenberg group*, Comm. Pure Appl. Math. **27** (1974), 429-522.
- [FS2] G. Folland and E. Stein, *Hardy Spaces on Homogeneous Groups*, Math. Notes **28** (1982), Princeton University Press.
- [FQ] M. Freedman and F. Quinn, *Topology of 4-Manifolds*. Princeton University Press, 1990.
- [Ge1] F. Gehring, *The L^p integrability of the partial derivatives of a quasiconformal mapping*, Acta Math. **130** (1973), 265-277.
- [Ge2] F. Gehring, *The Hausdorff measure of sets which link in Euclidean space*, "Contributions to Analysis: A Collection of Papers Dedicated to Lipman Bers", Academic Press, New York and London, 1974, p159-167.
- [GP] R. Greene and P. Petersen V., *Little topology, big volume*, Duke. Math. J. **67** (1992), 273-290.
- [GPW] K. Grove, P. Petersen V., and J.-Y. Wu, *Geometric finiteness theorems via controlled topology*, Invent. Math **99** (1990), 205-213; erratum, **104** (1991), 221-222.
- [Gr1] M.Gromov, *Structures Métriques pour les Variétés Riemanniennes* (J. Lafontaine, P. Pansu, ed.), Cedric/Fernand Nathan, Paris, 1981.
- [Gr2] M.Gromov, *Groups of polynomial growth and expanding maps*, Publ. Math. IHES **53** (1981), 183-215.
- [Gr3] M.Gromov, *Filling Riemannian manifolds*, J. Differential Geometry **18** (1983), 1-147.
- [Gr4] M.Gromov, *Asymptotic invariants of infinite groups*, Proceedings of the 1991 conference Sussex conference on Geometric Group Theory, Niblo and Koller, editors, London math. Soc. Lecture Notes **182**, 1993, Cambridge Univ. Press.
- [Gr5] M. Gromov, *Positive curvature, macroscopic dimension, spectral gaps and higher signatures*, IHES preprint M/95/36, 1995.
- [HaK] P. Hajłasz and P. Koskela, *Sobolev meets Poincaré*, C. R. Acad. Sci. Paris **320** (1995), 1211-1215.
- [HW] W. Hurewicz and H. Wallman, *Dimension Theory*, Princeton Univ. Press, 1941, 1969.
- [HY] J. Hocking, G. Young, *Topology*. Addison Wesley, 1961.
- [Je] D. Jerison, *The Poincaré inequality for vector fields satisfying Hörmander's condition*, Duke Math. J. **53** (1986), 503-523.
- [K] R. Kirby, *The Topology of 4-Manifolds*, Lecture Notes in Math. **1374** (1989), Springer-Verlag.

- [Ma] W. Massey, *A Basic Course in Algebraic Topology*, Springer-Verlag, 1991.
- [Mi] J. Milnor, *Topology from the Differentiable Viewpoint*, University of Virginia Press, Charlottesville, 1965.
- [Mo] E. E. Moise, *Geometric Topology in Dimensions 2 and 3*, Springer-Verlag, 1977.
- [N] L. Nirenberg, *Topics in Nonlinear Functional Analysis*, Courant Institute of Mathematical Sciences, 1974.
- [P1] P. Petersen V., *A finiteness theorem for metric spaces*, J. Diff. Geom. **31** (1990), 387-395.
- [P2] P. Petersen V., *Gromov-Hausdorff convergence of metric spaces*, Proc. Sym. Pure Math. **54**, Part 3, (1993), 489-504, Amer. Math. Soc.
- [R] W. Rudin, *Real and Complex Analysis*, McGraw-Hill, 1987.
- [Se1] S. Semmes, *Differentiable function theory on hypersurfaces in $\mathbf{R}^n$ (without bounds on their smoothness)*, Indiana Math. J. **39** (1990), 985-1004.
- [Se2] S. Semmes, *Bilipschitz mappings and strong A_∞ weights*, Ann. Acad. Sci. Fenn. A I **18** (1993), 211-248.
- [Se3] S. Semmes, *Finding structure in sets with little smoothness*, to appear, Proc. I.C.M. (Zürich, 1994), Birkhäuser.
- [Se4] S. Semmes, *On the nonexistence of bilipschitz parameterizations and geometric problems about A_∞ weights*, to appear, Revista Matemática Iberoamericana.
- [Se5] S. Semmes, *Good metric spaces without good parameterizations*, to appear, Revista Matemática Iberoamericana.
- [Se6] S. Semmes, to exist.
- [SS] L. Siebenmann and D. Sullivan, *On complexes that are Lipschitz manifolds*, "Geometric Topology", (J. Cantrell, ed.), Academic Press, 1979, 503-525.
- [St1] E. M. Stein, *Singular Integrals and Differentiability Properties of Functions*, Princeton University Press, 1970.
- [St2] E. M. Stein, *Harmonic Analysis : Real-Variable Methods, Orthogonality, and Oscillatory Integrals*, Princeton University Press, 1993.
- [Su] D. Sullivan, *Hyperbolic geometry and homeomorphisms*, "Geometric Topology", (J. Cantrell, ed.), Academic Press, 1979, 543-555.
- [V1] J. Väisälä, *Quasisymmetric embeddings in Euclidean spaces*, Trans. Amer. Math. Soc. **264** (1981), 191-204.
- [V2] J. Väisälä, *Metric duality in Euclidean spaces*, preprint.
- [Wa] F. Warner, *Foundations of Differentiable Manifolds and Lie Groups*, Scott, Foresman and Co., 1971.
- [Wh] H. Whitney, *On maps of an n -sphere into another n -sphere*, Duke Math. J. **3** (1937), 46-50.
- [Y] G. Yu, *Zero-in-the-spectrum conjecture, positive scalar curvature and asymptotic dimension*, preprint.