AD

IC/96/19

INTERNATIONAL CENTRE FOR THEORETICAL PHYSICS

CUBIC SCATTERING AMPLITUDES
FOR ALL MASSLESS REPRESENTATIONS
OF THE POINCARÉ GROUP
IN ANY SPACE-TIME DIMENSION

swg622

E.S. Fradkin

and

R.R. Metsaev

INTERNATIONAL
ATOMIC ENERGY
AGENCY

UNITED NATIONS
EDUCATIONAL,
SCIENTIFIC
AND CULTURAL
ORGANIZATION

MIRAMARE-TRIESTE

United Nations Educational Scientific and Cultural Organization
and
International Atomic Energy Agency
INTERNATIONAL CENTRE FOR THEORETICAL PHYSICS

**CUBIC SCATTERING AMPLITUDES
FOR ALL MASSLESS REPRESENTATIONS
OF THE POINCARÉ GROUP
IN ANY SPACE-TIME DIMENSION**

E.S. Fradkin¹
International Centre for Theoretical Physics, Trieste, Italy
and
Harvard University, Science Center 325,
Cambridge, MA 02138, USA

and
R.R. Metsaev²
P.N. Lebedev Physical Institute,
Leninsky prospect, 53, 117 924, Moscow, Russian Federation.

ABSTRACT

Using the language of highest weight representations and the light cone formalism we construct a full list of cubic amplitudes of scattering for all bosonic massless representations of the Poincaré group in any even space-time dimension

MIRAMARE – TRIESTE

February 1996

¹Permanent address: Department of Theoretical Physics, P.N. Lebedev Physical Institute, Leninsky prospect, 53, 117 924, Moscow, Russian Federation.

²E-mail: metsaev@lpi.ac.ru

The theory of massless higher spin fields is one of the promising branches of modern quantum field theory (see [1,2]). At present because of Refs. [3-6], a wide class of cubic interaction vertices for massless higher spin fields in flat space is known. These vertices turn out to be non-minimal for the case of interactions of higher spin fields with gravity. Cubic vertices for massless higher spin fields living in anti-de Sitter space-time were constructed in [7]. References [3-7] were devoted to the analysis of interactions in $D = 4$. A full list of cubic interaction vertices in the case of $D = 4$ flat space was given in [6]. In this paper we continue the investigation of the problem introducing the interaction for massless higher spin fields in higher space-time dimensions ($D > 4$) which was initiated in [8]. In Ref.[8] a full list of cubic interaction vertices for totally symmetric massless on mass shell representations of the Poincaré group for any D was obtained. Notice the results of Ref.[8] as well as of this paper do not contradict those of Refs.[9-12] because in our vertices for the case of interactions of higher spins with a graviton there are higher derivatives only. As is well known various types of symmetry properties of massless representations classify according to unitary irreps of the transversal $SO(D-2)$ group, i.e. in addition to the totally symmetric representations there are mixed symmetry ones [13-19]. It is likely that in studies of the self-consistency of higher (fourth,...) order interactions the former representations should be taken into consideration. We think that the higher order interaction vertices will be effectively nonlocal. The aim of this paper is to construct cubic interaction vertices for all massless representations of the Poincaré group in any even space-time dimension $D = 2n + 2$. To solve the problem it has been suggested [20] that the language of highest weight representations of the $SO(2n)$ group be used, and it was demonstrated how the procedure of construction of cubic interaction vertices works in the simplest cases $D = 5, 6$. The language of highest weight turns out to be extremely efficient for the analysis of cubic interaction vertices of any symmetry representations. Of course, use of this language leads to breaking of the manifest $SO(2n)$ transversal invariance. It is the price we pay in solving the problem under consideration. Efficiency of the different way of breaking of the manifest transversal invariance ($SO(8) \rightarrow SO(6) \times SO(2)$) has been demonstrated in Ref.[21] by

constructing the cubic vertices in the $D = 10$ superstring theory.

Before we proceed let us comment on the self-consistency of higher (fourth,...) order interaction vertices. It is known (see Ref.[4]) that cubic vertices do not guarantee self-consistency of fourth order interactions. However due to Refs.[22] the possibilities for constructing consistent higher order vertices increased. Namely, in [22] for the case of massless higher spin fields living in $D = 4$ anti-de Sitter space time the following remarkable fact was discovered: in order to solve the problem of higher order vertices it is necessary to use the spectrum of massless particles that contains every massless representation just once. Taking that into account one can suppose that the problem of higher order vertices in a flat space-time may also be solved by using the spectrum which contains every massless representation of Poincaré group just once. Another fact that supports this point of view is provided by the following reasoning. There exists the so called most singular representation (the oscillator representation) of the orthosymplectic algebra $osp(8)$ that has the following highly interesting property: when it is restricted to the Poincaré or anti-de Sitter algebra then it turns out to contain every massless bosonic representation of corresponding algebra just once (see [23]). In other words, the above-mentioned spectra can be obtained from the one and the same representation of the $osp(8)$! Because of that it is strongly believed that the difficulties of higher order vertices may be solved by incorporating an infinite number of different massless particles (every massless representation of the Poincaré algebra just once). The investigation of fourth order to interaction carried out in [24] supports this point of view.

Starting from the commutation relations (CR's) of the Poincaré algebra

$$\begin{aligned} [J^{\mu\nu}, J^{\rho\sigma}] &= i\eta^{\mu\rho}J^{\sigma\nu} + \dots, \quad [P^\mu, J^{\rho\sigma}] = i\eta^{\mu\rho}P^\sigma + \dots, \\ [P^\mu, P^\nu] &= 0, \quad \eta^{\mu\nu} = \text{diag}(+1, -1, \dots, -1), \end{aligned} \quad (1)$$

we introduce the light cone coordinates $x^\pm = (x^0 \pm x^{2n+1})/\sqrt{2}$, x^I , $I = 1, \dots, 2n$ and consider x^+ as the evolution parameter. Without loss of generality we analyse (1) for $x^+ = 0$. In the light cone coordinates the Poincaré algebra splits into a kinematical part spanned by the generators $P_-, P^I, J^{IK}, J^{+-}, J^{+I}$, which are usually realized quadratically in the physical

fields, and a dynamical part spanned by the generators $H \equiv P_+$ and J^{-I} realized nonlinearly.

Since we use extensively the constructions of Ref.[25], we would like to introduce the complex extension of the rotation group, denoted as $SO(2n, C)$. After that we transform the quadratical form $(x, x) = x^I x^I$ to the antidiagonal form $(x, x) = x^I s^{IJ} x^J$, where $s^{IJ} \equiv \delta^{I, 2n-J+1}$. It will be convenient to introduce the index $i = 1, \dots, n$ and split all vectors as $A^I = (A^i, A^{2n-i+1})$, after that we have the scalar product $(A, B) = \sum_{i=1}^n (A^i B^{2n-i+1} + A^{2n-i+1} B^i)$. For the case of small transversal indices $i, j, \dots$ we drop the summation over repeated indices. The reality condition can be written as $(A^I)^* = s^{IJ} A^J$.

Differential realization $r(G)$ of the on-shell representation $[p_+ = p^I s^{IK} p^K / (2p_-)]$ of the Poincaré algebra has the form (see Ref.[26])

$$\begin{aligned} r(P_-) &= \beta, \quad r(P_+) = h, \quad r(P^I) = p^I, \quad r(J^{+-}) = -\hat{x}^- \beta, \\ r(J^{IK}) &= \hat{x}^I p^K - \hat{x}^K p^I + M^{IK}, \quad r(J^{+I}) = -\hat{x}^I \beta, \\ r(J^{-I}) &= \hat{x}^- p^I - \hat{x}^I h - M^{IK} s^{KL} p^L / \beta, \end{aligned} \quad (2)$$

where $\beta \equiv p_-$, $h \equiv p^I s^{IK} p^K / (2\beta)$, $\hat{x}^I \equiv \iota s^{IK} \partial / \partial p^K$, $\hat{x}^- \equiv -\iota \partial / \partial \beta$, and M^{IK} are the generators of the $SO(2n)$ group.

To use the standard framework of the highest weight representation we should convert generators of $SO(2n)$ from the tensor form to the Cartan-Weyl form. To do that let us introduce anti-Hermitian generators of $SO(2n)$: $\mathcal{M}^{I,J} \equiv \iota M^{IJ}$ which satisfy the following CR's: $[\mathcal{M}^{I,J}, \mathcal{M}^{K,L}] = s^{I,K} \mathcal{M}^{L,J} + 3 \text{ perms.}$ Now we link the $\mathcal{M}^{I,J}$ with elements of $so(2n)$ algebra taken in the Cartan-Weyl form in the following fashion:

$$\begin{aligned} S_{l,k}^+ &\equiv \mathcal{M}^{2n-l+1, l+k}, \quad S_{l,k}^- \equiv \mathcal{M}^{n-l-k+1, l}, \\ \tilde{S}_{l,k}^+ &\equiv \mathcal{M}^{n+k+1, l+k}, \quad \tilde{S}_{l,k}^- \equiv \mathcal{M}^{2n-l-k+1, n-k}, \\ S_{l,k}^0 &\equiv \frac{1}{2}(M_l - M_{l+k}), \quad \tilde{S}_{l,k}^0 \equiv \frac{1}{2}(M_{n-k} + M_{2n-l-k+1}), \end{aligned} \quad (3)$$

where $M_l \equiv \mathcal{M}^{2n-l+1, l}$. Throughout this paper, unless otherwise specified, the indices l, k for $S_{l,k}^{0\pm}$ and $\tilde{S}_{l,k}^{0\pm}$ run over $l = 1, \dots, n-k-1$, $k = 1, \dots, n-1$, and $l = n-k, \dots, n$, $k = 1, \dots, n-1$ respectively. The generators $S_{l,k}^{0\pm}$ form the $su(n)$ subalgebra of $so(2n)$ while

$\tilde{S}_{n,1}^{\pm 0}$ and $S_{l,1}^{\pm 0}$ form Chevalley basis of $so(2n)$. The Cartan subalgebra is spanned by $\tilde{S}_{n,1}^0$ and $S_{l,1}^0$. The highest weight vector $|\mathbf{m}\rangle$ [where $\mathbf{m} = (m_1, \dots, m_n)$] is defined in the following way

$$\left[S_{l,1}^+ = \tilde{S}_{n,1}^+ = (2S_{l,1}^0 - p_l) = (2\tilde{S}_{n,1}^0 - p_n) \right] |\mathbf{m}\rangle = 0, \quad (4)$$

where $p_l \equiv m_l - m_{l+1}$, $l = 1, \dots, n-1$, $p_n \equiv m_{n-1} + m_n$. The numbers $m_1, \dots, m_n$, the so-called weight of the representation, satisfy the inequalities: $m_1 \geq m_2 \geq \dots \geq m_{n-1} \geq |m_n|$. To carry out calculations it is necessary to have a certain realization of representation. Let us briefly describe the realization we use (for more details see Refs.[25,27]). First, for any $g^c \in SO(2n, C)$ we have the Gauss decomposition

$$g^c = g_- g_0^{(\delta)} g_+^{(\eta, \xi)}, \quad g_+^{(\eta, \xi)} = \begin{pmatrix} \eta & \eta \xi s \\ 0 & s \eta^{(t)-1} s \end{pmatrix}, \quad (5)$$

where

$$g_0^{(\delta)} = \text{diag}(\delta_1, \dots, \delta_n, \delta_n^{-1}, \dots, \delta_1^{-1}),$$

$\eta_{i,j} = 0$, for $i > j$, $\eta_{i,i} = 1$, $\xi_{i,j} = -\xi_{j,i}$, $s_{i,j} = \delta_{i,n-j+1}$ and g_- is lower triangular matrix with units on the diagonal. The superscripts “(t)” and “-1” denote a matrix transposition and inversion respectively. Now the representation of $SO(2n)$ group of weight $\mathbf{m}$ is constructed on the space of the functions $\Phi(\eta, \xi)$ in the following way: for all $g \in SO(2n)$ one defines the operator T_g as

$$(T_g \Phi)(\eta, \xi) = \delta_1^{m_1} \dots \delta_n^{m_n} \Phi(\eta \cdot g, \xi \cdot g), \quad (6)$$

where $\eta \cdot g$, $\xi \cdot g$ as well as δ_i are defined from the decomposition $g_+^{(\eta, \xi)} g = g_- g_0^{(\delta)} g_+^{(\eta \cdot g, \xi \cdot g)}$. Using (5-6) one can calculate the generators of a representation. In the quite extensive literature we find expressions for the Chevalley basis or for the S^+ part of the algebra only. Since we will need manifest expressions for all generators let us write down the results of our calculations (here and henceforth α designates the set $\{\eta, \xi\}$):

$$S_{l,k}^+(\alpha) = \sum_j (\eta_{j,l} \frac{\partial}{\partial \eta_{j,l+k}} - \xi_{j,l+k} \frac{\partial}{\partial \xi_{j,l}}).$$

$$\begin{aligned}
\tilde{S}_{l,k}^+(\alpha) &= \frac{\partial}{\partial \xi_{n-k, 2n-l-k+1}}, \\
S_{l,k}^-(\alpha) &= \sum_j m_j \eta_{l,j}^{-1} \eta_{j,l+k} - \sum_j \xi_{j,l} \frac{\partial}{\partial \xi_{j,l+k}} \\
&\quad + \sum_{i,j,b} \varepsilon^{i,b} \eta_{l,b}^{-1} \eta_{b,j} \eta_{i,l+k} \frac{\partial}{\partial \eta_{i,j}}, \\
\tilde{S}_{l,k}^-(\alpha) &= \sum_j (\xi_{j, 2n-l-k+1} S_{n-k, j+k-n}^- \\
&\quad - \xi_{j, n-k} S_{2n-l-k+1, j+l+k-2n-1}^-) \\
&\quad + \sum_{i < j} (\xi_{i, n-k} \xi_{j, 2n-l-k+1} - \xi_{j, n-k} \xi_{i, 2n-l-k+1}) \frac{\partial}{\partial \xi_{i,j}}, \\
M_l(\alpha) &= m_l + \sum_j (\varepsilon^{l,l} \eta_{j,l} \frac{\partial}{\partial \eta_{j,l}} - \varepsilon^{l,j} \eta_{l,j} \frac{\partial}{\partial \eta_{l,j}} - \xi_{l,j} \frac{\partial}{\partial \xi_{l,j}}),
\end{aligned} \tag{7}$$

where $\varepsilon^{i,j} = 1(0)$ for $i < j$ ($i \geq j$). In expression for $\tilde{S}_{l,k}^-$ there are terms $S_{i,j}^-$ with $j \leq 0$ which should be rewritten according to the rules $S_{i,j,-j}^+ = S_{i,j}^+$, $S_{i,0}^- = M_i$ for $i, j > 0$. In (7) and below, unless otherwise specified, the indices i, j, b run over $1, \dots, n$.

From now on we consider η_{ij} for $i < j$ and ξ_{ij} as the creation operators, η_{ii} as the unit operator and make, in (7), the substitutions $\partial/\partial \eta_{i,j} \rightarrow \bar{\eta}_{i,j}$ for $i < j$, $\partial/\partial \xi_{ij} \rightarrow \bar{\xi}_{ij}$, where $\bar{\eta}_{i,j}$ and $\bar{\xi}_{i,j}$ are the annihilation operators ($\bar{\eta}_{i,j}|\mathbf{m}\rangle = 0 = \bar{\xi}_{i,j}|\mathbf{m}\rangle$) with CR's

$$[\bar{\eta}_{i,j}, \eta_{i',j'}] = \varepsilon^{i,j} \delta_{i,i'} \delta_{j,j'}, \quad [\bar{\xi}_{i,j}, \xi_{i',j'}] = \delta_{i,i'} \delta_{j,j'} - \delta_{i,j'} \delta_{j,i'}.$$

Using the method of Gelfand and Zetlin one can introduce the orthogonal basis system of the representation (in this work we rely only on the existence of such a basis (see Ref.[28])): $|\mathbf{m}, \{\lambda\}\rangle = N(\mathbf{m}, \{\lambda\}) \alpha^{\{\lambda\}} |\mathbf{m}\rangle$ where the set of numbers $\{\lambda\}$ designate the basis and N is a normalization factor. Now we introduce the generating function $|\Phi_{\mathbf{m}}(p, \alpha)\rangle \equiv \sum_{\{\lambda\}} \Phi_{\mathbf{m}}^{\{\lambda\}}(p) |\mathbf{m}, \{\lambda\}\rangle$, where, by definition, the fields $\Phi_{\mathbf{m}}^{\{\lambda\}}(p)$ (the dependence of fields on β is not shown explicitly) realize the highest weight representation of the $SO(2n)$ group. Writing the CR's for $\Phi_{\mathbf{m}}$ in the form

$$[\Phi_{\mathbf{m}}^{\{\lambda\}}(p), \Phi_{\mathbf{m}'}^{\{\lambda'\}}(p')]|_{equal\ x+} = \delta_{\mathbf{m}, \lambda; \mathbf{m}', \lambda'} \frac{\delta^{(2n+1)}(p+p')}{2\beta}, \tag{8}$$

one can construct quadratic level generators of the Poincaré algebra:

$$G_{\mathbf{m}} = \int \beta d\beta d^{(2n)}p \langle \Phi_{\mathbf{m}} | r(G_{\mathbf{m}}) | \Phi_{\mathbf{m}} \rangle.$$

Now let us study the cubic corrections to H and J^{-I} . Analyzing the CR's of H and J^{-I} with P_- , P^I , J^{+I} and the CR's $[H, J^{-I}]$ and $[J^{-I}, J^{-K}]$ in the cubic approximation we get the following structure for H and J^{-I} describing the interaction of the fields $\Phi_{\mathbf{m}_a}$ (here and henceforth $a = 1, 2, 3$ labels the three interacting fields)

$$H_3 = \int d\Gamma_3 \langle \Phi_3 | \mathbf{h}_3 \rangle. \quad (9a)$$

$$J_3^{-I} = \int d\Gamma_3 \left(\langle \Phi_3 | \mathbf{j}_3^{-I} \rangle - \frac{1}{3} \left(\sum_a \hat{x}_a^I \langle \Phi_3 | \right) | \mathbf{h}_3 \rangle \right), \quad (9b)$$

$$|\mathbf{h}_3\rangle = h_3(\alpha, \mathbf{P}, \beta) \prod_a |\mathbf{m}_a\rangle. \quad (10a)$$

$$|\mathbf{j}_3^{-I}\rangle = -(h^{(3)})^{-1} \sum_a r(J_a^{-I})^\dagger |\mathbf{h}_3\rangle, \quad (10b)$$

where $\langle \Phi_3 | \equiv \prod_a \langle \Phi_{\mathbf{m}_a}(p_a, \bar{\alpha}_a) |$, $h^{(3)} \equiv \sum_a h_a$,

$$d\Gamma_3 \equiv \delta^{(2n)} \left(\sum_a p_a^I \right) \delta \left(\sum_a \beta_a \right) \prod_a d\beta_a d^{(2n)}p_a, \quad (11)$$

$$h_a \equiv \frac{p_a^I s^{IJ} p_a^J}{2\beta_a}, \quad h^{(3)} = -\frac{\mathbf{P}^I s^{IJ} \mathbf{P}^J}{2\beta_1 \beta_2 \beta_3}, \quad \mathbf{P}^I \equiv \frac{1}{3} \sum_a \check{\beta}_a p_a^I, \quad (12)$$

$$r(J_a^{-I})^\dagger = p_a^I \hat{x}_a^- - h_a \hat{x}_a^I - M_a^{IK} s^{KL} p_a^L / \beta_a, \quad (13)$$

and $\check{\beta}_a \equiv \beta_{a+1} - \beta_{a+2}$, $\beta_4 \equiv \beta_1$, $\beta_5 \equiv \beta_2$. The CR of H with J^{+-} tells us that $|\mathbf{h}_3\rangle$ is a zero-degree homogeneity function with respect to β (the dependence of $\mathbf{P}$ on β should also be taken into account):

$$\left(\sum_a \beta_a \frac{\partial}{\partial \beta_a} + \Delta(\mathbf{P}) \right) h_3(\alpha, \mathbf{P}, \beta) = 0, \quad (14a)$$

$$\Delta(\mathbf{P}) \equiv \sum_i (\mathbf{P}^i \frac{\partial}{\partial \mathbf{P}^i} + \mathbf{P}^{2n-i+1} \frac{\partial}{\partial \mathbf{P}^{2n-i+1}}). \quad (14b)$$

From CR's of H with J^{IK} one gets the equations $\sum_a r(J_a^{IK}) |\mathbf{h}_3\rangle = 0$ which can be rewritten in terms of $h_3(\alpha, \mathbf{P}, \beta)$ (10a):

$$(\mathcal{L}^{I,K}(\mathbf{P}) + \sum_a \mathcal{M}_a^{I,K}(\alpha))h_3(\alpha, \mathbf{P}, \beta) = 0, \quad (15a)$$

$$\mathcal{L}^{I,K}(\mathbf{P}) \equiv \mathbf{P}^I \frac{\partial}{\partial \mathbf{P}^{2n-K+1}} - (K \leftrightarrow I). \quad (15b)$$

To simplify our equations we do not write the Fock vacuum explicitly and consider $\mathcal{M}$, S , and $\tilde{S}$ as differential operators (7). The operator $\mathcal{L}^{I,K}(\mathbf{P})$ is an anti-Hermitean form of the orbital part of the angular momentum and can be obtained from the equation $\mathcal{L}^{I,K}(\mathbf{P})|h_3\rangle = i \sum_a (\hat{x}_a^I p_a^K - \hat{x}_a^K p_a^I)|h_3\rangle$ where we take into account that $\mathbf{P}$ variables exhaust all p_a^I dependence of $|h_3\rangle$. The procedure of introducing the Cartan-Weyl basis for $\mathcal{L}$, denoted as $L_{l,k}^{\pm 0}$ and $\tilde{L}_{l,k}^{\pm 0}$, is identical to the one for $\mathcal{M}^{I,K}$ [in (3) it is necessary to make the substitutions L for S and $\mathcal{L}$ for $\mathcal{M}$]. In the case of a massless fields we can restrict ourselves to the interaction having κ th power derivatives in the vertex. The corresponding equation on the vertex is: $(\Delta(\mathbf{P}) - \kappa)h_3(\alpha, \mathbf{P}, \beta) = 0$.

In fact, the set of vertices h_3 satisfying Eqs.(14) and (15) is the solution to all CR's of the Poincaré algebra. However, from the set of solutions of physical interest are the vertices satisfying the additional condition which we call the locality condition. Let us formulate it. Expressions $|h_3\rangle$ and $|j_3^{-I}\rangle$ (10) must be regular in the limit $h^{(3)} \rightarrow 0$, and to have a nontrivial 3-point amplitude of scattering it is supposed that $h_{3(0)} \neq 0$ [for the definition of $h_{3(0)}$ see (16b)]. From (10b) it is clear that $|j_3^{-I}\rangle$ will have a singularity when $h^{(3)} \rightarrow 0$ due to $h_{3(0)} \neq 0$, and this singularity can be cancelled by an appropriate selection of $h_{3(0)}$. Thus we have the conditions

$$\sum_a r(J_a^{-I})^\dagger h_3(\alpha, \mathbf{P}, \beta) \Big|_{h^{(3)} \rightarrow 0} = 0, \quad (16a)$$

$$h_{3(0)} \equiv h_3(\alpha, \mathbf{P}, \beta) \Big|_{h^{(3)} \rightarrow 0} \neq 0. \quad (16b)$$

The eqs.(14-16) are the complete system of equations on the tree-level cubic interaction vertices which we are going to solve. We analyze Eqs.(14)–(16) in the following way.

(a) First, we are solving the “+” part of eqs.(15a):

$$\begin{aligned} (L_{l,k}^+(\mathbf{P}) + \sum_a S_{l,k}^{+(a)}(\alpha))h_3(\alpha, \mathbf{P}, \beta) &= 0, \\ (\tilde{L}_{l,k}^+(\mathbf{P}) + \sum_a \tilde{S}_{l,k}^{+(a)}(\alpha))h_3(\alpha, \mathbf{P}, \beta) &= 0. \end{aligned} \quad (17)$$

To solve Eqs.(17) it is convenient, in place of $\mathbf{P}^I$, to introduce new “momentum” variables:

$q_{i,i+1}$, $i = 1, \dots, n-1$; ρ_i , $i = 1, \dots, n$, and $\mathbf{P}^n$ where

$$q_{ij} \equiv \theta^{i,j} \frac{\mathbf{P}^i}{\rho_j}, \quad \rho_j \equiv \sum_i \mathbf{P}^{2n-i+1} q_{ij}.$$

and $\theta^{i,j} \equiv \delta^{i,j} + \varepsilon^{i,j}$. As a result we take the following solution of Eqs.(17)

$$h_3 = h_3(\alpha^{(1)}, \rho_1, \bar{\rho}_n, \beta), \quad (18a)$$

$$\xi_{ij}^{(1)(a)} \equiv (\tau^{-1} q^{-1} \xi^{(a)} q^{-1} \tau^{-1})_{ij} - \frac{\mathbf{P}^n}{\rho_1} (\delta_{i,1} \delta_{j,n} - \delta_{i,n} \delta_{j,1}), \quad (18b)$$

$$\eta_{ij}^{(1)(a)} \equiv (\eta^{(a)} q \tau)_{ij}, \quad (18c)$$

where

$$\tau_{i,j} \equiv \delta_{i,j} - \frac{\rho_j}{\rho_1} \delta_{i,1} \varepsilon^{1,j}, \quad \bar{\rho}_n \equiv \rho_n \mathbf{P}^n.$$

Recall that α designates the set $\{\eta, \xi\}$. Furthermore, there are additional equations on the $|\mathbf{h}_3\rangle$ [they are the remains of Eqs.(17)], which we would like to write after introducing the dimensionless variable $\bar{\rho}_n \equiv \rho_n \mathbf{P}^n / (\rho_1)^2$ (notice that the variables $\alpha^{(1)}$ are also dimensionless) and rewriting expression (18a) in the form $h_3 = (\rho_1)^\kappa h_{3\kappa}(\alpha^{(1)}, \bar{\rho}_n, \beta)$. We have written down the dimensionfull factor $(\rho_1)^\kappa$ explicitly. Now the relevant equations can be written as

$$\sum_a S_{l,k}^{+(a)}(\alpha^{(1)}) h_{3\kappa}(\alpha^{(1)}, \bar{\rho}_n, \beta) = 0, \quad (19a)$$

$$\sum_a \tilde{S}_{n+i-j+1,n-i}^{+(a)}(\alpha^{(1)}) h_{3\kappa}(\alpha^{(1)}, \bar{\rho}_n, \beta) = 0, \quad (19b)$$

where $l = 2, \dots, n-k$, $k = 1, \dots, n-1$ and $i, j = 2, \dots, n-1$.

(b) Second, we rewrite “+” and “0” parts of Eqs.(15a) in terms of the variables $\alpha^{(1)}$ and $\bar{\rho}_n$. The equations obtained are

$$\sum_a [S_{l,k}^{+(a)}(\alpha^{(1)}) + \delta_{l,1} \rho_n \tilde{S}_{n+1-k,n-1}^{+(a)}(\alpha^{(1)})] h_{3\kappa}(\alpha^{(1)}, \bar{\rho}_n, \beta) = 0, \quad (20a)$$

$$\begin{aligned} & \sum_a [\tilde{S}_{n+i-j+1,n-i}^{+(a)}(\alpha^{(1)}) + \bar{\rho}_n (\delta_{i,1} \varepsilon^{1,j} S_{1,j-1}^{+(a)}(\alpha^{(1)}) - (i \leftrightarrow j))] \\ & \times h_{3\kappa}(\alpha^{(1)}, \bar{\rho}_n, \beta) = 0, \quad i, j = 1, \dots, n-1, \end{aligned} \quad (20b)$$

$$(\sum_a M_l^{+(a)}(\alpha^{(1)}) + \delta_{l,1} (\kappa - 2\bar{\rho}_n \frac{\partial}{\partial \bar{\rho}_n})) h_{3\kappa}(\alpha^{(1)}, \bar{\rho}_n, \beta) = 0 \quad (20c)$$

(in eq (20c) $l = 1, \dots, n$).

(c) Third, writing Eqs.(16a) in terms of the same variables we obtain:

$$\sum_a \check{\beta}_a S_{1,j-1}^{-(a)}(\check{\alpha}^{(1)}) h_{3\kappa(0)}(\check{\alpha}^{(1)}, \beta) = 0, \quad (21a)$$

$$\sum_a \check{\beta}_a \check{S}_{j,n-1}^{-(a)}(\check{\alpha}^{(1)}) h_{3\kappa(0)}(\check{\alpha}^{(1)}, \beta) = 0, \quad (21b)$$

$$\sum_a \check{\beta}_a (\beta_a \frac{\partial}{\partial \beta_a} - M_1^{(a)}(\check{\alpha}^{(1)})) h_{3\kappa(0)}(\check{\alpha}^{(1)}, \beta) = 0, \quad (21c)$$

where $j = 2, \dots, n$. The $h_{3\kappa(0)}$ is defined by (16b) making there the substitutions $h_{3(0)} \rightarrow h_{3\kappa(0)}$ and $h_3 \rightarrow h_{3\kappa}$. The set of Eqs.(19)-(21) supplemented by Eq.(14) is the complete system equations for $h_{3\kappa}(\check{\alpha}^{(1)}, \bar{\rho}_n, \beta)$. Since $h_{3\kappa(0)}$ which is a dimensionless 3-point amplitude of scattering, is the object of prime physical interest we restrict ourselves to solutions of Eqs.(14),(19)-(21) for $h_{3\kappa(0)}(\check{\alpha}^{(1)}, \beta)$; i.e., from now on we analyze the relevant equations in $(\bar{\rho}_n)^0$ approximation.

Equations (19)-(20) express the invariance conditions with respect to the transversal rotations. In the approaches based on the tensor realization of the representations of the transversal group one would first analyze these equations and then the rest ones (see Refs.[8,29]) while in the approach based on the highest weight [20] it turns out to be more convenient to use the following procedure in solving (19)-(21).

(i) multiplying the $l = 1$ part of Eq.(20a) and the $l = 1$ part of Eq.(20b) by $\check{\beta}_1$, subtracting Eqs.(21a) and (21b) respectively from the resulting expressions, and repeating this procedure for cyclic permutations of the particle labels (1,2,3), one obtains the equations

$$\left(\frac{S_{1,j-1}^{-(1)}(\check{\alpha}^{(1)})}{\beta_1} - \frac{S_{1,j-1}^{-(2)}(\check{\alpha}^{(1)})}{\beta_2} = \frac{S_{1,j-1}^{-(3)}(\check{\alpha}^{(1)})}{\beta_3} \right) h_{3\kappa(0)}(\check{\alpha}^{(1)}, \beta), \quad (22a)$$

$$\left(\frac{\check{S}_{j,n-1}^{-(1)}(\check{\alpha}^{(1)})}{\beta_1} - \frac{\check{S}_{j,n-1}^{-(2)}(\check{\alpha}^{(1)})}{\beta_2} = \frac{\check{S}_{j,n-1}^{-(3)}(\check{\alpha}^{(1)})}{\beta_3} \right) h_{3\kappa(0)}(\check{\alpha}^{(1)}, \beta), \quad (22b)$$

where $j = 2 \dots n$. The general solution of eqs.(22) is

$$\begin{aligned} h_{3\kappa(0)}(\check{\alpha}^{(1)}, \beta) &= \prod_a \prod_{\nu=1}^{n-1} (\xi_{\nu}^{(a)} h_{\nu,a}^{(1,1+1)})^{m_{\nu}^{(a)} - m_{\nu+1}^{(a)}} \\ &\times (\xi_{n-1}^{(a)})^{2m_n^{(a)}} h_{\nu}(\check{\alpha}^{(1)}, X, \beta), \end{aligned} \quad (23)$$

where we have introduced the variables

$$\begin{aligned} X_\sigma &\equiv \sum_a \beta_a / \eta_\sigma^{(3)(a)}, \quad X_{2n-\sigma-1} \equiv \sum_a \beta_a / \xi_\sigma^{(3)(a)}, \\ \eta_{\sigma,\gamma}^{(3)} &\equiv \sum_\nu \Theta_{\sigma,\nu} \eta_{\nu,\gamma}^{(2)}, \quad \xi_{\sigma,\gamma}^{(3)} \equiv \xi_{\sigma,\gamma}^{(2)}, \end{aligned} \quad (24)$$

$$\begin{aligned} \frac{1}{\eta_\sigma^{(3)}} &\equiv \frac{1}{\eta_\sigma^{(2)}} - \sum_\nu \frac{1}{\xi_{1,\sigma}^{(2)}} \xi_{\sigma,\nu}^{(2)} \eta_{1,\nu}^{(2)}, \quad \xi_\sigma^{(3)} \equiv \frac{1}{\eta_{1,\sigma}^{(2)}} \xi_{1,\sigma}^{(2)}, \\ \xi_{\sigma,\gamma}^{(2)} &\equiv \xi_{\sigma+1,\gamma+1}^{(1)} - \frac{\xi_{1,\sigma+1}^{(1)}}{\eta_\gamma^{(2)}} + \frac{\xi_{1,\gamma+1}^{(1)}}{\eta_\sigma^{(2)}}, \end{aligned} \quad (25)$$

$$\begin{aligned} \eta_{\sigma,\gamma}^{(2)} &\equiv \sum_\nu \varepsilon^{\sigma,\nu} \frac{\eta_{1,\nu}^{(1)}}{\eta_{1,\sigma+1}^{(1)}} \eta_{\nu,\gamma+1}^{(1)}, \quad \xi_\sigma^{(2)} \equiv \sum_\nu \eta_{\sigma,\nu}^{(2)} \xi_{1,\nu+1}^{(1)}, \\ \frac{1}{\eta_\sigma^{(2)}} &\equiv - \sum_\nu \frac{\eta_{\sigma,\nu}^{(2)}}{\eta_{1,\nu+1}^{(1)}}, \quad \Theta_{\sigma,\nu} \equiv \delta_{\sigma,\nu} - \delta_{\sigma,\nu-1} \frac{\xi_{\nu-1}^{(2)}}{\xi_\nu^{(2)}} \end{aligned} \quad (26)$$

and $\eta^{\sigma,\beta}$ is the inversion of $\eta_{\sigma,\beta}^{(r)}$. In (24)–(26) the indices σ, γ, ν run over $1, \dots, n-1$.

Let us make a comment on the variables introduced. To solve Eqs.(22) we diagonalize the generators $S_{1,k}^{-(a)}$ first and then $\tilde{S}_{l,n-1}^{-(a)}$, i.e. we split the solution procedure in two stages. The superscripts (2) and (3) are used to indicate these stages.

(ii) now we rewrite eqs.(19) and $l = 1$ part of eqs.(20c) in terms of the new variables (24). The result of this procedure is

$$(L_{\sigma,\gamma}^{\pm 0}(X) + \sum_a S_{\sigma,\gamma}^{\pm 0(a)}(\alpha^{(3)})|_m) \bar{h}(\alpha^{(3)}, X, \beta) = 0, \quad (27a)$$

$$(\tilde{L}_{\sigma,\gamma}^{\pm 0}(X) + \sum_a \tilde{S}_{\sigma,\gamma}^{\pm 0(a)}(\alpha^{(3)})|_{\bar{m}}) \bar{h}(\alpha^{(3)}, X, \beta) = 0, \quad (27b)$$

$$(\Delta(X) + \kappa - \sum_a m_1^{(a)}) \bar{h}(\alpha^{(3)}, X, \beta) = 0, \quad (27c)$$

where $\Delta(X)$ and $L_{l,k}^{\pm 0}(X)$ can be obtained from (14b) and (15b) respectively making there the substitutions $n \rightarrow n-1$ and $P \rightarrow X$. The $S_{\sigma,\gamma}^{\pm 0}|_m$ and $\tilde{S}_{\sigma,\gamma}^{\pm 0}|_{\bar{m}}$ are obtained from (7), making there the shift $n \rightarrow n-1$ and then substituting $m_\sigma \rightarrow \bar{m}_\sigma = \varepsilon^{\sigma,n-1} m_{\sigma+1} - \delta_{\sigma,n-1} m_n$, $\sigma = 1, \dots, n-1$. In Eqs.(27) the indices σ, γ for $L_{\sigma,\gamma}^{\pm 0}, S_{\sigma,\gamma}^{\pm 0}|_m$ and $\tilde{L}_{\sigma,\gamma}^{\pm 0}, \tilde{S}_{\sigma,\gamma}^{\pm 0}|_{\bar{m}}$ run over $\sigma = 1, \dots, n-1, \gamma = 1, \dots, n-2$ and $\sigma = n-\gamma-1, \dots, n-1, \gamma = 1, \dots, n-2$ respectively.

(iii) rewriting Eq.(21c) and Eq.(14) in terms of (24) we get

$$\sum_a \check{\beta}_a (\beta_a \frac{\partial}{\partial \beta_a} + m_1^{(a)}) \bar{h}(\alpha^{(3)}, X, \beta) = 0, \quad (28a)$$

$$\sum_a (\beta_a \frac{\partial}{\partial \beta_a} + m_1^{(a)}) \bar{h}(\alpha^{(3)}, X, \beta) = 0. \quad (28b)$$

It should be emphasized that at this stage only Eq.(28a) expresses the locality condition while Eqs.(27) and (28b) reflect rotation invariance and homogeneity conditions. Thus at this stage we have reduced the $2n - 1$ locality condition Eqs.(21) to the single eq.(28a). The general solution of Eqs.(27) and (28) is

$$\bar{h}(\alpha^{(3)}, X, \beta) = \prod_a \beta_a^{-m_1^{(a)}} G(\alpha^{(3)}, X), \quad (29)$$

where $G(\alpha^{(3)}, X)$ is the generating function of the Clebsch-Gordan coefficients describing the coupling of four representations of the $SO(2n - 2)$ group: three representations with highest weights $\bar{\mathbf{m}}^{(a)}$ and one vector representation (X). The $G(\alpha^{(3)}, X)$ satisfies the equations obtained from eqs.(27) making there the substitution $\bar{h} \rightarrow G(\alpha^{(3)}, X)$.

Now collecting all steps in derivation of $h_{3(0)}$ we have our final result:

$$\begin{aligned} h_{3(0)} &= (\bar{\rho}_1)^\kappa \prod_a \beta_a^{-m_1^{(a)}} \prod_a \prod_{\nu=1}^{n-1} (\xi_{\nu}^{(2)(a)} \eta_{(a)}^{(1)1, \nu+1})^{m_{\nu}^{(a)} - m_{\nu+1}^{(a)}} \\ &\times (\xi_{n-1}^{(2)(a)})^{2m_n^{(a)}} G(\alpha^{(3)}, X). \end{aligned} \quad (30)$$

One can make sure that the $h_{3\kappa(0)}$ obtained satisfies the remainder of Eqs.(19)–(21).

For $h_{3\kappa(0)}$ to be well defined it must be a polynomial in primary operators α (5) or, what is the same, polynomial in $\alpha^{(1)}$ (18). As seen from (24)–(26) the nonpolynomial $\alpha^{(1)}$ dependence of $\alpha^{(3)}$ is the reason why the $h_{3\kappa(0)}$ is not polynomial generally. It turns out that, for the $h_{3\kappa(0)}$ to be polynomial, we must impose certain restrictions on the $\bar{\mathbf{m}}^{(a)}$ and κ [we must choose $\bar{\mathbf{m}}^{(a)}$ and κ in such a way that $\eta^{(1)1, \gamma}$ and $\xi_{\nu}^{(2)}$ coming from the prefactor of Eq.(30) suppress the maximal negative power of these variables coming from the G]. It is precisely the restrictions which leave no place for the minimal gravitational interaction of massless higher spin fields (see Refs.[20,21]).

Now let us briefly describe the progress which has been made in the paper. We have started with eq.(15) which expresses invariance condition with respect to rotations of the $SO(2n)$ group and with eq (16) which expresses locality condition. As a result, we are finishing with expression (30) where the G is the generating function of the Clebsch–Gordan coefficients. Thus we have satisfied the locality condition. The expression (30) is the main result of the paper.

As an illustration of (30) let us consider particular cases $D = 5, 6$. Five and six dimensions is an arena where the complex structure of (30) can be stripped of complications of higher dimensions while hopefully retaining many of the physical features of the problem of interest. Notice (30) is valid for even dimensions. It can be straightforwardly generalized to odd dimensions by using the method above demonstrated.

The case of $D = 5$. For the case $D = 5$ the transversal group is $SO(3)$. As is known its bosonic representations labeled by integer j . Let $\Phi_j^\lambda(p)$ be $(2j + 1)$ components of massless spin- j field: $\lambda = -j, -j + 1, \dots, j$. Now introducing one creation operator we collect all $\Phi_j^\lambda(p)$ in one Fock vector

$$|\Phi_j(p, \alpha)\rangle = \sum_{\lambda=-j}^j \Phi_j^\lambda(p) \frac{\alpha^{j-\lambda}}{((j+\lambda)!(j-\lambda)!)^{1/2}} |0\rangle. \quad (31)$$

For the cubic vertex describing the interaction of massless fields Φ_{j_a} (recall that $a = 1, 2, 3$) and having κ th power of derivatives we derive the expression

$$h_{3(0)} = \bar{P}^\kappa \prod_a \beta_a^{-j_a} \prod_a \zeta_a^{2j_a} \left(\sum_a \frac{j_a}{\zeta_a} \right)^{J-\kappa}, \quad (32)$$

where we have introduced the notations

$$\zeta_a = \alpha_a + \frac{1}{\sqrt{2}} \frac{P^3}{\bar{P}}, \quad \bar{P} = \frac{P^1 - i P^2}{\sqrt{2}}. \quad (33a)$$

$$J = \sum_a j_a. \quad (33b)$$

(for the definition of $\mathbf{P}$ see (12)). For $h_{3(0)}$ to be well defined it must be a polynomial of ζ_a . As readily seen from (32) this requirement leads to the following constraints on the allowed values of j_a and κ

$$J \geq \kappa, \quad 2j_a \geq J - \kappa. \quad (34a, b)$$

Now, having the inequalities (34) at hand we are ready to discuss the no-go theorem. The minimal gravitational interaction of massless higher spin fields corresponds to $\kappa = 2$ with one of j_a , say j_1 , taking the value $j_1 = 2$ (graviton), while $j_2, j_3 > 2$ (higher spins). It is easily seen that these values κ, j_a do not satisfy (34b). Since (32) and (34) form a complete list of interactions we conclude immediately that the above-mentioned interaction does not exist at all, a statement of no-go. This statement for the case of four dimensions has been obtained in [8-11]. In these works it was shown that the “naive” gravitational interaction of massless higher spin fields, introduced via covariantization of their free actions in the flat space time turned out to be inconsistent as this led to the loss of the higher spin gauge symmetries. The concrete obstacle was that the variation of the higher spin actions in a curved background contained the gravitational Weyl tensor which could not be compensated by some variation of the gravitation metric. In our approach it is the inequality (34b) that leaves no place for the minimal gravitational interaction of massless higher spin fields. Since the no-go theorem has been rigorously established in four dimensions, our result can be considered as a generalization of no-go to higher $D = 5$ dimension. Thus, (32) and (34) describe all possible Pauli-like interactions of massless higher spin fields. Note that spin- j massless field Φ_j can be associated with totally symmetric covariant tensor field $\Phi^{\mu_1 \dots \mu_j}$. As is well known in four dimension all massless fields can also be described by using totally symmetric tensor fields. However for the case $D \geq 6$, in addition to the totally symmetric massless representations, there exist so-called mixed symmetry ones. The latter can be associated with mixed symmetry covariant tensor fields only. It will be interesting to consider the question about minimal gravitational interaction for such fields. To do that we move to the

$D = 6$ case.

The case of $D = 6$

Now the transversal group is $SO(4)$. Its bosonic representations are labeled by two integers j_γ , $\gamma = 1, 2$. The subscript $\gamma = 1, 2$ refers to the two distinct $SU(2)_\gamma$ subgroups of the $SO(4)$ group, while j_1 and j_2 denote the Casimirs of $SU(2)_1$ and $SU(2)_2$ respectively. The j_γ are expressible in terms of higher weight ($\mathbf{m} = (m_1, m_2)$) of the representation $SO(4)$ group as $j_1 = (m_1 + m_2)/2$, $j_2 = (m_1 - m_2)/2$, where m_1, m_2 satisfy the inequality $m_1 \geq |m_2|$. To describe $(2j_1 + 1)(2j_2 + 1)$ components of massless field we use $\Phi_{j_1, j_2}^{\lambda_1, \lambda_2}(p)$, where $\lambda_\gamma = -j_\gamma, -j_\gamma + 1, \dots, j_\gamma$. Introducing two creation operators α_γ we collect all $\Phi_{j_1, j_2}^{\lambda_1, \lambda_2}(p)$ in one Fock vector

$$|\Phi_{j_1, j_2}(p, \alpha)\rangle = \sum_{\lambda_1 = -j_1}^{j_1} \sum_{\lambda_2 = -j_2}^{j_2} \Phi_{j_1, j_2}^{\lambda_1, \lambda_2}(p) \prod_{\gamma=1,2} \frac{\alpha_\gamma^{j_\gamma - \lambda_\gamma}}{((j_\gamma + \lambda_\gamma)!(j_\gamma - \lambda_\gamma)!)^{1/2}} |0\rangle. \quad (35)$$

The variables α_γ are expressible in terms of the variables η_{12}, ξ_{12} provided by Gauss decomposition (see Eq(5)) as: $\alpha_1 = \eta_{12}$, $\alpha_2 = \xi_{12}$. Note that massless representation Φ_{j_1, j_2} can be associated with mixed symmetry covariant tensor field. This covariant tensor field has the structure of a Young tableaux whose first and second row have length equal to m_1 and $|m_2|$ respectively. The case $m_2 = 0$ (i.e. $j_1 = j_2$) corresponds to totally symmetric fields while $m_2 \neq 0$ (i.e. $j_1 \neq j_2$) to mixed symmetry ones.

Now the $h_{3(0)}$ from (30) can be changed to the form

$$h_{3(0)} = (\mathbf{P}^1)^\kappa \prod_a \beta_a^{-J_a} \prod_{a, \gamma} \zeta_{a\gamma}^{2j_{a\gamma}} \prod_\gamma \left(\sum_a \frac{\beta_a}{\zeta_{a\gamma}} \right)^{J_\gamma - \kappa/2}, \quad (36)$$

where we introduce the notation

$$\zeta_{a1} = \alpha_{a1} - \frac{\mathbf{P}^2}{\mathbf{P}^1}, \quad \zeta_{a2} = \alpha_{a2} - \frac{\mathbf{P}^2}{\mathbf{P}^1}, \quad (37)$$

$$J_a = \sum_\gamma j_{a\gamma}, \quad J_\gamma = \sum_a j_{a\gamma}. \quad (38a, b)$$

Note that $J_a = m_1^{(a)}$. The momentum $\mathbf{P}^I$ is defined in complex coordinates $z^1 = (x^1 + ix^2)/\sqrt{2}$, $z^2 = (x^3 + ix^4)/\sqrt{2}$, $z^1 = (z^1)^*$, $z^2 = (z^2)^*$, $\mathbf{P}^1 = (\mathbf{P}^1)^*$, $\mathbf{P}^2 = (\mathbf{P}^2)^*$. An asterisk

is used to denote the complex conjugate. Again, for $h_{3(0)}$ to be well defined it must be a polynomial of $\zeta_{a\gamma}$. As readily seen from (36) this requirement leads to the following constraints on the allowed values of $j_{a\gamma}$ and κ

$$J_\gamma \geq \frac{1}{2}\kappa, \quad 2j_{a\gamma} \geq J_\gamma - \frac{1}{2}\kappa. \quad (39a, b)$$

Now we are ready to discuss the no-go theorem for the case of totally as well as mixed symmetry massless fields. Recall that the a th massless representation corresponds to totally (mixed) symmetric field if $j_{a1} = j_{a2}$ ($j_{a1} \neq j_{a2}$). The minimal gravitational interaction of massless higher spin fields corresponds to $\kappa = 2$ with one of $j_{a\gamma}$, say $j_{1\gamma}$, taking the values $j_{11} = j_{12} = 2$ (graviton), while $j_{2\gamma}, j_{3\gamma} > 2$ (higher spins). It is easily seen that these values $\kappa, j_{a\gamma}$ do not satisfy (39b). In other words totally symmetric as well as mixed symmetry massless higher spin fields do not have minimal gravitational interaction i.e. (36) and (39) describe Pauli-like interactions of massless higher spin fields for $D = 6$. As to totally symmetric fields, the above-mentioned generalization of no-go theorem to higher dimensions was expected.

We consider the fact that no-go theorem works for the case of mixed-symmetry massless fields are one of the essential new results obtained in this paper. Note that it is hard to put the mixed-symmetry fields in the context of commonly used covariant procedures of deriving the no-go theorem.

Acknowledgments

This investigation is partly supported by the NATO Linkage Grant # 931717. The work of one of the authors (R.R.M) is supported by Grant # MQM000 from International Science Foundation and by the Russian Fund of Fundamental Researches, Grant # 93-02-15541. E.S.F. would like to thank Professor M. Virasoro and the International Centre for Theoretical Physics, Trieste, for hospitality.

REFERENCES

- [1] S. Weinberg, *Phys. Rev.* **135**, (1964) B882, B1049.
- [2] C. Fronsdal, *Phys. Rev.* **D18** (1978) 3624
- [3] F. A. Berends, G. J. Burgers and J. van Dam, *Z. Phys.* **C24** (1984) 247; *Nucl. Phys.* **260** (1985) 295; **B271** (1986) 429.
- [4] S. Deser and Z. Yang, *Class. Quantum Grav.* **7** (1990) 149.
- [5] A. K. H. Bengtsson, I. Bengtsson and L. Brink, *Nucl. Phys.* **B227** (1983) 31.
- [6] A. K. H. Bengtsson, I. Bengtsson and N. Linden, *Class. Quantum Grav.* **4** (1987) 1333.
- [7] E. S. Fradkin and M. A. Vasiliev, *Ann. Phys.* **177** (1987) 63; *Phys. Lett.* **B189** (1987) 89
- [8] E. S. Fradkin and R. R. Metsaev, *Class. Quantum Grav.* **8** (1991) L89.
- [9] C. Aragone and S. Deser, *Phys. Lett.* **B86** (1979) 161.
- [10] S. Christensen and M. Duff, *Nucl. Phys.* **B154** (1979) 301.
- [11] F. A. Berends, J. W. van Holten, P. van Nieuwenhuizen and B. de Wit, *J. Phys.* **A3** (1980) 1643.
- [12] B. de Wit and D. Z. Freedman, *Phys. Rev.* **D21** (1980) 358.
- [13] T. Curtright and P. G. O. Freund, *Nucl. Phys.* **B172** (1980) 413.
- [14] T. Curtright, *Phys. Lett.* **B165** (1985) 304.
- [15] G. R. Black and B. G. Wyborne, *J. Phys. A: Math. Gen.* **16** (1983) 2405.
- [16] R. Delbourgo and P. Jarvis, *J. Phys. A: Math. Gen.* **16** (1983) 1275.
- [17] C. S. Aulakh, I. G. Koh and S. Ouvry, *Phys. Lett.* **B173** (1986) 284.
- [18] J. M. F. Labastida and T. R. Morris, *Phys. Lett.* **B180** (1987) 101.
- [19] J. M. F. Labastida, *Phys. Rev. Lett.* **58** (1987) 531
- [20] R. R. Metsaev, *Class. Quantum Grav.* **10** (1993) L39; *Phys. Lett.* **B309**(1993)39.
- [21] M. B. Green and J. H. Schwarz, *Nucl. Phys.* **B243** (1984) 475.

- [22] M. A. Vasiliev, *Phys. Lett.* **B238** (1990) 305; *Phys. Lett.* **B243** (1990) 378
- [23] C. Fronsdal, *Preprint UCLA/85/TEP/10*. Massless particles, orthosymplectic symmetry and another type of Kaluza-Klein theory.
- [24] R. R. Metsaev, *Mod. Phys. Lett.* **A6** (1991) 2411.
- [25] I. M. Gelfand and M. A. Naimark. Unitare Darstellungen der Klassischen Gruppen, (Academic-Verlag, Berlin 1957).
- [26] W. Siegel, *Introduction to string theory* (World Scientific, Singapore, 1988).
- [27] D. Zhelobenko, *Uspekhi Mat. Nauk.* **17** (1962) 27.
- [28] W. H. Klink and T. Ton-That, *Acta. Appl. Math.* **9** (1987) 137.
- [29] R. R. Metsaev, *Mod. Phys. Lett.* **A8** (1993) 2413.