Brunel
THE UNIVERSITY OF WEST LONDON

BRU/PH/213
September 1996

A Search for the $\xi(2235)$ in $\bar{p}p \rightarrow \pi^- \pi^+$

sw9646

A. Hasan¹ and D. V. Bugg²

(1) Department of Physics, Brunel University,
West London, Uxbridge UB8 3PH, UK
Email: Adil.Hasan@Brunel.ac.uk

(2) Department of Physics, Queen Mary & Westfield College,
London, E1 4NS
Email: Bugg@v2.rl.ac.uk

(Submitted to Phys. Lett. B)

Abstract

We present results of a search for the narrow $\xi(2235)$ in the $\bar{p}p \rightarrow \pi^-\pi^+$ channel. The search is carried out using an amplitude analysis of all the high statistics data currently available in the pion channel. No convincing evidence for a narrow state is observed.

The BES collaboration [1] has reported evidence for a narrow $\xi(2235)$ with a full width, Γ , in the range 15 to 20 MeV in the radiative J/ψ decay modes $\gamma\pi^+\pi^-$, $\gamma K\bar{K}$ and $\gamma\bar{p}p$. They find product branching ratios:

$$B(J/\psi \rightarrow \gamma\xi) \cdot B(\xi \rightarrow \pi^+\pi^-) = (5.6_{-1.6}^{+1.8} \pm 1.4) \times 10^{-5}, \quad (1)$$

$$B(J/\psi \rightarrow \gamma\xi) \cdot B(\xi \rightarrow K^+K^-) = (3.3_{-1.3}^{+1.6} \pm 1.1) \times 10^{-5}, \quad (2)$$

$$B(J/\psi \rightarrow \gamma\xi) \cdot B(\xi \rightarrow K_S K_S) = (2.7_{-0.9}^{+1.1} \pm 1.0) \times 10^{-5}, \quad (3)$$

$$B(J/\psi \rightarrow \gamma\xi) \cdot B(\xi \rightarrow \bar{p}p) = (1.5_{-0.5}^{+0.6} \pm 0.5) \times 10^{-5}. \quad (4)$$

A narrow ξ state around 2235 MeV has remained controversial for many years. Evidence for a narrow structure with a mass of 2230 MeV was first claimed by the Mark III collaboration [2] in J/ψ decays to γK^+K^- and $\gamma K_S^0 K_S^0$. The DM2 group [3] also studied the same J/ψ decay modes. However, they observed no evidence for narrow structure in either mode. They set upper limits at 95% confidence on the product branching ratios of 2.3×10^{-5} for Eq. (2) and 1.6×10^{-5} for Eq. (3). Instead, they reported a broad resonance with a mass of 2197 MeV and width 201 MeV. We refer to the Particle Data Group [4] for a summary of subsequent evidence concerning a narrow ξ state.

Three experiments have searched specifically for narrow structure in $\bar{p}p \rightarrow K\bar{K}$. The first measurement, by Bardin et al. [5], scanned the mass range 2220 to 2240 MeV in 0.6 MeV steps. The measurement depended on the ratio of cross sections $K\bar{K}/\pi^+\pi^-$ integrated over the centre of mass angular range 40° to 140° . No significant evidence for narrow structure was observed. For a resonance width of 30 MeV, a 3σ limit on the product branching ratio $BR2 = B(\xi \rightarrow \bar{p}p) \cdot B(\xi \rightarrow K^+K^-)$ was determined to be 5.4×10^{-4} for $J^{PC} = 0^{++}$ and 2.1×10^{-4} for $J^{PC} = 2^{++}$. The experimental group states that the sensitivity of these results varies roughly as $1/\sqrt{\Gamma}$, where Γ is the width of the resonance.

A similar measurement by Sculli et al. [6] with a centre of mass angular acceptance from 60° to 120° also measured the ratio of integrated cross sections for K^+K^- and $\pi^+\pi^-$ in the lab momentum range from 1290 MeV/c (~ 2180 MeV in mass) to 1550 MeV/c (~ 2268 MeV in mass) in 10 MeV/c momentum steps. Again, no evidence for narrow structure was observed. The upper limit on $BR2$ was 12×10^{-4} for 0^{++} and 3.8×10^{-4} for 2^{++} .

Barnes et al. [7] have measured the integrated cross section for $\bar{p}p \rightarrow K_S K_S$ at 18 masses from 2210 to 2260 MeV. Their mass resolution was ± 0.2 MeV. For a state with mass 2231 MeV and width 10 MeV they determined a 3σ limit on the product branching ratio to be $B(\xi \rightarrow \bar{p}p) \cdot B(\xi \rightarrow K^+K^-) = 2 \times B(\xi \rightarrow \bar{p}p) \cdot B(\xi \rightarrow K_S K_S) < 3.9 \times 10^{-4}/(2J+1)$. This high sensitivity arises because the background $K_S K_S$ signal is much smaller ($\sim 2\mu b$) than for K^+K^- ($60\mu b$) or $\pi^-\pi^+$ ($180\mu b$).

In the light of the new BES results, we examine what conclusions can be drawn from $\bar{p}p \rightarrow \pi^-\pi^+$ differential cross sections and polarisations measured by the PS172 experiment at LEAR [8]. The experiment has the advantage of covering the whole angular range from 0 to 180° . Unfortunately, the data are available only in steps of 6 to 12 MeV mass steps in the $\xi(2235)$ mass range (see Table 1). An extremely narrow resonance could hide in the gaps. We shall draw conclusions only for a resonance width comparable with the mass spacing of the data, $\Gamma \geq 5$ MeV. The experiment was carried out using a 3 cm long polarised target and the $\bar{p}$ beam suffered a momentum loss of 5.8 MeV/c (2.03 MeV in mass) at the ξ . The effect of this spread is almost negligible for the results presented here. However, a resonance with $\Gamma < 3$ MeV would be folded with the 2 MeV mass loss down the target.

A strong advantage of this channel is that an amplitude analysis exists spanning a wide mass range [9]. This analysis identifies the magnitudes and phases of broad background amplitudes. Any narrow resonance would interfere with these broad background amplitudes and gives rise to delicate perturbations in $d\sigma/d\Omega$ and A_{0N} . This gives information on the spin. Details of the amplitude analysis can be found in Ref. [9] and here we outline only the modification made for the present search.

A narrow Breit-Wigner resonance is added to our earlier solution with freely variable magnitude and phase. The mass of the narrow resonance is stepped by 5 MeV from 2225 to 2245 MeV using widths of 15, 10 or 5 MeV. The magnitude and phase of this resonance and all broad components are re-optimised. To obtain upper limits on the product branching ratio, the strength of the narrow state is increased until the χ^2 of the fit increases by 3σ . The relation between the peak cross section and the product branching ratio for the narrow resonance is

$$\sigma_{peak} = \pi(\hbar c/k)^2(2J+1)B(\bar{p}p)B(\pi^-\pi^+), \quad (5)$$

where B is the ratio of the channel width to the total width of the resonance, and k is the $\bar{p}$ centre of mass momentum. Note that one expects branching ratios to be proportional to phase space, i.e. centre of mass momenta, as well as to the squares of matrix elements for the decay. The available momenta at a mass of 2235 MeV are 1109 MeV/c for the pion, 1000 MeV/c for K_S and 607 MeV/c for the nucleon.

We shall show results in two forms. Firstly, Fig. 1 shows the peak cross sections, from the fit, for narrow resonances of $J = 0, 1, 2$ or 4 , assuming a width of 15 MeV. The observation of $\xi \rightarrow K_S K_S$ by BES, if confirmed, would rule out $J = 1$. However, limits for this value of J are particularly low and seem worth quoting. Those for $J = 3$ are similar to $J = 4$. Within the fit errors, there is no signal for $J = 0, 1, 2$ and 4 . For $J = 2$, there is a hint of something, but it is not centred at the BES mass of $2235 \pm 4 \pm 6$ MeV. We remark that our earlier analysis included a spin 2 resonance with a mass of 2226 MeV, a width of 226 MeV and a peak cross section of $28 \mu b$. Likewise, for $J = 4$ there is a hint in Fig. 1 of a 2σ effect. However, there could be some cross-talk between broad resonances and a possible narrow signal, and we do not consider that the evidence for a narrow signal in Fig. 1 is significant.

Secondly, Table 1 shows the products of branching ratios $B(\bar{p}p \rightarrow \xi) \cdot B(\xi \rightarrow \pi^+\pi^-)$ which give rise to 3σ effects for $\Gamma = 15, 10$ and 5 MeV. The BES data indicate a width in the range 15 to 20 MeV, but with statistical and systematic errors of the same general magnitude. For $\Gamma = 15$ MeV and a mass 2225-2240 MeV, our limits are in the range

$$B(\bar{p}p \rightarrow \xi) \cdot B(\xi \rightarrow \pi^+\pi^-) \leq (0.15 - 2.8) \times 10^{-4} \quad (6)$$

for $J = 0$ to 4 . They show a scatter which is compatible with statistical fluctuations in the data and provide new limits on the $\pi\pi$ channel. The 3σ limits for the $\Gamma = 5$ MeV wide state are less restrictive than those for $\Gamma = 15$ and 10 MeV because this width is close to the mass spacing of data points.

For $\Gamma = 15$ MeV, our sensitivity is comparable in the $\pi^-\pi^+$ channel with that of Barnes et al. in $K_S K_S$. This is a useful result, since the BES group claims that their $\pi^-\pi^+$ signal is a factor 2 larger than for $K_S K_S$.

Any interpretation of these results is a question of juggling the limits imposed by different processes. For purposes of illustration, we shall consider the implications of Eq. (6) for $\Gamma = 15$ MeV and $M = 2235$ MeV. Initially, we take the debatable assumption that decay widths

of $\xi(2235)$ are proportional to phase space in the $\bar{p}p$, $\pi\pi$ and $K\bar{K}$ channels, i.e. to decay momenta quoted above. The BES results of Eqs. (1)–(4) are roughly consistent with this within the sizable errors. Then the upper limit of Eq. (6) would give an upper limit on $B(\xi \rightarrow \pi^-\pi^+)$ of 2.3×10^{-2} at the 3σ level. From Eq. (1), this implies that

$$B(J/\Psi \rightarrow \gamma\xi) \geq 2.5 \times 10^{-3}. \quad (7)$$

This inclusive rate seems very high.

The Crystal Ball experiment has measured the inclusive gamma spectrum, Fig. 59 of Ref. [10], and we now deduce limits from it on $B(J/\Psi \rightarrow \gamma\xi)$. There is no evidence for a peak at 2235 MeV. The problem in interpreting this result quantitatively is knowing the normalisation of the spectrum. It may well contain contamination from photons following π^0 and η decays. We therefore normalise to the largest signal in the spectrum, due to the η' . This has a well known branching ratio $B(J/\Psi \rightarrow \gamma\eta')$ of $4.31 \pm 0.30 \times 10^{-3}$ and the peak contains ~ 2000 events. The authors state that the photon resolution is constant along the horizontal axis of their figure. This implies a mass resolution of $\sim \pm 27$ MeV at the ξ . A 3σ signal with this full width (~ 55 MeV) would correspond to ~ 150 events due to $J/\Psi \rightarrow \gamma\xi$, and an integrated signal, independent of the actual width of $\xi(2235)$ (as for the narrow η'), of $B(J/\Psi \rightarrow \gamma\xi) \leq 0.32 \times 10^{-3}$. We are able to check this result from the second largest signal in the γ spectrum, due to the $\iota(1440)$. The branching ratio for radiative production of $\iota(1440)$ and subsequent decay to $K\bar{K}\pi$ is 0.9×10^{-3} . The Crystal Barrel group [11] has determined the branching ratio of the iota to be $B(\iota \rightarrow K\bar{K}\pi)/B(\iota \rightarrow \eta\pi\pi) = 0.61 \pm 0.19$. These are the only likely decay modes of the ι in the narrow peak, although the $\rho\rho$ channel may well contribute at higher masses [12]. Then the total radiative decay rate for production of the ι is about 2.4×10^{-3} . This gives precisely the same normalisation as the η' signal, but with a larger error of $\pm 25\%$.

There could be some systematic uncertainty in these limits arising from variation of γ detection with energy. However, the discrepancy between the limits and Eq. (7) is nearly a factor 10. One must also remember that all limits we quote are stretching numerical values to 3σ . There seem to be only two escape routes from this dilemma. One possibility is that the ξ has a width much less than 15 MeV; results of Table 2, taken with BES results, firmly establish $\Gamma < 5$ MeV. But then the question arises: what selection rule makes it so narrow? One possible explanation of BES results is constructive interference between a cusp at the $\Lambda\bar{\Lambda}$ threshold and a broad resonance around 2200–2225 MeV such as DM2 and our own amplitude analysis have found. Alternatively, the link in the arithmetic which appears statistically weakest is the BES branching ratio of ξ to $\bar{p}p$, Eq. (4). If the ξ does not couple to $\bar{p}p$, the negative results from $\bar{p}p \rightarrow \pi^-\pi^+$ and $K\bar{K}$ are immediately explicable.

1 Figure Caption

Fig. 1. Peak cross-sections derived from the fit for $J = 0, 1, 2$ and 4 with $\Gamma(\xi) = 15$ MeV.

References

- [1] J.Z. Bai et al., Phys. Rev Lett. 76 (1996) 3502.
- [2] R.M. Baltrusaitis et al., Phys. Rev. Lett. 56 (1986) 107
- [3] J.E. Augustin et al., Phys. Rev. Lett. 60 (1988) 2238
- [4] Particle Data Group, Phys. Rev. 54 (1996) 1
- [5] G. Bardin et al., Phys. Lett. B195 (1987) 292
- [6] J. Sculli et al., Phys. Rev. Lett. 58 (1987) 1715
- [7] P.D. Barnes et al., Phys. Lett. B309 (1993) 469
- [8] A. Hasan et al., Nucl. Phys. B378 (1992) 3
- [9] A. Hasan and D. V. Bugg, Phys Lett B334 (1994) 215
- [10] L. Köpke and N.Wermes, Phys. Reports 174 (1989) 67.
- [11] C.Amsler et al., Phys. Lett. B358 (1995) 389.
- [12] D.V. Bugg et al., Phys. Lett. B353 (1995) 378.

Lab momentum (MeV/c)	Mass (MeV)
1351	2201.7
1400	2218.9
1416	2224.5
1449	2236.1
1467	2242.4
1500	2254.1
1550	2271.7

Table 1: Lab momenta of PS172 data versus ξ mass.

Γ (MeV)	Mass (MeV)	J = 0	J = 2	J = 4	J = 1
15	2220	0.6	0.7	1.1	0.5
	2225	1.1	1.6	2.5	0.2
	2230	0.3	2.6	1.8	0.15
	2235	0.7	2.8	1.7	1.1
	2240	0.8	2.2	1.6	1.9
	2245	0.8	2.5	1.4	1.5
10	2225	3.4	3.1	4.2	0.7
	2230	1.2	3.9	3.8	0.5
	2235	1.3	3.6	3.8	1.7
5	2225	6.5	7.3	8.2	2.3
	2230	6.6	6.9	9.7	4.4
	2235	4.2	7.9	8.3	2.7

Table 2: 3σ limits on $B(\bar{p}p \rightarrow \xi) \cdot B(\xi \rightarrow \pi^-\pi^+)$, all in units of 10^{-4} , as a function of the mass and width, Γ , of ξ .