ОБЪЕДИНЕННЫЙ
ИНСТИТУТ
ЯДЕРНЫХ
ИССЛЕДОВАНИЙ

Дубна

AC CERN LIBRARIES, GENEVA

SCAN-9703025

E2-96-376

Swg 711
V.V.Nesterenko, I.G.Pirozhenko*

ON THE CALCULATION
OF THE INTERQUARK POTENTIAL GENERATED
BY A STRING WITH MASSIVE ENDS

Submitted to «Physical Review D»

*Permanent address: Petrozavodsk State University, Petrozavodsk, 185640,
Russia

1996

1. Interquark potential generated by the Nambu-Goto string with point-like masses at ends was at first calculated in [1]. A standard method [3,4] based on variational estimation of the string functional integral in $D \rightarrow \infty$ limit was applied (D is the space-time dimension). However, the variational equations obtained in [1] were not solved exactly. Some approximation was used there. The aim of the present note is to justify the applicability of such approximation.

2. Let us remind briefly the derivation of the variational equations in the considered problem. The starting point is the well-known formula

$$\exp[-\beta V(R)] = \int [Du] \exp\{-S^\beta[u]\}, \quad \beta \rightarrow \infty, \quad (1)$$

relating interquark potential $V(R)$ with the string functional integral. The Euclidean action of the string with masses at its ends is given by the formula

$$\begin{aligned} S^\beta[u] = & M_0^2 \int_0^\beta dt \int_0^R dr \sqrt{\det(\delta_{ij} + \partial_i \mathbf{u} \partial_j \mathbf{u})} \\ & + \sum_{a=1}^2 m_a \int_0^\beta \sqrt{1 + \dot{\mathbf{u}}^2(t, r)} \quad i, j = 0, 1, \end{aligned} \quad (2)$$

where M_0^2 is the string tension, $\beta = T^{-1}$ is the inverse temperature, and string coordinates $\mathbf{u}(t, r)$ satisfy the periodicity condition

$$\mathbf{u}(0, r) = \mathbf{u}(\beta, r). \quad (3)$$

In the present case $1/(D-2)$ expansion is obtained by means of the following procedure [3]. The composite fields σ_{ij} for $\partial_i \mathbf{u} \partial_j \mathbf{u}$ are introduced, and the constraint $\delta_{ij} = \partial_i \mathbf{u} \partial_j \mathbf{u}$ is taken into account through the lagrange multipliers α^{ij} , $i, j = 0, 1$. To this end, the following presentation of a unit is inserted into the string functional integral (1)

$$1 = \int [D\sigma] \delta(\sigma_{ij} - \partial_i \mathbf{u} \partial_j \mathbf{u}).$$

After using the integral representation of δ -function we obtain

$$\exp[-\beta V(R)] = \int [D\mathbf{u}][D\alpha][D\sigma] \exp\{-S^\beta[\mathbf{u}, \alpha, \sigma]\}, \quad (4)$$

where

$$\begin{aligned} S^\beta[\mathbf{u}, \alpha, \sigma_i] = & \frac{M_0^2}{2} \int_0^\beta dt \int_0^R dr \left[2\sqrt{\det(\delta_{ij} + \sigma_{ij})} + \alpha^{ij}(\partial_i \mathbf{u} \partial_j \mathbf{u} - \sigma_{ij}) \right] \\ & + \sum_{a=1}^2 m_a \int_0^\beta dt \sqrt{1 + \dot{\mathbf{u}}^2(t, r_a)} \\ & i, j = 0, 1, \quad r_1 = 0, \quad r_2 = R. \end{aligned} \quad (5)$$

Functional integral over $\mathbf{u}$ can be done exactly as the action is quadratic in string coordinates. Functional integrals over α and σ are evaluated by variational method and in $(D-2) \rightarrow \infty$ limit they are equal to the integrand value at the stationary point of the action (5). The functional variables α_{ij} and σ_{ij} are diagonal matrices independent of t and r , $\alpha^{ij} = \delta^{ij} \alpha^j$, $\sigma_{ij} = \delta_{ij} \sigma_j$.

The action (5) leads to the following equations of motion

$$\Delta_\alpha = \alpha^0 \ddot{\mathbf{u}} + \alpha^1 \mathbf{u}'' = 0 \quad (6)$$

and boundary conditions

$$\frac{m_1}{\sqrt{1 + \sigma_0}} \ddot{\mathbf{u}} = -M_0^2 \alpha^1 \dot{\mathbf{u}} \quad r = 0, \quad (7)$$

$$\frac{m_2}{\sqrt{1 + \sigma_0}} \ddot{\mathbf{u}} = M_0^2 \alpha^1 \dot{\mathbf{u}} \quad r = R. \quad (8)$$

Integrating by parts the second term in (5) and taking into consideration the boundary conditions (7),(8) we arrive at the effective action

$$S^\beta[\mathbf{u}, \alpha, \sigma] = \frac{M_0^2}{2} \int_0^\beta dt \int_0^R dr \mathbf{u}(-\Delta_a) \mathbf{u}$$

$$+ M^2 \beta R \left[\sqrt{(1 + \sigma_0)(1 + \sigma_1)} - \frac{1}{2}(\alpha^0 \sigma_0 + \alpha^1 \sigma_1) \right], \quad (9)$$

which is quadratic in string coordinates. This action does not include exactly the string ends contributions. The latter are taken into account through the definition of the operator $(-\Delta_\alpha)$. Therefore after functional integration over $\mathbf{u}$ the term

$$\frac{D-2}{2} \text{Tr} \ln(-\Delta_\alpha) = \frac{1}{2} \sum_{k=1}^{\infty} \omega_k, \quad (10)$$

appears in the effective action [1]. Here ω_k are solutions of the frequency equation

$$\tan(\omega R) = \frac{2mM_0^2 \omega \alpha^0 \sqrt{1 + \sigma_0}}{m^2 \omega^2 - M_0^2 \alpha^0 \sqrt{1 - \sigma_0}}, \quad (11)$$

following from (6)-(8). Taking into account (10) we obtain the effective string action

$$S^\beta = M_0^2 \beta R \left\{ \sqrt{(1 + \sigma_0)(1 + \sigma_1)} - \frac{1}{2}(\alpha^0 \sigma_0 + \alpha^1 \sigma_1) \right\} \\ + \beta(D-2) \sqrt{\frac{\alpha^1}{\alpha^0}} E_c(\alpha^0, \sigma_0), \quad (12)$$

where

$$E_c(\alpha^0, \sigma_0) = \frac{1}{2} \sum_k \omega_k = \\ \frac{1}{2\pi} \int_0^\infty dy \ln \left\{ 1 - e^{-2Ry} \left(\frac{my - M_0^2 \alpha^0 \sqrt{1 + \sigma}}{my + M_0^2 \alpha^0 \sqrt{1 + \sigma_0}} \right) \right\} \quad (13)$$

is renormalised Casimir energy [1,2], depending on variational parameters α^0, σ_0

The stationary point of this action is defined from the variational equations

$$\alpha^0 = \sqrt{\frac{1 + \sigma_1}{1 + \sigma_0}} + \frac{D-2}{M_0^2 R} \sqrt{\frac{\alpha^1}{\alpha^0}} \frac{\partial E_c(\alpha_0, \sigma_0)}{\partial \sigma_0}, \quad (14)$$

$$\alpha^1 = \sqrt{\frac{1 + \sigma_0}{1 + \sigma_1}}, \quad (15)$$

$$\sigma_0 = -\frac{D-2}{M_0^2 R \alpha^0} \sqrt{\frac{\alpha^1}{\alpha^0}} E_c(\alpha^0, \sigma_0) + 2 \frac{D-2}{M_0^2 R} \sqrt{\frac{\alpha^1}{\alpha^0}} \frac{\partial E_c(\alpha^0, \sigma_0)}{\partial \alpha^0}, \quad (16)$$

$$\sigma_1 = \frac{D-2}{M_0^2 R} \frac{E_c(\alpha^0, \sigma_0)}{\sqrt{\alpha^0 \alpha^1}}. \quad (17)$$

Solution of the nonlinear equations (14)-(17) should be substituted in (12). This enables one to get the interquark potential in the string model under consideration.

3. When solving the variational equations (14)-(17) in paper [1], the following approximation has been used. The Casimir energy $E_c(\alpha^0, \sigma_0)$, entering the right-hand sides of Eqs. (14)-(17) was calculated at fixed values of variational parameters α^0 and σ_0 : $\alpha^0 = 1$ and $\sigma_0 = 0$. In definite sense this corresponds to the nonrelativistic approximation for quarks dynamics (1). In this approximation the derivatives of the Casimir energy over variational parameters α^0 and σ_0 are obviously equal to zero. As a result, the solution of the system (14)-(17) is

$$\alpha^0 = \sqrt{1 - 2\lambda}, \quad \alpha^1 = \frac{1}{\sqrt{1 - 2\lambda}}, \quad \sigma_0 = \frac{\lambda}{1 - 2\lambda}, \quad \sigma_1 = -\lambda, \quad (18)$$

where $\lambda = -(D-2) E_c / (M_0^2 R)$ [3].

In general case the following expressions

$$\begin{aligned} \frac{\partial E_c(\alpha^0, \sigma_0)}{\partial \alpha^0} &= \frac{2M_0^2 \sqrt{1 + \sigma_0}}{\pi m} \int_0^\infty dy \frac{(y - \eta) e^{-2y}}{(y - \eta)^3 - e^{-2y}(y - \eta)^2(y + \eta)}, \\ \frac{\partial E_c(\alpha^0, \sigma_0)}{\partial \sigma_0} &= \frac{\alpha^0 M_0^2}{\pi m \sqrt{1 + \sigma_0}} \int_0^\infty dy \frac{(y - \eta) e^{-2y}}{(y - \eta)^3 - e^{-2y}(y - \eta)^2(y + \eta)}, \end{aligned} \quad (19)$$

should be substituted in the right-hand sides of (14)-(17). Here $\eta = \alpha^0 \sqrt{1 + \sigma_0} M_0^2 R / m$.

Fig.1. Interquark potential obtained by means of the iterative solution of variational equations system (14)-(17). Curve's numbers are equal to the number of iterations. The curve 0 corresponds to zero approximation ($\alpha^0 = 1, \sigma_0 = 0$). The following dimensionless parameters are used: $\rho = M_0 R$, $\mu = m/M_0$.

Now we are going to construct an iterative solution of the exact equations (14)–(17). To this end we shall treat the terms $\partial E_c/\partial\sigma_0$ and $\partial E_c/\partial\alpha_0$ there as small corrections. To start with, we take the solution (18) as a zero approximation. Substituting it into the right-hand sides of (14)–(17), we obtain some values of α^0 , α^1 , σ_0 , σ_1 . We put them into the right-hand sides of (14)–(18) and so on. At each step we shall calculate the effective string action (12) obtaining the corresponding interquark potential and comparing it with the zero approximation.

The calculation results for different quark masses are presented in Fig.1. At large interquark distances the potential rises linearly and corrections to zero approximation are small. For heavy quarks ($\mu = m/M_0 = 100$) the iterative solution coincides with zero approximation at all range of available distances $R > R_c$. For average masses ($\mu \sim 1 - 10$) and not far from $R = R_c$ the calculated plots oscillate near zero approximation with increasing number of iteration. In the limit $R \rightarrow R_c$ the corrections are not small and the proposed iterative procedure may give no convergent result. For light quarks $\mu \sim 0.01$ iterative solution insignificantly displaces as a whole to smaller distances in comparison with zero approximation. It is important to note that consideration of Casimir energy dependence on α^0, σ_0 does not lead to significant modification of the critical radius R_c .

Taking all this into account, we infer that approximation (18) works sufficiently well for any quark mass and practically for all distances between quarks.

References

- [1] G. Lambiase, V.V. Nesterenko, Phys. Rev. **D54**, 1-12, (1996)
- [2] H. Kleinert, G. Lambiase, V.V. Nesterenko, Phys. Lett. B, (1996)
- [3] O. Alvarez, Phys. Rev. **D24**, 440, (1981).
- [4] M. Lüscher, K. Symanzik and P. Weisz, Nucl. Phys. **B173**, 365 (1980).

Received by Publishing Department
on October 14, 1996.

**The Publishing Department
of the Joint Institute for Nuclear Research
offers you to acquire the following books:**

Index	Title
E7-93-274	Proceedings of the International School-Seminar on Heavy Ion Physics. Dubna, 1993 (2 volumes, 498 and 590 pp., stiffened cover, in English)
Д14-93-325	Proceedings of the Workshop «Activation Analysis in Environment Protection». Dubna, 1992 (in Russian and English)
Д1,9,13-93-459	Proceedings of the II Workshop on JINR Tau-Charm Factory. Dubna, 1993 (266 p. in Russian and English)
E7-94-19	Proceedings of the International Conference on Dynamical Aspects of Nuclear Fission. Smolenice, 1993 (318 p. in English)
94-55	Proceedings of the International Bogoliubov Memorial Meeting. Dubna, 1993 (216 p. in Russian and English)
E2-94-95	Proceedings of the International Symposium «Dubna Deuteron-93». Dubna, 1993 (390 p. in English)
Д14-94-269	Abstracts of the XXX Conference on Low Temperature Physics. Dubna, 1994 (2 volumes in Russian and English)
E7-94-270	Proceedings of the Workshop on Physical Experiments and First Results on Heavy Ion Storage and Cooler Rings. Smolenice, 1992 (324 p. in English)
E13-94-290	Abstracts of the 17th International Conference «Nuclear Tracks in Solids». Dubna, 1994 (296 p. in English)
E2-94-347	International Workshop «Symmetry Methods in Physics». In Memory of Professor Ya.A.Smorodinsky. Dubna, 1993 (2 volumes, 602 p. in English)
E4-94-370	Proceedings of the IV International Conference on Selected Topics in Nuclear Structure. Dubna, 1994 (412 p. in English)
E4-94-386	Proceedings of the VI Trilateral German-Russian-Ukrainian Seminar on High-Temperature Superconductivity. Dubna, 1994 (340 p. in English)
D2-94-390	D.I.Blokhintsev. Proceedings of the Seminars. Dedicated to the 85th Anniversary of the Birthday of D.I.Blokhintsev. Dubna, 1995 (271 p. in Russian and English)

Index	Title
E3-94-419	Proceedings of the II International Seminar on Neutron-Nucleus Interactions (ISINN-2) «Neutron Spectroscopy, Nuclear Structure, Related Topics». Dubna, 1994 (363 p. in English)
D13-94-491	Proceedings of the XVI International Symposium on Nuclear Electronics and VI International School on Automation and Computing in Nuclear Physics and Astrophysics. Varna, 1994 (246 p. in Russian and English)
D13,14-95-49	Proceedings of the International Symposium on Muon and Pion Interactions with Matter. Dubna, 1994 (304 p. in Russian and English)
D3-95-169	Proceedings of the International Seminar ADVANCED PULSED NEUTRON SOURCES: Physics of/at ADVANCED Pulsed Neutron Sources. PANS-II. Dubna, 1994 (336 p. in Russian and English)
D1-95-305	Proceedings of the XVII Workshop on Neutrino Detector IHEP—JINR. Dubna, 1995 (178 p. in Russian and English)
E3-95-307	Proceedings of the III International Seminar on Interaction of Neutrons with Nuclei. Neutron Spectroscopy, Nuclear Structure, Related Topics. Dubna, 1995 (356 p. in English)
D4-95-308	V.G.Soloviev. Frontiers in Nuclear Physics. Dedicated to the 70th Anniversary of the Birthday of V.G.Soloviev. Dubna, 1995 (238 p. in Russian and English)
D3,14-95-323	VII School on Neutron Physics. Lectures. Vol.1. Dubna, 1995 (356 p. in Russian and English)
E10,11-95-387	Proceedings of the ESONE International Conference RTD'94 on REAL TIME DATA 1994 with Emphasis on Distributed Front-End Processing. Dubna, 1994 (358 p. in English)

Please apply to the Publishing Department of the Joint Institute for Nuclear Research for extra information. Our address is:

Publishing Department
Joint Institute for Nuclear Research
Dubna, Moscow Region
141980 Russia
E-mail: publish@pds.jinr.dubna.su.

Нестеренко В.В., Пирожено И.Г.

E2-96-376

К расчету межкваркового потенциала, генерируемого струной
с массами на концах

Предложена итерационная процедура решения системы вариационных уравнений, определяющих стационарную точку действия струны с массами на концах. Полученное итерационное решение используется для вычисления струнного функционального интеграла и оценки поправок к межкварковому потенциалу, генерируемому струной. Расчеты показали, что высшие поправки малы. Нулевое приближение достаточно хорошо воспроизводит межкварковый потенциал, генерируемый струной с массами на концах.

Работа выполнена в Лаборатории теоретической физики им. Н.Н. Боголюбова ОИЯИ.

Препринт Объединенного института ядерных исследований. Дубна, 1996

Nesterenko V.V., Pirozhenko I.G.

E2-96-376

On the Calculation of the Interquark Potential Generated
by a String with Massive Ends

It is proposed to solve the variational equations, defining the stationary point of the effective action of a string with massive ends, by means of iterative procedure. The obtained solution is used to evaluate the string functional integral. After that the corrections to interquark potential generated by a string with masses at the ends are calculated. The calculations show that the higher corrections prove to be small and zero approximation fairly well reproduces the interquark potential generated by the Nambu-Goto string with massive ends.

The investigation has been performed at the Bogoliubov Laboratory of Theoretical Physics, JINR.

Preprint of the Joint Institute for Nuclear Research. Dubna, 1996

Редактор Э.В.Ивашкевич. Макет Р.Д.Фоминой

Подписано в печать 30.10.96

Формат 60×90/16. Офсетная печать. Уч.-изд.листов 0,99

Тираж 460. Заказ 49469. Цена 1188 р.

Издательский отдел Объединенного института ядерных исследований
Дубна Московской области