CERN-TH/98-2
hep-ph/9801269

Heavy-Quark Effective Theory and Weak Matrix Elements

Matthias Neubert

Theory Division, CERN, CH-1211 Geneva 23, Switzerland

Abstract:

Recent developments in the theory of weak decays of heavy flavours are reviewed. Applications to exclusive semileptonic B decays, the semileptonic branching ratio and charm counting, beauty lifetimes, and hadronic B decays are discussed.

*Invited talk presented at the
International Europhysics Conference on High Energy Physics
Jerusalem, Israel, 19–26 August 1997*

CERN-TH/98-2
January 1998

19: Heavy-Quark Effective Theory and Weak Matrix Elements

Matthias Neubert (Matthias.Neubert@cern.ch)

Theory Division, CERN, CH-1211 Geneva 23, Switzerland

Abstract. Recent developments in the theory of weak decays of heavy flavours are reviewed. Applications to exclusive semileptonic B decays, the semileptonic branching ratio and charm counting, beauty lifetimes, and hadronic B decays are discussed.

1 Introduction and theoretical concepts

To discuss even the most significant recent theoretical developments in heavy-flavour physics in a single talk is a difficult task. Fortunately, this field is blooming and will continue to be of great importance in view of several new experimental facilities (B factories) to start operating in the near future. Below, I will review the latest theoretical developments in this field and discuss the most important phenomenological applications. They concern semileptonic B decays and the measurements of the CKM parameters $|V_{cb}|$ and $|V_{ub}|$, the semileptonic branching ratio and charm yield in inclusive B decays, the lifetimes of beauty mesons and baryons, and hadronic B decays, including the rare decays into two light mesons. I will start with an introduction to the main theoretical concepts used in the analysis of these processes.

The properties of hadrons containing a single heavy quark Q are characterized by the large separation of two length scales: the Compton wave length $1/m_Q$ of the heavy quark is much smaller than the typical size $1/\Lambda_{\text{QCD}}$ of hadronic bound states in QCD (see the left plot in Fig. 1). In the limit $m_Q \rightarrow \infty$, the configuration of the light degrees of freedom in the hadron becomes independent of the spin and flavour of the heavy quark. In that limit there is a global $\text{SU}(2n_h)$ spin-flavour symmetry of the strong interactions, where n_h is the number of heavy-quark flavours [1]. This symmetry helps in understanding the spectroscopy and decays of heavy hadrons from first principles. It does not allow us to solve QCD, but to parametrize the strong-interaction effects of heavy-quark systems by a minimal number of reduced matrix elements, thus giving rise to nontrivial relations between observables. In particular, all form factors for the weak $\bar{B} \rightarrow D^{(*)} \ell \bar{\nu}$ transitions are proportional to a universal function $\xi(w)$, where $w = v_B \cdot v_{D^{(*)}}$ is the product of the meson velocities. At the zero-recoil point $w = 1$, corresponding to $v_B = v_{D^{(*)}}$, this function is normalized to unity: $\xi(1) = 1$. The symmetry-breaking corrections to the heavy-quark limit can be organized in

an expansion in powers of the small parameters $\alpha_s(m_Q)$ and Λ_{QCD}/m_Q . A convenient way to do this is provided by the heavy-quark effective theory (HQET), whose purpose is to separate the short- and long-distance physics associated with the two length scales, making all dependence on the large mass scale m_Q explicit. This allows us to derive scaling laws relating different observables to each other. The philosophy behind the HQET is illustrated in the right plot in Fig. 1.

Fig. 1. Length scales of a heavy hadron, and construction of the HQET

The effective Lagrangian of the HQET is

$$\mathcal{L}_{\text{eff}} = \bar{h} i v \cdot D h + \frac{1}{2m_Q} \bar{h} (iD)^2 h + \frac{C(m_Q)}{4m_Q} \bar{h} \sigma_{\mu\nu} G^{\mu\nu} h + \dots,$$

where v is the velocity of the hadron containing the heavy quark, and h is the heavy-quark field. The Lagrangian incorporates the spin-flavour symmetry to leading order in $1/m_Q$. The only nontrivial short-distance coefficient has recently been calculated to next-to-leading and even next-to-next-to-leading order [2]–[4]. The result is

$$C(m_Q) = [\alpha_s(m_Q)]^{9/25} \left\{ 1 + 0.672 \alpha_s(m_Q) + (1.33 \pm 0.04) \alpha_s^2(m_Q) + \dots \right\}.$$

The long-distance physics is encoded in the hadronic matrix elements of HQET operators, e.g.

$$\mu_\pi^2(H) = -\langle H | \bar{h} (iD)^2 h | H \rangle, \quad \mu_G^2(H) = \frac{C(m_Q)}{2} \langle H | \bar{h} \sigma_{\mu\nu} G^{\mu\nu} h | H \rangle,$$

$$\bar{\Lambda}(H) = M_H - m_Q + \frac{\mu_G^2(H) - \mu_\pi^2(H)}{2M_H} + \dots,$$

which are referred to as the “kinetic energy”, the “chromomagnetic interaction”, and the “binding energy”, respectively. They are important parameters entering many applications of the heavy-quark expansion, the calculation of

inclusive decay rates and spectra in particular. Certain combinations of these parameters can be extracted from spectroscopy:

$$\begin{aligned}\mu_G^2(B) &= \frac{3}{4} (M_{B^*}^2 - M_B^2) \approx 0.36 \text{ GeV}^2, & \mu_G^2(\Lambda_b) &= 0, \\ \mu_\pi^2(\Lambda_b) - \mu_\pi^2(B) &= -\frac{M_B M_D}{2} \left(\frac{M_{\Lambda_b} - M_{\Lambda_c}}{M_B - M_D} - \frac{3}{4} \frac{M_{B^*} - M_{D^*}}{M_B - M_D} - \frac{1}{4} \right) \approx 0, \\ \bar{\Lambda}(\Lambda_b) - \bar{\Lambda}(B) &\approx 0.31 \text{ GeV}.\end{aligned}$$

The individual parameters μ_π^2 and $\bar{\Lambda}$ are scheme dependent, however, because of renormalon ambiguities in their definition. Once a scheme is chosen, they can be calculated nonperturbatively, or extracted from moments of inclusive decay spectra. Table 1 shows a collection of some experimental determinations of these parameters in the on-shell scheme.

Table 1. Determinations of the parameters $\bar{\Lambda}$ and μ_π^2 from inclusive decay spectra

Reference	Method	$\bar{\Lambda}(B)$ [GeV]	$\mu_\pi^2(B)$ [GeV ²]
Falk et al. [5]	Hadron Spectrum	≈ 0.45	≈ 0.1
Gremm et al. [6]	Lepton Spectrum	0.39 ± 0.11	0.19 ± 0.10
Chernyak [7]	$(\bar{B} \rightarrow X \ell \bar{\nu})$	0.28 ± 0.04	0.14 ± 0.03
Gremm, Stewart [8]		0.33 ± 0.11	0.17 ± 0.10
Li, Yu [9]	Photon Spectrum $(\bar{B} \rightarrow X_s \gamma)$	$0.65_{-0.30}^{+0.42}$	$0.71_{-0.70}^{+1.16}$

Using the operator product expansion, any inclusive decay rate of a beauty hadron can be expanded as

$$\Gamma(H) = \frac{G_F^2 m_b^5}{192\pi^3} \left(1 - \frac{\mu_\pi^2(H)}{2m_b^2} \right) \left\{ c_3 + c_5 \frac{\mu_G^2(H)}{m_b^2} + \sum_n c_6^{(n)} \frac{\langle O_n \rangle_H}{m_b^3} + \dots \right\},$$

where $\langle O_n \rangle_H$ are the matrix elements of local four-quark operators, which parametrize nonspectator effects in these decays, and c_i are calculable short-distance coefficients, which depend on CKM parameters, the ratios of quark masses, and the renormalization scheme. The free quark decay emerges as the leading term in a systematic $1/m_b$ expansion, with bound-state corrections suppressed by two inverse powers of the heavy-quark mass. Note that ratios of inclusive decay rates are independent of μ_π^2 and the common factor m_b^5 , as well as of most CKM parameters. The application of the operator product expansion to the calculation of inclusive decay rates relies on the assumption of quark-hadron duality. Strictly speaking, the theoretical description of such processes is thus not entirely from first principles.

2 Exclusive semileptonic decays

The most important applications of the HQET concern the description of exclusive semileptonic decays based on the quark transition $b \rightarrow c \ell \bar{\nu}$. This is where the theory is well tested, and theoretical uncertainties are best understood. The most important result is a precision determination of the CKM parameter $|V_{cb}|$ [10].

2.1 Determination of $|V_{cb}|$ from $\bar{B} \rightarrow D^{(*)} \ell \bar{\nu}$ decays

The differential semileptonic decay rates as a function of the kinematical variable $w = v_B \cdot v_{D^{(*)}}$ are given by

$$\begin{aligned} \frac{d\Gamma(\bar{B} \rightarrow D^* \ell \bar{\nu})}{dw} &= \frac{G_F^2 M_B^5}{48\pi^3} r_*^3 (1 - r_*)^2 \sqrt{w^2 - 1} (w + 1)^2 \\ &\quad \times \left[1 + \frac{4w}{w + 1} \frac{1 - 2wr_* + r_*^2}{(1 - r_*)^2} \right] |V_{cb}|^2 \mathcal{F}^2(w), \\ \frac{d\Gamma(\bar{B} \rightarrow D \ell \bar{\nu})}{dw} &= \frac{G_F^2 M_B^5}{48\pi^3} r^3 (1 + r)^2 (w^2 - 1)^{3/2} |V_{cb}|^2 \mathcal{G}^2(w), \end{aligned}$$

where $r_{(*)} = M_{D^{(*)}}/M_B$. In the heavy-quark limit, the form factors $\mathcal{F}(w)$ and $\mathcal{G}(w)$ coincide with the universal function $\xi(w)$ and are thus normalized to unity at $w = 1$. Much effort has gone into calculating the symmetry-breaking corrections to this limit, with the result that [11]

$$\begin{aligned} \mathcal{F}(1) &= 1 + c_A(\alpha_s) + 0 + \delta_{1/m^2} + \dots = 0.924 \pm 0.027, \\ \mathcal{G}(1) &= 1 + c_V(\alpha_s) + \delta'_{1/m} + \delta'_{1/m^2} + \dots = 1.00 \pm 0.07, \end{aligned}$$

where the short-distance coefficients c_A and c_V are known to two-loop order [12, 13]. These numbers include the leading-logarithmic QED corrections. The absence of first-order power corrections to $\mathcal{F}(1)$ is a consequence of Luke's theorem [14]. The theoretical errors quoted above include the perturbative uncertainty and the uncertainty in the calculation (and truncation) of power corrections, added in quadrature. If instead the errors are added linearly, the result for $\mathcal{F}(1)$ changes to 0.924 ± 0.041 . This value has recently been confirmed in a different regularization scheme, in which the separation between short- and long-distance contributions is achieved by means of a hard momentum cutoff [15].

A value for $|V_{cb}|$ can be obtained by extrapolating experimental data for the differential decay rates to the zero-recoil point, using theoretical constraints on the shape of the form factors. Model-independent bounds on the physical $\bar{B} \rightarrow D^{(*)} \ell \bar{\nu}$ form factors can be derived using analyticity properties of QCD correlators, unitarity and dispersion relations [16, 17]. Combining these methods with the approximate heavy-quark symmetry, very powerful

one-parameter functions can be derived, which approximate the physical form factors in the semileptonic region with an accuracy of better than 2% [18]. For instance, the function $\mathcal{G}(w)$ can be parametrized as

$$\frac{\mathcal{G}(w)}{\mathcal{G}(1)} \approx 1 - 8\rho_1^2 z(w) + (51.\rho_1^2 - 10.)z^2(w) - (252.\rho_1^2 - 84.)z^3(w),$$

where $z(w) = (\sqrt{w+1} - \sqrt{2})/(\sqrt{w+1} + \sqrt{2})$, and ρ_1^2 is the (negative) slope of the form factor at zero recoil. A similar parametrization can be given for the function $\mathcal{F}(w)$. At present, these constraints are only partially included in the analyses of experimental data.

The world-average results of such analyses are [19, 20]

$$\begin{aligned} |V_{cb}| \mathcal{F}(1) &= (35.2 \pm 2.6) \times 10^{-3}, \\ |V_{cb}| \mathcal{G}(1) &= (38.6 \pm 4.1) \times 10^{-3}. \end{aligned}$$

A good fraction of the present errors reflect the uncertainty in the extrapolation to zero recoil, which could be avoided by implementing the dispersive constraints mentioned above. When combined with the theoretical predictions for the normalization of the form factors at zero recoil, the data yield the accurate value

$$|V_{cb}| = (38.2 \pm 2.3_{\text{exp}} \pm 1.2_{\text{th}}) \times 10^{-3},$$

which is in good agreement with an independent determination from inclusive B decays [1, 19].

2.2 Tests of heavy-quark symmetry

In general, the decays $\bar{B} \rightarrow D \ell \bar{\nu}$ and $\bar{B} \rightarrow D^* \ell \bar{\nu}$ are described by four independent form factors: $\mathcal{G}(w)$ for the former process, and $h_{A1}(w)$, $R_1(w)$, $R_2(w)$ – a combination of which defines the function $\mathcal{F}(w)$ – for the latter one. In the heavy-quark limit, $\mathcal{G}(w)$, $\mathcal{F}(w)$ and $h_{A1}(w)$ become equal to the function $\xi(w)$, whereas $R_1(w)$ and $R_2(w)$ approach unity. The universality of the function $\xi(w)$ can be tested by measuring the ratio $\mathcal{G}(w)/\mathcal{F}(w)$ as a function of w . The ALEPH data for this ratio [21] are shown in Fig. 2; a similar measurement has also been reported by CLEO [22]. Within errors, the data are compatible with a universal form factor. At large recoil, where the experimental errors are smallest, this provides a test of heavy-quark symmetry at the level of 10–15%.

A more refined analysis of symmetry-breaking corrections has been done by measuring the ratios R_1 and R_2 close to zero recoil. There the HQET predicts that [1]

Fig. 2. Ratio of the two form factors $\mathcal{G}(w)$ and $\mathcal{F}(w)$, from Ref. [21]

$$R_1 \approx 1 + \frac{4\alpha_s(m_c)}{3\pi} + \frac{\bar{\Lambda}}{2m_c} = 1.3 \pm 0.1,$$

$$R_2 \approx 1 - \frac{\bar{\Lambda}}{2m_c} = 0.8 \pm 0.2,$$

$$\rho_{A_1}^2 - \rho_{\mathcal{F}}^2 = \frac{R_1 - 1}{6} + \frac{1 - R_2}{3(1 - r_*)} = 0.2 \pm 0.1.$$

The CLEO data for these three quantities are $R_1 = 1.24 \pm 0.29$, $R_2 = 0.72 \pm 0.19$, and $\rho_{A_1}^2 - \rho_{\mathcal{F}}^2 = 0.20 \pm 0.19$ [23], in good agreement with the theoretical predictions. With a little more precision, the data would start to test the pattern of symmetry-breaking effects in the heavy-quark expansion.

2.3 Total exclusive semileptonic rates

Summing up the average experimental results $\text{B}(\bar{B} \rightarrow D \ell \bar{\nu}) = (1.95 \pm 0.27)\%$, $\text{B}(\bar{B} \rightarrow D^* \ell \bar{\nu}) = (5.05 \pm 0.25)\%$, $\text{B}(\bar{B} \rightarrow D^{(*)} \pi \ell \bar{\nu}) = (2.30 \pm 0.44)\%$, and adding $\text{B}(\bar{B} \rightarrow X_u \ell \bar{\nu}) = (0.15 \pm 0.10)\%$ for the inclusive branching ratio for semileptonic decays into charmless final states, one gets a total of $(9.45 \pm 0.58)\%$ [19], which is not far below the average value for the inclusive semileptonic branching ratio, $\text{B}(\bar{B} \rightarrow X \ell \bar{\nu}) = (10.19 \pm 0.37)\%$, measured at the $\Upsilon(4s)$ resonance (see below). Thus, there is little room for extra contributions.

A solid theoretical understanding of B decays into p-wave charm meson resonances would be important in order to address the question of whether there are additional contributions not accounted for in the above sum, and also to understand the main source of background in the determination of $|V_{cb}|$ from $\bar{B} \rightarrow D^* \ell \bar{\nu}$ decays. The description of these processes in the context of the HQET involves, at leading order, two new universal functions: $\tau_{3/2}(w)$ for the decays into the narrow states (D_1, D_2^*), and $\tau_{1/2}(w)$ for those into the broad states (D_0^*, D_1^*). In the heavy-quark limit, the differential (in w) decay rates vanish at zero recoil, since in that limit the p-wave states are orthogonal

to the ground state. There is a sizable $1/m_c$ correction to the $\bar{B} \rightarrow D_1 \ell \bar{\nu}$ decay rate at zero recoil, which can be calculated in a model-independent way in terms of known charm meson masses [24]. It is important, since the kinematical region is restricted close to zero recoil ($1 < w < 1.3$). Detailed theoretical predictions for the semileptonic and hadronic decay rates into p-wave charm states, which incorporate the constraints imposed by heavy-quark symmetry, can be found in Refs. [24, 25].

3 Semileptonic $b \rightarrow u$ decays and $|V_{ub}|$

Two exclusive semileptonic B decay modes into charmless hadrons have been observed by CLEO; the corresponding branching ratios are [26]

$$\text{B}(\bar{B} \rightarrow \pi \ell \bar{\nu}) = (1.8 \pm 0.5_{\text{exp}} \pm 0.2_{\text{model}}) \times 10^{-4},$$

$$\text{B}(\bar{B} \rightarrow \rho \ell \bar{\nu}) = (2.5^{+0.6}_{-0.8_{\text{exp}}} \pm 0.5_{\text{model}}) \times 10^{-4}.$$

That the theoretical description of these processes involves heavy-to-light form factors implies a certain amount of model dependence, since heavy-quark symmetry does not help to fix their normalization in a precise way. To some extent, a discrimination between models can be obtained by requiring a simultaneous fit of both exclusive channels. From a χ^2 -weighted average of models, CLEO obtains [26]

$$|V_{ub}| = (3.3 \pm 0.4_{\text{exp}} \pm 0.7_{\text{model}}) \times 10^{-3}.$$

Even with the present, very limited statistics of the measurements, the theoretical uncertainties are the limiting factor in the determination of $|V_{ub}|$. In the future, the model dependence can and will be reduced mainly by combining different theoretical approaches, in particular: lattice calculations, which are restricted to the region of large q^2 [27]; light-cone QCD sum rules, including $O(\alpha_s)$ and higher-twist corrections [28, 29]; analyticity and unitarity constraints [30, 31]; dispersion relations [32]. Although the optimal strategy is not yet clear at present, I believe it will be possible to reach the level of 15% theoretical uncertainty.

The traditional way to extract $|V_{ub}|$ from inclusive $\bar{B} \rightarrow X \ell \bar{\nu}$ decays has been to look at the endpoint region of the charged-lepton energy spectrum, where there is a tiny window not accessible to $\bar{B} \rightarrow X_c \ell \bar{\nu}$ decays. However, this method involves a large extrapolation and is plagued by uncontrolled (and often underestimated) theoretical uncertainties. A reanalysis of the available experimental data (using the ISGW2 model) gives $|V_{ub}| = (3.7 \pm 0.6_{\text{exp}}) \times 10^{-3}$ [33], in agreement with the value quoted above. A better discrimination between $b \rightarrow u$ and $b \rightarrow c$ transitions should use vertex information combined with a cut on the invariant mass M_h (or energy E_h) of the hadronic final state [34]–[40]. Parton model calculations (with Fermi motion included) indicate that about 90% of all $\bar{B} \rightarrow X_u \ell \bar{\nu}$ decays have $M_h < M_D$,

as shown in Fig. 3. Ideally, this cut would thus provide for a very efficient discriminator. In practise, there will be some leakage so that presumably one will be forced to require $M_h < M_{\max}$ with some threshold $M_{\max} < M_D$. The task for theorists is to calculate the fraction of events with hadronic mass $M_h < M_{\max}$,

$$\Phi(M_{\max}) = \frac{1}{\Gamma} \int_0^{M_{\max}} dM_h \frac{d\Gamma(\bar{B} \rightarrow X_u \ell \bar{\nu})}{dM_h}.$$

To calculate this fraction requires an ansatz for the “shape function”, which describes the Fermi motion of the b quark inside the B meson [41, 42]. The first three moments of this function are determined in terms of known HQET matrix elements. Still, some theoretical uncertainty remains, mainly associated with the values of the b -quark mass and the kinetic energy μ_π^2 , as well as unknown $O(\alpha_s^2)$ corrections. As an example, Fig. 3 shows the dependence of $\Phi(M_{\max})$ on the value of m_b . It turns out that the resulting theoretical uncertainty strongly depends on the value of the threshold $M_{\max}$. First estimates yield $\delta|V_{ub}|/|V_{ub}| \approx 10\%$ for $M_{\max} = M_D$, and $\delta|V_{ub}|/|V_{ub}| \approx 20\%$ for $M_{\max} = 1.5 \text{ GeV}$ [38, 39]. This new method is challenging both for theorists and for experimenters, but it is superior to the endpoint method. I believe that ultimately a theoretical accuracy of 10% can be reached.

Fig. 3. Hadronic mass distribution (left) and fraction of $\bar{B} \rightarrow X_u \ell \bar{\nu}$ decays with invariant hadronic mass below $M_{\max}$ (right), from Ref. [38]

4 Semileptonic branching ratio and charm multiplicity

For many years, the apparent discrepancy between (some) measurements of the semileptonic branching ratio of B mesons and theoretical predictions for this quantity has given rise to controverse dispute and speculations about

deviations from the Standard Model. Another important input to this discussion has been the charm multiplicity n_c , i.e. the average number of charm (or anticharm) quarks in the hadronic final state of a B decay. From the theoretical point of view,

$$B_{\text{SL}} = \frac{\Gamma(\bar{B} \rightarrow X e \bar{\nu})}{\sum_{\ell} \Gamma(\bar{B} \rightarrow X \ell \bar{\nu}) + \Gamma_{\text{had}} + \Gamma_{\text{rare}}},$$

$$n_c = 1 + B(\bar{B} \rightarrow X_{cc}) - B(\bar{B} \rightarrow \text{no charm})$$

are governed by the same partial inclusive decay rates. The theoretical predictions for these two quantities depend mainly on two parameters: the quark-mass ratio m_c/m_b and the renormalization scale μ [43, 44]. The latter dependence reflects our ignorance about higher-order QCD corrections to the decay rates. The results obtained by allowing reasonable ranges for these parameters are represented by the dark-shaded area in Fig. 4. The two data points show the average experimental results obtained from experiments operating at the $\Upsilon(4s)$ resonance: $B_{\text{SL}} = (10.19 \pm 0.37)\%$ and $n_c = 1.12 \pm 0.05$, and at the Z resonance: $B_{\text{SL}} = (11.12 \pm 0.20)\%$ and $n_c = 1.20 \pm 0.07$ [19, 45]. At this conference, it has been emphasized that a dedicated reanalysis of the LEP data for the semileptonic branching ratio is necessary, because some sources of systematic errors had previously been underestimated [45]. To account for this, I have doubled the corresponding error bar on B_{SL} in Fig. 4. If we ignore the LEP point for the moment, it appears that the theoretical predictions for both B_{SL} and n_c lie significantly higher than the experimental results.

Fig. 4. Theory [44] versus experiment for the semileptonic branching ratio and charm multiplicity

At this conference, several new ingredients have been presented that shed some light on this problem. First, there are some nontrivial tests of the the-

ory. Table 2 shows details of the theoretical calculation in the form of the dominant partial decay rates normalized to $B(\bar{B} \rightarrow X_c \ell \bar{\nu})$, for $m_c/m_b = 0.29 \pm 0.03$, using the results of Refs. [43, 44, 46]. (Note that the values for $r_{c\bar{u}d'}$ are larger than the value 4.0 ± 0.4 often used in the literature.) It has been argued that the assumption of local quark–hadron duality, which underlies the theoretical treatment of inclusive decay rates, may fail for the decays $\bar{B} \rightarrow X_{c\bar{c}s'}$, because there is only little kinetic energy released to the final state. To test this hypothesis, one can eliminate the corresponding partial decay rate, in which case one obtains a linear relation between n_c and B_{SL} [47]. The result is shown as the light band in Fig. 4. The fact that, within this band, the original prediction (i.e. the dark-shaded area) is closest to the data indicates that there is no problem with quark–hadron duality. Another important check is provided by an experimental determination of the ratio $r_{c\bar{u}d'}$ using flavour-specific measurements of charm branching ratios [48]. The result is [49]–[51]

$$r_{c\bar{u}d'} = \frac{B(\bar{B} \rightarrow \text{open } c) - B(\bar{B} \rightarrow \text{open } \bar{c})}{B_{\text{SL}}} - (2 + r_{c\tau\bar{\nu}} - r_{u\bar{c}s'}) = 4.1 \pm 0.7,$$

in good agreement with the theoretical predictions given in Table 2. The theory input $r_{c\tau\bar{\nu}} - r_{u\bar{c}s'} = 0.19 \pm 0.03$ in this extraction has a very small uncertainty. In summary, it appears that the heavy-quark expansion works well for both of the hadronic decay rates.

Table 2. Predictions for ratios of partial inclusive decay rates

	$r_{c\tau\bar{\nu}}$	$r_{c\bar{u}d'}$	$r_{c\bar{c}s'}$	$r_{\text{no charm}}$
$\mu = m_b$	0.22 ∓ 0.03	4.21 ± 0.01	1.89 ∓ 0.44	0.14 ± 0.04
$\mu = m_b/2$	0.23 ∓ 0.03	4.75 ± 0.02	2.20 ∓ 0.49	0.19 ± 0.04

4.1 Is there a “missing charm puzzle”?

Several suggestions have been made to explain why the experimental value of $n_c = 1.12 \pm 0.05$ measured by CLEO [19] is smaller than the theoretical prediction $n_c = 1.20 \pm 0.06$ [44]. I have discussed above that the problem cannot be blamed on violations of quark–hadron duality, as was originally speculated by some authors. An interesting new proposal made by Dunietz et al. [49] is based on the fact that the theoretical definition of n_c refers to a fully inclusive quantity counting the number of charm and anticharm quarks per B decay, irrespective of whether they end up as “open” or “hidden” charm. If there were a sizable branching ratio for decays into hadronic final states containing undetected $(c\bar{c})$ pairs,

$$B(\bar{B} \rightarrow (c\bar{c})_{\text{undetected}} + X) \equiv b_{(c\bar{c})},$$

then the experimentally observed value of n_c would be lower than the theoretical one: $n_c^{\text{obs}} = n_c^{\text{th}} - 2b_{(c\bar{c})}$. Note that the conventional charmonium states J/ψ , ψ' , χ_{c1} , χ_{c2} , η_c are included in the charm counting and thus do not contribute to $b_{(c\bar{c})}$. However, it has been speculated that there may exist an enhanced production of exotic $(c\bar{c}g)$ hybrids, which decay into noncharmed light mesons and could yield a value of $b_{(c\bar{c})}$ of about 10% [49, 52].

Alternatively, enhanced flavour-changing neutral current (FCNC) processes such as $b \rightarrow sg$ could, simultaneously, lower the predictions for B_{SL} and n_c by a factor of $[1 + B(\text{new FCNC})]^{-1}$, thus providing for a “new physics explanation” of the missing charm puzzle [53, 54].

Fig. 5. Mechanisms for D and $\bar{D}$ production in $\bar{B}$ -meson decays

At least a partial answer to the question of whether there is any support for such speculations is provided by new, flavour-specific measurements of charm production. Using charm-particle-lepton correlations in tagged B decays, it is possible to measure the relative rate of the two mechanisms shown in Fig. 5. Whereas it was previously assumed that the first mechanism was responsible for all D -meson production, recent measurements indicate a rather large “wrong-charm” yield [48]:

$$\frac{B(\bar{B} \rightarrow \bar{D} X)}{B(\bar{B} \rightarrow D X)} = 0.100 \pm 0.026 \pm 0.016,$$

corresponding to a “wrong-charm” branching ratio $B(\bar{B} \rightarrow \bar{D} X) = (7.9 \pm 2.2)\%$. This observation is supported by measurements of both exclusive [55] and inclusive [56, 57] production of $D\bar{D}$ meson pairs. This effect was not included in Monte Carlo B -decay generators and may be partly responsible for the discrepancy in the CLEO and LEP measurements of B_{SL} and n_c [49]. When combined with previously measured decay rates, the new CLEO result implies [48]

$$B(\bar{B} \rightarrow X_{c\bar{c}}) - b_{(c\bar{c})} = (21.9 \pm 3.7)\%,$$

which is close to the theoretical prediction $B(\bar{B} \rightarrow X_{c\bar{c}}) = (22 \pm 6)\%$ [44], indicating that there is not much room for decays into undetected $(c\bar{c})$ pairs, represented by $b_{(c\bar{c})}$. This measurement is supported by a DELPHI result on the production of two open charm particles [58]. Furthermore, by measuring a double ratio of flavour-specific rates, CLEO obtains a bound on charmless modes that is largely independent of detection efficiencies and charm branching ratios. The result is $B(\bar{B} \rightarrow \text{no open charm}) = (3.2 \pm 4.0)\% < 9.6\%$ (90%

CL), which after subtraction of the known charmonium contributions implies [48]

$$B(\bar{B} \rightarrow \text{no charm}) + b_{(c\bar{c})} = (0.2 \pm 4.1)\% < 6.8\% \quad (90\% \text{ CL}).$$

Given that the Standard Model prediction for the charmless rate is $B(\bar{B} \rightarrow \text{no charm}) = (1.6 \pm 0.8)\%$ [46], it appears that there is little room for either hidden $(c\bar{c})$ production or new physics contributions. Again, this conclusion is supported by a DELPHI result obtained using impact parameter measurements: $B(\bar{B} \rightarrow \text{no open charm}) = (4.5 \pm 2.5)\% < 8.4\%$ (90% CL) [58].

Combining these two measurements, I conclude that $b_{(c\bar{c})} < 5\%$ (90% CL) cannot be anomalously large. The same bound applies to other, nonstandard sources of charmless B decays, i.e. $B(\text{new FCNC}) < 5\%$ (90% CL). This conclusion is supported by the DELPHI limit $B(\bar{B} \rightarrow X_{sg}) < 5\%$ (95% CL) obtained from a study of the $p_\perp$ spectrum of charged kaons produced in B decays [58]. It must be noted, however, that a preliminary indication of a kaon excess at large $p_\perp$, as expected from enhanced $b \rightarrow sg$ transitions [50], has been reported by SLD at this conference [59]. Even though a definite conclusion can therefore not be drawn before these measurements become final, at present there is no compelling experimental evidence of any nonstandard physics in inclusive B decays.

The two CLEO measurements quoted above can be combined to give a new determination of the charm multiplicity, in which the unknown quantity $b_{(c\bar{c})}$ cancels out. The result

$$n_c = 1 + \left[B(\bar{B} \rightarrow X_{c\bar{c}}) - b_{(c\bar{c})} \right] - \left[B(\bar{B} \rightarrow \text{no charm}) + b_{(c\bar{c})} \right] = 1.22 \pm 0.06$$

is significantly higher than (though consistent with) the value 1.12 ± 0.05 obtained using the conventional method of charm counting, and in excellent agreement with the theoretical prediction. I believe that, ultimately, this new way of measuring n_c will be less affected by systematic uncertainties than the traditional one. It seems that the “missing charm puzzle” is about to disappear.

5 Beauty lifetime ratios

The current world-average experimental results for the lifetime ratios of different beauty hadrons are [60]

$$\frac{\tau(B^-)}{\tau(B^0)} = 1.06 \pm 0.04, \quad \frac{\tau(B_s)}{\tau(B_d)} = 0.98 \pm 0.07, \quad \frac{\tau(\Lambda_b)}{\tau(B^0)} = 0.78 \pm 0.04.$$

Theory predicts that $|\tau(B_s)/\tau(B_d) - 1| < 1\%$ [61], and it will be very difficult to push the experimental accuracy to a level where one would become sensitive to sub-1% effects. The theoretical predictions for the other two ratios

have been analysed to third order in the heavy-quark expansion. Unfortunately, these predictions depend on some yet unknown “bag parameters” B_1 , B_2 , ε_1 , ε_2 , $\tilde{B}$, r parametrizing the hadronic matrix elements of local four-quark operators. In terms of these parameters, the results are [44]

$$\frac{\tau(B^-)}{\tau(B^0)} = 1 + 16\pi^2 \frac{f_B^2 M_B}{m_b^3} \left[k_1 B_1 + k_2 B_2 + k_3 \varepsilon_1 + k_4 \varepsilon_2 \right],$$

$$\frac{\tau(\Lambda_b)}{\tau(B^0)} = 0.98 + 16\pi^2 \frac{f_B^2 M_B}{m_b^3} \left[p_1 B_1 + p_2 B_2 + p_3 \varepsilon_1 + p_4 \varepsilon_2 + (p_5 + p_6 \tilde{B}) r \right],$$

where k_i and p_i are short-distance coefficients, whose values depend on the ratio m_c/m_b and on the renormalization scale. The large- N_c counting rules of QCD imply that $B_i = O(1)$ and $\varepsilon_i = O(1/N_c)$. The factorization approximation for the meson matrix elements suggests that $B_i \approx 1$ and $\varepsilon_i \approx 0$ [62]. Similarly, the constituent quark model suggests that $\tilde{B} \approx 1$ and $r \approx |\psi_{q\bar{q}}^{\Lambda_b}(0)|^2/|\psi_q^B(0)|^2$. The parameter r is the most uncertain one entering the predictions for the lifetime ratios. Existing theoretical estimates for this parameter range from 0.1 to 2. Some recent estimates can be found in Refs. [63, 64].

Without a reliable field-theoretical calculation of the bag parameters, no lifetime “predictions” can be obtained. However, one can see which ranges of the lifetime ratios can be covered using sensible values for the bag parameters [65]. To this end, I scan the following parameter space: $B_i, \tilde{B} \in [2/3, 4/3]$, $\varepsilon_i \in [-1/3, 1/3]$, $r \in [0.25, 2.5]$, $m_c/m_b = 0.29 \pm 0.03$, $f_B = (200 \pm 20)$ MeV. The resulting distributions for the lifetime ratios are shown in Fig. 6. The most important observation of this exercise is that it is possible to reproduce both lifetime ratios, within their experimental uncertainties, using the same set of input parameters. The theoretically allowed range for the ratio $\tau(B^-)/\tau(B^0)$ is centered around the experimental value; however, the width of the allowed region is so large that no accurate “prediction” of this ratio could have been made. Any value between 0.8 and 1.3 could be easily accommodated by theory. The predictions for the ratio $\tau(\Lambda_b)/\tau(B^0)$, on the other hand, center around a value of 0.95, and the spread of the results is much narrower. Only a small tail extends into the region preferred by experiment. Requiring that the theoretical results be inside the 2σ ellipse around the central experimental values, I find that the value of the parameter ε_1 must be close to zero, $\varepsilon_1 \approx -(0.1 \pm 0.1)$, in agreement with the expectation based on factorization. In addition, this requirement maps out the region of parameter space for ε_2 and r shown in the right lower plot of Fig. 6. Thus, the data indicate a large (with respect to most model predictions) value of r and a negative value of ε_2 . Ultimately, it will be important to perform reliable field-theoretical calculations of the hadronic parameters, for instance using lattice gauge theory or QCD sum rules, to see whether indeed the observed beauty lifetime ratios can be accounted for by the heavy-quark expansion (for a recent sum-rule estimate of the parameters B_i and ε_i , see Ref. [66]).

Fig. 6. Upper plots: Theoretical distributions for beauty lifetime ratios. The two curves in each figure correspond to different choices of the renormalization scale. The shaded areas show the experimental values with their 2σ error bands. Lower left: Two-dimensional distribution of lifetime ratios, together with the 1σ and 2σ contours around the central experimental values. Lower right: Distribution of the hadronic parameters ε_2 and r for simulations with results inside the 2σ ellipse.

This is particularly important for the parameter r , large values of which are not excluded a priori. Using a variant of the quark-model relation derived in Ref. [63], combined with a preliminary DELPHI measurement of the mass-splitting between the Σ_b^* and Σ_b baryons [67], I find

$$r \approx \frac{4}{3} \frac{M_{\Sigma_b^*}^2 - M_{\Sigma_b}^2}{M_{B^*}^2 - M_B^2} = 1.8 \pm 0.5.$$

With such a large value, it is possible to explain the short Λ_b lifetime without invoking violations of local quark-hadron duality.

Although at present there is thus no convincing evidence that the low value of the Λ_b lifetime could not be accommodated in the context of the heavy-quark expansion, it remains a possibility that in the future such a discrepancy may emerge, for instance if lattice calculations would show that $r \ll 1$. Then, one would have to blame violations of local quark-hadron duality to be responsible for the failure of the heavy-quark expansion for

inclusive, nonleptonic B -decay rates. Recently, several authors have studied in QCD-inspired models how quark-hadron duality may be violated. Based on a simple model for the difference of two heavy-light current correlators in the chiral and large- N_c limits, Shifman has argued that deviations from local duality are due to the asymptotic nature of the operator product expansion (OPE) [68]. More recently, Blok et al. have studied QCD₂ (i.e. QCD in 1+1 space-time dimensions) at next-to-leading order in $1/N_c$, where finite resonance widths provide for a dynamical smearing of correlation functions [69]. They find that, in the Euclidean region, deviations from local duality are indeed due to the divergence of the OPE, and are exponentially suppressed with Q^2 . In the physical (Minkowskian) region, on the other hand, they find that two mass scales, Λ_1 and Λ_2 , are dynamically generated. For $\Lambda_1^2 < q^2 < \Lambda_2^2$, there are strong oscillations of correlation functions. Global duality works, but local duality may be strongly violated. For $q^2 > \Lambda_2^2$, the oscillations become exponentially damped, and local duality is restored. They conjecture that a similar pattern may hold for real, four-dimensional QCD. Similar results have also been obtained by Grinstein and Lebed [70], and by Chibisov et al. studying the instanton vacuum model [71]. Quite different conclusions have been reached by Colangelo et al., who have considered a more “realistic” version of Shifman’s model, finding that global duality holds, but local duality is spoiled by $1/Q$ effects not present in the OPE [72]. This study gives some support for the conjecture by Altarelli et al. that there may be non-OPE terms of order $\bar{\Lambda}/m_Q$ increasing the nonleptonic decay rates of heavy hadrons [73]. Such terms could easily explain the short Λ_b lifetime. However, I stress that, although the model of Ref. [72] is interesting, it must not be taken as an existence proof of non-OPE power corrections. Nature may be careful enough to avoid power-like deviations from local duality.

6 Hadronic B decays

The strong interaction effects in hadronic decays are much more difficult to understand, even at a qualitative level, than in leptonic or semileptonic decays. The problem is that multiple gluon exchanges can redistribute the quarks in the final state of a hadronic decay. As a consequence, some phenomenological assumptions are unavoidable in trying to understand these processes.

6.1 Energetic two-body decays

It has been argued that energetic two-body decays with a large energy release are easier to understand because of the colour-transparency phenomenon: a pair of fast-moving quarks in a colour-singlet state, which is produced in a local interaction, effectively decouples from long-wavelength gluons [74]. This intuitive argument suggests that nonleptonic amplitudes factorize to a

good approximation. The main strong-interaction effects are then of a short-distance nature and simply renormalize the operators in the effective weak Hamiltonian [75]. For the case of $b \rightarrow c\bar{u}d$ transitions, for instance,

$$H_{\text{eff}} = \frac{G_F}{\sqrt{2}} V_{cb} V_{ud}^* \left\{ c_1(\mu) (\bar{d}u)(\bar{c}b) + c_2(\mu) (\bar{c}u)(\bar{d}b) \right\} + \dots,$$

where $(\bar{d}u) = \bar{d}\gamma^\mu(1 - \gamma_5)u$ etc. are left-handed, colour-singlet quark currents, and $c_1(m_b) \approx 1.1$ and $c_2(m_b) \approx -0.3$ are Wilson coefficients taking into account the short-distance corrections arising from the exchange of hard gluons. The effects of soft gluons remain in the hadronic matrix elements of the local four-quark operators. In general, a reliable field-theoretical calculation of these matrix elements is the obstacle to a quantitative theory of hadronic weak decays.

Using Fierz identities, the four-quark operators in the effective Hamiltonian may be rewritten in various forms. It is particularly convenient to rearrange them in such a way that the flavour quantum numbers of one of the quark currents match those of one of the hadrons in the final state of the considered decay process. As an example, consider the decays $\bar{B} \rightarrow D\pi$. Omitting common factors, the various amplitudes can be written as

$$\begin{aligned} A_{\bar{B}^0 \rightarrow D^+ \pi^-} &= \left(c_1 + \frac{c_2}{N_c} \right) \langle D^+ \pi^- | (\bar{d}u)(\bar{c}b) | \bar{B}^0 \rangle + 2c_2 \langle D^+ \pi^- | (\bar{d}t_a u)(\bar{c}t_a b) | \bar{B}^0 \rangle \\ &\equiv a_1 \langle \pi^- | (\bar{d}u) | 0 \rangle \langle D^+ | (\bar{c}b) | \bar{B}^0 \rangle, \\ A_{\bar{B}^0 \rightarrow D^0 \pi^0} &= \left(c_2 + \frac{c_1}{N_c} \right) \langle D^0 \pi^0 | (\bar{c}u)(\bar{d}b) | \bar{B}^0 \rangle + 2c_1 \langle D^0 \pi^0 | (\bar{c}t_a u)(\bar{d}t_a b) | \bar{B}^0 \rangle \\ &\equiv a_2 \langle D^0 | (\bar{c}u) | 0 \rangle \langle \pi^0 | (\bar{d}b) | \bar{B}^0 \rangle, \end{aligned}$$

where t_a are the SU(3) colour matrices. The two classes of decays shown above are referred to as class-1 and class-2, respectively. The factorized matrix elements in the last steps are known in terms of the meson decay constants f_π and f_D , and the transition form factors for the decays $\bar{B} \rightarrow D$ and $\bar{B} \rightarrow \pi$. Most of these quantities are accessible experimentally. Of course, the above matrix elements also contain other, nonfactorizable contributions. They are absorbed into the definition of the hadronic parameters a_1 and a_2 , which in general are process dependent. Recently, some progress in the understanding of these parameters has been made, leading to the “generalized factorization hypothesis” that [76]–[78]

$$a_1 \approx c_1(m_b), \quad a_2 \approx c_2(m_b) + \zeta c_1(m_b),$$

where ζ is a process-independent hadronic parameter (for energetic two-body decays only!), which accounts for the dominant nonfactorizable contributions to the decay amplitudes. To derive these results, one combines the $1/N_c$ expansion with an argument inspired by colour transparency [78].

Table 3. Experimental tests of the generalized factorization hypothesis

	a_1	a_2/a_1
$\bar{B} \rightarrow D^{(*)}\pi$	1.08 ± 0.06	0.21 ± 0.07
$\bar{B} \rightarrow D^{(*)}\rho$	1.07 ∓ 0.07	0.23 ± 0.14
$\bar{B} \rightarrow \psi^{(\prime)} K^{(*)}$		$ a_2 = 0.21 \pm 0.04$
$\bar{B} \rightarrow D^{(*)}\bar{D}_s^{(*)}$	1.10 ± 0.18	

Some experimental tests of the generalized factorization hypothesis are shown in Table 3, where I quote the values of a_1 and a_2 extracted from the analysis of different classes of decay modes, using data reported by CLEO [79]–[81]. Within the present experimental errors, there is indeed no evidence for any process dependence of the hadronic parameters. The generalized factorization prescription provides a simultaneous description of all measured Cabibbo-allowed two-body decays with a single parameter $\zeta = 0.45 \pm 0.05$ extracted from the data on class-2 decays [78]. So far, this theoretical framework is fully supported by the data.

The factorization hypothesis can be used to obtain rather precise values for the decay constants of the D_s and D_s^* mesons, by comparing the theoretical predictions for various ratios of $\bar{B}^0$ decay rates,

$$\begin{aligned} \frac{\Gamma(D^+ D_s^-)}{\Gamma(D^+ \pi^-)} &= 1.01 \left(\frac{f_{D_s}}{f_\pi} \right)^2, & \frac{\Gamma(D^{*+} D_s^-)}{\Gamma(D^{*+} \pi^-)} &= 0.72 \left(\frac{f_{D_s}}{f_\pi} \right)^2, \\ \frac{\Gamma(D^+ D_s^{*-})}{\Gamma(D^+ \rho^-)} &= 0.74 \left(\frac{f_{D_s^*}}{f_\rho} \right)^2, & \frac{\Gamma(D^{*+} D_s^{*-})}{\Gamma(D^{*+} \rho^-)} &= 1.68 \left(\frac{f_{D_s^*}}{f_\rho} \right)^2, \end{aligned}$$

with data. These predictions are rather clean for the following reasons: first, all decays involve class-1 transitions, so that deviations from factorization are probably very small; secondly, the parameter a_1 cancels in the ratios; thirdly, the two processes in each ratio have a similar kinematics, so that the corresponding decay rates are sensitive to the same form factors, however evaluated at different q^2 values. Finally, some of the experimental systematic errors cancel in the ratios (however, I do not assume this in quoting errors below). Combining these predictions with the average experimental branching ratios [79] yields

$$f_{D_s} = (234 \pm 25) \text{ MeV}, \quad f_{D_s^*} = (271 \pm 33) \text{ MeV}.$$

The result for f_{D_s} is in good agreement with the value $f_{D_s} = 250 \pm 37$ MeV extracted from the leptonic decay $D_s \rightarrow \mu^+ \nu$ [82]. The ratio of decay constants, $f_{D_s^*}/f_{D_s} = 1.16 \pm 0.19$, which cannot be determined from leptonic decays, is in good agreement with theoretical expectations [83, 84]. Finally, I note that, assuming SU(3)-breaking effects of order 10–20%, the established

value of f_{D_s} implies $f_D \gtrsim 200$ MeV, which is larger than most theoretical predictions.

6.2 B Decays into two light mesons

Since last year, CLEO has observed (or put strict upper limits on) a number of rare B decay modes, many of which have strongly suppressed tree amplitudes and are thus dominated by loop processes. The relevant quark diagrams for the transitions $b \rightarrow s(d)\bar{q}q$ are shown in Fig. 7, where I also give the powers of the Wolfenstein parameter $\lambda \approx 0.22$ associated with these diagrams. Because the penguin diagram shown on the right is a loop process, it is sensitive to new heavy particles and thus potentially probes physics beyond the Standard Model.

Fig. 7. Tree and penguin diagrams for rare B decays into light hadrons

Some of the experimental results that have caused a lot of excitement are (all in units of 10^{-5} and averaged over CP-conjugate modes; upper limits at 90% CL) [81, 85, 86]:

$$\mathcal{B}(B^0 \rightarrow K^+\pi^-) = 1.5_{-0.4}^{+0.5} \pm 0.1 \pm 0.1,$$

$$\mathcal{B}(B^+ \rightarrow K^0\pi^+) = 2.3_{-1.0}^{+1.1} \pm 0.3 \pm 0.2,$$

$$\mathcal{B}(B^0 \rightarrow \pi^+\pi^-) = 0.7 \pm 0.4 < 1.5,$$

$$\mathcal{B}(B^+ \rightarrow \pi^+\pi^0) = 1.0_{-0.5}^{+0.6} < 2.0,$$

$$\mathcal{B}(B^+ \rightarrow K^+\eta') = 7.1_{-2.1}^{+2.5} \pm 0.9,$$

$$\mathcal{B}(B \rightarrow \eta' + X_s) = 62 \pm 16 \pm 13 \quad (2.0 < p_{\eta'} [\text{GeV}] < 2.7).$$

In Table 4, I show the results of an analysis of these results in terms of SU(3)-invariant amplitudes representing diagrams having tree topology $T^{(\prime)}$, colour-suppressed tree topology $C^{(\prime)}$, penguin topology $P^{(\prime)}$, and SU(3)-singlet penguin topology $S^{(\prime)}$, where primed (unprimed) quantities refer to $b \rightarrow s(d)$

transitions [87]. The results of this analysis establish the presence of a significant SU(3)-singlet contribution S' active in the decays with an η' meson in the final state. They also indicate a rather large “penguin pollution” $|P/T| \approx 0.3 \pm 0.1$ in the decays $B \rightarrow \pi\pi$. This has important (and optimistic) implications for CP-violation studies using these decay modes [88].

Table 4. Amplitude analysis for some rare B decays into two light mesons. The dominant amplitudes are highlighted.

Decay mode	SU(3) invariant amplitudes	(in units of 10^{-6})
$B^0 \rightarrow K^+\pi^-$ $B^+ \rightarrow K^0\pi^+$	$-(T' + P') \left\{ \begin{array}{l} \\ P' \end{array} \right\}$	$ P' ^2 \approx 16 \pm 4$ $\Rightarrow P ^2 \approx 0.8 \pm 0.4$
$B^0 \rightarrow \pi^+\pi^-$ $B^+ \rightarrow \pi^+\pi^0$	$-(T + P) \left\{ \begin{array}{l} \\ -\frac{1}{\sqrt{2}}(T + C) \end{array} \right\}$	$ T ^2 \approx 8 \pm 4$ $\Rightarrow T' ^2 \approx 0.4 \pm 0.2$
$B^+ \rightarrow \eta' K^+$	$\frac{1}{\sqrt{6}}(T' + C' + 3P' + 4S')$	$ S' ^2 \approx 18 \pm 3$ no interference $ S ^2 \approx 5 \pm 2$ constr. interference

Several authors have discussed the question whether the penguin amplitudes in these transitions are anomalously large. There seems to be some agreement that a standard analysis, using next-to-leading order Wilson coefficients combined with factorized matrix elements, can accommodate the experimental results for the exclusive rare decay rates [89, 90]. However, it has been stressed that significant long-distance contributions to the charm penguin are required to fit the data [91, 92]. This is not too surprising, since the energy release in B decays is such that the $c\bar{c}$ pair in the charm penguin is not far from its mass shell, and hence this penguin is really more a long-distance than a short-distance process [93]. This is illustrated in Fig. 8. In the case of $B \rightarrow K\pi$ decays, the box may represent final-state rescattering processes such as $B \rightarrow D_s\bar{D} \rightarrow K\pi$ [94]–[97]. If the final state contains an η' meson, the box may represent an anomaly-mediated coupling of the η' to glue, or an intrinsic charm component in the η' wave function. At least to some extent these are different words for the same physics.

Fig. 8. Long-distance effects in “charming penguins”

In a way, the charm penguin is all there is in $b \rightarrow s$ FCNC processes [93], because the unitarity of the CKM matrix implies that

$$\sum_{q=u,c,t} V_{qs}^* V_{qb} P_q \approx V_{cs}^* V_{cb} (P_c - P_t).$$

The top penguin P_t simply provides the GIM cutoff for large loop momenta, whereas the up penguin P_u is strongly CKM suppressed. The construction of the effective weak Hamiltonian is used to separate the short- and long-distance contributions to the charm penguin, in a way that is illustrated in Fig. 9. The short-distance contributions from large loop momenta are contained in the local operators $Q_{3,\dots,6}$ and Q_8 , while long-distance effects from small momenta are contained in the matrix elements of the current-current operators $Q_{1,2}$. From the physical argument presented above, one expects sizable long-distance contributions, since the energy release in B decays is in the region of charm resonances. Therefore, purely short-distance estimates of the charm penguin may be very much misleading.

$$\begin{aligned} (P_c - P_t) &= \text{diagram with } b \text{ and } s \text{ lines, a top quark loop, and } k < m_t \\ &= \sum_{i=3}^6 c_i(\mu) \times \text{diagram with } b \text{ and } s \text{ lines meeting at a vertex } i \text{ with two } q \text{ lines} + c_8(\mu) \times \text{diagram with } b \text{ and } s \text{ lines and a gluon loop} \\ &+ \sum_{i=1,2} c_i(\mu) \times \text{diagram with } b \text{ and } s \text{ lines, a charm quark loop, and } k < \mu \\ &= P_{\text{SD}}(\mu) + P_{\text{LD}}(\mu) \quad (\text{only sum is RG-invariant!}) \end{aligned}$$

Fig. 9. Operator product expansion for the charm penguin

6.3 Explanations of the η' modes

To explain the yield of η' mesons in rare B decays, in particular the large inclusive η' production rate, is a challenge to theorists. Currently, the situation is still controversial, and several proposals are being discussed. It has been suggested that the process $b \rightarrow sg^*$ followed by $g^* \rightarrow \eta'g$ is enhanced, either by the anomalous coupling of the η' to glue [98]–[100], or because the initial $b \rightarrow sg$ rate is enhanced by some new physics [101]. Because of its three-body nature, this mechanism would mainly be responsible for multi-particle

inclusive processes containing η' mesons, but give a minor contribution to the two-body modes $B \rightarrow \eta' K^{(*)}$. It has also been argued that an enhanced η' production could result from a significant intrinsic charm component in the η' wave function, either through the colour-singlet mechanism $b \rightarrow (c\bar{c})_1 s$ followed by $(c\bar{c})_1 \rightarrow \eta'$ [102, 103], or through the colour-octet mechanism $b \rightarrow (c\bar{c})_8 s$ followed by $(c\bar{c})_8 \rightarrow \eta' X$ [104], which is strongly favoured by the structure of the effective weak Hamiltonian. The relevance of the intrinsic charm mechanism depends on the size of a “decay constant” $f_{\eta'}^{(c)}$, estimates of which range from 6 MeV to 180 MeV. A clarification of this issue is needed (see, e.g., the discussion in Refs. [89, 105]). Yet other possibilities to explain the large η' yield have been explored in Refs. [106, 107].

Table 5. Model predictions for ratios of rare B decay rates

	Ali, Greub [89]	Lipkin [109]	Cheng, Tseng [110]	Datta et al. [111]	Halperin et al. [102]
$B^+ \rightarrow \eta' K^{*+}$	0.01–0.2		0.20–0.24	0.36–0.38	≈ 2
$B^+ \rightarrow \eta' K^+$					
$B^+ \rightarrow \eta K^+$	0.04–0.14	0–0.02	0.03–0.08	0.02–0.09	
$B^+ \rightarrow \eta' K^+$					
$B^+ \rightarrow \eta K^{*+}$	0.6–2.9	0.5–8.2	0.26–0.48	1.0–1.4	
$B^+ \rightarrow \eta' K^{*+}$					

Future measurements of other decay channels, as well as of the distribution of the invariant hadronic mass $M(X_s)$ in the decay $B \rightarrow \eta' X_s$, will help to clarify the situation. Preliminary CLEO data indicate that relatively large values of $M(X_s)$ are preferred [81, 108]. Another good discriminator between models is provided by the ratios of the various $B \rightarrow \eta^{(\prime)} K^{(*)}$ decay rates. A collection of theoretical predictions for such ratios is shown in Table 5. In summary, the rare decays of B mesons into light mesons provide unique opportunities to extract many yet unknown hadronic matrix elements. At present, there are no convincing arguments that these processes could not be understood in a conventional Standard Model framework. In particular, because of the complexity of the dynamics of the η' meson, I believe it is premature to deduce evidence for new physics from the yield of η' production in B decays.

7 Summary and outlook

Owing to the combined efforts of experimenters and theorists, there has been significant progress in heavy-quark physics in the past few years. The exclusive and inclusive semileptonic decays mediated by the transition $b \rightarrow c \ell \bar{\nu}$ are understood from first principles, using the heavy-quark expansion. Heavy-quark effective theory (HQET) works well in describing these processes, and

starts being tested at the level of symmetry-breaking corrections. This, by itself, is a remarkable development, which puts the determination of $|V_{cb}| = 0.038 \pm 0.003$ on firm theoretical grounds. New ideas are being developed for getting a precise ($\sim 10\%$) determination of $|V_{ub}|$, combining exclusive and inclusive methods. The error in the current value, $|V_{ub}| = (3.3 \pm 0.8) \times 10^{-3}$, is still dominated by the theoretical uncertainty.

New flavour-specific measurements of semi-inclusive charm yields indicate that the “problem” of the charm deficit and low semileptonic branching ratio is disappearing. The theoretical predictions for $B(\bar{B} \rightarrow X_{c\bar{c}})$ and $B(\bar{B} \rightarrow \text{no charm})$ are confirmed by the data. There is no compelling evidence for violations of quark–hadron duality, nor for new physics or exotic decay modes in B decays. The low value of the A_b lifetime remains a surprise, but can be accommodated in a small corner of parameter space. Lattice calculations of baryon matrix elements of four-quark operators would help to clarify the situation. However, this may also be the first sign of departures from local quark–hadron duality.

Hadronic two-body decays of B mesons with a large energy release can be understood in terms of a generalized factorization hypothesis, which includes the leading nonfactorizable corrections. Finally, new data on B decays into two light mesons open a window to study the details of nonperturbative dynamics of QCD, including long-distance penguins and other exotica. So far, the Standard Model can account for the data, but we should be prepared to find surprises as measurements and theory become more precise. For now, it remains to stress that the observed penguin effects are large and thus raise hopes for having large CP asymmetries in these decays. Hence, there is more *Beauty* to come in a bright future.

References

- [1] For a review and a comprehensive list of references before 1996, see: M. Neubert, Phys. Rep. **245**, 259 (1994); Int. J. Mod. Phys. A **11**, 4173 (1996).
- [2] G. Amorós, M. Beneke and M. Neubert, Phys. Lett. B **401**, 81 (1997).
- [3] A. Czarnecki and A.G. Grozin, Phys. Lett. B **405**, 142 (1997).
- [4] A.G. Grozin and M. Neubert, Nucl. Phys. B **508**, 311 (1997).
- [5] A.F. Falk, M. Luke and M.J. Savage, Phys. Rev. D **53**, 6316 (1996).
- [6] M. Gremm et al. Phys. Rev. Lett. **77**, 20 (1996).
- [7] V. Chernyak, Nucl. Phys. B **457**, 96 (1995); Phys. Lett. B **387**, 173 (1996).
- [8] M. Gremm and I. Stewart, Phys. Rev. D **55**, 1226 (1997).
- [9] H. Li and H.-L. Yu, Phys. Rev. D **55**, 2833 (1997).
- [10] M. Neubert, Phys. Lett. B **264**, 455 (1991); **338**, 84 (1994).
- [11] The theoretical error in the prediction for $\mathcal{F}(1)$ has been reanalysed in *The BABAR Physics Books*, Chapter 8, to appear as a SLAC Report.

- [12] A. Czarnecki, Phys. Rev. Lett. **76**, 4124 (1996); A. Czarnecki and K. Melnikov, Nucl. Phys. B **505**, 65 (1997).
- [13] J. Franzkowski and J.B. Tausk, Preprint MZ-TH/97-37 [hep-ph/9712205].
- [14] M.E. Luke, Phys. Lett. B **252**, 447 (1990).
- [15] N. Uraltsev, Nucl. Phys. B **491**, 303 (1997); A. Czarnecki, K. Melnikov and N. Uraltsev, Preprint TPI-MINN-97-19 [hep-ph/9706311].
- [16] C.G. Boyd, B. Grinstein and R.F. Lebed, Phys. Lett. B **353**, 306 (1995); Nucl. Phys. B **461**, 493 (1996); Phys. Rev. D **56**, 6895 (1997).
- [17] I. Caprini and M. Neubert, Phys. Lett. B **380**, 376 (1996).
- [18] I. Caprini, L. Lellouch and M. Neubert, Preprint CERN-TH/97-91 [hep-ph/9712417].
- [19] P. Drell, Preprint CLNS-97-1521 [hep-ex/9711020], to appear in the Proceedings of the 18th International Symposium on Lepton-Photon Interactions, Hamburg, Germany, July 1997.
- [20] L.K. Gibbons, Proceedings of the 28th International Conference on High Energy Physics, Warsaw, Poland, July 1996, edited by Z. Ajduk and A.K. Wroblewski (World Scientific, Singapore, 1997), p. 183.
- [21] ALEPH Coll. (D. Buskulic et al.), Phys. Lett. B **395**, 373 (1997).
- [22] CLEO Coll. (M. Athanas et al.), Phys. Rev. Lett. **79**, 2208 (1997).
- [23] CLEO Coll. (J.E. Duboscq et al.), Phys. Rev. Lett. **76**, 3898 (1996); CLEO Coll. (A. Anastassov et al.), CLEO CONF 96-8, contributed paper to the 28th International Conference on High Energy Physics, Warsaw, Poland, July 1996.
- [24] A.K. Leibovich et al., Phys. Rev. Lett. **78**, 3995 (1997); Phys. Rev. D **57**, 308 (1998).
- [25] M. Neubert, Preprint CERN-TH/97-240 [hep-ph/9709327], to appear in Phys. Lett. B.
- [26] CLEO Coll. (J.P. Alexander et al.), Phys. Rev. Lett. **77**, 5000 (1996).
- [27] UKQCD Coll. (J.M. Flynn et al.), Nucl. Phys. B **461**, 327 (1996); for a review, see: J.M. Flynn and C.T. Sachrajda, Preprint SHEP-97-20 [hep-lat/9710057], to appear in *Heavy Flavours*, Second Edition, edited by A.J. Buras and M. Linder (World Scientific, Singapore).
- [28] A. Khodjamirian et al., Phys. Lett. B **410**, 275 (1997).
- [29] P. Ball and V.M. Braun, Phys. Rev. D **55**, 5561 (1997); E. Bagan, P. Ball and V.M. Braun, Preprint NORDITA-97-59 [hep-ph/9709243].
- [30] L. Lellouch, Nucl. Phys. B **479**, 353 (1996).
- [31] C.G. Boyd, B. Grinstein and R.F. Lebed, Phys. Rev. Lett. **74**, 4603 (1995); C.G. Boyd and M.J. Savage, Phys. Rev. D **56**, 303 (1997).
- [32] G. Burdman and J. Kambor, Phys. Rev. D **55**, 2817 (1997).
- [33] L.K. Gibbons, to appear in the Proceedings of the 7th International Symposium on Heavy Flavour Physics, Santa Barbara, California, July 1997.
- [34] V. Barger, C.S. Kim and R.J.N. Phillips, Phys. Lett. B **251**, 629 (1990).

- [35] J. Dai, Phys. Lett. B **333**, 212 (1994).
- [36] A.O. Bouzas and D. Zappala, Phys. Lett. B **333**, 215 (1994).
- [37] C. Greub and S.-J. Rey, Phys. Rev. D **56**, 4250 (1997).
- [38] R.D. Dikeman and N.G. Uraltsev, Preprint TPI-MINN-97-06-T [hep-ph/9703437]; I. Bigi, R.D. Dikeman and N. Uraltsev, Preprint TPI-MINN-97-21-T [hep-ph/9706520].
- [39] A.F. Falk, Z. Ligeti and M.B. Wise, Phys. Lett. B **406**, 225 (1997).
- [40] C. Jin, Preprint DO-TH-97-27 [hep-ph/9801230].
- [41] M. Neubert, Phys. Rev. D **49**, 3392 and 4623 (1994).
- [42] I.I. Bigi et al., Int. J. Mod. Phys. A **9**, 2467 (1994).
- [43] E. Bagan et al., Nucl. Phys. B **432**, 3 (1994); Phys. Lett. B **342**, 362 (1995) [E: **374**, 363 (1996)]; E. Bagan et al., Phys. Lett. B **351**, 546 (1995).
- [44] M. Neubert and C.T. Sachrajda, Nucl. Phys. B **483**, 339 (1997).
- [45] M. Feindt, these Proceedings.
- [46] A. Lenz, U. Nierste and G. Ostermaier, Phys. Rev. D **56**, 7228 (1997).
- [47] G. Buchalla, I. Dunietz and H. Yamamoto, Phys. Lett. B **364**, 188 (1995).
- [48] CLEO Coll. (T.E. Coan et al.), Preprint CLNS-97-1516 [hep-ex/9710028].
- [49] I. Dunietz et al., Eur. Phys. J. C **1**, 211 (1998).
- [50] A.L. Kagan and J. Rathsmann, Preprint [hep-ph/9701300]; A.L. Kagan, these Proceedings.
- [51] T. Browder, to appear in the Proceedings of the 7th International Symposium on Heavy Flavor Physics, Santa Barbara, California, July 1997.
- [52] F.E. Close et al., Preprint RAL-97-036 [hep-ph/9708265].
- [53] A. Kagan, Phys. Rev. D **51**, 6196 (1995).
- [54] L. Roszkowski and M. Shifman, Phys. Rev. D **53**, 404 (1996).
- [55] CLEO Coll., CLEO CONF 97-26, contributed paper eps97-337 to this Conference.
- [56] ALEPH Coll., contributed paper pa05-060 to the 28th International Conference on High Energy Physics, Warsaw, Poland, July 1996.
- [57] DELPHI Coll., contributed paper pa01-108 to the 28th International Conference on High Energy Physics, Warsaw, Poland, July 1996.
- [58] DELPHI Coll., DELPHI 97-80 CONF 66, contributed paper eps97-448 to this Conference; P. Kluit, these Proceedings.
- [59] M. Douadi, these Proceedings.
- [60] T. Junk, to appear in the Proceedings of the 2nd International Conference on B Physics and CP Violation, Honolulu, Hawaii, March 1997.
- [61] M. Beneke, G. Buchalla and I. Dunietz, Phys. Rev. D **54**, 4419 (1996).
- [62] B. Blok et al., Phys. Rev. D **49**, 3356 (1994) [E: **50**, 3572 (1994)]; I.I. Bigi et al., in: B Decays, edited by S. Stone, Second Edition (World Scientific, Singapore, 1994), p. 134.
- [63] J.L. Rosner, Phys. Lett. B **379**, 267 (1996).

- [64] P. Colangelo and F. De Fazio, Phys. Lett. B **387**, 371 (1996).
- [65] M. Neubert, Preprint CERN-TH/97-148 [hep-ph/9707217], to appear in the Proceedings of the 2nd International Conference on B Physics and CP Violation, Honolulu, Hawaii, March 1997.
- [66] M.S. Baek, J. Lee, C. Liu and H.S. Song, Preprint SNUTP-97-064 [hep-ph/9709386].
- [67] DELPHI Coll. (P. Abreu et al.), DELPHI 95-107 PHYS 542, contributed paper to the 28th International Conference on High Energy Physics, Warsaw, Poland, July 1996.
- [68] M. Shifman, in *Particles, Strings and Cosmology*, Proceedings of the Joint Meeting of the International Symposium on Particles, Strings and Cosmology and the 19th Johns Hopkins Workshop on Current Problems in Particle Theory, edited by J. Bagger et al. (World Scientific, Singapore, 1996), p. 69.
- [69] B. Blok, M. Shifman and D.-X. Zhang, Preprint TPI-MINN-13-97-T [hep-ph/9709333].
- [70] B. Grinstein and R.F. Lebed, Preprint UCSD/PTH 97-20 [hep-ph/9708396].
- [71] B. Chibisov et al., Int. J. Mod. Phys. A **12**, 2075 (1997).
- [72] P. Colangelo, C.A. Dominguez and G. Nardulli, Phys. Lett. B **409**, 417 (1997).
- [73] G. Altarelli et al., Phys. Lett. B **382**, 409 (1996).
- [74] J.D. Bjorken, Nucl. Phys. B (Proc. Suppl.) **11**, 325 (1989).
- [75] For a review, see: G. Buchalla, A.J. Buras and M.E. Lautenbacher, Rev. Mod. Phys. **68**, 1125 (1996).
- [76] H.-Y. Cheng, Phys. Lett. B **335**, 428 (1994); H.-Y. Cheng and B. Tseng, Preprint IP-ASTP-04-97 [hep-ph/9708211].
- [77] J.M. Soares, Phys. Rev. D **51**, 3518 (1995).
- [78] M. Neubert and B. Stech, Preprint CERN-TH/97-99 [hep-ph/9705292], to appear in *Heavy Flavours*, Second Edition, edited by A.J. Buras and M. Lindner (World Scientific, Singapore); M. Neubert, Preprint CERN-TH/97-169 [hep-ph/9707368], to appear in the Proceedings of the High-Energy Physics Euroconference on Quantum Chromodynamics (QCD 97), Montpellier, France, July 1997.
- [79] T.E. Browder, K. Honscheid and D. Pedrini, Ann. Rev. Nucl. Part. Sci. **46**, 395 (1996).
- [80] J.L. Rodriguez, to appear in the Proceedings of the 2nd International Conference on B Physics and CP Violation, Honolulu, Hawaii, March 1997.
- [81] D. Miller, these Proceedings.
- [82] D. Besson, these Proceedings.
- [83] S. Narison, Phys. Lett. B **322**, 247 (1994).
- [84] T. Huang and C.W. Luo, Phys. Rev. D **53**, 5042 (1996).

- [85] CLEO Coll. (R. Godang et al.), Preprint CLNS-97-1522 [hep-ex/9711010].
- [86] CLEO Coll., CLEO CONF 97-22, contributed paper eps97-333 to this Conference; CLEO CONF 97-13 (1997).
- [87] A.S. Dighe, M. Gronau and J.L. Rosner, Phys. Rev. Lett. **79**, 4333 (1997).
- [88] For a review, see: R. Fleischer, Int. J. Mod. Phys. A **12**, 2459 (1997).
- [89] A. Ali and C. Greub, Preprint DESY-97-126 [hep-ph/9707251]; A. Ali et al., Preprint DESY-97-235 [hep-ph/9712372].
- [90] N.G. Deshpande, B. Dutta and S. Oh, Preprint OITS-641 [hep-ph/9710354].
- [91] A.J. Buras and R. Fleischer, Phys. Lett. B **341**, 379 (1995).
- [92] M. Ciuchini et al., Nucl. Phys. B **501**, 271 (1997); M. Ciuchini et al., Preprint CERN-TH/97-188 [hep-ph/9708222].
- [93] J.D. Bjorken, Preprint SLAC-PUB-7521 [hep-ph/9706524], to appear in the Proceedings of the 2nd International Conference on *B* Physics and CP Violation, Honolulu, Hawaii, March 1997.
- [94] A.J. Buras, R. Fleischer and T. Mannel, Preprint CERN-TH/97-307 [hep-ph/9711262].
- [95] M. Neubert, Preprint CERN-TH/97-342 [hep-ph/9712224].
- [96] A.F. Falk et al., Preprint JHU-TIPAC-97018 [hep-ph/9712225].
- [97] D. Atwood and A. Soni, Preprint [hep-ph/9712287]; these Proceedings.
- [98] D. Atwood and A. Soni, Phys. Lett. B **405**, 150 (1997).
- [99] W.-S. Hou and B. Tseng, Preprint HEPPH-9705304 [hep-ph/9705304]; X.-G. He, W.-S. Hou and C.S. Huang, Preprint [hep-ph/9712478].
- [100] H. Fritzsch, Preprint CERN-TH-97-200 [hep-ph/9708348].
- [101] A.L. Kagan and A.A. Petrov, Preprint UCHEP-27 [hep-ph/9707354].
- [102] I. Halperin and A. Zhitnitsky, Phys. Rev. D **56**, 7247 (1997); Preprint HEPPH-9705251 [hep-ph/9705251].
- [103] E.V. Shuryak and A.R. Zhitnitsky, Preprint NI-97033-NQF [hep-ph/9706316].
- [104] F. Yuan and K.-T. Chao, Phys. Rev. D **56**, 2495 (1997).
- [105] A.A. Petrov, Preprint JHU-TIPAC-97016 [hep-ph/9712497].
- [106] M.R. Ahmady, E. Kou and A. Sugamoto, Preprint RIKEN-AF-NP-274 [hep-ph/9710509].
- [107] D. Du, C.S. Kim and Y. Yang, Preprint BIHEP-TH-97-15 [hep-ph/9711428]; D. Du and M. Yang, Preprint BIHEP-TH-97-17 [hep-ph/9711272].
- [108] CLEO Coll., CLEO CONF 97-13, contributed paper to this Conference.
- [109] H.J. Lipkin, Preprint ANL-HEP-CP-97-45A [hep-ph/9708253].
- [110] H.-Y. Cheng and B. Tseng, Preprint IP-ASTP-03-97 [hep-ph/9707316].
- [111] A. Datta, X.G. He and S. Pakvasa, Preprint UH-511-864-97 [hep-ph/9707259].