CM-P00056817

INTERPRETATION OF PION-NUCLEON SCATTERING RESULTS

William M. Layson

CERN - Geneva

A B S T R A C T

Recognizing that present information on pion-nucleon scattering in the range 300 - 1500 MeV is insufficient for a complete and unambiguous analysis in terms of angular momentum amplitudes, a study has been made employing restrictive assumptions related to the higher resonances. The π^-p angular distribution data are found consistent with resonances of $(T,J) = (1/2, 3/2^-)$ and $(1/2, 5/2^+)$ at 600 and 900 MeV. A plausible, but non-unique, set of amplitudes has been determined, and the only significant departure from a simple behaviour is in the amplitude for the $(1/2, 1/2^+)$ state, which could contain a resonance at 900 MeV. The $\Lambda^0 K^0$ production cross-section shows features in agreement with such a resonance. A proposed phenomenological model based on the important role of the $(3/2, 3/2^+)$ pion-nucleon state predicts the existence of eight resonances below 1500 MeV. Present experimental results are compared with the model and suggestions are made regarding possible tests.

I. INTRODUCTION

This paper is concerned with pion-nucleon collisions at intermediate energies, 300 - 1500 MeV. The term intermediate here excludes the energy range dominated by the first resonance designated by $(T, J) = (3/2, 3/2^+)$ ¹⁾, as well as the range of energies above 1500 MeV where there is little structure apparent in the cross-sections and where one can hopefully apply various high-energy approximations. In the middle range there is structure in the total cross-section and the differential cross-section is not adequately described by either the optical model or by the dominance of a single state.

Peierls ²⁾ in his analysis of pion photoproduction suggested that the $T = 1/2$ peaks at 600 and 900 MeV in scattering ³⁾ should be assigned the quantum numbers $(1/2, 3/2^-)$ and $(1/2, 5/2^+)$ respectively. Although this assignment is not completely established there is no contradictory evidence. Furthermore, Peierls presented an explanation for his result in terms of a model involving the $(3/2, 3/2^+)$ isobar. An extension of this model is developed in the present paper after a study of the available pion-nucleon scattering data. The spirit is then one of interpretation starting from observations rather than one of prediction from a basic theory.

Surveying the presently available experimental results, one finds that the differential elastic scattering cross-section $(\frac{d\sigma}{d\Omega})$ has been measured at some 20 points between 350 and 1200 MeV ⁴⁾, and the total cross-section for $\pi^\pm p$ has been determined to an accuracy of $\sim 5\%$. Cronin ⁵⁾ has calculated, using the forward dispersion relations, the real part of the forward scattering amplitude, which is an independent piece of information at a specific energy, since it depends on total cross-section at all energies. More recently, the recoil proton polarization has been measured at 90° in the c.m.s. for several energies near the higher resonances ⁶⁾.

In the range under consideration no complete phase shift analysis has been performed, because the amount of available information is insufficient to determine the phase shifts in this region of inelastic as well as elastic scattering ⁷⁾. It is possible to learn something about the contributions of different angular momentum states from the cosine power series expansion of $\frac{d\sigma}{d\Omega}$ relative to the angle in the c.m.s.:

$$\frac{d\sigma}{d\Omega} = \sum_{m=0}^{m_m} a_m \cos^m \theta \quad (\text{I.1})$$

Wood et al. ⁸⁾ and Layson ⁹⁾ have presented graphs of the a_n plotted against energy. Using this method it has been generally found that at no energy below 1050 MeV has $n_m > 6$ been required to fit the data. Since n_m is determined by the state of maximum angular momentum (L) involved in the scattering, one assumes that only $L \leq 3$ need be considered. From $\frac{d\sigma}{d\Omega}$ it is then possible to get not more than 7 experimental coefficients, a_n , to be fitted by the amplitudes of $(2L+1) \leq 7$ angular momentum states. Each such state can be completely described by a single complex phase shift, i.e., 14 real quantities in all. Of course, this reduces to 7 in the case of purely elastic processes, but there remains considerable ambiguity in solving for 7 phase shifts from 7 expansion coefficients or from $\frac{d\sigma}{d\Omega}$ directly ¹⁰⁾. In the general case the total cross-section (or imaginary part of forward scattering amplitude via optical theorem) and the real part of the forward scattering amplitude, $D_-(0)$, are measurements which give additional information since only the sum $[\sigma_T(4\pi\lambda)^{-1}]^2 + [D_-(0)]^2$ is known from $\frac{d\sigma}{d\Omega}$. At present one has 9 quantities to be fit by 14 unknowns, though eventually polarization measurements could give six more expansion coefficients, which would raise to 15 the number of experimental quantities. However, even if all this information were available, the difficulties of finding a solution would be formidable in view of the intrinsic ambiguities and the experimental uncertainties. Furthermore, the data at most energies do not give a very reliable indication as to the value of n_m . It should be added that

here only π^-p scattering is considered since much less information is available for π^+p or charge exchange processes, and at present there is little hope of finding the separate isospin components which make up each π^-p amplitude as follows :

$$A_{\pi^-p} = \frac{1}{3} \left(A_{T=3/2} + 2 A_{T=1/2} \right) . \quad (I.2)$$

The preceding paragraph makes one pessimistic toward the probability of understanding pion-nucleon collisions above 300 MeV, but the importance of the π -N interaction in general and the intriguing existence of the peaks in cross-section, demand that one get as much physics as possible from the abundant data, even though they are insufficient for a complete phase shift analysis. This is the purpose of the work reported here. More specifically, it is desired to answer the following questions :

- 1) are the two resonances indicated by the photoproduction analysis consistent with the π^-p data on $\frac{d\sigma}{d\Omega}$, σ_T and $D_-(0)$?
- 2) can one find a plausible set of scattering amplitudes, and do they show any new resonances or unusual behaviour ?
- 3) can the data for $\Lambda^0 K^0$ production be related to those for π^-p scattering ?
- 4) is there some model which explains or correlates the data analysis ?

The process of answering these questions is divided into several parts : Section II treats the elastic differential cross-section, Section III relates the $\Lambda^0 K^0$ production to the elastic scattering, Section IV is concerned with the total cross-section for π^+p , and Section V develops a phenomenological model for explaining the possible resonant states involved in the pion-nucleon collisions.

II. INTERPRETATION OF THE EXPERIMENTAL DIFFERENTIAL CROSS-SECTION

The procedure for answering the questions above begins with the expansion coefficients of Eq. (I.1). To interpret them it is necessary to know their dependence on the angular momentum amplitudes :

$$A_{LJ} = A_{L, L \pm 1/2} \equiv A_{L \pm} \equiv \frac{\gamma_{L \pm} e^{2i\delta_{L \pm}} - 1}{2i} \quad (\text{II.1})$$

where $\delta_{L \pm}$ is the real phase shift and $\gamma_{L \pm}$ is the real elasticity parameter such that $0 \leq \gamma_{L \pm} \leq 1$ (one could use an imaginary phase shift, $-\ln \eta$). The non-spin-flip and spin-flip amplitudes are respectively :

$$\begin{aligned} f(\theta) &= \lambda \sum_L [(L+1)A_{L+} + LA_{L-}] P_L(\cos \theta) \\ g(\theta, \phi) &= \lambda \sum_L [A_{L+} - A_{L-}] P'_L(\cos \theta) e^{i\phi} \end{aligned} \quad (\text{II.2})$$

where ϕ is the azimuthal angle, λ is the wave length and the $P_L^M(\cos \theta)$ are Legendre polynomials. In terms of the $A_{L \pm}$, the total cross-section and the real part of the forward scattering amplitude are given by :

$$D_-(0) + i \frac{\sigma_t}{4\pi\lambda} = \lambda \sum_L [(L+1)A_{L+} + LA_{L-}] \quad (\text{II.3})$$

Using Eq. (I.1) and the fact that $\frac{d\sigma}{d\Omega} = |f(\theta)|^2 + |g(\theta, \phi)|^2$, the a_n can be written :

$$a_m = \lambda^2 \operatorname{Re} \sum_{i \leq j} b_{ijn} A_i A_j^* \quad (\text{II.4})$$

where the b_{ijn} result from simple algebraic manipulation and are given in Table I.

Returning now to the "underdetermined" character of this problem mentioned in the introduction, one must rely on assumptions or educated guesses about the behaviour of the $A_{L\pm}$ if seven of them are to be treated using only nine experimental numbers. The value of equation (II.4) is clear from Table I, since it enables one to relate the observed behaviour of the a_n to the variation of a few amplitudes, $A_{L\pm}$, at least in the cases of a_5 and a_6 . Consider, for example, the zero or small negative values of a_6 found by a number of experimenters⁸⁾ at energies near 900 MeV. Assuming that the choice of n_m has been made correctly, this is a strong indication that $|A_{3+}|$ is small, though a fortuitous cancellation of two terms is possible.

From Table I one finds :

$$a_6 = \lambda^2 \left\{ 43.75 |A_{3+}|^2 + 262.5 [\operatorname{Re} A_{3-} - \operatorname{Re} A_{3+} + \operatorname{Im} A_{3-} - \operatorname{Im} A_{3+}] \right\}. \quad (\text{II.5})$$

By definition one has :

$$-\frac{1}{2} \leq \operatorname{Re} A_{L\pm} \leq \frac{1}{2}, \quad 0 \leq \operatorname{Im} A_{L\pm} \leq 1. \quad (\text{II.6})$$

Thus Eq. (II.5) shows that since $a_6 \leq 0$, $\operatorname{Re} A_{3-}$ and $\operatorname{Re} A_{3+}$ have different signs if $A_{3+} \neq 0$. In fact, only the data at 1020 actually demands a non-zero a_6 according to Wood et al.⁸⁾, so A_{3+} is probably zero within experimental error below 900 MeV. From this observation and Table I, one concludes that the peak in a_5 at 900 MeV is due to a resonance of $J = 5/2$. Thus the very reasonable assumption that A_{3+} does not have a strange behaviour, which permits a large A_{3+} but always a small a_6 , leads to a significant conclusion. Similar reasoning has been used throughout this work.

To find an answer to question 1) of the introduction, one now assumes $(1/2, 3/2^-)$ and $(1/2, 5/2^+)$ resonances at 600 and 900 MeV, respectively, giving them a Breit-Wigner form :

$$A_{L\pm} = \frac{2}{3} \frac{-\frac{1}{2} \Gamma_{\pi L\pm}}{(E - E_{L\pm}) + i \frac{1}{2} \Gamma_{L\pm}} = -\frac{1}{3} \Gamma_{\pi L\pm} \frac{(E - E_{L\pm}) - i \frac{1}{2} \Gamma_{L\pm}}{(E - E_{L\pm})^2 + \frac{1}{4} \Gamma_{L\pm}^2} \quad (\text{II.7})$$

where $\Gamma_{L\pm}$ is the total width, $\Gamma_{\pi L\pm}$ is the elastic scattering width, $E_{L\pm}$ is the resonant energy, and E is the pion c.m.s. energy. The factor $2/3$ accounts for the fact that the resonance occurs only in the $T = 1/2$ state. The parameters $E_{L\pm}$, $\Gamma_{\pi L\pm}$, and $\Gamma_{L\pm}$ of Eq. (II.7) are chosen to fit the peaks in $T = 1/2$ total cross-section, after an appropriate non-resonant contribution has been subtracted from the experimentally measured values of $\sigma_{T=1/2}$. Detailed discussion of this procedure is found in Section IV. With the above prescription for A_{3-} Eqs. (II.5), (II.7) show immediately that the $3+$ state has a positive phase shift if $A_{3+} \neq 0$, and that it is small ($\leq 6^\circ$) unless the scattering is very inelastic. This same conclusion results from similar reasoning based on the possibility that $n_m = 6$ (though $n_m = 5$ gives a sufficiently good fit below 1020 MeV). In this case a_6 passes from positive to negative at the resonance in good agreement with the expected effect of a resonance in the $3-$ state interfering with a $3+$ term of positive phase shift ⁹⁾. With or without a $\cos^6 \theta$ term in $d\sigma/d\Omega$ at 900 MeV, a value in the range $0^\circ \leq \delta_{3+} \leq 6^\circ$ fits the data.

Continuing with the same method, one can examine a_5 given by :

$$a_5 = \lambda^2 \left\{ 105 \operatorname{Re} A_{2-} A_{3+}^* + 112.5 \operatorname{Re} A_{2+} A_{3-}^* + 45 \operatorname{Re} A_{2+} A_{3+}^* \right\} \quad (\text{II.8})$$

Assuming 2- and 3- resonant states with A_{3+} chosen to fit a_6 , one readily determines $\text{Im } A_{2+}$ at 900 MeV where $\text{Re } A_{3-} = 0$. With the assumed values of A_{2-} , A_{3-} , and A_{3+} , the second term of Eq. (II.8) completely dominates the behaviour of a_5 consistent with the observed peak in a_5 at 900 MeV^{8),9)}. The positive sign and something of the magnitude of $\text{Re } A_{2+}$ can be found from the measured values of a_5 at lower energy. By demanding that A_{2+} behave in a "normal way", i.e., increase smoothly from zero at very low energies to the values specified by the above considerations at higher energies, a rough approximation to this amplitude is found. Of course, there are many ways to fit the data with A_{2-} , A_{2+} , A_{3-} and A_{3+} if the assumptions of the two resonances and "normal behaviour" for the non-resonant amplitudes are not made.

The next step toward an answer of question 1) is to find suitable values of A_0 , A_{1-} and A_{1+} . This is best accomplished using a computer to search for sets of amplitudes which agree with the a_n , δ_t and $D_-(0)$. Since a_5 and a_6 have been used in the selection of the four amplitudes of largest angular momentum, one now has six remaining unknowns and seven pieces of information. Thus, except for the uncertainty in the fitting of $\frac{d\delta}{d\mathcal{A}}$ and in choosing the higher $A_{L\pm}$, one can solve for A_0 , A_{1-} , and A_{1+} . Of the various solutions one hopes to find the correct one by using a trial and error procedure starting from values of $A_{L\pm}$ found at a lower energy (or from a_5 and a_6 as discussed above), and varying them cyclically until a "best fit" is obtained. The criterion is that one minimizes M as given by :

$$M = \sum_{m=0}^{m_m} \left| \frac{a_m^c - a_m}{\Delta a_m} \right|^2 + \left| \frac{\delta^c - \delta}{\Delta \delta} \right|^2 + \left| \frac{D_-(0)^c - D_-(0)}{\Delta D_-(0)} \right|^2 \quad (\text{II.9})$$

where the a_n^c , δ^c , and $D_-(0)^c$ are values calculated from a trial set of $A_{L\pm}$ and the Δa_n , $\Delta \delta$, and $\Delta D_-(0)$ are the uncertainties, directly or indirectly from experiment. In some cases slightly different input quantities give qualitatively different solutions, then the value of M and consistency with neighbouring energies determine the choice. No attempt has been made to study all the phase shift solutions rejected on this basis, but at each energy one consistent solution with a small M has been found.

The source material for the π^-p analysis described in the report is, for the most part, in the form of the coefficient a_n determined by various experimenters⁴⁾. Given the a_n together with ϵ_t and $D_-(0)$, the procedure of the above paragraph leads to solutions for the $A_{L\pm}$. Data at sixteen different energies have been treated, and the results are displayed in Figs. 1,2, where the values are marked by large dots and labelled "preferred". Also shown in these figures are results at thirteen energies which are marked by crosses and labelled "alternate". The reason for this latter nomenclature is that, in a particular case, either a value of n_m different from that chosen by the experimentalist has been used, or the data from a published histogram have been fitted by a simple least squares procedure without taking account of errors properly. With regard to the former reason, one should consider the variation of χ^2 with n_m when fitting data. For a function y measured in p intervals to be $y_1 \dots y_p$, χ^2 is defined by :

$$\chi^2 = \sum_{i=1}^p \left| \frac{y_i^c - y_i}{\Delta y_i} \right|^2 \quad (\text{II.10})$$

where y_i^c is a calculated point and Δy_i is the experimental uncertainty. In the present case the y_i^c are found from the set of a_n which minimizes χ^2 for a given value of n_m . To determine the n_m appropriate for a $\frac{d\epsilon}{d\Omega}$, one divides χ^2 by the number of degrees of freedom, $[p - (n_m + 1)]$, and chooses n_m at the value where

$$\frac{\chi^2}{p - (n_m + 1)}$$

ceases to vary appreciably as n_m increases. This means that the fit improves after this point only as much as one would expect from increasing the number of parameters regardless of the true form of the curve. This method of determining n_m is not always precise, and there appears, in some of the data, apparent inconsistency in n_m found at neighbouring energies. For example, the tabulation

of Wood¹⁰⁾ shows $n_m = 5$ at 427 MeV, while $n_m = 3$ at 600 MeV. In several such cases the present author has found additional sets of a_n with different n_m , and these have been used to find the solutions marked "alternate" in the figures. The term refers to the fact that the experimenter chose a different n_m , but the solution may be better in the sense of consistency with other data. Surveying Figs. 1,2, one sees that, in general, the dots and crosses for the $A_{L\pm}$ have the same appearance as functions of energy, despite the fact that the a_n may show effects which differ for the two cases. This insensitivity to the choice of n_m demonstrates that the solutions depend not only on the data but to a marked degree on the input trial values of $A_{L\pm}$, which in turn depend on the assumption of two resonances and "normal behaviour" of non-resonant phase shifts. The latter assumption is necessary in treating the limited amount of data in hand, but in the future more reliable results will be obtained by conventional methods of analysis in which one fits the measured values of $\frac{dG}{dN}$ and omits assumptions about $A_{L\pm}$. Also the accuracy of the fit to $\frac{dG}{dN}$ would be increased by dealing directly with measured points rather than with the a_n .

The results shown in Figs. 1,2 can now be used in answering the first two questions posed in the introduction. Since the data considered here have been fitted with the resonant amplitudes in such a way that $M \approx 1$, one is justified in saying that the $\pi^- p$ data are consistent with the photoproduction resonances. Moreover, the $A_{L\pm}$ have a smooth behaviour as energy increases, though the states of lower angular momentum show some "scatter".

The most unexpected result is that A_{1-} has a resonant appearance near 900 MeV. The calculated points are insufficient to define a clear peak, but it is evident that A_{1-} behaves quite differently from the other $A_{L\pm}$ except A_{2-} and A_{3-} , which are assumed to contain resonances in the present work. To check this behaviour adequately would require much more data, so one can only recognize here that in this limited analysis the computer found that the best solution showed large values of $\text{Im } A_{1-}$ near 900 MeV. Independent reasons for a resonance identification are given below. Beyond this, there does not seem to be anything very remarkable in Figs. 1,2. For example, $\text{Im } A_0$ increases from zero smoothly and seems to level off at a value of $1/2$ past 600 MeV. This is consistent with a large absorption ($\eta < 1$) for low angular momentum states, though $\text{Re } A_0$ is not zero, so $\eta \neq 0$.

III. INFORMATION FROM $\Lambda^0 K^0$ PRODUCTION

The differential cross-section, polarization, and total production cross-section ($\sigma_{\Lambda K}$) for $\Lambda^0 K^0$ from $\pi^- p$ have been measured in a number of experiments, many of which were designed to search for the cusp at $\Sigma^- K$ threshold^{11),12)}. Interesting features of the data are that they show a pronounced peak in $\sigma_{\Lambda K}$ near 900 MeV and that at this energy s+p waves do not seem sufficient to fit the experimental results; there is probably a $\cos^3\theta$ term in the angular distribution^{11),12)}. A resonant interpretation of $\sigma_{\Lambda K}$ has been suggested by B.T. Feld¹³⁾ to account for the rapid rise from threshold and the relatively strong maximum at incident pion energies of 900 MeV, but it has been difficult to associate this with the suspected $(1/2, 5/2^+)$ $\pi^- p$ resonance, because of the angular momentum barrier which strongly reduces any contribution from high angular momentum states near threshold. More recently, Kanazawa¹⁵⁾ has shown that a collection of older data are consistent with a resonance in either of the two states $J = 1/2, 3/2$ at ~ 950 MeV.

In view of the possible $(1/2, 1/2^+)$ resonance discussed in the previous section, it is interesting to see what agreement can be found with $\sigma_{\Lambda K}$ if one assumes a $\Lambda^0 K^0$ decay mode for the resonance. Fig. 3 shows a fit to the production cross-section from threshold to ~ 1.4 GeV. Curve A is computed as the sum of a smooth background, curve B, and a $(1/2, 1/2^+)$ resonance described by a two channel Breit-Wigner formula (elasticity 0.9) with a total width and energy of 125 MeV and 885 MeV (lab), respectively (70 MeV and 605 MeV, c.m.s.). By way of comparison, curve C shows an attempt to fit the data with an $(1/2, 5/2^+)$ resonance, which takes the unusual shape because of the angular momentum barrier factor¹⁶⁾. One concludes that a p wave (or s wave for even ΛK parity) resonance agrees with the data, but not an F wave. To actually decide which state, if any, is in resonance to produce the peak in $\sigma_{\Lambda K}$ one must analyze the differential cross-section and polarization results.

If $L \leq 1$ this is a simple procedure, but since there is a probable $\cos^3\theta$ term, one must include $L > 1$. Unfortunately, the most recent experimental data ¹¹⁾ are not accurate enough to permit a determination of the production amplitudes if higher waves are present. If only s and p waves are used four solutions can be found, two of these show a dominant $J = 1/2$ state, while the other two show a strong $J = 3/2$ state. In this respect the present results are consistent with those of Kanazawa ¹⁵⁾.

To summarize the information from $\Lambda^0 K^0$ production, one can say that there is strong evidence for a resonance of low angular momentum, but further specification must await improved data. In view of the $\pi^- p$ data, it is tempting to speculate that the production involves a sharp 1^- peak superposed on a predominantly s wave background with small $1+$ and $3-$ amplitudes. The last state would be significant only because of the 900 MeV $\pi^- p$ resonance.

IV. ANALYSIS OF THE TOTAL CROSS-SECTION

In the foregoing discussion only $\pi^- p$ collisions could be considered because of the paucity of data on the $\pi^+ p$ differential cross-section. However, if one is prepared to make a strong assumption, then the total cross-sections available for both $\pi^\pm p$ can be interpreted in a very interesting way. One must assume that the scattering in either isospin state $T = 1/2, 3/2$ changes slowly with energy except for the contributions of particular resonant states. In the case of $T = 1/2$ there is evidence for $(1/2, 3/2^-)$ and $(1/2, 5/2^+)$ resonances at 600 and 900 MeV as discussed earlier. The cross-section $\sigma_{1/2}$ is found using the principle of isospin independence from the cross-sections $\sigma_{\pi^\pm p}$ as follows :

$$\sigma_{1/2} = \frac{3}{2} \sigma_{\pi^- p} - \frac{1}{2} \sigma_{\pi^+ p} . \quad (\text{IV.1})$$

For $T = 3/2$ one has directly :

$$\sigma_{3/2} = \sigma_{\pi^+ p} . \quad (\text{IV.2})$$

Experimentally there are few results on the angular momentum states involved in $\pi^+ p$ scattering at energies above 300 MeV and no particular indication of pure resonances, although there is a broad bump in $\sigma_{\pi^+ p}$ near 1.3 GeV and a small irregularity at 850 MeV ¹⁷⁾. At 200 MeV there is the well-known $(3/2, 3/2^+)$ resonance, which is so beautifully explained by the Chew-Low theory. Above 1.5 GeV $\sigma_{\pi^- p} \approx \sigma_{\pi^+ p} \approx \text{constant}$.

In the spirit of the above assumption, one can divide the total cross-section into pure resonances and an isospin dependent background, which varies slowly with energy. At low energy the background is zero, but at high energy it gives the observed constant cross-section. This separation can be made very simply if the interferences between the resonant and non-resonant scattering is negligible. Although the $\pi^- p$ differential cross-section analysis has

indicated that the $A_{L\pm}$ may have a complex behaviour, the only available theory for the background is the optical model. Fernbach, Serber and Taylor¹⁸⁾ have described its use in nuclear physics, while Bethe and Wilson¹⁹⁾ have applied it to the scattering of pions on carbon. For lack of a better theory, various authors have fitted $\frac{d\sigma}{d\Omega}$ above 1 GeV²⁰⁾ using an optical model potential, though this is not a sensitive test of the model. In this approach one considers a sphere of radius a in which a pion wave is absorbed with probability per unit length K , and in which there is a refracting potential V causing a change in the wave number, $k_1 = (k_{r < a} - k)$. The scattering amplitude is :

$$A_{L\pm} = \frac{1}{2i} \left(e^{(-K + 2ik_1)S_L} - 1 \right) \quad (\text{IV.3})$$

where $2s_L$ is the distance traversed in the sphere by a classical particle at an impact parameter $(L+1/2)k^{-1}$. Thus one has :

$$S_L = \left[a^2 - (L+1/2)^2 k^2 \right]^{1/2} \quad (\text{IV.4})$$

For such a classical approximation to be good the wave number k must be large compared to k_1 and K , and $\frac{1}{k} = \lambda$ must be small compared to a , i.e., the following condition must hold :

$$\begin{aligned} ka &\gg 1 \\ k, k^{-1} &\ll 1 \\ K, K^{-1} &\ll 1 \end{aligned} \quad (\text{IV.5})$$

The total optical cross-section can be calculated from Eqs. (II.3) and (IV.3) :

$$\sigma_{\text{opt}} = 2\pi\lambda^2 \sum_{L=0}^{(L+1/2) < ka} (2L+1) \left(1 - e^{-K S_L} \cos 2k_1 S_L \right) \quad (\text{IV.6})$$

With the resonant contribution found from Eqs. (II.3) and (II.7), the total cross-section is then :

$$\sigma_T = \sigma_{opt} + \sum_{LJ} \pi k^2 (J+1/2) \frac{\Gamma_{\pi L\pm}}{\Gamma_{L\pm}} \frac{\Gamma_{L\pm}^2}{(E_{L\pm} - E)^2 + \frac{\Gamma_{L\pm}^2}{4}} \quad (IV.7)$$

In Figs. 4, 5 one sees the results of fitting formula (IV.7) to the $T = 1/2, 3/2$ cross-sections. Actually plotted is $\sigma_T/8\pi k^2$, a quantity which can be more easily associated with the amplitudes $A_{L\pm}$ than can σ_T . Resonance parameters have been chosen to fit the accurately known $(3/2, 3/2^+)$ phase shift, using Eq. (II.7) and the following definition of $\Gamma_{L\pm}$:

$$\Gamma_{L\pm} = \frac{4mM}{E_{L\pm} + E} \gamma_{L\pm}^2 [ka][V_L(ka)] \quad (IV.8)$$

where m and M are the pion and nucleon masses respectively, k is the pion c.m. momentum, a is the radius of the interaction volume, $v_L(ka)$ is the barrier penetration factor, and $\gamma_{L\pm}$ is the dimensionless reduced width. The energy dependent denominator comes from the relativistic normalization but is slowly varying compared to $(E_{L\pm} - E)$, so it is in practice a small change from the usual Breit-Wigner formula. The resulting values are :

$$a = 1.0 \text{ fermi} \quad \gamma_{L\pm}^2 = 0.133 \quad (IV.9)$$

It has been found that these same values of a and $\gamma_{L\pm}^2$ fit quite well the other peaks, though there is some latitude in choosing $\gamma_{L\pm}^2$ even for fixed a . It seems very unlikely, however, that $\gamma_{L\pm}^2$ could vary more than $\pm 25\%$ from the chosen value if a is fixed at 1.0 fermi. The resonant energies $E_{L\pm}$ and the elasticities, $\Gamma_{\pi L\pm}/\Gamma_{L\pm}$ (assumed to be constant), are the remaining

free parameters. The choice of J for each peak was not arbitrary, since those that are well defined (1, 3 and 5 in Table II) were given the assignment of Peierls consistent with his photoproduction results ²⁾ [see also Omnès and Valladas ²¹⁾], and the others (4, 7 and 8) were assigned according to the phenomenological model to be described in the next section. In the second case, resonance 4 is too poorly defined to allow any particular conclusions, but resonances 7 and 8 are interesting in that the assumption of two peaks agrees with the model and fits the cross-section slightly better than the assumption of a broad single peak with a width perhaps twice as great as the others.

The basic criteria for separating resonant and non-resonant contributions were that the latter be a smooth extrapolation to $E = 0$ of the high energy σ_T and that for a particular state

$$\sigma_{l,J} \leq (J + \frac{1}{2}) 4\pi \lambda^2 \frac{\Gamma_{\pi L \pm}}{\Gamma_{L \pm}} .$$

This last condition is important in the case of the large 900 MeV peak in $\sigma_{1/2}$. If $J = 5/2$ for this peak, then $\Gamma_{\pi L \pm} / \Gamma_{L \pm}$ must be large (~ 0.88) in order to fit the variation from a smooth background (Fig. 4), but this means there is little resonant elastic scattering (~ 5 mb), a statement which is barely consistent with the bump in the $T = 1/2$ inelastic cross-section data at 900 MeV ²¹⁾. If improved data show that there is as much as 10 - 15 mb resonant inelastic scattering (also consistent with present data), then there are two reasonable explanations : firstly, the resonant state may have $J = 7/2$, or, secondly, there may be a second resonance near 900 MeV. In the preceding section one saw that the small value of a_6 made it improbable that there is strong $J = 7/2$ scattering, but there was definite evidence for a $J = 1/2^+$ resonance near 900 MeV. Thus the need of more resonance cross-section for a consistent background elasticity would give added support to the possibility of another resonance.

Eqs. (IV.5) give conditions for the use of the optical model which can now be applied to the parameters used to fit the background scattering. They are given below for $T = 1/2, 3/2$ at two energies where complete data are available.

T	E_{lab}	K	k_1	$\frac{K}{k}$	$\frac{k_1}{k}$	ka
1/2	900	0.50	0.60	0.17	0.21	2.90
3/2	990	0.20	0.40	0.07	0.13	3.07

Thus the conditions are barely met at 900 MeV and the situation is worse at lower energy. This inadequacy emphasizes the fact that a complete phase shift analysis will be needed before truly authoritative statements can be made about the resonances.

One further point that should be checked is the neglect of the interference terms in Eq. (IV.7). For $a = 1.0$ fermi one has $ka = 2.28, 2.90$ at 600, 900 MeV respectively. For $L = 2, 3$ the condition $(L + 1/2) < ka$, which determines L_{max} for the optical model, excludes background amplitudes which could interfere with the resonant amplitudes.

An obvious check of the above analysis of the total cross-section is to try to fit the differential cross-section with a combination of optical model and resonant amplitudes. Unfortunately all such attempts fail, as one sees from the following considerations. In general $\frac{d\sigma}{d\Omega}$ near 900 MeV is characterized by a forward peak which can be fitted with the optical model and a bump in the region ⁸⁾ $90^\circ < \theta < 180^\circ$ with $\frac{d\sigma}{d\Omega} = 0$ at 180° . Since $\left(\frac{d\sigma}{d\Omega}\right)_{\text{optical}}$

essentially vanishes in the backward direction, $\frac{d\sigma}{d\Omega} (\theta > 90^\circ)$ should be given by the resonance contribution. For a single dominant resonance of $L \geq 2$, the $|P_L(x)|^2$ gives a distinctive peak at $\theta = 180^\circ$ which does not agree with the data and cannot be removed by the addition of smaller tail contributions of other resonances. Thus a more complicated set of amplitudes, such as found in Section II, must be used to explain $\frac{d\sigma}{d\Omega}$.

This investigation of the total cross-sections has shown that the separation of a smooth background from the experimentally measured values suggests the existence of a series of resonances. It seems unlikely that the background is due to optical potential scattering, but rather it is a complex combination of slowly varying amplitudes. It is, of course, still meaningful to speak of rapidly varying amplitudes, which produce peaks in total cross-sections, even if the background is not simple.

V. A PHENOMENOLOGICAL MODEL

It has been found that the rather large assortment of possible resonant states, which have been discussed above in connection with the analysis of the various cross-sections, can be correlated by a very simple phenomenological model. It is suggested by the almost complete dominance of the $(3/2, 3/2^+)$ π -N resonance at low energies, and it has been developed under the assumption that the state, which is dominant for a single pion with a nucleon, is the most important sub-state in the states of two or more pions with one nucleon. This idea has already been used by Peierls²⁾ in discussing the $(1/2, 3/2^-)$ and $(1/2, 5/2^+)$ peaks in photoproduction, and several calculations have been made^{22),23)} with the purpose of giving the idea a firm theoretical basis. The difficulties of such calculations will not be discussed here, instead the idea will be explored with the purpose of predicting experimental effects (resonances) other than those considered by previous authors.

The primary assumption of the model is that, of all multiple pion plus nucleon states, only those which possess a high probability of single pion plus nucleon sub-states of quantum numbers $(3/2, 3/2^+)$ will give resonances or isobars, which can be observed in pion-nucleon scattering. One is considering the following types of diagrams :

where N^* is the $(3/2, 3/2^+)$ isobar. It might be thought that many states could contain isobars, but it will be shown that parity conservation and symmetrization of pion wave functions severely restricts the role of N^* . It is appropriate to make several points in connection with this model :

- 1) the $(3/2, 3/2^+)$ isobar, as an unstable product in πN and NN collisions, has been successfully employed by Lindenbaum and Sternheimer²⁴⁾ and by Alles-Borelli et al.²⁵⁾ to explain πN correlations in reactions such as : $\pi + N \rightarrow N + \pi + \pi$.
- 2) the $(3/2, 3/2^+)$ resonance in $\pi - N$ scattering and pion photo-production is well described by the theory of Chew and Low which follows from a pseudovector $\pi - N$ interaction.
- 3) the fact that the higher resonances have the same reduced width as the $(3/2, 3/2^+)$ suggests that there may be a similar mechanism for all the resonances.

The quantitative features of this model are simply found from the quantum mechanical wave function for n free pions and a nucleon in the approximation that the nucleon is fixed. It is not necessary to consider what happens in the interaction volume, but merely to treat the angular momentum and isospin states of the free pions. This is sufficient to tell one which pion-nucleon sub-states have the quantum numbers appropriate for resonant interaction. It is useful to define an "enhancement factor", $f_p(T, J)$, which is the probability that a p wave pion and nucleon are in the $(3/2, 3/2^+)$ sub-state when the multiple pion-nucleon state has quantum numbers (T, J) ²⁶⁾. If more than one value of the probability is possible, then by definition $f_p(T, J)$ denotes the maximum. Symmetrization ensures that $f_p(T, J)$ is the same for all p wave pions, but s wave pions cannot have a resonant interaction.

To find $f_p(T, J)$ one first forms the symmetric wave function of two pions one of which is in the $(3/2, 3/2^+)$ sub-state.

$$S \psi_{T, T_3, J, M}^{(N12)} = (\text{constant}) \left\{ \psi_{T, T_3, J, M}^{(N12)} + \psi_{T, T_3, J, M}^{(N21)} \right\} \quad (V.1)$$

$$\psi_{T,T_3,T,M}^{(N12)} = \left\{ \sum_{t_3 \gamma_3} C_{3/2,1}(T, T_3; \gamma_3, t_3) N_{3/2}^{\gamma_3}(N, 1) \pi^{t_3}(2) \right\} \times \quad (V.2)$$

$$\left\{ \sum_{\mu m} C_{3/2, \ell}(\bar{T}, M; \mu, m) S_{3/2}^{\mu}(N, 1) Y_{\ell}^m(2) R_{\ell}(N12) \right\}.$$

The notation is as follows :

T_3, T_3, t_3 = 3rd components of total, isobar, and pion isospins;

M, μ, m = 3rd components of total, isobar, and pion angular momentum;

ℓ = pion orbital angular momentum.

$C_{\ell_1 \ell_2}(J, M; m_1, m_2)$ is the Clebsch-Gordan coefficient¹⁶⁾, $S_{3/2}^{\mu}(N1)$ and $N_{3/2}^{\gamma_3}(N1)$ are the angular and isospin functions of the isobar, $Y_{\ell}^m(2)$ and $\pi^{t_3}(2)$ are the angular and isospin functions of the pion and $R(N12)$ is the radial wave function for the particles. The symmetrized function $\psi_{T,T_3,J,M}^{(N12)}$ written above must be expanded into states of nucleon plus two pions, in order to determine the terms which are eliminated by symmetrization. This is important only in the case of $\ell = 1$ or two p wave mesons, since in the case of one s wave and one p wave meson no cancellation of terms can occur.

The last statement above becomes more clear if one considers the symmetrization of a general two free pion plus nucleon wave function :

$$\psi(N12) = \chi(N12) R(N12) \oplus (N12) \quad (V.3)$$

where $\chi(N12)$, $R(N12)$, and $\oplus(N12)$ are respectively the isospin, radial, and angular functions. The components of the general unsymmetrized function may always be divided into two parts, symmetric and antisymmetric under exchange of pions. Thus :

$$\begin{aligned}
\psi &= (\chi_S + \chi_A)(R_S + R_A)(\Theta_S + \Theta_A) \\
&= \chi_S R_S \Theta_S + \chi_S R_A \Theta_A + \chi_A R_S \Theta_A + \chi_A R_A \Theta_S \\
&\quad + \text{antisymmetric terms.}
\end{aligned} \tag{V.4}$$

The antisymmetric terms cancel out when one constructs ${}_S\psi$:

$${}_S\psi(N12) = C [\psi(N12) + \psi(N21)] \tag{V.5}$$

where C is a normalization constant. The R_A is perhaps unfamiliar, whereas symmetric isospin and angular functions are commonly used. From $R(N12)$, which is a product of two radial functions, $R_A(N12)$ and $R_S(N12)$ can be found.

$$\begin{aligned}
R(N12) &= R_1(1) R_2(2) \\
R_S(N12) &= \sqrt{1/2} [R_1(1) R_2(2) + R_1(2) R_2(1)] \\
R_A(N12) &= \sqrt{1/2} [R_1(1) R_2(2) - R_1(2) R_2(1)]
\end{aligned} \tag{V.6}$$

In general both R_S and R_A are non-zero, but in this model one is interested in functions for which both pions have a high probability of being in the vicinity of the nucleon, i.e., they are likely to be at the same radius. In this case R_S is the appropriate function, since for $R_1(i) \approx R_2(i)$, R_A is small in the vicinity of the nucleon, and $R_A = 0$ for two pions at exactly the same radius. Thus, neglecting R_A , only the first and third terms of Eq. (V.4) remain in ${}_S\psi$:

$${}_S\psi = C R_S (\chi_S \Theta_S + \chi_A \Theta_A) \tag{V.7}$$

To put Eq. (V.2) in this form, one has only to expand $N_{3/2}^{\gamma_3}$ and $S_{3/2}^{\mu}$ in terms of nucleon and pion functions, then form $\chi_s, \chi_A, \oplus_s$, and $\oplus_A$. In this way the effects of symmetrization are completely accounted for and one can finally determine $f_p(T, J)$ by recombining one pion and the nucleon to form functions $N_{1/2}^{\gamma_3}, N_{3/2}^{\gamma_3}, S_{1/2}^{\mu}$ and $S_{3/2}^{\mu}$, then one adds the squares of the coefficients for the $(3/2, 3/2^+)$ state. This simple method can give no information about the dynamics of the problem, but it does provide a definite measure of the probability of a $(3/2, 3/2^+)$ combination existing in a two pion plus nucleon state.

Since the two terms of Eq. (V.7) are separately possible wave functions, it might be thought that some other combination than the one given would give a larger value of $f_p(T, J)$. In fact, however, it can be shown by actual calculation that in starting from Eq. (V.2) the function with maximum $f_p(T, J)$ is found.

As an example of the calculation described above, consider the state $(1/2, 1/2^+)$. One treats χ and $\oplus$ separately [Eqs. (V.4) and (V.7)] in the initial steps and since both have total quantum number $1/2$ they are similar, so one need develop only $\chi_{1/2}^{1/2}$.

$$\chi_{1/2}^{1/2} = \sqrt{1/2} N_{3/2}^{3/2} \pi^- - \sqrt{1/3} N_{3/2}^{1/2} \pi^0 + \sqrt{1/6} N_{3/2}^{-1/2} \pi^+. \quad (V.8)$$

The functions $\chi_{s1/2}^{1/2}$ and $\chi_{A1/2}^{1/2}$ are found by expanding the functions $N_{3/2}^{\gamma_3}$ into pion plus nucleon functions, and then combining the pions in functions $Z_s^{\gamma_3}$ which are given by :

$$Z_s^{\gamma_3} = \sum_{t_3} C_{11}(s, s_3; s_3 - t_3, t_3) \pi^{s_3 - t_3} \pi^{t_3}. \quad (V.9)$$

The well-known functions Z_s^3 are symmetric if s is 0 or 2 and antisymmetric if s is 1. For the state $\chi_{1/2}^{1/2}$ one finds :

$$\begin{aligned}\chi_{s\frac{1}{2}}^{\frac{1}{2}} &= \sqrt{\frac{2}{3}} \rho Z_0^0 \\ \chi_{A\frac{1}{2}}^{\frac{1}{2}} &= \sqrt{\frac{1}{3}} (\sqrt{\frac{2}{3}} \rho Z_1^0 - \sqrt{\frac{2}{3}} m Z_1^1)\end{aligned}\quad (V.10)$$

The corresponding functions $\textcircled{H}_{s1/2}^{1/2}$ and $\textcircled{H}_{A1/2}^{1/2}$ are then used with $\chi_{s1/2}^{1/2}$ and $\chi_{A1/2}^{1/2}$ to give ${}_s\Psi$ as follows :

$${}_s\Psi = R_s (a \chi_{s\frac{1}{2}}^{\frac{1}{2}} \textcircled{H}_{s\frac{1}{2}}^{\frac{1}{2}} + b \chi_{A\frac{1}{2}}^{\frac{1}{2}} \textcircled{H}_{A\frac{1}{2}}^{\frac{1}{2}}) \quad (V.11)$$

where a and b have been introduced to prove in this case a point made above about the maximum of $f_p(T, J)$. The normalization [Eq. (V.10)] is such that we have the condition :

$$(\frac{2}{3} a)^2 + (\frac{1}{3} b)^2 = 1 \quad (V.12)$$

To find $f_p(T, J)$ one now changes the correctly symmetrized wave function to the isobar plus pion form to give :

$${}_s\Psi_{\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}}^{(N, 12)} = \sum_{T_3, M} D_{T_3, M}(a, b) N_{\frac{3}{2}}^{T_3}(N, 1) \pi_{(2)}^{\frac{1}{2}-T_3} S_{\frac{3}{2}}^{M(N, 1)} Y_{(2)}^{\frac{1}{2}-M} \quad (V.13)$$

+ (terms which are not $(3/2, 3/2)$)

where the coefficients $D_{T_3 M}(a, b)$ are actually rather complicated products of Clebsch-Gordan coefficients. Finally $f_p(T, J)$ is given by :

$$f_p(\frac{1}{2}, \frac{1}{2}) = \left(\sum_{T_3 M} |D_{T_3 M}(a, b)|^2 \right)_{\max} = \left[f'_p(\frac{1}{2}, \frac{1}{2}, a, b) \right]_{\max} \quad (V.14)$$

where the sub-script max indicates that $a + b$ are chosen to make $f_p(T, J)$ as large as possible. As stated earlier, this maximization is automatic when one starts from the pion plus isobar form [Eq. (V.8)]. In the present case this is easily shown using :

$$\frac{d}{da} \left[f'(\frac{1}{2}, \frac{1}{2}, a, b) \right] = 0 \quad (V.15)$$

and $b^2 = 9 - 4a^2$.

By comparing Eqs. (V.11) and (V.13), one finds :

$$f'(\frac{1}{2}, \frac{1}{2}, a, b) = \left(\frac{4a+b}{9} \right)^2 \quad (V.16)$$

this gives :

$$a = b = \sqrt{\frac{9}{5}}.$$

The fact that the two coefficients are equal shows that the two parts of Eq. (V.11) are correctly weighted (2:1) by the derivation from Eq. (V.8). It is clear,

however, that one could start from two pion plus nucleon wave functions and determine a and b without reference to Eq. (V.8). With the values of a, b above one has :

$$f_p(1/2, 1/2) = \frac{5}{9} .$$

$f_p(T, J)$ is given in Table III for those states which have the largest values and therefore, according to the model, could possibly have resonances due to the $(3/2, 3/2^+)$ mechanism.

In order to discuss experimental identification of the resonances, one must interpret the factor $f_p(T, J)$ in terms of a resonance position, but since the model is not a detailed theory, only a qualitative result is possible. In the case of a single pion and nucleon in the $(3/2, 3/2^+)$ state, $f_p(T, J) = 1$, and the peak kinetic energy is $E_{\pi \text{lab}} \approx 180 \text{ MeV}$ or ~ 1 pion mass in the c.m.s. For two p-wave pions and a nucleon in a $(5/2, 5/2^+)$ state, $f_p(T, J)$ is again 1 so the total energy above the nucleon mass should be approximately doubled :

$$\begin{aligned} E(3/2, 3/2^+) &= M + 2.04 m \\ E(5/2, 5/2^+) &= M + 4.08 m \end{aligned} \quad (\text{V.17})$$

For $f_p(T, J) < 1$, $E(T, J)$ should be larger since the strength of the interaction is less [one has in the Chew-Low formula $E_R \approx 1/f^2 \omega_{\text{max}}^2$ where f = the coupling constant, and ω_{max} is the cut-off parameter; in the Born approximation one finds that the phase shift is proportional to f^2]. Without further justification the following formula is suggested :

$$E(T, J) = M + \frac{E_R}{f_p(T, J)} m_p \quad (\text{V.18})$$

where n_p is the number of p-wave pions and E_R is determined from the $(3/2, 3/2^+)$ resonance. Eq. (V.18) predicts that there are five even parity resonances below 1500 MeV (numbers 1, 4, 5, 6, and 8 of Table III). To explain odd parity peaks, it is necessary to add an s wave pion to one or more p-wave pions in a multiple state, but the physical idea is the same. The position of the $(1/2, 3/2^-)$ peak gives an empirical value for the s wave pion energy if a term $n_s E_s$ is added to Eq. (V.18). Furthermore a third term $n_p E_p [1 - f_p(T, J)]$ may be added to account for non-resonant p-wave interaction. This last change is of little importance, but it can be considered a correction which permits one to fit three experimental peaks, $(3/2, 3/2^+)$, $(1/2, 3/2^-)$, $(1/2, 5/2^+)$ with a three parameter formula. The resonance positions of the other states which have large $f_p(T, J)$ are then predictions of this empirical formula :

$$E(T, J) = M + \frac{E_R}{f_p(T, J)} n_p + n_s E_s + n_p E_p [1 - f_p(T, J)] \quad (V.19)$$

where the parameters are :

$$\begin{aligned} E_R &= 2.04 \text{ m} \\ E_s &= 2.1 \text{ m} \\ E_p &= -1.0 \text{ m} . \end{aligned} \quad (V.20)$$

Table III lists the eight resonances expected on the basis of the model.

VI. SUMMARY AND CONCLUSION

The primary purpose of this work has been to answer the four precise questions posed in the introduction. Beyond this, some expansion of the questions has been included with the hope that the experimental suggestion of several resonances noted here for the first time will be considered in the planning of future experiments. The evidence reported is, of course, very tentative, but the possibility of a simple isobar mechanism producing a whole series of resonances is too interesting to be neglected. It should be noted that several authors have made calculations which add theoretical support. Carruthers²²⁾ found a $(3/2, 3/2^-)$ resonance in addition to the $(1/2, 3/2^-)$, Wong and Ross²²⁾ predicted $(1/2, 5/2^+)$, $(3/2, 3/2^+)$ peaks, and Tomozawa²³⁾ suggested a $(1/2, 1/2^+)$ resonance.

The results of the differential cross-section analysis are displayed for good fits to the a_n in the plots of $A_{L\pm}$ versus energy (Figs. 1,2). They confirm that $(1/2, 3/2^-)$ and $(1/2, 5/2^+)$ resonances at 600 and 900 MeV respectively are consistent with the measurements of $\frac{d\sigma}{d\Omega}$, though it is not possible to eliminate other possibilities $\overline{\Gamma}(1/2, 3/2^+)$ and $(1/2, 5/2^-)$, or no pure resonances, etc., on the basis of present information. Moreover, a plausible set of scattering amplitudes varying smoothly with energy between 300 and 1200 MeV has been found. The one variation from "normal behaviour" is in the amplitude A_{1-} , which shows irregularities consistent with a p-wave resonance near 900 MeV. This result is in striking agreement with the heretofore unexplained peak in $\Lambda^0 K^0$ production at this energy.

The analysis of the total cross-section is inherently unable to determine the angular momentum states involved in the scattering, at least if energies are far above threshold. On the other hand, the isospin states responsible for specific features of the data can be found. It has been shown that a system of resonances predicted from some specific assumptions (Section V.) is consistent with the variations from smooth behaviour observed in the $T = 1/2, 3/2$ total cross-sections. Furthermore, the data have been fitted by six resonances described by the Breit-Wigner formula and all having the same reduced width, a fact which indicates that they might be due to the same basic process.

The results quoted in this paper constitute significant circumstantial evidence for the resonance model based on the dominant role of the $(3/2, 3/2^+)$ isobar in multiple pion plus nucleon states. The evidence is in no sense incontrovertible, but the following points are of interest :

- 1) the $(3/2, 3/2^+)_{180}$, $(1/2, 3/2^-)$, and $(1/2, 5/2^+)$ peaks inspired the initial considerations by R.F. Peierls²⁾, but the $(1/2, 1/2^+)$ is an entirely new prediction which agrees with two and one half pieces of experimental evidence; namely, the peak found in A_{1-} , the peak in $\Lambda^0 K^0$ production, and the possible difficulty in explaining the inelastic resonance at 900 MeV.
- 2) the predicted $(3/2, 3/2^+)_{830}$ resonance at 830 MeV is in the correct place to explain the bump in $\pi^+ p$ total cross-section, and there may be a corresponding bump in A_{1+} though this is obscured in $\pi^- p$ scattering.
- 3) the $(3/2, 7/2^+)$ resonance is at the correct energy to explain the maximum in $\pi^+ p$ scattering near 1300 MeV, and it agrees with the positive phase shift found for A_{3+} .
- 4) the $(3/2, 5/2^-)$ resonance is not inconsistent with the $\pi^+ p$ total cross-section, but the $(3/2, 3/2^-)$ state shows no sign of a peak near that of $(1/2, 3/2^+)$ at 600 MeV. If this is a failure of the model, a reasonable explanation is that the transition process from one to two real meson states is inhibited for $\pi^+ p$ as indicated by the $T = 3/2$ cross-section being generally smaller than the $T = 1/2$. Also the $\pi^- \pi^-$ state for both $(3/2, 3/2^-)$ and $(3/2, 3/2^+)_{830}$ has $T = 1$, so one might expect that the formation of the ρ meson would interfere with the enhancement mechanism. One encouraging result is that other analyses²⁷⁾ have shown that the $(3/2, 3/2^-)$ phase shift is positive.

In conclusion three possible experimental methods for determining the validity of the model are suggested. The straightforward approach is to measure the $\pi^- p$ polarization at a number of angles for several energies near 900 MeV with the object of doing a true phase shift analysis. This would probably decide if the $(1/2, 5/2^+)$ resonance exists, and the $(1/2, 1/2^+)$ resonance prediction would be either refuted or admitted possible, though complete confirmation might be difficult because of ambiguities in the analysis. Secondly, one could obtain more accurate differential cross-section and polarization data for $\Lambda^0 K^0$ with the purpose of finding a peak in the $(1/2, 1/2^+)$ production amplitude. Thirdly, one could look for an isobar of quantum numbers $(5/2, 5/2^+)$ with mass of about 1510 MeV [Eq. (V.17)]. Observation of this object in some reaction, such as the one below, would be strong support for the model.

REFERENCES

- 1) T is the symbol for total pion-nucleon isospin, J is that for total pion-nucleon angular momentum, and $\pm$ indicates the total parity of the state.
- 2) R.F. Peierls, Phys.Rev. 118, 323 (1960).
- 3) B.J. Moyer, Rev.Mod.Phys. 33, 367 (1961).
- 4) I. Derado and N. Schmitz, Phys.Rev. 118, 309 (1960);
 J.I. Shonle, Phys.Rev. Letters 5, 156 (1960), UCRL 9362;
 A.R. Erwin and J.K. Kopp, Phys.Rev. 109, 1364 (1958);
 L. Baggett, UCRL 8302;
 S. Bergia, L. Bertocchi, V. Borelli, C. Brautti, L. Chervosani, L. Lavatelli, A. Minguzzi-Ranzi, R. Tosi, P. Waloschek and V. Zoboli, Nuovo Cimento 15, 551 (1960);
 M. Chrétien, J. Leitner, N.P. Samios, M. Schwarz and H. Steinberger, Phys.Rev. 108, 383 (1957);
 L. Bertanza, R. Carrara, A. Drago, P. Frazini, I. Mannelli, P.V. Silvestrini, and P.H. Stokes, Nuovo Cimento 19, 467 (1961);
 L.K. Goodwin, R. Kenney and V. Perez-Mendez, Phys.Rev. Letters 3, 522 (1959), UCRL 9119;
 F. Grard, G. Macleod, L. Montanet, M. Cresti, R. Barloutaud, C. Choquet, J.M. Gaillard, J. Heughebaert, A. Leveque, P. Lehmann, J. Meyer and D. Revel, Nuovo Cimento 22, 193 (1961).
- 5) J.W. Cronin, Phys.Rev. 118, 824 (1960).
- v6) E.F. Beall, B. Cork, P.G. Murphy, W.A. Wenzel, C.M.P. Johnson and L.J. Koester Jr., UCRL 9833.
- 7) Note, however, that some analysis in the lower part of the range was done by : W.D. Walker, J. Davis and W.D. Shepard, Phys.Rev. 118, 1612 (1960).
- 8) C.D. Wood, T.J. Devlin, J.A. Helland, M.J. Longo, B.J. Moyer and V. Perez-Mendez, Phys.Rev. Letters 6, 481 (1961).
- 9) W.M. Layson, thesis, M.I.T. Physics Dept. (1961), (unpublished).

34.

- 10) C.D. Wood, UCRL 9507.
- 11) F. Eisler, P. Franzini, J.M. Gaillard, A. Garfinkel, J. Keren, R. Plano, A. Prodel and M. Schwartz, Rev.Mod.Phys. 33, 436 (1961).
- 12) S.E. Wolf, N. Schmitz, L.J. Lloyd, W. Laskar, F.S. Crawford, J. Button, J.A. Anderson and G. Alexander, Rev.Mod.Phys. 33, 439 (1961); also : F.S. Crawford, private communication to B.T. Feld.
- 13) B.T. Feld, p. 15, Proceedings of the 1958 International Conference on High-Energy Physics, CERN, Geneva (1958).
- 14) J. Steinberger, Proceedings of the 1958 International Conference on High-Energy Physics, CERN, Geneva (1958).
- 15) A. Kanazawa, Phys.Rev. 123, 997 (1961).
- 16) J. Blatt and V.F. Weisskopf, "Theoretical Nuclear Physics", John Wiley & Sons, New York (1952).
- 17) T.J. Devlin, B.C. Barish, W.N. Hess, V. Perez-Mendez and J. Solomon, Phys.Rev. Letters 4, 242 (1960); UCRL 9017; T.J. Devlin, UCRL 9548.
- 18) S. Fernbach, R. Serber and T.B. Taylor, Phys.Rev. 75, 1352 (1949).
- 19) H.A. Bethe and R.R. Wilson, Phys.Rev. 83, 690 (1951).
- 20) For example, R.G. Thomas, UCRL 8965.
- 21) R. Omnès and G. Valladas, Proceedings of the Aix-en-Provence International Conference on Elementary Particles (1961).
- 22) P. Carruthers, Phys.Rev. Letters 4, 303 (1960); W. Wong and M. Ross, Phys.Rev. Letters 3, 398 (1959).
- 23) Y. Tomozawa, Nuclear Physics 28, 160 (1961).
- 24) S.J. Lindenbaum and R.M. Sternheimer, Phys.Rev. 123, 333 (1961).

- 25) V. Alles-Borelli, S. Bergia, E.P. Ferreira and P. Waloschek, Nuovo Cimento 14, 211 (1959).
- 26) In other notation $f_p(T, J)$ is found by projecting out the $(3/2, 3/2)$ combinations using an operator $p_{3/2, 3/2}(N \pi_i)$ in the relation :

$$f_p(T, J) \equiv \langle \pi_1 \cdots \pi_m, N, T, J | P_{3/2, 3/2}(N \pi_i) | \pi_1 \cdots \pi_m, N, T, J \rangle$$

- 27) J.H. Foote, O. Chamberlain, E.H. Rogers and H.M. Steiner, Phys.Rev. Letters 4, 30 (1960).

TABLE I

Tabulation of the b_{ijn}

i	j	n						
		0	1	2	3	4	5	6
0	0	1						
0	1-		2					
0	1+		4					
0	2-	-2		6				
0	2+	-3		9				
0	3-		-9		15			
0	3+		-12		20			
1-	1-	1						
1-	1+	-2		6				
1-	2-		4					
1-	2+		-9		15			
1-	3-	-3		9				
1-	3+	3		-30		35		
1+	1+	1		3				
1+	2-		-10		18			
1+	2+				12			
1+	3-	3		-36		45		
1+	3+	-3		-6		25		
2-	2-	1		3				
2-	2+	3		-36		45		
2-	3-				12			
2-	3+		21		-110		105	
2+	2+	9/4		-9/2		45/4		
2+	3-		45/2		-117		225/2	
2+	3+		9		-30		45	
3-	3-	9/4		-9/2		45/4		
3-	3+	-9/2		207/2		-675/2		525/2
3+	3+	9/4		45/4		-165/4		175/4

TABLE II
Experimental Resonances

Isobar No.	Kinetic Energy		T	J	$\frac{E(T,J)-M_n}{m}$	$(2\frac{M}{m} \gamma^2)$	Elasticity
	c.m.s.	lab.					
1	164	205	3/2	P _{3/2}	2.17	1.78	1.00
3	440	605	1/2	D _{3/2}	4.14	1.78	0.67
4	585	850	3/2	P _{3/2}	5.18	1.78	0.20
5	615	900	1/2	F _{5/2}	5.40	1.78	0.88
7	780	1215	3/2	D _{5/2}	6.58	1.78	0.33
8	880	1420	3/2	F _{7/2}	7.29	1.78	0.33

TABLE III
Predicted Isobars

Isobar	No. of Pions	Total Isospin	Angular Momentum	$f_p(T, J)$	$\frac{E(T, J) - M}{m}$	$E_{\pi \text{ lab}}$
1	1	$3/2$	$P_{3/2}$	1	2.04	180
2	2	$3/2$	$D_{3/2}$	1	4.14	600
3	2	$1/2$	$D_{3/2}$	1	4.14	600
4	2	$3/2$	$P_{3/2}$	13/18	5.10	830
5	2	$1/2$	$F_{5/2}$	2/3	5.46	900
6	2	$1/2$	$P_{1/2}$	5/9	6.46	1185
7	3	$3/2$	$D_{5/2}$	7/9	6.91	1310
8	3	$3/2$	$F_{7/2}$	7/9	7.19	1390

FIGURE CAPTIONS

Figs. 1,2 The π^-p scattering amplitudes for each angular momentum state are plotted against laboratory pion kinetic energy. The real and imaginary parts of A_{L+} deduced from a "least squares fit" of the experimental data are marked by points (•) for "preferred solutions" and by crosses (x) for "alternate solutions" ⁴⁾.

Fig. 3 The cross-section for $\Lambda-K$ production by π^-p is plotted at several points near threshold. Curve A shows $\sigma_{\Lambda K}$ composed of s wave background and p wave resonance contributions. Curve B shows only the former contribution, and curve C shows an attempted fit assuming an F wave resonance. The crosses (x) correspond to recent data ¹¹⁾ while dots (•) represent older results ¹⁴⁾.

Fig. 4 The decomposition of the $T = 1/2$ total cross-section ¹⁷⁾ into resonant and background contributions. $\frac{\sigma}{8\pi\lambda^2}$ is plotted against laboratory and c.m.s. pion kinetic energy.

Fig. 5 The decomposition of the $T = 3/2$ total cross-section ¹⁷⁾.

FIG. 2

Pion Lab Energy (BeV)

FIG. 3

FIG. 4

FIG. 5