CM-P00056908

ON THE SOLUTIONS OF THE BETHE-SALPETER EQUATION

A. Bastai

Istituto di Fisica dell'Università - Torino
Istituto Nazionale di Fisica Nucleare - Torino

L. Bertocchi

Istituto di Fisica dell'Università - Bologna
Istituto Nazionale di Fisica Nucleare - Bologna

G. Furlan *)

CERN - Geneva

M. Tonin

Istituto di Fisica dell'Università - Padova
Istituto Nazionale di Fisica Nucleare - Padova

A B S T R A C T

We study the solutions of the Bethe-Salpeter equation for bound states, in the ladder approximation, in the case of potentials both regular and singular near the origin. Two methods are presented, one in momentum space, the other in co-ordinate space, which allow the complete determination of eigenvalues and eigenfunctions, in the case where both the total energy E of the state and the mass of the exchanged particle are zero.

*) On leave from the Istituto di Fisica, Università di Trieste, Trieste - Italy.

1. INTRODUCTION

The aim of this paper is to discuss the mathematical properties of the bound state solutions of the Bethe-Salpeter equation, in ladder approximation, for certain classes of "potentials". This problem has been treated some years ago by Wick and Cutkosky ¹⁾ and subsequently by Ceolin et al. ²⁾ who were able to give a set of solutions in the case of regular potentials. We mean by this, those cases where the integral equations to be studied can be reduced to the Fredholm type or, in other words, where the singularities of the potential near the origin do not determine the behaviour of the wave function at $R = 0$. On the other hand, we will call singular potentials the cases where the equations are not of the Fredholm type, which corresponds to potentials which enter into the Fuchsian conditions at the origin. We do not consider here potentials for which a power behaviour of the wave function at $R = 0$ is not possible. This classification of the potentials, which is already possible in the non-relativistic case, has been discussed for the relativistic Bethe-Salpeter equation, together with its connection to field theory, in a deeper way in a preceding paper ³⁾. In this work we want to present in some detail the mathematical techniques which allow a unified treatment of both the regular and singular cases. Because the relevant features of the problem are independent of the total energy E of the state and of the mass of the exchanged particle, μ , we consider the simplified case where $E = \mu = 0$. We begin in Section 2 with a discussion of the regular case as an illustration of the method we will apply to the more general problem treated in Section 3. The procedure is based, in both cases, on the transition to momentum space through the expansion of the wave function

$$\varphi(R) = \int_0^\infty \varphi(q) J(qR) dq$$

where the J functions are the eigenfunctions of that part of the Hamiltonian which correctly describes the behaviour of the solution at the origin in co-ordinate space. The relevant point is that this part of the Hamiltonian coincides with the free one, only for regular potentials.

In the other case we have to collect in the free part, with a sort of canonical transformation, all the terms which influence the behaviour of the wave function at $R = 0$. In so doing we arrive, also in the singular case, at a Fredholm type equation which can be solved by standard methods. Section 4 contains a method of solution which works directly in co-ordinate space and which is a generalization of the factorization method for a second order differential equation. Finally in Appendix B, we introduce a more complicated method which has the advantage of an easier generalization to less simple potentials.

2. THE REGULAR CASE

We shall first briefly discuss the problem of the solution of the Bethe-Salpeter equation ⁴⁾ (in ladder approximation) with "regular" potentials, which means potentials which do not enter in the Fuchsian conditions at the origin.

The interest in discussing the regular case is twofold: first it is useful in order to introduce and clarify the method of solution we shall use in the "singular" case: secondly it will be interesting to compare the solutions of the regular and singular cases for those potentials for which exact solutions are obtainable.

As typical "regular" case, we shall consider the problem of two scalar mesons of mass $m = 1$, interacting via a scalar field of mass μ , and giving a bound state of mass $E < 2$.

The eigenvalues for the mass of the bound state can be obtained by solving the integral equation, which is the integral form of the Bethe-Salpeter equation in momentum space ^{*)}

$$\left[\left(p - \frac{E}{2} \right)^2 + 1 \right] \left[\left(p + \frac{E}{2} \right)^2 + 1 \right] \varphi(p) = \frac{g^2}{(2\pi)^4 i} \int \frac{d^4 k \varphi(k)}{(p-k)^2 + \mu^2} \quad (2.1)$$

We insist on the fact that Eq. (2.1) is an eigenvalue equation: in fact Wick ¹⁾ has shown that it is possible to continue the equation from the real to the imaginary energy (p_4) axis, passing therefore to an Euclidean space: it is then possible to show that Eq. (2.1) is a Fredholm integral equation, and therefore an eigenvalue equation.

Eq. (2.1) can be transformed into the fourth order differential equation in co-ordinate space

^{*)} As we have already said, we limit ourselves to the ladder approximation.

$$\left[\left(i \frac{\partial}{\partial x_\mu} - \frac{E}{2} \right)^2 + 1 \right] \left[\left(i \frac{\partial}{\partial x_\mu} + \frac{E}{2} \right)^2 + 1 \right] \psi(x) = g^2 D_F(x, \mu) \psi(x) \quad (2.2)$$

Of course, in order that Eq. (2.2) be equivalent to Eq. (2.1) it must be supplemented with suitable boundary conditions: now also in Eq. (2.2) it has been performed the Wick continuation from the real to the imaginary relative time axis; in this way the metric becomes Euclidean, and the problem is transformed from an hyperbolic to an elliptic one. It is therefore possible to impose boundary conditions for $R \rightarrow 0$ and $R \rightarrow \infty$, where R is the radius of the Euclidean four-dimensional space.

It has been shown by Wick that the boundary conditions, which select the solution of (2.2) equivalent to (2.1), are regularity conditions for $R \rightarrow \infty$, and the condition that among the four independent solutions at the origin $R = 0$ one must retain only the two regular and reject the two singular ones.

Of course, exact solutions of this regular case can be obtained only in very special cases.

A problem which admits an exact solution is the case in which both μ (the mass of the exchanged particle) and E (the mass of the bound state) vanish. This peculiar property is due to the high symmetry of the problem: but we notice that, also if the problem seems to be rather academic (the bound state has four-momentum equal to zero), it has a great interest since, as we shall see in the next Section, the difficulty that is found when one deals with singular potentials is independent both of the mass of the bound state and from the mass of the exchanged field, which generates the potential.

We shall therefore consider the simplified problem of two scalar particles of unit mass, interacting in ladder approximation via a scalar meson of zero mass, and giving a bound state of zero total four-momentum.

This problem has already been treated by Wick ¹⁾ and Cutkosky ¹⁾ and by Ceolin et al. ²⁾, who have obtained the eigenvalues of the problem ^{*)} starting from the integral form of the Bethe-Salpeter equation in momentum space and reducing it to a second order differential equation in momentum space, supplemented with regularity conditions. This equation turns out to be of the Papperitz kind and therefore its solutions can be obtained with standard methods.

We shall see now how the same equation can be obtained in a very simple way, starting from the fourth order differential equation in co-ordinate space.

In our hypothesis, the causal propagator $D_F(x)$ is simply given by

$$D_F(x) = \frac{1}{(2\pi)^4 i} \int d^4 q \frac{e^{iqx}}{q^2} = \frac{1}{4\pi^2 x^2} ; \quad x^2 = \sum_{i=1}^4 x_i^2$$

and the Bethe-Salpeter equation is

$$(\square - 1)^2 \varphi(x) = \frac{\lambda}{x^2} \varphi(x) \quad \lambda = \frac{g^2}{4\pi^2} \quad (2.3)$$

Let us now transform it in a differential equation in momentum space. In Eq. (2.3) the potential λ/x^2 depends only on $R = \sqrt{x^2}$; this fact allows the separation of the equation in four-dimensional polar co-ordinates. We can extract the angular dependence by putting

$$\varphi(x) = \sum_{\alpha \ell m} Y_{\alpha \ell m}(\theta_1, \theta_2, \varphi) \frac{\varphi_\alpha(R)}{R}$$

where $Y_{\alpha \ell m}$ are the hyperspherical harmonic functions ⁵⁾

^{*)} Having fixed the mass of the bound state, we shall consider the coupling constant g^2 as the eigenvalue parameter.

6.

$$Y_{\alpha \ell m}(\theta_1, \theta_2, \varphi) = N_{\alpha \ell m} e^{im\varphi} (\sin \theta_1)^\ell G_{\alpha-\ell}^{\ell+1}(\cos \theta_1)$$

$$P_\ell^m(\cos \theta_2), \quad \alpha \text{ integer } \geq 0, \quad \ell = 0, 1, \dots, \alpha, \quad |m| \leq \ell.$$

The G_μ^ν functions are Gegenbauer polynomials and $N_{\alpha \ell m}$ are normalization constants, which are not of interest for us.

We get

$$\left(\frac{d^2}{dR^2} + \frac{1}{R} \frac{d}{dR} - \frac{\nu^2}{R^2} - 1 \right)^2 \varphi_\alpha(R) = \frac{\lambda}{R^2} \varphi_\alpha(R) \quad (2.4)$$

$$\nu = \alpha + 1$$

The behaviours of $\varphi_\alpha(R)$ for $R \rightarrow 0$ are (except when $\alpha = 0$)

$$R^{-(\alpha+1)}; R^{-\alpha+1}; R^{\alpha+1}; R^{\alpha+3} \quad (2.4')$$

Our boundary condition for $R \rightarrow 0$ selects the two regular solutions

$$\varphi_\alpha^{(1)}(R) \sim R^{\alpha+3}; \quad \varphi_\alpha^{(2)}(R) \sim R^{\alpha+1} \quad (2.4'')$$

Therefore $\varphi_\alpha(R)$ vanishes at the origin at least as $R^{\alpha+1}$. On the other hand, the acceptable behaviours at $R \rightarrow \infty$ are

$$\varphi_\alpha(R) \sim e^{-R} R^{\pm \sqrt{\frac{1+\lambda}{4}}} \quad (2.4''')$$

We can go over to momentum space by multiplying both sides of Eq. (2.4) by R^2 and defining

$$\varphi_\alpha(R) = \int_0^\infty du J_\nu(\sqrt{u} R) \varphi_\alpha(u) \quad (2.5)$$

where $J_{\alpha+1}(\sqrt{u}R)$ is the Bessel function regular at the origin.

The transformed equation now reads

$$\left[4u \frac{d^2}{du^2} + 4 \frac{d}{du} - \frac{\nu^2}{u} \right] (u+1)^2 \varphi_\alpha(u) = -\lambda \varphi_\alpha(u) \quad (2.6)$$

if the following condition is satisfied :

$$\left[p \left((u+1)^2 \varphi_\alpha(u) \frac{d}{dp} J_\nu(\sqrt{u}R) - J_\nu(\sqrt{u}R) \frac{d}{dp} (u+1)^2 \varphi_\alpha(u) \right) \right]_0^\infty = 0 \quad (2.6')$$

which allows the following transformation of the operators

$$\frac{d^2}{dR^2} + \frac{1}{R} \frac{d}{dR} - \frac{\nu^2}{R^2} - 1 \rightarrow -(u+1); \quad R^2 \rightarrow - \left(4u \frac{d^2}{du^2} + 4 \frac{d}{du} - \frac{\nu^2}{u} \right) \quad (2.6'')$$

Putting now

$$(u+1)^2 \varphi_\alpha(u) = u^{\nu/2} \phi_\alpha(u)$$

we can rewrite Eq. (2.6) as

$$\left[\frac{d^2}{du^2} + \frac{\nu+1}{u} \frac{d}{du} + \frac{\lambda}{4u(u+1)^2} \right] \phi_\alpha(u) = 0 \quad (2.7)$$

The conditions for the existence of the Hankel transform (2.5) are

- $\varphi_\alpha(u)$ vanishes at infinity at least as $u^{-1/4-\epsilon}$;
- $\varphi_\alpha(u)$ can have at the origin a singularity of order not higher than $u^{-(\alpha+1)/2-1+\epsilon}$ that is $\lim_{u \rightarrow 0} u^{\alpha+2} \phi_\alpha(u) = 0$.

3.

We must now remember the boundary conditions on $\varphi_\alpha(R)$. At the origin $R = 0$ we have the condition that $\varphi_\alpha(R)$ vanishes at least as $R^{\alpha+1}$. It can be shown (see Appendix A) that this happens if $\varphi_\alpha(u)$ vanishes at infinity at least as $u^{-((\alpha+1)/2 + 1)}$, or more precisely if

$$\lim_{u \rightarrow \infty} u^{\frac{\alpha+1}{2} + 1} \varphi_\alpha(u) = 0 \quad (2.8)$$

At infinity $\varphi_\alpha(R)$ behaves as a decreasing exponential: therefore on $\varphi_\alpha(u)$ we have the condition at the origin that

$$\lim_{u \rightarrow 0} u^{-\frac{\alpha+1}{2} + \varepsilon} \varphi_\alpha(u) = 0 \quad (2.9)$$

We notice that conditions (2.8) and (2.9) are more stringent than the conditions a) and b) on the existence of the transform (2.5), which can therefore be discarded.

We can now observe that the solution of Eq. (2.7) which satisfies conditions (2.8) and (2.9) behaves like

$$\lim_{u \rightarrow \infty} \varphi_\alpha(u) \sim u^{-\nu} \quad (2.10)$$

$$\lim_{u \rightarrow 0} \varphi_\alpha(u) \sim \text{const}$$

and this solution satisfies also the condition (2.6) when $\alpha > -\frac{1}{2}$.

As it is extensively discussed in Ref. ²⁾, the eigenvalues of Eq. (2.7), taking into account conditions (2.10), are given by

$$\lambda_m = 4(\nu + m)(\nu + m + 1) \quad m = 0, 1, 2, \dots \quad (2.11)$$

and the eigenfunction corresponding to the lowest eigenvalue is

$$f_\alpha^{(0)}(u) = \frac{1}{(u+1)^\nu}$$

If we transform back this eigenfunction to co-ordinate space, we obtain

$$\varphi_{\alpha}^{(0)}(R) = \int_0^{\infty} du J_{\nu}(\sqrt{u}R) \frac{u^{1/2}}{(u+1)^{\nu+2}} = \frac{R^{\nu+1} K_1(R)}{2^{\nu} \Gamma(\nu+2)} \quad (2.12)$$

which does in fact satisfy the boundary conditions (2.4') and (2.4'') $\sqrt{K}_1(R)$ is the Bessel function of imaginary argument, which behaves as R^{-1} at $R \sim 0$ and $(e^{-R})/(R^{\frac{1}{2}})$ for $R \rightarrow \infty$.

Two alternative methods to solve Eq. (2.3) will be discussed respectively in Section 4 and Appendix B. The first method works in co-ordinate space and is a generalization of the factorization method of solution ⁶⁾ for a second order differential equation.

The second one consists in obtaining an integral equation of the Fredholm kind for the Bessel transform of the radial part of $\varphi(x)$: this method is particularly useful since it could be easily generalized to the cases in which the potential is not of the simple Coulomb form g^2/R^2 , as for instance when the mass of the exchanged boson does not vanish: it can therefore give the starting point of a series expansion for those classes of potentials for which no exact solutions are available.

3. THE SINGULAR CASE

In the previous Section we have discussed the problem of two scalar particles interacting through a scalar field, which is characterized by the fact that the potential term $D_F(x, \mu)$ does not enter into the Fuchsian conditions at $R = 0$ (regular potentials). Let us turn now to the study of the Bethe-Salpeter equation for scalar particles with potentials which behave at the origin as R^{-4} , as for instance a $D_F^2(x, \mu)$ term. In this last case the behaviour of the function $\varphi(x)$ at $R = 0$ depends on the potential and we are forced to include in that part of the equation, which has to be exactly solved, also the part of the potential which does enter into the indicial equation for $R = 0$.

We start directly with a Bethe-Salpeter equation in the case $E = 0$ $\mu = 0$ because the characteristic features of the problem (the behaviour at $R = 0$ of the wave function) are independent of these terms. After the separation in polar co-ordinates, we get for the radial function

$$O^2 \varphi_\alpha(R) = V(R) \varphi_\alpha(R) \quad (3.1)$$

with

$$0 = \frac{d^2}{dR^2} + \frac{1}{R} \frac{d}{dR} - \frac{V^2}{R^2} - 1 \quad V = \alpha + 1 \quad (3.2)$$

$$V(R) = \frac{\alpha}{R^4} + \bar{V}(R) \quad \lim_{R \rightarrow 0} R^4 \bar{V}(R) = 0$$

As in the regular case we will impose on $\varphi_\alpha(R)$ regularity conditions at the boundaries, namely

$$\lim_{R \rightarrow \infty} \varphi_\alpha(R) = 0 \quad (3.3)$$

$$\lim_{R \rightarrow 0} R^{\epsilon-1} \varphi_\alpha(R) = 0 \quad (3.4)$$

A complete discussion on the indicial equation (which is critically dependent on the coupling constant "a") at $R = 0$ is given in I. : for our purpose it is sufficient to say that the solution will be a linear combination of the two regular solutions at the origin (whose behaviour is R^{γ_1+1} , R^{γ_2+1} ; $\gamma_1, \gamma_2 > 0$) which goes at infinity as a decreasing exponential.

Now the crucial point in the solution of Eq. (3.3) is to find a procedure which generalizes the method discussed in the non-singular case. Namely, we have to include the singular part of the potential in the free term in order to arrive to an equation with regular potentials, similar to the one discussed in Section 2. This has been done in I. so that in this paper we shall simply start from the result of the previous work. Moreover, we will assume, from now on, for $\bar{V}(R)$ the simple form $\bar{V}(R) = b/R^2$. In I. it has been shown that Eq. (3.3) can be written in the form

$$\begin{aligned} \bar{O} \bar{\varphi}_\alpha &= \lambda R^{\beta-2} \bar{\varphi}_\alpha \\ \bar{O} \bar{\varphi}_\alpha &= R^{-\beta} \bar{\varphi}_\alpha \end{aligned} \quad (3.5)$$

or equivalently

$$R^{-\beta} \bar{O} R^\beta \bar{O} \bar{\varphi}_\alpha(R) = \frac{\lambda}{R^2} \bar{\varphi}_\alpha(R) \quad (3.6)$$

where

$$\bar{O} = \frac{d^2}{dR^2} + \frac{1}{R} \frac{d}{dR} - \frac{\bar{V}^2}{R^2} - 1$$

$$\bar{\varphi}_\alpha(R) = R^{-\beta/2} \varphi_\alpha(R)$$

$$\bar{v}^2 = v^2 + \beta - \beta^2/4 \quad (3.7)$$

$$\lambda = b - \beta(\beta - 2)$$

and β is given by

$$\beta = 2 - \left\{ 2[(\alpha+1)^2 + 1] - 2[\alpha^2(\alpha+2)^2 - a]^{1/2} \right\}^{1/2} \quad (3.8)$$

Using Eq. (3.8) we can write explicitly

$$\bar{v} = \left\{ \frac{1 + (\alpha+1)^2 + [\alpha^2(\alpha+2)^2 - a]^{1/2}}{2} \right\}^{1/2} \quad (3.8')$$

Real values for the considered quantities are understood and the coupling constant "a" is restricted to the range :

$$0 < a < \alpha^2(\alpha+2)^2$$

Moreover the behaviour at the origin is determined by $\bar{v}$ and β through the relations

$$\gamma_{1+1} = \bar{v} + 2 - \beta/2 \quad (3.9)$$

$$\gamma_{2+1} = \bar{v} + \beta/2$$

We want only to point out as in the limit $a \rightarrow 0$, $\beta \rightarrow 0$, $\bar{v} \rightarrow v$, $\gamma_{1+1} \rightarrow \alpha+3$, $\gamma_{2+1} \rightarrow \alpha+1$ so that for vanishing singular potentials our treatment will go back, continuously, to the regular case, as it does.

Let us discuss the solution of Eq. (3.5). We substitute for it the equation

$$R^2 R^{-\beta} \bar{O} R^{\beta} \bar{O} \bar{\varphi}_{\alpha}(R) = \lambda \bar{\varphi}_{\alpha}(R) \quad (3.10)$$

which can be rewritten as

$$\left[R^2 \bar{O} \bar{O} + \left(\beta^2 + 2\beta R \frac{d}{dR} \right) \bar{O} \right] \bar{\varphi}_{\alpha}(R) = \lambda \bar{\varphi}_{\alpha}(R) \quad (3.11)$$

As we did in the regular potential problem, we pass now to momentum space by putting

$$\bar{\varphi}_{\alpha}(R) = \int_0^{\infty} du J_{\bar{\nu}}(\sqrt{u} R) \bar{\varphi}_{\alpha}(u) \quad (3.12)$$

Then the operators transform as

$$R^2 \rightarrow -4u \left[\frac{d^2}{du^2} + \frac{1}{u} \frac{d}{du} - \frac{\bar{\nu}^2}{4u^2} \right]$$

$$R \frac{d}{dR} \rightarrow -2u \frac{d}{du} - 2 \quad ; \quad \bar{O} \rightarrow -(\bar{\nu} + 1)$$

and we get the equation

$$\left\{ 4u \left[\frac{d^2}{du^2} + \frac{1}{u} \frac{d}{du} - \frac{\bar{\nu}^2}{4u^2} \right] (u+1)^2 + (\beta^2 - 4)(u+1) - 4\beta u \frac{d}{du} (u+1) \right\} \bar{\varphi}_{\alpha}(u) = -\lambda \bar{\varphi}_{\alpha}(u) \quad (3.13)$$

under the condition

$$\left[b \left(J_{\bar{\nu}} \frac{d}{dp} (u+1)^2 \bar{\varphi}_{\alpha}(u) - (u+1)^2 \bar{\varphi}_{\alpha}(u) \frac{d}{dp} J_{\bar{\nu}}(pR) + 2\beta b (u+1) \bar{\varphi}_{\alpha} J_{\bar{\nu}} \right) \right]_0^{\infty} = 0 \quad (3.13')$$

We now observe that, with our choice of β , the boundary conditions on $\bar{\varphi}_{\alpha}(R)$ mean that $\bar{\varphi}_{\alpha}(R)$ must behave at the origin as $\bar{\varphi}_{\alpha}(R) \sim R^{\nu+1}$ and at infinity as a decreasing exponential: therefore the boundary conditions on $\bar{\varphi}_{\alpha}(u)$ are the same as (2.8) and (2.9), namely :

$$\begin{aligned} \lim_{u \rightarrow \infty} u^{\bar{\nu}/2+1} \bar{\varphi}_{\alpha}(u) &= 0 \\ \lim_{u \rightarrow 0} u^{-\bar{\nu}/2+\varepsilon} \bar{\varphi}_{\alpha}(u) &= 0 \end{aligned} \quad (3.14)$$

By putting now

$$\bar{\varphi}_{\alpha}(u) = u^{\bar{\nu}/2} (u+1)^{\beta/2-2} \bar{f}_{\alpha}(u)$$

we get the equation

$$\left[\frac{d^2}{du^2} + \frac{\bar{\nu}+1}{u} \frac{d}{du} + \frac{b}{4u(u+1)^2} \right] \bar{f}_{\alpha}(u) = 0 \quad (3.15)$$

[remember that $\beta/2(\beta/2-1) + \lambda/4 = b/4$].

The solution of Eq. (3.15) which satisfies conditions (3.14) must behave as

$$\bar{f}_{\alpha}(u) \underset{u \rightarrow \infty}{\sim} u^{-\bar{\nu}} ; \quad \bar{f}_{\alpha}(u) \underset{u \rightarrow 0}{\sim} \text{const.} \quad (3.16)$$

and this solution satisfies, as can be easily verified, also the conditions (3.13') since in all the interval $\alpha^2(\alpha+2)^2 > a > 0$ one has $\bar{\nu} - \beta - \frac{1}{2} > 0$.

Equation (3.15) and conditions (3.16) are formally identical with those encountered in Section 2 [Eqs. (2.7) and (2.10)] with the substitution

$$\bar{\nu} \rightarrow \nu \quad ; \quad b \rightarrow \lambda$$

We can therefore write down immediately the eigenvalues, which turn out to be

$$b_m = 4(\bar{\nu} + m)(\bar{\nu} + m + 1) \quad m = 0, 1, 2, \dots \quad (3.17)$$

and the eigenfunction corresponding to the lowest eigenvalue is

$$\bar{\varphi}_\alpha^{(0)}(u) = \frac{1}{(u+1)\bar{\nu}}$$

By retransforming this eigenfunction back to co-ordinate space, we obtain

$$\varphi_\alpha(R) = \frac{R^{\bar{\nu}+1} K_{1-\beta/2}(R)}{2^{\bar{\nu}-\beta/2} \Gamma(\bar{\nu}+2-\beta/2)} \quad (3.18)$$

which does indeed satisfy the right boundary conditions.

At this point the convenience of the choice for β ($\beta \rightarrow 0$, $a \rightarrow 0$) can be easily understood; in fact it allows to reduce the problem of solving Eq. (3.3) to a problem whose solution is already known, namely Eq. (3.15).

By looking at the form of the eigenvalues (3.17), we notice that we encounter here a situation which resembles the non-relativistic one: the form of the eigenvalues is the same as for a regular potential, and

the only effect of the singular potential is a modification of the centrifugal index, $\nu \rightarrow \bar{\nu}$. It is evident that in the absence of the b/R^2 potential there are no solutions for whichever a : this corresponds to the fact that the singular potential only redefines the centrifugal one and in this case we cannot have wave functions satisfying boundary conditions at the origin and at infinity at the same time, as in the Goldstein ⁷⁾ problem.

Also if we consider the problem of the bound states of two spin $\frac{1}{2}$ particles, we are faced with the difficulties deriving from a singular potential. In fact, if the interaction between the two fermions is mediated by a photon or a scalar meson, we have a second order differential equation with an $1/R^2$ potential ⁷⁾. We do not enter into details here: anyway we discuss a very particular case in order to get an insight into the problem.

In the electromagnetic potential case the Bethe-Salpeter equation will be of the form ($E = 0$ and ladder approximation)

$$(i \gamma \cdot \vec{\partial} - m) \psi(x) (i \gamma \cdot \overleftarrow{\partial} - m) = e^2 D_F(x) \gamma_\mu \psi(x) \gamma_\mu \quad (3.19)$$

If we put

$$\psi(x) = R^2 \varphi(x)$$

the equation for $\varphi(x)$ is

$$(i \gamma \cdot \vec{\partial} - m) R^2 \varphi(x) (i \gamma \cdot \overleftarrow{\partial} - m) = \lambda \gamma_\mu \varphi(x) \gamma_\mu$$

$$\lambda = \frac{e^2}{4\pi^2}$$

which in momentum space reads

$$(\not{p} + m) (\not{D}_p \varphi(p)) (\not{p} + m) = \lambda \gamma_\mu \varphi(p) \gamma_\mu \quad (3.20)$$

We do not try in this paper to find all the solutions of this equation, but we limit ourselves to verify a very simple and attractive solution. Let $\varphi(p)$ be

$$\varphi(p) = \frac{1}{p + m} \quad (3.21)$$

still

$$\partial_\mu \varphi(p) = -\varphi(p) \gamma_\mu \varphi(p)$$

$$\square \varphi(p) = 2\varphi(p) \gamma_\mu \varphi(p) \gamma_\mu \varphi(p)$$

and since $(p+m) = \varphi^{-1}(p)$, Eq. (3.21) gives a solution corresponding to $\lambda = 2$.

It is interesting to notice that the eigenvalue $\lambda = 2$ and the form of the solution

$$\varphi = \frac{1}{p + m} = \frac{p - m}{p^2 + m^2} \quad (3.21')$$

which corresponds to the free propagator, are strictly analogous to the lowest eigenvalue and eigenfunction of the scalar problem with an $1/R^2$ potential, which were

$$\lambda = 2 \quad \varphi(p) = \frac{1}{p^2 + m^2}$$

Going back to co-ordinate space we get for the $\varphi(x)$ corresponding to (3.21), aside unimportant constants,

$$\varphi(x) = m R K_1(m R) + i \gamma \cdot x m K_0(m R) \quad (3.22)$$

4. THE FACTORIZATION METHOD

Equation (3.2) can be also solved directly in co-ordinate space, using for instance a simple extension of the old factorization method ⁶⁾ to the case of a fourth order differential equation.

This method enables us to obtain very rapidly the solutions, and gives us directly the eigenfunctions in co-ordinate space, which is an advantage over the methods previously used in momentum space. We think therefore that it can be useful to rederive in this way the solutions of Eq. (3.2).

The factorization method has been often used in the past to solve Schrödinger's equation with central potentials: let us recall briefly how it works. The idea upon which the method is based can be expressed physically in the following way : we consider a differential operator, which represents a physical quantity, and transform it in a product of creation and annihilation operators for the physical quantity in question. The eigenfunctions can then be written, with the help of these annihilation and creation operators, and we then obtain the corresponding eigenvalues.

Let us state all this more exactly : suppose we have a differential equation, of fourth order for instance, which can be written in the two equivalent forms

$$+H^{m+1} - H^{m+1} \varphi(\lambda, m; R) = [\lambda - L(m+1)] \varphi(\lambda, m; R) \quad (4.1)$$

$$-H^m + H^m \varphi(\lambda, m; R) = [\lambda - L(m)] \varphi(\lambda, m; R) \quad (4.1')$$

where $\pm H^m$ are second order differential operators depending on a parameter, m , and $L(m)$ is a function of m . We can say that $\pm H^m$ are

annihilation and creation operators, since, if $\varphi(\lambda, m; R)$ is a solution of Eq. (4.1), then the functions

$$\begin{aligned} -H^{m+1} \varphi(\lambda, m, R) &= \varphi(\lambda, m+1, R) \\ +H^m \varphi(\lambda, m, R) &= \varphi(\lambda, m-1, R) \end{aligned} \quad (4.2)$$

are also solutions of (4.1), corresponding to the same eigenvalue λ , but to values of the parameter m shifted by a unity.

Suppose now that $\pm H^m$ are one the adjoint of the other, in the sense that

$$\int d\tau (\varphi, -H^m, \psi) = \int d\tau (+H^m \varphi, \psi)$$

where φ and ψ are continuous differentiable functions which vanish with their first derivatives at the boundaries. If, moreover, $L(m)$ is an increasing function of m and with suitable boundary conditions for $\varphi(\lambda, m; R)$, it can be easily shown that the eigenfunctions of (4.1) are given by

$$\varphi(\lambda, m; R) = \left(\prod_{i=0}^m +H^{m+i} \right) Y(m+u, R) \quad (4.3)$$

where $Y(m+u; R)$ are solutions of the second order differential equation

$$-H^{m+u+1} Y(m+u; R) = 0 \quad (4.4)$$

The eigenvalues corresponding to these eigenfunctions are

$$\lambda_m = L(m+u+1) \quad (4.5)$$

If we can determine the boundary conditions for Eq. (4.4) in such a way that the functions (4.3) satisfy the right boundary conditions for Eq. (4.1), the problem is completely solved.

After this introduction we can go back to Eq. (3.6)

$$\left(\frac{d^2}{dR^2} + \frac{1+\beta}{R} \frac{d}{dR} + \frac{\beta^2/4 - \bar{\nu}^2}{R^2} - 1 \right) \varphi_\alpha(R) = \frac{\lambda}{R^2} \varphi_\alpha(R) \quad (4.6)$$

We must write it in the two equivalent forms (4.1) and (4.1'). Let us write down Eq. (4.6) in the form

$$R \left(\frac{d^2}{dR^2} + \frac{\beta}{R} \frac{d}{dR} + \frac{1+\beta^2/4 - \bar{\nu}^2}{R^2} - 1 \right) \varphi_\alpha(R) = \lambda \frac{\varphi_\alpha(R)}{\sqrt{R}} \quad (4.7)$$

With the positions

$$\begin{aligned} Q^+(\bar{\nu}) &= \frac{d}{dR} \pm \frac{\bar{\nu} - 1/2}{R} + \frac{\beta/2}{R} \\ P^+(\bar{\nu}) &= \frac{d}{dR} \pm \frac{\bar{\nu} + 1/2}{R} - \frac{\beta/2}{R} \end{aligned} \quad (4.8)$$

Eq. (4.7) becomes

$$\begin{aligned} (R P^+ Q^+ + (\beta - 2\bar{\nu}) Q^+ - R) \varphi_\alpha(R) &= \lambda \frac{\varphi_\alpha(R)}{\sqrt{R}} \\ (R Q^+ P^+ + (2 - 2\bar{\nu} - \beta) P^+ - R) \varphi_\alpha(R) &= \lambda \frac{\varphi_\alpha(R)}{\sqrt{R}} \end{aligned} \quad (4.9)$$

or equivalently

$$\begin{aligned} & \left\{ \left[R P^+ Q^+ + (\beta - 2\bar{\nu}) Q^+ - R \right] R \left[Q^+ P^+ - 1 \right] - \right. \\ & \quad \left. - R P^+ \left[Q^+ P^+ - 1 \right] \cdot (2 - 2\bar{\nu} - \beta) + (\beta - 2\bar{\nu})(2 - 2\bar{\nu} - \beta) \cdot \right. \\ & \quad \left. \cdot \left[Q^+ P^+ - 1 \right] + 4 \left[\bar{\nu}(\bar{\nu} - 1) - \beta/2(\beta/2 - 1) \right] \right\} \frac{\varphi_\alpha(R)}{\sqrt{R}} = \lambda \frac{\varphi_\alpha(R)}{\sqrt{R}} \end{aligned} \quad (4.10)$$

and from here

$$\begin{aligned} & R \left\{ \left[P^+ + \frac{\beta - 2\bar{\nu}}{R} \right] \left[Q^+ - \frac{2 - 2\bar{\nu} - \beta}{R} \right] - 1 \right\} R \left\{ Q^+ P^+ - 1 \right\} \frac{\varphi_\alpha(R)}{\sqrt{R}} \\ & = \left\{ \lambda - 4 \left[\bar{\nu}(\bar{\nu} - 1) - \beta/2(\beta/2 - 1) \right] \right\} \frac{\varphi_\alpha(R)}{\sqrt{R}} \end{aligned} \quad (4.11)$$

and in the end

$$\begin{aligned} & R \left[Q_{\bar{\nu}}^+ P_{\bar{\nu}}^+ - 1 \right] R \left[Q_{\bar{\nu}}^+ P_{\bar{\nu}}^+ - 1 \right] \frac{\varphi_\alpha(R)}{\sqrt{R}} = \\ & = \left\{ \lambda - 4 \left[\bar{\nu}(\bar{\nu} - 1) - \beta/2(\beta/2 - 1) \right] \right\} \frac{\varphi_\alpha(R)}{\sqrt{R}} \end{aligned} \quad (4.12)$$

The equation remains unchanged if we interchange $\nu \rightarrow -\nu$.

Moreover

$$Q^\pm(\bar{\nu}) \rightarrow Q^\mp(\bar{\nu}+1)$$

$$P^\pm(\bar{\nu}) \rightarrow P^\mp(\bar{\nu}+1)$$

We can therefore write (4.6) also in the following form

$$\begin{aligned} & R \left[Q_{\bar{\nu}+1}^+ P_{\bar{\nu}+1}^+ - 1 \right] R \left[Q_{\bar{\nu}+1}^- P_{\bar{\nu}+1}^- - 1 \right] \frac{\varphi_\alpha(R)}{\sqrt{R}} \\ & = \left\{ \lambda - 4 \left[\bar{\nu}(\bar{\nu}+1) - \beta/2(\beta/2 - 1) \right] \right\} \frac{\varphi_\alpha(R)}{\sqrt{R}} \end{aligned} \quad (4.13)$$

We have now succeeded in writing Eq. (4.6) in the two forms (4.1') and (4.1), namely Eqs. (4.12) and (4.13). In our case

$$H_{\pm}^{\bar{\nu}} = R [Q^{\pm}(\bar{\nu}) P^{\pm}(\bar{\nu}) - 1] \quad (4.14)$$

$$L(\bar{\nu}) = 4 [\bar{\nu}(\bar{\nu}-1) - \beta/2 (\beta/2-1)]$$

It is easily verified that the operators $H_{\pm}^{\bar{\nu}}$ are one the adjoint of the other.

We notice now that the solution of the second order differential equation

$$H_{-}^{\bar{\nu}+m} \varphi(R) = R \left[\left(\frac{d}{dR} - \frac{\bar{\nu}+m-1/2}{R} + \frac{\beta/2}{R} \right) \left(\frac{d}{dR} - \frac{\bar{\nu}+m+1/2}{R} - \frac{\beta/2}{R} \right) - 1 \right] \varphi(R) = 0$$

which vanishes at the origin and at infinity is

$$\varphi(R) = R^{\bar{\nu}+m+1/2} K_{1-\beta/2}(R)$$

where $K_{1-\beta/2}(R)$ are Bessel functions of imaginary argument. The eigenvalues of (4.6) shall therefore be

$$\lambda_m = 4 [(\bar{\nu}+m)(\bar{\nu}+m+1) - \beta/2 (\beta/2-1)] \quad (4.15)$$

and the corresponding eigenfunctions

$$\varphi_m(R) = R^{1/2} \left(\frac{m}{0!}; H_{+}^{\bar{\nu}+1} \right) R^{\bar{\nu}+m+1/2} K_{1-\beta/2}(R) \quad (4.16)$$

It is easy now to verify that, with the expressions (3.8), for β and $\bar{\nu}$, the functions (4.16) satisfy the right boundary conditions, that have been imposed to the solutions of Eq. (4.6), according to the discussion of Section 2.

We can therefore conclude that (4.15) and (4.16) represent really the eigenvalues and eigenfunctions of our problem.

Obviously the same procedure can be applied to the regular potential case, and the corresponding results can be directly obtained from the previous ones in the limit $\beta \rightarrow 0$.

We give here explicitly the first few "wave functions" in the regular potential case :

$$\varphi_{\alpha}^{(0)}(R) = R^{\alpha+2} k_1(R)$$

$$\varphi_{\alpha}^{(1)}(R) = 4(\alpha+1)(\alpha+2) R^{\alpha+2} k_1(R) - 4(\alpha+2) R^{\alpha+3} k_0(R)$$

$$\varphi_{\alpha}^{(2)}(R) = 16(\alpha+1)(\alpha+2)^2(\alpha+3) R^{\alpha+2} k_1(R) - 16(\alpha+2)(\alpha+3) \quad (4.17)$$

$$(2\alpha+5) R^{\alpha+3} k_0(R) + 8(\alpha+3)(2\alpha+5) R^{\alpha+4} k_1(R).$$

A P P E N D I X A

We briefly discuss here the connection between the behaviour of $\varphi_\alpha(R)$ for $R \rightarrow 0$ and that of its Hankel transform for $u \rightarrow \infty$. We define

$$\varphi_\alpha(R) = \int_0^\infty du \, J_\nu(\sqrt{u}R) \varphi_\alpha(u) \quad (\text{A.1})$$

and we want to see under what conditions on $\varphi_\alpha(u)$, the function behaves at $R = 0$ as R^ν . To this end we use the following representation

$$J_\nu(x) = \frac{1}{2\pi i} \int_{-i\infty}^{+i\infty} \frac{\Gamma(-s)}{\Gamma(s+\nu+1)} \left(\frac{x}{2}\right)^{\nu+2s} ds \quad (\text{A.2})$$

and Eq. (A.1) becomes

$$\varphi_\alpha(R) = \left(\frac{R}{2}\right)^\nu \frac{1}{2\pi i} \int_{-i\infty}^{+i\infty} \frac{\Gamma(-s)}{\Gamma(s+\nu+1)} \left(\frac{R}{2}\right)^{2s} \lambda(s) ds \quad (\text{A.3})$$

$$\lambda(s) = \int_0^\infty du \, \varphi_\alpha(u) u^{\nu/2+s} \quad (\text{A.4})$$

It is easy to see that the integral converges when $\Re s = \beta - 1$, while it does not converge in the case $\Re s = 1 - \beta$. In fact, the condition of convergence is

$$\Re(2\bar{\nu} + 1) > \Re(-s) > -1$$

We shall first evaluate the regular kernel, then extract explicitly the singularities of the irregular one, and show that the total kernel $H(\bar{\nu}, \beta, u_1, u_2)$ is perfectly defined and convergent. We have

$$K(\bar{\nu}, \beta - 1, u, u') = \frac{\Gamma(\bar{\nu} + \beta/2)}{\Gamma(\bar{\nu} + 1)\Gamma(1 - \beta/2)2^{1-\beta}} \begin{cases} \left(\frac{u'}{u}\right)^{\bar{\nu}/2} u^{-\beta/2} F(\bar{\nu} + \beta/2, \beta/2, \bar{\nu} + 1, u'/u) & u' < u \\ \left(\frac{u}{u'}\right)^{\bar{\nu}/2} u'^{-\beta/2} F(\bar{\nu} + \beta/2, \beta/2, \bar{\nu} + 1, u/u') & u' > u \end{cases} \quad (\text{B.6})$$

where $F(a, b, c; z)$ is the hypergeometric function. Using for it the following integral representation

$$F(a, b, c; z) = \frac{\Gamma(c)}{\Gamma(b)\Gamma(c-b)} \int_0^1 \frac{dt \cdot t^{b-1}}{(1-tz)^a} (1-t)^{c-b-1}$$

$$\Re c > \Re b > 0$$

we get, with some manipulations

$$K(\bar{\nu}, \beta - 1, u, u') = \frac{1}{2^{1-\beta} \Gamma^2(1 - \beta/2)} \left(\frac{u}{u'}\right)^{\bar{\nu}/2} \int_0^1 \frac{dx \cdot e(u' - ux) x^{\bar{\nu} + \beta/2 - 1}}{(1-x)^{\beta/2} (u' - ux)^{\beta/2}} \quad (\text{B.7})$$

Now, in order to define correctly $K(\bar{\nu}, 1 - \beta, u, u')$ we shall give a small imaginary part to the arguments of the Bessel functions and let it tend to zero at the end of the calculation. To this end we will remember the relations

$$J_{\bar{\nu}}(\sqrt{u'} R) = \frac{1}{2} [H_{\bar{\nu}}^{(1)}(\sqrt{u'} R) + H_{\bar{\nu}}^{(2)}(\sqrt{u'} R)]$$

and

$$H_{\bar{\nu}}^{(1)}[(\sqrt{u'} + i\eta) R] = \frac{2}{i\pi} e^{-i\eta/2 \bar{\nu}} K_{\bar{\nu}}[(\eta - i\sqrt{u'}) R]$$

$$H_{\bar{\nu}}^{(2)}[(\sqrt{u'} - i\eta) R] = -\frac{2}{i\pi} e^{i\eta/2 \bar{\nu}} K_{\bar{\nu}}[(\eta + i\sqrt{u'}) R]$$

and we finally obtain

$$K(\bar{\nu}, 1-\beta, u, u') = \frac{1}{2\pi} 2^{1-\beta} \frac{\Gamma(\bar{\nu}+1-\beta/2)}{\Gamma(\bar{\nu}+1) \Gamma(\beta/2)} \times$$

$$\theta(u-u') u'^{\bar{\nu}/2} \left\{ \frac{e^{-i\eta\beta/2} F(a, b, c, u'/(u+2i\eta\sqrt{u'}))}{(\sqrt{u'} + i\eta)^{\bar{\nu}+2-\beta}} - \frac{e^{i\eta\beta/2} F(a, b, c, u'/(u-2i\eta\sqrt{u'}))}{(\sqrt{u'} - i\eta)^{\bar{\nu}+2-\beta}} \right\} \quad (B.8)$$

$$+ \theta(u'-u) u^{\bar{\nu}/2} \left\{ \frac{e^{-i\eta\beta/2} F(a, b, c, u/(u'+2i\eta\sqrt{u'}))}{(\sqrt{u'} + i\eta)^{\bar{\nu}+2-\beta}} - \frac{e^{i\eta\beta/2} F(a, b, c, u/(u'-2i\eta\sqrt{u'}))}{(\sqrt{u'} - i\eta)^{\bar{\nu}+2-\beta}} \right\}$$

where $a = \bar{\nu}+1-\beta/2$, $b = \beta/2$, $c = \bar{\nu}+1$. $a+b-c = 1$ and η is an infinitesimal real positive number.

If we remember now the formula

$$F(a+\sigma, 1+a+\sigma-c, c; z) = F(a, 1+a-c, c; z) (1-z)^{-\sigma}$$

we can extract the singularities of our kernel, which are simply the cuts of the hypergeometric function

$$K(\bar{\nu}, 1-\beta, u, u') = \frac{2^{1-2\beta}}{\pi} \frac{\Gamma(\bar{\nu}+1-\beta/2) \Gamma(\beta/2)}{\Gamma(\bar{\nu}+\beta/2)} K(\bar{\nu}, \beta-1, u, u').$$

$$\left\{ \frac{e^{-i\eta\beta/2}}{(u-u'+i\epsilon)^{1-\beta}} - \frac{e^{i\eta\beta/2}}{(u-u'-i\epsilon)^{1-\beta}} \right\} \quad (B.9)$$

We are now in a condition to evaluate the kernel $H(\bar{\nu}, \beta, u_1, u_2)$. Let us first consider the case $u_1 < u_2$. Using Eqs. (B.7) and (B.9) we have

$$H(\bar{\nu}, \beta, u_1, u_2) = \frac{\Lambda}{2\pi} \left(\frac{u_1}{u_2}\right)^{\bar{\nu}/2} \int_0^1 \frac{dx \cdot x^{a-1}}{(1-x)^{\beta/2}} \int_0^1 \frac{dy \cdot y^{a-1}}{(1-y)^{\beta/2}} \cdot \frac{1}{u_2+1} \int_{u_1 x}^{u_2/y} \frac{du'}{(u'+1)(u'-u_1 x)^{\beta/2}(u_2-u'y)^{\beta/2}} \left\{ \frac{e^{-i\pi\beta/2}}{(u_1-u'+i\epsilon)^{1-\beta}} - \frac{e^{i\pi\beta/2}}{(u_1-u'-i\epsilon)^{1-\beta}} \right\} \quad (\text{B.10})$$

where

$$\Lambda = \frac{\Gamma(\bar{\nu} + 1 - \beta/2)}{\Gamma^2(\beta/2) \Gamma(\bar{\nu} + \beta/2)}$$

If we consider first the integration over u' we see that the singularities of the integrand are three cuts in $(-\infty, u_1 x)$, (u_1, ∞) , $(u_2/y, \infty)$ and the pole in $u' = -1$, as shown in Fig. 1

Fig. 1

The dotted line shows the integration path.

By reversing the cut of the function $(u'-u_1 x)^{-\beta/2}$, bringing it to the interval $(u_1 x, \infty)$, we have

$$H(\bar{\nu}, \beta, u_1, u_2) = \frac{\Lambda}{2\pi} \left(\frac{u_1}{u_2}\right)^{\bar{\nu}/2} \frac{1}{u_2+1} \int_0^1 \frac{dx \cdot x^{a-1}}{(1-x)^{\beta/2}} \int_0^1 \frac{dy \cdot y^{a-1}}{(1-y)^{\beta/2}} \times \int_c \frac{du'}{u'+1} \cdot \frac{1}{(u_1 x - u')^{\beta/2} (u_2 - u'y)^{\beta/2} (u_1 - u')^{1-\beta}} \quad (\text{B.11})$$

and the disposition of the cuts is the following : Fig. 2

Fig. 2

The three cuts in the interval $(u_2/y, \infty)$ cancel: that is, if we start from a point which lies outside the interval $(u_1x, u_2/y)$ of the real axis, and come back to the same point, following a path which encircles the segment $(u_1x, u_2/y)$, the function gets multiplied by a factor $e^{2\pi i(-\beta/2 - (\beta/2 - 1) + \beta)} = 1$, and returns therefore to its original value. We can then close the integration path at the point u_2/y : we have now a closed contour, as shown in Fig. 3 :

Fig. 3

The integrand vanishes at infinity like $1/u'^2$: the integral is therefore simply equal to the residue in the pole at $u' = -1$, and we get

$$H(\bar{\nu}, \beta, u_1, u_2) = \Lambda \left(\frac{u_1}{u_2} \right)^{\bar{\nu}/2} \int_0^1 \frac{dx \cdot x^{a-1}}{(1-x)^{\beta/2} (u_1x+1)^{\beta/2} (u_2+1)} \int_0^1 \frac{dy \cdot y^{a-1}}{(1-y)^{\beta/2} (u_2+y)^{\beta/2} (u_1+1)} \quad (\text{B. 12})$$

The same method can be used in the case $u_2 > u_1$ and, remembering the relation

$$F(\bar{\nu} + \beta/2, \beta/2, \bar{\nu} + 1, z) = \frac{z^{-\bar{\nu}/2}}{(1-z)^{\beta/2}} P(\bar{\nu} + 1) P_{-\beta/2}^{-\bar{\nu}} \left(\frac{1+z}{1-z} \right)$$

the resulting integrals can be evaluated in terms of Legendre functions, giving

$$H(\bar{\nu}, \beta, u_1, u_2) = \frac{(\bar{\nu} - \beta/2) \Gamma(\bar{\nu} + \beta/2) \Gamma(\bar{\nu} - \beta/2)}{(1+u_2)^{\beta/2} (1+u_1)^{1-\beta/2}} \begin{cases} P_{-\beta/2}^{-\bar{\nu}} \left(\frac{1-u_1}{1+u_1} \right) P_{-\beta/2}^{-\bar{\nu}} \left(\frac{u_2-1}{u_2+1} \right) & u_1 < u_2 \\ P_{-\beta/2}^{-\bar{\nu}} \left(\frac{u_1-1}{u_1+1} \right) P_{-\beta/2}^{-\bar{\nu}} \left(\frac{1-u_2}{1+u_2} \right) & u_1 > u_2 \end{cases} \quad (\text{B.13})$$

This kernel is a regular function: therefore our equation, of the Fredholm type, can be solved in order to obtain the eigenvalues.

It can be easily verified that, by transforming Eq. (B.3) with the kernel (B.12) back to a differential equation in momentum space, we get the same differential equation (3.15). In fact, introducing the new function

$$\phi(u) = u^{-\bar{\nu}/2} (1+u)^{1-\beta/2} \mathcal{P}^{(1)}(u) \quad (\text{B.14})$$

and remembering that

$$\left(\frac{d^2}{du^2} + \frac{1}{u} \frac{d}{du} + \frac{\beta/2(\beta/2-1)}{u(u+1)^2} - \frac{\bar{\nu}^2}{4u^2} \right) P_{-\beta/2}^{-\bar{\nu}} \left(\pm \frac{1-u}{1+u} \right) = 0$$

we find, after some manipulations, that

$$\left(\frac{d^2}{du^2} + \frac{\bar{\nu}+1}{u} \frac{d}{du} + \frac{b}{4u(u+1)^2} \right) \phi(u) = 0 \quad (\text{B.15})$$

To obtain the boundary conditions implied by the starting integral equation, we insert there

$$\frac{\lambda \varphi(u)}{4(u+1)^2} = -u \varphi'' - (\bar{\nu}+1) \varphi' - \frac{\beta/2 (\beta/2-1)}{(u+1)^2} \varphi$$

and integrating by parts we get

$$\begin{aligned} \lim_{u \rightarrow 0} u^{\bar{\nu}+1} \varphi'(u) &= 0 \\ \lim_{u \rightarrow \infty} (\bar{\nu} \varphi(u) + u \varphi'(u)) &= 0 \end{aligned} \quad (\text{B.17})$$

which select the solutions of Eq. (B.15) whose behaviour is

$$\varphi(u) \sim \cos t \quad ; \quad \varphi(u) \sim u^{-\bar{\nu}} \\ u \rightarrow 0 \quad \quad \quad u \rightarrow \infty$$

Obviously the same procedure can be applied to the regular potential case. In the limit $\beta \rightarrow 0$ we get, for instance, from (B.13) the kernel of the regular case and its expression is

$$H(\alpha+1, 0, u_1, u_2) = \frac{1}{4(\alpha+1)(u_1+1)(u_2+1)} \cdot \begin{cases} \left(\frac{u_1}{u_2}\right)^{\frac{\alpha+1}{2}} & u_1 < u_2 \\ \left(\frac{u_2}{u_1}\right)^{\frac{\alpha+1}{2}} & u_1 > u_2 \end{cases}$$

R E F E R E N C E S

- 1) G.C. Wick, Phys.Rev. 96, 1124 (1954);
R.E. Cutkosky, Phys.Rev. 96, 1135 (1954).
- 2) C. Ceolin, F. Duimio, R. Stroffolini, S. Fubini, Nuovo Cimento 26, 247 (1962).
- 3) A. Bastai, L. Bertocchi, S. Fubini, G. Furlan and M. Tonin, preceding paper, from now on cited as I.
- 4) H.A. Bethe and E.E. Salpeter, Phys.Rev. 84, 1232 (1951);
M. Gell-Mann and F.E. Low, Phys.Rev. 84, 350 (1951).
- 5) See, e.g. : A. Erdélyi et al., Bateman Manuscript Project, McGraw Hill (1953), Vol. II.
- 6) L. Infeld and T.E. Hull, Rev.Mod.Phys. 23, 21 (1951).
- 7) J.S. Goldstein, Phys.Rev. 91, 1516 (1953).