CERN LIBRARIES, GENEVA

CM-P00056972

S

SINGULAR POTENTIALS IN COORDINATE SPACE

H. Cornille

CERN - Geneva

9608/TH.479

24 September, 1964

ABSTRACT

With the help of the Laplace transform previous works have shown the possibility of useful approximate methods for the Jost functions in the case of Yukawa generalized potentials, where the most singular term near the origin is like $G^2 r^{-2n}$ ($n \geq 1$, $G^2 > 0$ for $n > 1$). Now, we want to study directly in coordinate space the same problem of the construction of the Jost function for families of singular potentials where the most singular term near the origin is repulsive. In this first paper we consider the same family of short-range potentials, as previously made with the use of Laplace transform, and also a family of arbitrary power potentials more singular than r^{-2} . The key (as previously made) is to define "new Jost solutions" asymptotically ingoing (or outgoing) and going to constant (Jost function) near the origin. If we consider the perturbation expansion of these new Jost functions we find, that if we connect in a certain manner the order of the perturbation p and the radial coordinate, then the Jost function is the limit of convergent sequences. More explicitly, we find that there exists a limiting dependence $r_L(p) = \text{const}/(p^{(1/n-1)-\epsilon'})$, ($\epsilon' > 0$ arbitrary small) such that if $r(p)$ goes to zero less rapidly than $r_L(p)$ these sequences are convergent. We have never used arbitrary cut-off or hard-core procedure and only when p is infinite one has that these $r(p)$ dependences are 0.

I. INTRODUCTION

In two preceding papers^{1,2)} we have studied the problem of the explicit determination of the Jost function for generalized Yukawa potentials with a behaviour near the origin for the most singular term like $G^2 r^{-2n}$ ($n \geq 1$). In the first work¹⁾ the case $G^2 > -(\ell + \frac{1}{2})^2$ was considered, whereas in the second one for the S wave we have studied $n > 1$ and $G^2 > 0$.

For both cases the starting point was the same. We want to factorize in an exact manner the most singular parts of the functions near the origin, in such a way that we do not affect the asymptotic behaviour at infinity. This can be made for the Jost solutions as well as for the regular solutions (and we can also impose as a further requirement, that the remaining solutions behave as in the usual case, but this is a secondary point).

These procedures define "new Jost solutions" or "new regular solutions." But there is still a big difference between the cases $n = 1$ and $n > 1$. For $n = 1$ we always have a Fuchs' type equation, whereas for $n > 1$ we have still a singular equation not of the Fuchs' type. Thus, in the case $n = 1$ this factorization of the most singular part is sufficient in order to resolve the problem encountered (Jost function, S matrix etc.), but this is not the case for $n > 1$. In Refs. 1 and 2 the tool used was the Laplace transform and these features were illustrated.

We now consider the more delicate case $n > 1$. Suppose that we have been able to reduce the problem to the following form. We want to determine a finite quantity $\lim_{x \rightarrow x_0} \Phi(x) = \Phi(x_0)$ (for simplicity we suppose $x_0 = 0$) defined by

$$a) \quad \Phi(x) = \sum_0^{\infty} \psi_q(x) \quad \text{or} \quad b) \quad \Phi(x) = \int_0^{\infty} \kappa(x,y) dy ,$$

where we can determine $\psi_q(x)$ or $\kappa(x,y)$ independently of $\Phi(x)$ from integral equations or differently. Further, suppose that we know

2.

$$1) \quad \lim_{x \rightarrow 0} \psi_q(x) \quad \text{or} \quad \lim_{z \rightarrow \infty} \left(\int_0^z \left(\lim_{x \rightarrow 0} \kappa(x, y) \right) dy \right) \text{ is divergent.}$$

$$2) \quad \lim_{x \rightarrow 0} \left(\sum_0^\infty \psi_q(x) \right) \quad \text{or} \quad \lim_{x \rightarrow 0} \left(\int_0^\infty \kappa(x, y) dy \right) \text{ exist and are equal to } \Phi(0).$$

Then we have the feeling that if we connect the variable $x \rightarrow 0$ and, in a) the order of the perturbation expansion, or in b) the range of the integration, it is perhaps possible to find $x(p)$ or $x(z)$ such that

$$3) \quad \Phi(0) = \lim_{p \rightarrow \infty} \left(\sum_0^p \psi_p(x(p)) \right) \quad \text{or} \quad \Phi(0) = \lim_{z \rightarrow \infty} \int_0^z \kappa(x(z), y) dy.$$

Because of the fact that we cannot inverse $x \rightarrow 0$ and $p \rightarrow \infty$ in a) or $x \rightarrow 0$ and $z \rightarrow \infty$ in b) we have also the feeling that if these $x(p) \xrightarrow[p \rightarrow \infty]{} 0$ or $x(z) \xrightarrow[z \rightarrow \infty]{} 0$ exist, they cannot be entirely arbitrary.

In Ref. 2 the integral in b) was connected with the Laplace transform and $\Phi(0)$ was the Jost function. We have shown that for $n \geq 2$ properties 1), 2) give 3), if $x(z)$ goes to zero less rapidly than limiting dependence. For $1 < n \leq 2$ where we can inverse the two limits $x \rightarrow 0$ and $z \rightarrow \infty$ in 1) there are no limiting dependences $x(z)$ in order to obtain $\Phi(0)$ as the limit of sequences 3).

It is the purpose of this work to show that for the perturbation expansion a) of the new Jost solution in radial coordinate, where properties 1 and 2 occur, we can find limiting dependence $r_L(p)$ in 3) such that 3) is satisfied if $r(p)$ goes to zero less rapidly than a limiting dependence $r_L(p)$.

In Section II we consider only S waves, where we explain the method for singular Yukawa generalized potentials and the family of arbitrary power singular potentials more singular than r^{-2} . In Section III, we extend briefly the results to $\ell \neq 0$ and, in Section IV we consider the peculiar case of the Bethe-Salpeter equation reduced in a differential form very similar to the Schrödinger one.

II. S WAVE SCHRÖDINGER EQUATION

a) Short-range potentials

We consider the Schrödinger equation for $\ell = 0$

$$\left(\frac{d^2}{dr^2} + k^2 \right) u = Vu . \quad (1)$$

We have studied previously the case of singular potentials with short range in two cases:

a) potential singular like r^{-2} near the origin¹⁾

$$V = G^2 \frac{e^{-\mu r}}{r^2} \quad (G^2 > -1/4) \quad (2.a)$$

b) potentials more singular than r^{-2} near the origin²⁾

$$V = e^{-\mu r} \left[\frac{G^2}{r^{2n}} - \sum_{2n-1}^{N_0} \frac{A_p}{r^p} \right] \quad (n > 1, G^2 > 0) \quad (2.b)$$

where n is arbitrary > 1 . For integer n , $N_0 = 2$. For non-integer n , N_0 is the smallest number deduced from $2n$ by substitution of successive integers, such that $2 < N_0 < 3$.

We have found that for potentials like (2.a) and (2.b), in order to obtain the Jost functions and the S matrix, it was very convenient to use "new Jost solutions" $F(\pm k, r)$ such that if $R(\pm k, r)$ is the solution of (1) asymptotically ingoing or outgoing

$$R(k, r) = Z(r) F(k, r) \underset{r \rightarrow \infty}{\simeq} e^{-ikr} , \quad (3)$$

We want to impose to $Z(r)$ conditions such that the "new Jost solutions" $F(k, r)$ near the origin have a behaviour when $r \rightarrow 0$ similar to the usual case.

Then Z is a singular k independent factor of the kind
 $Z(r) = e^{\int_{-\infty}^r \psi(r') dr'}$ with

$$1) \quad Z(r) \rightarrow 1 \quad \text{or} \quad F(k, r) \sim e^{-ikr} \quad \begin{matrix} r \rightarrow \infty \\ r \rightarrow \infty \end{matrix}$$

$$2) \quad Z(r) \simeq \text{singular part of } R(k, r) \quad \text{or} \quad F(k, r) \rightarrow \text{const.} \quad \begin{matrix} r \rightarrow 0 \\ r \rightarrow 0 \end{matrix}$$

Then $F(k, r)$ satisfies the differential equation

$$\left(\frac{d^2}{dr^2} + k^2 \right) F = \left(-2\psi \frac{d}{dr} + W \right) F \quad (W = V - \psi' - \psi^2) \quad (4)$$

and for the S matrix we get

$$S(k) = \frac{F(k, 0)}{F(-k, 0)} \quad (5)$$

It was also shown that we can choose ^{*)} ψ of the kind

$$\psi = e^{-\lambda r} \left(\sum_n^{n_0} \frac{a_n}{r^n} + \frac{b}{r} \right) \quad \lambda > 0 \quad (6)$$

where all $a_q = 0$ for $n = 1$; $n_0 = 2$ for n integer > 1 , n_0 for n non-integer > 1 is the smallest number deduced from n by subtraction of successive integers such that $1 < n_0 < 2$. The parameters a_q, b can be determined with the condition that $W \simeq 0(1/r^m)$ where $m = 1$ if $n = 1$, $m = 2$ if $1 < n \leq 3/2$, $m = n$ if n integer and $m = n + n_0 - 1$ for non-integer $n \geq 3/2$. The presence of the exponential ($\exp - \lambda r$) ensures that the condition $Z(r) \rightarrow 1$ is satisfied. Many choices of λ are possible but for instance ^{1, 2)} $\lambda = \mu$ or $\lambda = \mu/2$ are more convenient. We recall that it was shown that the condition $W \simeq 0(1/r^m)$ gives two families of parameters a_n , entering in the expression of ψ , and then two factors $Z_{\epsilon}(r)$ [see Ref. 2, Eq. (2.7)]

^{*)} (6) is convenient for families of singular potentials (2.a,b) with a power behaviour near the origin. For another family, for instance logarithmic singular potentials, the corresponding form of ψ will be modified⁸⁾.

$$a_n = \epsilon G, \quad a_{n-1} = -\epsilon \frac{A_{2n-1}}{2G}, \quad b = n/2 - \epsilon \frac{B_1}{2G} \quad \epsilon = \pm 1$$

$$Z_\epsilon(r) \underset{r \rightarrow 0}{\simeq} Y_\epsilon(r) X_\epsilon(A_{p,\mu}), \quad X_\epsilon = \text{const.}$$

$$B_1 = B_1(G, \mu, A_p)$$

It was also shown that with these choices of $\epsilon = \mp 1$, the corresponding $Y_\epsilon(r)$ behave, near the origin, like the "singular" part and "regular" part of the solutions of (1). For Eq. (3) we choose, of course, $\epsilon = -1$. Now if we consider a "regular" solution u of (1), we can also define another "regular" solution with the same asymptotic behaviour as u by the change $u = Z_{\epsilon=+1} \Phi$ and $\Phi \underset{r \rightarrow 0}{\rightarrow} \text{const.}$ (Or, if we want $\Phi(r) \simeq \text{const } r$ similarly to the usual case where the potential is like $r^{-\eta}$ ($\eta < 2$), we take $\epsilon = +1$ and modify $b \rightarrow b-1$ in (6).) However, we remark that for $n > 1$ with the new regular solution or new Jost solutions, there is still a big difference in comparison with the usual case, because the corresponding differential equation (4) is not of the Fuchs' type near the origin. This is why it is perhaps more simple to work in the first stage with the new Jost solutions F . In fact, while the left-hand side of (4) gives the possibility of starting asymptotically with a purely outgoing or ingoing wave, the right-hand side appears, for such solutions as a perturbation, with an exponential decreasing factor. Then we can use an integral equation for F starting from infinity with a Green function, as in the usual case (see Eq. (7) below). This was also the reason why we have studied in Refs. 1 and 2 Eq. (4) by means of Laplace transform:

$$F(k, r) = e^{-ikr} \int_0^\infty e^{-\alpha r} \rho(\alpha) d\alpha.$$

It was shown that ρ satisfies a Volterra type equation $\rho(\alpha) = \delta(\alpha) + \Sigma \rho_q(\alpha)$. For the corresponding expansion of the Jost function as a series whose p -th term comes from the contribution of order p of the inverse Laplace transform we have found

$$a) \quad F(k, 0) = \sum_0^\infty \int_0^\infty \rho_q(\alpha) d\alpha \quad \text{for } n = 1 \text{ and } G^2 > -1/4$$

b) $\sum_0^\infty \int_0^\infty f_q(\alpha) d\alpha$ was always divergent for $n > 1$.

Thus, for $n = 1$, $Z(r)$ was sufficient in order to obtain the Jost function similarly as in the usual case ($V(r) \underset{r \rightarrow 0}{\simeq} r^{-\eta}$ $\eta < 2$).

For $n > 1$, we are in a very different situation because the considered expansion always diverges. It was then necessary to consider other methods²⁾. We have found that there exist sequences $G(x, r(x), k)$:

$$G(x, r(x), k) = \int_0^x e^{-\alpha r(x)} \rho(\alpha, k) d\alpha \quad \text{such that} \quad \lim_{x \rightarrow \infty} G(x, r(x), k) = F(k, 0)$$

if $r(x)$ goes to zero less rapidly than a limiting dependence $r_L(x)$.

It is the aim of this paper to show that these kinds of convergences encountered in Ref. 2 (concerning the possibility of connecting two limiting dependences in order to obtain Jost functions and S matrix) are not peculiar to the use of the Laplace transform, but can arise in other treatments of problems connected with singular potentials and with non-analytic expansions with respect to the coupling constant. We want to show here that indeed this is also the case in coordinate space.

From (4) and the asymptotic behaviour of F we can write an equivalent integral equation

$$F(k, r) = e^{-ikr} + \frac{1}{k} \int_r^\infty \sin k(r - r') \left(-2\psi(r') \frac{d}{dr'} + W(r') \right) F(k, r') dr' \quad (7)$$

but it is more convenient to study another related function $F = e^{-ikr} f$ which satisfies

$$f(k, r) = 1 + \frac{1}{2ik} \int_r^\infty \left(e^{2ik(r-r')} - 1 \right) \left(-2\psi(r') \frac{d}{dr'} + W(r') \right) f(k, r') dr'. \quad (8)$$

We iterate (8) and write formally

$$f(k, r) = 1 + \sum_{q=1}^{\infty} f_q(k, r) . \quad (9)$$

The Jost function is defined as

$$f(k, 0) = 1 + \lim_{r \rightarrow 0} \left(\sum_{q=1}^{\infty} f_q(r) \right) . \quad (10)$$

For the case $n = 1$ we find $f_q(r) \rightarrow \text{const.}$ as $r \rightarrow 0$. Thus, in this case, each iteration gives a finite contribution to the sum. Of course, this is not sufficient for the sum $\sum f_q(0)$ to be convergent. But it can be shown by investigating the behaviour of $f(k, r)$ near $r = 0$ in the integral equation (8), that $\lim_{r \rightarrow 0} f(k, r) \rightarrow \text{const}$ for G^2 in a certain range ($G^2 > -1/4$) which includes $G^2 = 0$. All this is similar to what was found in Ref. 1. This is easy to understand because the Laplace transform of (8) was the equation studied, and the results are clearly connected.

For the case $n > 1$ we find for the most singular term (leading term) of the q -th iteration of f in (8)

$$f_q(r) \simeq \text{const } r^{-[q(n-1) + m - n - 1]} . \quad (11)$$

At this stage we consider only the r dependence, but of course the q dependence is also very important, and this will be illustrated later. Then in this case ($n > 1$) each iteration is divergent when r goes to 0. Now, if we try in (8) a power behaviour for $f \simeq \text{const } r^{-\gamma}$ as $r \rightarrow 0$ we find by comparing the most singular terms at both sides of (8)

$$\text{const}_1 r^{-\gamma} \simeq \text{const}_2 r^{-(\gamma + n - 1)} .$$

This means that for $n > 1$ a power behaviour of f when $r \rightarrow 0$ is not allowed. This reflects the fact that $f(k, r)$ has essential singularities near $r = 0$.

We must then investigate the behaviour of the whole function f near the origin in order to show that the limit in (10) exists. We can use two methods.

1) Firstly we note that (8) is also the inverse Laplace transform of the equation studied in Ref. 2 [Eq. (3.3)]. If we consider the iteration of the Laplace transform $\rho = \sum \rho_q$ we have identically for each f_q of (8) $f_q(r) = \int_0^\infty e^{-\alpha r} \rho_q d\alpha$ and the behaviour (11) was found. Furthermore, in Ref. 2 it was shown that $\lim_{r \rightarrow 0} \left(\int_0^\infty e^{-\alpha r} \rho(x) d\alpha \right)$ exists, but

$$\begin{aligned} \lim_{r \rightarrow 0} \left(\sum_{q=0}^{\infty} f_q(r) \right) &= \lim_{r \rightarrow 0} \left(\sum_{q=0}^{\infty} \int_0^\infty e^{-\alpha r} \rho_q(\alpha) d\alpha \right) = \lim_{r \rightarrow 0} \left(\int_0^\infty \sum_{q=0}^{\infty} \rho_q e^{-\alpha r} d\alpha \right) = \\ &= \lim_{r \rightarrow 0} \left(\int_0^\infty e^{-\alpha r} \rho(\alpha) d\alpha \right) < \infty. \end{aligned}$$

2) In order to have a self-consistent study (independent of the Laplace transform) in coordinate space, we note that the integral equation (7) is equivalent to the differential equation (4) (and the same is true for (8) with a slight change from (4) for the corresponding differential equation.) This slight change does not affect the behaviour of the solutions of (4) near the origin. In Ref. 2 (Appendix) it was shown that $f(k, r)$ near the origin behaves like

$$f(k, r) \underset{r \rightarrow 0}{\sim} \text{const}_1 + \text{const}_2 \bar{f}(r)$$

where the singular part $\bar{f}(r)$ is

$$\begin{aligned} \bar{f}(r) = \exp \left[- \frac{2G}{(n-1)r^{n-1}} + \frac{1}{(n-2)r^{n-2}} \left(\frac{A_{2n-1}}{G} + \mu G \right) + \dots + \right. \\ \left. + \frac{\text{const}}{r^{n_0-1}} + \text{const} \log(r) \right]. \end{aligned} \quad (12)$$

The behaviour (12) ensures that the limit in (10) exists. Thus we can write

$$f(k,0) - 1 = \lim_{r \rightarrow 0} \left(\lim_{p \rightarrow \infty} \sum_{q=1}^p f_q(r) \right) \quad (13)$$

$$f(k,0) - 1 = \lim_{r \rightarrow 0} \left(\lim_{x \rightarrow \infty} \int_0^x e^{-\alpha r} \rho(\alpha) d\alpha \right). \quad (14)$$

In Ref. 2, where (14) has been proved, we have connected the two limiting procedures $x \rightarrow \infty$ and $r \rightarrow 0$ by replacing the double limit by a single one and attributing an $x(r)$ or $r(x)$ dependence. We have also been able to show that there exists a limiting dependence $x_L(r)$ or $r_L(x)$ such that for $r(x) \rightarrow 0$ less rapidly than $r_L(x)$ (or $x(r) \rightarrow \infty$ more rapidly than $x_L(r)$), (14) is satisfied and the Jost function is the limit of convergent sequences.

Here, we consider (13). The fact that we cannot inverse the two limiting procedures is illustrated by the fact that

$$\lim_{p \rightarrow \infty} \left(\sum_{q=1}^p \lim_{r \rightarrow 0} f_q(r) \right) \quad (15)$$

is divergent for any $n > 1$.

We define the sequence

$$G(p,r,k) = 1 + \sum_{q=1}^p f_q(r). \quad (16)$$

In this paper, we want to see if it is possible to connect the order of the perturbation expansion ($p \rightarrow \infty$) and the radial coordinate ($r \rightarrow 0$), in order to have the following identities for $r(p) \rightarrow 0$ $p \rightarrow \infty$

$$\lim_{p \rightarrow \infty} G(p, r(p), k) = f(k, 0) = \lim_{r \rightarrow 0} \left(\lim_{p \rightarrow \infty} G(p, r, k) \right). \quad (17)$$

Since the two limits $p \rightarrow \infty$ and $r \rightarrow 0$ cannot be interchanged, (15), we argue that $r(p)$ cannot be completely arbitrary for $n > 1$. This is in contrast with the situation which occurs in (14) for peculiar ranges of n . We recall that in Ref. 2, for the cases $1 < n \leq 2$ ($n = 2$ for special values of the potential), it was shown that the two limits $x \rightarrow \infty$ and $r \rightarrow 0$ can be interchanged and, correspondingly, for these values of n , $r(x)$ can be chosen arbitrarily going to zero.

We then have the feeling that for the same n , $n > 1$ in (13) and $n \geq 2$ in (14), the limiting dependence $r_L(p)$ cannot be the same as $x_L(r)$ previously found in Ref. 2.

We note that for any p we have identically

$$f(k, r(p)) = G(p, r(p), k) + R(p, r(p), k) \quad (18)$$

where

$$R(p, r, k) = \sum_{p+1}^{\infty} f_q(r) . \quad (19)$$

Suppose now that we give an arbitrary dependence $r(p)$ such that $r(p) \rightarrow 0$ when $p \rightarrow \infty$. Then from (18) one has

$$f(k, 0) = \lim_{p \rightarrow \infty} G(p, r(p), k) + \lim_{p \rightarrow \infty} R(p, r(p), k) . \quad (20)$$

Of course, although the sum of the two limits of G and R is finite, this does not ensure that separately each of these limits is finite. In fact, we can have, for instance

- 1) the two limits are infinite and cancel, in order that the sum be finite;
- 2) the two limits are not defined (for instance oscillating like $\lim_{p \rightarrow \infty} \sin(\text{const } p)$);
- 3) the two limits are finite.

We want to find a situation similar to the third case such that $\lim_{p \rightarrow \infty} R \rightarrow 0$. Then the problem is reduced to find, if possible, limiting dependence such that

$$\lim_{p \rightarrow \infty} R(p, r(p), k) \rightarrow 0 \quad (r(p) \geq r_L(p)) . \quad (21)$$

For this we have only to investigate the singular part of $f(k, r)$, because for the regular part, where we can inverse the two limits ($p \rightarrow \infty$ and $r \rightarrow 0$), we can attribute an arbitrary $r(p)$ dependence such that $r(p) \rightarrow 0$ when $p \rightarrow \infty$.

In order to investigate the singular part of the remaining terms $R(p, r, k)$ of the perturbative expansion $\sum_q f_q(r)$, we must, in principle, for each f_q , investigate all the singular terms. Of course we cannot investigate all the singular terms coming in $f_q(r)$ when q is very large. But, because we know the whole singular part of $f(k, r)$ from (12), we shall see that in fact it is sufficient to investigate for each q the two most singular terms. In the Appendix it is shown from the determination of these two singular terms when p is large:

a) for $n > 2$

$$R(p, r, k) \simeq \text{const } r^n \left[\sum_{q=p+1}^{\infty} \left(\frac{-2G}{(n-1)r^{n-1}} \right)^q \frac{1}{q!} \left[1 + \frac{qr \delta_a}{(n-2)} \left(\frac{n-1}{-2G} \right) + \dots \right] \right] \quad (22.a)$$

b) for $1 < n \leq 2$

$$R(p, r, k) \simeq \text{const } r^n \left[\sum_{q=p+1}^{\infty} \left(\frac{-2G}{(n-1)r^{n-1}} \right)^q \frac{1}{q!} \left[1 - q \delta_b \log r \left(\frac{-2G}{n-1} \right) r^{n-1} \right] + \dots \right] \quad (22.b)$$

where for $n > 2$ $\delta_a = \frac{A_{2n-1}}{G} + \mu G$, for $n = 2$ $\delta_b = n + \frac{A_{2n-1}}{G} + \mu G$, for $1 < n < 2$ $\delta_b = n$.

We recall that it was shown in Ref. 2 (Appendix) that for $i < n < i+1$, (i integer) we have for the singular part of $f(k,r)$ near the origin, a product of i essential singularities and a power behaviour. This is the reason why, when we consider the two first leading singular terms we must distinguish between $n > 2$ and $1 < n \leq 2$. If we want to consider the third singular term of each $f_q(r)$ we still consider $n > 3$ and $2 < n \leq 3$. We recall now that in Ref. 2 (see also (12)) we have shown for the whole singular part of $f(k,r)$

a) $n > 2$

$$\begin{aligned} \bar{f}(r) \underset{r \rightarrow 0}{\simeq} \text{const } r^n \exp \left[\left(\frac{-2G}{(n-1)r^{n-1}} \right) \left(1 + \frac{r \delta_a}{(n-2)} \left(\frac{n-1}{-2G} \right) + \dots + \right. \right. \\ \left. \left. + \text{const } r^{n-n_0} + (\text{const}-n) \left(\frac{(n-1)r^{n-1}}{-2G} \right) \log(r) \right) \right] \end{aligned} \quad (23.a)$$

b) $1 < n < 2$

$$\begin{aligned} \bar{f}(r) \underset{r \rightarrow 0}{\simeq} \text{const } r^{-(\delta_b - n)} \exp \left(\frac{-2G}{(n-1)r^{n-1}} \right) \underset{r \rightarrow 0}{\simeq} \text{const } r^n \exp \left[\left(\frac{-2G}{(n-1)r^{n-1}} \right) \times \right. \\ \left. \times \left(1 + \delta_b \left(\frac{n-1}{2G} \right) r^{n-1} \log(r) \right) \right] . \end{aligned} \quad (23.b)$$

Thus we see from (22) and (23), when p is large, that the singular part of R is of the kind

a) $n > 2$

$$R(p,r,k) \simeq \text{const } r^n \sum_{q=p+1}^{\infty} \left(\frac{-2G}{(n-1)r^{n-1}} \right)^q \frac{1}{q!} \left[1 + o(r) \right]^q \quad (24.a)$$

b) $1 < n \leq 2$

$$R(p,r,k) \simeq \text{const } r^n \sum_{q=p+1}^{\infty} \left(\frac{-2G}{(n-1)r^{n-1}} \right)^q \frac{1}{q!} \left[1 + o(r^{n-1} \log(r)) \right]^q . \quad (24.b)$$

Now we attribute an $r(p)$ dependence and use the Stirling formula

$$q! \underset{q \rightarrow \infty}{\simeq} q^q e^{-q} \sqrt{2\pi q} . \quad (25)$$

From (24), if $r(p)$ goes to zero less rapidly^{*)} than $1/p^{(1/n-1)-\epsilon}$ ($\epsilon > 0$ arbitrary small), we get

$$\lim_{p \rightarrow \infty} R(p, r(p), k) \rightarrow 0 \quad \text{if} \quad r(p) \geq \frac{\text{const}}{p^{(1/n-1)-\epsilon}} \quad (\epsilon > 0) . \quad (26)$$

Note that as previously thought the two limiting dependences $r_L(p)$ and $r_L(x)$ are not the same (only for $n = 2$ are they similar). Although (22), (23) and (24) are not valid for $n = 1$, if we take formally $r \simeq 1/p^{(1/n-1)-\epsilon}$ for $n = 1$, we find, in this case, that there is no limiting dependence. This is clearly correct from the above consideration about the $n = 1$ case. What is the significance of this $r_L(p)$? It means, roughly speaking, that if we consider an approximation up to a certain order p of the perturbation expansion, we must take into account at most, only the part of the interaction which is outside a limiting radius $r_L(p)$. When p grows up, this radius decreases and only when p is infinite does it reduce to zero. Now this radius depends mainly on two things. Firstly, it is connected with the most singular part of the interaction. On the other hand, it depends also on the singular integral equation considered, and on the other variable coupled to the radial coordinate. For instance, in Ref. 2, we have considered the inverse Laplace transform and the other variable x was the limit of integration in the inverse Laplace transform connected for potentials like (2.b) to the order of perturbation of the inverse Laplace transform. It can also be (see Ref. 2 for $1 < n < 2$), that there is no limiting dependence, but we note that in all cases considered in Ref. 2 and here, for $n > 1$, it appears that it is convenient to construct the Jost function as the limit of convergent sequences.

^{*)} Note that $r(p) \simeq 1/p^{(1/n-1)-\epsilon}$ is not the finest limiting dependence. We can choose $r(p) \simeq \text{const}/p^{(1/n-1)} r_1(p)$ and determine the const and $r_1(p)$, which is a less important dependence with respect to p than a power, but the difference is then not very significant.

We find that $Y_\epsilon(r)$, independent of λ , is a product of essential singularities and a power

$$Y_\epsilon(r) = r^\alpha \exp \left(\sum_{i=1}^j \frac{\beta_i(\epsilon)}{r^{\gamma_i}} \right)$$

and

$$\begin{aligned} \gamma_1 &= n-1, \gamma_2 = n-2, \dots, \gamma_j = n_0-1 \\ \beta_1 &= -\frac{\epsilon G}{(n-1)}, \beta_2 = +\frac{\epsilon A_{2n-1}}{2G(n-2)}, \dots \end{aligned} \quad (29)$$

In Appendix B we investigate directly the behaviour of R near the origin and we find that $Y_{\epsilon=-1}$ and $Y_{\epsilon=+1}$ corresponds respectively to the "singular" and the "regular" part of R . Thus in this case also Z_ϵ can be used in order to define "new Jost solutions" and new "regular solutions".

Now, we consider the new Jost solution $F(k,r)$ such that $R(k,r) = Z_{-1}(r)F(k,r)$ and try to construct the Jost function $F(k,0)$.

1) If we use the Laplace transform we obtain the same type of convergence as in Ref. 2. In the case of power behaviour (27), the only difference with the short range (2.b) case is that we do not have the peculiar iterative properties for the iterations of the inverse Laplace transform (i.e. we have not $p_q(\alpha) = 0$ for $\alpha < q\lambda$). But all other things go parallel in the two cases and we have, for instance, the same kind of $x_L(r)$ dependence for $n \geq 2$.

2) On the other hand, we can also consider directly the problem in coordinate space. From the differential equation (4), similarly to the short range case, we get

$$f(k,0) = \lim_{r \rightarrow 0} \left(\sum_{q=0}^{\infty} f_q(r) \right).$$

All the arguments previously given are still valid and, for instance, we can still connect the order of the perturbation expansion and the radial coordinate. We find that if $r(p) \rightarrow 0$ less rapidly than $1/p^{(1/n-1)-\epsilon}$, the corresponding sequences $G(p, r(p), k)$ converge to the Jost function.

III. CASE OF $\ell \neq 0$ SCHRÖDINGER EQUATION

In this case

$$\left(\frac{d^2}{dr^2} + k^2 - \frac{\ell(\ell+1)}{r^2} \right) R_\ell = V R_\ell. \quad (30)$$

We define similarly to the $\ell = 0$ case

$$R_\ell(k, r) = Z_\ell(r) F_\ell(k, r) \underset{r \rightarrow \infty}{\sim} e^{-ikr} \quad (31)$$

$$Z_\ell(r) = \exp \int_r^\infty \psi_\ell(r') dr', \quad F_\ell(k, r) \underset{r \rightarrow \infty}{\sim} e^{-ikr}$$

ψ_ℓ of the type Eq. (6) and F_ℓ satisfied:

$$\left(\frac{d^2}{dr^2} - \frac{\ell(\ell+1)}{r^2} + k^2 \right) F_\ell = \left(-2\psi \frac{d}{dr} + W_\ell \right) F_\ell \left(W_\ell = V - \dot{\psi}_\ell - \psi_\ell^2 \right). \quad (32)$$

1) In the first case we consider briefly the case of short-range potential $V = -Ge^{-\mu r}/r^2$. In this case, (30) is of the Fuchs' type near the origin and the centrifugal potential modifies the power behaviour of the "regular" and "singular" solutions near the origin. In Ref. 1 this case has been investigated with the use of the Laplace transform for Eq. (32). We recall that in Ref. 1 we have defined Z_ℓ in such a way that the Jost function be $f_\ell(k) = \lim_{r \rightarrow 0} r^\ell F_\ell(k, r)$. The corresponding study in coordinate space is straightforward, and we give only some arguments. In Ref. 1 (Eq. 27), the above equation is given explicitly. Then we have the extension of Eq. (7)

$$F_\ell(k, r) = kr h_\ell^{(2)}(kr) + \frac{1}{2ik} \int_r^\infty k^2 r r' \left(h_\ell^{(1)}(kr) h_\ell^{(2)}(kr') - h_\ell^{(2)}(kr) h_\ell^{(1)}(kr') \right) \times$$

$$\times \left(-2\psi(r') \frac{d}{dr'} + W_\ell(r') \right) F_\ell(k, r') dr'. \quad (33)$$

If we iterate (33) each term is, near the origin, singular like $r^{-\ell}$ and

$$f_{\ell}(k) = \sum_{q=0}^{q=\infty} \lim_{r \rightarrow 0} r^{\ell} F_{\ell, q}(k, r)$$

is convergent if $G < (\ell + 1/2)^2$.

2) For $n > 1$ we consider now the family of short-range (2.b) or arbitrary power (27). But in these cases the centrifugal potential is not sufficiently singular in order to modify the singularities of R_{ℓ} near the origin. In other words, the "singular part" and the "regular part" of R_{ℓ} are ℓ independent. Thus we have for all ℓ , the same singular factors:

$$Z_{\ell, \epsilon}(r) = Z_{\ell=0, \epsilon}(r) \quad \text{or} \quad \psi_{\ell=0, \epsilon} = \psi_{\ell, \epsilon}.$$

So, as in the $\ell = 0$ case, consider $R_{\ell}(k, r) = Z_{\ell=0, \epsilon=-1}(r) F_{\ell}(k, r)$ and $F_{\ell}(k, r) \simeq \text{const.}$ as $r \rightarrow 0$. Thus, we consider as in (7) the integral equation

$$f_{\ell}(k, r) = 1 + \frac{1}{2ik} \int_r^{\infty} \left(e^{2ik(r-r')} - 1 \right) \left(-2\psi_{\ell=0}^{(r')} \frac{d}{dr'} + W_{\ell=0}^{(r')} - \frac{\ell(\ell+1)}{r'^2} \right) f_{\ell}(k, r') dr'. \quad (34)$$

We iterate Eq. (34):

$$f_{\ell}(k, r) = \sum_{q=0}^{\infty} f_{\ell, q}(r)$$

and all the previous analysis is still valid. In fact, only for $1 < n \leq 3/2$ and $n = 2$ do we have $W(r)$ and $\ell(\ell+1)/r^2$ of the same order of singularity. In all other cases for the family (2.b) and (27) $\ell(\ell+1)/r^2$ is less singular near the origin than $W(r) \simeq O(1/r^n)$. But we recall (Appendix A) that $W(r)$ plays no role in the method of convergence considered, the essential reason being that it does not affect the behaviour

of the singularities of $F(k, r)$ near the origin (see Appendix of Ref. 2).

Here also the Jost function can be defined by the limit of convergent sequences:

$$f_\ell(k, 0) = \lim_{p \rightarrow 0} \left(\sum_{q=0}^p f_{\ell, q}(r(p), k) \right)$$

for $r(p)$ going to zero less rapidly than the $\ell = 0$ limiting dependence $1/p^{(1/n-1)-\epsilon}$.

IV. BETHE-SALPETER EQUATION

We recall that Domokos³⁾ and Sawyer⁴⁾ have recently investigated the Bethe-Salpeter equation for two fermions in Euclidean-coordinate space. They have reduced their studies of the n -th projection of the Bethe-Salpeter wave function to a differential equation. (We consider always in this paper, the case where the most singular part of the potential is repulsive.) If we consider the form given in Ref. 3.b (Eq. 2) of this differential equation, the only change with the Schrödinger equation (30) is that $\ell(\ell+1)$ must be changed by $(n^2 - 1/4)$. For the family (2.b) or (27) of potentials, the formalism previously developed for $\ell \neq 0$ Schrödinger case is still valid. In particular, if we iterate Eq. (34) (with $\ell + 1/2 \rightarrow n$), we can still construct convergent sequences with the same limiting dependence $r_L(p)$.

V. CONCLUSION

Thus, we have shown in this paper for the potentials (2.b) and (27), that from the perturbative expansion of "new Jost solutions" we can construct the Jost functions by connecting the radial coordinate and the order of the perturbation expansion. We have seen that if we attribute an $r(p)$ dependence, the Jost function is the limit of convergent sequences if $r(p)$ goes to zero less rapidly than $r_L(p)$. In fact (see Appendix A and Section III), this $r_L(p)$ dependence depends mainly on the most singular term of the potential near the origin and, consequently, on the leading essential singularity of the new Jost solution near the origin. Then for other families of potentials where the most singular term near the origin is not like a power $G^2 r^{-2n}$, the corresponding $r_L(p)$ can be different.

Recently^{5,6,7)}, other families of singular potentials (logarithmic) have been studied in coordinate space, mainly in order to test the so-called "peratization technique". In a further paper we will give the extension of the present work to this family of singular potentials⁸⁾. The main thing to investigate for these potentials, in the present method, is to find the correct behaviour near the origin of the singular and regular parts. But, even if for two different families of leading singular repulsive potentials these $r_L(p)$ are different, we can choose in both cases an $r(p) \xrightarrow{p \rightarrow \infty} 0$ dependence such that $r(p) \geq r_{L_1}(p)$ where $r_{L_1}(p)$ is the **highest** of $r_{L_1}(p)$, $r_{L_2}(p)$.

Here, although we have often referred to the previous work (Ref. 2), where the Laplace transform was used, all results and proofs of this paper are obtained directly in coordinate space. However, we thought that it would be interesting to show as much as possible the analogies between the difficulties encountered in both cases, and the similitude of the methods given, in order to resolve these difficulties.

In coordinate space there is another case very similar to the Schrödinger one. It is the coupled Dirac equation which is equivalent to a second order differential equation. The extension of the present work to this case will be investigated later.

From a practical point of view we want to add that it is perhaps interesting now to test formalisms which do not require the introduction of arbitrary cut-off procedures (see also, the recent work of Bertocchi⁹). It is also our hope that we can extend directly to the S matrix some of the methods presented here and in Refs. 1 and 2.

ACKNOWLEDGEMENTS

It is a great pleasure to thank Dr. E. Predazzi for his constant interest in this work.

A P P E N D I X A

We want to investigate the singular part of the remaining sequences $R(p, r, k)$ when p is large and $r \rightarrow 0$. We start from the integral equation (8) and iterate. We recall that for the potential (2.b) it was shown in Ref. 2 that near the origin

$$\psi(r) \underset{r \rightarrow 0}{\simeq} \frac{G}{r^n} - \frac{1}{2} \left(\frac{A_{2n-1}}{G} + \mu G \right) \frac{1}{r^{n-1}} + \dots - \frac{n}{2r} \quad (A.1)$$

The behaviour near the origin of the singular part of the q -th iteration of $f(r)$ is given from (8)

$$f_q(r) \underset{r}{\simeq} \int_r^a (r-r') 2\psi(r') \left(\frac{d}{dr'} + W(r') \right) f_{q-1}(r') dr' \quad (A.2)$$

where a is a small finite constant.

Firstly, for each $f_q(r)$, we consider the leading singular term

$$f_q^{(1)}(r) \simeq C_q r^{-[(q(n-1)+m-n-1)]}$$

and from (A.2 and (A.1) it is easy to find the recurrence relation between C_q and C_{q-1} . We put $\alpha = 1/(n-1)$ and get

$$C_q = \left(\frac{-2G}{n-1} \right)^q \frac{1}{(q-2+(m-2)\alpha) \prod_{j=1}^{q-1} (q-j+(m-1)\alpha)}$$

From (A.7,8) we see that the second singular leading term of the perturbation expansion is connected with the second singularity of $\bar{f}(r,k)$. We note that the correspondence relation (A.6') gives a relation between r and q very similar to the $r_L(p)$ found. This is easily understood because the singular part of the perturbation expansion factorizes the highest essential singularity.

We note also that terms coming from $W(r)$ do not play a role for the most singular terms. Consider for instance n integer, then $m = n$ and $W(r)$ gives for each $f_q(r)$ a singularity in r like $f_q^{(2)}(r)$ as quoted above. However, its q dependence is much smaller than D_q and the equivalent relations (A.6,6') ensure the neglect of the corresponding terms. The considerations given for $W(r)$ are still valid for the approximations made in the kernel of (A.2) when we have taken $\sin k(r-r') \simeq \text{const } (r-r')$. This can be easily verified (see, for instance, the similar case in the Laplace transform Ref. 2 Eq. (3.9 a,b)).

These considerations illustrate the fact that for a given singular term of a certain order we must consider both its r and q dependence. With the help of (A.5,6,6') we can compare two such terms. This was of major importance in Ref. 2 where we investigate the asymptotic behaviour of the Laplace transform $\rho(\alpha)$.

A P P E N D I X B

We want to find the "singular" and the "regular" part of $R(k, r)$ near the origin for the family of singular potentials (27). We start from

$$\left(\frac{d^2}{dr^2} + k^2 - \frac{G^2}{r^{2n}} - \frac{A_{2n-1}}{r^{2n-1}} - \frac{A_{2n-2}}{r^{2n-2}} \dots - \frac{A_{N_0}}{r^{N_0}} \right) R = 0 \quad n > 1, N_0 > 2. \quad (B.1)$$

(B.1) not being of the Fuchs' type, similarly to what was made in the Appendix of Ref. 2, we consider an ansatz of products of essential singularities, and we want to show that we can determine all the parameters

$$R(k, r) \simeq \text{const } r^\alpha \exp \sum_i \left[\frac{\beta_i}{r^{\gamma_i}} \right] \quad (B.2)$$

$$(\gamma_i > 0, \gamma_1 > \gamma_2 \dots).$$

We substitute (B.2) in (B.1) and consider the most singular terms near the origin

$$\begin{aligned} \exp \left(\sum_i \frac{\beta_i}{r^{\gamma_i}} \right) r^{\alpha-2} \left[\sum_i (\gamma_i \beta_i)^2 r^{-2\gamma_i} + 2 \sum_{i \neq j} \gamma_i \gamma_j \beta_i \beta_j r^{-(\gamma_i + \gamma_j)} + \right. \\ \left. + \sum_i (2\alpha - \gamma_i - 1)(-\gamma_i \beta_i) r^{-\gamma_i} + \alpha(\alpha - 1) + \right. \\ \left. + -G^2 r^{-(2n-2)} + A_{2n-1} r^{-(2n-3)} + \dots \right] \simeq 0. \quad (B.3) \end{aligned}$$

We determine the parameters γ_i , β_i , α by the condition that the coefficients of the most singular terms in the bracket of (B.3) are zero.

1) From the most singular term in the bracket we find

$$\begin{cases} \gamma_1 = n - 1 \\ \beta_1 = -\epsilon G / (n - 1) \quad \epsilon = \pm 1 \end{cases}$$

Firstly, we see that the condition $\gamma_1 > 0$ is equivalent to $n > 1$. Secondly, for $\beta_1 > 0$ ($\beta_1 < 0$) corresponds the singular (regular) part of the solution. The corresponding power behaviour of this most singular term in the bracket is of the order $r^{-(2n-2)}$. Now we consider the term of order $r^{-(2n-3)}$ and so on.

2) From the next singular term of the bracket we find

$$\begin{cases} \gamma_2 = n - 2 \\ \beta_2 = A_{2n-1}/2 \in G(n-2) . \end{cases}$$

This gives another essential singularity in (B.2) for $n > 2$.
 If $1 < n \leq 2$ we have only one essential singularity and a power behaviour. We can go on in this method and we find that for $i < n < i+1$ (i integer) we have a product of i essential singularities and a power behaviour. We remark that the term k^2 never comes into the determination of these "singular" or "regular" dependences of the solutions near the origin. Finally we get

$$R(k,r) \underset{r \rightarrow 0}{\simeq} Y_{\text{sing}}(r) \text{ const}_1 + Y_{\text{reg}}(r) \text{ const}_2 .$$

REFERENCES

- 1) H. Cornille, Nuovo Cimento 33, 434 (1964).
- 2) H. Cornille and E. Predazzi, "Singular potentials with short range", CERN Preprint 9125/TH.441. A brief report of this work has been given in Physics Letters 10, 149 (1964).
- 3) a. G. Domokos, P. Suranyi and A. Vancura, Dubna Preprint P1512 (1963).
b. G. Domokos, P. Suranyi and A. Vancura, CERN Preprint 9011/TH.432 (1964).
- 4) R.F. Sawyer, "Non-renormalizability and the short range force in some field theoretic models", Preprint 1963.
- 5) B.A. Arbuzov, A.T. Filippov and O.A. Khrustalev, Physics Letters 8, 205 (1964).
- 6) T.T. Wu, "Scattering by the singular potential $-gr^{-4} \log r$ ", Preprint.
- 7) H.H. Aly, Riazuddin and A.M. Zimmerman, "Singular logarithmic potentials and peratization", Preprint.
"Scattering by singular logarithmic potentials", Preprint.
- 8) H. Cornille and E. Predazzi, to be published.
- 9) L. Bertocchi, S. Fubini and G. Furlan, "On the theory of scattering by singular potentials", CERN Preprint 9564/TH.477.