CM-P00057006

A NUCLEON POTENTIALG. Fast ^{*)}, G. Ranft ⁺⁾ and J.J. de Swart ^{*)}

CERN - Geneva

A B S T R A C T

Potentials to describe the hyperon-nucleon interaction have been calculated some time ago with the prescription of Brueckner and Watson. They treat the Λ and Σ channels on equal footing. We have calculated from them an effective potential for the Λ -nucleon interaction that takes only implicitly the effect of the closed Σ channel into account. This procedure, however, is not restricted to the hyperon-nucleon case, although we only applied it there.

The potentials we used were in accordance with an even $\Lambda - \Sigma$ parity and with unitary symmetry for the pseudo-vector coupling constants, which they contain as parameters.

It is shown that the phase shifts, resulting from the effective ΛN potential, coincide, up to about the threshold for Σ production, with the ones obtained from the original hyperon-nucleon potentials. Calculations were done both for the singlet and triplet cases. In this way, one gets an impression about the effective amounts of central, spin-orbit and tensor force in the ΛN interaction.

^{*)} Present address: University of Nijmegen, Nijmegen, The Netherlands.

⁺⁾ On leave from the Karl-Marx-Universität, Leipzig.

I. INTRODUCTION

Consider the elastic scattering of two particles A and B. When one describes the interaction with a central potential, one has to solve a radial Schrödinger equation for every channel characterized by a specific value of ℓ , the orbital angular momentum. If there is not only a central potential but also a tensor potential between the two spin $1/2$ particles, the two channels with $\ell = J \pm 1/2$ (where J is the total angular momentum) will be coupled. This is the case for example in the low energy nucleon-nucleon scattering.

If the two particles can scatter not only into the final state A + B, but also into a final state with particles C and D, then again one obtains sets of coupled radial equations. This is for example the case in lambda-nucleon scattering. The final state may be here a channel with a nucleon and a sigma hyperon, if sufficiently high energy lambdas are used.

Suppose now that one has N coupled channels, but that some of them are closed (because their thresholds are higher than the available energy).

Clearly the closed channels will still influence the scattering into the open ones and one still should solve N coupled equations, only the closed channels have a damped wave as boundary condition. An example of this is again the Λ nucleon scattering below the threshold for Σ production.

Solving a set of N coupled second order differential equations, which describe such a system of two-body scattering channels, is quite laborious but possible. In many other cases, however, especially when more particles are present, the additional complications of the coupled channels are a little

too much to handle at present. Examples are the interactions of Λ hyperons in hyperfragments, the final state interactions in processes like $K^- + d \rightarrow Y + N + \pi$, etc.

For a system of coupled open and closed channels, where the coupling can conveniently be described by potentials, we have tried to find a way of re-defining the potentials for the open channels such that one can completely ignore the closed channels. The effective potential so constructed takes, then implicitly the effect of the closed channels into account.

As a model to demonstrate the procedure, we take the case of lambda-nucleon scattering below the Σ threshold. The ΛN channel is strongly coupled to the ΣN channel. Potentials, which treat both channels on an equal footing, have been calculated ^{1),2)}. They come from a Brueckner-Watson type of calculation ³⁾ of the one and two pion exchange potentials, to which a hard core is added. The coupling constants, which these potentials contain as parameters, are chosen in accordance with unitary symmetry for the pseudo-vector coupling constants. The hard cores are so adjusted as to give the Λ nucleon scattering lengths as determined from the analyses of the light hyperfragments ^{4),5),6)}. Using these potentials, one can calculate all scattering data by solving the multi-channel Schrödinger equation numerically ^{7),8)}. It is from these potentials, that treat the ΛN and ΣN channels on an equal footing, that we shall construct an effective one channel ΛN potential, that gives the same low energy scattering parameters as the original potentials. Such a procedure is not unique. We choose, therefore, the simplest ^{*}) formula that gives the correct scattering length and the correct effective range. The effective potential so obtained turns out to be energy

^{*}) There are, of course, many simple ways to obtain the wanted result. What we call the simplest is therefore very subjective.

and ℓ dependent. It will be seen, however, that the dependence of this effective potential on ℓ and E is small. Actually, the effective potential obtained at $E = 0$ for $\ell = 0$ reproduces the scattering data for all ℓ waves up to about the threshold for Σ production.

In section II we will treat the singlet Λ nucleon scattering and we show how to obtain the effective potential. In section III we treat the more complicated case of triplet Λ nucleon scattering with $\ell = J \pm 1/2$. The numerical method is given in section IV and in section V we give and discuss the results.

II. EFFECTIVE Λ N POTENTIAL

We shall consider first the case of Λ N scattering in the singlet state ($S = 0$). It is well known that the Λ N channel is strongly coupled to the $T = 1/2$ Σ N channel, as can be seen from the existence of the reaction $\Lambda + N \rightarrow \Sigma + N$. This means that a description of the low energy Λ N scattering with a one-channel Schrödinger equation ⁹⁾

$$\frac{1}{2m_\Lambda} \chi'' + \left(E_\Lambda - V - \frac{1}{2m_\Lambda} \frac{L^2}{r^2} \right) \chi = 0 \quad (1)$$

has to break down for E_Λ (lab) > 167 MeV (the threshold for Σ production) and possibly much earlier.

A more realistic way to describe the Λ N interaction is, therefore, to take the Σ N channel explicitly into account. That can be done by describing the scattering with two coupled Schrödinger equations, which we shall write in matrix notation ⁷⁾:

$$\frac{1}{2m} \psi'' + \left(E - V - \frac{1}{2m} \frac{L^2}{r^2} \right) \psi = 0 \quad (2)$$

Here, ψ is a column vector, while E , M and V are matrices:

$$\psi = \begin{pmatrix} u_\Lambda \\ u_\Sigma \end{pmatrix}, \quad E = \begin{pmatrix} E_\Lambda & 0 \\ 0 & E_\Lambda - \Delta M \end{pmatrix}, \quad m = \begin{pmatrix} m_\Lambda & 0 \\ 0 & m_\Sigma \end{pmatrix},$$

and

$$V = \begin{pmatrix} V_{\Lambda\Lambda} & V_{\Lambda\Sigma} \\ V_{\Sigma\Lambda} & V_{\Sigma\Sigma} \end{pmatrix} \quad (3)$$

with u_Λ and u_Σ being the radial wave functions, m_Λ and m_Σ the reduced masses in the Λ N and Σ N channels; $\Delta M = M_\Sigma - M_\Lambda$ is the difference between the Λ and Σ rest masses.

In solving (2) for an energy below the Σ threshold, one has to impose the boundary conditions:

$$\psi(x_0) = 0$$

$$\psi(r) \rightarrow k_\lambda^e \begin{pmatrix} k_\lambda r j_\ell(k_\lambda r) \cot \delta_\ell - k_\lambda r n_\ell(k_\lambda r) \\ b_\ell i k_\Sigma r h_\ell^{(1)}(i k_\Sigma r) \end{pmatrix} \text{ if } r \rightarrow \infty \quad (4)$$

where

$$E_\lambda = \frac{k_\lambda^2}{2m_\lambda} = \Delta M - \frac{k_\Sigma^2}{2m_\Sigma}$$

and where j_ℓ , n_ℓ and h_ℓ are the spherical Bessel, Neumann and Hankel functions. x_0 is the singlet hard core radius. Our problem will be: given a potential V for the two-channel case (3), to find an effective potential V_{eff} , that, when used in the one-channel equation (1) will reproduce as well as possible the phase shifts δ_ℓ obtained with the two-channel potential V and equation (2).

The solution is the following: with the given potential V we can solve the two-channel equation (2) and thus obtain the wave function Ψ . From this Ψ we then construct the wave function χ for the one-channel case. This χ will give the same phase shift if it satisfies $\chi \rightarrow u_\lambda$ if $r \rightarrow \infty$; furthermore, one wants $\chi(x_0) = 0$ to make it a reasonable radial wave function. Clearly, the simple choice $\chi = u_\lambda$ would give the proper phase shift at the same energy for which we have solved equation (2); it would not, however, give the same effective range as in the two-channel case. Therefore, we take χ to be

$$\chi = \left(1 + \frac{u_\Sigma^2}{u_\Lambda^2}\right)^{1/2} u_\Lambda \quad (5)$$

it will then also give the same effective range at the energy considered. From (1) we conclude then:

$$V_{\text{eff}} = \frac{1}{2m_\Lambda} \frac{\chi''}{\chi} - \frac{L^2}{2m_\Lambda r^2} + E_\Lambda \quad (6a)$$

or using (2) and (5):

$$\begin{aligned} V_{\text{eff}} = & \frac{u_\Lambda^2}{u_\Lambda^2 + u_\Sigma^2} V_\Lambda + \frac{2u_\Lambda u_\Sigma}{u_\Lambda^2 + u_\Sigma^2} \frac{m_\Lambda + m_\Sigma}{2m_\Lambda} V_{\Lambda\Sigma} \\ & + \frac{u_\Sigma^2}{u_\Lambda^2 + u_\Sigma^2} \frac{m_\Sigma}{m_\Lambda} (V_{\Sigma\Sigma} + \Delta M) \\ & + E_\Lambda \left(1 - \frac{m_\Sigma}{m_\Lambda} \frac{u_\Sigma^2}{u_\Lambda^2 + u_\Sigma^2} - \frac{u_\Lambda^2}{u_\Lambda^2 + u_\Sigma^2}\right) \end{aligned} \quad (6b)$$

We note here that this method is limited to cases where u_Λ has no zero points in the range where the potentials are important. The ΛN system has no bound state and the method is therefore applicable for low energies.

We still have to show that the choice (5) really yields the right effective range. Therefore we note that the effective range for the one-channel case is given by ⁸⁾

$$R_e = 2 \int_0^\infty (\sqrt{v^2 - \chi^2} - F_e) dr \quad (7)$$

where $v = k_{\Lambda}^l (k_{\Lambda} r j_l(k_{\Lambda} r) \cot \delta_l - k_{\Lambda} n_l(k_{\Lambda} r))$ and F_l ⁸⁾ subtracts out the pole terms in the integrand resulting from the Neumann function in v . For the two-channel case, we find

$$R_e = 2 \int_0^{\infty} (v^2 - u_{\Lambda}^2 - u_{\Sigma}^2 - F_e) dr \quad (8)$$

a formula which holds for every l .

Comparing (7) and (8), it is clear that the choice (5) indeed gives the right result.

III. THE TRIplet CASE

For scattering in the triplet state ($S = 1$), things are a bit more complicated. The states with the orbital angular momentum $\ell = J \pm 1$ are coupled by the tensor force; J being the total angular momentum.

The radial wave equation describing the ΛN scattering without considering the ΣN channel, however, can still be written in the form (1), where now χ , V , and L^2 are 2×2 matrices. L^2 is diagonal with elements $\ell(\ell + 1)$.

Likewise, taking again the ΣN channel into account, (2) can retain its old form; however, Ψ must now be interpreted as a 4×2 matrix of which the odd rows refer to channels with $\ell = J - 1$ and the even ones to those with $\ell = J + 1$. The first two rows will correspond to the ΛN channels and the last two to the ΣN channels. Correspondingly, m , V , E and L^2 are now 4×4 matrices.

The boundary conditions for (2) are now

$$\Psi(x_1) = 0$$

$$\Psi \rightarrow \begin{pmatrix} v \\ w \end{pmatrix} \quad \text{if } r \rightarrow \infty$$

where $v = k_\Lambda^\ell (JK^{-1} - N)$ and $w = HB$. Here J is a diagonal matrix with elements $k_\Lambda r j_\ell(k_\Lambda r)$, while N and H are defined analogously in terms of the functions n_ℓ and $h_\ell^{(1)}$. x_1 is the radius of the triplet hard core.

K^{-1} is the inverse of the reaction matrix, it has the eigen values $\cot \delta_{\ell J}$, where $\delta_{\ell J}$ are the eigen phase shifts. The matrix B is a constant matrix. The effective range R and the scattering length A are now 2×2 matrices defined by

$$k_{\lambda}^{l+1/2} K^{-1} k_{\lambda}^{l+1/2} = -A^{-1} + \frac{1}{2}(k_{\lambda}^2 - k_0^2) R \quad (9)$$

where $E_0 = \frac{k_0^2}{2m_{\lambda}}$ is the energy at which we perform the effective range expansion. The effective ranges are given for the two-channel case (1) by

$$R = 2 \int_0^{\infty} (v^T v - \chi^T \chi - F) dr \quad (10a)$$

and for the four-channel case (2) by

$$R = 2 \int_0^{\infty} (v^T v - u_{\lambda}^T u_{\lambda} - u_{\Sigma}^T u_{\Sigma} - F) dr \quad (10b)$$

Here F is a diagonal matrix, whose diagonal elements are equal to the functions F_{ℓ} mentioned in the singlet case.

To obtain the correct scattering length, χ has again to fulfil the condition

$$\chi \rightarrow u_{\lambda} \quad \text{if} \quad r \rightarrow \infty \quad (11)$$

while to obtain the correct effective range we need

$$\chi^T \chi = u_{\lambda}^T u_{\lambda} + u_{\Sigma}^T u_{\Sigma} \quad (12)$$

The conditions (11) and (12) are fulfilled if we take

$$\chi = \left(1 + u_{\Lambda}^T u_{\Sigma}^T u_{\Sigma} u_{\Lambda}^{-1}\right)^{1/2} u_{\Lambda} \quad (13)$$

The potential V_{eff} is then given by

$$V_{\text{eff}} = \frac{1}{2m_{\Lambda}} \chi'' \chi^{-1} - \frac{L^2}{2m_{\Lambda} r^2} + E_{\Lambda} \quad (14)$$

Unfortunately, this V_{eff} will not in general fulfil the requirement imposed by time reversal invariance

$$V_{\text{eff}} = V_{\text{eff}}^T.$$

This condition is equivalent to the Wronskian condition on the wave function χ

$$\chi^T \chi' - \chi'^T \chi = 0 \quad (15)$$

Time reversal invariance for the four-channel case gives

$$m_{\Lambda} u_{\Lambda}^T u'_{\Lambda} + m_{\Sigma} u_{\Sigma}^T u'_{\Sigma} - m_{\Lambda} u'_{\Lambda} u_{\Lambda} - m_{\Sigma} u'_{\Sigma} u_{\Sigma} = 0$$

We were unable to find an expression for χ in terms of u_{Λ} and u_{Σ} , which satisfies at the same time the conditions (11), (12) and (15). Therefore we calculated V_{eff} via (13) and afterwards symmetrized it by taking for both the new off diagonal elements half the sum of the old ones.

Although the conditions (11) and (12) are then no longer rigorously fulfilled, their violation turned out to be negligible in the cases considered.

The effective potential being a 2×2 matrix, we have chosen to represent in the following way ⁷⁾

$$V = V_c + V_T S_{12} + V_{so} \underline{L} \cdot \underline{S} + V_{LL} Q_{12} \quad (16)$$

where

$$S_{12} = \frac{3(\underline{\sigma}_1 \cdot \underline{r})(\underline{\sigma}_2 \cdot \underline{r})}{r^2} - \underline{\sigma}_1 \cdot \underline{\sigma}_2$$

$$Q_{12} = \frac{(\underline{\sigma}_1 \cdot \underline{L})(\underline{\sigma}_2 \cdot \underline{L}) + (\underline{\sigma}_2 \cdot \underline{L})(\underline{\sigma}_1 \cdot \underline{L})}{2} - (\underline{\sigma}_1 \cdot \underline{\sigma}_2) L^2$$

The functions V_c , V_T , V_{so} and V_{LL} give the radial dependences of the central, tensor, spin-orbit and quadratic spin-orbit potentials. They will in general be ℓ dependent. From the effective potentials, say for the cases 3D_1 , 3D_2 and 3D_3 one obtains via (16) the radial functions. By definition we took the off diagonal element of the ${}^3D_3 - {}^3G_3$ effective potential as $V_T(D)$. Likewise $V_T(P)$ was identified with ${}^3P_2 - {}^3F_2$.

IV. NUMERICAL METHOD

As a starting point for our calculations we used the hyperon-nucleon potentials $V_{\Lambda\Lambda}$, $V_{\Lambda\Sigma} = V_{\Sigma\Lambda}$ and $V_{\Sigma\Sigma}$ calculated according to the Brueckner-Watson prescription ^{1),2),3)}. These potentials have as parameters the coupling constants $f_{NN\pi}$, $f_{\Lambda\Sigma\pi}$ and $f_{\Sigma\Sigma\pi}$, and the hard core radii: X_0 for the singlet states and X_1 for the triplet states. For the coupling constants we choose values consistent ^{8),10)} with the experimental data obtained in the analysis of the light hyperfragments, the capture of Σ^- hyperons at rest in hydrogen and deuterium and with the analysis of the recently obtained double hyperfragment $\Lambda\Lambda\text{Be}^{10}$. The coupling constants are

$$f_{NN\pi} = 0.285$$

$$f_{\Sigma\Sigma\pi} = 0.100$$

$$f_{\Lambda\Sigma\pi} = 0.271$$

These values are in accordance with unitary symmetry for the pseudo-vector coupling constants. The cores $X_0 = 0.3064 \mu^{-1}$ and $X_1 = 0.3550 \mu^{-1}$ are so chosen as to reproduce the nucleon scattering lengths.

For the nucleon, hyperon and pion masses we used the values

$$M_N = 938.9 \text{ MeV}/c^2$$

$$M_\Lambda = 1115.4 \text{ MeV}/c^2$$

$$M_\Sigma = 1192.3 \text{ MeV}/c^2$$

$$\mu = 138.1 \text{ MeV}/c^2$$

The wave equations (2) were solved numerically on a high-speed electronic computer using the Noumanoff method ⁷⁾. This

was done for all S, P and D waves and for several energies using the potentials mentioned above. Thus the corresponding phase shifts and wave functions were obtained. From the "zero" energy ($E_{\Lambda}(\text{lab}) = 0.1 \text{ MeV}$) solutions the effective potentials were then calculated according to the formula (16). The wave equation (1) was then solved using these effective potentials. In all cases we reproduced the scattering lengths and effective ranges within the accuracy of the calculation.

V. RESULTS

The calculated effective potentials are exhibited in figures 1 and 2. For convenience, we have drawn them on a normal scale in figure 1 and on a logarithmic scale in figure 2. In the singlet states it was found that there was a small difference in the effective potential for the 1S_0 and 1P_1 states. This difference is too small, however, to be shown properly in our graphs. The effective potentials in the 1P_1 , 1D_2 and 1F_3 states are essentially the same. It has been pointed out by Dalitz ¹⁰⁾ that the influence of the Σ channel is quite unimportant in the singlet case for small values of $f_{\Sigma\pi}$. Therefore, it is interesting to compare the effective potential with $V_{\Lambda\Lambda}$ for $S = 0$. We found that these potentials are indeed quite similar; their difference is again too small to show in our graphs. The effective potential however has a slightly weaker tail than $V_{\Lambda\Lambda}$, while its innermost part is somewhat stronger. The effective potential is, therefore, somewhat stronger and of shorter range than $V_{\Lambda\Lambda}$.

In the triplet states it was found that there was a small difference between the S, P and D central potentials, too small to draw. The triplet central potential looks quite similar to the singlet potential as can be seen in the graphs ¹¹⁾. It has, of course, a larger core than in the singlet case. The tensor potential V_T is rather weak and has a slight ℓ dependence. It has an attractive inner part followed by a repulsive tail. The strength of the tail increases with ℓ . The spin-orbit potential V_{so} is rather weak and repulsive and almost independent of ℓ .

The quadratic spin-orbit potential V_{LL} is very weak and enormously ℓ dependent. It is important to stress that these potentials will of course have all the deficiencies of the potentials $V_{\Lambda\Lambda}$, $V_{\Lambda\Sigma}$ and $V_{\Sigma\Sigma}$ which were used as input.

The input potentials for instance contained no L.S force, so one should not expect the L.S force we find, and which is generated by the closed Σ channel, to be even slightly realistic.

It is well known in the nucleon-nucleon case that there the L.S force plays an important rôle. These deficiencies of the input potentials could perhaps be overcome if one includes the exchange of vector mesons, K mesons and scalar mesons, instead of only one and two pion exchange contributions.

For several values of the energy and l we have calculated the phase shifts and effective ranges as well with the original potentials, as with the effective ones. The results are shown in Tables 1 to 4. One sees that the values obtained with the original potentials (1) coincide up to about threshold with the ones obtained with the effective potential (2). We may thus conclude that the effective potentials reproduce the scattering data rather well.

ACKNOWLEDGEMENTS

We would like to express our thanks to the DD Division in CERN for running the programmes.

One of the authors (G.R.) wishes to thank Professor Weisskopf and Professor Van Hove for their hospitality at CERN.

	$\ell = 0$		$\ell = 1$		$\ell = 2$		$\ell = 3$	
E	$\delta_0(2)$	$\delta_0(1)$	$\delta_1(2)$	$\delta_1(1)$	$\delta_2(2)$	$\delta_2(1)$	$\delta_3(2)$	$\delta_3(1)$
0,1	7,22	7,22						
1,0	21,1	21,1	0,02	0,02				
5,0	36,5	36,5	0,20	0,20				
10,0	41,6	41,5	0,57	0,57				
25,0	43,6	43,0	2,13	2,12	0,05	0,05		
50,0	40,4	40,0	5,51	5,49	0,23	0,23	0,02	0,02
100,0	32,6	31,5	13,0	12,9	0,93	0,92	0,10	0,10
150,0	26,1	24,1	19,8	19,1	1,97	1,94	0,25	0,24
160,0	25,1	22,8	20,9	20,2	2,21	2,17	0,28	0,28
167,0	24,6	21,8	21,8	20,9	2,39	2,34	0,31	0,30

TABLE 1

The singlet phase shifts δ_ℓ in degrees for different values of E_Λ in MeV and ℓ for the cases (2) and (1).

J = 1						
E	$\delta_{01}(1)$	$\delta_{21}(1)$	$\epsilon_1(1)$	$\delta_{01}(2)$	$\delta_{21}(2)$	$\epsilon_1(2)$
0,1	1,06			1,06		
1,0	3,29		- 0,03	3,29		- 0,03
5,0	6,64		- 0,18	6,64		- 0,18
10,0	8,30	0,02	- 0,36	8,29	0,02	- 0,36
25,0	9,25	0,17	- 0,99	9,13	0,17	- 1,01
50,0	7,25	0,86	- 2,61	6,85	0,81	- 2,85
100,0	1,04	4,80	3,46	- 0,14	3,81	6,35
160,0	- 6,02	26,9	- 4,54	- 8,29	10,0	1,25
167,0	- 6,72	41,3	- 6,14	- 9,19	10,9	1,10

TABLE 2a

The triplet phase shifts $\delta_{\ell J}$ in degrees for different values of E_{Λ} in MeV and ℓ and J, and the coupling parameters ϵ_J for the cases (2) and (1).

J = 2						
E	$\delta_{12}(1)$	$\delta_{32}(1)$	$\epsilon_2(1)$	$\delta_{12}(2)$	$\delta_{32}(2)$	$\epsilon_2(2)$
1,0	0,01		- 0,02	0,01		- 0,02
5,0	0,16		- 0,10	0,16		- 0,12
10,0	0,44		- 0,20	0,44		- 0,23
25,0	1,55	0,01	- 0,49	1,55	0,01	- 0,52
50,0	3,60	0,05	- 0,96	3,56	0,05	- 0,94
100,0	7,01	0,38	- 2,01	6,81	0,32	- 1,82
160,0	9,28	1,88	- 4,42	8,75	1,07	- 3,16
167,0	9,44	2,34	- 5,10	8,87	1,18	- 3,35

TABLE 2b

ℓ	R(1)	R(2)
0	1.35	1.35
1	4.06	4.07
2	278	280

TABLE 3

The singlet effective range R in pion - Compton wavelengths for S, P and D scattering for both cases (2) and (1).

	R(1)	R(2)
R_{11}	5.59	5.59
$R_{12} = R_{21}$	2.11	2.13
R_{22}	62.8	65.3

TABLE 4

The triplet effective range matrix R_{ij} in pion - Compton wavelengths for $J = 1$ for cases (2) and (1).

REFERENCES

- 1) D.B. Lichtenberg and M.H. Ross, Phys. Rev. 107, 1714 (1957).
- 2) J.J. de Swart and K. Iddings, Phys. Rev. 128, 2810 (1962).
- 3) K.A. Brueckner and K.M. Watson, Phys. Rev. 92, 1023 (1953).
- 4) R.H. Dalitz and B.W. Downs, Phys. Rev. 110, 958 (1958);
111, 967 (1958); 114, 593 (1959).
- 5) K. Dietrich, R. Folk and H.J. Mang in Proceedings of the
Rutherford Jubilee International Conference, Manchester,
1961.
- 6) J.J. de Swart, Physics Letters 5, 58 (1963).
- 7) J.J. de Swart and C. Dullemond, Ann. of Phys. 16, 263 (1961).
- 8) J.J. de Swart and C. Dullemond, Ann. of Phys. 19, 458 (1962).
- 9) We use units such that $\hbar = c = 1$.
- 10) R.H. Dalitz, Physics Letters 5, 53 (1963).
- 11) R. Schrills and B.W. Downs, Phys. Rev. 131, 390 (1963).

FIGURE CAPTIONS

Figure 1 The effective potentials.

Figure 2 Absolute values of the effective potentials.

FIG. 1

FIG. 2