Analysis of CP -Noninvariant Interactions and the K_1^0, K_2^0 System*

T. D. LEE

Department of Physics, Columbia University, New York, New York

AND

L. WOLFENSTEIN†

CERN, Geneva, Switzerland

(Received 10 February 1965)

In order to explain the recent observation of $K_2^0 \rightarrow \pi^+ + \pi^-$ by Christenson *et al.*, we assume that the relevant interaction Hamiltonian can be separated into two parts: $H_G + H_F$, where H_G is the usual CP -conserving and T -invariant weak interaction and H_F is a new interaction which is not invariant under CP . Depending on the selection rule of H_F with respect to the strangeness quantum number S , this new interaction H_F can be much stronger than H_G , or somewhat weaker than H_G , or very much weaker than H_G . The properties of these three different classes of possibilities are discussed and some possible experimental tests are suggested. The system of K_1^0 and K_2^0 is analyzed for these different possibilities without assuming the validity of CPT invariance. Efforts are made to separate out the various experimental consequences that follow from different symmetry requirements which may be satisfied by this new CP -noninvariant interaction H_F .

I. GENERAL DISCUSSIONS

RECENTLY, Christenson *et al.*¹ observed that the long-lived component of the K^0 , the K_2^0 , decays into two π mesons with a rate given by

$$|\epsilon| \equiv [\text{Rate}(K_2^0 \rightarrow \pi^+ + \pi^-) / \text{Rate}(K_1^0 \rightarrow \pi^+ + \pi^-)]^{1/2} \cong 2.2 \times 10^{-3}. \quad (1)$$

Throughout this paper we assume that these reactions can occur in a vacuum which has a definite $CP = +1$ value.² The operators C and P denote, respectively, the charge conjugation and the space inversion which are defined by the strong and the electromagnetic interactions. Thus, the final state $(\pi^+ + \pi^-)$ must have a definite $CP = +1$ value. Since out of the states K^0 and $\bar{K}^0$ only one initial state with $CP = +1$ can be constructed, the observation that both K_1^0 and K_2^0 can decay into $(\pi^+ + \pi^-)$ shows that CP cannot be conserved in the $K_{\pi 2}$ decay. If CPT invariance is assumed, then the time-reversal invariance must also be violated in the $K_{\pi 2}^0$ decay. From the magnitude of the observed value $|\epsilon|$ and the already existing experimental limits on the possible CP violation in other weak processes,³ it appears that the amount of either CP violation or the

time-reversal violation, at least in these particular reactions, is relatively small.

We assume that it is possible to separate the relevant interaction Hamiltonian into two parts:

$$H_G + H_F, \quad (2)$$

where H_G is the usual weak interaction which satisfies the $\Delta S \neq \pm 2$ rule and is invariant under CP and T separately, where S is the strangeness quantum number and T is the time reversal operator. The remainder H_F is a new interaction which does not conserve CP , nor is it invariant under T . For order-of-magnitude estimations, we may regard H_G and H_F as characterized, respectively, by the coupling constants G and F where G can be either the Fermi coupling constant G_V in β decay, or the usual strangeness-nonconserving weak coupling constant $\sim \frac{1}{4}G_V$, and F is a new coupling constant which has the same dimension as G_V .

Clearly, with only one observed CP -violating reaction, a great variety of different CP -noninvariant Hamiltonian H_F can be proposed. It is important to notice that, at present, not only is the detailed form of H_F unknown, but even the approximate order of magnitude of its coupling constant F is limited only within an extremely wide range by the existing experiments. As we shall see, there are three very different classes of possibilities depending on the particular selection rule that H_F satisfies with respect to the strangeness quantum number S . These three classes are listed in Table I. Some specific properties of these three classes of possibilities will be discussed in the following:

(1) If H_F satisfies the $\Delta S = 0$ rule and if it does not contain any leptonic operators, then (F/G) can be $\approx 10^3$. Thus, H_F is much stronger than the usual weak interaction H_G . The dimensionless coupling constant of H_F is given by (within a factor ~ 10)

$$(4\pi)^{-1} F m_p^2 \approx |\epsilon| \cong 2.2 \times 10^{-3}, \quad (3)$$

* This research was supported in part by the U. S. Atomic Energy Commission.

† National Science Foundation Postdoctoral Fellow on leave from the Carnegie Institute of Technology, Pittsburgh, Pennsylvania.

¹ J. H. Christenson, J. W. Cronin, V. L. Fitch, and R. Turlay, Phys. Rev. Letters **13**, 138 (1964).

² The alternative possibility has been considered elsewhere. See, e.g., J. Bernstein, N. Cabibbo, and T. D. Lee, Phys. Letters **12**, 146 (1964); J. S. Bell and J. K. Perring, Phys. Rev. Letters **13**, 348 (1964); T. D. Lee, Phys. Rev. (to be published); F. Gürsey and A. Pais (unpublished).

³ M. T. Burg, V. E. Krohn, T. B. Novey, G. R. Ringo and V. L. Telegdi, Phys. Rev. Letters **1**, 324 (1958). See, also, J. W. Cronin and O. E. Overseth, in *Proceedings of the 1962 Conference on High Energy Physics at CERN*, edited by F. J. M. Farley and M. E. Meyer (North-Holland Publishing Company, Amsterdam, 1963), p. 453.

TABLE I. Three different classes of CP -noninvariant interaction H_F .

F/G (within a factor ~ 10)	Conditions satisfied by H_F	Remarks
$\sim 10^3$	$\Delta S=0$ transitions only P is conserved C invariance violated T invariance violated	Both strong and weak processes have C -noninvariant and T -noninvariant amplitudes which are $\sim \epsilon $ times the C -, T -invariant amplitudes.
$\sim 10^{-2}$ – 10^{-3}	$\Delta S=\pm 1$ transitions allowed $\Delta S=\pm 2$ transitions forbidden CP invariance violated	Only weak processes have CP -violating amplitudes.
$\sim 10^{-9}$	$\Delta S=\pm 2$ transitions allowed CP invariance violated	Neglecting terms of $O(10^{-9})$, but keeping those of $O(\epsilon)$, only $K_{1,2}^0$ decays violate CP invariance. These violations are due entirely to the mass operator of the $K_{1,2}^0$ system.

where m_p is the mass of the proton. Assuming that CPT invariance holds,^{4,5} the observation of the $(\pi^+\pi^-)$ decay mode of K_2^0 and the present limit⁶ on the electric dipole moment of the neutron require that H_F must conserve P but violate C invariance and T invariance. The question whether H_F may or may not contain terms that violate the conservation of isotopic spin $\mathbf{I}$ can be left open. We note that if H_F satisfies the $|\Delta\mathbf{I}|=0$ rule, then it is convenient to regard H_F as a small C -, or T -, noninvariant part of the usual strong interaction; otherwise, it is more convenient to regard H_F as a completely new interaction independent of the existing strong interactions.

The decay of K_2^0 to 2π , as well as other CP , or T , violating weak processes, can occur through the second-order expression $H_G H_F$, which contains terms that violate the strangeness conservation, P invariance, C invariance, PC invariance, and T invariance. The possible existence of such a H_F can also be tested by examining the limits of T invariance and C invariance in the strong interactions. If Eq. (3) holds, then all nuclear matrix elements could have a small admixture of T -noninvariant amplitude which is $\sim |\epsilon|$ times the T invariant amplitude. Such a T -noninvariant amplitude can be observed by studying, e.g., the E2, M1 interference terms in an appropriate nuclear transition.⁷

Another possible test of such a relatively strong H_F was pointed out by Friedberg, Lee, and Schwartz.⁸ The decay of η^0 is known to violate the G parity. In

particular, the decay mode

$$\eta^0 \rightarrow 3\pi \quad (4)$$

can occur⁹ via the virtual electromagnetic interactions to a final (3π) state with $I=1$ and $C=+1$. It can also decay, now, through H_F to a final state with $C=-1$ and $I=0$, or 2, with a relative amplitude $\sim r(\epsilon/\alpha)$, where α is the fine structure constant and r is a reduction factor depending on the final-state pion interactions and the angular momentum barrier effects. The approximate order of magnitude of r is expected to be $\lesssim 1$, if H_F contains a $|\Delta\mathbf{I}|=2$ part; otherwise, it could be much smaller than 1. The interference between the $C=+1$ amplitudes can result in an asymmetry between π^+ and π^- in the Dalitz plot of the final $(\pi^+\pi^-\pi^0)$ distribution. Because of the known strong pion interactions in the final states, such an asymmetry does not violate CPT invariance. The detection of such an asymmetry would be an unequivocal proof of violation of C invariance in the η^0 decay; it would also establish that the C -noninvariant interaction H_F is much stronger than the usual weak interaction H_G .

(2) If H_F allows the $\Delta S=\pm 1$ transitions, but forbids the $\Delta S=\pm 2$ transitions, then F must be of the same order of magnitude as, or somewhat smaller than, G . (The H_F may contain, in addition, also some $\Delta S=0$ transition amplitudes.) If the observed slow rate¹ of $K_2^0 \rightarrow \pi^+\pi^-$ is regarded as a typical example of CP noninvariance, then F should be smaller than G and, within a factor ~ 10 , is given by

$$F \approx 10^{-2}G - 10^{-3}G. \quad (5)$$

Such a H_F can be easily described by assigning some appropriate phase factors to the various coupling constants in the usual weak interactions. The general observable effects in such a case have already been discussed in the literature.¹⁰ Recently, in connection

⁴ The general theoretical arguments for expecting CPT invariance have been given by W. Pauli, *Niels Bohr and the Development of Physics* (Pergamon Press, Inc., London, 1955) and J. Schwinger, *Phys. Rev.* **91**, 720, 732 (1953); **94**, 1366 (1953). See, also, G. Lüders, *Kgl. Danske Videnskab. Selskab Mat. Fys. Medd.* **28**, No. 5 (1954).

⁵ If H_F is not invariant under CPT , then there would be a mass difference between any strongly interacting particle and its anti-particle; e.g., $\langle K^0 | H_F | K^0 \rangle$ could be different from $\langle \bar{K}^0 | H_F | \bar{K}^0 \rangle$. Such a difference is clearly ruled out if F is $\approx 10^3 G$.

⁶ E. M. Purcell and N. F. Ramsey, *Phys. Rev.* **78**, 807 (1950); N. F. Ramsey, *Molecular Beams* (Oxford University Press, London, 1956).

⁷ See, e.g., B. Jacobsohn and E. Henley, *Phys. Rev.* **113**, 225 (1959).

⁸ R. Friedberg, T. D. Lee, and M. Schwartz (unpublished).

⁹ If H_F contains a C -conserving but $|\Delta\mathbf{I}|=1$, or 3, part then η^0 can also decay via H_F to a 3π final state with $C=+1$ and $I=1$, or 3.

¹⁰ See, e.g., T. D. Lee and C. N. Yang, Brookhaven National Laboratory BNL 443 (T-91), 1957, Chaps. IV and V (unpublished).

with the observation¹ of Christenson *et al.*, many specific models have also been proposed.¹¹ In this paper, we will not investigate further this well-studied case.

(3) The final possibility is that H_F may be very much weaker¹² than H_G ; i.e.,

$$F \approx 10^{-9}G \quad (6)$$

or

$$F \approx 10^{-15}m_p^{-2}. \quad (7)$$

In this case, H_F must allow a $\Delta S = \pm 2$ transition and, in particular, the matrix element

$$\langle \bar{K}^0 | H_F | K^0 \rangle \neq 0. \quad (8)$$

It will be shown that, in this case, to the accuracy of keeping $|4\pi F/G^2 m_p^2| \approx |\epsilon|$, but neglecting $(F/G) \sim 10^{-9}$ as compared to 1, the CP -violating effects in all known weak processes are *already completely determined*, provided the validity of the CPT theorem⁴ is assumed; otherwise, an additional new real parameter is needed. We note that if one neglects $F/G \sim 10^{-9}$, then in all known weak processes, except those involving K_1^0 and K_2^0 , there is no observable CP -violating effect. For the K_1^0 and K_2^0 states, terms of the order F/G^2 can occur, but only through the mass operator. Assuming the validity of the CPT theorem, the eigenstates K_1^0 and K_2^0 , each of which has a definite lifetime, are given by¹³

$$K_1^0 = (1 + |\epsilon|^2)^{-1/2} [(1 + \epsilon)K^0 + (1 - \epsilon)\bar{K}^0] \quad (9)$$

and

$$K_2^0 = (1 + |\epsilon|^2)^{-1/2} [(1 + \epsilon)K^0 - (1 - \epsilon)\bar{K}^0], \quad (10)$$

where ϵ is a complex parameter,

$$\epsilon = |\epsilon| e^{i\delta}. \quad (11)$$

The magnitude of ϵ is proportional to F/G^2 . As will be shown in Sec. III, to the accuracy of neglecting F/G , the parameter ϵ has already been measured. The absolute magnitude of ϵ is given by Eq. (1), and the phase δ is given by

$$\tan \delta \approx -2[(m_1 - m_2)/(\gamma_1 - \gamma_2)], \quad (12)$$

where m_i and γ_i are the observed mass and width of K_i^0 ($i=1$ or 2). To the same accuracy, this parameter ϵ determines all the CP violating effects in $K_{1,2}^0$ decays.

Within the present experimental techniques, while it is possible to measure weak-interaction amplitudes to the accuracy of $|\epsilon| \sim 2 \times 10^{-3}$, it seems rather unlikely that any known weak-interaction amplitude can be determined to the accuracy of $F/G \sim 10^{-9}$. Thus, in the case that H_F is very much weaker than H_G , we may consider a simple *phenomenological model*, in which,

except for the mass operator of the K_1^0, K_2^0 system, all matrix elements are assumed to conserve CP and to be invariant under T . All CP -violating phenomena in this model are due to the CP noninvariance of the mass operator of the K_1^0 and K_2^0 states. The further assumption of CPT invariance requires all these CP violations to be described by one single phenomenological parameter ϵ , which has already been measured [except that Eq. (12) has two discrete solutions δ and $(\pi + \delta)$]. If the underlying CP -violating interaction H_F does satisfy the general conditions given by Eqs. (7) and (8), then to the assumed accuracy of neglecting F/G but keeping F/G^2 , all observable effects of such an extremely weak interaction are identical with this simple phenomenological model. If CPT invariance is violated, an additional real parameter η is needed to characterize the mass operator. To the same assumed accuracy, all observable CP -violating or CPT -violating effects of H_F contain only this single unknown parameter η .

In the above, we summarize the three different possibilities of the CP -noninvariant interaction H_F . The remaining part of the paper concerns, mainly, the $K_{1,2}^0$ system. In Sec. II, we review the general properties of the $K_{1,2}^0$ system, which are valid for all three classes of H_F , and without assuming the validity of the CPT theorem. Applications to the simple phenomenological model, in which only the mass operator violates CP invariance, are given in Sec. III, if CPT invariance holds, and in Sec. IV, if CPT invariance is also violated.

II. GENERAL DESCRIPTION OF K_1^0 AND K_2^0 STATES

In this section, we review the general description of the K_1^0 and K_2^0 system which is valid for *any* interaction Hamiltonian and *without* the assumption of CPT invariance.¹⁴ We consider a coherent mixture of K^0 and $\bar{K}^0$ whose amplitude at time t is described by the wave function

$$\psi(t) = a(t)|K^0\rangle + b(t)|\bar{K}^0\rangle \quad (13)$$

or simply as

$$\psi(t) = \begin{pmatrix} a(t) \\ b(t) \end{pmatrix}, \quad (14)$$

where t is measured in the rest system of the K meson, and the state $|\bar{K}^0\rangle$ is defined to be

$$|\bar{K}^0\rangle = CP|K^0\rangle. \quad (15)$$

The CP operator can only be determined by the strong and the electromagnetic interactions up to a phase factor

$$e^{iS\chi}, \quad (16)$$

where S is the strangeness operator and χ an arbitrary real number. However, since H_G is, by definition, the usual CP -conserving weak interaction, and since it must allow the $\Delta S = \pm 1$ transitions, this phase factor

¹¹ R. G. Sachs, Phys. Rev. Letters **13**, 286 (1964); N. Cabibbo, Phys. Letters **12**, 137 (1964); T. N. Truong, Phys. Rev. Letters **13**, 358 (1964); S. Glashow, *ibid.* **14**, 35 (1965).

¹² L. Wolfenstein, Phys. Rev. Letters **13**, 562 (1964). Our estimate of the order of magnitude of F is somewhat lower than in this reference. We obtain it from $(Fm_p^2/4\pi) \approx |\epsilon|(G_V m_p^2/4\pi)^2 (\frac{1}{2})^2$. A similar result is obtained from $(Fm_p^2/4\pi) \approx |\epsilon| \times [(m_1 - m_2)/m_K]$.

¹³ T. D. Lee, R. Oehme, and C. N. Yang, Phys. Rev. **105**, 1671 (1957).

¹⁴ Cf. R. Sachs, Ann. Phys. (N. Y.) **22**, 239 (1963). Sachs' notation is related to ours by $r = e^{i\phi}$, $s = \cot(\theta/2)$. He uses a different normalization condition for the eigenstates, however.

χ can be fixed by H_G . In the following, we will use this choice of phase factor and regard the CP operator as completely determined.

By using the Weisskopf-Wigner method¹⁵ of solving the time-dependent Schrödinger equation, the differential equation for $\psi(t)$ is given by

$$-d\psi/dt = (\Gamma + iM)\psi, \quad (17)$$

where Γ and M are two (2×2) Hermitian matrices,

$$\Gamma = \Gamma^\dagger \quad (18)$$

and

$$M = M^\dagger. \quad (19)$$

The matrix elements of Γ are related to that of the relevant perturbation Hamiltonian $\mathcal{H}$ by (neglecting $\mathcal{H}^3$)

$$\Gamma_{11} = 2\pi \sum_j \rho_j |\langle j | \mathcal{H} | K^0 \rangle|^2, \quad (20)$$

$$\Gamma_{22} = 2\pi \sum_j \rho_j |\langle j | \mathcal{H} | \bar{K}^0 \rangle|^2, \quad (21)$$

and

$$\Gamma_{21} = \Gamma_{12}^* = 2\pi \sum_j \rho_j \langle \bar{K}^0 | \mathcal{H} | j \rangle \langle j | \mathcal{H} | K^0 \rangle, \quad (22)$$

where the sum extends over all possible decay channels j , and ρ_j denotes the density of states per unit energy in the j th channel. The matrix elements of M are given by (neglecting $\mathcal{H}^3$)

$$M_{11} = m_{K^0} + \langle K^0 | \mathcal{H} | K^0 \rangle + \sum_\alpha \mathcal{P} [|\langle \alpha | \mathcal{H} | K^0 \rangle|^2 / (m_\alpha^0 - m_{K^0}^0)], \quad (23)$$

$$M_{22} = m_{\bar{K}^0} + \langle \bar{K}^0 | \mathcal{H} | \bar{K}^0 \rangle + \sum_\alpha \mathcal{P} [|\langle \alpha | \mathcal{H} | \bar{K}^0 \rangle|^2 / (m_\alpha^0 - m_{\bar{K}^0}^0)], \quad (24)$$

and

$$M_{21} = M_{12}^* = \langle \bar{K}^0 | \mathcal{H} | K^0 \rangle + \sum_\alpha \mathcal{P} [\langle \bar{K}^0 | \mathcal{H} | \alpha \rangle \langle \alpha | \mathcal{H} | K^0 \rangle / (m_\alpha^0 - m_{K^0}^0)], \quad (25)$$

where $\mathcal{P}$ stands for the principal value. In the case that H_F is much stronger than H_G (i.e., $F \approx 10^3 G$), the perturbation Hamiltonian $\mathcal{H}$ in all the above equations is given by

$$\mathcal{H} = H_G; \quad (26)$$

the sum α stands for the appropriate sums and integrations over all (unperturbed) eigenstates of the strong interaction, the electromagnetic interaction, and the new interaction H_F ; and $m_\alpha^0, m_{K^0}^0$ are the corresponding (unperturbed) eigenvalues of the states α and K^0 (or $\bar{K}^0$). In the other two cases ($F \lesssim G$ or $F \ll G$), the perturbation Hamiltonian is given by

$$\mathcal{H} = H_G + H_F, \quad (27)$$

the states α, K^0 (or $\bar{K}^0$) are eigenstates of the strong and electromagnetic interactions only, and $m_\alpha^0, m_{K^0}^0$ are the corresponding unperturbed eigenvalues.

We may express $(\Gamma + iM)$ in terms of the usual Pauli matrices $\sigma_1, \sigma_2, \sigma_3$:

$$(\Gamma + iM) = D + i(E_1 \sigma_1 + E_2 \sigma_2 + E_3 \sigma_3), \quad (28)$$

where

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

and D, E_1, E_2 , and E_3 are four complex numbers. It is convenient to introduce the complex variables E, θ , and ϕ :

$$E = (E_1^2 + E_2^2 + E_3^2)^{1/2}, \quad (29)$$

$$E_3 = E \cos \theta, \quad (30)$$

$$E_2 = E \sin \theta \sin \phi, \quad (31)$$

and

$$E_1 = E \sin \theta \cos \phi. \quad (32)$$

The eigenstates K_1^0 and K_2^0 , defined by

$$(\Gamma + iM) |K_j^0\rangle = [\frac{1}{2} \gamma_j + i m_j] |K_j^0\rangle, \quad (33)$$

where $j=1$ or 2 can be readily obtained; these solutions are identical in form to that of a spin- $\frac{1}{2}$ magnet in a magnetic field, and are given by

$$|K_1^0\rangle = N_1 \begin{pmatrix} \cos \frac{1}{2} \theta \\ e^{i\phi} \sin \frac{1}{2} \theta \end{pmatrix}, \quad (34)$$

and

$$|K_2^0\rangle = N_2 \begin{pmatrix} \sin \frac{1}{2} \theta \\ -e^{i\phi} \cos \frac{1}{2} \theta \end{pmatrix}, \quad (35)$$

where

$$N_1 = [|\cos \frac{1}{2} \theta|^2 + |e^{i\phi} \sin \frac{1}{2} \theta|^2]^{-1/2} \quad (36)$$

and

$$N_2 = [|\sin \frac{1}{2} \theta|^2 + |e^{i\phi} \cos \frac{1}{2} \theta|^2]^{-1/2}. \quad (37)$$

The observed mass m_j and width γ_j of K_j^0 are related to D and E by

$$D = \frac{1}{4}(\gamma_1 + \gamma_2) + \frac{1}{2}i(m_1 + m_2) \quad (38)$$

and

$$iE = \frac{1}{4}(\gamma_1 - \gamma_2) + \frac{1}{2}i(m_1 - m_2). \quad (39)$$

A useful parameter is

$$\begin{aligned} \alpha &= \langle K_1^0 | K_2^0 \rangle \\ &= N_1 N_2 [(\cos \frac{1}{2} \theta)^* (\sin \frac{1}{2} \theta) \\ &\quad - (e^{i\phi} \sin \frac{1}{2} \theta)^* (e^{i\phi} \cos \frac{1}{2} \theta)]. \end{aligned} \quad (40)$$

From the property that Γ is a positive-definite Hermitian matrix, it follows that (independent of the validity of the CPT theorem)

$$|\alpha|^2 \leq \gamma_1 \gamma_2 / [(m_1 - m_2)^2 + \frac{1}{4}(\gamma_1 + \gamma_2)^2], \quad (41)$$

which is $\sim 1.3 \times 10^{-3}$, if $|m_1 - m_2|$ is set to be about equal to γ_1 .

¹⁵ V. F. Weisskopf and E. P. Wigner, Z. Physik **63**, 54 (1930); **65**, 18 (1930).

The states K^0 and $\bar{K}^0$ can be expressed as linear combinations of the eigenstates K_1^0 and K_2^0 :

$$|K^0\rangle = N_1^{-1} \cos \frac{1}{2} \theta |K_1^0\rangle + N_2^{-1} \sin \frac{1}{2} \theta |K_2^0\rangle \quad (42)$$

and

$$e^{i\phi} |\bar{K}^0\rangle = N_1^{-1} \sin \frac{1}{2} \theta |K_1^0\rangle - N_2^{-1} \cos \frac{1}{2} \theta |K_2^0\rangle. \quad (43)$$

Let us consider a neutral K -meson beam which, at $t=0$, is in a pure K^0 state (or a pure $\bar{K}^0$ state). The fractional number of all K mesons decaying per unit time at time t is given by the familiar expression

$$\begin{aligned} \mathcal{N}_{\pm}(t) = & A_{\pm} \gamma_1 \exp(-\gamma_1 t) + B_{\pm} \gamma_2 \exp(-\gamma_2 t) \\ & + \{C_{\pm} [\frac{1}{2}(\gamma_1 + \gamma_2) - i(m_1 - m_2)] \\ & \times \exp[-\frac{1}{2}(\gamma_1 + \gamma_2)t + i(m_1 - m_2)t] + \text{c.c.}\}, \quad (44) \end{aligned}$$

where the plus (or minus) sign in the subscripts denotes the case that the neutral K meson is, at $t=0$, in a K^0 state (or a $\bar{K}^0$ state). The coefficients A_+ , A_- , B_+ , B_- are real and C_+ , C_- are complex. These eight real coefficients depend only on two complex numbers θ and ϕ :

$$\begin{aligned} A_+ &= N_1^{-2} |\cos \frac{1}{2} \theta|^2, & B_+ &= N_2^{-2} |\sin \frac{1}{2} \theta|^2, \\ A_- &= N_1^{-2} |e^{-i\phi} \sin \frac{1}{2} \theta|^2, & B_- &= N_2^{-2} |e^{-i\phi} \cos \frac{1}{2} \theta|^2, \\ C_+ &= (N_1 N_2)^{-1} (\cos \frac{1}{2} \theta)^* (\sin \frac{1}{2} \theta) \alpha, \end{aligned}$$

and

$$C_- = -(N_1 N_2)^{-1} (e^{-i\phi} \sin \frac{1}{2} \theta)^* (e^{-i\phi} \cos \frac{1}{2} \theta) \alpha. \quad (45)$$

The integrated value of $N_{\pm}(t)$ over all t must be 1; i.e.,

$$A_{\pm} + B_{\pm} + C_{\pm} + C_{\pm}^* = 1. \quad (46)$$

The constants $A_{\pm}$, $B_{\pm}$, and $C_{\pm}$ can be directly measured by studying the time development of such a K -meson beam. These measurements can be used to test the various symmetry or symmetry-violation properties of the K_1^0 , K_2^0 system.

Remarks

(1) In general, the (2×2) matrix $(\Gamma + iM)$ depends on eight real parameters, of which four parameters can be determined by the observed widths γ_1 , γ_2 and masses m_1 , m_2 of K_1^0 and K_2^0 . The remaining four parameters can be chosen to be the real and the imaginary parts of θ and ϕ :

$$\theta = \theta_R + i\theta_I \quad (47)$$

and

$$\phi = \phi_R + i\phi_I. \quad (48)$$

Three of these, θ_R , θ_I , and ϕ_I , can be determined by measuring $A_{\pm}$, $B_{\pm}$, $C_{\pm}$ and by using Eq. (45). The real part of ϕ depends on the relative phase between K^0 and $\bar{K}^0$, which, as remarked earlier, is determined only if we use the usual CP -conserving weak interaction H_G and the definition of $\bar{K}^0$ given by Eq. (15).¹⁶

¹⁶ For the simple phenomenological model, the effects of H_G and H_F are clearly separated and ϕ_R can be determined from experiments, as discussed in Sec. III and IV. For the other classes, this is less simple. Sometimes, it may be more convenient to define

(2) If CPT invariance holds, then¹³

$$\Gamma_{11} = \Gamma_{22} \quad (49)$$

and

$$M_{11} = M_{22}. \quad (50)$$

By using Eq. (28), it follows that

$$E_3 = 0. \quad (51)$$

Thus,

$$\theta = \frac{1}{2} \pi, \quad (52)$$

$$N_1 = N_2 = [\frac{1}{2}(1 + |e^{i\phi}|^2)]^{-1/2}, \quad (53)$$

and α is real. The validity of the CPT theorem can be tested by measuring the constants $A_{\pm}$, $B_{\pm}$, $C_{\pm}$ in Eq. (44) and observing that

$$A_+ = B_+ = A_- e^{-2\phi_I} = B_- e^{-2\phi_I} \quad (54)$$

and

$$C_+ = -C_- e^{-2\phi_I} \text{ is real.} \quad (55)$$

If CPT invariance holds, the eigenstates K_1^0 and K_2^0 can also be expressed in the forms given by Eqs. (9) and (10) which depend only on one complex parameter

$$\epsilon = (1 - e^{i\phi}) / (1 + e^{i\phi}). \quad (56)$$

(3) Independent of CPT invariance, if T invariance holds, then

$$\Gamma_{12}^* / \Gamma_{12} = M_{12}^* / M_{12}. \quad (57)$$

Thus, the imaginary part of ϕ must be zero; i.e.,

$$\phi_I = 0. \quad (58)$$

By using Eqs. (36), (37), and (45), we find

$$N_1 = N_2,$$

α is pure imaginary,

$$A_{\pm} = B_{\mp}, \quad (59)$$

and

$$C_+ = C_-^*. \quad (60)$$

Equations (59) and (60) can be used as tests of T invariance (without any assumption concerning the invariance of CPT , or CP , or C , or P).

(4) Independent of CPT invariance, if F/G is $\sim 10^{-9}$, then by using Eqs. (20)–(25) and Eq. (27), it is easy to see that to the accuracy of keeping F/G^2 , but neglecting F/G , all CP violations and CPT violations are due to the mass operator M . As already discussed in the previous section, to this assumed accuracy, such a theory is identical with the simple phenomenological model in which, except the mass operator M of the K_1^0 , K_2^0 system, all other transition matrix elements satisfy the consequences of CP invariance and T invariance. Thus

$$\Gamma_{12} = \Gamma_{21} \text{ is real and } \Gamma_{11} = \Gamma_{22},$$

the relative phase between K^0 and $\bar{K}^0$ such that the amplitudes for K^0 and $\bar{K}^0$ going to the $I=0$ two-pion state are relatively real. [A method of determining ϕ_R by using this convention has been discussed by T. T. Wu and C. N. Yang, Phys. Rev. Letters 13, 380 (1964).] The two definitions of the relative phase between K^0 and $\bar{K}^0$ are the same for the simple phenomenological model.

which correspond to

$$E_2, \text{ real} \quad (61)$$

and

$$E_3, \text{ real.} \quad (62)$$

The further consequences of this simple phenomenological model will be studied in the next two sections.

III. CONSEQUENCES OF THE SIMPLE PHENOMENOLOGICAL MODEL AND CPT INVARIANCE

In the case of the simple phenomenological model, if CPT invariance holds, the eigenstates K_1^0 and K_2^0 are given by Eqs. (9) and (10). By using Eqs. (31), (39), (52), and (56), the parameter ϵ is given by

$$2\epsilon/(1-\epsilon^2) = (E_2/|E|)e^{i\beta}, \quad (63)$$

where the phase β is defined by

$$iE = |E|e^{-i\beta} = \frac{1}{4}(\gamma_1 - \gamma_2) + \frac{1}{2}i(m_1 - m_2). \quad (64)$$

According to Eq. (61), E_2 is real. Thus, the phase δ and the magnitude $|\epsilon|$ of the parameter ϵ satisfy

$$\tan\delta = \frac{\sin\beta - |\epsilon|^2 \sin(2\delta + \beta)}{\cos\beta - |\epsilon|^2 \cos(2\delta + \beta)}. \quad (65)$$

In this model, all the transition matrix elements for the decays of K^0 and $\bar{K}^0$ into any continuum states, such as 2π , 3π , etc., satisfy the consequences of CP invariance. The magnitude $|\epsilon|$ is, therefore, given by Eq. (1). Consequently, both $|\epsilon|$ and δ have been experimentally measured. If we neglect $|\epsilon|^2$ in Eq. (65), the phase δ is given by Eq. (12):

$$\tan\delta \cong \tan\beta = -2(m_1 - m_2)/(\gamma_1 - \gamma_2),$$

which has always two solutions, $\delta = \beta$ and $\delta = \pi + \beta$. There are, at present, some discrepancies between the existing measurements¹⁷ of the mass difference $(m_1 - m_2)$; therefore, the exact value of β , or δ , is not yet definite.

The following is a list of some CP -violating effects in this simple model under the assumption of CPT invariance¹⁸:

¹⁷ The experimental value of the mass difference between K_1^0 and K_2^0 varies from $0.50 \gamma_1$ to $1.9 \gamma_1$. See B. Aubert *et al.*, Phys. Letters **10**, 215 (1964); U. Camerini *et al.*, Phys. Rev. **128**, 362 (1962); J. H. Christenson *et al.*, Conference on Fundamental Aspects of Weak Interactions, BNL, New York (1963); J. H. Christenson, Tech. Rept. 34, Princeton University (1964); V. L. Fitch, Nuovo Cimento **22**, 1160 (1961); M. L. Good, *et al.*, Phys. Rev. **124**, 1223 (1961). In Ref. 12, the lowest of these values was used to obtain a numerical value for the ratio in Eq. (70).

¹⁸ While the equalities given by Eqs. (66)–(72) can be used to test the validity of the simple model (i.e., $F \approx 10^{-9}G$), it is important to notice that some of these results are also valid in other cases. For example, if we consider the case $F \approx 10^{-2}G$ and assume that H_F and H_G both satisfy the $|\Delta I| = \frac{1}{2}$ rule for the strangeness-nonconserving amplitudes. The $|\Delta I| = \frac{3}{2}$ amplitude is, then, due to the combined effect of electromagnetic interaction and either H_G or H_F . Thus, neglecting $O(\epsilon\alpha)$, the equality

$$\frac{\text{Rate}(K_2^0 \rightarrow \pi^+ + \pi^-)}{\text{Rate}(K_1^0 \rightarrow \pi^+ + \pi^-)} = \frac{\text{Rate}(K_2^0 \rightarrow 2\pi^0)}{\text{Rate}(K_1^0 \rightarrow 2\pi^0)}$$

remains valid. Assuming CPT invariance and the $\Delta Q = \Delta S$ rule

$$\frac{\text{Rate}(K_2^0 \rightarrow 2\pi^0)}{\text{Rate}(K_1^0 \rightarrow 2\pi^0)} = \frac{\text{Rate}(K_2^0 \rightarrow \pi^+ + \pi^-)}{\text{Rate}(K_1^0 \rightarrow \pi^+ + \pi^-)} = |\epsilon|^2, \quad (66)$$

$$\frac{\text{Rate}(K_1^0 \rightarrow 3\pi^0)}{\text{Rate}(K_2^0 \rightarrow 3\pi^0)} = |\epsilon|^2. \quad (67)$$

The decay of $K_1^0 \rightarrow \pi^+ + \pi^- + \pi^0$ can occur with the final 3π system in either a $CP = -1$ state (which shows CP violation) or a $CP = +1$ state (which is compatible with CP conservation). The amplitude of $K_1^0 \rightarrow \pi^+ + \pi^- + \pi^0$ in a final $CP = -1$ state is ϵ times the corresponding amplitude of K_2^0 decay. The amplitude of $K_1^0 \rightarrow \pi^+ + \pi^- + \pi^0$ in a final $CP = +1$ state does not violate CP conservation, but is reduced from the corresponding $CP = -1$ amplitude of K_2^0 decay only by the centrifugal barrier effects.

(3) For leptonic decays, we assume that the $\Delta Q = \Delta S$ rule holds for the transition matrix elements between the K^0 , $\bar{K}^0$ states and any leptonic states [i.e., the $\Delta Q = \Delta S$ rule holds for H_G].

$$\begin{aligned} \text{Rate}(K_1^0 \rightarrow \pi^+ + l^- + \bar{\nu}_l) \\ = \text{Rate}(K_2^0 \rightarrow \pi^+ + l^- + \bar{\nu}_l), \end{aligned} \quad (68)$$

$$\begin{aligned} \text{Rate}(K_1^0 \rightarrow \pi^- + l^+ + \nu_l) \\ = \text{Rate}(K_2^0 \rightarrow \pi^- + l^+ + \nu_l), \end{aligned} \quad (69)$$

and

$$\frac{\text{Rate}(K_1^0 \rightarrow \pi^- + l^+ + \nu_l)}{\text{Rate}(K_1^0 \rightarrow \pi^+ + l^- + \bar{\nu}_l)} = \left| \frac{1 + \epsilon}{1 - \epsilon} \right|^2 \cong 1 + 4|\epsilon| \cos\delta, \quad (70)$$

where $l = e$ or μ . Identical relations also hold for K_{14} decays, provided we replace π^+ and π^- in the above formulas by $(\pi^+ + \pi^0)$ and $(\pi^- + \pi^0)$, respectively.

(4) By using Eqs. (52) and (56), the coefficients $A_{\pm}$, $B_{\pm}$, and $C_{\pm}$ in Eq. (44) are given by

$$A_{\pm} = B_{\pm} = (1 + |\epsilon|^2)/2 |1 \pm \epsilon|^2 \cong \frac{1}{2} [1 \mp 2|\epsilon| \cos\delta] \quad (71)$$

and

$$C_{\pm} = \pm(\epsilon + \epsilon^*)/2 |1 \pm \epsilon|^2 \cong \pm |\epsilon| \cos\delta. \quad (72)$$

IV. CONSEQUENCES OF THE SIMPLE MODEL WITHOUT CPT INVARIANCE

In this section, we list the CP -violating and CPT -violating phenomena for the same simple phenomenological model, but without assuming CPT invariance. According to Eqs. (61) and (62), these violations depend only on two real parameters E_2 and E_3 . The observed rate of $K_2^0 \rightarrow \pi^+ + \pi^-$ gives one relations between these two parameters. The whole problem, therefore, con-

one finds, in addition, for this case that $|\epsilon|$ and δ are given within 10% by Eqs. (1) and (12). Thus, except for Eq. (67), Eqs. (66) and (68)–(72) also hold to a fair degree of approximation for the case $F \approx 10^{-2}G$, provided that H_F and H_G both satisfy the $|\Delta I| = \frac{1}{2}$ rule. [The results for δ can be most conveniently obtained by using Eqs. (20) and (23) of Wu and Yang (Ref. 16).]

tains only one real parameter which has not yet been measured.

It is convenient to represent the K_1^0 and K_2^0 states by

$$|K_1^0\rangle = [2(1 + |\epsilon_1|^2)]^{-1/2} \begin{pmatrix} 1 + \epsilon_1 \\ 1 - \epsilon_1 \end{pmatrix} \quad (73)$$

and

$$|K_2^0\rangle = [2(1 + |\epsilon_2|^2)]^{-1/2} \begin{pmatrix} 1 + \epsilon_2 \\ -(1 - \epsilon_2) \end{pmatrix}, \quad (74)$$

where ϵ_1 and ϵ_2 are related to θ and ϕ by

$$(1 - \epsilon_1)/(1 + \epsilon_1) = e^{i\phi} \tan \frac{1}{2}\theta \quad (75)$$

and

$$(1 - \epsilon_2)/(1 + \epsilon_2) = e^{i\phi} \cot \frac{1}{2}\theta, \quad (76)$$

which can, in turn, be expressed in terms of the real parameters E_2 and E_3 .

We note that

$$\begin{aligned} \frac{\text{Rate}(K_2^0 \rightarrow \pi^+ + \pi^-)}{\text{Rate}(K_1^0 \rightarrow \pi^+ + \pi^-)} &= \frac{\text{Rate}(K_2^0 \rightarrow 2\pi^0)}{\text{Rate}(K_1^0 \rightarrow 2\pi^0)} \\ &= \frac{1 + |\epsilon_1|^2}{1 + |\epsilon_2|^2} |\epsilon_2|^2 \end{aligned} \quad (77)$$

and

$$\frac{\text{Rate}(K_1^0 \rightarrow 3\pi^0)}{\text{Rate}(K_2^0 \rightarrow 3\pi^0)} = \frac{1 + |\epsilon_2|^2}{1 + |\epsilon_1|^2} |\epsilon_1|^2. \quad (78)$$

From Eq. (77) and the experimental result¹ of Christenson *et al.*, it follows that

$$\frac{|\epsilon_2|^2}{1 + |\epsilon_2|^2} \leq |\epsilon|^2 \cong 4.8 \times 10^{-6},$$

or

$$|\epsilon_2| \leq |\epsilon| \times [1 - |\epsilon|^2]^{-1/2} \cong 2.2 \times 10^{-3}, \quad (79)$$

which, together with the inequality (41), implies that

$$|\epsilon_1| \leq [\sim 3.6 \times 10^{-2}]. \quad (80)$$

Thus, $|\epsilon_1|$ and $|\epsilon_2|$ must both be small numbers, and $|\epsilon_1|^2$, $|\epsilon_2|^2$ and $|\epsilon_1 \epsilon_2|$ may be neglected as compared to 1. In this *approximation*, all formulas become considerably simpler. The results are given below:

(1) Let

$$\epsilon_j = |\epsilon_j| \exp(i\delta_j); \quad (81)$$

then

$$|\epsilon_1| = |\epsilon_2| = |\epsilon| \quad (82)$$

and

$$\frac{1}{2}(\delta_1 + \delta_2) = \delta, \quad (83)$$

where

$$\delta = \beta \quad \text{or} \quad (\pi + \beta). \quad (84)$$

The constants $|\epsilon|$ and β have both been measured, and are given, respectively, by Eqs. (1) and (64).

The entire problem contains only one unknown real parameter which may be chosen to be

$$\frac{1}{2}(\delta_1 - \delta_2) = \eta. \quad (85)$$

(2) The coefficients $A_{\pm}$, $B_{\pm}$, and $C_{\pm}$ in Eq. (44) are given by

$$A_{\pm} = \frac{1}{2}[1 \mp 2|\epsilon| \cos(\delta - \eta)], \quad (86)$$

$$B_{\pm} = \frac{1}{2}[1 \mp 2|\epsilon| \cos(\delta + \eta)], \quad (87)$$

and

$$C_{\pm} = \pm |\epsilon| (\cos \delta) e^{-i\eta}. \quad (88)$$

(3) For the K^0 decay into pions, Eqs. (66) and (67) remain valid. For the K^0 decay into leptons, if $\Delta Q = \Delta S$ rule holds, we have

$$\begin{aligned} \text{Rate}(K_j^0 \rightarrow \pi^- + l^+ + \nu_l) / \text{Rate}(K_j^0 \rightarrow \pi^+ + l^- + \bar{\nu}_l) \\ = 1 + 4|\epsilon| \cos \delta_j, \end{aligned} \quad (89)$$

where $l = e$ or μ and $j = 1$ or 2 . The same formula holds for K_{14} decays, provided $\pi^{\mp}$ is replaced by $(\pi^{\mp} + \pi^0)$.

(4) If T invariance holds (but not CPT invariance), then by using Eqs. (58), (75), and (76), and since $\cos \delta \neq 0$, we find

$$\eta = \frac{1}{2}\pi. \quad (90)$$

(5) It has been suggested² that the observation of $K_2^0 \rightarrow \pi^+ + \pi^-$ can be compatible with CP invariance, provided there exists an energy difference ΔE between the K^0 and the $\bar{K}^0$ states due to some cosmological background. The apparent CP noninvariance is attributed not to the basic violation of the interaction, but is regarded as the result of the initial (or boundary) condition of our universe which gives rise to this background field. As remarked in the paper by Bernstein *et al.*,² the amount of this energy difference ΔE is proportional to E^{2J} , where $J = \text{spin of the cosmological background field (or, rank of the corresponding field tensor)}$ and E is the energy of the K meson measured in the rest system of the cosmological background field.

From a phenomenological point of view, except for this energy dependence of ΔE , the description of the K_1^0 and K_2^0 system in such a cosmological background field is the same as the simple model (without CPT invariance) considered in this paper, provided

$$\epsilon_2 = -\epsilon_1; \quad (91)$$

i.e.,

$$\eta = \frac{1}{2}\pi,$$

which is identical with Eq. (90). If $J = 0$, then there is no difference between the cosmological possibility with CP invariance and the case of an extremely weak CP -noninvariant, CPT -noninvariant, but T -invariant H_F with $F \sim 10^{-9}G$, unless either the measurement of the appropriate transition amplitudes can be improved to include terms of the order of 10^{-9} , or the observation of a real emission of the cosmological field quantum can become feasible.

ACKNOWLEDGMENT

One of us (L.W.) wishes to thank Professor L. Van Hove and Professor V. F. Weisskopf for the hospitality of CERN.