REHIVES

FIAT THEORY OF GRAVITATION

A. Capella
CERN - Geneva

A B S T R A C T

We construct a Lorentz covariant theory of gravitation with linear field equations which accounts for all the observational facts, i.e., the three tests of general relativity, with only one adjustable parameter apart from the gravitational coupling constant. The equations of motion of the theory arise both from a variational principle and from the conservation law of the total energy-momentum tensor of the system. The conservation law of mass is also satisfied.

*) On leave of absence from the Centre National de la Recherche Scientifique, France.

1. INTRODUCTION

For many years attempts to construct Lorentz covariant linear theories of gravitation have been made ¹⁾. The first of these was Birkhoff's theory in 1944 ²⁾. As is well known the equations of motion in this theory are postulated "ad hoc" and cannot be obtained either from a variational principle or from the conservation law of the total energy-momentum tensor of the system. The conservation law of the mass is also lacking in this theory.

In most of the subsequent theories, although the conservation law of the total energy-momentum tensor is satisfied, the conservation law of the mass is still lacking. The reasons for this are different in each theory. In the theories ^{3),4)}, for instance, the equations of motion are postulated to be given by the conservation law of the total energy-momentum tensor. However, this conservation law does not agree with the Euler equations arising from the Lagrangian of the system. This can happen because the total energy-momentum tensor in the conservation law is not taken to be the "metric" energy-momentum tensor of the system ⁵⁾. In the theory ⁶⁾, on the contrary, the equations of motion are obtained from a variational principle. However, the total Lagrangian depends not only on the velocity variables and the field variables but also on other variables such as the momenta. The latter are not trivially related to the former because of the interaction term in the Lagrangian. For this reason the conservation law of the mass is also lacking in this theory when a continuous scheme for matter is used.

Some time ago two theories ^{7),8)} were proposed in which both the conservation law of the energy and the conservation law of mass are satisfied and which account for the three tests of general relativity using only one adjustable parameter apart from the gravitational coupling constant. As the author's theory ⁸⁾ has been published in a rather compact form, the details, including some previously unpublished results, are given in Sections 2 and 3. Section 3 also includes the calculation of the predictions of the theory for the fourth test of general relativity recently suggested by Shapiro ⁹⁾. In Section 4 the comparison between the two theories ^{7),8)} is made.

2.

2. FIELD EQUATIONS AND EQUATIONS OF MOTION

Let us consider the following Lagrangian density for a system of n particles of masses m_N ($N = 1, \dots, n$) interacting with the gravitational field described by the symmetric potentials $h_{\alpha\beta}$ ¹⁰⁾

$$\mathcal{L}^{(p)}(x) + \mathcal{L}^{(i)}(x) = \sum_N m_N \int ds_N \delta(x - z_N) \left[- (g_{\alpha\beta} \dot{z}_N^\alpha \dot{z}_N^\beta)^{1/2} + \right. \\ \left. + \chi h_{\alpha\beta}(x) \dot{z}_N^\alpha \dot{z}_N^\beta (g_{\rho\sigma} \dot{z}_N^\rho \dot{z}_N^\sigma)^{-1/2} \right] \quad (2.1)$$

where $z_N^\alpha(s_N)$ is the trajectory of the N^{th} particle, $\dot{z}_N^\alpha = dz_N^\alpha / ds_N$, and χ the gravitational coupling constant.

We add to (2.1) the following Lagrangian density for the free gravitational field

$$\mathcal{L}^{(f)}(x) = \frac{1}{2} \left(\partial_\alpha h_{\beta\gamma} \partial^\alpha h^{\beta\gamma} + a \partial_\alpha h \partial^\alpha h \right) \sqrt{-g} \quad (2.2) \quad (11)$$

where a is a real constant.

The Lagrangians (2.1) and (2.2) yield the field equations

$$\square (h_{\alpha\beta} + a g_{\alpha\beta} h) = \chi T_{\alpha\beta}^{(p)} \quad (2.3)$$

where $\square \equiv \partial_\rho \partial^\rho$, $h = g^{\alpha\beta} h_{\alpha\beta}$ and

$$\sqrt{-g} T^{(p)\alpha\beta} = \sum_N m_N \int ds_N \delta(x - z_N) \dot{z}_N^\alpha \dot{z}_N^\beta. \quad (2.4)$$

The equations of motion of a particle are obtained from the action principle

$$\delta W = 0, \quad W = \int d^4x [\mathcal{L}^{(p)}(x) + \mathcal{L}^{(i)}(x)] = \int ds \Lambda(s),$$

where, according to (2.1), we have

$$\Lambda(s) = -m (g_{\alpha\beta} \dot{z}^\alpha \dot{z}^\beta)^{1/2} + m \chi h_{\alpha\beta}(z) \dot{z}^\alpha \dot{z}^\beta (g_{\rho\sigma} \dot{z}^\rho \dot{z}^\sigma)^{-1/2}. \quad (2.5)$$

The Euler-Lagrange equations arising from (2.5) are the following

$$\frac{d}{ds} (\dot{z}_\lambda - 2\chi h_{\alpha\lambda} \dot{z}^\alpha + \chi h_{\alpha\beta} \dot{z}^\alpha \dot{z}^\beta \dot{z}_\lambda) = -\chi \dot{z}^\alpha \dot{z}^\beta \partial_\lambda h_{\alpha\beta}. \quad (2.6)$$

It is well known ⁵⁾ that, using the field equations (2.3), the equations of motion (2.6) are equivalent to the conservation law of the total energy-momentum tensor of the system

$$\partial_\alpha T^{\alpha\beta} \equiv \partial_\alpha [T_{metric}^{(f)\alpha\beta} + T^{(p)\alpha\beta} + T^{(c)\alpha\beta}] = 0, \quad (2.7)$$

where

$$\begin{aligned} \sqrt{-g} T^{\alpha\beta} &= \frac{\delta \mathcal{L}^{(f)}}{\delta g_{\alpha\beta}} - \frac{\delta (\mathcal{L}^{(p)} + \mathcal{L}^{(c)})}{\delta g_{\alpha\beta}} \equiv \\ &\equiv \sqrt{-g} [T_{metric}^{(f)\alpha\beta} + T^{(p)\alpha\beta} + T^{(c)\alpha\beta}], \end{aligned} \quad (2.8)$$

the variational derivatives being given by

$$\frac{\delta \mathcal{L}}{\delta g^{\alpha\beta}} = 2 \left(\frac{\partial \mathcal{L}}{\partial g^{\alpha\beta}} - \partial_\rho \frac{\partial \mathcal{L}}{\partial \partial_\rho g^{\alpha\beta}} \right) \quad 12) \quad (2.9)$$

The energy-momentum tensor of the system consists of three terms as shown by (2.8). The first term is called the "metric" energy-momentum tensor of the field. According to (2.2), (2.8) and (2.9) we have explicitly

4.

$$\begin{aligned}
T_{metric}^{(f)\alpha\beta} = & \partial^\alpha h_{\rho\sigma} \partial^\beta h^{\rho\sigma} + a \partial^\alpha h \partial^\beta h - \frac{1}{2} g^{\alpha\beta} (\partial_\rho h_{\sigma\tau} \partial^\rho h^{\sigma\tau} + a \partial_\rho h \partial^\rho h) + \\
& h^{\rho\nu} \partial_\rho \partial^{(\alpha} h^{\beta)\nu} - h^{\nu(\beta} \partial^\alpha \partial_\rho h^{\rho\nu)} + \partial^{(\alpha} h^{\beta)\nu} \partial_\rho h^{\rho\nu} - \partial_\rho h^{\nu(\beta} \partial^\alpha h^{\rho\nu)} - \\
& - \partial_\rho \partial^\rho (h^{\alpha\nu} + a g^{\alpha\nu} h) h^\beta_\nu - \partial_\rho \partial^\rho (h^{\beta\nu} + a g^{\beta\nu} h) h^\alpha_\nu. \quad (2.10)
\end{aligned}$$

Using (2.3) the divergence of $T_{metric}^{(f)}$ becomes

$$\partial_\alpha T_{metric}^{(f)\alpha\beta} = T^{(\rho)\rho\sigma} \partial_\beta h_{\rho\sigma} - 2 T^{(\rho)\rho\sigma} \partial_\rho h_{\beta\sigma} - 2 \partial_\rho T^{(\rho)\rho\sigma} h_{\beta\sigma}. \quad (2.11)$$

With the aid of this relation it is easy to prove that (2.6) and (2.7) are equivalent.

The second term in (2.8) is the energy-momentum tensor of the particles. Its explicit expression is given by Eq. (2.4) (no sum now because we are considering only one particle).

The third term is the interaction energy-momentum tensor, given by

$$\sqrt{-g} T^{(i)\alpha\beta} = m \int ds \delta(x-z) \dot{z}^\rho \dot{z}^\sigma h_{\rho\sigma}(x) \dot{z}^\alpha \dot{z}^\beta. \quad (2.12)$$

As shown by (2.10), $T_{metric}^{(f)}$ contains terms which can be considered to belong either to the energy-momentum tensor of the field or to the interaction part of the total energy-momentum tensor. Let us define

$$\begin{aligned}
T^{(f)\alpha\beta} &= T_{metric}^{(f)\alpha\beta} + \partial_\rho \partial^\rho (h^{\alpha\nu} + a g^{\alpha\nu} h) h_\nu^\beta + \partial_\rho \partial^\rho (h^{\beta\nu} + a g^{\beta\nu} h) h_\nu^\alpha = \\
&= T_{metric}^{(f)\alpha\beta} + \frac{m}{\sqrt{-g}} \int ds \delta(x-z) \dot{z}^\rho \dot{z}^{(\alpha} h^{\beta)}_\rho \quad (2.13)
\end{aligned}$$

The canonic energy-momentum tensor arising from (2.2), when symmetrized according to the rule suggested by Rosenfeld and Belinfante coincides with the tensor (2.13).

Using (2.13) we can rewrite Eq. (2.7) in the following way

$$\partial_\alpha T^{\alpha\beta} \equiv \partial_\alpha [T^{(f)\alpha\beta} + T^{(p)\alpha\beta} + T'^{(i)\alpha\beta}] = 0, \quad (2.7')$$

where

$$\sqrt{-g} T'^{(i)}_{\alpha\beta} = \sqrt{-g} T^{(i)}_{\alpha\beta} - m \int ds \delta(x-z) \dot{z}^\rho \dot{z}_{(\alpha} h_{\beta)\rho}(x). \quad (2.14)$$

As will be shown in Section 3, the tensor $T^{(f)}$ is more suitable than $T_{metric}^{(f)}$ to describe the energy and momentum of the field.

Until now we have used a pointlike scheme for matter. If we use a continuous scheme [with $T_{\alpha\beta}^{(p)} = \mu u_\alpha u_\beta$ instead of (2.4)], the resulting equations of motion are equivalent to (2.6). Indeed, Eqs. (2.7') are replaced by

$$\partial_\alpha [T^{(f)\alpha\beta} + \mu u^\alpha u^\beta + \mu u^\rho u^\sigma h_{\rho\sigma} u^\alpha u^\beta - \mu u^\rho u^{(\alpha} h^{\beta)}_\rho] = 0, \quad (2.7'')$$

where $u^\alpha = dx^\alpha/ds$. From (2.7'') we deduce both the conservation law of mass $\partial_\alpha (\mu u^\alpha) = 0$, and the equations of motion (2.6) (as expected since (2.1) is an homogeneous polynomial of degree one in the velocity variables depending only on these variables and on the field variables).

6.

3. CALCULATION OF THE OBSERVATIONAL EFFECTS

- a) Let us first calculate the $h_{\alpha\beta}$ originating from a mass point with mass M at the origin of a co-ordinate system. From (2.3) we obtain

$$h'_{00} = \frac{\chi}{4\pi} \frac{M}{r}, \quad h'_{\alpha i} = 0 \quad (3.1)$$

where

$$h'_{\alpha\beta} = h_{\alpha\beta} + a g_{\alpha\beta} h. \quad (3.2)$$

From (3.1) and (3.2) we get

$$\begin{aligned} h_{00} &= \frac{\chi}{4\pi} \frac{1+3a}{1+4a} \frac{M}{r} \equiv \frac{b}{\chi} G \frac{M}{r}, \\ h_{ii} &= \frac{\chi}{4\pi} \frac{a}{1+4a} \frac{M}{r} \equiv \frac{b'}{\chi} G \frac{M}{r} \end{aligned} \quad (3.3)$$

G being Newton's constant of gravitation. All the non-diagonal components of $h_{\alpha\beta}$ are zero.

Substituting (3.3) in (2.6) and neglecting terms of order higher than v^2 (because their contribution to the observational effects is negligible) we find

$$\begin{aligned} \frac{d^2 x^i}{ds^2} &= -G \frac{M}{r^3} x^i [b + (b+b')v^2] + G \frac{M}{r^2} \frac{dx^i}{ds} \frac{dr}{ds} (b+2b') + \\ &+ G^2 \frac{M^2}{r^4} x^i b (b+2b'). \end{aligned} \quad (3.4)$$

Agreement with Newton's theory is obtained by putting $b = 1$. Equations of motion of type (3.4) have been studied by several authors. As is well known ¹⁴⁾ the closed trajectories are ellipses whose perihelion advances at each revolution an angle $\delta\omega$ given by

$$\delta w = (3+4b') \frac{\pi G M}{A(1-e^2)} \quad (3.5)$$

where A is the major semi-axis of the ellipse and e its excentricity. Formula (3.5) agrees with Einstein's formula for the perihelion shift when $b' = 3/4$. We then obtain from (3.3)

$$\chi = \sqrt{7\pi G}^{15), \quad \alpha = -\frac{3}{5}$$

- b) With these values of the parameters we shall now calculate the bending of light rays at the edge of the sun and the red shift. We are going to use the same approach as Moshinsky¹⁶⁾.

We start from the Lagrangian for the gravitational field interacting with the electromagnetic field. This Lagrangian is the following

$$L = \frac{1}{2} (\partial_\alpha h_{\beta\gamma} \partial^\alpha h^{\beta\gamma} + \alpha \partial_\alpha h \partial^\alpha h) + \chi h_{\alpha\beta} \tau^{\alpha\beta} + \frac{1}{4} F_{\alpha\beta} F^{\alpha\beta}, \quad (3.6)$$

where $F_{\alpha\beta} = \partial_\alpha \varphi_\beta - \partial_\beta \varphi_\alpha$ is the electromagnetic field tensor and $\tau_{\alpha\beta}$ the Maxwell tensor. Indeed, the Euler equations obtained by varying in (3.6) the $h_{\alpha\beta}$, are the Eqs. (2.3) except that $T_{\alpha\beta}^{(p)}$ is replaced by $\tau_{\alpha\beta}$.

Varying in (3.6) the electromagnetic potentials, it follows that

$$\partial_\alpha [F^{\alpha\beta} + 2\chi h^{\alpha\rho} F_\rho{}^\beta + 2\chi h^{\beta\rho} F^\alpha{}_\rho - \chi h F^{\alpha\beta}] = 0. \quad (3.7)$$

Using the values of $h_{\alpha\beta}$ given by (3.3), Eqs. (3.7) coincide with Maxwell's equation, except that the electromagnetic constants ϵ and μ are no longer equal to one but are the following functions of the gravitational potentials

$$\epsilon = 1 + \chi(h_{00} + h_{11}) = 1 + (1 + b') \frac{GM}{r},$$

$$\mu = \frac{1}{1 - \chi(h_{00} + h_{11})} = \frac{1}{1 - (1 + b') \frac{GM}{r}}.$$

This corresponds to a medium with a refractive index

$$n = \sqrt{\epsilon\mu} \approx 1 + (1 + b') \frac{M}{r} G. \quad (3.8)$$

The path of the light rays is then obtained from Fermat's principle. The resulting formula for the deflection angle is

$$\alpha = \frac{7}{2} \frac{GM}{R} \quad (3.9)$$

M being the mass of the sun and R the distance from the centre of the sun to the unperturbed light path. Einstein's formula coincides with (3.9) except that a factor four replaces our $7/2$ factor. Consequently the predicted value is 12% smaller than Einstein's value. Such a value is not in disagreement with the observed value ¹⁶⁾.

For the red shift of the energy levels of a hydrogen atom, we start from the Lagrangian obtained by adding to the Dirac Lagrangian for the electron free field, the interaction Lagrangian $L^{(i)} = h_{\alpha\beta} t^{\alpha\beta}$. Here $t^{\alpha\beta}$ is the energy-momentum tensor of the Dirac field. We write down the corresponding Euler equations and with these equations we calculate the energy levels of a hydrogen atom. (Of course, the electromagnetic potential $\phi = e/r$ must be changed to $\phi' = e/\epsilon r$.) The result is the following

$$E'_n = E_n (1 - 2\chi h_{00}) = E_n \left(1 - \frac{GM}{r}\right), \quad (3.10)$$

in perfect agreement with Einstein's formula.

- c) Let us now calculate the prediction of the theory for the fourth test of general relativity recently suggested by Shapiro⁹⁾. This test involves measuring the time delays between transmission of radar pulses to one of the inner planets and detection of the echoes. Using Eq. (3.8) we obtain for this time delay the following expression

$$t = 2 \int_0^{x_e + x_p} \frac{dx}{c} = 2 \int_0^{x_e + x_p} dx \left[1 + (1 + b') \frac{GM}{rc} \right] =$$

$$2(x_e + x_p) + 2(1 + b') GM \ln \left[\frac{x_p + (x_p^2 + d^2)^{1/2}}{-x_e + (x_e^2 + d^2)^{1/2}} \right] \equiv t_0 + \Delta t, \quad (3.11)$$

where d is the distance from the centre of the sun to the radar path and x_e , x_p are the distances along the path from the earth antenna to the point of closest approach to the sun and from this point to the planet, respectively. The measurable quantity is the difference $\Delta \equiv \Delta t_s - \Delta t_i$, where Δt_s is the value of Δt when the sun is between the earth and the planet close to the radar path, and Δt_i the corresponding value when the planet is between the earth and the sun.

For Mercury, with d equal to two times the radius of the sun, we obtain from (3.11)

$$\Delta = \Delta_{G.R.} + \frac{4}{100} \Delta_{G.R.}, \quad (3.12)$$

where $\Delta_{G.R.}$ is the value obtained by Shapiro from general relativity. The expected experimental error is 5 to 10%.

Let us compare (3.11) with the formula obtained by Shapiro from Einstein's theory⁹⁾

$$\Delta t = 4GM \left\{ \ln \left[\frac{x_p + (x_p^2 + d^2)^{1/2}}{-x_e + (x_e^2 + d^2)^{1/2}} \right] - \frac{1}{2} \left[\frac{x_p}{(x_p^2 + d^2)^{1/2}} + \frac{2x_e + x_p}{(x_e^2 + d^2)^{1/2}} \right] \right\}. \quad (3.13)$$

We see that the second term in (3.13) is missing in (3.11). This term is quite important. For Mercury, with $d = 2 R_0$, it is 17% of the first term. However, the numerical factor in front of (3.11) is smaller than the corresponding one in (3.13). This partially compensates the contribution of the missing term, yielding finally the result (3.12).

Comparison of (3.11) and (3.13) shows, however, that a flat theory of gravitation, with a refractive index given by (3.8), cannot yield the same values as general relativity for both the bending of light rays and the fourth test. In Birkhoff's theory, for instance, b' is equal to one ¹⁾ and (3.11) yields

$$\Delta_{\text{BIRKHOFF}} = \Delta_{\text{G.R.}} + \frac{18}{100} \Delta_{\text{G.R.}}$$

A precise measurement of both the bending of light rays and the fourth test could then support general relativity over flat theories or vice versa.

- d) Let us calculate finally the total energy of the system in the quasi-static approximation. Using the static solutions (3.3) extended to n particles and Eqs. (2.10) and (2.13) we find

$$P_0^{(f)} = \int T_{00}^{(f)} d^3X = \frac{1}{2} \int d^3X \left[(\partial_i h_{00})^2 + 3(\partial_i h_{11})^2 + a(\partial_i h)^2 \right] = \frac{G}{2} \sum_{M,N=1}^n \frac{m_M m_N}{r_{MN}} \quad (3.14)$$

$$P_0^{(f)} = \int T_{\text{metric}}^{(f)00} d^3X = \int T^{(f)00} d^3X - 2 \sum_N m_N \int d^3x \delta(x-z_N) (\dot{z}_N^0)^2 h^{00}(x) d^3x \simeq -\frac{3}{4} G \sum_{M,N=1}^n \frac{m_M m_N}{r_{MN}} \quad (3.15)$$

These two expressions (3.14) and (3.15) show that $T^{(f)}$ (and not $T_{\text{metric}}^{(f)}$) has to be used as the energy-momentum tensor of the gravitational field.

For the total energy of the system we obtain from (2.7), (2.12), (2.4) and (3.15) the following result

$$P^0 = \int T^{00} d^3X = \int \left[T_{\text{metric}}^{(f)00} + T^{(r)00} + T^{(c)00} \right] d^3X \approx \sum_{N=1}^n m_N \left(1 + \frac{v_N^2}{2} \right) - \frac{G}{2} \sum_{M,N=1}^n \frac{m_M m_N}{r_{MN}}. \quad (3.16)$$

This is the usual expression for an attractive force. (In writing the last equalities in (3.15) and (3.16) we have used a non-relativistic approximation.)

4. COMPARISON BETWEEN THE THEORIES ^{6),7)}

The main difference between the theory ⁶⁾ and ours, is that the Lagrangian density $\mathcal{L}^{(p)} + \mathcal{L}^{(i)}$ in the former is a homogeneous polynomial of degree two in the velocity variables, so the integrals (2.1) are not independent of the parameter s_N . Two different sets of parameters must be chosen depending on whether matter is described by a pointlike scheme or by a continuous one, in order to obtain in both cases the same equations of motion.

The conservation law (2.7) is replaced here by the following one

$$\partial_\alpha (T_{metric}^{\alpha\beta} + T^{(p)\alpha\beta}) = 0, \quad (4.1)$$

$T^{(i)}$ being equal to zero. As for the conservation law of mass

$\partial_\alpha (\mu u^\alpha) = 0$, it is replaced here by the conservation of a mass μ' modified by the field

$$\partial_\alpha (\mu' u^\alpha) \equiv \partial_\alpha [\mu (1 + h_{\rho\sigma} u^\rho u^\sigma)^{1/2} u^\alpha] = 0. \quad (4.2)$$

Note, however, that in Eq. (4.1) the mass which appears is the mass μ and not the modified mass μ' .

In our theory, on the contrary, the mass μ is conserved, but a "modified mass" appears in the conservation law of the total energy-momentum tensor, as can be seen by rewriting Eq. (2.7'') in the following way

$$\partial_\alpha [T_{metric}^{\alpha\beta} + T^{(p)\alpha\beta} (1 + h_{\rho\sigma} u^\rho u^\sigma)] = 0. \quad (4.3)$$

As for the observational facts the theory ⁶⁾ (with $\chi = \sqrt{6\pi G}$ and $a = -1$, as put by the authors) yields the same formulae as our theory, except that formula (3.9) is replaced by

$$\chi = 3 G \frac{M}{R}. \quad (4.4)$$

The predicted value for the bending of light rays is thus 25% smaller than in Einstein's theory.

For the fourth test one obtains instead of (3.12)

$$\Delta = \Delta_{GR} - \frac{1}{10} \Delta_{G.R.}.$$

Although agreement with the observational value for the bending of light rays is perhaps somewhat more doubtful in theory ⁶⁾ than in ours, the value of the parameter a in the former is, from an aesthetic point of view, more plausible than in ours. What seems to be a rather weak point in theory ⁶⁾ however, is that one obtains for the total energy of the system in the quasi-static approximation, instead of (3.16), the value

$$P^0 = \int (T_{metric}^{(f)00} + T^{(r)00}) d^3x \simeq \sum_{N=1}^n \left(1 + \frac{v_N}{2}\right)^2 - \frac{3}{2} G \sum_{M,N=1}^n \frac{m_M m_N}{r_{MN}}, \quad (4.5)$$

which is a rather unfamiliar expression.

CONCLUSIONS

We have constructed a Lorentz covariant theory of gravitation, with linear field equations and containing, apart from the gravitational coupling constant κ , only one arbitrary parameter a . Putting $\kappa = \sqrt{7\pi G}$ and $a = -3/5$, the theory yields the same formulae as general relativity for the perihelion shift, the red shift and the bending of light rays (excepting the numerical factor in the latter which is 12% smaller than in Einstein's theory). For the fourth test of general relativity, suggested by Shapiro, the predicted value for the measurable quantity is 4% greater than the value obtained by Shapiro from Einstein's theory.

The theory satisfies the consistency requirements of classical field theories since the field equations and the equations of motion are both deduced from a variational principle, and the latter agrees with the conservation law of the total energy-momentum tensor obtained from the Lagrangian of the system. The conservation law of mass is also satisfied.

In the non-relativistic limit the theory agrees with Newton's theory with respect to both the equations of motion and the total energy of the system.

ACKNOWLEDGEMENTS

I would like to thank very much the Institut des Hautes Etudes Scientifiques (where most of this work has been made) and its Director, Mr. L. Motchane, for his hospitality. I also thank Professor L. Van Hove for a critical reading of the manuscript.

REFERENCES AND FOOTNOTES

- 1) We restrict ourselves to spin two theories. As is well known, scalar theories do not yield the correct values for the observational effects and spin one theories give a repulsive force.
- 2) G.D. Birkhoff, Collected Math.papers, Am.Math.Soc. (New York, 1950), Vol. 2, pp. 737, 920, 929, 967 and 973.
- 3) A. Papapetroli, Praktika Acad. Athens 19, 224 (1944).
- 4) A. Capella, Cahiers de Phys. 144, 341 (1962) and 158, 446 (1964).
- 5) See, for instance, A. Trautman, in "Gravitation" - John Wiley, (New York, 1962). See also our Eqs. (2.8) and (2.9).
- 6) F.J. Belinfante and J.C. Swihart, Ann.Phys. 1, 168 (1957).
- 7) S. Lederer and M.A. Tonnelat, Nuovo Cimento 34, 884 (1964).
See also S. Mavridès, Nuovo Cimento 34, 1530 (1964).
- 8) A. Capella, Compt.rend. 260, 1341 (1965).
- 9) I.I. Shapiro, Phys.Rev.Letters 13, 789 (1964).
- 10) The signature of the metric is +, -, -, -. We use units such that $c = 1$. From now on the symbols (p), (i), (f) mean particles, interaction and field, respectively.
- 11) g is the determinant of $g_{\alpha\beta}$.
- 12) To calculate such an expression one must use a curvilinear system of co-ordinates in order to take into account the first derivative of $g_{\alpha\beta}$ contained in the covariant differentiation.
- 13) As usual the symbol $(\alpha\beta)$ means symmetrization in α and β .
- 14) See, for instance, V.J. Chazy, La Théorie de la Relativité et la Mécanique Céleste, Tome 1, p. 114, edited by Gauthier-Villars (Paris, 1928).
- 15) In general relativity $\chi = \sqrt{8\pi G}$.
- 16) M. Moshinsky, Phys.Rev. 80, 514 (1950).
- 17) For the sun the observations made in 1952 give the following value (as corrected by Mikhailov) $\chi = 1''.43 \pm 0''.18$. Formula (3.9) gives $\chi = 1''.53$. The mean value of the observation between 1919 and 1952 is, however, bigger than Einstein's value.