CURRENT COMMUTATORS AND THE MESON BOOTSTRAP

Chan Hong-Mo
CERN - Geneva

A B S T R A C T

The commutation relations between the time components and divergences of axial vector currents are applied to the meson spectrum suggested by bootstrap. In an idealized system, a unique solution is obtained for the masses and coupling constants of spin 0 mesons, which is in crude agreement with experiment. Of particular interest is the structural similarity between the relations here obtained and the usual bootstrap equations.

1. It has been suggested by Gell-Mann ¹⁾ that current operators satisfy certain commutation relations at equal times, generating a Lie algebra. These commutation relations are supposed to be exact, at least as far as strong interaction is concerned, although the Hamiltonian itself is not invariant under the related group. This assumption is a strong restriction on hadron dynamics. We shall investigate here some of its consequences on the meson system in connection with the meson bootstrap.

We consider only the space integrals of the time components of vector and axial vector currents ; these generate the "chiral" algebra of $U_3 \times U_3$ ¹⁾. The axial current in weak interaction, following Gell-Mann, is here identified as a member of the axial vector octet, though this is not necessary for most of our arguments. In addition, we assume ²⁾

- (A) the current operators exhaust themselves within a range of several hundred MeV, concentrating on resonances, so that only matrix elements connecting nearby "pole states" are appreciable ;
- (B) these matrix elements obey rapidly convergent dispersion relations, thus implying the Goldberger-Treiman approximation.

(A) and (B) are similar in effect respectively to the OPE and to the low-threshold dominance assumptions in bootstrap models ³⁾.

We take as our meson spectrum that suggested by bootstrap ³⁾. This gives at low energies ($\lesssim 1.5$ GeV) an octet and a singlet each of $0^\pm$ and $1^\pm$ mesons, roughly in that order ⁴⁾.

(s) $0^\pm(1^+)$, $0^\pm(8^+)$, $1^\pm(1^-)$, $1^\pm(8^-)$.

The spectrum is reasonably well supported by experiment and coincides with that contained in the $(\bar{6}, 6) - (6, \bar{6})$ representation of "chiral" $U_6 \times U_6$ ⁵⁾.

2.

Also, as in the bootstrap model ³⁾, we simplify the problem by assuming SU_3 symmetry and zero widths for resonances, although these should not be necessary.

2. Define ¹⁾

$$A_i(t) = -i \int \mathcal{A}_{i4}(\underline{x}, t) d^3 \underline{x}, \quad (1)$$

$$V_i(t) = -i \int \mathcal{V}_{i4}(\underline{x}, t) d^3 \underline{x},$$

where $\mathcal{A}_{i\mu}$ and $\mathcal{V}_{i\mu}$ for $i = 1, \dots, 8$ and $\mu = 1, \dots, 4$ are the axial vector and vector current densities. They obey the following commutation relations ¹⁾

$$[A_i(t), A_j(t)] = i f_{ijk} V_k(t) \quad (2)$$

$$[V_i(t), A_j(t)] = i f_{ijk} A_k(t) \quad (3)$$

$$[V_i(t), V_j(t)] = i f_{ijk} V_k(t) \quad (4)$$

Taking the relation (2) between O^- octet states ($|Pap\rangle$, $a = 1, \dots, 8$), and expanding the commutator over a complete set, one has

$$\begin{aligned} \sum_n \langle Pap | A_i | n \rangle \langle n | A_j | Pbp' \rangle - \sum_n \langle Pap | A_j | n \rangle \langle n | A_i | Pbp' \rangle \\ = i f_{ijk} \langle Pap | V_k | Pbp' \rangle \end{aligned} \quad (5)$$

By assumption (A), since the O^- mesons have a low mass, we need only insert intermediate states of low energies (≤ 1 GeV), namely states in our spectrum (S). Because of (B), and parity and C conservation, $|n\rangle$ in (5) can only be a $O^+(1^+)$, $O^+(8^+)$ or $1^-(8^-)$ state, denoted respectively by $|Sp\rangle$, $|Sap\rangle$, and $|Vap_r\rangle$, where $r = 1, 2, 3$ represents the polarization of the 1^- meson.

The integral A_i in (1) may be written,

$$A_i(0) = -i \int d^4x \theta(t) \partial_\mu \mathcal{A}_i^\mu(x, t) \quad (6)$$

Then from the assumed SU_3 and Lorentz invariance the matrix element $\langle Pap | A_i(0) | Vbp'r \rangle$, for example, may be expressed as

$$\begin{aligned} \langle Pap | A_i(0) | Pbp' \rangle &= \\ &= (-i)(2\pi)^3 \delta(\underline{p}-\underline{p}') \frac{1}{p_0-p'_0} \langle Pap | \partial_\mu \mathcal{A}_i^\mu(0) | Pbp' \rangle \\ &= (-i)(2\pi)^3 \delta(\underline{p}-\underline{p}') \frac{1}{p_0-p'_0} f_{\pi b} p_\mu \varepsilon^{\mu} F_1(s) \end{aligned} \quad (7)$$

The "form factor" $F_1(s)$ is a function only of the momentum transfer $s = (p-p')^2$, and by (B) is dominated by the $O^-(8^+)$ poles whose residues are proportional to G_1/f ⁶⁾. Other matrix elements are similar.

Substitution of (7) and similar formulae into (5) yields a p dependent relation. Following Fubini et al. ⁷⁾ we take the limit $|\underline{p}| \rightarrow \infty$ or $s \rightarrow 0$ to ensure minimum leakage from our intermediate states. This gives the equation ⁶⁾

4.

$$\begin{aligned}
 & \frac{G_1^2}{4m^2} \left(\sum_c f_{aic} f_{cjb} - \sum_c f_{ajc} f_{cib} \right) \\
 & + (-1) \frac{G_0^2}{(\tilde{\mu}^2 - \mu^2)^2} \left(\sum_c d_{aic} d_{cjb} - \sum_c d_{ajc} d_{cib} \right) \\
 & + (-1) \frac{G_0'^2}{(\mu^2 - \tilde{\mu}'^2)^2} (d_{aio} d_{oib} - d_{ajo} d_{oib}) \\
 & = (f^2) \left(\sum_k f_{ijk} f_{akb} \right)
 \end{aligned} \tag{8}$$

The three terms on the left correspond respectively to $1^-(8^-)$, $0^+(8^+)$ and $0^+(1^+)$ intermediate states. On the right, we have made use of the non-renormalization of the conserved vector form factors. The constant f measures the universal coupling of the $0^-(8^+)$ meson fields to $\partial_\mu A_i^\mu$ and may be estimated from $\pi^\pm$ decay. Equation (8) is thus a relation between the masses and coupling constants of mesons with known coefficients.

It is convenient to represent equations such as (8) by diagrams. Thus representing 0^- mesons by dotted lines and 1^- mesons by wavy lines, we have Fig. 1⁸⁾ which is strongly reminiscent of bootstrap equations.

Figure 1

Equation (8) is supposed to hold for all $a, b, i, j = 1, \dots, 8$ and may be regarded as a 64×64 matrix equation of the form

$$M_{ab,ij} = 0 \quad (9)$$

where M is Hermitian. For M to be zero it is necessary and sufficient to have all its eigenvalues zero. The eigenvectors of M , however, are known, being the Clebsch-Gordan coefficients in the series

$$8 \times 8 = 1 + 8 + 8' + 10 + \bar{10} + 27 \quad (10)$$

whose eigenvalues are linear combinations of SU_3 crossing matrix elements (Racah coefficients) ⁹⁾. By contracting (8) with Clebsch-Gordan coefficients, g_{ij}^X for $X = 1, 8, 8', 10, \bar{10}, 27$, one then obtains scalar equations. Since $M_{ab,ij}$ is antisymmetric in i and j , only contraction with g_{ij}^X for $X = 8', 10, \bar{10}$ gives non-trivial relations. Thus multiplying (8) with $g_{ij}^{8'} \cong f_{ijk}$ and summing over i, j one has

$$\frac{G_1^2}{4m^2} + \frac{5}{9} \frac{G_0^2}{(\mu^2 - \mu'^2)^2} + \frac{4}{9} \frac{G_2^{1/2}}{(\mu^2 - \mu'^2)^2} = f^2 \quad (11)$$

Similarly, by contracting with g_{ij}^X , $X = 10, \bar{10}$, one has

$$\frac{G_0^2}{(\mu^2 - \mu'^2)^2} = \frac{G_2^{1/2}}{(\mu^2 - \mu'^2)^2} \quad (12)$$

Equations (11) and (12) together are completely equivalent to (8).

Further conditions between masses and coupling constants can be obtained from (2) by taking it between meson states other than $0^-(8^+)$ mesons. We shall consider here only $0^\pm$ mesons as external particles, these being the lowest in mass, so as to ensure by (A) that the intermediate

6.

states remain within our spectrum (S). Taking (2) between $0^+(8^+)$ states, an analogous calculation to that described above gives an equation represented by Fig. 2⁸⁾. This in turn by contraction with Clebsch-Gordan coefficients yields the following scalar conditions⁶⁾

$$\frac{\tilde{G}_1^2}{4\tilde{m}^2} + \frac{5}{9} \frac{G_0^2}{(\tilde{\mu}^2 - \mu^2)^2} + \frac{4}{9} \frac{\tilde{G}_0^{1/2}}{(\mu^{1/2} - \tilde{\mu}^2)^2} = f^2 \quad (13)$$

$$\frac{G_0^2}{(\tilde{\mu}^2 - \mu^2)^2} = \frac{\tilde{G}_0^{1/2}}{(\mu^{1/2} - \tilde{\mu}^2)^2} \quad (14)$$

All other conditions obtained by taking (2) between $0^\pm$ meson states in (S) are trivially satisfied because of SU_3 symmetry or parity conservation.

Figure 2

One can of course do the same for the relations (3) and (4) as we have done for (2). However, since we have assumed invariance under SU_3 with V_i as its generators, Eqs. (3) and (4) only express the fact that V_i and A_i transform like octets, and are readily seen to give only trivial relations.

From (2)-(4) then, one obtains by this procedure only Eqs. (11)-(14). These are in a sense bootstrap equations connecting dynamical parameters of the meson system though they are yet insufficient to determine these parameters.

3. We consider next the commutation relations between the "chiralities" (1) and the divergences (time derivatives) of the currents. These involve the transformation properties of the Hamiltonian.

We note first the identity

$$[A_i, H] = -i \frac{\partial}{\partial t} A_i(t) = -i D_i(t) \quad (15)$$

where

$$D_i(t) = \int \partial_\mu A_i^\mu(x, t) d^3x$$

Then, following Gell-Mann ¹⁾, we assume that in the SU_3 limit

$$H = H_0 + \lambda u_0 \quad (16)$$

where H_0 is invariant under both V_i and A_i , while u_0 transforms like the zeroth component of the $(\bar{3}, 3) - (3, \bar{3})$ representation. Explicitly ¹⁾

8.

$$\begin{aligned}
 [V_i, u_j] &= i f_{ijk} u_k \\
 [V_i, v_j] &= i f_{ijk} v_k \\
 [A_i, u_j] &= -i d_{ijk} v_k \\
 [A_i, v_j] &= i d_{ijk} u_k
 \end{aligned}
 \tag{17}$$

$j, k = 0, 1, \dots, 8$

Thus we have

$$D_k = i [A_k, H] = d_{kol} \lambda v_l = \lambda \sqrt{\frac{2}{3}} v_k \tag{18}$$

which means that D_k also transform like components of $(\bar{3}, 3) - (3, \bar{3})$.
In particular

$$[A_j, D_k] = i d_{jkl} \sqrt{\frac{2}{3}} \lambda u_l \tag{19}$$

The unknown quantities u_l may be eliminated from (19) by commuting again with A_i , giving

$$\begin{aligned}
 [A_i, [A_j, D_k]] &= A_i A_j D_k - A_i D_k A_j - A_j D_k A_i + D_k A_j A_i \\
 &= d_{jkl} d_{ilm} D_m.
 \end{aligned}
 \tag{20}$$

$l, m = 0, 1, \dots, 8$

Similar relations can of course be obtained by commuting D 's with V 's, but these are trivial in the SU_3 limit.

Equation (20) is a non-linear relation between A's and D's which by the Goldberger-Treiman relations are essentially 0^- meson fields. It can thus be exploited as (2) in 2. to give "bootstrap conditions" on dynamical parameters. Since (20), unlike (2), depends on the assumption (16), these new conditions will in general be independent of (11)-(14).

We restrict the external states again to $0^\pm$ mesons so that intermediate states remain in (S). The computation proceeds in the same way as in 2. One needs only note that

$$\langle \alpha p | D_i | b p' \rangle = i(p_0 - p'_0) \langle \alpha p | A_i | b p' \rangle \quad (21)$$

Taking (20) between $0^-(1^+)$ and $0^+(1^+)$ states, one has, for example,

$$\begin{aligned} \langle P_p | A_i A_j D_k | S p' \rangle \\ = \sum_n \sum_m \langle P_p | A_i | n \rangle \langle n | A_j | m \rangle \langle m | D_k | S p' \rangle \end{aligned} \quad (22)$$

where because of SU_3 symmetry and C and P conservation the only possible intermediate states from (S) are $|n\rangle = 0^+(8^+)$ and $|m\rangle = 0^-(8^+)$. This is represented by Fig. 3⁸⁾. Equation (20) in the limit $|\underline{p}| = |\underline{p}'| \rightarrow \infty$ or $s \rightarrow 0$ then yields⁶⁾

$$\begin{aligned} (-1) \frac{\tilde{G}_0'}{(\mu'^2 - \tilde{\mu}^2)} \cdot \frac{G_0}{(\tilde{\mu}^2 - \mu^2)} - \frac{G_0'}{(\mu^2 - \tilde{\mu}'^2)} \left\{ (\mu'^2 - \tilde{\mu}^2) + (\mu^2 - \tilde{\mu}'^2) - 2(\tilde{\mu}^2 - \mu^2) \right\} \\ = G_0'' f^2 \end{aligned} \quad (23)$$

Figure 3

With $0^-(8^+)$ and $0^+(1^+)$ mesons as external particles, the possible intermediate states are represented by the diagrams of Fig. 4⁸⁾. The limit $|p| \rightarrow \infty$ yields a matrix condition on the dynamical parameters, depending on four indices, a, i, j, k . One may, as in 2. extract scalar conditions from this by contracting with Clebsch-Gordan coefficients, e.g., by multiplying with g_{jk}^X for $X = 1, 8, 8', 10, \bar{10}, 27$, and summing over j and k . One obtains the following equations :

i) $X = 1$

$$-\frac{10}{3} \frac{G_0^2}{(\tilde{\mu}^2 - \mu^2)^2} \frac{(\mu^2 - \tilde{\mu}^2)}{(\mu^2 - \tilde{\mu}^{1/2})} + 12 \frac{G_0^{1/2}}{(\mu^2 - \tilde{\mu}^{1/2})^2} + 6 \frac{G_1^2}{4m^2} \frac{(m^2 - \mu^2)}{(\tilde{\mu}^{1/2} - \mu^2)} = \frac{16}{3} f^2 \quad (24)$$

ii) $X = 8$

$$\frac{5}{3} \frac{G_0^2}{(\tilde{\mu}^2 - \mu^2)^2} - \frac{2}{3} \frac{G_0^2}{(\tilde{\mu}^2 - \mu^2)^2} \frac{(\mu^2 - \tilde{\mu}^2)}{(\tilde{\mu}^{1/2} - \mu^2)} + \frac{4}{3} \frac{G_0^{1/2}}{(\tilde{\mu}^{1/2} - \mu^2)^2} + 3 \frac{G_1^2}{4m^2} \frac{(m^2 - \mu^2)}{(\tilde{\mu}^{1/2} - \mu^2)} = \frac{5}{3} f^2 \quad (25)$$

iii) $X = 8'$

$$\frac{G_1^2}{4m^2} (m^2 - \tilde{\mu}^{1/2}) = 0 \quad (26)$$

iv) $X = 10, \bar{10}$, trivial

v) $X = 27$

$$-\frac{2}{3} \frac{G_0^2}{(\tilde{\mu}^2 - \mu^2)^2} \frac{(\mu^2 - \tilde{\mu}^2)}{(\tilde{\mu}^{1/2} - \mu^2)} - 2 \frac{G_1^2}{4m^2} \frac{(m^2 - \mu^2)}{(\tilde{\mu}^{1/2} - \mu^2)} + \frac{4}{3} \frac{G_0^{1/2}}{(\tilde{\mu}^{1/2} - \mu^2)^2} = 0 \quad (27)$$

Figure 4

With $0^+(8^+)$ and $0^-(1^+)$ mesons as external particles, the calculation is very similar and yields four further non-trivial conditions ;

$$-\frac{10}{3} \frac{G_0^2}{(\tilde{\mu}^2 - \mu^2)^2} \frac{(\tilde{\mu}^2 - \mu^2)}{(\mu^{12} - \tilde{\mu}^2)} + 12 \frac{\tilde{G}_0^{1/2}}{(\mu^{12} - \tilde{\mu}^2)^2} + 6 \frac{\tilde{G}_1^2}{4\tilde{m}^2} \frac{(\tilde{m}^2 - \mu^2)}{(\mu^{12} - \tilde{\mu}^2)} = \frac{16}{3} f^2 \quad (28)$$

$$\begin{aligned} \frac{5}{3} \frac{G_0^2}{(\tilde{\mu}^2 - \mu^2)^2} - \frac{2}{3} \frac{G_0^2}{(\tilde{\mu}^2 - \mu^2)^2} \frac{(\tilde{\mu}^2 - \mu^2)}{(\mu^{12} - \tilde{\mu}^2)} + \frac{4}{3} \frac{\tilde{G}_0^{1/2}}{(\mu^{12} - \tilde{\mu}^2)^2} + \\ + 3 \frac{\tilde{G}_1^2}{4\tilde{m}^2} \frac{(\tilde{m}^2 - \mu^2)}{(\mu^{12} - \tilde{\mu}^2)} = \frac{5}{3} f^2 \end{aligned} \quad (29)$$

$$\frac{\tilde{G}_1^2}{4\tilde{m}^2} (\tilde{m}^2 - \mu^{1/2}) = 0 \quad (30)$$

$$-\frac{2}{3} \frac{G_0^2}{(\tilde{\mu}^2 - \mu^2)^2} \frac{(\tilde{\mu}^2 - \mu^2)}{(\mu^{12} - \tilde{\mu}^2)} - 2 \frac{\tilde{G}_1^2}{4\tilde{m}^2} \frac{(\tilde{m}^2 - \mu^2)}{(\mu^{12} - \tilde{\mu}^2)} + \frac{4}{3} \frac{\tilde{G}_0^{1/2}}{(\mu^{12} - \tilde{\mu}^2)^2} = 0 \quad (31)$$

Again the relation (20) may be taken between $0^-(8^+)$ and $0^+(8^+)$ states. This yields an equation dependent on five indices, which on contraction gives a large number of scalar conditions. These, however, all turn out to be dependent on Eqs. (11)-(14) and (23)-(31), and will be omitted. Other combinations of $0^\pm$ external particles from (S) yield only trivial conditions. By applying our procedure to (20) therefore, we obtain only the eight preceding equations which are of present interest.

4. The two sets of bootstrap Eqs. (11)-(14) and (23)-(31), deduced respectively from (2) and (20), together imply the following solution :

$$(\mu'^2 - \tilde{\mu}^2) = \frac{1}{2} (\tilde{\mu}^2 - \mu^2) = (\mu^2 - \mu'^2) = \Delta \quad (a)$$

$$\frac{\tilde{G}_0'^2}{(\mu'^2 - \tilde{\mu}^2)^2} = \frac{G_0^2}{(\tilde{\mu}^2 - \mu^2)^2} = \frac{G_0'^2}{(\mu^2 - \mu'^2)^2} = f^2 \quad (b)$$

$$G_1^2/4m^2 = \tilde{G}_1^2/4\tilde{m}^2 = 0 \quad (c) \quad (32)$$

$$G_0'' = \pm G_0 \quad (d)$$

$$G_0' G_0 < 0 \quad (e)$$

If one identifies the axial current in weak interaction with components of $\mathcal{A}_1^\mu$, f is a quantity estimable independently from $\pi^\pm$ decay¹⁰⁾. The solution then depends on the single parameter Δ , the "mass-splitting".

The relation (32.a) has been deduced by Gell-Mann as a consequence of approximate $U_3 \times U_3$ symmetry for small λ , (16)¹⁾. It is interesting that this approach, which does not explicitly assume λ to be small, gives the same result. Assigning σ (~ 350 MeV) to $0^+(1^+)$, π, K, η (410 MeV) to $0^-(8^+)$, ε_0, κ (730 MeV) to $0^+(8^+)$, and X_0 (960 MeV) to $0^-(1^+)$ ¹¹⁾, one finds that the formula (32.a) gives the correct ordering of masses, but the numerical agreement is poor.

Equation (32.b) implies that the coupling constants $G_0^2, G_0'^2, \tilde{G}_0'^2$ are all of order λ , and therefore small. This is in qualitative agreement with the narrow widths of ε_0 and κ inspite of their 0^+ assignments. We note that bootstrap calculations also give a small value for G_0 ³⁾.

The vanishing of G_1 (connected to ρ decay) is of course in poor agreement to experiment. Because of the large denominators dividing G_1^2 in sum rules such as (11), however, a small distortion from our idealized system is most likely sufficient to remedy this inconsistency.

Inspite of the poor agreement of this present calculation to experiment, it has shown some very interesting structural points, especially in its similarity to usual bootstrap models. It is hoped that by putting in more realistic dynamical approximations such as SU_3 splitting and finite widths for resonances, the situation will be much improved.

ACKNOWLEDGEMENT

The author wishes to thank Drs. E. Abers and G. Segrè for several instructive conversations.

A P P E N D I X

List of symbols and conventions.

m	=	mass of $1^-(8^-)$ mesons,
$\tilde{m}$	=	mass of $1^+(8^-)$ mesons,
μ	=	mass of $0^-(8^+)$ mesons,
$\tilde{\mu}$	=	mass of $0^+(8^+)$ mesons,
μ'	=	mass of $0^-(1^+)$ meson,
$\tilde{\mu}'$	=	mass of $0^+(1^+)$ meson.

The various couplings between the mesons are :

$$\begin{aligned}
 [0^-(8^+), 0^-(8^+), 1^-(8^-)] &: f_{ijk} p_\mu \varepsilon^{r\mu} G_1 \\
 [0^-(8^+), 0^+(8^+), 1^+(8^-)] &: f_{ijk} p_\mu \varepsilon^{r\mu} \tilde{G}_1 \\
 [0^-(8^+), 0^-(8^+), 0^+(8^+)] &: d_{ijk} G_0 \\
 [0^-(8^+), 0^-(8^+), 0^+(1^+)] &: d_{ijo} G'_0 \\
 [0^-(8^+), 0^+(8^+), 0^-(1^+)] &: d_{ijo} \tilde{G}'_0 \\
 [0^-(1^+), 0^-(1^+), 0^+(1^+)] &: \sqrt{2/3} G''_0.
 \end{aligned}$$

f_{ijk} and d_{ijk} are the same as in Gell-Mann ¹⁾

$$\begin{aligned}
 a^\mu b_\mu &= a_0 b_0 - \underline{a} \cdot \underline{b}. \\
 \varepsilon^\mu \varepsilon_\mu &= -1 ; \quad \varepsilon^\mu_r \varepsilon^\nu_r = -g_{\mu\nu} + p_\mu p_\nu / m^2.
 \end{aligned}$$

REFERENCES AND FOOTNOTES

- 1) M. Gell-Mann, Physics 1, 63 (1964).
- 2) R. Dashen and M. Gell-Mann, CALTECH preprints 68/36, 68/37.
- 3) See, e.g., Chan Hong-Mo and C. Wilkin, Phys.Letters 13, 263 (1964) ;
CERN preprint TH. 569 (1965).
- 4) Meson multiplets are denoted by $J^P(X^C)$ where
X = dimension of SU_3 representation,
C = charge conjugation parity.
- 5) R.P. Feynman, M. Gell-Mann and G. Zweig, Phys.Rev.Letters 13, 678
(1964) ;
K. Bardacki, J.M. Cornwall, P.G.O. Freund and B.W. Lee, Phys.Rev.
Letters 13, 698 (1964).
- 6) For a list of notations, see the Appendix.
- 7) S. Fubini and G. Furlan, Physics 1, 229 (1965).
A more elegant, relativistic covariant way of obtaining the same
result has been developed by S. Fubini, G. Furlan and C. Rossetti,
CERN preprint TH. 598 (1965). We use this non-covariant approach
here for simplicity.
- 8) Dotted lines denote 0^- mesons, full lines 0^+ mesons, wavy lines
 1^- mesons, springs 1^+ mesons.
- 9) See, e.g., Chan Hong-Mo, P.C. DeCelles and J.E. Paton, Nuovo Cimento
33, 70 (1964).
- 10) See, e.g., M. Gell-Mann, Phys.Rev. 125, 1067 (1962).
- 11) See, e.g., G. Goldhaber, UCRL report, review of meson resonances.