CM-P00057371

ARCHIVES

THE ABSORPTION MODEL AND PRODUCTION OF 2^+ MESONSH. Högaasen, J. Högaasen ^{*)}, R. Keyser and B.E.Y. Svensson

CERN -- Geneva

A B S T R A C T

The absorption model for high energy peripheral production of resonances is applied to the production of spin parity 2^+ mesons in pion-nucleon interaction with the twofold purpose of testing the model against the experimental data and of discussing some approximations made within the model. By a detailed numerical investigation of the partial wave expansion we show how the impact parameter approximation, often used in the application of the model, may significantly affect both the predicted cross-section and the decay distribution. Moreover, the cross-section, but not so much the decay distribution, is sensitive to the detailed way in which each partial wave in the model is modified by the absorption. We trace these features back to the fact that the low partial waves, although strongly suppressed as compared to the unmodified Born approximation, still play a major role. The comparison with experiment for production of f_0 (pion exchange) shows that the slope of the cross-section is correctly predicted but that the absolute magnitude is too big by a factor of around two. For $A_2(1310)$ production (rho exchange) the slope of the cross-section can be made to agree with experiment only after the introduction of a fairly strong form factor; the absolute magnitude and the energy dependence of the cross-section is wrong. These results strengthen the conclusion that the absorption model is unable to reproduce the experimental data for production of high spin resonances, especially if vector meson exchange has to be assumed.

The spin 2 formalism, including decay widths and decay distributions, is presented in the Appendices.

*) On leave from Laboratoire de Physique Théorique, Département de Physique Nucléaire, Orsay (S. et O.), France.

1. INTRODUCTION

Among the boson resonances presently known, there are three classified as having spin 2 and positive parity. These are the pion-pion resonance f_0 of mass 1253 MeV and width 100 MeV ¹⁾, the rho-pion resonance A_2 of mass 1310 MeV and width 80 MeV ¹⁾, and the recently discovered kaon-pion resonance K^{**} of mass 1400 MeV and width 145 MeV ²⁾. Although there still remain some doubts as to the proper spin parity assignment of these resonances, we will, in our considerations, assume that the 2^+ assignment is the correct one.

One purpose of this paper is to see how well the production mechanism for the two first of these resonances, as they appear in pion-nucleon interaction, can be understood in terms of some currently used models. Experimentally, the production cross-section for these resonances is strongly peaked in the direction of the incoming meson momentum, indicating a peripheral interaction and suggesting a one-particle exchange (OPE) production mechanism ³⁾. However, as is now well known, the simplest form of the OPE model, the unmodified Born term model (BTM), fails to reproduce the experimentally observed peaking and has to be corrected, either by an over-all form factor ^{4),5)} or by inclusion of effects due to competition from other open channels, as is done in the absorption model ⁶⁻⁹⁾ and in the K matrix model ¹⁰⁾. Here we shall only consider the BTM and the absorption model.

The other purpose of the paper is to discuss some approximations made within the absorption model. First, we investigate the impact parameter approximation (IPA) to the exact partial wave summation (EPWS), which has frequently been used in the application of the model so far. We do this by making a thorough numerical discussion of the partial wave expansion and by comparing the predicted cross-section and the decay distribution (i.e., the resonance spin space density matrix) in the IPA with the results in the EPWS. Second, we discuss the sensitivity of the model to the detailed way in which each partial wave is modified by the absorption, i.e., to the angular momentum dependence of the initial and final state elastic scattering phase shifts producing the absorption.

The plan of the paper is the following: in Section 2 we give the expressions for the production cross-section and the spin space density matrix elements of the 2^+ resonance in the unmodified Born approximation assuming exchange of a pseudoscalar or a vector meson. Section 3 contains a short description of the absorption model, but is mainly devoted to a numerical investigation of the impact parameter approximation and of the detailed angular momentum dependence assumed for the absorption. The results of our calculations are compared with experiment in Section 4, while Section 5 contains the conclusions. In Appendix A we have collected the relevant formulae for the helicity states of a spin 2 particle, as well as the couplings used and the corresponding decay widths. Appendix B contains the angular distributions in the decays $2^+ \rightarrow 0^-0^-$ and $2^+ \rightarrow 1^-0^-$, while Appendix C lists the helicity amplitudes in the Born approximation.

2. CROSS-SECTION AND SPIN DENSITY MATRIX ELEMENTS IN THE BORN APPROXIMATION

In this section we give the expressions for the cross-section and the polarization, expressed in terms of its spin space density matrix elements as defined in Appendix B, for a 2^+ boson produced in pseudoscalar meson-nucleon interaction, assuming that the process occurs through exchange of a pseudoscalar or a vector meson.

The relevant Feynman diagram is shown in Fig. 1, which also indicates our notations. With the four-momentum of particle a denoted by a_μ , its mass by m_a (etc. for the other particles), we have $e_\mu = c_\mu - a_\mu = b_\mu - d_\mu$. The momentum transfer squared is $-t = e_\mu e_\mu^{(11)}$. Moreover, a nucleon of mass m is described by the usual Dirac spinor u , normalized to $\bar{u}u = 2m$, on which

act the conventional Dirac matrices γ_5 and γ_μ . Finally, the spin 2 particle is characterized by a symmetrical traceless second order tensor (see Appendix A). Using the couplings of Appendix A we arrive at the following expressions for the matrix element of the process in Fig. 1:

for pseudoscalar meson exchange:

$$M^{(PS)} = \frac{4 G g_2}{m_c} \cdot \bar{u}(d) \gamma_5 u(b) \cdot \frac{1}{m_c^2 - t} \cdot a_\rho \varphi_{\rho\sigma}^* a_\sigma, \quad (2.1)$$

for vector meson exchange

$$M^{(V)} = \frac{i f_2 (G_V + G_T)}{m_c^2} \cdot \bar{u}(d) \left[\gamma_\mu + i y \frac{1}{m_b + m_d} d_\mu \right] u(b) \cdot \frac{1}{m_c^2 - t} \cdot \varepsilon_{\mu\rho\tau\omega} \varphi_{\rho\sigma}^* a_\sigma a_\tau c_\omega, \quad (2.2a)$$

$$y = \frac{2 G_T}{G_V + G_T}. \quad (2.2b)$$

The normalization of the amplitudes in Eqs. (2.1 - 2.2) is such that the cross-section is given by

$$\frac{d\sigma}{dt} = \frac{1}{64 \pi s q^2} \cdot \frac{1}{2} \cdot \sum |M|^2, \quad (2.3)$$

where, for vanishing polarization in the initial state and no polarization measurements in the final state, the summation extends over all spin states. Furthermore, s is the square of the total c.m.s. energy and q is the magnitude of the three-momentum for either particle in the initial state.

Performing the summation using the appropriate projection operators ¹²⁾ gives the following expression for the cross-section:

for pseudoscalar meson exchange:

$$\frac{d\sigma_{\text{Born}}^{(\text{PS})}}{dt} = \frac{G^2}{4\pi} \cdot \frac{g_a^2}{4\pi} \cdot \frac{8\pi}{3s q^2} \cdot \frac{a_c^4 \cdot [(m_b - m_d)^2 - t]}{m_c^2 \cdot (m_e^2 - t)^2} \quad (2.4)$$

for vector meson exchange:

$$\frac{d\sigma_{\text{Born}}^{(\text{V})}}{dt} = \frac{(G_V + G_T)^2}{4\pi} \cdot \frac{f_q^2}{4\pi} \cdot \frac{\pi}{4s q^2} \cdot \frac{a_c^2}{m_c^4} \cdot \frac{1}{(m_e^2 - t)^2} \quad (2.5a)$$

$$\cdot \left[\{ (m_b - m_d)^2 - t \} \cdot a_c^2 \cdot m_c^2 + \frac{1}{2} \hat{S} \cdot \left\{ v^2 - y^2 \cdot \frac{t}{(m_b + m_d)^2} \right\} \right],$$

$$v = q - y \quad (2.5b)$$

Here, a_c is the magnitude of the three-momentum of particle a in the rest frame of particle c , given by

$$a_c^2 = \frac{1}{4m_c^2} \cdot [t - (m_c + m_a)^2] \cdot [t - (m_c - m_a)^2], \quad (2.6)$$

and, with Θ denoting the c.m.s. production angle, q' the magnitude of the c.m.s. three-momentum of either particle in the final state,

$$\hat{S} = s q^2 q'^2 \sin^2 \Theta \quad (2.7)$$

With the definition of the spin space density matrix elements $\rho_{\ell,m}$ for the 2^+ meson (particle c in the notation of Fig. 1) as given in Appendix B, we find:

for pseudoscalar meson exchange:

$$\rho_{0,0} = 1, \quad (2.8a)$$

$$\text{all others} = 0. \quad (2.8b)$$

for vector meson exchange:

$$\rho_{1,1} = \rho_{-1,-1} = \frac{1}{2}, \quad (2.9a)$$

$$\begin{aligned} \rho_{1,-1} = \rho_{-1,1} = & \frac{\hat{S} \cdot \left\{ v^2 - y^2, \frac{t}{(m_b + m_d)^2} \right\}}{\frac{i}{4} \cdot \left\{ (m_b - m_d)^2 - t \right\} \cdot a_c^2 m_c^2 + \frac{1}{2} \hat{S} \cdot \left\{ v^2 - y^2, \frac{t}{(m_b + m_d)^2} \right\}} \end{aligned} \quad (2.9b)$$

$$\text{all others} = 0. \quad (2.9c)$$

The simple values for $\rho_{0,0}$ and $\rho_{1,1}$ are explained in Ref. 13); the value of $\rho_{1,-1}$ has to be derived explicitly.

The similarity between the formulae (2.4 - 2.9) and the corresponding formulae for the production of vector mesons⁵⁾ should be noted. The essential difference is the occurrence of an extra factor a_c^2 as compared to the Jackson-Pilkuhn formulae. This is due to the fact that the incoming pseudo-scalar particle and the exchanged particle now have a relative orbital angular

momentum one unit higher than for vector meson production. We also note another common feature with vector meson production: there is no interference between the BTM amplitudes with pseudoscalar and with vector meson exchange, so that the two cross-sections (2.4) and (2.5) add incoherently.

As is by now well known, the unmodified Born approximation has several defects, that we recall here briefly:

- i) the slope of the production cross-section is not steep enough;
- ii) the absolute magnitude of the production cross-section is too big in most cases;
- iii) the unitarity bounds are badly violated for low angular momenta;
- iv) the energy dependence is often wrong, in particular for vector meson exchange.

The point i) may be seen explicitly from Eqs. (2.4) and (2.5): the high power of t in the numerator dominates over the propagator, especially in the case of vector meson exchange, where also m_ρ is large. The introduction of rapidly changing form factors, mostly chosen in a pure phenomenological way, can remedy this point, and at the same time the points ii) and iii), but casts doubt on the whole approach. One way out of this dilemma was the suggestion of the absorption model⁶⁻⁸⁾, to which we now turn. We will, however, in Section 4b return to the question of the necessity of form factors.

3. THE ABSORPTION MODEL. EXACT PARTIAL WAVE SUMMATION VERSUS THE IMPACT PARAMETER APPROXIMATION

Concentrating on the point iii) in the list at the end of the previous section by introducing a strong damping of the low partial waves, the absorption model also gives a better, although not always a good, angular distribution i) and an over-all reduction ii) ¹⁴⁾. It also appears less phenomenological, in that the damping of the low partial waves is related to the elastic scattering data, known at least in the initial state.

Here we shall only give a brief description of the absorption model; for a full account of the theory we refer to Refs. ^{6-9), 15)}. To this end, let us write the partial wave expansion of the helicity amplitudes ¹⁶⁾ in the Born approximation, for a process like that of Fig. 1, as

$$\begin{aligned} \langle \lambda_c \lambda_d | B(\cos \Theta) | \lambda_a \lambda_b \rangle = \\ = \langle \lambda_c \lambda_d | A | \lambda_a \lambda_b \rangle \cdot \sum_{j_{\min}}^{\infty} (2j+1) \cdot c_{\lambda, \lambda'}^j(z) \cdot d_{\lambda, \lambda'}^j(\Theta) \cdot k(j) + \\ + \sum_{j_{\min}}^{\infty} \langle \lambda_c \lambda_d | T^j | \lambda_a \lambda_b \rangle \cdot d_{\lambda, \lambda'}^j(\Theta) \cdot k(j) , \end{aligned} \quad (3.1a)$$

$$\lambda = \lambda_a - \lambda_b, \quad \lambda' = \lambda_c - \lambda_d, \quad (3.1b)$$

$$j_{\min} = \max(|\lambda|, |\lambda'|), \quad (3.1c)$$

$$z = \frac{1}{2q_1 q_1'} \left[m_e^2 - m_b^2 - m_d^2 + 2E_b E_d \right]. \quad (3.1d)$$

We have here followed the notation of Section 2 and further denoted the energy of particle a by E_a , its helicity by λ_a , etc. (in the particular reaction of Fig. 1, $\lambda_a = 0$). The whole angular dependence of the helicity amplitudes is contained in the rotation coefficients $d_{\lambda, \lambda'}^j(\theta)$ ¹⁷⁾, i.e., the coefficients $\langle \lambda_c \lambda_d | A | \lambda_a \lambda_b \rangle$ and the "extraordinary terms" $\langle \lambda_c \lambda_d | X^j | \lambda_a \lambda_b \rangle$, which in fact vanish for all but the very lowest values of j , are functions only of the energy (and of the masses of the particles involved). Moreover, for the rotation coefficients of the second kind $c_{\lambda, \lambda'}^j(z)$, we use the definition¹⁸⁾

$$\left(\sin \frac{\theta}{2}\right)^{|\lambda - \lambda'|} \cdot \left(\cos \frac{\theta}{2}\right)^{|\lambda + \lambda'|} \cdot \frac{1}{2 - \cos \theta} = \sum_{j_{\min}}^{\infty} (2j+1) \cdot c_{\lambda, \lambda'}^j(z) \cdot d_{\lambda, \lambda'}^j(\theta). \quad (3.2)$$

In the unmodified Born approximation, finally, one has $k(j) \equiv 1$ for all j values. The absorption model, on the other hand, consists in using for "the absorption factor" $k(j)$ the expression

$$k(j) = \exp[i\delta_i(j)] \cdot \exp[i\delta_f(j)], \quad (3.3)$$

where $\delta_i(j)$ is the elastic scattering phase shift for the process $a+b \rightarrow a+b$ while $\delta_f(j)$ is the corresponding quantity for $c+d \rightarrow c+d$, and where it is assumed that the elastic scattering is helicity conserving.

To determine these phase shifts, it is assumed that the elastic cross-section has the form (σ_{tot} is the total cross-section)

$$\frac{d\sigma_{\text{el}}}{dt} = \frac{\sigma_{\text{tot}}^2}{16\pi} \cdot \exp[\Omega t]. \quad (3.4)$$

This parametrization is a rather good approximation to the experimental data at small momentum transfer. One then derives

$$\exp[i\delta(j)] \approx 1 - C \cdot \exp[-\gamma j^2], \quad (3.5a)$$

$$C = \frac{\sigma_{tot}}{4\pi A} \quad (3.5b)$$

$$\gamma^{-1} = 2 q^2 A, \quad (3.5c)$$

where $C \leq 1$ for consistency.

Thus the absorption factor $k(j)$ is determined from the elastic scattering, which however is known experimentally only in the initial state in general. Although the derivation of the model seems to require the elastic scattering in the initial and in the final state to be equal, in the application of the model one often treats the "absorption parameter" C_f and A_f of the elastic scattering in the final state as free parameters, varying within reasonable limits.

Here we shall not discuss any more the physical motivation for the absorption model, but only investigate the impact parameter approximation (IPA), frequently used in the application of the model, as opposed to the exact partial wave summation (EPWS) of Eq. (3.1). At the same time, we shall study the sensitivity of the results to the choice of the absorption factor, i.e., to the particular function of j used for $\exp[i\delta(j)]$ in Eq. (3.3).

Consider the first (infinite) sum of Eq. (3.1)

$$S = \sum_{j=\min}^{\infty} (g_{j+1}) \cdot c_{\lambda, \lambda'}^j(z) \cdot d_{\lambda, \lambda'}^j(\theta) \cdot k(j), \quad (3.6)$$

where for simplicity we shall suppose $\lambda \geq |\lambda'|$. Following Durand and Chiu¹⁵⁾, we make the approximations

$$\begin{aligned} c_{\lambda, \lambda'}^j(z) &\approx R \cdot x^\alpha \cdot \left(\frac{\varepsilon}{g}\right)^\alpha \cdot \left(1 + \frac{\varepsilon^2}{4}\right)^{\beta/2} \cdot K_\alpha(\varepsilon x) \approx \\ &\approx R \cdot x^\alpha \cdot \left(\frac{\varepsilon}{g}\right)^\alpha \cdot K_\alpha(\varepsilon x), \end{aligned} \quad (3.7a)$$

$$\begin{aligned} d_{\lambda, \lambda'}^j(\theta) &\approx R^{-1} \cdot (x^2 - \alpha)^{-\alpha/2} \cdot J_\alpha(\omega \sqrt{x^2 - \alpha}) \approx \\ &\approx R^{-1} \cdot x^{-\alpha} \cdot J_\alpha(\omega x), \end{aligned} \quad (3.7b)$$

$$R^2 = \frac{(j - \lambda)! \cdot (j + \lambda')!}{(j + \lambda)! \cdot (j - \lambda')!}, \quad (3.7c)$$

$$\varepsilon = \sqrt{g(z-1)}, \quad \omega = g \sin \theta/2, \quad (3.7d)$$

where $J_n(y)$ and $K_n(y)$ denote the ordinary and the hyperbolic Bessel function respectively, and where, with $\alpha = \lambda - \lambda'$ and $\beta = \lambda + \lambda'$, we have put

$$x^2 = (j + 1/2)^2 - \frac{1}{4}(\alpha - 1)^2 - \frac{1}{4}\beta^2. \quad (3.7e)$$

The approximations (3.7a) and (3.7b) are valid for small values of ξ and ω , respectively, but do not necessarily require j to be large. Furthermore, in converting the sum in Eq. (3.6) into an integral, an approximation which is good when many partial waves contribute, and noticing that $(2j+1)dj = 2x dx$, we find

$$S \approx I = 2 \cdot \left(\frac{\xi}{2}\right)^\alpha \cdot \int_{x_{\min}}^{\infty} x dx \cdot K_\alpha(\xi x) \cdot J_\alpha(\omega x) \cdot k(j(x)), \quad (3.8)$$

where j as a function of x is given by Eq. (3.7e) and $x_{\min}$ is obtained from Eq. (3.7e) for $j = j_{\min}$.

Further simplifications consist in replacing $x_{\min}$, the lower limit of integration in Eq. (3.8) by $j_{\min}$, or even by 0¹⁹⁾, and in writing $j = x - \frac{1}{2}$, arriving at the formula first used in this context by Gottfried and Jackson⁷⁾. Of these approximations, it is not difficult to keep track of the first. Consequently, for the simplicity of discussion, we shall from now on replace $x_{\min}$ in Eq. (3.8) by 0, unless otherwise stated. The second simplification has the effect of shifting the argument in the Bessel functions and in the absorption factor slightly from its more correct value (3.7e).

One sees this shift between the arguments j and x , especially well from Fig. 2, where we have plotted, for $\alpha = 0$ and $\beta = 1$ (i.e., for $\lambda = \lambda' = \frac{1}{2}$) the corresponding rotation and Bessel functions as functions of j and x respectively, for the values $z = 1.2$ and $\cos \theta = 0.8$, which for the actual expansion parameters ξ^2 and ω^2 means $\xi^2 = \omega^2 = 0.4$. In this figure we have also plotted the absorption factors

$$k_a(j) = 1 - \exp[-0.03 \cdot j^2], \quad (3.9a)$$

$$k_b(j) = 1 - \exp[-0.03 \cdot (j - 1/2)^2], \quad (3.9b)$$

i.e., for (3.9a) we have in Eq. (3.3) and (3.5) chosen $C_i = C_f = 1$, $\gamma_i = \gamma_f = 0.03$, corresponding to a slightly stronger absorption than dictated by experiment (cf. Section 4). The form (3.9b) is chosen to allow for a comparison of the results obtained by the EPWS and by the IPA. At the same time, this will give us the opportunity to investigate the effects of a change in the absorption factor on the amplitudes, since the expressions (3.6) and (3.8) are proportional to the helicity amplitudes, if for the time being we may neglect the extraordinary terms in (3.1).

The effect of the approximations leading to Eq. (3.8) can be seen from Fig. 3, where we have plotted the functions

$$u(j) = (2j+1) \cdot c_{1/2, 1/2}^j(z) \cdot c_{1/2, 1/2}^j(\theta) \cdot k(j) \quad (3.10)$$

$$v(x) = 2x \cdot K_0(\epsilon x) \cdot J_0(\omega x) \cdot k(x) \quad (3.11)$$

first without absorption, i.e., with $k(j) \equiv 1$ (Fig. 3a), then with the two different forms of $k(j)$ defined in Eq. (3.9); the area under the respective curves gives the value of the sum S and the integral I . The absorption reduces the first positive peak (observe the different scales in Fig. 3a and 3b) considerably, the second negative peak less, so that they become of the same order of magnitude, resulting more or less in a cancellation between them both

in the sum and in the integral. As a consequence, even a small change $\Delta k(j)$ in the absorption factor $k(j)$, which means large values of $(\Delta k(j)/k(j))$ for the smallest j values where $k(j)$ is very small, will produce large changes in $u(j)$ and $v(x)$, resulting in very different values for the total area under the curves. This can be seen from Table 1, in which we compare, for all values of α and β relevant to production of 2^+ mesons, the results obtained from the EPWS (3.6) and the IPA (3.8), both for the two forms (3.9) of the absorption factor. As before, we have taken $z = 1.2$ and $\cos \theta = 0.8$, but give in the table also the values of the only non-vanishing sum and integral for $\cos \theta = 1.0$.

Before we consider in detail the results collected in Table 1, we want to make a few other remarks with respect to the result of Fig. 3.

Although the low partial waves in the absorption model are very much suppressed as compared to the unmodified Born approximation, their contribution to the sum (integral) is still considerable. As a consequence, the omission of the extraordinary terms of Eq. (3.1), always contributing to the very lowest partial waves, might not be a good approximation. We have left them out from the partial wave considerations of this section, but we always keep them when we calculate experimental quantities (cross-section, density matrix elements).

	$\alpha \beta$ ($\lambda \lambda'$)	S_a	S_b	I_a	I_b	$\frac{S_a - S_b}{S_a} \%$	$\frac{S_a - I_b}{S_a} \%$	$\frac{S - I_a}{S_a} \%$
$\cos \theta = 1$	0 1 ($\frac{1}{2} \frac{1}{2}$)	0.853	0.664	0.994	0.781	22	8.5	-16.5
	0 1 ($\frac{1}{2} \frac{1}{2}$)	-2.02 $\cdot 10^{-2}$	-5.72 $\cdot 10^{-2}$	4.82 $\cdot 10^{-2}$	-1.48 $\cdot 10^{-2}$	183	-27	-334
$\cos \theta = 0.8$	1 0 ($\frac{1}{2} -\frac{1}{2}$)	7.57 $\cdot 10^{-2}$	4.61 $\cdot 10^{-2}$	11.2 $\cdot 10^{-2}$	7.56 $\cdot 10^{-2}$	39	0.1	-48
	1 2 ($\frac{3}{2} \frac{1}{2}$)	9.79 $\cdot 10^{-2}$	6.21 $\cdot 10^{-2}$	11.2 $\cdot 10^{-2}$	7.56 $\cdot 10^{-2}$	37	23	-14
	2 1 ($\frac{3}{2} -\frac{1}{2}$)	5.30 $\cdot 10^{-2}$	3.97 $\cdot 10^{-2}$	5.95 $\cdot 10^{-2}$	4.60 $\cdot 10^{-2}$	25	13	-11
	2 3 ($\frac{5}{2} \frac{1}{2}$)	6.41 $\cdot 10^{-2}$	5.00 $\cdot 10^{-2}$	5.95 $\cdot 10^{-2}$	4.60 $\cdot 10^{-2}$	22	27	7
	3 2 ($\frac{5}{2} -\frac{1}{2}$)	2.60 $\cdot 10^{-2}$	2.13 $\cdot 10^{-2}$	2.49 $\cdot 10^{-2}$	2.05 $\cdot 10^{-2}$	18	21	4

Table 1

Values obtained from the EPMS compared to the IPA. $S_{a,b}$ are given by Eq. (3.6) $I_{a,b}$ by Eq. (5.8) with $x_{\min}$ replaced by 0, for the two forms (3.9a and b) of the absorption factor. The value of z is 1.2, the values of $\cos \theta$ are 1.0 and 0.8. Note that for $\cos \theta = 1$ there is only one non-vanishing sum and integral.

To this observation should, however, be added the remark that the partial waves contributing to the sum (integral) now extend to $j \approx 10$, while the corresponding figure in the unmodified Born approximation is $j \approx 5$. This reduction in the relative importance of the lowest partial waves is the reason for the sharpening in the production angular distribution predicted by the absorption model as compared to the unmodified Born model.

Our results complement those of Bander and Shaw²⁰⁾, who showed that using an absorption factor affecting only the lowest partial waves and turning it off at $j = 3/2$ or $j = 5/2$ did change the cross-section but turning it off at $j = 5\frac{1}{2}$ or at $j = 10\frac{1}{2}$ had not much effect. They had, however, a much smaller ξ value than we have; also, when $\alpha > 0$ the second (negative) peak is less important (cf. below).

Now we shall undertake a more detailed discussion of the numerical results presented in Table 1. Let us first concentrate on the values of $(S_a - S_b)/S_a$, giving the relative change on the EPWS generated by a modification of the absorption factor; the corresponding quantity $(I_a - I_b)/I_a$ for the IPA, not listed in the table, takes roughly the same values. The change is enormous for $\alpha = 0$, $\beta = 1$, but as α and β increases, the change decreases, having its smallest value still around 20 %. Alluding to Fig. 3, we can understand this trend by observing²¹⁾ that with increasing α , the first (positive) maximum of $K_\alpha(\xi x)J_\alpha(\omega x)$ is shifted towards larger x values, its first (negative) minimum loosing in importance because of the damping from $K_\alpha(\xi x)$. Moreover, since $K_\alpha(\xi x)J_\alpha(\omega x)$ behaves as $\log 1/x$ for small x values when $\alpha = 0$, but as a constant when $\alpha > 0$, combined with the fact that $j_{\min}$ increases as λ increases, the very lowest partial waves, especially sensitive to the modification, become less important.

Since the derivation of the absorption model meets its greatest difficulties in the lowest partial waves the strong dependence of the results on the specific choice of the absorption factor is slightly discouraging. To see the effect on the production cross-section and the density matrix we have in Fig. 4, as an illustration, plotted $d\sigma/dt$ and some of the matrix elements for the reaction $\pi^+ p \rightarrow A_2^+ p$ with rho exchange at 4 GeV/c using the absorption factors (3.9) [curves marked (a) and (b)]. Observe that in this case $z = 1.27$, slightly bigger than the z value used above. As can be seen, the cross-section is very much changed, especially for large values of $\cos \theta$; on the other hand, the density matrix elements do not change very much.

In this context, let us also remark that if one uses, instead of Eq. (3.9), still another form for $k(j)$ given by Eq. (3.3) with

$$\exp[2i\delta(j)] = 1 - C \exp[-\gamma \cdot j(j+1)], \quad (3.12)$$

which perhaps is a more physical choice, one obtains a weaker absorption factor and consequently a still larger cross-section.

Next, we consider the value of $(S_a - I_b)/S_a$, which give the relative change from the EPWS to the IPA, if one identifies x with $j + \frac{1}{2}$ in the absorption factor. From Table 1, this value fluctuates but is never bigger in absolute magnitude than about 25%. Note the excellent agreement for $\alpha = 1$, $\beta = 0$, in which case Eq. (3.7) reads exactly $x = j + \frac{1}{2}$. This suggests that the IPA could be very much improved relative to the EPWS if the exact relation (3.7e) were used for the argument of the absorption factor.

Another point worth noticing concerns the complete omission in the IPA of the dependence on β . From Eq. (3.8) with $x_{\min}$ replaced by 0 (cf. Reference ¹⁹⁾), one might think that this omission is justified in the small angle approximation. However, in the case of absorption, the factor $k(j)$ modifies individually each partial wave, so the smallness of ϵ is now as relevant for the approximation as the smallness of ω [cf. Eq. (3.7a)]. We see this numerically from Table 1: while the value of, e.g., I_b is the same for $\alpha = 1$, $\beta = 0$ or 2, and also for $\alpha = 2$, $\beta = 1$ or 3, that of S_a changes, the difference increasing for increasing β . The corrections obtained by using more correct values for $x_{\min}$ as the lower limit of integration in Eq. (3.8) usually decrease the value of the integral, which only occasionally results in a shift in the right direction.

The influence of the IPA on the cross-section and the density matrix elements can be seen from Fig. 4, where the curves (c) are calculated using the IPA (3.8) with $x_{\min}$ replaced by 0 (for (c') $x_{\min}$ is replaced by $j_{\min}$) and with the absorption factor (3.9b). Not only is there now a change in the cross-section, but also the density matrix elements change [the density matrix elements for the curve (c') are practically the same as those marked (c)].

Finally, we have in Table 1 a column for $(S_a - I_a)/S_a$, which gives the relation between the EPWS and the IPA if one identifies x with j in the absorption factor. Note the improved agreement for increasing α and β . This is due to the fact that for the relevant values of j , i.e., the first few bigger than $j_{\min}$, the approximation of putting $x = j$ in the absorption factor becomes better for bigger α and β .

In conclusion, the various ways of treating the partial wave expansion in the absorption model differ considerably, the choice of an impact parameter approximation instead of an exact partial wave summation amounting to roughly the same differences as the various choices of the absorption factor. Since

from a computational point of view it is by no means more difficult to use the EPWS (in fact, using the recurrence formulae for the rotation coefficients¹⁸⁾, the EPWS is much faster on a computer than the IPA), we will in the comparison with experiment, to be undertaken in the next section, always use the EPWS.

Before we close this section, let us make a few remarks concerning some other features of the absorption model.

The effect of absorption, although remedying partially the points i) - iii) mentioned at the end of Section 2, does not help concerning the energy dependence $\sqrt{\text{point iv)}}^{18),22)}$ and does not necessarily ensure suppression below the unitary limit¹⁸⁾; thus, in the case of 2^+ meson production, some of the partial wave amplitudes with vector meson exchange will increase indefinitely with increasing energy, no matter which values of the absorption parameters $C_{i,f}$ and $A_{i,f}$ we choose. For pseudoscalar meson exchange, and with $C_i < 1$ as imposed by experiment, we have to choose $C_f = 1$ to ensure boundedness, which, though, does not necessarily force the amplitudes to respect the unitary limit.

However, in the range of energy which we shall consider here (3-8 GeV/c incident laboratory momentum), we are presumably still at such low energies, that we may forget about these asymptotic considerations and apply the absorption model as the simplest available model, which has had some success.

4. COMPARISON WITH EXPERIMENT

a) Production of $f_0(1253)$

As f_0 has G parity +1 and isospin zero, the only meson that can be exchanged in an OPE model is a pion. From the expression (A.10) for the width, applied to the decay of f_0 with mass 1253 MeV, we find for an experimental

width $\Gamma(f_0 \rightarrow \eta^+ \eta^-) = 60 \text{ MeV}^{-1}$ a coupling constant

$$\frac{g_a^2}{4\pi} = 1.6. \quad (4.1)$$

For the coupling of the charged pion at the nucleon vertex we use $(G^2/4\pi) = 27$.

On Fig. 5 are shown the calculated cross-sections as compared to the experimental results for the reactions $\pi^- p \rightarrow f_0 n$ at 4 GeV/c²³⁾ and $\pi^+ n \rightarrow f_0 p$ at 6 GeV/c²⁴⁾ incident laboratory momentum. The curves marked (a) are the result of the BTM, Eq. (2.4); it fails completely to reproduce the experimental data. The results of the absorption model, using Eqs. (3.1), (C.5) and (2.3), are marked (b). The values of the absorption parameters used in Eqs. (3.3)-(3.5)^{25), 26)} are $A_i = 7.6(\text{GeV}/c)^{-2}$, $C_i = 0.76$ and $C_f = 1$ at both energies, while at 4 GeV/c $A_f = 14(\text{GeV}/c)^{-2}$ and at 6 GeV/c $A_f = 9(\text{GeV}/c)^{-2}$; taking smaller values for A_f implies only a slight increase in the absolute value of the cross-section. Apart from the fact that the magnitude of the cross-section is too big by a factor of around 2 (observe the different scales used for the theoretical and the experimental results), the agreement is fair; in particular, one has a great improvement in the angular distribution²⁷⁾.

The density matrix elements are more difficult to compare to experiment. We can certainly not from our simple model explain the highly asymmetric decay distribution of the pions in the rest frame of f_0 at 4 GeV/c²³⁾. The experiment at 6 GeV/c, however, indicates that $\rho_{0,0} \sim 0.6$ and $\rho_{1,1} \sim 0.2$ when averaged over all production angles²⁴⁾. Our theoretical results are shown in Fig. 6; the fit to the (uncertain) data is perhaps not good but in any case better than the BTM²⁷⁾.

b) Production of $A_2(1310)$

Since the A_2 meson has G parity -1 , the only candidate among the presently well-established bosons of spin less than 2 for the exchanged meson in a OPE model is the rho meson. From the experimentally determined width of 80 MeV ¹⁾ for the decay $A_2 \rightarrow \rho \pi$, we find for an A_2 mass of 1310 MeV and a rho mass of 763 MeV the coupling constant from the expression (A.11) for the width:

$$\frac{f_\rho^2}{4\pi} = 190. \quad (4.2)$$

For the coupling constants of the rho meson to the nucleon, we use $G_V^2/4\pi = 1.2$, $G_T/G_V = 3.7$ ⁵⁾.

We first compare the results of our calculation for the cross-section of the reaction $\pi^+ p \rightarrow A_2^+ p$ to the experimental one at 4 GeV/c incident momentum ²⁸⁾ in Fig. 7. The curve marked (a) is the result in the unmodified Born approximation, as given by Eq. (2.5); it is much too big and does not show any decrease with increasing momentum transfer, the reason for this being the high powers of t in the numerator of Eq. (2.5) and also the fact that the rho meson mass is fairly big.

The absorption model result from Eqs. (3.1), (C.6) and (2.3), using $A_i = A_f = 7.6(\text{GeV}/c)^{-2}$, $C_i = 0.76$ and $C_f = 1.0$ as values for the absorption parameters ^{25), 26)}, is marked (b). Even if considerably reduced, as compared to the unmodified Born approximation, this curve also fails to reproduce the experimental data. By increasing A_f the theoretical curve decreases in absolute magnitude and becomes slightly more peaked, but cannot give a good fit for any reasonable value of A_f .

In looking for other effects that could improve the theoretical results, we tried to include a form factor into the absorption model calculations²⁹⁾; we choose the form

$$F(t) = \frac{\Lambda^2 - m_e^2}{\Lambda^2 - t}, \quad (4.3)$$

which is normalized to unity at $t = m_e^2$. Denoting by $B_{\{\lambda\}}(m_e)$ the value of a helicity Born amplitude without form factor, the inclusion of the form factor (4.3) into this amplitude before one introduces the absorption implies that the amplitude now reads $B_{\{\lambda\}}(m_e) = B_{\{\lambda\}}(\Lambda)$, a form which then remains valid even with absorption.

Keeping the absorption parameters at the values given above, we varied Λ and found rough agreement with the experimental slope only for rather small values of Λ ; the results for $\Lambda = 400$ MeV is plotted as curve (c) in Fig. 7, which also shows the same curve normalized to the experimental points at $-t \sim 0.2(\text{GeV}/c)^2$ [curve marked (c')] and the curve (b) multiplied by $F^2(t)$ for $\Lambda = 400$ MeV [curve marked (b')]. A comparison between (c) and (b') shows that the value of the cross-section is about the same, either the form factor is introduced before or after the absorption. For a higher value of A_F the theoretical curve is more peaked, but it requires a value $A_F \sim 5A_i$ to reproduce the experimental slope for $\Lambda = 400$ MeV.

Concerning the other possible test of the theoretical results, namely the distribution (B.12) in the decay of the A_2 meson as expressed in terms of its spin space density matrix elements there is not much data available, partly due to difficulties in treating the interference between the two reactions $\pi N \rightarrow \rho N^*$ and $\pi N \rightarrow A_2 N$. However, from the data at 2.75 GeV/c incident laboratory momentum³⁰⁾ we conclude that $\rho_{2,2} \approx 0$ and $\rho_{1,1} \approx 0.5$, averaged over all momentum transfer. Our corresponding theoretical values (including a form factor) are $\rho_{2,2} \approx 0.05$, $\rho_{1,1} \approx 0.4$.

To see the modifications on the decay distribution caused by introducing absorption alone or together with a form factor, we have in Fig. 8 plotted our theoretical results for some of the density matrix elements for A_2 in the same reaction as above, using the same parameters as those used in fitting the cross-section. As has been noted before ^{14),20)}, the effect of a form factor (4.3) is to shift the values of the matrix elements back towards their values in the unmodified Born approximation.

As to the energy dependence it again turns out to be completely wrong in this case of vector meson exchange even with a form factor included. While our theoretical model predicts a cross-section about 10 times as big as the experimental one at 4 GeV/c, the corresponding ratio is 43 at 8 GeV/c ($\sigma_{\text{theor}} \approx 8 \text{ mb}$, $\sigma_{\text{exp}} \approx 190 \mu\text{b}$ ³¹⁾). Also the predicted slope is not steep enough at this energy.

c) Production of $K^{**}(1400)$

Since there is no restriction from G parity conservation on the exchanged particle in the reaction $KN \rightarrow K^{**}N$, implying that both a pion and a rho meson, if isospin permits also an omega meson, could be exchanged and since the data available are relatively meager, we think it is unfair to try to fit our model to the experimental results. The greater number of free parameters could hide the bad features of the model, already found above in those cases where only one meson could be exchanged.

5. SUMMARY AND CONCLUSIONS

The results of Section 3, collected in Table 1 and Figs. 2-4, are that the impact parameter approximation, as it has previously been used in the application of the absorption model, changes both the predicted cross-section and the decay distribution significantly as compared to the exact partial wave summation. Furthermore, to use slightly different absorption factors has large effects on the cross-section but not on the decay distribution. These results are due to the fact that the absorption model, although introducing a large damping of the low partial waves as compared to the unmodified Born approximation, still has a major contribution from these waves.

The comparison with experiment of the predictions of the absorption model for the production of spin parity 2^+ mesons in pion-nucleon interactions has not been very successful. Not only is the absolute magnitude of the cross-sections too big, but in the case of vector meson exchange the slope of the cross-section is also wrong, a feature which can be repaired only by introducing a rather rapidly changing form factor. The energy dependence in case of vector meson exchange is again completely wrong. The predicted decay distribution have to wait for more experiment.

Putting this together with previous results ³²⁾ we are thus led to the following conclusion: although very good in reproducing the experimental data for those reactions where a pion is exchanged and where the spins of the outgoing particles are low (an excellent example is the reaction $\pi p \rightarrow \rho p$ ¹⁴⁾), the absorption model becomes worse as soon as the spins increase, either of the exchanged particle or of the particles in the final state. The model, although remedying partly the diseases of the unmodified Born approximation, has thus itself a limited scope. We still lack an applicable theoretical model for high energy two-body resonance production, which is able to account for the experimental results over a wider range of energy and of spin.

A P P E N D I X A

HELICITY STATES, COUPLINGS AND DECAY WIDTHS FOR A SPIN PARITY 2^+ PARTICLE

a) Helicity States for a Spin 2 Particle

In general, the wave function for a particle of spin $2^{12)}$, independent of its parity, is a symmetric and traceless second order tensor $\varphi = (\varphi_{\rho\sigma})$ which in momentum space fulfils the condition

$$p_{\rho} \varphi_{\rho\sigma} = 0, \quad (A.1)$$

where p_{ρ} is the four-momentum. These conditions reduce the number of independent tensor components to five, as it should be for a particle of spin 2.

According to the definition of a helicity state ¹⁶⁾, the wave function $\psi(s, \lambda; \bar{p})$ for a particle of spin s , helicity λ and momentum p , in a co-ordinate system with z axis parallel to $\bar{p}$, is given by ³³⁾

$$\psi(s, \lambda; \bar{p}) = Z(v) \psi(s, \lambda; 0) \quad (A.2)$$

where $Z(v)$ is an ordinary Lorentz transformation along the z axis with the velocity $v = (|\bar{p}|/E)$, $E = \sqrt{p^2 + m^2}$, m being the mass of the particle, and where $\psi(s, \lambda; 0)$ is the rest system wave function with spin component λ along the z axis.

The helicity states for spin 2³⁴⁾ may be built up from the states for spin 1 (see below). Let us therefore consider the case $s = 1$ first. Here $\psi(s = 1, \lambda; 0)$ is a four-vector given by³⁵⁾

$$\begin{aligned}\psi(1, \lambda; 0) &= \mp \frac{1}{\sqrt{2}} (1, \pm i, 0, 0) \quad \text{for } \lambda = \pm 1, \\ &= (0, 0, 1, 0) \quad \text{for } \lambda = 0,\end{aligned}\tag{A.3}$$

which, using (A.2), gives

$$\begin{aligned}\psi(1, \lambda; \bar{p}) &\equiv \varepsilon^{(\lambda)}(\bar{p}) = \mp \frac{1}{\sqrt{2}} (1, \pm i, 0, 0) \quad \text{for } \lambda = \pm 1 \\ &= (0, 0, \frac{E}{m}, \frac{|\bar{p}|}{m}) \quad \text{for } \lambda = 0\end{aligned}\tag{A.4}$$

The rest frame wave function for $s = 2$ is now obtained from the $s = 1$ states using Clebsch-Gordan coefficients

$$\begin{aligned}\psi(s=2, \lambda; 0) &= \left(\varphi_{\rho\sigma}^{(\lambda)}(\bar{p}=0) \right) = \\ &= \sum_{\lambda', \lambda''} \psi(1, \lambda'; 0) \psi(1, \lambda''; 0) \langle 1, \lambda'; 1, \lambda'' | 2, \lambda \rangle\end{aligned}\tag{A.5}$$

which from (A.2) means

$$\varphi_{\rho\sigma}^{(\lambda)}(\vec{p}) = \sum_{\lambda', \lambda''} \varepsilon_{\rho}^{(\lambda')}(\vec{p}) \varepsilon_{\sigma}^{(\lambda'')}(\vec{p}) \langle 1, \lambda'; 1, \lambda'' | 2, \lambda \rangle \quad (\text{A.6})$$

or explicitly ³⁴⁾

$$\begin{aligned} \varphi_{\rho\sigma}^{(\pm 2)} &= \varepsilon_{\rho}^{(\pm 1)} \varepsilon_{\sigma}^{(\pm 1)} \\ \varphi_{\rho\sigma}^{(\pm 1)} &= \frac{1}{\sqrt{2}} \left[\varepsilon_{\rho}^{(\pm 1)} \varepsilon_{\sigma}^{(0)} + \varepsilon_{\rho}^{(0)} \varepsilon_{\sigma}^{(\pm 1)} \right] \\ \varphi_{\rho\sigma}^{(0)} &= \frac{1}{\sqrt{6}} \left[\varepsilon_{\rho}^{(1)} \varepsilon_{\sigma}^{(-1)} + \varepsilon_{\rho}^{(-1)} \varepsilon_{\sigma}^{(1)} + 2 \varepsilon_{\rho}^{(0)} \varepsilon_{\sigma}^{(0)} \right] \end{aligned} \quad (\text{A.7})$$

It is easily verified that these helicity states for a spin 2 particle fulfil the general conditions on the wave function stated above.

b) Couplings and Decay Widths

For the most general couplings of interest to us, we write in the notation of Fig. 1,

$$0^- 0^- 2^+ \quad \frac{g_2}{m_c} (a-e)_{\rho} \varphi_{\rho\sigma} (a-e)_{\sigma} = \frac{4g_2}{m_c} a_{\rho} \varphi_{\rho\sigma} a_{\sigma} \quad (\text{A.8})$$

$$0^- 1^- 2^+ \quad i \frac{f_2}{m_c} \varepsilon_{\mu\nu} \varepsilon_{\mu\rho\tau\omega} \varphi_{\rho\sigma} a_{\sigma} a_{\tau} c_{\omega} \quad (\text{A.9})$$

Here g_2 and f_2 are dimensionless coupling constants and $\epsilon_{\alpha\beta\gamma\omega}$ is the completely antisymmetric fourth order tensor with non-vanishing components ± 1 . For the couplings at the nucleon vertex we use the notations and conventions of Ref. 5).

Using Eq. (A.8) we calculate the width Γ_0 for the decay $2^+ \rightarrow 0^- 0^-$ to be

$$\Gamma_0 = \frac{g_2^2}{4\pi} \cdot \frac{16}{15} \cdot \frac{P^5}{m_c^4} \quad (\text{A.10})$$

where P is the magnitude of the three-momentum of either decay particle in the rest frame of the 2^+ meson.

In the same way, the width Γ_1 for the decay $2^+ \rightarrow 1^- 0^-$ is

$$\Gamma_1 = \frac{f_2^2}{4\pi} \cdot \frac{1}{10} \cdot \frac{Q^5}{m_c^4} \quad (\text{A.11})$$

where Q is the magnitude of the three-momentum for the vector meson in the 2^+ rest frame.

A P P E N D I X B

THE ANGULAR DISTRIBUTION IN THE DECAY OF A 2^+ PARTICLE

As has become general use ³⁶⁾, we choose in the rest frame of the 2^+ particle a right-hand co-ordinate system, having its z axis along the beam direction in this frame and with its x axis in the production plane. This system we call the "canonical system". The z axis in that system is chosen as the axis of quantization for the spin 2 particle; its spin space density matrix elements in the canonical system is denoted by $\rho_{\ell,m}$, $\ell, m = 0, \pm 1, \pm 2$. This is a Hermitian matrix, which fulfils the symmetry relation

$$\rho_{-\ell,-m} = (-1)^{\ell-m} \rho_{\ell,m} \quad (\text{B.1})$$

due to parity conservation in the production process.

a) The Decay $2^+ \rightarrow 0^- 0^-$

Let the momentum of one of the decay particles be specified by the polar angles ϑ , φ in the canonical system defined above. The distribution in these angles is then (η is a normalization constant, independent of ϑ and φ) ¹³⁾

$$\begin{aligned} W(\vartheta, \varphi) = & \eta \left[\sin^4 \vartheta \cdot \{ \rho_{2,2} + \rho_{2,-2} \cdot \cos 4\varphi \} + \right. \\ & + \sin^2 \vartheta \cdot \{ \rho_{1,1} - \rho_{1,-1} \cdot \cos 2\varphi \} + 3 \rho_{0,0} \cdot (\cos^2 \vartheta - 1/3)^2 - \\ & - 4 \sin^2 \vartheta \cdot \cos \vartheta \cdot \{ \operatorname{Re} \rho_{2,1} \cdot \cos \varphi - \operatorname{Re} \rho_{2,-1} \cdot \cos 3\varphi \} + \\ & + 2\sqrt{6} \cdot \operatorname{Re} \rho_{2,0} \cdot \sin^2 \vartheta \cdot (\cos^2 \vartheta - 1/3) \cdot \cos 2\varphi - \\ & \left. - 2\sqrt{6} \cdot \operatorname{Re} \rho_{1,0} \cdot \sin 2\vartheta \cdot (\cos^2 \vartheta - 1/3) \cdot \cos \varphi. \right] \quad (\text{B.2}) \end{aligned}$$

This distribution determines all but the imaginary part of the spin density matrix (observe that with our choice of co-ordinate system, $\rho_{2,-2}$ and $\rho_{1,-1}$ are real). In the absorption model, as well as in the BTM, all $\rho_{\ell,m}$ are predicted to be real.

b) The Decay $2^+ \rightarrow 1^- 0^-$ Followed by $1^- \rightarrow 0^- 0^-$

There exist already in the literature several treatments of the decay of a boson into three pseudoscalar mesons³⁷⁻⁴⁰⁾, with or without an intermediate state containing a vector meson. However, the specific reaction of interest to us is often, in these treatments, hidden in the generalities of the approach and we give here a, as we hope, reasonably self-contained derivation, especially suited for the particular case under study.

The three-body decay (cf. Fig. 9 for the notation)

$$\begin{array}{l} c \longrightarrow E + V \\ \quad \quad \quad \searrow \quad \quad \quad \nearrow \\ \quad \quad \quad F + H \end{array} \quad (B.3)$$

is in general described by five independent kinematical variables, which we conveniently choose as the two energy variables¹¹⁾ $s_1 = -(F+H)^2$ and $s_2 = -(E+H)^2$, i.e., two squared masses defining the Dalitz plot, and three angles, defining completely the orientation of the decay plane in the rest frame of particle c . As these angles we choose the polar angles $\Omega_N = (\vartheta_N, \varphi_N)$ of the normal $\vec{N} = \vec{E} \times \vec{F}$ to the decay plane and an angle ψ defining the direction of the three-momentum $\vec{E}$ of particle E in the decay plane, all angles being referred to the canonical system defined above.

We shall here only be interested in the distribution of the normal $\bar{N}$ to the decay plane ⁴¹⁾. Thus we must perform integrations over the energy variables s_1 and s_2 , i.e., over suitable regions of the Dalitz plot, and over the angle ψ . The derivation below proceeds, then, in several steps: we start by writing down the decay matrix element for the decay (B.3), at first omitting complications from the fact that two of the final particles in (B.3) might be identical. We then give the decay distribution in a co-ordinate system chosen such that the integration over the angle ψ is trivial, and transform afterwards from that system to the canonical one. Finally, we discuss the integration over the variables s_1 and s_2 and in this context also the influence of identical particles.

The decay matrix element for the decay (B.3) is obtained from the coupling (A.9) by replacing the vector meson polarization vector ε_{μ} by

$$\frac{\delta_{\mu\lambda} + m_v^{-2} \cdot (E+F)_{\mu} (E+F)_{\lambda}}{s_1 - M_v^2} \cdot g g_1 \cdot F_{\lambda} , \quad (\text{B.4})$$

where we have used the decay matrix element

$$g_1 \varepsilon_{\lambda} (F-H)_{\lambda} = g g_1 \cdot \varepsilon_{\lambda} F_{\lambda} \quad (\text{B.5})$$

for the decay of the vector meson. The mass M_v in the propagation denominator of Eq. (B.4) is the complex mass of the vector meson which in terms of its real mass m_v and width Γ_v is given by

$$M_v = m_v - i \frac{1}{2} \Gamma_v . \quad (\text{B.6})$$

For $m_V \gg \Gamma_V$, insertion of (B.4) into (A.9) gives the matrix element $M^{(k)}$ for the decay in rest of a spin parity 2^+ particle, having spin component k along the axis of quantization, in the form

$$M^{(k)} \propto \varphi_{ns}^{(k)} \tilde{E}_s N_n, \quad (\text{B.7a})$$

$$\tilde{E}_s = E_s (s_1 - m_V^2 + i m_V \Gamma_V)^{-1}, \quad (\text{B.7b})$$

where we have suppressed uninteresting constant factors and where as usual Latin indices imply summation over the space components only. In terms of these decay matrix elements, the decay distribution reads

$$W(\Omega_N, \psi; s_1, s_2) = \mathcal{N} \cdot \int_{k,l} M^{(k)} M^{(l)*}, \quad (\text{B.8})$$

where $\mathcal{N}$ is a normalization constant, independent of the five variables specifying the decay.

In writing down the distribution (B.8) we have made no assumption concerning the choice of the co-ordinate system in the spin 2 particle rest frame; with a proper definition of the density matrix elements and the angles, it is therefore valid in any such system. In order to perform the integration over the angle ψ , it is particularly convenient to consider the distribution referred to the co-ordinate system $\Sigma' = (x', y', z')$ of Fig. 10. Denoting all quantities referred to this system by a prime, we have the following connection between the density matrix elements in Σ' and in the canonical system

$$P'_{k', l'} = D^{(2)}_{k, k'}(\varphi_N, \psi_N, 0) P_{k, l} D^{(2)}_{l, l'}(\varphi_N, \psi_N, 0) \quad (\text{B.9a})$$

$$D^{(2)}_{l, l'}(\varphi, \psi, 0) = e^{-i l \cdot \varphi} \cdot d^{(2)}_{l, l'}(\psi) \quad (\text{B.9b})$$

where $d^{(2)}_{l, l'}(\psi)$ are the rotation coefficients (cf. Section 3).

The reason to consider the system Σ' becomes evident when we write down the matrix elements (B.7) in this system: using the states (A.7), we find

$$M^{(k')} \begin{cases} \propto e^{i k' \psi} & \text{for } k' = \pm 1 \\ = 0 & \text{otherwise} \end{cases} \quad (\text{B.10})$$

Thus, the decay distribution, after integration over the angle ψ , reads in the system Σ'

$$W(\Omega_N; s_1 s_2) = \int d\psi W(\Omega_N, \psi; s_1 s_2) \propto P'_{1,1} + P'_{-1,-1} \quad (\text{B.11})$$

We have now only to use Eq. (B.9) to transform back to the canonical system. The distribution reads explicitly ⁴²⁾

$$\begin{aligned}
W(\Omega_N; s_1, s_2) = \mathcal{H}' \cdot [& p_{3,2} \cdot \sin^2 \vartheta_N \cdot (1 + \cos^2 \vartheta_N) + \\
& + p_{1,1} \cdot \{1 + \cos^2 \vartheta_N (1 - 4 \sin^2 \vartheta_N)\} + 3 p_{0,0} \cdot \sin^2 \vartheta_N \cdot \cos^2 \vartheta_N - \\
& - p_{3,-2} \cdot \sin^4 \vartheta_N \cdot \cos 4 \varphi_N - p_{1,-1} \cdot \sin^2 \vartheta_N \cdot (1 - 4 \cos^2 \vartheta_N) \cdot \cos 2 \varphi_N - \\
& - 4 \operatorname{Re} p_{2,1} \cdot \cos^3 \vartheta_N \cdot \sin \vartheta_N \cdot \cos \varphi_N - \\
& - 4 \operatorname{Re} p_{2,-1} \cdot \sin^3 \vartheta_N \cdot \cos \vartheta_N \cdot \cos 3 \varphi_N - \\
& - 2 \sqrt{6} \cdot \operatorname{Re} p_{2,0} \cdot \cos^2 \vartheta_N \cdot \sin^2 \vartheta_N \cdot \cos 2 \varphi_N - \\
& - \sqrt{6} \cdot \operatorname{Re} p_{1,0} \cdot \sin 2 \vartheta_N \cdot (1 - 2 \cos^2 \vartheta_N) \cdot \cos \varphi_N.
\end{aligned} \tag{B.12}$$

The proportionality factor $\mathcal{H}'$ is independent of the angles ϑ_N and φ_N , but depends on s_1 and s_2 (and the production angle Θ for the 2^+ particle).

The distribution (B.12) is the angular distribution of the normal to the decay plane for that part of the Dalitz plot which corresponds to the values s_1 and s_2 . Integration over s_1 and s_2 , which will almost always be necessary in order to improve statistics, will only affect the value of the proportionality factor $\mathcal{H}'$, to the extent that all interference effects arising from the identity of the decay particles can be neglected. Thus, with this assumption, all the events in the Dalitz plot have the same weight in the decay angular distribution.

What happens then, if two of the three particles E, F and H of the decay (B.3) are identical ⁴⁰⁾? This is the case, e.g., in the decay $A_2^+ \rightarrow \pi^+ \rho^0$, $\rho^0 \rightarrow \pi^+ \pi^-$. To be definite, let us suppose, that particles E and F are identical. The decay matrix element $M^{(k)} \equiv M^{(k)}(E)$ of Eq. (B.7a) has then to be symmetrized with respect to the interchange of E and F, i.e., we have to make the replacement

$$\begin{aligned}
M^{(k)}(E) &\rightarrow \\
\rightarrow M^{(k)}(E, F) = M^{(k)}(E) + M^{(k)}(F) &\propto \varphi_{\pi s}^{(k)} (\tilde{E}_s - \tilde{F}_s) N_{\pi}, \quad (B.13a)
\end{aligned}$$

$$\tilde{F}_G = F_G \cdot (s_2 - m_v^2 + i m_v \Gamma_v)^{-1} \quad (B.13b)$$

Remember here that $\bar{N}$ is defined as $\bar{E} \times \bar{F}$. The decay distribution replacing (B.8) then reads

$$\begin{aligned}
W_E &\equiv W(\Omega_N, \psi; s_1, s_2) \rightarrow \\
\rightarrow W_{E,F}(\Omega_N, \psi; s_1, s_2) &= \Re \cdot \rho_{k,l} M^{(k)}(E, F) M^{(l)*}(E, F) = \\
= \Re \rho_{k,l} [M^{(k)}(E) M^{(l)*}(E) + M^{(k)}(F) M^{(l)*}(F) - & \quad (B.14) \\
- \{M^{(k)}(E) M^{(l)*}(F) + M^{(k)}(F) M^{(l)*}(E)\}] = \\
= W_E + W_F + W_{int}.
\end{aligned}$$

After integration over the angle ψ , the two distributions W_E and W_F will both reduce to the form (B.12), the only difference being an interchange of s_1 and s_2 in $\Re$. This is due to the symmetry under the interchange of E and F . Moreover, integration in the energy variables s_1 and s_2 over the vector meson bands in the Dalitz plot outside the region of interference will give the two distributions the same proportionality factor, i.e., all the events outside that region have the same weight in the decay angular distribution. The events inside the interference region, however, have a different weight and have to be discarded in the experimental analysis.

We conclude that in the experimental study of the angular distribution of the normal to the decay plane in the decay (B.3), in which two of the three decay particles are identical, all events outside the interference region in the Dalitz plot may be used; the distribution takes the form (B.12)⁴¹⁾. We also remark that the restriction of the events to certain regions of the Dalitz plot does not introduce any bias in the angular distribution simply because the Dalitz plot variables and the angles are independent. This is in contrast to a three-body production process in which two resonances can be formed in the final state (e.g., $K^+p \rightarrow K^0 \pi^+p$); in this case the Dalitz plot variables and the decay angles are not independent.

A P P E N D I X C

HELICITY AMPLITUDES IN THE BORN APPROXIMATION

The helicity matrix elements, calculated in the Born approximation, are denoted by $\langle \lambda_c \lambda_d | B | \lambda_b \rangle$, where λ_b is the helicity of particle b , and so on, in the notation of Fig. 1. They fulfil the relation

$$\langle -\lambda_c, -\lambda_d | B | -\lambda_b \rangle = (-)^{\lambda_c - \lambda_d + \lambda_b + 1} \langle \lambda_c \lambda_d | B | \lambda_b \rangle \quad (C.1)$$

due to parity conservation.

We introduce the following notation besides that already used in Section 2

$$\xi(\pm 1) \equiv \xi_{\pm} = \sqrt{(E_b + m_b)(E_d + m_d)} \left[\frac{q}{E_b + m_b} \pm \frac{q'}{E_d + m_d} \right], \quad (C.2a)$$

$$\eta(\pm 1) \equiv \eta_{\pm} = \sqrt{(E_b + m_b)(E_d + m_d)} \left[1 \pm \frac{q q'}{(E_b + m_b)(E_d + m_d)} \right], \quad (C.2b)$$

$$v_a = \frac{q}{E_a}, \quad v_c = \frac{q'}{E_c}, \quad \alpha = v_c / v_a \quad (C.2c)$$

$$\kappa = \frac{\sqrt{s} v_c}{m_b + m_c} \quad (C.2d)$$

where E_a is the c.m.s. energy of particle a , and so on. By using helicity spinors and the helicity states (A.7), we find using the amplitudes given in Section 2:

for pseudoscalar meson exchange

$$\begin{aligned} \langle 0, \lambda_d | B^{(PS)} | \lambda_b \rangle &= \\ &= \frac{4Gg_2}{\sqrt{6}m_c} \cdot \frac{q^2}{m_e^2 - t} \cdot \left\{ (-4\lambda_b\lambda_d) \cdot (-)^{\frac{1}{2}\lambda_b} \cdot d_{\lambda_b, \lambda_d}^{1/2}(\theta) \cdot \left\{ \sin^2\theta - \frac{2E_c^2}{m_c^2}(\cos\theta - \alpha) \right\} \right\} \quad (C.3a) \end{aligned}$$

$$\begin{aligned} \langle 1, \lambda_d | B^{(PS)} | \lambda_b \rangle &= \\ &= -\frac{4Gg_2}{m_c} \cdot \frac{q^2}{m_e^2 - t} \cdot \frac{E_c}{m_c} \cdot \left\{ (-4\lambda_b\lambda_d) \cdot (-)^{\frac{1}{2}\lambda_b} \cdot d_{\lambda_b, \lambda_d}^{1/2}(\theta) \cdot \sin\theta \cdot (\cos\theta - \alpha) \right\} \quad (C.3b) \end{aligned}$$

$$\begin{aligned} \langle 2, \lambda_d | B^{(PS)} | \lambda_b \rangle &= \quad (C.3c) \\ &= -\frac{2Gg_2}{m_c} \cdot \frac{q^2}{m_e^2 - t} \cdot \left\{ (-4\lambda_b\lambda_d) \cdot (-)^{\frac{1}{2}\lambda_b} \cdot d_{\lambda_b, \lambda_d}^{1/2}(\theta) \cdot \sin^2\theta \right\} \end{aligned}$$

for vector meson exchange

$$\begin{aligned} \langle \lambda_c \lambda_d | B^{(V)} | \lambda_b \rangle &= \\ &= (G_V + G_T) \cdot \frac{f_2}{m_c^2} \cdot \frac{q^2 E_c}{2(m_e^2 - t)} \cdot (-)^{\lambda_d + \lambda_b} \cdot \\ &\quad \cdot \left\{ \langle \lambda_c \lambda_d | B_1 | \lambda_b \rangle + \gamma \cdot n \cdot \langle \lambda_c \lambda_d | B_2 | \lambda_b \rangle \right\} \quad (C.4a) \end{aligned}$$

$$\begin{aligned} \langle 2, \lambda_d | B_1 | \lambda_b \rangle &= \sin\theta \cdot \left[\delta_{\lambda_d, 1/2} \cdot 4\lambda_b \cdot \left\{ (2\lambda_b 2\lambda_d) \cdot d_{-\lambda_d, \lambda_b}^{1/2}(\theta) \cdot (\cos\theta - \alpha) \right\} + \right. \\ &\quad \left. + \left\{ 2\lambda_b 2\lambda_d \cdot \left\{ (2\lambda_b 2\lambda_d) + v_c \cdot \gamma(2\lambda_b 2\lambda_d) \right\} \cdot d_{\lambda_d, \lambda_b}^{1/2}(\theta) \sin\theta \right\} \right] \quad (C.4b) \end{aligned}$$

$$\langle 1, \lambda_d | B_1 | \lambda_b \rangle =$$

$$= \frac{E_c}{m_c} \cdot \left[\delta_{\lambda_d, 1/2} \cdot 4\lambda_b \cdot \left\{ (2\lambda_b 2\lambda_d) \cdot d_{-\lambda_d, \lambda_b}^{1/2}(\theta) \cdot (\cos\theta - \alpha)^2 + \right. \right. \\ \left. \left. + \left\{ 2\lambda_b 2\lambda_d \cdot \left\{ (2\lambda_b 2\lambda_d) + v_c \cdot \eta(2\lambda_b 2\lambda_d) \right\} \cdot d_{\lambda_d, \lambda_b}^{1/2}(\theta) \cdot \sin\theta \cdot (\cos\theta - \alpha) - \right. \right. \right. \\ \left. \left. \left. - 2\lambda_b 2\lambda_d \cdot \left\{ (2\lambda_b 2\lambda_d) \cdot \frac{m_c^2}{E_c^2} \cdot d_{-\lambda_d, \lambda_b}^{1/2}(\theta) \cdot \sin^2\theta \right\} \right] \right. \quad (C.4c)$$

$$\langle 0, \lambda_d | B_1 | \lambda_b \rangle =$$

$$= -\sqrt{6} \cdot 2\lambda_b 2\lambda_d \cdot \left\{ (2\lambda_b 2\lambda_d) \cdot d_{-\lambda_d, \lambda_b}^{1/2}(\theta) \cdot \sin\theta \cdot (\cos\theta - \alpha) \right. \quad (C.4d)$$

$$\langle 2, \lambda_d | B_2 | \lambda_b \rangle = -\eta(-2\lambda_b 2\lambda_d) \cdot d_{\lambda_d, \lambda_b}^{1/2}(\theta) \cdot \sin^2\theta$$

(C.4e)

$$\langle 1, \lambda_d | B_2 | \lambda_b \rangle =$$

$$= -\eta(-2\lambda_b 2\lambda_d) \cdot d_{\lambda_d, \lambda_b}^{1/2}(\theta) \cdot \sin\theta \cdot (\cos\theta - \alpha) \quad (C.4f)$$

$$\langle 0, \lambda_d | B_2 | \lambda_b \rangle = 0$$

(C.4g)

For the application of the absorption model, the helicity amplitudes are written in the form (3.1). We list here the coefficients $\langle \lambda_c \lambda_d | A | \lambda_b \rangle$ and $\langle \lambda_c \lambda_d | X^j | \lambda_b \rangle$ obtained from Eqs. (C.2) and (C.3) respectively.

For pseudoscalar meson exchange we write

$$\langle \lambda_c \lambda_d | B^{(PS)} | \lambda_b \rangle = \frac{G g_2}{m_c} \cdot \frac{q}{q'} \cdot \langle \lambda_c \lambda_d | B^{(PS)} | \lambda_b \rangle \quad (C.5a)$$

The coefficients $\langle \lambda_c \lambda_d | A | \lambda_b \rangle$ and $\langle \lambda_c \lambda_d | X^j | \lambda_b \rangle$ below are then obtained by writing $\langle \lambda_c \lambda_d | B^{(PS)} | \lambda_b \rangle$ in the form (3.1):

$$\langle 0, \frac{1}{2} | A | \pm \frac{1}{2} \rangle = \frac{1}{2\sqrt{6}} \cdot \xi_{\mp} \cdot \left\{ 1 - z^2 - 2 \frac{E_c^2}{m_c^2} (\alpha - z)^2 \right\} \quad (C.5b)$$

$$\langle 0, \frac{1}{2} | X^{\frac{1}{2}} | \pm \frac{1}{2} \rangle = \frac{1}{2\sqrt{6}} \cdot \xi_{\mp} \cdot \left\{ (1 + 2 \frac{E_c^2}{m_c^2})(z \pm \frac{1}{3}) - 4\alpha \cdot \frac{E_c^2}{m_c^2} \right\} \quad (C.5c)$$

$$\langle 0, \frac{1}{2} | X^{\frac{3}{2}} | \pm \frac{1}{2} \rangle = \frac{1}{3\sqrt{6}} \cdot \xi_{\mp} \cdot \left(1 + 2 \frac{E_c^2}{m_c^2} \right) \quad (C.5d)$$

$$\langle 1, \frac{1}{2} | A | \pm \frac{1}{2} \rangle = \frac{1}{2} \cdot \frac{E_c}{m_c} \cdot \xi_{\mp} \cdot (1 \pm z) \cdot (\alpha - z) \quad (C.5e)$$

$$\langle 1, \frac{1}{2} | \bar{X}^{\frac{1}{2}} | \pm \frac{1}{2} \rangle = \pm \frac{1}{9} \cdot \frac{E_c}{m_c} \cdot \left\{ \mp \cdot (2 \pm \frac{2}{3} - \alpha) \right\}, \quad (C.5f)$$

$$\langle 1, \frac{1}{2} | \bar{X}^{\frac{3}{2}} | \pm \frac{1}{2} \rangle = \pm \frac{1}{3} \cdot \frac{E_c}{m_c} \cdot \left\{ \mp \right\} \quad (C.5g)$$

$$\langle 1, -\frac{1}{2} | \bar{H} | \pm \frac{1}{2} \rangle = \pm \frac{E_c}{m_c} \cdot \left\{ \pm \cdot (\alpha - 2) \right\} \quad (C.5h)$$

$$\langle 1, -\frac{1}{2} | \bar{X}^{\frac{3}{2}} | \pm \frac{1}{2} \rangle = \pm \frac{1}{\sqrt{3}} \cdot \frac{E_c}{m_c} \cdot \left\{ \pm \right\} \quad (C.5i)$$

$$\langle 2, \frac{1}{2} | \bar{H} | \pm \frac{1}{2} \rangle = -\frac{1}{2} \cdot \left\{ \mp \cdot (1 \pm 2) \right\} \quad (C.5j)$$

$$\langle 2, \frac{1}{2} | \bar{X}^{\frac{3}{2}} | \pm \frac{1}{2} \rangle = \frac{1}{2\sqrt{3}} \cdot \left\{ \mp \right\} \quad (C.5k)$$

$$\langle 2, -\frac{1}{2} | \bar{H} | \pm \frac{1}{2} \rangle = \mp \cdot \left\{ \pm \right\} \quad (C.5l)$$

All $\langle \lambda_c \lambda_d | X^j | \lambda_b \rangle$ not listed vanish.

For vector meson exchange we write

$$\begin{aligned} \langle \lambda_c \lambda_d | B^{(V)} | \lambda_b \rangle &= \\ &= \frac{f_2}{m_c} \cdot (G_V + G_T) \cdot \frac{q E_c}{g q'} \cdot (-)^{\lambda_b + \lambda_d} \langle \lambda_c \lambda_d | B^{(V)} | \lambda_b \rangle \end{aligned} \quad (C.6a)$$

The coefficients $\langle \lambda_c \lambda_d | A | \lambda_b \rangle$ and $\langle \lambda_c \lambda_d | X^j | \lambda_b \rangle$ below are then obtained by writing $\langle \lambda_c \lambda_d | B^{(V)} | \lambda_b \rangle$ in the form (3.1).

$$\langle 0, \frac{1}{2} | A | \pm \frac{1}{2} \rangle = \sqrt{\frac{3}{2}} \cdot \xi_{\pm} \cdot (1 \pm z) \cdot (z - \alpha) \quad (C.6b)$$

$$\langle 0, \frac{1}{2} | X^j | \pm \frac{1}{2} \rangle = \pm \sqrt{\frac{3}{2}} \cdot \xi_{\pm} \cdot \left(\pm \frac{q}{3} + z - \alpha \right) \quad (C.6c)$$

$$\langle 0, \frac{1}{2} | X^{3/2} | \pm \frac{1}{2} \rangle = \pm \sqrt{\frac{2}{3}} \cdot \xi_{\pm} \quad (C.6d)$$

$$\begin{aligned} \langle 1, \frac{1}{2} | A | \pm \frac{1}{2} \rangle &= \frac{1}{2} \cdot \frac{E_c}{m_c} \cdot \left[-2 \xi_{\pm} \cdot (z - \alpha)^2 + \xi_{\pm} \cdot \frac{m_c^2}{E_c^2} \cdot (1 - z)^2 + \right. \\ &\quad \left. + (\pm \xi_{\pm} + v_c \eta_{\pm}) \cdot (1 \pm z) \cdot (z - \alpha) - y \cdot \eta_{\mp} \cdot (1 \pm z) \cdot (z - \alpha) \right] \end{aligned} \quad (C.6e)$$

$$\begin{aligned} \langle 1, \frac{1}{2} | X^{1/2} | \pm \frac{1}{2} \rangle &= \\ &= \frac{1}{2} \cdot \frac{E_c}{m_c} \cdot \left[2 \xi_{\pm} \cdot \left(\pm \frac{1}{3} + z - \alpha \right) + \xi_{\pm} \cdot \frac{m_c^2}{E_c^2} \cdot \left(\pm \frac{1}{3} + z \right) + \right. \\ &\quad \left. + (\pm \xi_{\pm} + v_c \eta_{\pm}) \cdot \left(\pm \frac{q}{3} + z - \alpha \right) \pm y \cdot \eta_{\mp} \cdot \left(\pm \frac{q}{3} + z - \alpha \right) \right] \end{aligned} \quad (C.6f)$$

$$\begin{aligned}
\langle 1, 1/2 | \bar{X}^{3/2} | \pm 1/2 \rangle &= \\
&= \frac{1}{3} \cdot \frac{E_c}{m_c} \cdot \left[\left(2 + \frac{m_c^2}{E_c^2} \right) \cdot \xi_{\pm} \mp (\pm \xi_{\pm} + v_c \cdot \eta_{\pm}) \pm y \cdot n \cdot \eta_{\mp} \right] \quad (C.6g)
\end{aligned}$$

$$\begin{aligned}
\langle 1, -1/2 | \bar{H} | \pm 1/2 \rangle &= \\
&= \pm \frac{E_c}{m_c} \cdot \left[\xi_{\mp} \cdot \frac{m_c^2}{E_c^2} \cdot (1 \pm z) - (\mp \xi_{\mp} + v_c \cdot \eta_{\mp}) \cdot (z - \alpha) + y \cdot n \cdot \eta_{\pm} \cdot (z - \alpha) \right] \quad (C.6h)
\end{aligned}$$

$$\begin{aligned}
\langle 1, -1/2 | \bar{X}^{3/2} | \pm 1/2 \rangle &= \\
&= \frac{1}{\sqrt{3}} \cdot \frac{E_c}{m_c} \left[-\xi_{\mp} \cdot \frac{m_c^2}{E_c^2} \pm (\mp \xi_{\mp} + v_c \cdot \eta_{\mp}) \mp y \cdot n \cdot \eta_{\pm} \right] \quad (C.6i)
\end{aligned}$$

$$\begin{aligned}
\langle 2, 1/2 | \bar{H} | \pm 1/2 \rangle &= \\
&= - \left[2 \xi_{\pm} \cdot (z - \alpha) - (\pm \xi_{\pm} + v_c \cdot \eta_{\pm}) \cdot (1 \pm z) + y \cdot n \cdot \eta_{\mp} \cdot (1 \pm z) \right] \quad (C.6j)
\end{aligned}$$

$$\begin{aligned}
\langle 2, 1/2 | \bar{X}^{3/2} | \pm 1/2 \rangle &= \\
&= \mp \frac{1}{\sqrt{3}} \left[\mp 2 \xi_{\pm} + (\pm \xi_{\pm} + v_c \cdot \eta_{\pm}) - y \cdot n \cdot \eta_{\mp} \right] \quad (C.6k)
\end{aligned}$$

$$\langle 2, -1/2 | \bar{H} | \pm 1/2 \rangle = \mp 2 \cdot \left[(\mp \xi_{\mp} + v_c \cdot \eta_{\mp}) - y \cdot n \cdot \eta_{\pm} \right] \quad (C.6l)$$

All $\langle \lambda_c \lambda_d | X^j | \lambda_b \rangle$ not listed vanish.

REFERENCES AND FOOTNOTES

- 1) A.H. Rosenfeld, A. Barbaro-Galtieri, W.H. Barkas, P.L. Bastien, J. Kirz and M. Roos, Revs. Modern Phys. 36, 977 (1964).
- 2) Birmingham - Glasgow - London (I.C.) - Oxford - Chilton (R.H.E.L.) collaboration, Phys. Letters 14, 338 (1965); and private communication from S. Misbahuddin, J.M. Scarr and A.M. Segar.
- 3) For a recent review of the OPE model including references to earlier work see:
J.D. Jackson, invited paper at the topical Conference on Correlation of Particles Emitted in Nuclear Reactions, Gatlinburg, Tennessee, October 15-17, 1964, an elaborate version of which is to be published in Revs. Modern Phys.
- 4) E. Ferrari and F. Selleri, Suppl. Nuovo Cimento 24, 453 (1962) and Nuovo Cimento 27, 1450 (1963).
- 5) J.D. Jackson and H. Pilkuhn, Nuovo Cimento 33, 906 (1964) and errata, Nuovo Cimento 34, 1841 (1964).
- 6) N.J. Sopkovich, Nuovo Cimento 26, 186 (1962).
- 7) K. Gottfried and J.D. Jackson, Nuovo Cimento 34, 735 (1964).
- 8) L. Durand and Y.T. Chiu, Phys. Rev. Letters 12, 399 (1964) and erratum, Phys. Rev. Letters 13, 45 (1964).
- 9) Recent attempts to derive the absorption model outside the framework of potential theory are given in J.S. Ball and W.R. Frazer, Phys. Rev. Letters 14, 746 (1965); and H.D.D. Watson, Phys. Letters 17, 72 (1965);
See also A. Białas and L. Van Hove, CERN preprint TH. 537 (1965); and R. Omnès, Phys. Rev. 137, B649 (1965).

- 10) K. Dietz and H. Pilkuhn, *Nuovo Cimento* 37, 1561 (1965);
R.C. Arnold, *Phys. Rev.* 136, B1388 (1964).
- 11) Our metric is $(+ + + -)$.
- 12) See, e.g., K.J. Barnes, *J. Math. Phys.* 6, 788 (1965),
- 13) J.D. Jackson, *Nuovo Cimento* 34, 1644 (1964).
- 14) For some typical results obtained within the absorption model see, e.g.,
J.D. Jackson, J.T. Donohue, K. Gottfried, R. Keyser and B.E.Y. Svensson,
"Peripheral Production and Decay of Resonances in High Energy Meson-
Nucleon Collisions", to be published in *Phys. Rev.*
- 15) L. Durand and Y.T. Chiu, Yale University preprint (February, 1965),
to be published in *Phys. Rev.*
- 16) M. Jacob and G.C. Wick, *Ann. Phys.* 7, 404 (1959).
- 17) We use the phase conventions of Edmonds, *Angular Momentum in Quantum
Mechanics*, Princeton University Press, 1957; cf. also Ref. 18).
- 18) H. Högaasen and J. Högaasen, CERN preprint TH. 554 (1965), to be
published in *Nuovo Cimento*.
- 19) This last choice produces a strict equality between S and I in the
absence of absorption ($k(j) \equiv 1$) if either β or ϵ vanish.
- 20) M. Bander and G.L. Shaw, Stanford preprint ITP 160 (1965),
to be published in *Phys. Rev.*
- 21) For simplicity we make the discussion in terms of the Bessel functions;
the results are equally valid for the rotation coefficients as functions
of j .
- 22) V. Barger and M. Ebel, *Phys. Rev.* 138, B1148 (1965);
G.A. Ringland and R.J.N. Phillips, *Phys. Letters* 12, 62 (1965).

- 23) Aachen - Birmingham - Bonn - Hamburg - London (I.C.) - München collaboration, Nuovo Cimento 31, 729 (1964).
- 24) CERN - Ecole Polytechnique collaboration, Dubna Conference 1964 and private communication from H. Winzeler.
- 25) W. Galbraith, E.W. Jenkins, T.F. Kycia, B.A. Leontic, R.H. Phillips and A.L. Read, Phys. Rev. 138, B913 (1965).
- 26) S.J. Lindenbaum, Proceedings of the International Conference on Nucleon Structure, Stanford 1963 (edited by R. Hofstadter and L.I. Schiff).
- 27) We have also tried a moderate form factor of the type described in Section 4b, the effect of which is to suppress the theoretical cross-section at large production angles and shifting the values of the density matrix elements towards their values in the unmodified Born approximation.
- 28) Aachen - Berlin - Birmingham - Bonn - Hamburg - London (I.C.) - München collaboration, Phys. Rev. 138, B897 (1965).
- 29) Form factor and absorption model have previously been considered in Refs. 14) and 20).
- 30) Saclay - Orsay - Bari - Bologna collaboration, Phys. Letters 15, 69 (1965).
- 31) Aachen - Berlin - CERN collaboration, private communication from D.R.O. Morrison.
- 32) Without giving a complete list of references, we refer to Refs. 3), 14), 18), 22), B.E.Y. Svensson, Nuovo Cimento 37, 714 (1965); and CERN preprint TH. 549, to be published in Nuovo Cimento; and L. Durand and Y.T. Chiu, Phys. Rev. 137, B1530 (1965), which together cover most reactions treated in the absorption model so far.

- 33) A general helicity state is obtained from the state (A.2) by a rotation¹⁵⁾. Since this is essentially trivial, at least for the integer spin case under study here, we prefer to leave it out.
- 34) Helicity states for spin 2 particles have also been given by Y. Frishman and E. Gotsman, Israel Atomic Energy Commission preprint IA-1040 (1965), to be published in Phys. Rev.
- 35) The notation is such that a four-vector a is written (a_x, a_y, a_z, a_0) .
- 36) K. Gottfried and J.D. Jackson, Nuovo Cimento 33, 309 (1964).
- 37) C. Zemach, Phys. Rev. 133, B1201 (1964);
S.U. Chung, Phys. Rev. 138, B1541 (1965).
- 38) P. Dennery and A. Krzywicki, Phys. Rev. 136, B839 (1964).
- 39) S.M. Berman and M. Jacob, SLAC-PUB-73 (1965); and SLAC-43 (1965), to be published in Phys. Rev.
- 40) W.R. Frazer, J.R. Fulco and F.R. Halpern, Phys. Rev. 136, B1207 (1964).
- 41) Due to the symmetry between the direction of the momentum $\vec{E}$ of particle E and the normal $\vec{N}$ in the decay matrix element (B.7a), the derivation leading to the decay distribution (B.12) can be repeated to give the distribution in the polar angles $\Omega_E = (\vartheta_E, \varphi_E)$ of $\vec{E}$, integrated over the direction angle of the normal around $\vec{A}$. Furthermore, the considerations due to identical particles can also be taken over, i.e., Eq. (B.12), with (ϑ_N, φ_N) replaced by (ϑ_E, φ_E) , gives the distribution in the canonical system of the momentum of that particle, which does not form the vector meson.
- 42) This is the same distribution as for uncorrelated three-pion decay of a 2^+ boson after integration over the whole Dalitz plot^{38),39)}.

FIGURE CAPTIONS

- Figure 1 : Feynman diagram, in the Born approximation, for the process considered in Section 2.
- Figure 2 : The rotation coefficients for $\lambda = \lambda' = \frac{1}{2}$ compared to the corresponding Bessel functions according to Eq. (3.7), and the absorption factors (3.9). The rotation functions are represented by staircase curves through their values at half-integer j values.
- Figure 3 : Comparison in terms of the functions (3.10)-(3.11) of the EPWS and the IPA for $z = 1.2$ and $\cos \theta = 0.8$. The staircase curves representing the EPWS are drawn through the values of the function at half-integer j values. The sum (3.6) and integral (3.8) for $\lambda = \lambda' = \frac{1}{2}$ is given by the area under the respective curves.
- a) No absorption ($k(j) \equiv 1$). The area under the staircase curve is 2.37, under the continuous curve 2.50 units.
 - b) With absorption as given by Eq. (3.9); indices a and b refer respectively to Eqs. (3.9a) and (3.9b). The areas are given in Table 1.
- Figure 4 : Illustrating the effect of the IPA and the different choices of the absorption factor on the cross-section and some density matrix elements in the reaction $\pi^+ p \rightarrow A_2^+ p$ at 4 GeV/c (rho-exchange, $z = 1.27$). The curves marked (a) and (b) are calculated from Eqs. (3.1) and (C.6), using the absorption factors (3.9a), (3.9b) respectively, the curves marked (c) from the same formulae but using the IPA (3.8) with $x_{\min}$ replaced by 0 (for (c') with $x_{\min}$ replaced by $j_{\min}$) and with $k(x)$ given by Eq. (3.9b).

Figure 5 : Differential cross-section for production of f_0 . Experimental results are from Ref. ²³⁾ and Ref. ²⁴⁾. Curves marked (a) are obtained from the BTM, those marked (b) are the absorption model results as described in the text. The left (right) scale applies to the experimental (theoretical) cross-section.

Figure 6 : Some density matrix elements as functions of momentum transfer squared for f_0 in the reaction $\pi^+ n \rightarrow f_0 p$ at 6 GeV/c. Notation as in Fig. 5.

Figure 7 : Differential cross-section for the reaction $\pi^+ p \rightarrow A_2^+ p$ at 4 GeV/c. Experimental results are from Ref. ²⁸⁾. Curve marked (a) is the BTM prediction, curve (b) the absorption model result without form factor, curve (b') is curve (b) multiplied with the square of the form factor (4.3) for $\Lambda = 400$ MeV, curve (c) includes both absorption and a form factor (4.3) as described in the text and curve (c') is curve (c) normalized to the experimental cross-section.

Figure 8 : Some density matrix elements as functions of momentum transfer squared for A_2^+ in the reaction $\pi^+ p \rightarrow A_2^+ p$ at 4 GeV/c. Notation as in Fig. 7.

Figure 9 : Illustrating the decay $2^+ \rightarrow 1^- 0^-$, $1^- \rightarrow 0^- 0^-$.

Figure 10 : Co-ordinate systems used in deriving the decay distribution (B.12).

Σ is the "canonical system", Σ' is obtained from Σ by an Euler rotation $(\varphi_N, \psi_N, 0)$.

FIG.1

FIG.2

a

b

FIG. 3

FIG. 4

b

a

FIG. 6

FIG.7

FIG.8

FIG. 9

FIG.10