CM-P00057295

IMPROVEMENT OF THE GREENBERG-LOW BOUND

A. Martin

CERN -- Geneva

ABSTRACT

It is shown that the Greenberg-Low upper bound on the scattering amplitude $|F(s, \cos\theta=1)| < s^2 \log^2 s$ which is derived from the axiomatic analyticity domain of the scattering amplitude can be replaced by

$$\int \frac{\text{Im } F(s, \cos\theta=1)}{s^3} ds$$

convergent and $|F(s, \cos\theta=1)|/|s|^2 \rightarrow 0$ for $s \rightarrow \infty$ $\varepsilon < \text{Arg } s < \pi - \varepsilon$. Further it is shown that the polynomial boundedness condition inside the Lehmann ellipse can be replaced by the weaker condition $|F(s, \cos\theta=1 + \frac{c}{s^2})| < \exp |s|^\alpha$ with $\alpha < \frac{1}{6}$.

I. INTRODUCTION

From axiomatic field theory which tells us that the fixed energy scattering amplitude $F(s, \cos \theta)$ (s square of c.m. energy, θ c.m. scattering angle) is analytic inside an ellipse with foci -1 and $+1$ and a semi-major axis $\cos \theta_0 = 1 + C/s^{2/3}$, Greenberg and Low²⁾ have shown that if one adds the polynomial boundedness condition

$$|F(s, \cos \theta = 1 + \frac{C'}{s^{2/3}})| < s^N$$

for some $C' < C$ ³⁾ then one finds the following upper bound for the scattering amplitude :

$$|F(s, \cos \theta = 1)| < s^2 (\log s)^2 \quad (1)$$

This result comes from the fact that analyticity and polynomial boundedness limit the number of partial waves contributing effectively to the scattering to

$$L_{\max} = C'' s \log s \quad (2)$$

This cut-off is not sharp. However, by choosing C'' big enough, one can show that the error committed in neglecting the tail of the partial wave distribution is less than s^{-N} , N as large as one wishes.

Now let us show that it is rather unlikely that inequality (1) could be saturated uniformly. First, Khuri and Kinoshita⁴⁾ and Jin and MacDowell⁵⁾ have shown that if for $s \rightarrow \infty$

$$\frac{|\operatorname{Re} F(s, \cos \theta = 1)|}{\operatorname{Im} F(s, \cos \theta = 1)} < \infty$$

the integral

$$\int \frac{\operatorname{Im} F(s, \cos \theta = 1)}{s^3} ds$$

2.

converges and forward dispersion relations contain at most two subtractions. Second, consider now the fixed t region $[t = -2k^2(1 - \cos \theta)]$. It is easy to prove that there the Greenberg-Low bound becomes

$$|F(s, t)| < \left(\frac{s}{t}\right)^{1/4} (L_{\max})^{3/2} = \frac{s^{7/4} (\log s)^{3/2}}{t^{1/4}} \quad (3)$$

So, for any fixed negative t the scattering amplitude satisfies a twice subtracted dispersion relation and we expect by continuity that this will persist for $t = 0$. However, continuity is a dangerous thing in mathematical physics because it could be non-uniform. Therefore we need a more rigorous argument which will be presented in the next Section.

II. IMPROVEMENT OF THE BOUND

What we need to work with is a quantity $\bar{F}(s, t)$ which, like $F(s, t=0)$, possesses some positivity property. We know that since, in a strip of values of t , $F(s, t)$ satisfies dispersion relations,

$$\int_{-|t|}^0 F(s, t) w(t) dt$$

satisfies again dispersion relations. [However, if F possesses the symmetry $s \leftrightarrow u$ where $s+t+u = \sum M^2$ the averaged F will no longer possess this symmetry.]

Now, a simple averaging which possesses this positivity property is

$$\bar{F}(s, t) = \frac{1}{\pi} \int_t^0 \frac{dt' F(s, t')}{\sqrt{t'(t-t')}} \quad , \quad t < 0 \quad (4)$$

Notice that

$$\frac{1}{\pi} \int_t^0 \frac{dt'}{\sqrt{t'(t-t')}} = 1 \quad (5)$$

Then

$$\Im F(s, t) = \frac{1}{\pi} \int_t^0 \frac{dt' \Im F(s, t')}{\sqrt{t'(t-t')}}$$

If we re-express t in terms of $\cos \theta$ we get ⁶⁾

$$\begin{aligned} \frac{\theta[t'(t-t')]}{\sqrt{t'(t-t')}} &= \frac{\theta[(1-\cos \theta')(\cos \theta - \cos \theta')]}{2k^2 \sqrt{(1-\cos \theta')(\cos \theta' - \cos \theta)}} \\ &= \frac{1}{2k^2} \frac{\pi}{2} \sum_0^\infty (2\ell+1) P_\ell(\cos \theta') [P_\ell(\cos \bar{\theta})]^2 \end{aligned} \quad (6)$$

where $\cos \bar{\theta} = (\cos \theta + 1)/2$.

Then, using the variable $\cos \theta'$ and the partial wave expansion for s physical :

$$F(s, \cos \theta) = \frac{\sqrt{s}}{k} \sum (2\ell+1) f_\ell(s) P_\ell(\cos \theta)$$

we get for s physical :

$$\Im \bar{F}(s, t) \equiv \Im F(s, \cos \theta) = \frac{\sqrt{s}}{k} \sum (2\ell+1) \Im f_\ell(s) (P_\ell(\cos \bar{\theta}))^2 \quad (7)$$

which, by unitarity, is positive.

We notice that $\bar{F}(s, t)$, as well as $\text{Im} \bar{F}(s, t)$, for any fixed t are again bounded by $C(t)s^{7/4}(\log s)^{3/2}$. From

$$F(s_1, t) - F(s_2, t) = \frac{(s_1 - s_2)(s_1 - 4 + t + s_2)}{\pi} \int_{4\mu^2}^{\infty} \frac{\text{Im} F(s', t) (2s' - 4 + t)}{(s' - s_1)(s' - s_2)(s' - 4 + t + s_1)(s' - 4 + t + s_2)} ds' \quad (8)$$

$0 < s_1$ and $s_2 < 4\mu^2$, t small enough, we get, using absolute convergence properties to invert orders of integration :

$$\begin{aligned} \bar{F}(s_1, t) - \bar{F}(s_2, t) &= \frac{(s_1 - s_2)(s_1 - 4 + t + s_2)}{\pi} \int_{4\mu^2}^{\infty} \frac{\text{Im} \bar{F}(s', t) (2s' - 4)}{(s' - s_1)(s' - s_2)(s' - 4 + s_1)(s' - 4 + s_2)} ds' \\ &+ \int_t^0 \frac{t' \phi(s_1, s_2, t') dt'}{\sqrt{t'(t - t')}} \end{aligned} \quad (9)$$

$\phi(t', s_1, s_2)$ represents the correction due to the presence of various t' dependent factors outside $A_s(s', t')$. Clearly for $-|t_0| \leq t < 0$, $\phi(t', s_1, s_2)$ is bounded by $(C(s_1, s_2))/(t')^{1/4}$ according to (4) and therefore the whole correction goes to zero as $t \rightarrow 0$.

Now let us take s_1 and $s_2 < 4\mu^2$ but sufficiently close to the threshold. Glaser, Epstein and Bros⁷⁾ have shown that $F(s, t)$ is analytic in a sphere $|t| < \eta$, $|s - 4| < \eta$ minus the physical cut. From this result it is easy to deduce that if s_1, s_2 and t are chosen inside this region of analyticity,

$$\lim_{t \rightarrow 0} \bar{F}(s_1, t) \quad \text{and} \quad \lim_{t \rightarrow 0} \bar{F}(s_2, t)$$

exist and are in fact equal to $F(s_1, 0)$ and $F(s_2, 0)$ respectively.

Hence, since

$$\lim_{t \rightarrow 0} \int_t^0 \frac{t' \phi(s_1, s_2, t') dt'}{\sqrt{t'(t-t')}} = 0$$

we deduce that

$$\lim_{t \rightarrow 0} \left[\int_{4\mu^2}^{\infty} \frac{\text{Im } \bar{F}(s', t) [2s' - 4] ds'}{(s' - s_1)(s' - s_2)(s' - 4 + s_1)(s' - 4 + s_2)} \right] \quad (10)$$

exists.

On the other hand we have for any finite s

$$\lim_{t \rightarrow 0} \bar{A}_s(s', t) = A_s(s', 0) \quad (11)$$

with $A_s(s', t) \equiv \text{Im } F(s', t)$ for real t ,

because we are inside the domain of analyticity of $A_s(s', t)$. Let us show that this, combined with the positivity of $\bar{A}_s(s', t)$, implies that

$$\int_{4\mu^2}^{\infty} \frac{\text{Im } F(s', 0) (2s' - 4) ds'}{(s' - s_1)(s' - s_2)(s' - 4 + s_1)(s' - 4 + s_2)} \quad (12)$$

converges.

Indeed, let us suppose it diverges. Since the integrand is positive, this means that given M arbitrarily large, we can find $s(M)$ such that :

$$\int_{4\mu^2}^{s(M)} \frac{\text{Im } F(s', 0) [2s' - 4] ds'}{(s' - s_1)(s' - s_2)(s' - 4 + s_1)(s' - 4 + s_2)} > M$$

But then from the continuity property (11) we get

$$\int_{4\mu^2}^{s(M)} \frac{\text{Im } \bar{F}(s', t) (2s' - 4) ds'}{(s' - s_1)(s' - s_2)(s' - 4 + s_1)(s' - 4 + s_2)} > M - \varepsilon$$

However, from (10), we have

$$\int_{4\mu^2}^{\infty} \frac{\text{Im } \bar{F}(s', t) (2s' - 4) ds'}{(s' - s_1)(s' - s_2)(s' - 4 + s_1)(s' - 4 + s_2)} < C$$

for $-|t_0| < t \leq 0$.

Hence, taking $M \gg C$ we get that the integral over $\bar{A}_s(s', t)$ from $s(M)$ to ∞ is negative, which is impossible since $\bar{A}_s(s', t)$ is positive. Therefore (12) converges and actually coincides with (10). Hence

$$\int_{4\mu^2}^{\infty} \frac{\text{Im } F(s', t=0) ds'}{s'^3} \quad (13)$$

converges, which corresponds to an average growth less fast than $s^2/\log s$. We have therefore gained essentially a factor $(\log s)^3$ on the Greenberg-Low bound. This, of course, is a rather weak improvement. What matters really is that Eq. (9) holds in the limit of $t=0$, the limit on $\bar{A}_s(s, t)$ being now taken under the integral. Therefore forward dispersion relations need at most two subtractions, and it is easy to show that in all complex directions :

$$\lim_{s \rightarrow \infty} \frac{F(s, t=0)}{s^2} = 0$$

This result is essential in all the work of Khuri and Kinoshita.

Other minor results can be obtained : from

$$\operatorname{Im} F(s, t=0) < \frac{s^2}{\log s}$$

(on an infinite sequence of points) one gets

$$|\operatorname{Im} F(s, t)| < \frac{s^{7/4}}{(\log s)^{3/4}} \quad \text{and} \quad |F(s, t)| < s^{7/4}$$

for fixed t , and

$$|\operatorname{Im} F(s, \cos \theta)| < \frac{s^{3/2}}{(\log s)^{3/4}} \quad \text{and} \quad |F(s, \cos \theta)| < s^{3/2}$$

for fixed angle at the corresponding set of values of s .

III. REMOVAL OF THE POLYNOMIAL BOUNDEDNESS CONDITION

There is very good hope that the polynomial boundedness condition will be established from basic axiomatic postulates (however, maybe not in the strict sense where we use it; an averaging over energies might be necessary). For safety, however, we want to present the weakest conditions under which

$$\int \frac{\operatorname{Im} F(s', t=0) ds'}{s'^3}$$

converges.

8.

Instead of assuming $A_s(s, t = \frac{C}{s}) < s^N$ we now try $\text{Im } F(s, t = \frac{C}{s}) < \exp s^\alpha$.
Then the "maximum" angular momentum is given by

$$\exp s^\alpha / P_{L_{\max}}(1 + \frac{\epsilon}{s^2}) = 1$$

i.e., $L_{\max} = s^{\alpha+1}$.

Then the fixed t bound is given by

$$|F(s, t)| < \frac{s^{\frac{1}{4} + \frac{3}{2}(\alpha+1)}}{t^{\frac{1}{4}}}$$

and it is easy to see that the exponent of s is less than two for $\alpha < \frac{1}{6}$. On the other hand, we have $|F(s, t=0)| < s^{2\alpha+2} = s^{2+1/3}$. Therefore we can repeat in this case all the arguments again and prove that the forward dispersion relations have at most two subtractions, assuming of course, as usual, that in complex directions $|F(s, t=0)|$ is bounded by $C(\epsilon) \exp \epsilon |s|$ for any arbitrarily small ϵ .

ACKNOWLEDGEMENTS

The author wishes to thank Professor V. Glaser and Dr. J.D. Bessis for fruitful discussions.

R E F E R E N C E S

- 1) H. Lehmann, Nuovo Cimento Suppl. 14, 153 (1959).
- 2) O.W. Greenberg and F.E. Low, Phys.Rev. 124, 2047 (1961).
- 3) This version of the argument can be found in
A. Martin, Phys.Rev. 129, 1432 (1963).
- 4) N.N. Khuri and T. Kinoshita, Phys.Rev. 137, B720 (1965):
Phys.Rev.Letters 14, 84 (1965):
A. Martin, Phys.Letters 15, 76 (1965).
- 5) Y.S. Jin and S.W. MacDowell, Phys.Rev. 138, B1279 (1965).
- 6) See, for instance, M.L. Goldberger and K.M. Watson, "Collision
Theory", Wiley, New York (1964), p. 595.
- 7) J. Bros, H. Epstein and V. Glaser, Nuovo Cimento 31, 1265 (1964).