CM-P00057759

175chives .

Ref. TH. 844

ON ENERGY AND MOMENTUM TRANSFER DEPENDENCEIN $\pi p \rightarrow \rho N^*$

J.T. Donohue +)

CERN - Geneva

A B S T R A C T

The effects on energy and momentum transfer dependence of cross-sections, caused by the variation of $\Delta_{\min}^2$ with incident momentum and external masses, are investigated using a simple model for the differential cross-section. It is found that at intermediate momenta, $\sim 3-6$ GeV/c the incident momentum dependence of the total cross-section for $\pi p \rightarrow \rho N^*$ differs appreciably from that of the differential cross-section. The consequences for the Δ^2 distributions of a zero near $\Delta^2 = 0$ are examined, and it is found that plotting in $\Delta^2 - \Delta_{\min}^2$ should be used to resolve this question.

+) National Science Foundation Postdoctoral Fellow.

Recently experimental data on the quasi-two-body reactions : $\pi^+ p \rightarrow \rho^0 N^{*++}$ and ωN^{*++} in the energy range 3-4 GeV/c have been reported ¹⁾. Reference 1) also contains a comparison with experiments at other energies on $\rho^0 N^{*++}$ production, and the authors conclude that the energy dependence of the total cross-section is inconsistent with meson exchange as the dominant production mechanism. Three experimental results are :

1. in the range of incident momenta from 2 to 8 GeV/c, the $\rho^0 N^{*++}$ total cross-section decreases approximately as $P_{inc}^{-0.5}$;
2. the exponential slope in Δ^2 of $d\sigma/d\Delta^2$ is found to increase from ~ 9 (GeV/c)² at 2.5 GeV/c to ~ 20 at 7 GeV/c;
3. the differential cross-section rounds off at small momentum transfer, $\Delta^2 \leq 0.1$ (GeV/c)².

According to the compilation of Morrison ²⁾, the energy dependence of the total cross-sections for reactions believed to be dominated by meson exchange is typically between $P_{inc}^{-1.5}$ and $P_{inc}^{-2.0}$ ^{*)}. The decay angular distributions in the $\rho^0 N^{*++}$ reaction are reasonably close to the predictions of simple pion exchange, and it seems difficult to reconcile the observed energy dependence with that suggested by pion exchange, namely P_{inc}^{-2} .

The $\rho^0 N^{*++}$ reaction has been studied using the absorption model ^{3),4),5)} and more recently in Regge pole analyses ^{6),7),8),9)}. In the latter analyses, the reaction is of considerable importance, since it should provide evidence for problems associated with reggeized pion exchange, such as conspiracy or kinematical zeros. Frautschi and Jones ⁷⁾ have analyzed the reaction using the results of Cohen-Tannoudji, Morel and Navelet ¹⁰⁾, and suggest that the dominant amplitude has a zero at the

^{*)} This is an empirical conclusion, based on the fact that quasi-two-body reactions in which the quantum numbers of some mesons can be exchanged show similar energy dependence.

It is clear from Fig. 1 that the momentum dependence of the total cross-section can differ appreciably from that of the differential cross-section, at least for intermediate momenta, $\sim 3-6$ GeV/c. This effect, which is caused by the kinematic inaccessibility of small Δ^2 , is enhanced when A , the exponential Δ^2 slope, is large. This sensitivity to A can be seen by comparing curves a. and c., and may explain why ωN^* and K^*N^* production cross-sections show $\sim P_{\text{inc}}^{-1.7}$ dependence ¹⁾. The exponential slopes for these reactions are ~ 5 (GeV/c)⁻² for ωN^* at 8 GeV/c, while for K^*N^* $A = 6$ (GeV/c)⁻² at 3 GeV/c and 11 (GeV/c)⁻² at 5 GeV/c ²⁾. The phase space effects are thus less severe in these reactions than in ρN^* ³⁾.

The differential cross-section $d\sigma/d\Delta^2$ is plotted v. Δ^2 in Figs. 2a and 2b for the incident momenta 3, 5, 8 and 16 GeV/c. In Fig. 2a we have taken $F(\Delta^2) = 1$, while in Fig. 2b, $m_0^2 = m_\pi^2$, which corresponds to the amplitude having a zero at $\Delta^2 = 0$ and a pole at $\Delta^2 = -m_\pi^2$. The vertical scale is arbitrary and we have chosen $N=2$, $A=13$ (GeV/c)⁻². As expected, the dependence of Δ_{min}^2 on the external masses causes the differential cross-section to turn over at small Δ^2 , even though the amplitude has no zero there. A close comparison of Figs. 2a and 2b shows that there is a characteristic difference between the curves calculated without and with the zero and pole structure. In the former the shape of the turn-over at small Δ^2 becomes narrower as the incident momentum increases. This is understood from Eq. (1). The width of the turn-over is essentially a measure of the variation of Δ_{min}^2 between low and high $\pi\pi$ and πp masses. This variation decreases as P_{inc}^{-1} , so the Δ^2 interval in which the turn-over occurs shrinks as the momentum increases. If the amplitude has a zero, the shape of the rounding off is roughly independent of incident momentum as seen from Fig. 2b. However, in view of the small intervals of Δ^2 in which the turn-over occurs, ~ 0.05 (GeV/c)², experiments of high precision would be needed to distinguish the shapes of the curves in Fig. 2a from those in Fig. 2b.

One method which has been suggested to overcome the phase space difficulties is to plot $d\sigma/d\Delta^{2'}$, where $\Delta^{2'}$ is $\Delta^2 - \Delta_{\min}^2$, $\Delta_{\min}^2$ being the minimum Δ^2 for the external particle masses. In this variable the phase space does not cause curves to round off, since $\Delta^{2'} = 0$ is allowed for all values of momenta and external masses. The price paid for this is that a fixed $\Delta^{2'}$ corresponds to some interval in Δ^2 , and any Δ^2 singularities of the amplitude will be slightly smeared out in $\Delta^{2'}$. We have investigated numerically the problem of plotting $d\sigma/d\Delta^{2'}$ vs $\Delta^{2'}$ for the same curves as appear in Figs. 2a and 2b¹⁴⁾. These are shown in Fig. 3. Those curves with $F(\Delta^2) = 1$ remain pure exponentials in $\Delta^{2'}$, and hence appear as straight lines. Those curves with the zero and pole structure show a rounding off at small $\Delta^{2'}$, the rounding becoming more apparent with increasing incident momentum. From Fig. 3 we conclude that the test of whether the amplitude has a zero near $\Delta^2 = 0$ becomes feasible for incident moment ~ 8 GeV/c, and that $\Delta^{2'}$ is the relevant variable for such tests.

Finally it should be noted that the variation of $\Delta_{\min}^2$ with P_{inc} may cause an apparent increase in the exponential slope A . If one assumes that the actual $d\sigma/d\Delta^2$ shape is concave, i.e., the curvature is negative, in the usual semi-log plot, then as P_{inc} increases, the decrease of $\Delta_{\min}^2$ allows more of the sharply peaked region to contribute. In the linear approximation this would simulate an increase of A with P_{inc} , as is observed experimentally for ρN^* .

Just as this work was completed there appeared a paper by Wolf¹⁵⁾, in which the reaction $\pi p \rightarrow \rho N^*$ is studied in a one-pion-exchange model, with form factors. In the paper the widths of the ρ and N^* are used, and good fits to the differential cross-section are obtained at various energies. Thus Wolf finds a conclusion identical to ours, that the energy dependence of the total cross-section can differ appreciably from that of the differential cross-section.

The author would like to thank Dr. D.R.O. Morrison for helpful discussions.

R E F E R E N C E S

- 1) D. Brown, G. Gidal, R.W. Birge, R. Bacastow, S.Y. Fung, W. Jackson and R. Pu, Phys.Rev.Letters 19, 664 (1967).
- 2) D.R.O. Morrison, Proceedings of the Stony Brook Conference on High Energy Two-Body Reactions, Stony Brook, April 1966 (unpublished); Phys.Letters 22, 528 (1966).
- 3) J.D. Jackson, J.T. Donohue, K. Gottfried, R. Keyser and B.E.Y. Svensson, Phys.Rev. 139, B428 (1965).
- 4) Aachen-Berlin-CERN Collaboration, Phys.Letters 19, 608 (1965).
- 5) P.C.M. Yock and D. Gordon, Phys.Rev. 157 1362 (1967).
- 6) I. Bender, V. Linke and H.J. Rothe, "Study of Angular Decay Correlations for the Process $\pi + N \rightarrow \rho + N^*$ in the Regge Pole Model I", University of Heidelberg preprint (1967).
- 7) S. Frautschi and L. Jones, "The Reggeization of Pion Exchange in Production Processes", California Institute of Technology preprint 68-135 (1967).
- 8) M. Le Bellac, "Pion Conspiracy in $\pi N \rightarrow \rho \Delta$ ", Argonne National Laboratory preprint (1967).
- 9) B. Haber, U. Maor, G. Yekutieli and E. Gotsman, "Investigation of the Pion Trajectory in a Regge Pole Model", Weizmann Institute preprint (1967).
- 10) G. Cohen-Tannoudji, A. Morel and H. Navelet, "Kinematical Singularities, Crossing Matrix and Kinematical Constraints for Two-Body Helicity Amplitudes", Saclay preprint (1967).
- 11) This form allows us to investigate the effect on Δ^2 distributions of a zero and pole structure in the amplitude. Less violent behaviour as well as factors like $(\cos^2 \theta/2)$ and p wave Breit-Wigner curves would be over-refinements in this crude model.
- 12) It should be remarked that the large widths of the ρ and N^* do not have much influence on the shapes in Fig. 1. Very similar energy dependence is obtained by using the central values of $m_{\pi\pi}$ and $m_{p\pi}$ instead of integrating over masses.

8.

- 13) For $Kp \rightarrow K^*N^*$ there is some disagreement between experimental results for the total cross-section. The experiments of the Brussels-CERN collaboration at 3, 3.5 and 5 GeV/c, R. George et al., Nuovo Cimento 49, 9 (1967), find a momentum dependence $\sim P_{\text{inc}}^{-1}$. But as is seen in Fig. 1 of Ref. 1), experiments at higher energies suggest a more rapid decrease with energy.
- 14) To obtain the dependence on Δ^2 , one replaces Δ^2 in Eq. (2) by $(\Delta^2 + \Delta_{\text{min}}^2(P_{\text{inc}}, m_{\pi\pi}^2, m_{p\pi}^2))$, and integrates over the $\pi\bar{\pi}$ and $p\pi$ masses.
- 15) G. Wolf, Phys.Rev.Letters 19, 925 (1967).

FIGURE CAPTIONS

Figure 1 σ v. P_{inc} for the reaction $\pi p \rightarrow \rho^0 N^{*++}$, as calculated using Eq. (2). The parameters used are :
 $m_\rho = 0.765$ GeV, $m_{N^*} = 1.236$ GeV, $\Gamma_\rho = \Gamma_{N^*} = 0.125$ GeV,
 $0.680 \leq m_{\pi\pi} \leq 0.860$ GeV, $1.16 \leq m_{p\pi} \leq 1.13$ GeV. The quantities (A, N) have the values :
 curve a : $(13(\text{GeV}/c)^{-2}, 2)$,
 curve b : $(13(\text{GeV}/c)^{-2}, 1.7)$,
 curve c : $(6(\text{GeV}/c)^{-2}, 2)$.
 The curves have been normalized to 0.5 mb at 8 GeV/c, and straight lines corresponding to $P_{inc}^{-0.5, -1, -2.0}$ are shown for comparison.

Figure 2a $d\sigma/d\Delta^2$ v. Δ^2 as calculated from Eq. (2) with $F(\Delta^2) = 1$, at $P_{inc} = 3, 5, 8$ and 16 GeV/c. The parameters are as for curve a. of Fig. 1. The vertical scale is arbitrary, but the same for all incident momenta.

Figure 2b $d\sigma/d\Delta^2$ v. Δ^2 as in Fig. 2a, but with $F(\Delta^2) = (1 - m_\pi^2 / (m_\pi^2 + \Delta^2))^2$.

Figure 3 The quantity $d\sigma/d\Delta'^2$ v. Δ'^2 . The straight lines correspond to the curves in Fig. 2a, while the solid curves correspond to those of Fig. 2b.

FIG.1

FIG. 3